Colonial choanoflagellate isolated from Mono Lake harbors a microbiome

Hake, K. H.[1, 2], West, P.T.[3], McDonald, K.[4], Laundon, D.[5], Feng, C.[1], Burkhardt, P.[5, 6], Richter, D.[7], Banfield, J.F.[3], and King, N.[1,*]

[1]Howard Hughes Medical Institute and Department of Molecular and Cell Biology, University of California, Berkeley, CA, USA

[2]Present address: Calico Life Sciences, South San Francisco, CA, USA

[3]Department of Environmental Science, Policy, & Management, University of California, Berkeley, CA, USA

[4]Electron Microscopy Laboratory, University of California, Berkeley, CA, USA

[5]Marine Biological Association of the United Kingdom, Plymouth, United Kingdom

[6]Sars International Centre for Molecular Marine Biology, University of Bergen, Bergen, Norway

[7]Institut de Biologia Evolutiva (CSIC-Universitat Pompeu Fabra), Barcelona, Spain

[*]Send correspondence to nking@berkeley.edu

ABSTRACT

Choanoflagellates offer key insights into bacterial influences on the origin and early evolution of animals. Here we report the isolation and characterization of a new colonial choanoflagellate species, *Salpingoeca monosierra,* that, unlike previously characterized species, harbors a stable microbiome. *S. monosierra* was isolated from Mono Lake, California and forms large spherical colonies that are more than an order of magnitude larger than those formed by the closely related *S. rosetta*. By designing fluorescence *in situ* hybridization probes from metagenomic sequences, we found that *S. monosierra* colonies are colonized by members of the halotolerant and closely related *Saccharospirillaceae* and *Oceanospirillaceae,* as well as purple sulfur bacteria (*Ectothiorhodospiraceae*) and non-sulfur *Rhodobacteraceae.* This relatively simple microbiome in a close relative of animals presents a new experimental model for investigating the evolution of stable interactions among eukaryotes and bacteria.

IMPORTANCE

The animals and bacteria of Mono Lake (California) have evolved diverse strategies for surviving the hypersaline, alkaline, arsenic-rich environment. We sought to investigate whether the closest living relatives of animals, the choanoflagellates, exist among the relatively limited diversity of organisms in Mono Lake. We repeatedly isolated members of a single species of choanoflagellate, which we have named *Salpingoeca monosierra,* suggesting that it is a stable and abundant part of the ecosystem. Characterization of *S. monosierra* revealed that it forms large spherical colonies that each contain a microbiome, providing an opportunity to investigate the evolution of stable physical associations between eukaryotes and bacteria.

OBSERVATION

A newly identified choanoflagellate species forms large colonies that contain a microbiome

Choanoflagellates are the closest living relatives of animals and, as such, provide insights into the origin of key features of animals, including animal multicellularity and cell biology (1, 2). Over a series of four sampling trips to Mono Lake, California (Fig. 1A; Table S1) we collected single-celled choanoflagellates and large spherical choanoflagellate colonies, many of which were hollow (Fig. 1B) and resembled the blastula stage of animal development. In colonies and single cells, each cell had the typical collar complex observed in other choanoflagellates: an apical flagellum surrounded by a collar of microvilli (1, 2). In these "rosette" colonies, the cells were oriented with the basal pole of each cell pointing inwards and the apical flagellum facing out (Fig. 1). To study the Mono Lake choanoflagellates in greater detail, we established clonal cultures from ten independent isolates, two of which were each started from a single-celled choanoflagellate and the remaining eight of which were each started from a single colony (Table S1). The two strains started from single-celled choanoflagellates, isolates ML1.1 and ML1.2, took on the colonial morphology observed in the other isolates after culturing in the laboratory, suggesting that the colonies and single cells isolated from Mono Lake could belong to the same species. We are aware of no prior reports of choanoflagellates having been cultured from any alkaline soda lake, including Mono Lake.

The 18S rDNA genes for six of the Mono Lake isolates were sequenced and found to be >99% identical (Table S1). Phylogenetic analysis revealed that all of the isolates are members of a single choanoflagellate species and closely related to species in the genus *Salpingoeca* (Fig. 1C and Fig. S1). Therefore, we propose the name *Salpingoeca monosierra,* with the species name inspired by the location of Mono Lake in the Sierra Nevada mountain range. (See Supplemental Methods for a formal species description.) In phylogenetic analyses based either on 18S rDNA (Fig. 1C and Fig. S1) or on six highly conserved genes (Fig. S2)(3)(3)(3), we found that the closest relatives of *S. monosierra* are the emerging model choanoflagellate *S. rosetta* (4-8)(4–8)(4–8) and the largely uncharacterized choanoflagellate *Microstomoeca roanoka* (3, 9).

Although *S. monosierra* and *S. rosetta* form rosette-shaped spherical colonies, they differ greatly in size. *S. rosetta* colonies range from 10-30 μm in diameter while *S. monosierra* forms among the largest choanoflagellate colonies observed (1, 10), with a single culture exhibiting colony sizes spanning from 10-120 μm in diameter (Fig. 1D-F). Unlike the rosettes of *S. rosetta*, in which the basal poles of cells are closely apposed in the rosette center (5, 7), cells in large *S. monosierra* rosettes form a shell on the surface of a seemingly hollow sphere. Inside the ostensibly hollow sphere, a branched network of extracellular matrix connects the basal poles of all cells (Fig. S3.)

Upon staining *S. monosierra* with the DNA dye Hoechst 33342, we observed the expected toroidal nuclei in each choanoflagellate cell (11), but were surprised to detect

Hoechst-positive material in the interior of *S. monosierra* colonies (Fig. 2A, A'). Transmission electron microscopy revealed the presence of 1 µm and smaller cells with diverse morphologies bounded by cell walls in the centers of rosettes (Fig. 2B, B'; Fig. S4). Together, these observations led us to hypothesize that the centers of *S. monosierra* colonies contain bacteria.

By performing hybridization chain reaction fluorescence *in situ* hybridization (HCR-FISH (12–14)) with a broad-spectrum probe of bacterial 16S rRNA, EUB338 (15), we confirmed that the cells in the center of colonies are bacteria (Fig. 2C). A second probe that specifically targeted 16S rRNA sequences from Gammaproteobacteria,

110   GAM42a, revealed that the majority of the bacteria inside the colonies are Gammaproteobacteria (Fig. 2C')(16). Finally, by incubating *S. monosierra* cultures with fluorescently labeled D-amino acids, which are specifically incorporated into the cell walls of growing bacteria, we found that the bacteria in *S. monosierra* colonies are alive

115   and growing (Fig. S5)(17). Therefore, *S. monosierra* contains a microbiome (18).

To visualize the spatial distribution of choanoflagellate and bacterial cells in a representative colony, we generated a 3D reconstruction from serial sections imaged by TEM. The colony contained 70 choanoflagellate cells that were tightly packed, forming a

120   largely continuous monolayer of cells (Fig. 2D). As observed by immunofluorescence microscopy (Figs. 2A and 2A'), all cells were highly polarized and oriented with their apical flagella and collars extending away from the centroid of the rosette. Many cells were connected by fine intercellular bridges (Fig. S6) that have been previously observed in other colonial choanoflagellates, including *S. rosetta* (5, 19).

The 3D reconstruction also revealed at least 200 bacterial cells in the center of the rosette (Fig. 2D', 2D''), some of which were physically associated with and wrapped around the choanoflagellate ECM (Fig. S7). A small number of bacterial cells were observed between the lateral surfaces of choanoflagellate cells, although it was not

130   possible to determine whether they were entering or exiting the colony (Fig. 2D''; Fig. S8). Colonies failed to incorporate bacteria-sized bovine serum albumin (BSA)-coated latex microspheres (0.2 µm and 1 µm) into their centers, suggesting that environmental bacteria may not be capable of passively accessing the centers of *S. monosierra* colonies (Fig. S9).

Gammaproteobacteria and Alphaproteobacteria in the *S. monosierra* microbiome

We next sought to identify which bacteria comprise the microbiomes of *S. monosierra.* To identify candidate bacteria for which to design FISH probes, we first

140   sequenced and assembled metagenomes and 16S rDNA sequences from choanoflagellate-enriched samples and from environmental bacteria-enriched samples. These samples were derived from two co-cultures of *S. monosierra* with Mono Lake bacteria, ML2.1E and ML2.1G (Fig. S10), with the enrichment for choanoflagellates or bacteria performed by centrifugation. A total of 24 different bacterial species were

145   identified via two complementary bioinformatic approaches (EukRep Metagenomic Analysis and EMIRGE 16S rRNA Analysis; Table S2), of which 22 species were present

in fractions enriched with *S. monosierra* colonies (Table S3). The phylogenetic relationships among these and other bacterial species were determined based on analysis of 16S rDNA sequences (Fig. 2E and S11).

The 22 bacterial species detected in cultures with *S. monosierra* may have co-sedimented with the *S. monosierra* colonies due their community-structure densities (e.g. biofilms), a transient association with the choanoflagellate colonies (e.g. as prey), or through a stable association with the choanoflagellate colonies. Upon investigation by FISH microscopy, we detected ten or eleven of these species in the centers of *S. monosierra* colonies (Table S4, Fig. S12). (The uncertainty regarding the precise number of choanoflagellate-associated bacterial species stems from the inability to disambiguate 16S rDNA sequences corresponding to one or two of the species.) Of these microbiome bacteria, nine were Gammaproteobacteria from the families *Oceanospirillaceae* (Fig. S12A; OceaML1, OceaML2, OceaML3, OceaML4, OceaML4)*, Ectothiorhodospiraceae* (Fig. S12B; EctoML1, EctoML2, EctoML3, EctoML4), and *Saccharospirillaceae* (Fig. S12C; SaccML), matching our original observation that the majority of the bacteria were Gammaproteobacteria (Fig. 2C, C'). The remaining species was a *Roseinatronobacter* sp*. (RoseML; Alphaproteobacteria) (Fig. S12D). The microbiome bacteria exhibited an array of morphologies, from long and filamentous to rod shaped (Fig. S4 and Fig. S13). Intriguingly, with the exception of OceaML3, which was exclusively detected inside *S. monosierra* colonies of ML2.1E (Fig. S14), all other microbiome species identified in this study were detected both inside and outside the colonies.

In animals, the microbiome often contains a core set of host-adapted bacteria that are present in many or all individuals in a host species, as well as a more flexible set of bacteria that may be found in only a subset of individuals (20–22). To identify core members of the *S. monosierra* microbiome, we measured the frequency with which colonies of strain ML2.1EC contained or lacked a number of representative microbiome bacteria (Fig. S15A) and estimated the abundance of each species relative to the total microbiome (Fig. S15B). Only one bacterium tested, OceaML1, was found in the microbiome of all *S. monosierra* colonies (Fig. S15A). The other most frequently observed members of the microbiome were SaccML (93.3% of colonies), EctoML3 (91.8% of colonies) and EctoML1 (82.4% of colonies; Fig. S15A). Two other Gammaproteobacterial species, OceaML2 and EctoML2, were found in ~50 - 60% of colonies, while the Alphaproteobacterium RoseML was found in only 13.9% of rosettes.

The most common resident of the *S. monosierra* microbiome, OceaML1, was also the most abundant, representing on average 66.4% of the total bacterial load per colony (Fig. S15B). The next most abundant bacteria, found in >80% of colonies, represented approximately one-third of the bacterial biomass in the *S. monosierra* microbiomes in which they were found: SaccML (30.3% of total bacteria), EctoML2 (36.7%), and EctoML3 (32.5%). When present, the remaining bacterial species -- OceaML2, EctoML1, and RoseML -- averaged 10% of the microbiome or less. As the species with the highest frequency and abundance, OceaML1 appears to be a core member of the *S. monosierra* microbiome. However, other symbionts detected in *S. monosierra* (e.g.

SaccML and OceaML2-4) are close relatives of OceaML1 and may engage in similar metabolic or developmental functions in their interactions with *S. monosierra* (23).

Discussion

Interactions with bacteria are essential to choanoflagellate nutrition and life
200 history. Bacteria are the primary food source for choanoflagellates, and the choanoflagellate *S. rosetta* responds to different secreted bacterial cues to undergo either multicellular developmental or mating (24–26). We report the isolation and characterization of a new choanoflagellate species, *S. monosierra,* that forms large colonies and contains a microbiome consisting of at least ten different bacterial
205 symbionts. To our knowledge, this is the first example of a stable interaction between choanoflagellates and ectosymbiotic bacteria. Future studies will be important to determine how the *S. monosierra* colonies and their bacterial symbionts interact.

We detected 10 – 11 bacterial species from four different families
210 (*Saccharospirillaceae, Oceanospirillaceae, Ectothiorhodospiraceae,* and *Rhodobacteraceae)* in the *S. monosierra* microbiome. Comparisons with other symbioses suggest that their interactions with *S. monosierra* may relate to metabolism and detoxification of environmental sulfur. Such functions have been reported for *Oceanospirillaceae* species that are symbionts of diverse marine animals, from corals to
215 snails (27–34) and for *Ectothiorhodospiraceae* that are symbionts of diverse ciliates and animals (35–38).

Although the composition of the animal gut microbiome often varies between individuals in a species, many host species harbor a core set of microbes, with which
220 they participate in stable metabolic interactions and may coevolve (21, 22). Indeed, choanoflagellates express homologs of Toll-like receptors that, in animals, mediate interactions with gut bacteria to maintain homeostasis (9). In *S. monosierra*, OceaML1 was found in 100% of rosettes assessed and was the most abundant bacterium in the microbiome, suggesting that its interactions with *S. monosierra* may be essential for *S.*
225 *monosierra* biology. Interestingly, the related bacterium *Endozoicomonas sp.* (Order Oceanospirillales), is a core member of the gut microbiome of the ascidian *Ciona intestinalis*, where it represents up to 54% of the bacterial biomass (30).

S. monosierra and its associated microbiome provide a unique opportunity to
230 characterize a new symbiotic interaction between a single choanoflagellate species and a microbial community. Due to the phylogenetic relevance of choanoflagellates, this symbiotic relationship has the potential to illuminate the ancestry of and mechanisms underlying stable associations between animals and bacteria, colonization of organisms by diverse microbial communities, and insights into one of the most complex animal-
235 bacterial interactions, the animal gut microbiome.

Data Availability: Genbank accession numbers for bacterial 16S rDNA sequences are listed in Supplementary Data File 1. Sequences for *S. monosierra* 18S rDNA, EFL, and Hsp90 are currently in submission to Genbank and can be found in Supplementary Data File 2.

Author contributions: *P.W. performed metagenomic analysis, K.M. did the transmission electron microscopy, D.L. and P.B. did the TEM-reconstruction, and C.F. helped culture M. monosierra. K.H. performed all other experiments and analysis. D.J.R. originally isolated S. monosierra and contributed to manuscript editing. J.B. and N.K. contributed to project leadership, experimental design, figure design, writing, and editing.*

Acknowledgements

We thank Tarja Hoffmeyer for help with establishing some of the original *S. monosierra* cultures. We thank Emil Ruff and Julia Schwartzman for assistance with early CARD-FISH experiments through the Physiology Course and Microbial Diversity Course at the Marine Biological Laboratory in Woods Hole. We thank Michael VanNieuwenhze for the fluorescent D-amino acids. We thank the following for helpful discussions and research support: Reef Aldayafleh, David Booth, Ben Larson, Frank Nitsche, Monika Sigg, and Arielle Woznica. We thank Alain Garcia De Las Bayanos and Karen Carniol for valuable feedback on the manuscript.

Supplement Contents

265   **Text S1:** Supplemental Materials and Methods

       Figure S1: Independent choanoflagellate isolates from Mono Lake are the same
       species
       Figure S2: Six-gene phylogenetic tree of *S. monosierra* and other choanoflagellates.
270   **Figure S3.** *S. monosierra* colonies contain an extracellular matrix (ECM).
       Figure S4: Bacterial symbionts in *S. monosierra* colonies exhibit a range of
       morphologies.
       Figure S5: Bacteria inside *S. monosierra* colonies are alive and growing.
       Figure S6: Intercellular bridges connect cells in *S. monosierra* colonies.
275   **Figure S7:** Bacterial symbionts physically associate with and wrap around the
       choanoflagellate ECM.
       Figure S8: Bacteria are found wedged between the lateral surfaces of *S. monosierra*
       cells.
       Figure S9: *S. monosierra* colonies cannot be passively penetrated by sub-micron
280   particles.
       Figure S10: Schematic describing the establishment of *S. monosierra* strains.
       Figure S11: Phylogenetic analysis of 16S rRNA tree of EMIRGE sequences.
       Figure S12: Diverse bacteria are found in the microbiome of *S. monosierra.*
       Figure S13: *S. monosierra* symbionts exhibit filamentous and rod morphologies.
285   **Figure S14:** The symbiont OceaML3 is present exclusively inside rosettes.
       Figure S15: OceaML1 is a core member of the *S. monosierra* microbiome.

       Table S1: Date, location, phenotype, and isolate designations for *Salpingoeca
       monosierra* isolates.
290   **Table S2:** Nine genera of bacteria identified in *S. monosierra* cultures through two
       independent analyses: comparison of ribosomal proteins detected through
       metagenomic assembly and by 16s rRNA assembly and analysis.
       Table S3: Predicted targets of HCR-FISH probes based on 16S rRNA sequences.
       Bacteria in bold were identified inside *S. monosierra* colonies.
295   **Table S4:** Full length probes, with spacer and initiator sequences, used for HCR-FISH.
       Table S5: Growth Media Recipes.

Figure 1

Figure 1. A new colony-forming choanoflagellate isolated from Mono Lake.
(A) Choanoflagellates were collected from two sampling sites (asterisks) near the shore of Mono Lake, California. (Modified from map at monolake.org.) (B) *S. monosierra* forms
305 large colonies (DIC image). Scale bar = 50µm. (C) *S. monosierra* (shown in bold) is a craspedid choanoflagellate and closely related to *S. rosetta* and *M. roanoka.* Phylogeny based on 18S sequences. A phylogeny with 18S sequences from other colonial choanoflagellate isolates from Mono Lake show they fall within a single clade (Figure S1). Animal branches were collapsed to save space. Bootstrap values indicated.
310 *Previous identifier: *Savillea micropora* (3)*. (D-E) Two representative colonies reveal the extremes of the *S. monosierra* colony size range (D, 58 µm diameter; E, 19 µm diameter; scale bar = 20 µm). In *S. monosierra* colonies, each cell is oriented with its apical flagellum (white; labeled with anti-tubulin antibody) and the apical collar of microvilli (red; stained with phalloidin) pointing out. Nuclei (cyan) were visualized with
315 the DNA-stain Hoechst 33342. (F) Colonies of *S. monosierra* span from 10 µm in diameter, a size comparable to that of small *S. rosetta* colonies, to 120 µm, over an order of magnitude larger. Diameters of *S. monosierra* and *S. rosetta* colonies were plotted as a violin plot; median indicated as thick black line. Diameters of representative colonies indicated as colored bars behind violin plot (D, red bar; E, blue bar).
320

Figure 2

Figure 2. *S. monosierra* colonies are filled with bacteria.

(A, A') The center of a representative *S. monosierra* colony, shown as a maximum
intensity projection (A) and optical z-section (A'), contains DNA (revealed by Hoechst
33342 staining; cyan). Apical flagella were labeled with anti-tubulin antibody (white);
microvilli were stained with phalloidin (red). Hoechst 33342 staining (cyan) revealed the
toroidal choanoflagellate nuclei along the colony perimeter and an amorphous cloud of
DNA sitting within the central cavity formed by the monolayer of choanoflagellate cells.
(B-B') Thin section through a representative *S. monosierra* colony, imaged by
transmission electron microscopy (TEM), revealed the presence of small cells in the
central cavity. (B') Inset (box; panel B') reveals that the interior cells are each
surrounded by a cell wall. (C-C''') The small cells inside *S. monosierra* colonies are
bacteria, as revealed by hybridization with a broad spectrum 16S rRNA probe (C,
green) and a probe targeting Gammaproteobacteria (C', red). Choanoflagellate nuclei
and bacterial nucleoids were revealed by staining with Hoechst (C'', cyan). (C''') Merge
of panels C – C''. Scale bar for all = 5 μm. (D-D'') 3D reconstruction of a 70-cell *S.
monosierra* choanoflagellate colony from transmission electron micrographs of serial
ultrathin sections revealed that the bacteria are closely associated with and wrapped
around the ECM inside the colony. (D) Whole colony view. (D') Cut-away view of colony
center. Color code: cell bodies (cyan); microvilli (orange); flagella (green); bacteria (red);
ECM (white); intercellular bridges (yellow, see also Fig. S6); filopodia (purple). (D'')
Setting all features of the colony transparent, except the bacteria and ECM, revealed
the presence of bacteria positioned between the lateral surfaces of choanoflagellate
cells (brackets, see also Fig. S8). (E) Unrooted phylogenetic tree representing bacterial
diversity modified from (39), illustrated to indicate the phylogenetic placement of
bacteria co-cultured from Mono Lake with *S. monosierra.* The bacteria belonged to four
major classes: Spirochaetia, Alphaproteobacteria, Gammaproteobacteria, and
Bacteroidetes, however the bacteria found associated with *S. monosierra* colonies
came only from Alphaproteobacteria and Gammaproteobacteria. Circles represent the
phylogenetic placement of non-symbionts (white) and symbionts (*Oceanospirillaceae
sp.,* magenta*; Saccharospirillaceae sp.,* green*; Ectothiorhodospiraceae sp.,* blue*;
Roseinatronobacter sp.,* orange). See also Figs. S11 and S12.

References

1. Leadbeater BSC. 2015. The Choanoflagellates: Evolution, Ecology, and Biology. Cambridge University Press, Cambridge, UK.

2. Brunet T, King N. 2017. The origin of animal multicellularity and cell differentiation. Dev Cell 43:124–140.

365   3. Carr M, Richter DJ, Fozouni P, Smith TJ, Jeuck A, Leadbeater BSC, Nitsche F. 2017. A six-gene phylogeny provides new insights into choanoflagellate evolution. Mol Phylogenet Evol 107:166–178.

4. Booth DS, Szmidt-Middleton H, King N. 2018. Transfection of choanoflagellates illuminates their cell biology and the ancestry of animal septins. Mol Biol Cell
370   https://doi.org/10.1091/mbc.e18-08-0514.

5. Dayel MJ, Alegado RA, Fairclough SR, Levin TC, Nichols SA, McDonald K, King N. 2011. Cell differentiation and morphogenesis in the colony-forming choanoflagellate< i> Salpingoeca rosetta</i>. Dev Biol 357:73–82.

6. Fairclough SR, Chen Z, Kramer E, Zeng Q, Young S, Robertson HM, Begovic E,
375   Richter DJ, Russ C, Westbrook MJ, others. 2013. Premetazoan genome evolution and the regulation of cell differentiation in the choanoflagellate Salpingoeca rosetta. Genome Biol 14:R15.

7. Levin TC, Greaney AJ, Wetzel L, King N, Sánchez Alvarado A. 2014. The rosetteless gene controls development in the choanoflagellate S. rosetta. Elife 3.

380   8. Wetzel LA, Levin TC, Hulett RE, Chan D, King GA, Aldayafleh R, Booth DS, Sigg MA, King N. 2018. Predicted glycosyltransferases promote development and prevent spurious cell clumping in the choanoflagellate S. Rosetta. Elife https://doi.org/10.7554/eLife.41482.

9. Richter DJ, Fozouni P, Eisen M, King N. 2018. Gene family innovation,
385   conservation and loss on the animal stem lineage. Elife 7:e34226.

10. Larson BT, Ruiz-Herrero T, Lee S, Kumar S, Mahadevan L, King N. 2019. Biophysical principles of choanoflagellate self-organization. bioRxiv 659698.

11. Burkhardt P, Grønborg M, McDonald K, Sulur T, Wang Q, King N. 2014. Evolutionary insights into premetazoan functions of the neuronal protein Homer.
390   Mol Biol Evol msu178.

12. Choi HMT, Calvert CR, Husain N, Huss D, Barsi JC, Deverman BE, Hunter RC, Kato M, Lee SM, Abelin ACT, Rosenthal AZ, Akbari OS, Li Y, Hay BA, Sternberg PW, Patterson PH, Davidson EH, Mazmanian SK, Prober DA, van de Rijn M, Leadbetter JR, Newman DK, Readhead C, Bronner ME, Wold B, Lansford R,

395   Sauka-Spengler T, Fraser SE, Pierce NA. 2016. Mapping a multiplexed zoo of mRNA expression. Development 143:3632–3637.

13.   DePas WH, Starwalt-Lee R, Van Sambeek L, Kumar SR, Gradinaru V, Newman DK. 2016. Exposing the three-dimensional biogeography and metabolic states of pathogens in cystic fibrosis sputum via hydrogel embedding, clearing, and rRNA
400   labeling. MBio 7:e00796----16.

14.   Pernthaler A, Pernthaler J, Amann R. 2002. Fluorescence in situ hybridization and catalyzed reporter deposition for the identification of marine bacteria. Appl Environ Microbiol 68:3094–3101.

15.   Amann RI, Binder BJ, Olson RJ, Chisholm SW, Devereux R, Stahl DA. 1990.
405   Combination of 16S rRNA-targeted oligonucleotide probes with flow cytometry for analyzing mixed microbial populations. Appl Environ Microbiol 56:1919–1925.

16.   Manz W, Amann R, Ludwig W, Wagner M, Schleifer KH. 1992. Phylogenetic Oligodeoxynucleotide Probes for the Major Subclasses of Proteobacteria: Problems and Solutions. Syst Appl Microbiol 15:593–600.

410 17.   Kuru E, Hughes HV, Brown PJ, Hall E, Tekkam S, Cava F, De Pedro MA, Brun Y V, Vannieuwenhze MS. 2012. In situ probing of newly synthesized peptidoglycan in live bacteria with fluorescent D-amino acids. Angew Chemie - Int Ed 51:12519–12523.

18.   WHIPPS JM. 1991. Effects of Mycoparasites on Sclerotia-Forming Fungi, p. 129–
415   140. In BEEMSTER, ABR, BOLLEN, GJ, GERLAGH, M, RUISSEN, MA, SCHIPPERS, B, TEMPEL, A (eds.), Biotic Interactions and Soil-Borne Diseases. Elsevier.

19.   Laundon D, Larson BT, McDonald K, King N, Burkhardt P. 2019. The architecture of cell differentiation in choanoflagellates and sponge choanocytes. PLOS Biol
420   17:e3000226.

20.   Shapira M. 2016. Gut Microbiotas and Host Evolution: Scaling Up Symbiosis. Trends Ecol Evol 31:539–549.

21.   Roeselers G, Mittge EK, Stephens WZ, Parichy DM, Cavanaugh CM, Guillemin K, Rawls JF. 2011. Evidence for a core gut microbiota in the zebrafish. ISME J
425   5:1595–1608.

22.   Huse SM, Ye Y, Zhou Y, Fodor AA. 2012. A core human microbiome as viewed through 16S rRNA sequence clusters. PLoS One 7.

23.   Doolittle WF, Booth A. 2017. It's the song, not the singer: an exploration of holobiosis and evolutionary theory. Biol Philos 32:5–24.

430 24.   Alegado RA, Brown LW, Cao S, Dermenjian RK, Zuzow R, Fairclough SR, Clardy

J, King N. 2012. A bacterial sulfonolipid triggers multicellular development in the closest living relatives of animals. Elife 1:e00013.

25. Woznica A, Cantley AM, Beemelmanns C, Freinkman E, Clardy J, King N. 2016. Bacterial lipids activate, synergize, and inhibit a developmental switch in choanoflagellates. Proc Natl Acad Sci 113:7894–7899.

26. Woznica A, Gerdt JP, Hulett RE, Clardy J, King N. 2017. Mating in the closest living relatives of animals is induced by a bacterial chondroitinase. Cell 170:1175–1183.

27. Bayer T, Neave MJ, Alsheikh-Hussain A, Aranda M, Yum LK, Mincer T, Hughen K, Apprill A, Voolstra CR. 2013. The microbiome of the red sea coral Stylophora pistillata is dominated by tissue-associated Endozoicomonas bacteria. Appl Environ Microbiol 79:4759–4762.

28. Beinart RA, Nyholm S V, Dubilier N, Girguis PR. 2014. Intracellular Oceanospirillales inhabit the gills of the hydrothermal vent snail Alviniconcha with chemosynthetic, γ-Proteobacterial symbionts. Environ Microbiol Rep 6:656–664.

29. Carlos C, Torres TT, Ottoboni LMM. 2013. Bacterial communities and species-specific associations with the mucus of Brazilian coral species. Sci Rep 3.

30. Dishaw LJ, Flores-Torres J, Lax S, Gemayel K, Leigh B, Melillo D, Mueller MG, Natale L, Zucchetti I, De Santis R, Pinto MR, Litman GW, Gilbert JA. 2014. The gut of geographically disparate Ciona intestinalis harbors a core microbiota. PLoS One 9:e93386.

31. Goffredi SK, Orphan VJ, Rouse GW, Jahnke L, Embaye T, Turk K, Lee R, Vrijenhoek RC. 2005. Evolutionary innovation: A bone-eating marine symbiosis. Environ Microbiol 7:1369–1378.

32. Kurahashi M, Yokota A. 2007. Endozoicomonas elysicola gen. nov., sp. nov., a γ-proteobacterium isolated from the sea slug Elysia ornata. Syst Appl Microbiol 30:202–206.

33. Morrow KM, Moss AG, Chadwick NE, Liles MR. 2012. Bacterial Associates of Two Caribbean Coral Species Reveal Species-Specific Distribution and Geographic Variability. Appl Environ Microbiol 78:6438–6449.

34. Vezzulli L, Pezzati E, Huete-Stauffer C, Pruzzo C, Cerrano C. 2013. 16SrDNA Pyrosequencing of the Mediterranean Gorgonian Paramuricea clavata Reveals a Link among Alterations in Bacterial Holobiont Members, Anthropogenic Influence and Disease Outbreaks. PLoS One 8:e67745.

35. Bayer C, Heindl NR, Rinke C, Lücker S, Ott JA, Bulgheresi S. 2009. Molecular characterization of the symbionts associated with marine nematodes of the genus

Robbea. Environ Microbiol Rep 1:136–144.

36. Payne SH, Loomis WF. 2006. Retention and loss of amino acid biosynthetic
470   pathways based on analysis of whole-genome sequences. Eukaryot Cell 5:272–
        276.

37. Rinke C, Schmitz-Esser S, Stoecker K, Nussbaumer AD, Molnar DA, Vanura K,
        Wagner M, Horn M, Ott JA, Bright M. 2006. "Candidatus Thiobios zoothamnicoli",
        an ectosymbiotic bacterium covering the giant marine ciliate Zoothamnium
475   niveum. Appl Environ Microbiol 72:2014–2021.

38. Tian RM, Wang Y, Bougouffa S, Gao ZM, Cai L, Bajic V, Qian PY. 2014. Genomic
        analysis reveals versatile heterotrophic capacity of a potentially symbiotic sulfur-
        oxidizing bacterium in sponge. Environ Microbiol 16:3548–3561.

39. Hug LA, Baker BJ, Anantharaman K, Brown CT, Probst AJ, Castelle CJ,
480   Butterfield CN, Hernsdorf AW, Amano Y, Ise K, Suzuki Y, Dudek N, Relman DA,
        Finstad KM, Amundson R, Thomas BC, Banfield JF. 2016. A new view of the tree
        of life. Nat Microbiol 1:nmicrobiol201648.