Title

The ascending arousal system shapes low-dimensional neural dynamics to mediate awareness

of intrinsic cognitive states

Authors

Brandon Munn[1,2,†*], Eli J. Müller[1,2,†], Gabriel Wainstein[1,2], and James M. Shine[1,2,*]

Affiliations

1   Complex Systems Research Group, The University of Sydney, Sydney, NSW, Australia

2   Brain and Mind Centre, The University of Sydney, Sydney, NSW, Australia

*   Corresponding authors: brandon.munn@sydney.edu.au & mac.shine@sydney.edu.au

†   Denotes equal contribution

Abstract

Models of cognitive function typically focus on the cerebral cortex and hence overlook

functional links to subcortical structures. This view neglects the highly-conserved ascending

arousal system's role and the computational capacities it provides the brain. In this study, we

test the hypothesis that the ascending arousal system modulates cortical neural gain to alter

the low-dimensional energy landscape of cortical dynamics. Our analyses of spontaneous

functional magnetic resonance imaging data and phasic bursts in both locus coeruleus and

basal forebrain demonstrate precise time-locked relationships between brainstem activity,

low-dimensional energy landscapes, network topology, and spatiotemporal travelling waves.

We extend our analysis to a cohort of experienced meditators and demonstrate locus

coeruleus-mediated network dynamics were associated with internal shifts in conscious

awareness. Together, these results present a novel view of brain organization that highlights

the ascending arousal system's role in shaping both the dynamics of the cerebral cortex and

conscious awareness.

Main Text

It is often difficult to see the forest for the trees, but to fully understand a concept typically

involves an accurate depiction of both. That is, we need to comprehend not only the detailed

workings of a specific system, but also how that system functions within a broader context

of interacting parts. Modern theories of whole-brain function exemplify this challenge. For

instance, activity in the brain has been shown to incorporate signatures of both local

computational specificity (e.g., specialized regions within the cerebral cortex) as well as

system-wide integration (e.g., the interactions between the cortex and the rest of the brain)[1,2].

Anatomical evidence suggests that the balance between integration and segregation is

mediated in part by the relatively fixed white matter connections between cerebral cortical

regions[1] – local connectivity motifs support segregated activity, whereas the axonal, re-

entrant connections between regions act to integrate the distributed signals via a highly

interconnected structural backbone[3]. However, how the human brain is also capable of

remarkable contextual flexibility given this relatively fixed connectivity remains poorly

understood.

During cognitive tasks, neural activity rapidly reconfigures the functional large-scale network

architecture of the brain to facilitate coordination between otherwise segregated cortical

regions. Precisely how this flexibility is implemented in the brain without altering structural

connectivity remains an open question in systems neuroscience. Although it is often

overlooked in theories of whole brain function, the neuromodulatory ascending arousal

system is well-placed to mediate this role[4]. The arousal system is comprised of a range of

nuclei spread across the brainstem and forebrain that send wide-reaching axons to the rest of

the central nervous system[5]. At their target sites, arousal neurons release neuromodulatory

neurotransmitters that shape and constrain a region's processing mode – altering their

excitability and responsivity without necessarily causing them to fire an action potential[4,6]. As

a result, subtle changes in the concentration of neuromodulatory chemicals can cause

massive alterations in the dynamics of the target regions, leading to nonlinear effects on the

coordinated patterns of activity that emerge from 'simple' neuronal circuits[4].

The ascending arousal system also contains substantial heterogeneity – unique cell

populations project in diverse ways to the cerebral cortex and release distinct

neurotransmitters. One key dichotomy is the distinction between adrenergic

neuromodulation (predominantly via the locus coeruleus [LC]), which promotes arousal and

exploratory behaviour[7], and cholinergic neuromodulation (such as via the basal nucleus of

Meynert [BNM]), which is associated with attentional focus and vigilance[8]. These highly

interconnected[9] structures both promote wakefulness and arousal[10,11], albeit via distinct

topological projections to the cerebral cortex: the LC projects in a diffuse manner that

crosses typical specialist boundaries, whereas the BNM projects in a more targeted, region-

specific manner[12] (Fig. 1A). The two systems have also been linked with distinct and

complimentary computational principles: the noradrenergic LC is presumed to modulate

interactions between neurons (response gain)[13], whereas the cholinergic BNM is presumed to

facilitate divisive normalization (multiplicative gain)[14]. Based on these anatomical and

computational features, we have hypothesized that the interaction between these two

neuromodulatory systems is crucial for mediating the dynamic, flexible balance between

integration and segregation in the brain[15].

Another crucial feature of the ascending arousal system is that the number of neurons that

project to the cerebral cortex is several orders of magnitude smaller than those that project

back to the brainstem and forebrain[16–18]. Based on this feature, we further hypothesize that

shifts in arousal are realized through a low-dimensional modulation of the ongoing neural

activity ('brain state')[17]. Conceptually, low-dimensional neural dynamics can be depicted as

evolving on a brain state energy landscape[19], where the energy of a given state corresponds

to the occurrence probability, e.g. high energy brain states have a low occurrence probability

(and *v. v.*). That is brain states evolve along the energy landscape topography, much like a ball

rolls under the influence of gravity down a valley and requires energy to traverse up a hill,

this corresponds to an evolution towards an attractive or repulsive brain state, respectively.

This technique can resolve what might otherwise be obscured states of attraction (and

repulsion) in a multi-stable system and has been successfully applied to the dynamics of

spiking neurons[20,21], blood oxygenation level dependent (BOLD) functional magnetic

resonance imaging (fMRI)[22,23], and magnetoencephalography (MEG)[24]. The approach offers

several conceptual advances, but perhaps most importantly, it renders the otherwise

daunting task of systems-level interpretation relatively intuitive. Importantly, this framework

is not a mere analogy[30], as the topography of the energy landscape shares a 1-to-1

correspondence with the generative equations required to synthesize realistic neural
timeseries data[25]. In this manuscript, we test these ideas by combining high-resolution resting
state fMRI data with analytic techniques from the study of complex systems.
96
Results

To begin with, we extracted time series data from major subcortical hubs within the
noradrenergic LC[9] (Fig. 1A, red) and cholinergic BNM[26] (Fig. 1A, green) systems from 59
healthy participants who had undergone high-resolution, 7T resting-state functional
magnetic resonance imaging (fMRI; 2 mm^3 voxels; TR = 586 ms repetition time). Given the
known spatiotemporal interactions between the ascending arousal system and fluctuations in
cerebrospinal fluid, we first controlled for activity fluctuations in the nearby fourth ventricle,
which contains no neural structures, but nonetheless can cause alterations in the BOLD
signal over time. We next accounted for nearby gray-matter signals, by regressing the signal
from the nearby pontine nuclei. Using the residuals from these regressions from the LC
signal, τ_{LC}, and the BMN signal, τ_{BNM}, we focused on the difference between these signals
(τ_{LC-BNM} and τ_{BNM-LC}; concatenated across subjects) and then identified time points associated
with phasic bursts of LC activity that led to sustained adrenergic (versus cholinergic)
influence over evolving brain state dynamics (and *v.v.* for phasic bursts of BNM; see
Methods). Importantly, the phasic mode of firing within the noradrenergic arousal system
has been specifically linked to systemic influences that occur on time-scales relevant to
cognitive function[8,27]. Tracking the mean cortical BOLD response around these peaks
identified a spatiotemporal travelling wave (Fig. 1B; velocity = 0.13ms^{-1}) that propagated
from frontal to sensory cortices and tracked closely with the known path of the dorsal
noradrenergic bundle[9], albeit with a preserved 'island' within the parietal operculum (Fig.
1B). A block-resampling null model was applied to ensure that the results were not due to
spatial-autocorrelation (p < 0.05; see Methods). These results can be inverted for BNM
activity (relative to LC) as $\tau_{BNM-LC} = -\tau_{LC-BNM}$. Furthermore, these results confirm that
coordinated macroscale activity patterns align to fluctuations in activity within the ascending
arousal system of the brainstem[28].

Figure 1. Sympathetic activity precedes network-level integration. A) regional time series were extracted

from the locus coeruleus (red), which is thought to alter response gain, and the basal nucleus of Meynert

(green), which is thought to alter multiplicative gain, and compared to BOLD signal and topological signatures

during the resting state; B) we observed a anterior-to-posterior traveling wave (velocity ~ 0.13ms⁻¹) following

peaks in $\tau_{LC\text{-}BNM}$, which are shown on both the left (LH) and right (RH) hemispheres of a cortical flat map; C)

the lagged cross-correlation between   and PC – dotted line depicts the zero-lag correlation, and the

black lines depict the upper (lower) bounds of a block-resampled null model (95% CI); D) mean cortical

participation coefficient (PC) preceding (left) and following (right) the zero-lagged   value, only the

right hemisphere is shown (mirrored for 'Post'); E) the participation coefficient following peak   was

higher in the right- (red) vs. the left- (blue) hemisphere (p < 0.001; green bar).

Time-varying network topology

Based on previous empirical[29], modelling[30] and theoretical[15] work, we predicted that phasic

bursts in $\tau_{LC\text{-}BNM}$ would facilitate network-level integration by modulating increased neural

gain among regions distributed across the cerebral cortex. As predicted, we observed a

strong positive correlation between $\tau_{LC\text{-}BNM}$ and network-level integration (p < 0.05, block-

resampling null model; Fig. 1C) across the brain (Fig. 1D). An increase in phasic activity

within the LC (relative to the BNM) preceded an increase in the mean level of integration

within the cerebral cortex that was dominated by the frontoparietal cortices (Fig. 1D;

parcellated according to the 17 resting-state networks identified in[31]). Interestingly, this

global integration was opposed by a relative topological segregation of limbic, visual, and

motor cortices (Fig. S2). This increase in the synchronisation of the frontoparietal cortices

following an increase in sensory-limbic coordination and LC activity may reflect arousal-

enhanced processing of sensory stimuli[32,33]. Furthermore, regional integration occurred

earlier in the right- vs. the left-hemisphere (p<0.001; Fig. 1E), which is consistent with the

known anatomical bias of the LC system[34,35]. Together, these findings provide robust

evidence for the hypothesis that the balance between ascending noradrenergic and

cholinergic tone facilitates a transition towards topological integration across the

frontoparietal network of the brain[15].

Neuromodulation of the Energy Landscape

The results of our initial analysis demonstrate that coordinated distributed activity in the

cortex align with changes in small groups of neuromodulatory cells activity in the brainstem

and forebrain, which in turn are proposed to constrain brain dynamics onto a low-

dimensional energy landscape (Fig. 2A). The effects of noradrenaline and acetylcholine can

also be easily viewed through this lens: by integrating the brain, noradrenaline should flatten

the energy landscape (Fig. 2A, red) facilitating otherwise unlikely brain state transitions,

whereas in contrast, the segregative nature of cholinergic activity should act to deepen

energy valleys (Fig. 2A, green) decreasing the likelihood of a brain state transition. In

previous work, we have shown a correspondence between low-dimensional brain state

dynamics across multiple cognitive tasks and the heterogenous expression of metabotropic

neuromodulatory receptors[17]. This implies that neuromodulators act similar to catalysts in

chemical reactions, which lower (or raise) the activation energy (E_A) required to transform

chemicals from one steady state (or energy well) to another (Fig. 2B). In the context of the

interconnected, heterarchical networks that comprise the cerebral cortex, this would have

the effect of flattening (or deepening) the energy landscape, promoting variable (or rigid)

brain states[36] (Fig. 2A).

Figure 2 – LC and BNM mediated shifts in energy landscape brain state space dynamics. A) an energy

landscape, which defines the energy required to move between different brain states: by increasing response

gain, the LC should flatten the energy landscape (red); by increasing multiplicative gain, the BNM should

deepen the energy wells (green); B) the topography of the energy landscape can be conceptualized as similar to

the activation energy (E_A) that must be overcome in order to convert one chemical to another; C) Empirical

BOLD trajectory energy as a function of MSD and TR of the baseline activity (E_A, black) and after phasic

bursts in LC (E_{LC}, red) and BNM (E_{BNM}, green) – relative to the baseline energy landscape phasic bursts in LC

(E_{LC} - E_A, red inset) lead to a flattening or reduction of the energy landscape, whereas peaks in BNM (E_{BNM} -

E_A, green inset) lead to a raising of the energy landscape. D) Empirical activation energy as a function of MSD

averaged over lags   TR during base baseline activity (E_A, Left) and following phasic bursts in LC

(E_{LC}, red) and BNM (E_{BNM}, green).

To elucidate the role of phasic activity from the neuromodulatory system in modifying the
energy landscape, we first estimated the energy of BOLD signal transitions across the
cerebral cortex. Importantly, the term 'energy' here is used in reference to its definition in
statistical physics and hence does not represent the biological use of the term, which instead
stands for the energy used by the brain to maintain or change neural activity. Specifically, we
define the energy landscape, E, as the natural logarithm of the inverse probability of
observing a given BOLD MSD at a given time-lag t, $P(MSD, t)$, calculated as $E =$
$\ln \frac{1}{P(MSD,t)}$, where $MSD_{t,t_0} = \langle |x_{t_0+t} - x_{t_0}|^2 \rangle_r$ is the mean-squared displacement (MSD) of
BOLD signal, $x_t = [x_{1,t}, x_{2,t}, ..., x_{r,t}]$ across r voxels and t is the number of time-lags of
size TR from a reference timepoint t_0 [20]. The probability of a BOLD signal transition,
$P(MSD, t)$, was estimated from the sampled MSD_{t,t_0}, and we used a Gaussian kernel
density estimation $P(MSD, t) = \frac{1}{4n} \sum_{i=1}^{n} K\left(\frac{MSD_{t,i}}{4}\right)$, where $K(u) = \frac{1}{2\sqrt{\pi}} e^{-\frac{1}{2}u^2}$ (see
Methods). Our analysis is consistent with the statistical mechanics interpretation that the
energy of a given state, E_σ, and its probability are related $P_\sigma = \frac{1}{Z} e^{-\frac{E_\sigma}{T}}$, where Z is the
normalisation function and T is a scaling factor equivalent to temperature in
thermodynamics, where we set $T = 1$ and $Z = 1$ [20]. In this framework, a highly probable
relative change in BOLD (as quantified by the MSD) corresponds to a relatively low energy
transition (i.e., low E_A), whereas an infrequently visited state will require the most energy
(i.e., high E_A).
203
By treating energy as inversely proportional to the probability of brain state occurrence, our
approach resembles other studies that have been applied to spiking dynamics of neuronal
populations, spiking neurons[20,21], BOLD fMRI[22,23], MEG[24], and natural scene[37]. However,
these studies binarized continuous signals to reduce the brain state space (to 2^r states),
however this approach requires the fitting of a threshold, which can be problematic in
continuously recorded data. In contrast, our approach reduces the dimensionality by
analysing the likelihood of a change in BOLD activity (i.e., the MSD), and thus retains the
dimensionality of the underlying signal without the need for thresholding. This approach
overcomes a major limitation inherent to previous energy landscape studies that require a
large sample size to sufficiently sample the brain state space.

With this in mind, we turned our attention to the relationship between the ascending arousal

system dynamics and the MSD energy landscape. To test the hypothesis that the

neuromodulatory system alters the topography of the energy landscape, we calculated BOLD

MSD energetics following phasic bursts of both LC relative to BNM (τ_{LC-BNM}), E_{LC}, and

BNM relative to LC (τ_{BNM-LC}), E_{BNM}, i.e., t_0 was the onset of a phasic burst, and compared

these to sampled brain evolutions without large changes in LC and BNM arousal i.e., t_0 was

all timepoints outside of a phasic burst in LC and BNM, analogous to the baseline energy

landscape E_A. We identified phasic bursts as peaks in the second derivative of the arousal

BOLD signals τ_{LC-BNM} and τ_{BNM-LC} that lead to a sustained increase in BOLD activity for

each individual (see Methods) and using these criteria, we identified 148 τ_{LC-BNM} time points

and 130 τ_{BNM-LC} time points.

The energy landscapes for these three states are defined by the energy for a given BOLD

MSD at a given TR delay. Figure 2C demonstrates the baseline energy landscape (Fig. 2C,

black), which corresponds to the reaction pathway in Fig. 2B, and the MSD energy

landscape following phasic bursts in the LC (Fig. 2C, red) and BNM (Fig. 2C, green). These

figures demonstrate an MSD energy landscape across displacement and time, wherein the

energy relates to the likelihood of seeing a given mean change in bold activity (i.e., MSD) at a

given temporal displacement (i.e., TR). For example, all the MSD energy landscapes have a

high energy peak for large MSD at a short timescale as it is extremely unlikely that the bold

activity would change significantly (quantified by a large MSD) in one TR (~0.5s), and

neuromodulation increases the energy of such an initial change. The utility of the MSD

energy landscape can be seen when comparing large phasic bursts of LC and BNM relative

to baseline fluctuations. We found the largest change occurs around 10-15 TR (~6-9s)

following a phasic burst that typically corresponds to a peak in the LC or BNM BOLD

signal. At this ~6-9s temporal delay we see direct evidence that a phasic burst of LC

flattened the energy landscape (decreased the energy relative to baseline Fig. 2C, red inset),

thus making previously unlikely large MSD trajectories far more probable (Fig. 2D red),

whereas a phasic burst of BNM activity (increased energy relative to baseline Fig. 2C, green

inset) caused the energy landscape to be elevated, thus promoting local trajectories, and

making large MSD deviations unlikely (Fig. 2D green). These patterns are analogous to

modulating a physical landscape in which towns sit within valleys separated by impassable

mountains – when BNM is high, the towns remain isolated, whereas when LC is high, the

towns are separated by easily navigated rolling plains and transitions between towns (novel

combinations of consecutive brain states) can be easily realised.

We next asked whether LC and BNM combined synergistically to alter the energy landscape.

To achieve this, we isolated simultaneous phasic peaks in both LC and BNM (τ_{LC+BNM}). We

found that the LC + BNM energy landscape differed from either independent LC or BNM

activation, shifting the brain state into divergent regimes than could be explained by the

HRF. By comparing the MSD energy topography for a given TR slice we found that the

landscape switched from an anti- to de-correlation with the HRF. In other words, the

cooperative behaviour between the noradrenergic and cholinergic systems allowed the brain

to reach unique BOLD MSDs that neither could facilitate individually. To examine how

simultaneous LC+BNM activity altered the energy landscape, we compared the energy

relative to the two individual landscapes. As demonstrated in Fig. S3, the energy landscape

following phasic bursts of LC+BNM differed in magnitude from that expected from a linear

superposition of the LC and BNM energy landscape – i.e., LC+BNM ≠ (LC) + (BNM).

Furthermore, to explore the dominance of either LC or BNM in this signal, we minimised

the relationship LC+BNM = αLC + βBNM (conditional upon α and β being positive

constants) and found that $\alpha = 0.16$ and $\beta = 0.84$ gave the best match to the LC+BNM

energy landscape. That is, the BNM dynamics dominates the simultaneous LC+BNM energy

landscape, which is consistent with the unidirectional synaptic projections from the LC that

synapse upon the BNM on their way through to the cortex[12], and suggests that phasic

LC+BNM bursts may be initiated by the LC in order to elicit a cascade of BNM activity.

Conscious awareness of shifts in BOLD state

Interpreting the relationship between neuroimaging data and conscious awareness is

notoriously challenging. For instance, it is currently not possible to directly determine the

contents of self-directed thought without intervening, and thus altering, the contents of

consciousness[38]. Although we can't determine the contents of consciousness directly, we can

use task designs to modulate the state of consciousness. To this end, we leveraged data from

a group of 14 expert meditators who were asked to meditate during an fMRI scanning

session[39], and to press a button when they noticed that their focus had drifted from their

breath (Fig. 3A). At this point, there is a mismatch between expectation and conscious

awareness, which is an internal state that has been previously linked to the activation of the

noradrenergic system, both in theoretical[40,41] and computational[42] work. Based on these

studies, we predicted that the switch in internal conscious awareness would be facilitated by

increases in locus coeruleus-mediated integration and subsequent reconfiguration of low-

dimensional brain states. Analysing time-resolved network data with a finite impulse

response model, we observed a peak in locus coeruleus activity (Fig. 3B), TR-to-TR mean

squared displacement (Fig. 3C) and elevated network-level integration (Fig. 3D) surrounding

the change in conscious awareness (all p_{PERM} < 0.05; 95% CI of null distribution). These

results confirm that the locus coeruleus mediates energy landscape reconfigurations and that

these changes modulate internal states of conscious awareness.

Figure 3 – Awareness of intrinsic state changes. A) participants performing breath-awareness meditation

(Focus; blue) were trained to respond with a Button Press (orange) when they became Aware (purple) that they

had become Distracted (i.e., their attention had wandered from their breath) and to then re-focus their

Attention (blue) on their breath; B) we observed a peak in τ_{LC-BNM} (i.e., LC > BNM; red) ~4 seconds before
the button press, which then returned to low levels (i.e., BNM > LC) in the 2-4 seconds following the button
press; C) the Mean Squared Displacement (MSD; dark orange) of TR-to-TR BOLD signal was increased above
null values around the peak in τ_{LC-BNM}, as well as following the re-establishment of attentional focus (in
panels B & C, grey shading depicts 95% CI of block-resampled null distribution); D) we observed a peak in
mean participation coefficient (PC) ~4 seconds (2 TRs) prior to the Button Press during the task. All: grey
shading depicts 97.5th and 2.5th percentile of null distribution i.e., outside grey shading indicates a value
different than null [p < 0.05]; and lower: red shading represents SEM error bars).

Discussion

Our results provide evidence for an arousal-mediated macroscopic network and energy
landscape reconfiguration which tracks with moment-to-moment alterations in conscious
awareness. By tracking fluctuations in BOLD signal within the noradrenergic LC and the
cholinergic BNM, we were able to demonstrate fundamental ways in which the low-
dimensional, dynamic, and topological signature of cortical dynamics was related to changes
within the ascending arousal system. Furthermore, we demonstrated a link between these
dynamic reconfigurations and alterations in conscious awareness in a cohort of experienced
meditators. In this way, our results provide a novel, systems-level perspective on the
distributed dynamics of the human brain.

There is growing evidence that distributed neural dynamics in the brain are well described by
relatively low-dimensional models[17,18,43–45], however the biological constraints that impose
these features on the brain remain poorly understood. Due to the low number of cells in the
arousal system and their broad projections to the rest of the brain, we theorized that
neuromodulatory regions are well-placed to shape and constrain the vast number of neurons
in the cerebral cortex into low-dimensional dynamic modes. Our results support this
prediction by showing that patterns of activity in key regions within the brainstem and
forebrain relate to fundamental alterations in a dynamically evolving energy landscape. In
other words, neural state space trajectories are a powerful framework that extends beyond
that of mere analogy, and the ascending arousal system is well-placed to mediate
deformations in the energy landscape.

Much in the same way that there are many different reference frames for navigation – e.g.,
egocentric (i.e., straight, left, right directions), which is independent of the environment; and
allocentric (i.e., following compass directions and visual cues), which is dependent on the
environment – we can interrogate energy landscapes using different vantage points on
BOLD dynamics. Our displacement framework is consistent with an egocentric (or 'first-
person') frame of reference, wherein MSD is used to track BOLD trajectories from an initial
state which maps out the topology of the energy landscape (i.e., a BOLD MSD implies a
BOLD trajectory). Nonetheless, the method does not distinguish between two different
neural trajectories that possess the same MSD. In comparison, other methods have evaluated
the energy landscape for a given pre-defined state estimated from thresholded BOLD
timeseries[22,23] a framework consistent with an allocentric (or 'third-person') reference frame.
This framework has the advantage of calculating energy for a given state, however, it also
requires substantial exploration of the state-space – which is typically unfeasible – or the
need to resort to severe coarse-graining (such as the binarization of BOLD activity) which
further diminishes interpretability. Furthermore, the allocentric view does not provide
insights into the transitions between each energy state, whereas this information is inherent
to the egocentric reference. Along these lines, we found that the egocentric reference frame
clearly demonstrated the flattening and deepening of the energy landscape, providing indirect
evidence that the ascending arousal system is well set-up to control brain-state dynamics
'egocentrically' (as opposed to specific neural activity patterns). Nevertheless, given
improvements in recording length and novel analytic techniques to probe the brains
dynamical landscape, we expect that the field will ultimately discover even more optimal
mappings between neurobiology and low-dimensional brain state dynamics.

The results of our state-space analysis have important implications for the biological
mechanisms underlying cognition. For instance, the concept of locus coeruleus-mediated
energy landscape flattening is reminiscent of the α1 receptor-mediated notion of a 'network
reset'[40]. By increasing response gain (Fig. 1A) through the modulation of second-messenger
cascades[4], noradrenaline released by the LC would augment inter-regional coordination[30].
Importantly, this capacity could confer adaptive benefits across a spectrum, potentially
facilitating the formation of flexible coalitions in precise cognitive contexts[46], while also
forcing a broader landscape flattening (i.e., a 'reset') in the context of large, unexpected

changes[27,40].   Similarly, the idea that phasic cholinergic bursts deepens energy wells is
consistent with the idea that the cholinergic system instantiates divisive normalization within
the cerebral cortex[14]. Numerous cognitive neuroscience studies have shown that heightened
acetylcholine levels correspond to improvements in attentional precision[8,8]. By deepening
energy wells, acetylcholine from the BNM could ensure that the brain remains within a
particular state and is hence not diluted by other (potentially distracting) brain states.
Determining the specific rules that govern the links between the neuromodulation of the
energy landscape and cognitive function[47–49] is of paramount importance, particularly given
the highly integrated and degenerate nature of the ascending arousal system[50].

Our results also provide a systems-level perspective on an emerging corpus of work that
details the microscopic circuit level mechanisms responsible for conscious phenomena[51]. In
particular, a number of recent studies have highlighted the intersection between the axonal
projections of the ascending arousal system and pyramidal cell dendrites in the supragranular
regions of the cerebral cortex as a key site for mediating conscious awareness. For instance,
optogenetic blockage of the connections between the cell bodies and dendrites of thick-
tufted layer V pyramidal cells in the sensory cortex causally modulated conscious arousal in
mice[52]. Other work has shown that both the noradrenergic[53] and cholinergic[54] systems alter
this mechanism, albeit in distinct ways: noradrenaline would promote burst firing due to the
α2a receptor-mediated closure of Ih HCN leak-channels[53], whereas the cholinergic system
instead prolongs the time-scale of firing via M1 cholinergic receptor activation on pyramidal
cell dendrites[54]. In this way, coordinated activity in the ascending arousal system can mediate
alterations in microcircuit processing that ultimately manifest as alteration in macroscopic
brain network dynamics.

The vascular nature of the T2* fMRI signal is such that it is impossible to rule out the role of
haemodynamics in the results we obtained in our analysis. Indeed, there is evidence that
noradrenaline causes a targeted hyperaemia through the augmentation of G-protein-coupled
receptors on vascular smooth muscle cells[55,56]. However, it is also clear that the
haemodynamics and massed neural action in the cerebral cortex are inextricably linked[57,58]. In
addition, there is evidence that stimulation of the locus coeruleus leads to the high-
frequency, low-amplitude electrophysiological activity patterns characteristic of the awake

state[10]. Together, these results argue that the locus coeruleus mediates a combination of
haemodynamic and neural responses that facilitate integrative neural network interactions
and subsequently mediate alterations in conscious awareness.

In this manuscript, we have argued that the ascending arousal system provides crucial
constraints over normal brain function, however there are numerous examples wherein
pathology within the ascending arousal system leads to systemic impairments in cognition. In
addition to disorders of consciousness[59], dementia syndromes are also crucially related to
dysfunction within the ascending arousal system. For instance, Alzheimer's disease has been
linked to tau pathology within the BNM[26], however individuals with Alzheimer's disease also
often have pathological involvement of the LC as well[60]. Similarly, individuals with
Parkinson's disease often have extra-dopaminergic pathology in the LC[61], as well as in the
cholinergic tegmentum[62]. Given the pathological processes at play in these disorders, we
expect that other neuromodulatory systems will also be impaired, and in turn effect the
macroscopic dynamics of the system in ways that remain to be elucidated.

In conclusion, we leveraged a high-resolution 7T resting state fMRI dataset to test the
hypothesis that activity within the ascending arousal system shapes and constrains patterns
of systems-level network reconfiguration. Our results support specific predictions from a
recent hypothetical framework[15], and further delineate the manner in which the autonomic
nervous system shapes and constrains ongoing, low-dimensional brain state dynamics in the
central nervous system in a manner that supports changes in conscious awareness.

Methods

7T resting state fMRI

Sixty-five healthy, right-handed adult participants (mean, 23.35 years; SD, 3.6 years; range

18–33 years; 28 females) were recruited, of whom 59 were included in the final analysis (four

participants were excluded due to MR scanning issues, one participant was excluded due to

an unforeseen brain structure abnormality, and one was excluded due to inconsistent BOLD

dynamics following global-signal regression). Participants provided informed written consent

to participate in the study. The research was approved by The University of Queensland

Human Research Ethics Committee. These data were originally described in Hearne et al.,

2017[63]. 1050 (~10 minutes) whole-brain 7T resting state fMRI echo planar images were

acquired using a multiband sequence (acceleration factor = 5; 2 mm^3 voxels; 586 ms TR;

23 ms TE; 40^0 flip angle; 208 mm FOV; 55 slices). Structural images were also collected to

assist functional data pre-processing (MP2RAGE sequence – 0.75 mm^3 voxels 4,300 ms TR;

3.44 ms TE; 256 slices).

DICOM images were first converted to NIfTI format and realigned. T1 images were

reoriented, skull-stripped (FSL BET), and co-registered to the NIfTI functional images using

statistical parametric mapping functions. Segmentation and the DARTEL algorithm were

used to improve the estimation of non-neural signal in subject space and the spatial

normalization. From each grey-matter voxel, the following signals were regressed: linear

trends, signals from the six head-motion parameters (three translation, three rotation) and

their temporal derivatives, white matter, and CSF (estimated from single-subject masks of

white matter and CSF). The aCompCor method (Behzadi et al., 2007) was used to regress

out residual signal unrelated to neural activity (i.e., five principal components derived from

noise regions- of-interest in which the time series data were unlikely to be modulated by

neural activity). Participants with head displacement > 3 mm in > 5% of volumes in any one

scan were excluded (n = 5). A temporal band pass filter (0.01 < f < 0.15 Hz) was applied to

the data.

Brain parcellation

Following pre-processing, the mean time series was extracted from 400 pre-defined cortical

parcels using the Schaefer atlas (Schaefer et al., 2018). Probabilistic anatomical atlases were

used to define the location of the noradrenergic LC[64] and the cholinergic BNM (Ch4 cell
group)[65]. The mean signal intensity from each region was extracted and then used for
subsequent analyses. To ensure that the BOLD data were reflective of neuronal signals, we
statistically compared LC and BNM time series with a number of potential nuisance signals
from: i) the cerebrospinal fluid; ii) the cortical white matter; iii) mean framewise
displacement; and iv) a 2mm^3 sphere in the fourth ventricle (centred at MNI co-ordinates: 0
-45 -30)[66]. All signals were unrelated to LC and BNM activity ($|r| < 0.05$ in each case),
however given the spatial proximity of the LC to the fourth ventricle, we opted to use a
linear regression to residualize the signal from the fourth ventricle. To ensure that BOLD
signals from nearby grey matter structures were not influencing the locus coeruleus
timeseries, we extracted the mean activity of the locus coeruleus mask after shifting the mask
anteriorly such that it overlapped with an area of the pons that harbours the nuclei (i.e.,
+8mm in the Y direction). In the same manner in which we previously regressed the
dynamics of the fourth ventricle, we regressed the activity of this non-LC pontine region,
and then re-analysed our data. Each of the results was statistically identical following this
approach, providing confidence that the original conclusions were not biased by a lack of
regional specificity.
462
Phasic increases in neuromodulatory BOLD signal
To identify phasic increases in neuromodulatory BOLD signal, we calculated the second
derivative (i.e., the acceleration) of the LC and BNM time series, and then identified points
in time that fulfilled three criteria: 1) value greater than or equal to 2 s.d. above the mean
acceleration; 2) value of the original time series, i.e., LC or BNM, was greater than or equal
to 2 s.d. above the mean of the time series within the following 10 TRs (i.e., 5.8 seconds);
and 3) the time point was not present within the first or last 20 TRs of an individual subjects'
trial (so as to avoid potential boundary effects). Using these criteria, we identified 148 $\tau_{\text{LC-BNM}}$
time points, 130 $\tau_{\text{BNM-LC}}$ time points and 316 $\tau_{\text{LC+BNM}}$ time points across all 59 subjects. To
ensure that the choice of 2 s.d. threshold was reflective of the underlying dynamics, we
altered this threshold between 1-3 s.d. and found robustly similar patterns. For subsequent
analyses, we identified time points in the 21 TR window surrounding these peaks, and then
used these to conduct statistical comparisons of the low-dimensional, complex network
signature of brain network dynamics as a function of phasic ascending arousal system

activity. Each of these patterns was confirmed using a lag-based cross-correlation analysis,

which demonstrated similar phenomena to those that we present in the manuscript.

To monitor the propagation of cortical signals with respect to $\tau_{\text{LC-BNM}}$, $\tau_{\text{BNM-LC}}$ and $\tau_{\text{LC+BNM}}$, we

extracted the time-to-peak of the cross-correlation between these signals and each of the 400

cortical parcels within the 10 TR (i.e., 5.8 second) windows following each identified phasic

peak. These patterns were mapped onto the cortex (Fig. 1B) for visualization and clearly

demonstrated anterior-to-posterior direction for the wave. We then used the volumetric

MNI co-ordinates of the Schaefer parcellation scheme to calculate the average velocity of the

travelling wave (0.13m s^{-1}).

In order to obtain an appropriate null model against which to compare our data, we

identified 5,000 random timepoints within the concatenated dataset that did not substantially

overlap with the already identified $\tau_{\text{LC-BNM}}$, $\tau_{\text{BNM-LC}}$ and $\tau_{\text{LC+BNM}}$ time series, and used these to

populate a null distribution[67]. Outcome measures were deemed significant if they were more

extreme than the 95$^{\text{th}}$ (or 5$^{\text{th}}$) percentile of the null distribution. Crucially, this ensured that

our data could not be explained by the characteristic spatial and temporal autocorrelation

present in BOLD timeseries data.

Time-resolved functional connectivity.

To estimate functional connectivity between the 400 regions of interest, we used the

multiplication of temporal derivatives (MTD) technique. Briefly, MTD is computed by

calculating the point-wise product of temporal derivative of pair-wise time series. The

resultant score is then averaged over a temporal window, w (a window length of 20 TRs was

used in this study, though results were consistent for w = 10–50 TRs).

Modularity Maximization

The Louvain modularity algorithm from the Brain Connectivity Toolbox (BCT[73]) was used

on the neural network edge weights to estimate community structure. The Louvain

algorithm iteratively maximizes the modularity statistic, Q, for different community

assignments until the maximum possible score of Q has been obtained:

$$Q_T = \frac{1}{v^+}\sum_{ij}\left(w_{ij}^+ - e_{ij}^+\right)\delta_{M_iM_j} - \frac{1}{v^+ + v^-}\sum_{ij}\left(w_{ij}^- - e_{ij}^-\right)\delta_{M_iM_j},$$

where v is the total weight of the network (sum of all negative and positive connections), w_{ij}

is the weighted and signed connection between regions i and j, e_{ij} is the strength of a

connection divided by the total weight of the network, and δ_{MiMj} is set to 1 when regions are

in the same community and 0 otherwise. '+' and '−' super-scripts denote all positive and

negative connections, respectively. The modularity of a given network is therefore a

quantification of the extent to which the network may be subdivided into communities with

stronger within-module than between-module connections.

For each epoch, we assessed the community assignment for each region 500 times and a

consensus partition was identified using a fine-tuning algorithm from the Brain Connectivity

Toolbox (BCT; http://www.brain-connectivity-toolbox.net/). We calculated all graph

theoretical measures on un-thresholded, weighted and signed connectivity matrices[73]. The

stability of the γ parameter was estimated by iteratively calculating the modularity across a

range of γ values (0.5-2.5; mean Pearson's r = 0.859 +-0.01) on the time-averaged

connectivity matrix for each subject – across iterations and subjects, a γ value of 1.0 was

found to be the least variable, and hence was used for the resultant topological analyses.

Participation Coefficient

The participation coefficient, PC, quantifies the extent to which a region connects across all

modules (i.e., between-module strength) and has previously been used to successfully

characterize hubs within brain networks (e.g. see [75]). The PC for each region was calculated

within each temporal window as,

$$PC = 1 - \sum_{s=1}^{n_M}\left(\frac{\kappa_{isT}}{\kappa_{iT}}\right)^2$$

530   where k_{isT} is the strength of the positive connections of region i to regions in module s at

531   time T, and k_{iT} is the sum of strengths of all positive connections of region i at time T.

532   Negative connections were discarded prior to calculation. The participation coefficient of a

533   region is therefore close to 1 if its connections are uniformly distributed among all the

534   modules and 0 if all of its links are within its own module.

Brain State Displacement and the Energy Landscape

To quantify the change in BOLD activity following phasic bursts of neuromodulation we calculated the BOLD mean-squared displacement (MSD). The MSD is a measure of the deviation in BOLD activity, $\boldsymbol{x}_t = [x_{1,t}, x_{2,t}, \dots, x_{r,t}]$ for r parcels, with respect to the activity at the phasic onset, t_0. The MSD is calculated as the average change of each voxel

$$MSD_{t,t_0} = \langle |\boldsymbol{x}_{t_0+t} - \boldsymbol{x}_{t_0}|^2 \rangle_r,$$

and it is calculated for different t_0, where t_0 are the onset of a subcortical phasic burst, across t TRs. We are interested in the probability, P_{MSD}, that we will observe a given displacement in BOLD at a given time-lag t. We estimated the probability distribution function $P(MSD, t)$ from n MSD_{t,t_0} samplings, — e.g., the identified n phasic bursts of subcortical structures (as above) – using a Gaussian kernel density estimation $P(MSD, t) = \frac{1}{4n} \sum_{i=1}^{n} K\left(\frac{MSD_{t,i}}{4}\right)$, where $K(u) = \frac{1}{2\sqrt{\pi}} e^{-\frac{1}{2}u^2}$ and we display the results for t between 1 to 15 TR and MSD between 0 to 50. As is typical in statistical mechanics the energy of a given state, E_σ, and its probability are related $P(\sigma) = \frac{1}{Z} e^{-\frac{E_\sigma}{T}}$, where Z is the normalisation function and T is a scaling factor equivalent to temperature in thermodynamics [20]. In our analysis $\sum_\sigma P_\sigma = 1$ by construction and we can set $T = 1$ for the observed data. Thus, the energy of each BOLD MSD for a given at a given time-lag t, E, is then equal to the natural logarithm of the inverse probability, $P(MSD, t)$, of its occurrence:

$$\mathrm{E} = \ln \frac{1}{P(MSD, t)}.$$

Meditation Dataset

Fourteen healthy right-handed non-smoking meditation practitioners (11 female; age 28-66) underwent Siemens 3T MRI scanning (T1: TR = 2600 msec, TE = 3.9 msec, TI = 900 msec, FOV = 24 cm, 256 x 256 matrix, voxel dimensions = 1 x 1 x 1 mm³; T2*: weighted gradient-echo pulse sequence, TR = 1500 msec, TE = 30 msec, flip angle = 90 deg, FOV = 192 cm, 64 x 64 matrix, voxel dimensions = 3 x 3 x 4 mm³). All participants signed a consent form approved by the Institutional Review Board at Emory University and the Atlanta Veterans Affairs

Research and Development Committee as an indication of informed consent. Participants

were asked to meditate for 20 min in the MRI scanner by maintaining focused attention on

the breath and keeping the eyes closed. They were instructed to press a button whenever

they realized their mind had wandered away from the breath, and then return their focus to

the breath. The epoch of time immediately prior to the button press was thus the moment in

time in which each individual recognized that their focus had deviated from their breath.

This information was used to construct a finite impulses response model that mapped the 5

TRs prior-to and following each button press. We then modelled LC>BNM activity, low-

dimensional dynamics and network topology around this epoch to construct a model of

state-space reconfiguration as a function of intrinsic conscious awareness. Non-parametric,

block-resampling null distributions were utilized for statistical testing ($p < 0.05$).

Data availability

The BOLD data was obtained from (Hearne et al., 2017)[63] and

The BOLD data that support the findings of this study were obtained from (Hearne et al.,

2017)[63] and they are available from the authors upon reasonable request. The subcortical

timeseries (τ_{LC} and τ_{BNM}) that support the findings of this study are available at

(github.com/Bmunn/BSI).

Code availability

All the code required to conduct the analysis can be found on Github at

(github.com/Bmunn/BSI).

Supplementary Figures

Figure S1. The locus coeruleus. Left: The anatomical locus coeruleus mask projected onto MNI 0.5mm

standard brain (orange); Right: The anatomical locus coeruleus mask down-sampled onto an example 7T Echo

Planar Image from a single subject (red).

Figure S2. Time-varying correlations. Average correlation preceding (left) and following (middle) the zero-

lagged　　　　value, along with the difference between the two (right); squares represent eight pre-defined

sub-networks: Vis – visual, SM – somatomotor, DAN – dorsal attention, VAN – ventral attention, LIM –

limbic, CON – control, DMN – default and TP – temporal pole.

Figure S3 – LC and BNM move dynamics to differing regimes than unaroused activity and their simultaneous combination LC+BNM. The energy landscape of simultaneous LC+BNM phasic bursts relative to their linear superposition, suggesting the simultaneous combination may allow the system to reach particularly unique brain-states that neither individually could reach.

References

1. Sporns, O. Network attributes for segregation and integration in the human brain. *Current Opinion in Neurobiology* **23**, 162–171 (2013).

2. Shine, J. M. The thalamus integrates the macrosystems of the brain to facilitate complex, adaptive brain network dynamics. *Progress in Neurobiology* 101951 (2020) doi:10.1016/j.pneurobio.2020.101951.

3. Bullmore, E. & Sporns, O. The economy of brain network organization. *Nature Reviews Neuroscience* **13**, 336–349 (2012).

4. Shine, James M *et al.* Computational models link cellular mechanisms of neuromodulation to large-scale brain dynamics. *Nature Neuroscience* **Ahead of Print**,.

5. Brown, E. N., Purdon, P. L. & Van Dort, C. J. General Anesthesia and Altered States of Arousal: A Systems Neuroscience Analysis. *Annu. Rev. Neurosci.* **34**, 601–628 (2011).

6. Salinas, E. & Sejnowski, T. J. Book Review: Gain Modulation in the Central Nervous System: Where Behavior, Neurophysiology, and Computation Meet. *Neuroscientist* **7**, 430–440 (2001).

7. Sara, S. J. The locus coeruleus and noradrenergic modulation of cognition. *Nature Reviews Neuroscience* **10**, 211–223 (2009).

8. Hasselmo, M. E. & Sarter, M. Modes and models of forebrain cholinergic neuromodulation of cognition. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology* **36**, 52–73 (2011).

9. Samuels, E. R. & Szabadi, E. Functional neuroanatomy of the noradrenergic locus coeruleus: its roles in the regulation of arousal and autonomic function part II: physiological and pharmacological manipulations and pathological alterations of locus coeruleus activity in humans. *Current neuropharmacology* **6**, 254–285 (2008).

10. Carter, M. E. *et al.* Tuning arousal with optogenetic modulation of locus coeruleus

neurons. *Nature Neuroscience* **13**, 1526–1533 (2010).

11. Lin, S.-C., Brown, R. E., Hussain Shuler, M. G., Petersen, C. C. H. & Kepecs, A.

Optogenetic Dissection of the Basal Forebrain Neuromodulatory Control of Cortical

Activation, Plasticity, and Cognition. *The Journal of neuroscience : the official journal of the*

Society for Neuroscience **35**, 13896–13903 (2015).

12. Zaborszky, L. *et al.* Neurons in the basal forebrain project to the cortex in a complex

topographic organization that reflects corticocortical connectivity patterns: an

experimental study based on retrograde tracing and 3D reconstruction. *Cerebral cortex*

(New York, N.Y. : 1991) **25**, 118–137 (2015).

13. Servan-Schreiber, D., Printz, H. & Cohen, J. A network model of catecholamine effects-

Gain, signal-to-noise ratio, and behavior. *Science* **249**, 892–895 (1990).

14. Schmitz, T. W. & Duncan, J. Normalization and the Cholinergic Microcircuit: A Unified

Basis for Attention. *Trends in Cognitive Sciences* **22**, 422–437 (2018).

15. Shine, J. M. Neuromodulatory Influences on Integration and Segregation in the Brain.

Trends in Cognitive Sciences **23**, 572–583 (2019).

16. Cunningham, J. P. & Yu, B. M. Dimensionality reduction for large-scale neural

recordings. *Nature Neuroscience* **17**, 1500–1509 (2014).

17. Shine, J. M. *et al.* Human cognition involves the dynamic integration of neural activity

and neuromodulatory systems. *Nat Neurosci* **22**, 289–296 (2019).

18. Gallego, J. A., Perich, M. G., Chowdhury, R. H., Solla, S. A. & Miller, L. E. Long-term

stability of cortical population dynamics underlying consistent behavior. *Nat Neurosci*

(2020) doi:10.1038/s41593-019-0555-4.

19. Miller, P. Dynamical systems, attractors, and neural circuits. *F1000Res* **5**, 992 (2016).

20. Tkačik, G. *et al.* Thermodynamics and signatures of criticality in a network of neurons.

Proc Natl Acad Sci USA **112**, 11508–11513 (2015).

21. Bialek, W. Perspectives on theory at the interface of physics and biology. *Rep. Prog. Phys.*

81, 012601 (2018).

22. Watanabe, T. *et al.* A pairwise maximum entropy model accurately describes resting-state

human brain networks. *Nat Commun* **4**, 1370 (2013).

23. Ezaki, T., Watanabe, T., Ohzeki, M. & Masuda, N. Energy landscape analysis of

neuroimaging data. *Phil. Trans. R. Soc. A.* **375**, 20160287 (2017).

24. Krzemiński, D. *et al.* Energy landscape of resting magnetoencephalography reveals

fronto-parietal network impairments in epilepsy. *Network Neuroscience* **4**, 374–396 (2020).

25. Breakspear, M. Dynamic models of large-scale brain activity. *Nature Neuroscience* **20**, 340–

352 (2017).

26. Mesulam, M. M. Cholinergic circuitry of the human nucleus basalis and its fate in

Alzheimer's disease. *Journal of Comparative Neurology* **521**, 4124–4144 (2013).

27. Aston-Jones, G. & Cohen, J. D. An Integrative Theory of Locus Coeruleus-

Norepinephrine Function: Adaptive Gain and Optimal Performance. *Annual Review of*

Neuroscience **28**, 403–450 (2005).

28. Muller, L., Chavane, F., Reynolds, J. & Sejnowski, T. J. Cortical travelling waves:

mechanisms and computational principles. *Nat Rev Neurosci* **19**, 255–268 (2018).

29. Shine, J. M. *et al.* The Dynamics of Functional Brain Networks: Integrated Network

States during Cognitive Task Performance. *Neuron* **92**, 544–554 (2016).

30. Shine, J. M., Aburn, M. J., Breakspear, M. & Poldrack, R. A. The modulation of neural

gain facilitates a transition between functional segregation and integration in the brain.

Elife **7**, e31130 (2018).

31. Thomas Yeo, B. T. *et al.* The organization of the human cerebral cortex estimated by

intrinsic functional connectivity. *Journal of Neurophysiology* **106**, 1125–1165 (2011).

32. Mather, M. & Sutherland, M. R. Arousal-Biased Competition in Perception and

Memory. *Perspect Psychol Sci* **6**, 114–133 (2011).

33. Mather, M., Clewett, D., Sakaki, M. & Harley, C. W. Norepinephrine ignites local

hotspots of neuronal excitation: How arousal amplifies selectivity in perception and

memory. *Behav Brain Sci* **39**, e200 (2016).

34. Robinson, R. Differential behavioral and biochemical effects of right and left

hemispheric cerebral infarction in the rat. *Science* **205**, 707–710 (1979).

35. Tona, K.-D. *et al.* In vivo visualization of the locus coeruleus in humans: quantifying the

test–retest reliability. *Brain Struct Funct* **222**, 4203–4217 (2017).

36. Kanamaru, T., Fujii, H. & Aihara, K. Deformation of attractor landscape via cholinergic

presynaptic modulations: a computational study using a phase neuron model. *PloS one* **8**,

e53854 (2013).

37. Munn, B. & Gong, P. Critical Dynamics of Natural Time-Varying Images. *Phys. Rev. Lett.*

121, 058101 (2018).

38. Kucyi, A., Hove, M. J., Esterman, M., Hutchison, R. M. & Valera, E. M. Dynamic Brain

Network Correlates of Spontaneous Fluctuations in Attention. *Cerebral cortex (New York,*

N.Y.□: 1991) **27**, 1831–1840 (2016).

39. Hasenkamp, W., Wilson-Mendenhall, C. D., Duncan, E. & Barsalou, L. W. Mind

wandering and attention during focused meditation: a fine-grained temporal analysis of

fluctuating cognitive states. *NeuroImage* **59**, 750–760 (2012).

40. Sara, S. J. & Bouret, S. Orienting and reorienting: the locus coeruleus mediates cognition

through arousal. *Neuron* **76**, 130–141 (2012).

41. Jahn, C. I., Varazzani, C., Sallet, J., Walton, M. E. & Bouret, S. Noradrenergic But Not

Dopaminergic Neurons Signal Task State Changes and Predict Reengagement After a

Failure. *Cerebral Cortex* bhaa089 (2020) doi:10.1093/cercor/bhaa089.

42. Sales, A. C., Friston, K. J., Jones, M. W., Pickering, A. E. & Moran, R. J. Locus

Coeruleus tracking of prediction errors optimises cognitive flexibility: an Active

Inference model. *PLoS computational biology* **15**, e1006267 (2019).

43. Gotts, S. J., Gilmore, A. W. & Martin, A. Brain networks, dimensionality, and global

signal averaging in resting-state fMRI: Hierarchical network structure results in low-

dimensional spatiotemporal dynamics. *NeuroImage* **205**, 116289 (2020).

44. Shine, J. M. *et al.* The Low-Dimensional Neural Architecture of Cognitive Complexity Is

Related to Activity in Medial Thalamic Nuclei. *Neuron* **104**, 849-855.e3 (2019).

45. Mante, V., Sussillo, D., Shenoy, K. V. & Newsome, W. T. Context-dependent

computation by recurrent dynamics in prefrontal cortex. *Nature* **503**, 78–84 (2013).

46. Müller, E. J., Munn, B. R. & Shine, J. M. Diffuse neural coupling mediates complex

network dynamics through the formation of quasi-critical brain states. *Nat Commun* **11**,

6337 (2020).

47. de Gee, J. W. *et al.* Pupil-linked phasic arousal predicts a reduction of choice bias across

species and decision domains. *eLife* **9**, e54014 (2020).

48. Pfeffer, T. *et al.* Catecholamines alter the intrinsic variability of cortical population

activity and perception. *PLoS Biology* **16**, e2003453 (2018).

49. Kosciessa, J. Q., Lindenberger, U. & Garrett, D. D. *Thalamocortical excitability adjustments*

guide human perception under uncertainty.

http://biorxiv.org/lookup/doi/10.1101/2020.06.22.165118 (2020)

doi:10.1101/2020.06.22.165118.

50. Brezina, V. Beyond the wiring diagram: signalling through complex neuromodulator

networks. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* **365**,

2363–2374 (2010).

51. Aru, J., Suzuki, M. & Larkum, M. E. Cellular Mechanisms of Conscious Processing.

Trends in Cognitive Sciences **24**, 814–825 (2020).

52. Suzuki, M. & Larkum, M. E. General Anesthesia Decouples Cortical Pyramidal

Neurons. *Cell* **180**, 666-676.e13 (2020).

53. Labarrera, C. *et al.* Adrenergic Modulation Regulates the Dendritic Excitability of Layer 5

Pyramidal Neurons In Vivo. *Cell reports* **23**, 1034–1044 (2018).

54. Williams, S. R. & Fletcher, L. N. A Dendritic Substrate for the Cholinergic Control of

Neocortical Output Neurons. *Neuron* **101**, 486-499.e4 (2019).

55. Bekar, L. K., Wei, H. S. & Nedergaard, M. The Locus Coeruleus-Norepinephrine

Network Optimizes Coupling of Cerebral Blood Volume with Oxygen Demand. *J Cereb*

Blood Flow Metab **32**, 2135–2145 (2012).

56. Lecrux, C. & Hamel, E. Neuronal networks and mediators of cortical neurovascular

coupling responses in normal and altered brain states. *Phil. Trans. R. Soc. B* **371**, 20150350

(2016).

57. Moore, C. I. & Cao, R. The Hemo-Neural Hypothesis: On The Role of Blood Flow in

Information Processing. *Journal of Neurophysiology* **99**, 2035–2047 (2008).

58. Chow, B. W. *et al.* Caveolae in CNS arterioles mediate neurovascular coupling. *Nature*

579, 106–110 (2020).

59. Snider, S. B. *et al.* Disruption of the ascending arousal network in acute traumatic

disorders of consciousness. *Neurology* **93**, e1281–e1287 (2019).

60. Robertson, I. H. A noradrenergic theory of cognitive reserve: implications for

Alzheimer's disease. *Neurobiology of Aging* **34**, 298–308 (2013).

61. Rommelfanger, K. S. & Weinshenker, D. Norepinephrine: The redheaded stepchild of

Parkinson's disease. *Biochemical pharmacology* **74**, 177–190 (2007).

62. Bohnen, N. I. & Albin, R. L. The cholinergic system and Parkinson disease. *Behavioural*

Brain Research **221**, 564–573 (2011).

63. Hearne, L. J., Cocchi, L., Zalesky, A. & Mattingley, J. B. Reconfiguration of Brain

Network Architectures between Resting-State and Complexity-Dependent Cognitive

Reasoning. *J. Neurosci.* **37**, 8399–8411 (2017).

64. Keren, N. I., Lozar, C. T., Harris, K. C., Morgan, P. S. & Eckert, M. A. In vivo mapping

of the human locus coeruleus. *NeuroImage* **47**, 1261–1267 (2009).

65. Eickhoff, S. B. *et al.* A new SPM toolbox for combining probabilistic cytoarchitectonic

maps and functional imaging data. *NeuroImage* **25**, 1325–1335 (2005).

66. Li, R. *et al.* A pulse of transient fMRI increases in subcortical arousal systems during

transitions in attention. 63.

67. Nichols, T. E. & Holmes, A. P. Nonparametric permutation tests for functional

neuroimaging: a primer with examples. *Human brain mapping* **15**, 1–25 (2002).

Author contributions

764 MS conceived, funded, and directed the project. MS Curated the data. BM, EM, MS
765 Conducted the analysis. BM, MS Wrote the original draft. BM, EM, GW, MS Reviewed and
766 edited the manuscript.