Are narrow-ranging species doomed to extinction? Projected dramatic decline in future climate

suitability of two highly threatened species

Running title: Narrow-ranging species and climate change

Nicolas Dubos[1, 2], Frederique Montfort[3], Clovis Grinand[3], Marie Nourtier[3], Gregory Deso[4], Jean-Michel

Probst[5], Julie Hanta Razafimanahaka[6], Raphali Rodlis Andriantsimanarilafy[6], Eddie Fanantenana

Rakotondrasoa[6], Pierre Razafindraibe[6], Richard Jenkins[6,7] and Angelica Crottini[8]

9
10

1 Centre d'Ecologie et des Sciences de la Conservation (CESCO UMR 7204), Sorbonne Université, MNHN,
55 rue Buffon, 75005 Paris, France.

2 INRAE (UMR TETIS), Maison de la télédétection, 500 rue Jean-François Breton 34093 Montpellier
Cedex 5, France.

3 N'Lab, Nitidæ, Maison de la télédétection, 500 rue Jean-François Breton 34093 Montpellier Cedex 5

4 Association Herpétologique de Provence Alpes Méditerranée, Maison des Associations, 384 route de
Caderousse, F-84100 Orange, France.

5 Association Nature and Patrimoine, 1 rue des amarantes résidence Valeriane 2, Bat C appartement 15.
97490 Sainte Clotilde, Île de La Réunion, France.

6 Madagasikara Voakajy, B.P. 5181, Antananarivo (101), Madagascar.

7 Durrell Institute of Conservation and Ecology, School of Anthropology and Conservation, Marlowe
Building, University of Kent, Canterbury, CT2 7NR ; Current address: IUCN Global Species Programme,
219c Huntingdon Road, Cambridge, CB3 0DL, UK.

8 CIBIO, Research Centre in Biodiversity and Genetic Resources, InBIO, Universidade do Porto, Campus

Agrário de Vairão, Rua Padre Armando Quintas, No 7, 4485-661 Vairão, Portugal.

Abstract

Narrow-ranging species are usually omitted from Species distribution models (SDMs) due to statistical constraints, while they are predicted to be particularly vulnerable to climate change. The recently available high-resolution environmental predictors, along with recently developed methods enable to increase the eligibility of narrow-ranging species for SDMs, provided their distribution is well known. We fill a gap of knowledge on the effect of predicted climate change on narrow-ranging species. We modelled the distribution of the golden mantella frog *Mantella aurantiaca* and the Manapany day gecko *Phelsuma inexpectata,* for which the distribution of their occurrence records is well documented. Our modelling scheme included a range of processes susceptible to address statistical issues related to narrow-ranging species. We predict an alarming decline in climate suitability in the whole current distribution area of both species by 2070, potentially leading to a complete extinction in most scenarios. We identified the areas with the best climate suitability in the future, but these remain largely suboptimal regarding species climatic niche. The high level of habitat fragmentation suggests that both species likely need to be at least partly translocated. Climate change may not only drive range contractions or distribution shifts in narrow-ranging species, but may lead to the complete extirpation of suitable environments across their entire region. This study suggests that the level of threats of narrow-ranging species already identified as threatened may be underestimated, especially in heterogeneous tropical areas. We stress the need to develop sampling campaigns and implement proactive actions for narrow-ranging species in the tropics.

Keywords: Climate change, Ensemble of Small Models, Jackknife, Madagascar, *Mantella aurantiaca*, *Phelsuma inexpectata*, Reunion Island, Species distribution models, Translocation

Introduction

Climate change is predicted to become the main driver of biodiversity loss it in the next decades (Bellard *et al.*, 2012). Species Distribution Models (SDMs) are probably the most common approach used to predict the impact of future climate change on species. They are used to predict current and future environmental suitability, and provide guidelines for the identification of priority areas for protection (Leroy *et al.*, 2014), habitat restoration and species (re)introduction/translocations (e.g., Bellis et al., 2020; Draper et al., 2019; Butt et al., 2020; Westwood et al., 2020). Habitat restoration and translocations are two ecological engineering techniques enabling the restoration of depleted populations. Translocation programs will become increasingly needed in the face of climate change, especially for species with small distribution ranges (Thomas, 2011). In this regard, SDMs help identifying suitable receptor sites that meet the species' habitat requirements while accounting for climate suitability (Bellis et al., 2020). In highly degraded environments a combination of habitat restoration and translocation may be needed to avoid species extinctions. This may be the case in highly fragmented tropical systems, where the number of narrow-ranging species is higher and where climate change effects are expected to be stronger (Tewksbury et al., 2008).

The impact of future climate change is largely understudied in endangered narrow-ranging species. This is mainly due to the difficulty to model their current distribution and project their future because of low sample sizes and subsequent little spatial replicates when fitted on climate data (Botts *et al.*, 2013; Platts et al., 2014; Breiner et al., 2015; Galante et al., 2018). Both these factors lead to statistical constraints that withdraw these species from eligibility for SDMs. However, the omission of narrow-ranging species in SDMs may be problematic in terms of conservation planning, because the area encompassing their distribution may be downplayed (Platts *et al.*, 2014). Although narrow-ranging species are known to be more vulnerable to climate change (Pearson *et al.*, 2014), few studies have provided quantitative assessment of climate change impacts on these (but see Alamgir *et al.*, 2015; Zhang *et al.*, 2020).

The recent availability of high-resolution climatic data (e.g., Fick & Hijmans, 2017; Karger et al., 2017), along with high-resolution land cover data (e.g., Vieilledent et al., 2018) is offering new opportunities for modelling the distribution of these species (Lannuzel et al., 2021). However, despite a probable increase in

statistical power, there is still the possibility for SDMs to produce misleading results due to spatial sampling

bias (Phillips *et al.*, 2009). The effect of sample bias may be particularly strong in rare (or poorly known)

species with small sample sizes, because models are more influenced by each occurrence data that is used

(Pearson *et al.*, 2007). A number of techniques were recently developed for data-poor species. For instance,

jackknife procedures (i.e. leave-one-out; Pearson et al. 2007) improves model assessments and Ensemble of

Small Models (ESMs) enable to deal with model complexity while keeping sufficient explanatory power

(Breiner et al. 2015). Other techniques are dedicated to sampling bias correction, often implying the filtering

of occurrence or environmental data (Gábor *et al.*, 2019) or non-random pseudo-absence selection (Phillips

et al., 2009). However, data filtering can become problematic for species with low sample sizes (Vollering et

al., 2019), especially when species distribution is highly localised (Inman et al., 2021) and is not

recommended in absence of evidence of bias in occurrence data (Gábor et al., 2019). Similarly, non-random

pseudo-absence selection is not always effective (Dubos et al., 2021) and tends to make predictions worse in

narrow-niche species (Inman et al., 2021). However, the restriction of the background area (where pseudo-

absences are generated), i.e. background thickening, can be advantageous for sample bias correction when

data is scarce without discarding presence records (Acevedo et al. 2012; Vollering et al. 2019). On the other

hand, the reliability of an SDM is more driven by the quality of the data than the implementation of models

(Araùjo et al., 2019). Therefore, the best option for narrow-ranging SDMs may be to (1) use all the available

methodological tools to deal with statistical issues, and (2) select species for which the distribution is well

known.

Given the predicted magnitude of climate change, along with the narrow thermal tolerance of tropical species

(Tewksbury et al., 2008), we may not only expect a reduction or a geographical shift in suitable conditions

for narrow-ranging species, but the extirpation of suitable conditions across the entirety of their distribution

range. Here we fill a gap of knowledge regarding the impact of future climate change on narrow-ranging

species using two species for which the distribution is particularly well documented. These were the

Manapany day gecko *Phelsuma inexpectata,* classified as Critically Endangered (IUCN & MNHN, 2010)

and the golden mantella frog *Mantella aurantiaca* formerly classified as Critically Endangered (Vences &

Raxworthy, 2008), now classified as Endangered after the inclusion of one locality record which increased its extent of occurrence (IUCN SSC Amphibian Specialist Group 2020). Both species are in continued decline (Probst & Turpin, 1997; Crottini et al., 2019), live in highly fragmented areas (respectively in Reunion Island and central Madagascar) and are in urgent need for conservation actions. Given the important, long-term efforts invested to document their distribution, we assume that the geographic information for these species is nearly comprehensive and unbiased. We tested whether climate change will 'only' drive range reductions/shifts, or lead to a total extirpation of their suitable areas. We eventually identify the most suitable candidate areas for habitat restoration and translocation across their respective regions.

Methods

Occurrence data

Phelsuma inexpectata – The Manapany day gecko is endemic to the south of Reunion Island. We retrieved 31 occurrence data from literature (Bour *et al.*, 1995). Since then, the surroundings of the known distribution range of the species were regularly visited (an 11 km-long coastal band; Fig. 1). Two localities corresponding to introduced populations were identified west of the current range, which we added to the data (Deso, 2001; Porcel *et al.*, 2021). Recent sampling campaigns enabled to find additional occupied habitats (Dubos 2010) but did not add any occurrence point after aggregation at the resolution of the environmental variables (30 arc seconds). Therefore, we assume that the sample occurrence of the species is nearly comprehensive. The total number of 30 sec. occupied pixels resulted in 15 presence points.

Mantella aurantiaca – The golden mantella frog is distributed in central-Eastern Madagascar (region of Moramanga; Fig. 1). We obtained 131 occurrence data from Piludu *et al.* (2015). Those included compiled published data from surveys conducted between 2001 and 2007 (Bora *et al.*, 2008; Randrianavelona *et al.*, 2010) and new locations from additional surveys conducted between 2008 and 2013. More recent surveys conducted between 2014 and 2019 enabled to add 39 occurrences. The region has been extensively surveyed

and it is very likely that most occupied habitats were identified. After aggregation to match the resolution of

environmental data, sample size resulted in 101 occurrence points.

Fig. 1 Study area and species, with occurrence points.

Climate data

We used the 19 bioclimatic variables for 30 arc sec (approximately 900m) resolution of the current climate

data and of the 2070 projections from CHELSA (Karger et al., 2017; Fig. S1, S2). We decided to include all

the 19 variables because both temperature and precipitation are related to the species' biology, including

those related to indices of variability (e.g., cyclones drive mortality in *Phelsuma* and heavy rains drive

reproduction in *Mantella*; Vinson, 1975; Randrianavelona et al., 2010). We used three Global Circulation

Models (GCMs; i.e., BCC-CSM1-1, MIROC5 and HadGEM2-AO) and two greenhouse gas emission

scenarios (the most optimistic RCP26 and the most pessimistic RCP85; Fig. S3, S4). We also ran sets of

models using the WorldClim Global Climate Data (Fick & Hijmans, 2017), with the same GCMs and

scenarios to account for potential effect of the baseline data in our predictions.

Distribution modelling

We modelled and projected species distributions with the biomod2 R package (Thuiller et al. 2009), using 10

modelling techniques: generalised linear model (GLM), generalised additive model (GAM), classification

tree analysis (CTA), artificial neural network (ANN), surface range envelop (SRE, as known as BIOCLIM),

flexible discriminant analysis (FDA) and random forest (RF), Multiple Adaptive Regression Splines

(MARS), Generalised Boosting Model (GBM) and Maximum Entropy (MaxEnt). For machine learning

modelling techniques, hyperparameters were set to their default values implemented in biomod2 (see

https://www.rdocumentation.org/packages/biomod2/versions/3.5.1/topics/BIOMOD_ModelingOptions). We

generated five different sets of randomly-selected pseudo-absences (random generation is recommended for

rare and specialised species; Inman et al., 2021). We selected one variable per group of inter-correlated

variables to avoid collinearity (Pearson's r > 0.7) and assessed the relative importance of each variable kept

with 10 permutations per model replicate (total = 500). The variables included in the final models were those

with a relative importance > 0.2 across at least 50% of model runs (these varied with the baseline climate and

the background extent; Table 1). We show the fitted predicted values for each individual model, and a mean

smoothed response curve. We predicted species distributions with an ensemble of small models approach

(ESM; Breiner et al., 2015). We ran sets of bivariate models, i.e. including all pairwise combinations of the

selected variables, and produced an ensemble model with the mean predictions across all models weighted

by their respective AUC (see below). This method is advocated for rare species and enables to reduce model

complexity without reducing the explanatory power. We set three runs of cross validation (except for the

jackknife procedure; see below). We ran a first set of models (1), hereafter referred to as 'Base' setting, with

1000 pseudo-absences selected from a background covering the entire respective island of each species (i.e.

all island cells). We ran a second set (2) where the 1000 pseudo-absences were down-weighted to equal

presence data (setting prevalence to 0.5) referred to as 'equal total weight' (ETW; Liu et al. 2019). We

reperformed the first two model sets but with a background covering the southern part of Reunion Island and

the eastern forested part of Madagascar (see 'Land use data' section), without, and with pseudo-absence

down-weighting, respectively referred to as (3) 'Restricted background' and (4) 'ETW-Restricted

background'. Eventually, we ran a (5) Base model set with the Worldclim baseline climate instead of Chelsa,

referred to as 'Worldclim baseline'. For both species we present five ensemble models of current distribution

(of varying background, prevalence and baseline climate), and 30 ensemble models of projected future

distribution (of varying background, prevalence, baseline climate, GCM and RCP). We did not account for

species dispersal ability, because the purpose was not to predict potential shifts, but to assess how suitable

will be the climate in the future in order to identify candidate sites for restoration and translocation.

Model evaluations – Ideally, performance evaluations are based on block-cross validation to limit spatial

autocorrelation at large scales. In our case, species distributions are highly localised and the use of spatial

splits would result in strong unbalances between blocks. We therefore used a random partitioning for *M.*

aurantiaca and a jackknife for *P. inexpectata*. The jackknife procedure is appropriate for small sample sizes

(Pearson *et al.*, 2007; Galante *et al.*, 2018). This approach consists in running n iterations corresponding to

the number of occurrence data, removing one occurrence for each run of calibration. Models are evaluated

with the withheld occurrence. We did not used it for *M. aurantiaca* given that the number of occurrence data

was sufficient for a standard procedure, and used an 80% calibration subset (random partitioning), with five

cross-validation runs. We assessed model performance using the Area Under the Operative Curve (AUC), the

True Skill Statistics (TSS) and the Boyce index (the latter was not computed for the jackknife procedure,

because this index is based on Spearman's coefficients and cannot be estimated on the basis of a single

location). Poorly performing models could result from calibration subsets falling into different conditions

than evaluation subsets. For ensemble models (mean predictions), we excluded models for which the AUC

was below 0.9 and for which the Boyce index was below 0.5 (Gillard et al., 2017).

We performed a multivariate environmental similarity surfaces analysis (MESS) to determine whether

models are well informed for projections on novel (future) data. We eventually quantified the uncertainty

related to model input parameters (modelling technique, pseudo-absence distribution, number of pseudo-

absences generated, pseudo-absence down-weighting, GCMs, RCPs and baseline climate data) by computing

the standard deviation of the suitability scores between model predictions.

Land use data

We accounted for the habitat requirements of our model species by applying a filter to the projected climate

suitability based on land use and land cover data (e.g., Gillard *et al.*, 2017). This enabled to minimise model

complexity while remaining biologically realistic and relevant for conservation applications.

Reunion Island – We used very high-resolution land cover categories (Urban, agricultural, natural, water) at

1.5m resolution (resampled at 15m for computing purposes) derived from remote sensing (Dupuy and

Gaetano, 2019; Fig. S5). Since *P. inexpectata* can be found in both urbanised and natural areas and avoids

agricultural areas (Probst & Turpin, 1997), we excluded the latter land use type only. This is relevant also for

conservation applications because habitat restoration is more difficult to implement in agricultural areas.

Madagascar – *Mantella aurantiaca* is exclusively found in swampy forested areas (Randrianavelona *et al.*,

2010). We filtered our suitability maps to include only cells corresponding to rainforests. We obtained forest

cover data for two periods (1990 and 2017) from a study that combined global tree cover loss data with

historical national forest (resolution 30m; Vieilledent et al., 2018; Fig. S6). We used forest cover of the year

1990 for current distribution models because it corresponds approximately to the period of the oldest

occurrence data (some areas were deforested since then). For future projections, we used the latest forest

cover data available, i.e. 2017.

Results

Variable selection and model performance

Three to four variables were selected in every cases (no change between prevalence settings; see details in

Table 1; Fig. S7, S8). The current distribution of both species was well predicted (across five model sets: *P.*

inexpectata: mean AUC = 0.96, mean TSS = 0.91; *M. aurantiaca*: mean Boyce = 0.82, mean AUC = 0.97,

mean TSS = 0.94; Fig. S9, S10). We excluded 5076 poorly performing models out of 18750 for *P.*

inexpectata, and 343 out of 4140 for *M. aurantiaca*. The median suitability score at the presence points for *P.*

inexpectata was 836 (911 when removing the two introduced populations) and 783 for *M. aurantiaca*.

Models consistently identified the current distribution range between all runs for both species, as shown by

the uncertainty map (SD intermodel suitability scores at the presence points of 168 for *P. inexpectata* and 113

for *M. aurantiaca*; Fig. S11, S12).

Table 1. Selected variables for two narrow-ranging species in Reunion Island and Madagascar. Variables

were obtained from CHELSA (Karger et al., 2017) and Worldclim (Fick & Hijmans, 2017), and selected on

the basis of 10 permutations per modelling technique and pseudo-absence set (total = 500).

Species	Model set	Selected variables
Phelsuma inexpectata	Base	Bio1, Bio15, Bio18
	Restricted background	Bio1, Bio15, Bio18
	Worldclim baseline	Bio1, Bio3, Bio14, Bio16
Mantella aurantiaca	Base	Bio1, Bio3, Bio4, Bio5
	Restricted background	Bio1, Bio18, Bio19
	Worldclim baseline	Bio10, Bio11, Bio14, Bio16

Future climate suitability

All scenarios indicated an important decrease in climate suitability in the entirety of the current range of both

species by 2070 (Fig. 2–7). On average across scenarios and GCMs, suitability scores decreased by 59% for

P. inexpectata (Fig. 8) and 73% for *M. aurantiaca* at the presence points (Fig. 9). For *M. aurantiaca*, only

one scenario out of 30 showed high suitability values within the current range (i.e. the MIROC5 GCB,

RCP85, Chelsa baseline, restricted background).

In Reunion Island, predictions were the most variable between GCMs, but most ensemble models predicted a

low climate suitability across the island. The highest suitability was found for the MIROC5 GCM,

distributed along a band at higher altitudes (Fig. 2b). However, the MESS analyses indicated novel

conditions in the most suitable areas, suggesting model extrapolations in these areas for this GCM (Fig.

S13). According to the response curves (Fig. S14) and the predicted decrease in summer precipitations (Fig.

S3c-d), climate suitability may have been overestimated.

In Madagascar, predictions varied the most with the background extent. Apart from the restricted background

sets, none of the ensemble models identified any suitable area in the future. Nevertheless, we consistently

identified a thin forest band in central Madagascar, south the current distribution, as the most suitable area

across GCMs and scenarios (Fig. 3b). Models identified clear climatic windows, with no issue related to

model extrapolation (Fig. S15, S16).

Fig. 2 Current (a) and projected future (b) environmental suitability for *Phelsuma inexpectata* using Chelsa

as baseline climate data, no prevalence setting and the whole island as a background. Predictions were

filtered to remove agricultural areas. X and Y axes represent the coordinates (WGS84). Black points

represent the occurrence data. Future projections (2070) were estimated from three Global Circulation

models (GCM) and two RCP scenarios. Left panels represent the most optimistic scenario (RCP26) and right

panels represent the most pessimistic scenario (RCP85). Models produced similar results when setting

prevalence to 0.5 (Equal total weights).

Fig. 3 Current (a) and projected future (b) environmental suitability for *Phelsuma inexpectata* using Chelsa

as baseline climate data, no prevalence setting and restricted background. Predictions were filtered to remove

agricultural areas. X and Y axes represent the coordinates (WGS84). Black points represent the occurrence

data. Future projections (2070) were estimated from three Global Circulation models (GCM) and two RCP

scenarios. Left panels represent the most optimistic scenario (RCP26) and right panels represent the most

pessimistic scenario (RCP85). Models produced similar results when setting prevalence to 0.5 (Equal total

weights).

Fig. 4 Current (a) and projected future (b) environmental suitability for *Phelsuma inexpectata* using

Worldclim as baseline climate data. Predictions were filtered to remove agricultural areas. X and Y axes

represent the coordinates (WGS84). Black points represent the occurrence data. Future projections (2070)

were estimated from three Global Circulation models (GCM) and two RCP scenarios. Left panels represent

the most optimistic scenario (RCP26) and right panels represent the most pessimistic scenario (RCP85).

Fig. 5 Current (a) and projected future (b) environmental suitability for *Mantella aurantiaca* using the Chelsa

baseline climate, no prevalence setting and the whole island as a background. Predictions were filtered to

include only forested areas. X and Y axes represent the coordinates (WGS84). Black points represent the

occurrence data. Future projections (2070) were estimated from three Global Circulation models (GCM) and

two RCP scenarios. Top panels represent the most optimistic scenario (RCP26) and bottom panels represent

the most pessimistic scenario (RCP85). Models produced similar results when setting prevalence to 0.5

(Equal total weights).

Fig. 6 Current (a) and projected future (b) environmental suitability for *Mantella aurantiaca* using the Chelsa

baseline climate, no prevalence setting and a restricted background. The background was filtered to include

only the eastern rainforest corridor. X and Y axes represent the coordinates (WGS84). Black points represent

the occurrence data. Future projections (2070) were estimated from three Global Circulation models (GCM)

and two RCP scenarios. Top panels represent the most optimistic scenario (RCP26) and bottom panels

represent the most pessimistic scenario (RCP85). Models produced similar results when setting prevalence to

0.5 (Equal total weights).

Fig. 7 Current (a) and projected future (b) environmental suitability for *Mantella aurantiaca* using the

Worldclim baseline climate. Predictions were filtered to include only forested areas. X and Y axes represent

the coordinates (WGS84). Black points represent the occurrence data. Future projections (2070) were

estimated from three Global Circulation models (GCM) and two RCP scenarios. Top panels represent the

most optimistic scenario (RCP26) and bottom panels represent the most pessimistic scenario (RCP85).

Fig. 8 Variation in climate suitability scores at the occurrence points of *Phelsuma inexpectata* under current

climate and 2070 climate. We show the variability in model predictions related to the climate scenario (RCP

2.6 versus RCP 8.5), the Global circulation model (3 GCMs), the background extent (wide versus restricted),

prevalence setting (no setting versus Equal Total Weights, ETW) and baseline climate (Chelsa versus

Worldclim). Each boxplot is composed of the first decile, the lower hinge, the median, the upper hinge and

the nineth decile.

Fig. 9 Variation in climate suitability scores at the occurrence points of *Mantella aurantiaca* under current
climate and 2070 climate. We show the variability in model predictions related to the climate scenario (RCP
2.6 versus RCP 8.5), the Global circulation model (3 GCMs), the background extent (wide versus restricted),
prevalence setting (no setting versus Equal Total Weights, ETW) and baseline climate (Chelsa versus
Worldclim). Each boxplot is composed of the first decile, the lower hinge, the median, the upper hinge and
the nineth decile.

Discussion

Dramatic and widespread decline in climate suitability

We predict a strong decline of climate suitability in the whole current distribution area of both species by
2070. The subsequent high extinction risk in these species is not surprising, since small occupied areas are
known to be good predictors of vulnerability to climate change (Pearson *et al.*, 2014), but forecasting models

were lacking and we contribute to fill this gap. We also predict that few or no zone will be suitable in the

future across the entirety of their respective regions if those species are not given the opportunity to adapt.

Depending on the Global Circulation model and the scenario, future projections still identified areas with

suboptimal climatic conditions. In Reunion Island, the higher suitability found at ca. 20km west from the

current distribution was presumably driven by an increase in precipitation. *Phelsuma inexpectata* is mostly

present where precipitation is the lowest (< 500mm during the wet season, Fig., S10). The identified area

coincides with the region with the driest conditions in Reunion Island in the future, according to these GCMs

(Fig. S1). In all model sets, the MIROC5 GCM showed a higher—but suboptimal—climate suitability

around the southern coast of the island, with the highest suitability along a band following the upper lands.

This prediction is likely driven by the increase in overall temperature, along with reduced precipitation. The

species lives under the hottest and driest conditions of the island, which fits the thermal requirements of the

species for reproduction in captivity (28°C; McKeown, 1993). However, we believe this GCM is

questionable because (1) the reduction in precipitation does not reflect the possible global increase in cyclone

risk, and (2) the MESS analysis and response curves suggest an overestimation of the suitability scores. Note

that this scenario (i.e. RCP 8.5)—which was the least pessimistic for *P. inexpectata*—is of interest to explore

the widest range of possibilities in the future, but is considered unlikely (Hausfather & Peters, 2020).

In Madagascar, the current distribution is mainly explained by both winter (i.e. the dry season) and summer

(i.e., the wet season) conditions with a short interval of suitability (Fig. S15). An important decline in climate

suitability may therefore be driven by small changes in either winter or summer conditions. During winter,

individuals migrate up the hills to shelter under dead woods and leaves and presumably enter a state of torpor

or hibernation (Randrianavelona *et al.*, 2010; Edmonds *et al.*, 2020). The narrow climatic window may

represent optimal conditions that minimise the risk of desiccation during this period of low activity. The

species is also dependent on summer conditions, with a narrow window of suitable temperature and

precipitations. This is highly consistent with a field study which recorded a surface temperature of 20-23°C

at the occupied sites (Edwards et al., 2019) and an experimental design which found a decrease in activity

when temperature deviates from 21.5°C (Edwards, 2019) with warm and rainy conditions (Fig. S12).

Summer corresponds to the period of reproduction, where females lay eggs under dead leaves on slopes,

which are likely washed down into temporary pond during heavy rain episodes. This may explain the

dependence with summer high precipitation and temperatures. This combination of climatic conditions is

unlikely to be met in the future, except for the MIROC5 GCM under the RCP85 scenario (but this scenario is

unlikely, Hausfather & Peters, 2020). However, we consistently identified potential suboptimal areas in the

central-south of the eastern forest corridor and in one small area in the north east (corresponding to the top of

mountains). For both species *P. inexpectata* and *M. aurantiaca*, the high level of habitat fragmentation

associated to agricultural areas may limit potential distribution shifts, which calls the need for human

intervention.

Sources of uncertainty

The largest source of uncertainty was related to the GCM for *P. inexpectata*, and to the background extend

for *M. aurantiaca*. We followed a protocol that attempted to mitigate most sources of uncertainty,

corresponding to the acceptable standards defined in Araújo *et al.* (2019). The long term, extensive and

repeated efforts dedicated to species sampling enabled to define the current distributions of both species with

high accuracy. Uncertainty map showed a high level of agreement between model replicates (i.e. modelling

techniques, cross-validation runs and pseudo-absence runs; Fig. S17, S18).

In absence of information regarding the dispersal capacity of both species, we used two background

corresponding to past dispersal hypotheses. The background covering the whole respective islands seems the

most appropriate when assuming that only the sea could represent a dispersal barrier and that species could

have dispersed throughout the whole island with regard to past climate and forest cover (Crottini et al. 2019),

as assumed in Fieldsend et al. (2021). This might be the case for both species, especially *P. inexpectata* since

the island is smaller and mostly occupied by a closely related species (*P. borbonica*; Dubos et al. 2021a). The

restricted background corresponds to a hypothesis where *M. aurantiaca* dispersal is limited by current forest

cover, and where *P. inexpectata* is limited to the driest part of Reunion Island. This procedure enabled to

encompass the widest range of possibilities with respect to two extreme hypotheses.

We included the 19 bioclimatic variables for the selection process, which is not recommended in most cases.

However, we reduced the number of variables by removing the intercorrelated ones, and then selected the

most biologically meaningful variables. We used the finest resolution available for climate data (i.e. 30 arc

sec) which seems sufficient to discriminate suitable to unsuitable areas at the scale of Reunion Island and

Madagascar. The inclusion of high-resolution land use variables enabled to improve the realism of both

distributions and provides specific guidelines for conservation applications. In addition, we found a causal

interpretation for the selected variables, which supports the biological significance of our models (Fourcade

et al. 2018; Dubos et al. 2021b). An important limitation is that we did not use scenarios of future land use.

This may lead to an overestimation of the available habitat in Madagascar, since the country is under

important rates of deforestation (Veilledent et al., 2018). In Reunion Island, most of the natural habitats is

incorporated in private properties or public gardens and in steep zones with limited access for agricultural

practices. We believe that the apparent stability in agricultural areas would maintain the applicability of our

results in the future. The inclusion of a range of different greenhouse-gas emission scenarios and GCMs

showed an important uncertainty in the future suitability for *P. inexpectata*. However, this uncertainty is

mostly related to one GCM (i.e. MIROC5), which we assume to be doubtful. The remaining ones

consistently identified the most suitable area by 2070. Uncertainty related to model design was mitigated by

the limitation of model complexity (with the ESM approach and the post-filtering technique), the removal of

collinearity and the testing of a range of input parameters (number of pseudo-absences sets and cross-

validation subsets). Model performance was assessed with random partitions for *M. aurantiaca*, while spatial

partitions are recommended. However, we argued that this methods is not appropriate for highly localised

species due to strong imbalance between spatial blocks. We used multiple evaluation metrics, including

discrimination (AUC and TSS) and reliability (i.e. calibration; Boyce index) metrics, all showing a high

performance overall. We did not account for species dispersal ability, because the purpose was not to predict

potential shifts, but to assess how suitable will be the climate in the future in order to identify candidate sites

for restoration and translocation. Both species have low dispersal ability and their habitat is highly

fragmented. Therefore, the potential for distributional shifts may be strongly limited and future projections

must not be interpreted as the future distributions of our study species. An important limitation may be the

absence of empirical knowledge on species thermal tolerances or other features of their climatic niche,

adaptability and plasticity, which prevents us from determining whether our future projections

underestimated the environmental suitability. Similarly, the extent of the species' fundamental niche is

unknown, and differs from the realised niche as a result of competition, habitat loss and environmental

history, which may particularly important in endemic narrow-ranging species.

A glimmer of hope

Despite the predicted low suitability in climate conditions, it is possible that species persist under changing

conditions through adaptation or plasticity (Chevin *et al.*, 2010; Hoffmann & Sgró, 2011). This hypothesis is

supported by the persistence of two introduced populations of *P. inexpectata* away from their current range in

suboptimal environments (Fig. 2). Their persistence may result either from physiological or behavioural

adaptation while benefiting from a combination of urban island effect and access to microclimate refuges in

anthropogenic structures, as it is the case for *Phelsuma grandis* in Florida (Fieldsend *et al.*, under review). It

is also frequent that the current distribution of a given species represents only a fragment of its climatic niche

(e.g., Guisan *et al.*, 2014). For instance, experimental design on *M. aurantiaca* showed no expression of a

thermal stress during periods of extreme heat (Edmonds *et al.*, 2015). Therefore, predictions of climate

suitability may underestimate the bounds of our model species niches. Further studies are needed to better

characterise their climatic niche, and explore potential adaptive and plastic responses to changing conditions

in these species and, more generally, in other threatened narrow-ranging species. Meanwhile, we encourage

practitioners to implement conservation measures to grant those species a chance to adapt and persist.

Nevertheless, the rate of climate change is generally faster than that of animal adaptive responses (Radchuk

et al., 2019), which stresses the need for urgent actions. In the tropics, extinction risks may be greatly

reduced with the development of land conservation programs provided climate change is mitigated by inter-

governmental actions (Hannah et al., 2020).

Conservation application

In Reunion Island, we identified two potential areas suitable for habitat restoration of *Phelsuma inexpectata*

around the south-western coast and along a mid-altitude band. Despite climate conditions are predicted to be

suboptimal, we believe these might represent the best options to ensure the long-term persistence of the

species. Habitat restoration should be focussed on natural habitats invaded by non-native plants and in

urbanised areas. Habitat restoration will consist in promoting the spread of native species (*Pandanus utilis*,

Latania lontaroides, *Scaevola taccada* and *Psiadia retusa*) for which the species depends on (Bour, 1995).

Finally, natural colonisation of newly suitable environments will be likely impossible for the species due to

its poor dispersal ability and the high level of habitat fragmentation. We therefore encourage the design of

translocation programs in management schemes.

In Madagascar, the area with the highest future climate suitability is located at the south of the current

distribution, along the eastern rainforest corridor between the current distribution area and the north-west of

the Vatovavy-Fitovinany region. The forest cover within the distribution range of *M. aurantiaca* has

experienced a continued decline, regardless of the conservation status of the inhabited area (Piludu *et al.*,

2015; Vieilledent *et al.*, 2018). We recommend to reinforce the level of protection and to improve

governance and management of the Mangabe (Moramanga region) and the Marolambo (Vatovavy-

Fitovinany region) reserves. This can be achieved by promoting the development of alternative economic

solutions through the development of valuable and sustainable activities that mitigate the rate of conversion

of natural areas, the long-term management of soil fertility, and by considering the development of

ecotourism. This species was included into a program that enabled to develop amphibian husbandry

capacities in Madagascar, and that succeeded to establish captive bred colonies for this species

(Rakotonanahary et al. 2017). Efforts of captive husbandry should be maintained (and possibly expanded to

other species) and we encourage the design of translocation programs accounting for both local habitat

characteristics and future climate suitability. Further study of biotic interaction between *M. aurantiaca* and

other amphibians is needed to assess the consequences of its introduction.

Are all endangered narrow-ranging species doomed to extinction?

Narrow-ranging species usually live under very specific environmental conditions and are the most

vulnerable to climate change (Botts *et al.*, 2013). Not only they are more prone to face distribution shifts and

range contractions, but they might also disappear due to the strong alteration of their climatic envelope

throughout their entire region. This may be the case for most narrow-ranging species with a specialised niche

in regions with heterogeneous climates and high levels of endemism such as Madagascar but also Central

America, South East Asia, tropical islands and more generally in tropical rainforests and mountains (Kier *et*

al., 2009). In such regions, small shifts in climatic conditions may induce important changes in local

environmental suitability for endemic—often specialised—species (e.g., Raxworthy *et al.*, 2008). The risk is

greater in tropical regions where species live closer to the upper bound of their thermal tolerance (Tewksbury

et al., 2008; Şekercioĝlu *et al.*, 2012; Dubos *et al.*, 2019). Species that are already under threats are at the

greatest risk because of a large array of synergistic effects (Şekercioĝlu *et al.*, 2012). This might be the case

for our two model species, which are threatened by habitat destruction and invasive species (Dubos, 2013;

Piludu *et al.*, 2015). Among the few Critically Endangered species for which the impact of climate has been

investigated, most predict severe reductions in the climate suitability (e.g., Alamgir *et al.*, 2015; Zhang *et*

al., 2020), with the suitable range of the giant salamander *Andrias davidianus* predicted to decrease by more

than two thirds by 2050 (Zhang *et al.*, 2020). This species has faced a number of threats and climate change

may be leading to imminent extinction despite a larger distribution range. Despite recent efforts developed

for the monitoring of rare species in the tropics (e.g., Dubos et al., 2020), there is still an important lack of

species occurrence data in these regions (Feeley & Silman, 2011). With the newly available high-resolution

climate and land use data, the spectre of eligible species for SDMs has enlarged. We urge filling this data

void by starting to assess the effect of climate change for narrow-ranging species at broader taxonomic

scales, promote field investigations, assess species thermal requirements, and develop proactive conservation actions.

Acknowledgements

We are grateful to all the field workers and their helpers who collected the data. We thank Boris Leroy for useful discussions and for his continued support.

Conflict of Interest Statement

The authors declare no conflict of interests.

Data availability statement

The dataset and the scripts used in this analysis are available as R objects in the supporting information.

References

Acevedo, P., Jiménez-Valverde, A., Lobo, J.M., Real, R., 2012. Delimiting the geographical background in species distribution modelling. J. Biogeogr. 39, 1383–1390. https://doi.org/10.1111/j.1365-2699.2012.02713.x

Alamgir M, Mukul SA, Turton SM, 2015. Modelling spatial distribution of critically endangered Asian elephant and Hoolock gibbon in Bangladesh forest ecosystems under a changing climate. Applied Geography, 60, 10–19.

Araújo MB, Anderson RP, Barbosa AM et al., 2019. Standards for distribution models in biodiversity assessments. Science Advances, 5, 1–12.

Bellard C, Bertelsmeier C, Leadley P, Thuiller W, Courchamp F., 2012. Impacts of climate change on the

future of biodiversity. Ecology Letters, 15, 365–377.

Bellis J, Bourke D, Maschinski J, Heineman K, Dalrymple S., 2020. Climate suitability as a predictor of

conservation translocation failure. Conservation Biology, 1–9.

Bora P, Dolch R, Jenkins R et al.., 2008. Geographical distribution of three species of Malagasy poison frogs

of high conservation priority: Mantella aurantiaca , M . crocea and M . milotympanum. Herpetology

Notes, 1, 39–48.

Botts EA, Erasmus BFN, Alexander GJ., 2013. Small range size and narrow niche breadth predict range

contractions in South African frogs. Global Ecology and Biogeography, 22, 567–576.

Bour R, Probst J, Ribes S., 1995. Phelsuma inexpectata Mertens, 1966, le lézard vert de Manapany-les-Bains

(La Réunion): données chorologiques et écologiques (Reptilia, Gekkonidae). Dumerilia, 2, 99–124.

Butt, N., Chauvenet, A. L., Adams, V. M., Beger, M., Gallagher, R. V., Shanahan, D. F., ... & Possingham, H.

P., 2020. Importance of species translocations under rapid climate change. Conservation Biology.

https://doi.org/10.1111/cobi.13643

Chevin LM, Lande R, Mace GM., 2010. Adaptation, plasticity, and extinction in a changing environment:

Towards a predictive theory. PLoS Biology, 8.

Crottini, A., Orozco-Terwengel, P., Rabemananjara, F.C.E., Susanne Hauswaldt, J., Vences, M., 2019.

Mitochondrial introgression, color pattern variation, and severe demographic bottlenecks in three

species of malagasy poison frogs, genus mantella. Genes (Basel). 10.

https://doi.org/10.3390/genes10040317

Deso G., 2001. Note sur le transport insolite de Geckos verts, le cas du Phelsuma inexpectata. Bulletin

Phaethon, 13, 56.

Draper D, Marques I, Iriondo JM., 2019. Species distribution models with field validation, a key approach

for successful selection of receptor sites in conservation translocations. Global Ecology and

Conservation, 19.

Dubos, N., 2010. Analyse des habitats et déclin du Gecko vert de Manapany, Phelsuma inexpectata Mertens,

1966 sur l'île de La Réunion: implications pour sa conservation. Muséum national d'Histoire Naturelle.

Dubos, N., 2013. New locality record for Phelsuma grandis (Sauria: Gekkonidae) in Reunion, in sympatry

with the Critically Endangered Phelsuma inexpectata. Herpetology Notes, 6, 309–311.

Dubos, N., Dehorter, O., Yves, P., & Viol, I. Le., 2019. Thermal constraints on body size depend on the

population's position within the species' thermal range in temperate songbirds. Global Ecology and

Biogeography, 28(2), 96–106. https://doi.org/10.1111/geb.12805

Dubos, N., Morel, L., Crottini, A., Freeman, K., Honoré, J., Lava, H., Noël, J., Porton, I., Rendryrendry, G.,

Rosa, G., & Andreone, F., 2020. High interannual variability of a climate-driven amphibian community

in a seasonal rainforest. Biodiversity and Conservation, 29, 893–912. https://doi.org/10.1007/s10531-

019-01916-3

Dubos, N., Augros, S., Deso, G., Probst, J., Notter, J., Markus, A., Associations, M., 2021. Here be dragons :

important spatial uncertainty driven by climate data in forecasted distribution of an endangered insular

reptile. BioRxiv. https://doi.org/10.1101/2021.06.14.448338.

Dubos N, Preau C, Lenormand M et al., 2021. Assessing the effect of sample bias correction in species

distribution models when independent data is unavailable. ArXiv:2103.07107.

Dupuy S, Gaetano R (2019. Reunion island - 2017, Land cover map (Pleiades).

Edmonds D, Rakotoarisoa JC, Rasoanantenaina S et al., 2015. Captive husbandry, reproduction, and

fecundity of the golden mantella (Mantella aurantiaca) at the Mitsinjo breeding facility in Madagascar.

Salamandra, 51, 315–325.

Edmonds D, Adamovicz L, Rakotoarisoa JC, Soamiarimampionona J, Harris R., 2020. Seasonal Activity

Patterns of Golden Mantellas (Mantella aurantiaca). Journal of Herpetology, 54, 325–330.

Edwards, W. M., 2019. Conservation of the golden mantella in Madagascar : Integrating in situ and ex situ

research. Kent University.

Edwards, W. M., Griffiths, R. A., Bungard, M. J., Rakotondrasoa, E. F., Razafimanahaka, J. H.,

Razafindraibe, P., Andriantsimanarilafy, R. R., & Randrianantoandro, J. C., 2019. Microhabitat

preference of the critically endangered golden mantella frog in Madagascar. Herpetological Journal,

29(4), 207–213. https://doi.org/10.33256/hj29.4.207213

Feeley KJ, Silman MR., 2011. The data void in modeling current and future distributions of tropical species.

Global Change Biology, 17, 626–630.

Fick SE, Hijmans RJ., 2017. WorldClim 2: new 1-km spatial resolution climate surfaces for global land

areas. International Journal of Climatology, 37, 4302–4315.

Fieldsend T, Dubos N, Krysko K, Raxworthy CJ, Malone S., 2021. In situ adaptation and ecological release

facilitate the occupied niche expansion of a non-native reptile. Ecology and Evolution, 11, 9410–9422.

Fourcade, Y., Engler, J.O., Rödder, D., Secondi, J., 2014. Mapping species distributions with MAXENT

using a geographically biased sample of presence data: A performance assessment of methods for

correcting sampling bias. PLoS One 9, 1–13. https://doi.org/10.1371/journal.pone.0097122

Gábor L, Moudrý V, Barták V, Lecours V., 2019. How do species and data characteristics affect species

distribution models and when to use environmental filtering? International Journal of Geographical

Information Science, 00, 1–18.

Galante PJ, Alade B, Muscarella R, Jansa SA, Goodman SM, Anderson RP., 2018. The challenge of

modeling niches and distributions for data-poor species: a comprehensive approach to model

complexity. Ecography, 41, 726–736.

Gillard M, Thiébaut G, Deleu C, Leroy B., 2017. Present and future distribution of three aquatic plants taxa

across the world: decrease in native and increase in invasive ranges. Biological Invasions, 19, 2159–

2170.

Guisan A, Petitpierre B, Broennimann O, Daehler C, Kueffer C., 2014. Unifying niche shift studies: Insights

from biological invasions. Trends in Ecology and Evolution, 29, 260–269.

Hausfather, Z., & Peters, G. P., 2020. Emissions–the 'business as usual'story is misleading. Nature 577, 618–

620.

Hoffmann AA, Sgró CM., 2011. Climate change and evolutionary adaptation. Nature, 470, 479–485.

IUCN, MNHN, 2010. La Liste rouge des espèces menacées en France. Premiers résultats pour la faune de La

Réunion. 26pp pp.

IUCN SSC Amphibian Specialist Group, 2020. Mantella aurantiaca. IUCN Red List Threat. Species.

https://doi.org/https://dx.doi.org/10.2305/IUCN.UK.2020-3.RLTS.T12776A508612.en

Karger DN, Conrad O, Böhner J et al., 2017. Climatologies at high resolution for the earth's land surface

areas. Scientific Data, 4, 1–20.

Kier G, Kreft H, Tien ML et al., 2009. A global assessment of endemism and species richness across island

and mainland regions. Proceedings of the National Academy of Sciences of the United States of

America, 106, 9322–9327.

Lannuzel G, Balmot J, Dubos N, Thibault M, Fogliani B., 2021. High-resolution topographic variables

accurately predict the distribution of rare plant species for conservation area selection in a narrow-

endemism hotspot in New Caledonia. Biodiversity and Conservation, 30, 963–990

Leroy B, Bellard C, Dubos N et al., 2014. Forecasted climate and land use changes, and protected areas: the

contrasting case of spiders (ed Bradley B). Diversity and Distributions, 20, 686–697.

Liu, C., Newell, G., White, M., 2019. The effect of sample size on the accuracy of species distribution

models: considering both presences and pseudo-absences or background sites. Ecography (Cop.). 42,

535–548. https://doi.org/10.1111/ecog.03188

Mckeown S. (1993. The general care and maintenance of day geckos. In: Advanced Vivarium Systems.

Pearson RG, Raxworthy CJ, Nakamura M, Townsend Peterson A., 2007. Predicting species distributions

from small numbers of occurrence records: A test case using cryptic geckos in Madagascar. Journal of

Biogeography, 34, 102–117.

Pearson RG, Stanton JC, Shoemaker KT et al., 2014. Life history and spatial traits predict extinction risk due

to climate change. Nature Climate Change, 4, 217–221.

Phillips SJ, Dudík M, Elith J, Graham CH, Lehmann A, Leathwick J, Ferrier S., 2009. Sample selection bias

and presence-only distribution models: Implications for background and pseudo-absence data.

Ecological Applications, 19, 181–197.

Piludu N, Dubos N, Razafimanahaka JH, Razafindraibe P, Christian J, Jenkins RKB., 2015. Distribution ,

threats and conservation of a Critically Endangered amphibian (Mantella aurantiaca) in eastern

Madagascar. Herpetology Notes, 8, 119–123.

Platts PJ, Garcia RA, Hof C, Foden W, Hansen LA, Rahbek C, Burgess ND., 2014. Conservation

implications of omitting narrow-ranging taxa from species distribution models, now and in the future.

Diversity and Distributions, 20, 1307–1320.

Porcel, X., Deso, G., Probst, J., & Dubos, N., 2021. Sympatrie entre le Gecko vert de Manapany Phelsuma

inexpectata endémique de la Réunion et le Gecko vert poussière d'or P. laticauda introduits au Domaine

du café grillé : peuvent-ils cohabiter ? Bulletin Phaethon, 53, 36–38.

Probst J-M, Turpin A., 1997. Disparition d'une population de Gecko de Manapany dans le secteur littoral de

Saint-Joseph. Bulletin Phaethon, 6, 104.

Radchuk V, Reed T, Teplitsky C et al., 2019. Adaptive responses of animals to climate change are most likely

insufficient. Nature Communications, 10, 1–14.

Randrianavelona R, Rakotonoely H, Ratsimbazafy J, Jenkins RKB, 2010. Conservation assessment of the

critically endangered frog Mantella aurantiaca in Madagascar. African Journal of Herpetology, 59, 65–

78.

Rakotonanahary, T. F., Dawson, J., Andreone, F., Rakotoarisoa, J. C., Edmonds, D., 2017. Amphibian

Captive Breeding Programs for Madagascar: Where We Are Now and the Road Ahead. FrogLog, 118,

37–39.

Raxworthy CJ, Pearson RG, Rabibisoa N et al., 2008. Extinction vulnerability of tropical montane endemism

from warming and upslope displacement: a preliminary appraisal for the highest massif in Madagascar.

Global Change Biology, 14, 1703–1720.

Şekercioĝlu çaĝan H, Primack RB, Wormworth J., 2012. The effects of climate change on tropical birds.

Biological Conservation, 148, 1–18.

Tewksbury JJ, Huey RB, Deutsch CA., 2008. Putting the Heat on Tropical Animals The Scale of Prediction.

Science (New York, N.Y.), 320, 1296–1297.

Thomas CD., 2011. Translocation of species, climate change, and the end of trying to recreate past ecological

communities. Trends in Ecology and Evolution, 26, 216–221.

Vences, M. and Raxworthy, C.J., 2008. Mantella aurantiaca. 2008: e.T12776A3381123.

http://dx.doi.org/10.2305/IUCN.UK.2008.RLTS.T12776A3381123.en.

Vieilledent G, Grinand C, Rakotomalala FA, Ranaivosoa R, Rakotoarijaona JR, Allnutt TF, Achard F, 2018.

Combining global tree cover loss data with historical national forest cover maps to look at six decades

of deforestation and forest fragmentation in Madagascar. Biological Conservation, 222, 189–197.

Vinson J.M., 1975. Notes on the reptiles of Round Island. Mauritian Institute Bulletin, 8, 49–67.

Vollering, J., Halvorsen, R., Auestad, I. & Rydgren, K., 2019. Bunching up the background betters bias in

species distribution models. Ecography, 42, 1717–1727.

Westwood R, Westwood AR, Hooshmandi M, Pearson K, LaFrance K, Murray C., 2020. A field-validated

species distribution model to support management of the critically endangered Poweshiek skipperling

(Oarisma poweshiek) butterfly in Canada. Conservation Science and Practice, 2, 1–15.

Zhang Z, Mammola S, Liang Z et al., 2020. Future climate change will severely reduce habitat suitability of

the Critically Endangered Chinese giant salamander. Freshwater Biology, 65, 971–980.