Seizures initiate in zones of relative hyperexcitation in a zebrafish epilepsy model

James E. Niemeyer[1], Poornima Gadamsetty[1], Chanwoo Chun[2], Sherika Sylvester[2], Jacob P. Lucas[1], Hongtao Ma[1,3], Theodore H. Schwartz[1,3,#], Emre Aksay[2#]

1. Department of Neurological Surgery, Weill Cornell Medicine, New York, NY

2. Institute for Computational Biomedicine and the Department of Physiology and Biophysics, Weill Cornell Medicine, New York, NY

3. Feil Family Brain and Mind Research Institute, Weill Cornell Medicine, New York, NY

#. Correspondence may be addressed to

Theodore H Schwartz, MD
Department of Neurological Surgery
Weill Cornell Medicine
New York, NY 10065
Email: schwarh@med.cornell.edu

Emre Aksay, PhD
Department of Physiology and Biophysics
Weill Cornell Medicine
New York, NY 10021
Email: ema2004@med.cornell.edu

Abstract

Seizures are thought to arise from an imbalance of excitatory and inhibitory neuronal activity. While most classical studies suggest excessive excitatory neural activity plays a generative role, some recent findings challenge this view and instead argue that excessive activity in inhibitory neurons initiates seizures. We investigated this question of imbalance in a zebrafish seizure model with multi-regional two-photon imaging of excitatory and inhibitory neuronal activity using a nuclear-localized calcium sensor. We found that seizures consistently initiated in circumscribed zones of the midbrain before propagating to other brain regions. Excitatory neurons were both more prevalent and more likely to be recruited than inhibitory neurons in initiation as compared with propagation zones. These findings support a mechanistic picture whereby seizures initiate in a region of hyper-excitation, then propagate more broadly once inhibitory restraint in the surround is overcome.

Introduction

Excitation and inhibition are countervailing forces of brain activity that, in normal brains, operate in balance. However, in certain pathological brain states, such as epileptic seizures, this excitation:inhibition (E:I) balance goes awry. While traditional doctrine held that seizures resulted from hyperactive excitatory drive[1-4], recent studies in various models have suggested a generative role for inhibitory neuronal subtypes in ictal onset. This revised picture has been supported by observations of increased inhibitory activity in pre-seizure periods[5-8]. For example, low-voltage fast seizures in humans are marked by early inhibitory cell firing prior to peak excitatory cell activity[7]. Evidence for a direct causal role of inhibitory cells in seizure onset is also reported[9-11]. For example, blocking the effects of inhibitory activity with a GABA antagonist (picrotoxin) can paradoxically prevent spontaneous seizures in a genetic mouse model of human autosomal-dominant frontal lobe epilepsy[9]. Another study showed that optogenetic stimulation of a specific interneuron subtypes (parvalbumin- or somatostatin-positive cells) can initiate seizures[11]. Together, these finding challenge the traditional concept of excitatory neuron hyperactivity as the primary driver of seizure initiation.

However, the role of inhibitory cells in seizure onset is still widely debated. Not only is there data showing that inhibitory cell activation can inhibit seizure onset[12], but activation of inhibitory cells has been clearly shown to shorten seizure duration[13, 14]. While differences in seizure models, experimental preparations, and recording techniques could explain these disparate results, the explanations may be more subtle. While one recent study found that inhibitory cells can exert either anti- or pro-seizure effects depending on the timing of activation[15], another showed that the effects of excitatory and inhibitory cells may be spatially dependent[16]. These findings suggest that the role of E:I balance in seizures may not be as simple as "more E" or "earlier I", but rather that the effects of E:I balance on ictal onset and evolution may be spatiotemporally contingent.

Understanding how E:I balance plays into ictogenesis has several experimental requirements. First, a large region of brain, inclusive of various interconnected structures, must be analyzed during seizure periods. Ideally, this areal sampling would include both the site of seizure initiation as well as sites of later seizure propagation. Second, the neural activity analyzed should simultaneously include both E and I information that can be separated with

single-cell resolution. Finally, the neural activity should be recorded in awake animals whose
activity can best recapitulate seizures lacking the confounding effects of anesthesia.
These criteria can be relatively easily met in the zebrafish preparation[17]. These
vertebrates exhibit significant homology with humans at the genetic[18], neurochemical, and brain
structural[19] levels. Their transparency, rapid development, and small size in the first few weeks
after fertilization allows whole brain optical access at sub-cellular resolution throughout the
brain. Their genetic tractability permits tissue and cell-type specific experimentation. Further,
their ease-of-handling facilitates both high-throughput drug screening[20] and high-resolution
imaging and electrophysiological studies in the unanesthetized animal. Prior work on seizures
using this animal model has already shown similar electrographic profiles[21], glial involvement[22],
and responses to seizure treatments[23] as found in human epilepsy.
Here we used the zebrafish model to investigate spatiotemporal variations in E:I balance
during seizure initiation and propagation. We monitored seizure dynamics using both
electrophysiological recording and two-photon calcium imaging that enabled tracking of
neuronal dynamics across the brain at the single cell level. We found specific regions of the
brain consistently associated with seizure initiation, while others with seizure propagation. Using
transgenic specification to separate excitatory and putative inhibitory populations, we found that
the ratio of E:I activation at ictal onset was greatest in initiation zones. This finding demonstrates
the importance of early excitation at triggering ictal events in the zebrafish model and
challenges recent reports arguing that early inhibitory neuron activity is always the key driver in
initiating seizures.

Materials and Methods

Data collection

69   All experiments were performed in accordance to the guidelines and approval of Weill Cornell
70   Medicine's Institutional Animal Care and Use Committee.

Zebrafish Preparation:
Mutant zebrafish larva (*Tg(HuC:h2b-GCaMP6f) x Tg(Vglut2a:dsRed)*)[24, 25] were used at 5 - 8
days after fertilization - the h2b-tag localizes GCaMP to the nucleus to disambiguate the site of
origin of the signal by eliminating contamination from axonal and dendritic sources. After
anesthetizing for 1 min in 0.1 g/L tricaine-methanesulfonate (MS222, VWR TCT0941- 025G),
zebrafish were mounted dorsal up in a 35 mm petri dish (35–1008, BD Falcon) lid containing a
bed of 1.5 ml hardened 1% agarose (Invitrogen 15510-027) by embedding with droplets of 1.7%
low temperature agarose (Sigma A0701-100G). Once the agarose hardened, a small incision in
the skin was made (typically) above the neuropil in the right optic tectum for electrophysiology
(see below). The agarose around the tail of the zebrafish was removed to monitor tail
movement. The larva was then paralyzed by point incubation in droplets of 300 µM
pancuronium bromide (Sigma P1918-10MG) and covered in 2.5-3 ml of glucose-free 10%
(vol/vol) Evans medium (2.9 mM KCl, 1.2 mM MgCl2, 2.1 mM CaCl2, 134 mM NaCl, and 10mM
HEPES, pH 7.8) once tail movement ceased Epileptiform activity was induced with bath
application of 15 mM pentylenetetrazol (PTZ) (Sigma P6500-50G), which is more common and
effective than alternatives such as pilocarpine[22] and 4-Aminopyridine[26]. Recordings were
performed within 2 hours after embedding zebrafish in agarose.

Electrophysiology
Local-Field Potential (LFP) recordings were acquired with metal microelectrodes during initial
seizure characterization. One ~2-3 MΩ tungsten microelectrode (WE30013.0F3, Microprobes)
was inserted in the optic tectum record LFP and a silver wire was placed in the Evans medium
to server as a reference. In some experiments, a second tungsten microelectrode was placed in
either telencephalon or hindbrain region to confirm the global synchronous of ictal events.
Electrophysiological recordings were amplified 1000x with a microelectrode amplifier (Model
1800, A-M Systems, Carlsborg, WA), digitized at 10K Hz using CED Power 1401, and stored
with a computer runningSpike2 software (Cambridge Electronic Design).
During calcium imaging, the LFP was recorded using a glass microcapillary to avoid
obstructing fluorescence excitation and light collection. Capillaries (1B120F-4, WPI) were pulled
to a fine tip with vertical puller (P30, Sutter Instruments) to a tip diameter of 2-3 µm. The
capillaries were filled with 4M NaCl after front-loading molten agarose (1.7% Ultrapure Agarose,
Thermo) to the tip to prevent diffusion of the electrode solution once inserted into the brain.
When tested in Evans medium, these electrodes had a resistance of 1-3 MΩ. The electrode was
inserted through incision in the skin into the neuropil of the right optic tectum. Recordings were
made with a microelectrode amplifier (Model 1800, A-M Systems, Carlsborg, WA) using
bandpass filtering (between 0.1 Hz and 1 kHz) at 10 kHz sampling rate and digitized using
custom MATLAB code.
108
Two-photon Imaging
A custom-built two photon microscope was used for imaging calcium activity and fluorescence
in the embedded zebrafish. A MaiTai HP laser (Newport) was used to focus a 915 nm light
beam through a water immersion 40x/0.8 NA objective (OBJ1; LUMPlanFl40XW/IR2, Olympus)
onto the fish. Multiple separate horizontal planes of the fish were acquired sequentially in the
dark at 2 Hz (256×256 pixels, 550x550 µm). The emitted light from the zebrafish larva was
collected by the objective and directed towards two dichroic mirrors (DM1, 720dcxruv; DM2,
565dcxr; Chroma Technology) which reflected this light into two separate channels of green (F3;
ET525\50m-2p; Chroma) and red (F2; ET605\70m-2p, Chroma) to be then amplified by the
photomultiplier tubes (PMTs, R3896; Hamamatsu). A substage photodetector (PD, PDA36A,
Thorlabs) detected light scattered from the focal plane and together with signal from the PMTs
was processed by MATLAB based software ScanImage[27] to generate fluorescence images.
121

Analysis

Electrophysiology
Electrographic data was used to validate and examine ictal events during 2-photon imaging.
The spectrogram of the LFP was calculated using short-time Fourier transform (*spectrogram* in
MATLAB), with a window width of 0.1 s. Time of onset in the epileptiform events was
determined using a statistical method based on the mean and 3 times the standard deviation
(SD) of the power between 1-60 Hz during a 5 s interictal window. For electrographic analyses
(**Fig1**, **Fig S1**), the first increase in power > 3 SD above the baseline was established as the
onset of the event. The termination was set at the moment the LFP power returned within 3 SD
of baseline.
132
133

134 **Image analysis**
135
Preprocessing
Images were collected by ScanImage and stored in tiff format. These data files were
subsequently de-interleaved in Fiji[28] to extract the green (GCaMP) and red (DsRed) channels.
Rigid motion correction was applied in MATLAB 2018a using the NoRMCorre algorithm and
toolbox[29] on the red channel images, which did not contain activity-dependent fluorescence. The
transformation matrix calculated on the red channel in each fish imaging series was then
applied to the green GCaMP6f channel, yielding two motion-corrected tiff image series.
We used these data sets to examine ictal event activity across brain regions with two
overarching analysis methods: First, we used pixelwise methods to characterize ictal event
initiation, spread, and involvement across brain regions. Second, we performed single-cell
analyses to observe ictal event development simultaneously in VGlut2a excitatory neurons and
putative inhibitory interneurons, paying particular attention to E:I ratio and differential
involvement of these two cell classes as well as network-level differences between them.
149
Correlation Coefficient Masks
During ictal events, GCaMP6f fluorescence in the image increases as the ictal event initiates
and propagates through the brain. We first extracted relevant brain areas by using the
preprocessed GCaMP6f data to calculate pixelwise Pearson's correlation with the whole image
average activity in the green GCaMP6f channel. To achieve this, we initially performed 2-
dimensional Gaussian filtering on the imaging data ($\sigma=2$ pixels, ~4 micrometers) in MATLAB
(*imgaussfilt*) and then smoothed individual pixels over time using a lowpass Butterworth filter (1st
order, cutoff frequency of .2 Hz). These data are hereafter referred to as "smoothed images".
The individual pixel traces were then compared with the average of all pixel values in the
image and a cutoff of .6 correlation (Pearson's *r*) was used to remove any pixels not sufficiently
correlated with the average image activity. To remove any remaining noise, any smoothed
pixels not connected in groups of at least 80 smoothed pixels were removed. The resulting
image mask, called the correlation coefficient mask, simply represents the extent of brain that
was involved in seizure events over individual files. This mask was applied in subsequent
analyses of ictal event development.
165
Lag analyses for individual ictal events
Individual ictal events were next detected in smoothed image series with a semi-automated
method. First, general ictal event identification was performed: GCaMP6f fluorescence change
was calculated as the frame-by-frame difference in average fluorescence, and peaks were
extracted with a minimum peak height of average plus 2 SD. A minimum peak-to-peak distance
of 8 seconds was also included. Due to slight variations in noise or atypically spreading ictal
events in some data sets the standard deviation cutoff was adjusted or performed on the
average fluorescence activity instead of the difference. When average activity was used to find
peaks, we then used 10 sec windows around those extracted times to find the peak first
derivative activity. We note that these adjustments to the automated process are only included
to produce accurate windows for finding ictal onsets and would not change the onset times. This
general identification produced the time at which the ictal event was increasing most rapidly in
the fluorescence data. All individual ictal events were verified by visual examination of both the
average GCaMP6f activity and the GCaMP6f image series videos.
Next, we used these ictal event times to create a window of whole-image ictal event
activity of 6 seconds prior to this point, called the global template. This ictal event onset

template was then compared with all individual pixels from the correlation coefficient mask in
time windows around the individual ictal events: a 16 s window was used, starting at 10 s prior
to peak and lasting 6 s past peak, a large time window that always captured the ictal event
onset.
Then, the time of maximum correlation between the pixel trace and global template was
found for each pixel from the mask yielding a matrix, for each ictal event, of the time when each
pixel was most correlated to the global ictal event template. Pixels not significantly correlated
(Pearson's r, p<.05) with the ictal event average or with a peak correlation below r=.6 were
excluded. When visualized as a heatmap this matrix provides a "lag map" of where the ictal
event activity began and spread across the imaging field of view. To remove any remaining
noise, these lag maps were subsequently smoothed by a 2-D Gaussian kernel (σ=6 pixels, ~11
micrometers). To account for the line-scanning nature of our 2-photon images, a temporal
transformation matrix was applied to shift every pixel by its true time in milliseconds (e.g., the
first pixel of a single tiff image was scanned 256 ms prior to the pixel in the middle of the image).
Finally, when displayed, heat color indicates time relative to the average onset time calculated
across pixels, negative values indicating pixels that were active earlier, positive values
indicating pixels that were active later. A total of 10 separate animals produced N=177 ictal
events.
From these lag maps, we used MATLAB's edge detection functions to extract the
regions of the fish brain that were ictal event-active at the earliest 500 ms. We also applied an
area requirement of 300 connected pixels (~.5% of the 256x256 images) to exclude noise.
These regions were used to measure which anatomical brain areas were most commonly
invaded first by the ictal events. These regions were labelled the "initiation sites" in later single
cell analyses. Other regions of the brain that were invaded later by the seizure are referred to as
"propagation sites".
To examine brain region involvement over different data sets, we compared the edges of
initiation sites, per event, with the anatomical fish images (average fluorescence) and recorded
the amount of times that an initiation site contained a particular brain region, given that the brain
region was present (e.g., if the habenula was not visible in a specific ictal series then it was not
included in the analyses of early onsets).
Correlations of these maps within fish were calculated as Pearson's r (across seizures
within the same FOV) to measure similarity of seizure propagation within individual animals.
These correlations were thus only calculated when more than one event occurred in a particular
FOV within a fish.
A summary map (**Fig. 3d**) was created within an individual fish by setting a temporal
cutoff (500 ms) for early activity in the lag map, creating a binary mask from this cutoff, and then
averaging these binary masks over different ictal events. Thus, if a pixel is involved in 100% of
the ictal events of one fish then it will receive a 1, if a pixel is involved in half of the ictal events it
will receive a .5, etc. The correlation matrix (**Fig. 3e**) was calculated on ictal series with multiple
seizures in FOVs containing all listed brain regions.
222
Single cell analysis
Single cells were extracted from the unsmoothed GCaMP6f images using an automated custom
MATLAB algorithm with an added optional step that allowed for manually marking any single
cells missed by our algorithm. First, within a single data set, the lowest quartile of GCaMP6f
activity frames were used to create an average GCaMP6f image. Background subtraction was
performed using structural element disks in MATLAB (with *imadjust*) and then an algorithm was

used to iteratively step through all pixel luminance values (0 to 255) down to the lowest 20%
and extract all connected regions containing at least 4 pixels (~2 microns/pixel), but removing
any cells containing more than 40 pixels. All images were visually examined and any missed
cells that could be visually determined were added manually in MATLAB with a custom script
(using *imfreehand*), but typically fewer than 5 cells were added at this step. Finally, we applied a
correlation threshold cutoff to remove any single cells whose activity was not at least r=.3
correlated with the average activity of all the extracted cells. On average, our analyses extracted
351 seizure-active cells per image following these steps.
In separate cellular composition analyses, where we were interested in examining any
detectable cells without requiring that they be seizure-active, we followed the above methods
but used a lower Pearson's cutoff of r=.1, which was low enough to include more cells than the
r=.3 cutoff but high enough to exclude artifacts (such as occur on skin edges and near the
animal's eyes).
To separate putative inhibitory (expressing only GCaMP) and excitatory cell types (co-
expressing DsRed and GCaMP in VGlut2.1+ cells) we took the red intensity values from
individual cells and plotted their histogram. These histograms are characterized by two peaks:
the first peak, lower red values, denotes the GCaMP-positive cells that do not co-express
DsRed (putative inhibitory interneurons); the second peak, higher red intensity values, denotes
the GCaMP-positive cells that also co-express DsRed (putative excitatory neurons). These
histograms were fit by a dual gaussian mixture model, and the mean and standard deviation of
both component gaussians were extracted (*fitgmdist* in MATLAB). The cutoff used in our study
for labeling a cell as excitatory was two standard deviations past the first gaussian's mean. In
some cases, artifacts from the zebrafish skin, eyes, or a tilt in the animal's body could affect
these histograms. In those cases, the histogram fitting was performed regionally by manually
selecting anatomical regions (applying MATLAB's *imfreehand*) and then performing the dual
gaussian fit. All images were visually examined after cell type dissociation by plotting cell types
on composite red/green images of the larval zebrafish to verify appropriate labelling by the
algorithm. In two data sets we could not extract E vs I cell type differences (due to either
imaging or fluorophore labelling technical difficulties), so the 10 ictal events in those data series
were not included in our E:I single cell analyses leaving N=167 ictal events.
259
Seizure onsets
In order to compare results across animals and seizure events, we extracted ictal onset times
from each ictal event for all single cells. General ictal events were found above (see Lag
Analyses), but for single cell analyses we sought to determine the exact time of cell onsets
relative to a population onset. Thus, we applied a similar correlation method as the pixelwise lag
analyses above but with global templates that were centered around the peak second derivative
of single cell population activity, the time of fastest recruitment of individual neurons to a seizure
event.
Single cell involvement in seizures was determined by identifying the times at which
individual cells became significantly involved in the ictal events (**Fig. S4**). To find these times in
single neurons we filtered the cell GCaMP6f profiles by temporally smoothing in 2 s windows
(boxcar average) and then compared these traces in time windows around ictal event onsets
that were found in the lag analyses above.
Global response templates in this analysis were created by taking the average cell
response from a period of equal length between the first and second peak derivative of the
average single cell fluorescence traces. This created a template centered on the peak

acceleration point of the ictal event, a time that more accurately represents true ictal event onset
compared to other methods of cutoffs such as half-height or 2-standard deviations above the
mean. We refer to this peak acceleration time of the population activity as the ictal onset.
279   Next, we found the peak linear correlation (Pearson's *r*) between this template and
individual cell fluorescence traces in a window from 8 s before and 3 s after the period of the
global template. This wide temporal window was important because our method of global
template creation depends on the rise time of the ictal event: slowly evolving ictal events will
have longer templates than faster events. (Note: the correlation window used for single cell
onsets was 11 seconds plus the length of the global template, making it a window approximately
equal to the 16 sec window used in lag analyses above). Our algorithm required that a single
cell be at least *r*=.8 correlated with the global template to be included as having an onset in the
ictal event, a strict requirement to reduce the chance of erroneous single cell onset
classifications. The onset time for a single cell was classified as the time where it reached 5% of
peak correlation (above the .8 cutoff) with the global template. This 5% requirement was
included because many cells displayed wide peaks of correlation with the global template:
taking the earliest peak correlation most accurately represented the onset of a single cell in the
ictal event. Because some cells displayed small fluctuations in fluorescence that coincided with
ictal onsets, when cells exhibited more than one *r*=.8 peak during the time window analyzed, we
took the later *r*=.8 peak to prevent erroneous early-onset classifications. Finally, all individual
cell onsets were temporally adjusted for the 2-photon system's scanning (as done for the
pixelwise lag analysis, shifting by the pixel scan time from the cell centroid), and the population
activity was temporally corrected by 256 ms (the midpoint time of the frame scan). This method
was applied to all data sets and all cells regardless of E or I classification.
299   When temporally aligning data we used the average ictal onset time calculated across all
300   cells significantly recruited to each individual seizure, a procedure analogous to the relative
301   times calculated in our lag analyses above. Thus, data are displayed relative to this time zero
302   point.
303   Inter-ictal periods were defined as all times outside of ictal events (after PTZ
304   application). We classified the termination of ictal events as the point at which the average
305   fluorescence activity returned to its value at the second derivative onset, which was used to
306   separate ictal vs interictal time periods for later analyses.
307
Excitation:Inhibition (E:I) balance
We examined E:I relationships by calculating an E:I ratio index. We use this method instead of
simple E/I ratio because in some cases we observed low numbers of cells in our initiation sites.
E:I index values in initiation regions were calculated by summing these cell types within the
earliest active brain areas (see above, Lag analyses). E:I ratios in propagation zones were
calculated the same way, using remaining areas of the brain that were significantly involved in
seizures (propagation zones). The index value was calculated as
315
316    $$EI_{index} = \frac{E-I}{E+I} \ ,$$
317
where E and I are counts of E and I cells. The E:I ratio index provides a measure of the relative
differences in amounts of E and I cells located in the initiation and propagation regions.

We measured excitatory and inhibitory cell recruitment throughout ictal events by
temporally aligning all events and then calculating the fraction of E and I cells significantly
involved in the ictal events before and after the onset.

Ictal vs interictal brain state and network analyses
Dimensionality metric
A population of N neurons makes an activity trajectory in the N dimensional space over some
time points (T). The effective dimensionality of the trajectory can be quantified with the
uniformity of the singular values (λ_i) of the trajectory matrix (NxT, mean-subtracted).

330
$$q = \frac{\left(\sum_{i=1}^{\min(N,T)} \lambda_i\right)^2}{\sum_{i=1}^{\min(N,T)} \lambda_i^2}$$

If the singular values are perfectly uniform, all principal components explain equal amount of
variance, and therefore the effective dimensionality should maximal. If only one singular value is
non-zero, then the effective dimensionality should be 1. For our dimensionality analysis, the
number of cells (N) was variable but always greater than the number of time points
(T=10s*2Hz=20 samples) of our time window, so the upper limit of the dimensionality was fixed
at T. This allowed us to compare the effective dimensionality across trials with different number
of cells without further adjustments. A dimensionality at an instantaneous time-point t is
estimated by calculating the dimensionality over the time window centered around t. The
instantaneous estimation was used to construct a time-series from which we could visualize the
time evolution of dimensionality.

Excitatory & inhibitory correlations
Populations of excitatory cell pairs (e-e) and inhibitory cell pairs (i-i) were analyzed for their
correlations around ictal events. The correlation metric (*C*) is an average pairwise correlation
value of the calcium traces over a time window (1.5 s),

348
$$C_{J,K} = \frac{1}{|J||K|} \sum_{j \in J} \sum_{k \in K} corr(X)_{j,k} \, ,$$

where X is a matrix whose rows are a time series of a single cell within the rolling time window,
and J and K are sets of indices for excitatory or inhibitory cells. *corr*(X) returns a correlation
matrix. Self-correlations (diagonal elements) are ignored. A correlation at an instantaneous
time-point t is estimated by calculating the correlation over the time window centered around t.
The instantaneous estimation was used to construct a time-series from which we could visualize
the time evolution of correlation.

Results

We present our investigation of the spatiotemporal dynamics of E:I balance during seizures as
follows. First, we introduce the experimental preparation and describe electrophysiological
characterization of inter-ictal and ictal events in this epilepsy model (**Fig. 1**). Next, we present a
pixel-based timing analysis (**Fig. 2**) that is used to create maps of the spatiotemporal dynamics

in the brain during seizures (**Fig. 3**). We then examine timing at a cellular level in excitatory and
inhibitory populations in regions designated as 'initiation' and 'propagation' zones (**Fig. 4**).
Finally, we confirm that findings from our single-cell analysis hold at the network level (**Fig. 5**)
and when examined with spatial specificity using correlation analysis (**Fig. 6**).
Experiments were performed using larval zebrafish with all neurons expressing
GCaMP6f and VGlut2.1+ cells expressing DsRed (**Fig. 1a**). The first fluorophore, H2B-tethered
GCaMP6f, is a fast nuclear-localized calcium sensor that is localized to the nucleus to eliminate
contamination from neuropil signal and enable better separation of signals between cells[25]. The
second fluorophore, DsRed, was used to help separate excitatory from inhibitory neurons.
VGlut2.1 is a vesicular glutamate transporter gene found in the vast majority of glutamatergic
cells in the larval zebrafish[24]. The largest proportion of VGlut2.1- cells are inhibitory GABAergic
neurons, with non-trivial contribution from excitatory VGlut2.2+ cells (primarily cerebellar
granule neurons) and inhibitory glycinergic neurons. Due to the purely excitatory nature of the
VGlut2.1+ population and the primarily inhibitory nature of the VGlut2.1- populations, below we
refer to the former as 'excitatory (E)' neurons and the latter as 'inhibitory (I)' neurons.
Our first goal was to characterize seizures measured electrographically. Seizures were
induced using bath applied PTZ in larval zebrafish that were held in agar. While the definition of
a "seizure" can be flexible and the exact features of seizures likely vary between species and
models, our electrophysiology data below demonstrates separable ictal-like and interictal-like
events with electrographic signatures similar to those observed in other animal models and
humans. For simplicity, here we use the word "seizure" to refer to the longer ictal events as
others have done for PTZ-induced ictal activity in zebrafish[22, 26, 30]. In initial two-electrode
experiments, we placed electrodes in different brain regions (telencephalon, optic tectum, or
hindbrain) and found that ictal events developed concurrently after ~15 minutes of PTZ
application. Electrodes placed in the optic tectum consistently captured the largest set of ictal
events (unpublished observations)—hence, in subsequent experiments, we used only one
electrode placed in the neuropil of the optic tectum. **Figure 1b** shows an example
electrophysiology recording, demonstrating ictal-like events lasting for several seconds ('ictal',
filled circle symbol), followed by quiescent periods and occasional brief interictal-like events
('interictal', open symbol), as well as their corresponding power spectrograms (**Fig. 1b)**. Ictal
events (left) were marked by an initial low frequency power increase (orange), a brief cross-
frequency spike with most power <20 Hz (red), a sustained phase with a more uniform cross-
frequency distribution (often up to 100 Hz) that lasted for several seconds (green), and a
recovery phase at lower frequency with gradual relaxation (blue). In contrast, interictal events
(right) only exhibited the short cross frequency event (red) before moving to a briefer recovery
phase than seen with ictal events (blue). Both forms of event then returned to baseline. While
more complicated classification schemes could be used, separation of ictal from interictal events
based on duration provided a robust method as ictal events typically lasted 4 times longer than
interictal ones (**Fig. 1c**; Wilcoxon signed-rank test, N=9 fish, 1685 events, p=0.003).
We next employed simultaneous cellular-resolution 2-photon imaging of larval zebrafish
along with the electrophysiological recordings. Ictal events were clearly observable in population
activity and within specific brain regions (**Fig. 1d**; **Fig. S1**). Interictal events, on the other hand,
did not lead to global changes in activity. Examining global activity spatially, we found that ictal
events did not typically initiate in or recruit the telencephalon (**Fig. 1e).** While activity in the optic
tectum (OT) and rhombomere 1 (R1) correlate strongly with the whole-brain average calcium
response, the telencephalon (Tel) does not. A correlation matrix built with average calcium
activity in different brain regions from multiple series of seizures from different fish (N=5 fish, 22
seizures) indicates that activity in the telencephalon and habenula is largely uncorrelated from
the rest of the brain during ictal events.
412

Figure 1: Ictal and interictal activity in the larval zebrafish. **a** Top: experimental setup for 2-photon imaging and electrophysiology in larval zebrafish during seizures. Bottom: single imaging plane revealing distribution of VGlut+ (magenta) and VGlut- (green) cells in a variety of brain regions (R: rhombomere). **b** Electrophysiological signatures of ictal- and interictal-like events in the zebrafish. (i) Top: Local Field Potential (LFP, top); Middle: spectrogram; Bottom: total power during a period of ictal (closed circle) and interictal (open circle) events; (ii) Similar organization as above but at finer time resolution, with indication of distinct sections (orange, red, green, blue) characterizing the time-course of the two types of events. **c** Top, Middle: histogram of event duration times for two fish (green) and best fit (black) with a gaussian mixture model; Bottom: for all fish, duration time for ictal and interictal events. **d** Top: LFP, Middle: global change in GCaMP6f fluorescence; Bottom: regional change in fluorescence during a series of ictal (closed circle) and interictal events (OT, optic tectum; R1, rhombomere 1; Tel, telencephalon). **e** Matrix of correlations in the activity among various brain regions (Tel: telencephalon, Hab: habenula, Thal: thalamus, PrT: pretectum, OT: optic tectum, R1 rhombomere 1).

To examine the involvement and timing of different brain areas in PTZ seizures we next developed a method to extract seizure propagation maps within individual fish. We first used correlation to global average ictal activity to create a mask that excluded pixels where there were no changes in fluorescence related to the seizure (**Fig. 2a**). The activity in the retained pixels was not monolithic, with some regions beginning to show elevations in fluorescence well before other regions (**Fig. 2b,** red arrows). To examine these temporal variations more carefully, we correlated pixel-level activity to that of a template of ictal activity initiation defined from a segment of the global fluorescence event (**Fig. 2c,** *middle*), incrementally shifting the template in time to identify when activity in particular pixel began to rise (see Methods). The template shift times for each pixel defined a propagation map for each ictal event (**Fig. 2c**, *right*). This template matching procedure allowed identification of the earliest portions of the ictal activity in

439   each pixel and was less susceptible to errors than procedures that relied upon threshold
440   crossings or features such as peak fluorescence.
441

442
443   **Figure 2. Ictal event mapping. a** Left: average fluorescence over a series of ictal events in one brain;
444   Middle: pixel-wise correlation with brain average; Right: the correlation mask. **b** A single seizure's
445   fluorescence activity is shown over time; red arrows highlight early activity. **c** Left: sample pixels and their
446   activity profiles (*middle*) over a single seizure are shown; Right: the corresponding smoothed lag map for
447   this seizure shows the relative activation times for each pixel with respect to the brain average.
448
We applied this algorithm to create a lag map for each individual ictal event (N=10 fish,
N=177 ictal events). **Figure 3a** shows the lag maps of 7 ictal events in one larval zebrafish (the

same animal as **Fig. 2**). This revealed a similarity between neighboring ictal events (e.g., Events
1-4 vs 5-7). In this animal the left cerebellum and the optic tectum showed consistent early
involvement in every ictal event. To quantify how similar ictal propagation appeared between
ictal events, we measured the correlation coefficient of the 2-D spatiotemporal lag maps of all
pairs of ictal events for this animal (**Fig. 3b**). This similarity was quantified for the entire data set
by pooling within-animal correlations, revealing a positively skewed distribution (**Fig. 3c,**
average Pearson's r=0.42 +/- 0.25, N=177 seizures).
458

461   ***Figure 3. Ictal events evolve similarly within fish, but initiate in different regions across fish***. **a** Lag
462   maps for seven seizures in one animal. **b** Correlations between the events in (**a**). **c** correlations of all pairs
463   of ictal events recorded. **d** A summary map of the ictal events in (**a**). Violin plots (**e**) and a schematic fish
464   brain map (**f**) of early-onset regions show where ictal events initiated most commonly across different fish.
465

We next developed a scheme to summarize these spatiotemporal patterns and identify
regions consistently involved in seizure initiation. Lag maps for individual ictal events were
binarized by assigning a value of 1 to all pixels where neuronal activity initiated at least 500 ms
in advance of the population event, and a value of 0 to all other pixels. Averaging these mask

arrays across all events within a single fish provided a 2-dimensional summary showing where
ictal events initiated most frequently within an individual animal (**Fig. 3d**). For the animal whose
seizures are displayed in **Figure 3a**, we see that the left cerebellum and portions of the left
tectum exhibit the highest probability of involvement in ictal initiation. Across all animals (N=10),
the most common initiation zones were the optic tectum, thalamus, pre-tectum, and
cerebellum/R1 (**Fig. 3e-f**). Propagation zones were defined as the remainder of the brain to
which the seizure spread outside the initiation zone.
Having identified the ictal initiation and propagation zones, we next focused on the issue
of E:I balance in these different regions during seizures. We first asked if there were differences
between the initiation and propagation zones in the relative numbers of excitatory and inhibitory
cells. Taking advantage of the ease with which single neurons can be distinguished with
nuclear-localized GCaMP6f, we labelled individual soma using standard computer vision
algorithms and separated them into excitatory and inhibitory cells based on the intensity of the
DsRed fluorophore (total N=19,434 cells across 9 fish; example event and seizure-recruited
cells shown in **Fig. 4a**). An event-specific comparison of the differences between the initiation
and propagation regions in composition was performed by calculating for each ictal event a
value for an E:I composition index; the index was defined such that a region with only excitatory
cells had a composition index value of 1, all inhibitory cells an index of -1, and an equal number
of excitatory and inhibitory cells an index of 0. E:I composition in initiation regions was -0.07 +/-
0.29 (SEM), while E:I composition in propagation regions -0.30 +/- 0.29 (**Fig 4b**; at cutoff
requiring at least 10 E and 10 I cells per initiation site, N=49 seizures, Wilcoxon signed-rank test
p<0.001). This difference corresponds to a roughly 61% increase in the E:I ratio in the initiation
zone relative to the propagation zone calculated over all animals and all seizures. This finding
was robust to variation in the minimal number of cells of each type allowed in each region (**Fig.
S3**).
We next investigated if there were differences between the initiation and propagation
zones in the relative numbers of excitatory and inhibitory cells that were *recruited* into the ictal
event. Cells were deemed to be recruited into the seizure if they exhibited of correlation of 0.8 or
better with the global ictal event (**Fig. S3**). The locations of such cells during one seizure event
are shown in **Figure 4c**, revealing a much larger E:I ratio of recruited cells within the initiation
zone than in the propagation zone. To quantify this observation across, we calculated an E:I
recruitment index, defined as above but only for recruited cells. E:I recruitment in initiation
regions was 0.04 +/- 0.25 (SEM), while E:I recruitment in propagation regions was -0.20 +/-
0.28, a significant difference (p<.0001, N=167, Wilcoxon signed-rank test; **Fig. 4d** at cutoff
requiring 10 E and 10 I cells per initiation site). This corresponds to a roughly 64% increase in
E:I ratio in the initiation zone relative to the propagation zone across seizures, and was robust
across varying cutoffs of cell counts (**Fig. S3**). Thus, on average there is both a greater fraction
of excitatory cells in initiation zones relative to propagation zones, and these excitatory cells are
also recruited at a higher fraction within the initiation zone.

509
Figure 4. Ictal events preferentially initiate in excitatory cell-rich regions of the zebrafish brain. a A
sample fish image with E and I cells differentially labelled and its associated lag map and initiation zone for
a single ictal event analyzed in this figure. **b** E:I index in initiation and propagation zones (i) and within-
seizure E:I comparisons (ii). **c** Single cell locations in a seizure, split by initiation vs propagation zone
location (i) and cell counts from initiation and propagation zones (ii). **d** E:I index of cells recruited to seizures
in initiation and propagation zones (i) and within-seizure E:I index comparisons**. e** Individual cell activity
traces (smoothed and interpolated for clarity) are shown for one ictal event. Color groups denote onset
times from the earliest 50% (red) and latest 50% (blue) of onsets. Boxplots show distributions of onset
times. **f** Left: sample excitatory and inhibitory cells from initiation zones; Right: their associated calcium
traces during this event. Detected onset times are plotted on traces. **g** Left: average number of cells at
onset across all ictal events grouped by initiation or propagation zone membership; Right: the ratio of cells
at onset (E:I) over time, normalized to by the number of inhibitory cells at onset. **h** E:I ratio index calculated
in initiation (orange) and propagation (blue) zones. All error bars are mean +/- SEM.

We next examined how E:I recruitment during the ictal event varied over time in the
initiation and propagation zones. We observed asynchronous single cell onsets during ictal
events: **Figure 4e** shows smoothed and interpolated single cell fluorescence traces for one ictal
event, colored by initiation time, demonstrating the temporal spread of single cell activity that
these seizures typically displayed. As expected, across our data set we observed significantly
earlier onsets in the initiation zones compared to propagation zones (**Fig. 4g**, **Fig. S3c**).
Interestingly, in the propagation region, the number of newly recruited excitatory cells was
always less than the number of newly recruited inhibitory cells (**Fig. 4g,** *left*, solid lines, N=167
seizures). In contrast, in the initiation zone, the number of newly recruited excitatory cells was
comparable to the number of newly recruited inhibitory cells at the beginning and end of the ictal
event, but was notably higher at a time approximately 1 second ahead of the ictal onset. To see
this difference more clearly, we normalized these recruitment curves to the peak of the number
of newly recruited inhibitory cells (**Fig. 4g**, *right*), revealing 1) that the number of newly recruited
cells peaked earlier in the initiation zone, and 2) a difference of over 50% at peak in the ratios of
newly activated excitatory vs inhibitory cells between the initiation and propagation sites (white
arrow, **Fig. 4g**). Quantifying these observations using the E:I recruitment index introduced
above, we see that significant differences in the pattern of recruitment between the initiation and
propagation regions are present in the period -1.5 to 0 seconds before ictal onset (N=36
seizures with at least 10 E and 10 I cells in initiation zone, Wilcoxon signed-rank test p<0.0001,
Fig. 4h).
We next examined if this privileged position for excitatory neurons during the initiation of
seizures was also apparent in higher order relationships between excitatory and inhibitory cells.
We began by using Principal Components Analysis to look at the overall degree of coordination
in excitatory versus inhibitory populations during the seizure initiation process. For both groups,
the number of dimensions needed to explain population dynamics dropped dramatically at the
time of ictal onset (**Fig. 5a-b**). This collapse to lower dimensionality, an indication of greater
coordination within a population, was most pronounced for the excitatory population (**Fig. 5c**;
pre-PTZ N=2781, post-PTZ N=167, right: I-E dimensionality differences, Wilcoxon signed-rank
test, *** p<0.001, **p<0.01, n.s. not significant).

Figure 5. Dimensionality of the excitatory and inhibitory cell population activities. **a** Left: Eigenspectrum of the excitatory population activity over 10 sec.-wide rolling window, normalized to the greatest eigenvalue per time point. Vertical line at time 0 is the ictal onset point. Right: same analysis done for inhibitory cell population. **b** Average trends of the dimensionality. Vertical line at time 0 is the average ictal onset point. Dimensionality over 10 sec moving window (magenta: e-e; green: i-i). **c** Left: Dimensionalities of i-i pairs and e-e pairs shown separately for pre-PTZ, -12~-2 second and -2~8 second periods relative to the average ictal onset point. Right: differences in I-E dimensionalities at -2~8 seconds.

To examine this increased coordination of E cells vs I cells with greater temporal specificity and regional context, we next analyzed correlation structure within these cell types by calculating a correlation metric (*C*) in a rolling time window (**Fig. 6a**). We focused on correlations between excitatory-excitatory (E-E) and inhibitory-inhibitory (I-I) cell pairs. Average correlation was generally low for both pairings during baseline conditions and inter-ictal periods (**Fig. 6b**, *top*). Correlation increased dramatically for both pairings around the time of ictal initiation, with correlation change in the initiation zone on average larger and more advanced in time than that in the propagation zone. We found that E-E correlation was slightly elevated relative to I-I correlation during ictal onset in both regions (**Fig. 6b**, insets); this elevation was clearer when looking at the correlation difference, which was largest around the time of ictal onset in the initiation zone (**Fig. 6b-c,** Wilcoxon signed-rank test, N=36 seizures with 10 E and 10 I cells in initiation zones, *** p<0.001, ** p<0.01, n.s. not significant; **Fig 6d**, within-seizure correlation differences in initiation and propagation sites in events with 10 E and 10 I cells). Thus, these correlation patterns further suggest that the coordinated activation of excitatory neurons in the initiation zone plays an important role in triggering ictal events.

583
Figure 6. Excitatory and inhibitory cell network correlations. a Visualization of activity correlations at the cross-section of the whole brain. Pairs with correlation values higher than 0.999 are shown. Top: average normalized fluorescence trace. Middle: correlations during interictal period. Magenta: highly correlated e-e; Green: i-i. Bottom: correlations during ictal period. **b** Average trends of interictal to ictal transition for the initiation and propagation zones. Top: correlation metric calculated over 1.5 sec.-wide rolling window (magenta: e-e; green: i-i). Bottom: difference of e-e and i-i correlations; **c** Correlation metric values of i-i pairs and e-e pairs shown separately for -6~-2 second and -2~2 second periods relative to the average ictal onset point. **d** Within-seizure correlations of e-e cell pairs (top) and i-i cell pairs (bottom) between initiation and propagation zones.

Discussion

In this study, we addressed the question of E:I balance in seizure initiation and propagation using the zebrafish model. The main technical limitation afflicting prior studies of this topic has generally been the need for broad, cell-specific sampling of the widespread neuronal substrate inclusive of the ictal onset zone and all areas of potential propagation. We overcame these challenges using a global imaging strategy in the larval zebrafish, an optically transparent vertebrate preparation allowing for simultaneous imaging of multiple cell types. We demonstrated a privileged role for excitation over inhibition in the seizure initiation zone, while regions that were invaded later were biased towards stronger inhibitory cell recruitment.

Seizure dynamics in the larval zebrafish
One specific relevance of our findings will be to aid efforts to identify seizure medications with high-throughput screening. Zebrafish are increasingly used to test potential drug treatments for diseases such as epilepsy[23, 31]. High-throughput drug screening is a primary method of therapeutic discovery[32], and the zebrafish offers significant advantages in this regard due to the ease with which drugs can be delivered uniformly to the brain while behavioral changes are

monitored *en masse*. However, while such high-throughput studies can be quite useful in identifying promising classes of drugs, optimization of both the therapeutic and its dosing will benefit greatly from a mechanistic understanding of the pathophysiology being addressed. The larval zebrafish affords several advantages for understanding the cellular and circuit mechanisms that have been used to study the normal brain, including high-resolution optical access globally, a suite of genetic tools for monitoring and manipulating specific cell types, and the capacity to study all constitutive elements of an intact circuit. Here we have harnessed some of this power to begin understanding the pathophysiology of seizures. The preponderance of excitatory activity in the initiation zone points towards exploiting this model to screen, with greater granularity, classes of drugs that can preferentially regulate the activity of glutamatergic neurons in midbrain structures.

We triggered seizure events in the zebrafish model using PTZ, a chemoconvulsant commonly used in therapeutic screening. PTZ is the only pharmacological seizure-inducing drug employed at the initial identification stage by the NINDS Epilepsy Therapy Screening Program[33]. PTZ is known to induce seizures in humans[34, 35] and lower seizure threshold in epileptic patients[36]. PTZ-induced seizures in the zebrafish occurred spontaneously, repeatedly, and throughout the brain, consistent with earlier reports[22, 26, 37, 38]. Furthermore, the ictal LFP had spectral features[39] similar to those observed in rodent models[40] and human patients[41]: at initiation, ictal events displayed an abrupt broad-frequency power increase, followed by a longer period where power was restricted to a slightly lower frequency range, concluding with a return to baseline power at the time of termination. This commonality supports the idea that the PTZ model of ictogenesis in the zebrafish is of general relevance for the study of seizures in other species.

Spatio-temporal consistency of seizure initiation and propagation

Regions of ictal onset and propagation were generally conserved within fish, but somewhat varied between animals. Rodent slice experiments have shown that the seizure foci can change slowly over time: one focus can dominate for several minutes before being replaced by another region[42]. However, such studies have also found that certain brain regions (e.g., hippocampal CA2-3) and cell layers (neocortical layer 5 pyramidal cells) are predisposed to act as pacemakers in seizure-generating activity[43, 44]. Our findings extend these results to an awake, whole-brain model. We found a significant level of consistency in initiation and propagation patterns within fish, although correlations weakened over tens of minutes. While there was some variation between animals, we found that midbrain structures had a quite high propensity to serve as initiation zones. We also confirmed the surprising finding that the telencephalon (homologous with human cortex) was one of the last regions to be recruited in the seizing zebrafish brain[22]. Thus, our results provide further evidence that certain subcortical sites may serve a prominent role in generalized seizure initiation[45-48] though these results do not negate a role for cortical contributions.

E:I balance differs in initiation and propagation zones

Synchronous excitation has long been thought to drive epileptic seizures[3], while recent cell-specific studies have suggested a role for inhibitory neurons in both seizure initiation[6, 7] and termination[12, 14]. Our finding adds to the modern E:I discussion by reaffirming the classic distinction between ictal initiation and propagation. Specifically, we observed a higher ratio of recruited excitatory:inhibitory cells near the seizure initiation site compared to locations that were recruited later by the same seizure. A recent study in anesthetized mice reported a similar finding: the E:I activity ratio, calculated on averaged parvalbumin-positive (PV) and non-PV interneuron activity, was higher in the initiation zone compared to propagation zones during the pre-seizure period[16]. However, the experiment involved focal injection of 4-AP, where the assumption that the injection and onset sites are superimposed may not always be correct. Our

experiments circumvent this ambiguity by allowing seizures to develop spontaneously in
anatomic regions with lower threshold in a model that permits whole-brain imaging so that we
could identify distinct initiation zones on an event-by-event basis.
Some studies in rodents[6, 13] and humans[7] have reported earlier activity from inhibitory
cell populations before recruitment to seizure events. We did not observe significantly different
onset timing between E and I cell types. We speculate that this could be due to differences in
recording location (initiation vs propagation zones), or possibly due to variations in technique.
For example, Khoshkoo et al. used an optogenetic kindling model[13], and the firing rate changes
between inhibitory and excitatory cells observed by Miri et al. were on the order of just ~2 Hz[6],
which our calcium imaging may not have been sufficiently sensitive to detect. Our findings
therefore do not rule out the possibility of earlier inhibitory cell activation, but do support a model
of ictogenesis that depends on higher-weighted excitatory cell recruitment at initiation sites.
Numerous studies have shown that surround inhibition, present outside of an ictal focus,
can play a role in restraining seizures[16, 49-53]. This restraining penumbra is a region characterized
by massive synaptic barrages, but little neuronal firing, until a temporal point at which the
seizure annexes this region[16, 50, 52-54]. In our study using a nuclear-localized GCaMP to focus
purely on somatic activity, we did not observe significant differences in E and I cell timing, but
did observe that seizure initiation zones were heavily weighted towards excitation (higher E:I)
while later propagation regions were weighted towards inhibition (lower E:I). In this drug-bathed
model, initiation zones are selected based on inherent increased E:I whereas propagation
zones transiently resist rapid spread, presumably through a mechanism based on higher
relative local inhibition. Future work in this model may shed further light on this inhibitory
restraint by combining imaging and intracellular recording[53], allowing simultaneous
measurement of synaptic barrages, spiking, and whole-brain activity during seizure dynamics.
686
Conclusion
The goal of epilepsy research is to understand how normal brain activity transitions to an ictal
state and how the event spreads across adjacent normal brain regions. In our model we found
that midbrain and diencephalic regions were the most common sites of seizure initiation.
Further, during individual seizures we observed significantly higher E:I ratios in initiation
compared with propagation zones, supporting a model of seizures in which excitatory activity
overruns inhibitory restraint to initiate the seizure while distal regions remain more protected by
an inhibitory veto[50, 51, 54]. The differential E:I balance we report across brain regions provides a
framework for designing and interpreting drug screening studies. Further, our finding of
hyperexcitation in initiation zones could be leveraged to perturb neural activity in a cell-specific
manner to attenuate seizure initiation or even arrest ongoing seizures.
698
699
Acknowledgments

701 The authors thank Catherine Schevon, Michael Wenzel, Jyun-you Liou, and Jonathon Victor as
702 well as members of the Aksay and Schwartz labs for constructive discussion of this manuscript.

Supplemental data figures

Figure S1. Identical timing of ictal events in calcium and electrophysiological signals. A sample series of ictal events is shown from **Figure 1**, with two seizures highlighted (pink). The first event is shown in higher temporal magnification (i), as well as the fifth event (ii). A 2 Hz notch filter was applied to the electrophysiological signals in the bottom panels, and the calcium activity is taken from the entire imaging frame. Our onset detection method was applied to all seizures (red dashed line).

714
715
716 ***Figure S2. Single cell activity.*** **a** Individual cell activity is shown (rows) for the same cells prior to PTZ
717 administration (*left*) and interictal activity prior to an ictal event (*right*) after PTZ administration. Excitatory
718 and inhibitory cell types are shown separately and sorted by detected onset time within this ictal event. **b**
719 Top: cell population activity demonstrates a series of four ictal events; Bottom: single cell activity during
720 these events is shown sorted by average onset across events.
721

Figure S3. Associated with Figure 4 E:I and onset timing. a Left: the difference in average onset time
between excitatory and inhibitory cell populations is shown (positive values indicate earlier excitatory cell
onset) for whole brain Right: same analysis for initiation zones. Initiation zones are divided into varying
sizes (All seven violins: N=167, N=167, N=101, N=62, N=36, N=18, N=12 ictal events; none were significant
by α=.05, Wilcoxon signed-rank test). **b** Probability of participation was calculated across all E and I cells
that participated in more than one seizure (N=40 data series with at least 2 seizures, 159 seizures total;
N.S. α=.05, Wilcoxon rank-sum test). **c** Recruitment times between E and I cell types (N=1264, N=1135,
N=12143, N=14992; N.S., α=.05, Wilcoxon rank-sum test) and between initiation and propagation sites

(**** p<.0001, Wilcoxon signed-rank test). **d** Number of E and I cells detected in regions that were
determined to be initiation and propagation zones, regardless of whether these cells participated in seizure
events, shown across multiple size cutoffs (N=49, N=18, Wilcoxon signed-rank test, *** p<.001). For all
data (167 seizures), with no cell count cutoffs, E:I composition in initiation regions was -0.069 +/- 0.042
(SEM), while E:I composition in propagation regions -0.232 +/- 0.024, corresponding to a roughly 29%
increase in E:I ratio in the initiation zone relative to the propagation zone across all seizures. **e** Number of
E and I cells recruited in initiation and propagation zones shown across multiple size cutoffs (N=36, N=12,
p<.001, Wilcoxon signed-rank test, ** p<.01, *** p<.001). For all data (167 seizures), with no cell count
cutoffs, E:I recruitment in initiation regions was 0.034 +/- 0.043 (SEM), while E:I recruitment in propagation
regions was -0.162 +/- 0.022, a significant difference corresponding to a roughly 40% increase in E:I ratio
in the initiation zone relative to the propagation zone across all seizures.

750
Figure S4. Single cell onset detection. Ictal events were detected from global (all cells) activity, and a
template of global activity was created around the peak second derivative of this trace. Top: this template
was slid along single cell activity traces to determine the estimated onset of the single cell based on its
correlation with the global template (see Methods). Bottom, left to right: the single cell location in the fish,
the correlation at each step of the analysis with the best fit location marked, and the template aligned with
the single cell onset estimate. This single cell's onset was ~1.8 seconds earlier than the population ictal
onset.

References

1.   Ayala GF, Dichter M, Gumnit RJ, Matsumoto H, Spencer WA. Genesis of Epileptic Interictal Spikes - New Knowledge of Cortical Feedback-Systems Suggests a Neurophysiological Explanation of Brief Paroxysms. Brain Res 1973;52:1-17.

2.   Dichter MA, Ayala GF. Cellular Mechanisms of Epilepsy - a Status-Report. Science 1987;237:157-164.

3.   Matsumoto H, Marsan CA. Cortical Cellular Phenomena in Experimental Epilepsy - Ictal Manifestations. Exp Neurol 1964;9:305-&.

4.   Treiman DM. GABAergic mechanisms in epilepsy. Epilepsia 2001;42:8-12.

5.   Lillis KP, Kramer MA, Mertz J, Staley KJ, White JA. Pyramidal cells accumulate chloride at seizure onset. Neurobiol Dis 2012;47:358-366.

6.   Miri ML, Vinck M, Pant R, Cardin JA. Altered hippocampal interneuron activity precedes ictal onset. Elife 2018;7.

7.   Elahian B, Lado NE, Mankin E, et al. Low-voltage fast seizures in humans begin with increased interneuron firing. Ann Neurol 2018;84:588-600.

8.   Toyoda I, Fujita S, Thamattoor AK, Buckmaster PS. Unit Activity of Hippocampal Interneurons before Spontaneous Seizures in an Animal Model of Temporal Lobe Epilepsy. J Neurosci 2015;35:6600-6618.

9.   Klaassen A, Glykys J, Maguire J, Labarca C, Mody I, Boulter J. Seizures and enhanced cortical GABAergic inhibition in two mouse models of human autosomal dominant nocturnal frontal lobe epilepsy. P Natl Acad Sci USA 2006;103:19152-19157.

10.   Yekhlef L, Breschi GL, Lagostena L, Russo G, Taverna S. Selective activation of parvalbumin- or somatostatin-expressing interneurons triggers epileptic seizurelike activity in mouse medial entorhinal cortex. J Neurophysiol 2015;113:1616-1630.

11.   Shiri Z, Manseau F, Levesque M, Williams S, Avoli M. Interneuron Activity Leads to Initiation of Low-Voltage Fast-Onset Seizures. Ann Neurol 2015;77:541-546.

12.   Calin A, Stancu M, Zagrean AM, Jefferys JGR, Ilie AS, Akerman CJ. Chemogenetic Recruitment of Specific Interneurons Suppresses Seizure Activity. Front Cell Neurosci 2018;12.

13.   Khoshkhoo S, Vogt D, Sohal VS. Dynamic, Cell-Type-Specific Roles for GABAergic Interneurons in a Mouse Model of Optogenetically Inducible Seizures. Neuron 2017;93:291-298.

14.   Krook-Magnuson E, Armstrong C, Oijala M, Soltesz I. On-demand optogenetic control of spontaneous seizures in temporal lobe epilepsy. Nat Commun 2013;4.

15.   Magloire V, Cornford J, Lieb A, Kullmann DM, Pavlov I. KCC2 overexpression prevents the paradoxical seizure-promoting action of somatic inhibition. Nat Commun 2019;10.

16.   Wenzel M, Hamm JP, Peterka DS, Yuste R. Acute Focal Seizures Start As Local Synchronizations of Neuronal Ensembles. J Neurosci 2019;39:8562-8575.

17.   Kalueff AV, Echevarria DJ, Stewart AM. Gaining translational momentum: More zebrafish models for neuroscience research Preface. Prog Neuro-Psychoph 2014;55:1-6.

18.   Howe K, Clark MD, Torroja CF, et al. The zebrafish reference genome sequence and its relationship to the human genome. Nature 2013;496:498-503.

19.   Kalueff AV, Stewart AM, Gerlai R. Zebrafish as an emerging model for studying complex brain disorders. Trends Pharmacol Sci 2014;35:63-75.

20.   Steenbergen PJ, Richardson MK, Champagne DL. The use of the zebrafish model in stress research. Prog Neuro-Psychoph 2011;35:1432-1451.

21.   Baraban SC, Taylor MR, Castro PA, Baier H. Pentylenetetrazole induced changes in zebrafish behavior, neural activity and c-FoS expression. Neuroscience 2005;131:759-768.

22.   Verdugo CD, Myren-Svelstad S, Aydin E, et al. Glia-neuron interactions underlie state transitions to generalized seizures. Nat Commun 2019;10.

23.   Baraban SC, Dinday MT, Hortopan GA. Drug screening in Scn1a zebrafish mutant identifies clemizole as a potential Dravet syndrome treatment. Nat Commun 2013;4.

24.   Higashijima S, Mandel G, Fetcho JR. Distribution of prospective glutamatergic, glycinergic, and GABAergic neurons in embryonic and larval zebrafish. J Comp Neurol 2004;480:1-18.

25.   Dunn TW, Mu Y, Narayan S, et al. Brain-wide mapping of neural activity controlling zebrafish exploratory locomotion. Elife 2016;5.

26.   Liu J, Baraban SC. Network Properties Revealed during Multi-Scale Calcium Imaging of Seizure Activity in Zebrafish. Eneuro 2019;6.

27.   Pologruto TA, Sabatini BL, Svoboda K. ScanImage: flexible software for operating laser scanning microscopes. Biomed Eng Online 2003;2:13.

28.   Schindelin J, Arganda-Carreras I, Frise E, et al. Fiji: an open-source platform for biological-image analysis. Nat Methods 2012;9:676-682.

29.   Pnevmatikakis EA, Giovannucci A. NoRMCorre: An online algorithm for piecewise rigid motion correction of calcium imaging data. J Neurosci Methods 2017;291:83-94.

30.   Turrini L, Fornetto C, Marchetto G, et al. Optical mapping of neuronal activity during seizures in zebrafish. Sci Rep-Uk 2017;7.

31.   Eimon PM, Ghannad-Rezaie M, De Rienzo G, et al. Brain activity patterns in high-throughput electrophysiology screen predict both drug efficacies and side effects. Nat Commun 2018;9.

32.   Macarron R, Banks MN, Bojanic D, et al. Impact of high-throughput screening in biomedical research. Nat Rev Drug Discov 2011;10:188-195.

33.   Pitkanen A, Buckmaster PS, Galanopoulou AS, Moshe SL. Models of Seizures and Epilepsy Second edition Preface. Models of Seizures and Epilepsy, 2nd Edition 2017:Xxv-Xxvi.

34.   Cooper K, Fink M. The Chemical Induction of Seizures in Psychiatric Therapy Were Flurothyl (Indoklon) and Pentylenetetrazol (Metrazol) Abandoned Prematurely? J Clin Psychopharm 2014;34:602-607.

35.   de Meduna L. New methods of medical treatment of schizophrenia. Arch Neuro Psychiatr 1936;35:361-363.

36.   Friedlander WJ. Relation of Metrazol Eeg-Convulsant Threshold and Alpha Index. Electroen Clin Neuro 1962;14:291-&.

37.   Rosch RE, Hunter PR, Baldeweg T, Friston KJ, Meyer MP. Calcium imaging and dynamic causal modelling reveal brain-wide changes in effective connectivity and synaptic dynamics during epileptic seizures. Plos Comput Biol 2018;14.

38.   Hadjiabadi D, Lovett-Barron M, Raikov I, et al. Maximally selective single cell target for circuit control in epilepsy. bioRxiv 2020:2020.2010.2020.340364.

39.   Hunyadi B, Siekierska A, Sourbron J, Copmans D, de Witte PAM. Automated analysis of brain activity for seizure detection in zebrafish models of epilepsy. J Neurosci Meth 2017;287:13-24.

40.   Li YH, Li JJ, Lu QC, Gong HQ, Liang PJ, Zhang PM. Involvement of Thalamus in Initiation of Epileptic Seizures Induced by Pilocarpine in Mice. Neural Plast 2014;2014.

41.   Bin Altaf MA, Yoo J. A 1.83 mu J/Classification, 8-Channel, Patient-Specific Epileptic Seizure Classification SoC Using a Non-Linear Support Vector Machine. Ieee T Biomed Circ S 2016;10:49-60.

42.   Tsau Y, Guan L, Wu JY. Initiation of spontaneous epileptiform activity in the neocortical slice. J Neurophysiol 1998;80:978-982.

43.   Pinto DJ, Patrick SL, Huang WC, Connors BW. Initiation, propagation, and termination of epileptiform activity in rodent neocortex in vitro involve distinct mechanisms. J Neurosci 2005;25:8131-8140.

44.   Prince DA, Connors BW. Mechanisms of Epileptogenesis in Cortical Structures. Ann Neurol 1984;16:S59-S64.

45.   Bernasconi A, Bernasconi N, Natsume J, Antel SB, Andermann F, Arnold DL. Magnetic resonance spectroscopy and imaging of the thalamus in idiopathic generalized epilepsy. Brain 2003;126:2447-2454.

46.   Avoli M, Gloor P. Role of the Thalamus in Generalized Penicillin Epilepsy - Observations on Decorticated Cats. Exp Neurol 1982;77:386-402.

47.   Avoli M, Rogawski MA, Avanzini G. Generalized epileptic disorders: An update. Epilepsia 2001;42:445-457.

48.   Yaksi E, Jamali A, Verdugo CD, Jurisch-Yaksi N. Past, present and future of zebrafish in epilepsy research. Febs J 2021.

49.   Prince DA, Wilder BJ. Control Mechanisms in Cortical Epileptogenic Foci - Surround Inhibition. Arch Neurol-Chicago 1967;16:194-+.

50.   Trevelyan AJ, Schevon CA. How inhibition influences seizure propagation. Neuropharmacology 2013;69:45-54.

51.   Trevelyan AJ, Sussillo D, Watson BO, Yuste R. Modular propagation of epileptiform activity: Evidence for an inhibitory veto in neocortex. J Neurosci 2006;26:12447-12455.

52.   Liou JY, Ma HT, Wenzel M, et al. Role of inhibitory control in modulating focal seizure spread. Brain 2018;141:2083-2097.

53.   Jayant K, Wenzel M, Bando Y, et al. Flexible Nanopipettes for Minimally Invasive Intracellular Electrophysiology In Vivo. Cell Rep 2019;26:266-+.

54.   Schevon CA, Weiss SA, McKhann G, et al. Evidence of an inhibitory restraint of seizure activity in humans. Nat Commun 2012;3.