TITLE: A human antibody reveals a conserved site on beta-coronavirus spike proteins and confers protection against SARS-CoV-2 infection

SUMMARY: A human mAb isolated from a COVID-19 donor defines a protective cross-neutralizing epitope for pan-β-CoV vaccine design strategies

AUTHORS:
Panpan Zhou[1,3,4,*], Meng Yuan[2,*], Ge Song[1,3,4,*], Nathan Beutler[1,*], Namir Shaabani[1,*], Deli Huang[1], Wan-ting He[1,3,4], Xueyong Zhu[2], Sean Callaghan[1,3,4], Peter Yong[1,3,4], Fabio Anzanello[1,3,4], Linghang Peng[1], James Ricketts[1], Mara Parren[1], Elijah Garcia[1], Stephen A. Rawlings[5], Davey M. Smith[5], David Nemazee[1], John R. Teijaro[1], Thomas F. Rogers[1,5,†], Ian A. Wilson[2,3,4,6,†], Dennis R. Burton[1,3,4,7,†], Raiees Andrabi[1,3,4,†]

AUTHOR AFFILIATIONS:
[1]Department of Immunology and Microbiology, The Scripps Research Institute, La Jolla, CA 92037, USA.
[2]Department of Integrative Structural and Computational Biology, The Scripps Research Institute, La Jolla, CA 92037, USA.
[3]IAVI Neutralizing Antibody Center, The Scripps Research Institute, La Jolla, CA 92037, USA
[4]Consortium for HIV/AIDS Vaccine Development (CHAVD), The Scripps Research Institute, La Jolla, CA 92037, USA.
[5]Division of Infectious Diseases, Department of Medicine, University of California, San Diego, La Jolla, CA 92037, USA.
[6]Skaggs Institute for Chemical Biology, The Scripps Research Institute, La Jolla, CA 92037, USA.
[7]Ragon Institute of Massachusetts General Hospital, Massachusetts Institute of Technology, and Harvard University, Cambridge, MA 02139, USA.
[*]These authors contributed equally to this work.
[†]Corresponding author. Email: trogers@scripps.edu (T.F.R.); wilson@scripps.edu (I.A.W); burton@scripps.edu (D.R.B.); andrabi@scripps.edu (R.A.).

ABSTRACT

Broadly neutralizing antibodies (bnAbs) to coronaviruses (CoVs) are valuable in their own right as prophylactic and therapeutic reagents to treat diverse CoVs and, importantly, as templates for rational pan-CoV vaccine design. We recently described a bnAb, CC40.8, from a coronavirus disease 2019 (COVID-19)-convalescent donor that exhibits broad reactivity with human beta-coronaviruses (β-CoVs). Here, we showed that CC40.8 targets the conserved S2 stem-helix region of the coronavirus spike fusion machinery. We determined a crystal structure of CC40.8 Fab with a SARS-CoV-2 S2 stem-peptide at 1.6 Å resolution and found that the peptide adopted a mainly helical structure. Conserved residues in β-CoVs interacted with CC40.8 antibody, thereby providing a molecular basis for its broad reactivity. CC40.8 exhibited in vivo protective efficacy against SARS-CoV-2 challenge in two animal models. In both models, CC40.8-treated animals exhibited less weight loss and reduced lung viral titers compared to controls. Furthermore, we noted CC40.8-like bnAbs are relatively rare in human COVID-19 infection and therefore their elicitation may require rational structure-based vaccine design strategies. Overall, our study describes a target on β-CoV spike proteins for protective antibodies that may facilitate the development of pan-β-CoV vaccines.

MAIN TEXT

Introduction

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has led to the current global pandemic (*1-3*). SARS-CoV-2 is a virus that belongs to the coronaviridae family of which six members have previously crossed into humans from animal reservoirs and established widespread infections (*4, 5*). These include four endemic human coronaviruses (HCoVs) (HCoV-229E, HCoV-HKU1, HCoV-OC43, HCoV-NL63) responsible for non-severe, seasonal infections (*4*) as well as SARS-CoV-1 and MERS-CoV (Middle East Respiratory Syndrome CoV) that are associated with high morbidity and mortality in humans (*6, 7*). Among the seven HCoVs, SARS-CoV-2 closely resembles SARS-CoV-1 and, to lesser degree, MERS-CoV. Together with HCoV-HKU1 and HCoV-OC43, these viruses belong to the *β-coronavirus* genus (*4, 5*). SARS-CoV-2 is highly transmissible in humans and causes coronavirus disease-2019 (COVID-19), associated with severe respiratory failure leading to high morbidity and a reported mortality of about 0.7 to 2% of infected individuals worldwide (*2, 8, 9*). There are considerable concerns that future coronavirus spillovers will trigger new pandemics (*10-15*).

Coronavirus pandemic preparedness may consider responses through establishment of techniques for rapid generation of specific reagents to counter the emerging coronavirus and control spread. An alternative is to seek to identify broadly neutralizing antibodies (bnAbs) to coronaviruses and use molecular information gleaned on their epitopes to rationally design pan-coronavirus vaccines (*16-18*). Pan-coronavirus vaccines and

antibodies could be stockpiled ahead of the emergence of a new coronavirus and used

to rapidly contain the virus. BnAbs and pan-coronavirus vaccines that target more

conserved regions of the virus may also be more effective against antigenically variant

viruses, such as have been described for the variants of concern in the COVID-19

pandemic (*19-22*).

All HCoVs possess a surface envelope spike glycoprotein that mediates interaction with

host cell receptors and enables virus fusion (*4, 23*). SARS-CoV-2 (similar to SARS-CoV-

1) utilizes the receptor binding domain (RBD) in the S1 subunit of the spike protein to

engage human angiotensin converting enzyme 2 (hACE2) on host cells for cell entry and

infection (*23-27*). The SARS-CoV-2 spike glycoprotein is the primary target of neutralizing

antibodies (nAbs) (*28-31*). On the spike protein, the RBD is highly immunogenic and is

recognized by the majority of nAbs (*28, 32-43*), and thus is a major focus of current nAb-

based vaccine design efforts (*28, 44, 45*). However, due to sequence diversity, cross-

reactivity to the RBD region is limited, especially among emerging coronaviruses with

pandemic potential (*10-13*). The most potent nAbs in humans during natural infection are

typically raised to epitopes overlapping the ACE2 binding site (*32, 33, 42, 45, 46*). As the

rapid spread of the SARS-CoV-2 virus continues, these epitopes are coming under strong

immune selection pressure at the population level, leading to the selection of SARS-CoV-

2 neutralization escape variants (*19-22, 47-49*). The relevant mutations may result in

reduced effectiveness of vaccine-induced antibody responses in humans since such

responses also tend to target RBD epitopes overlapping the ACE2 binding site, and

because all currently approved vaccines are based on the wild-type virus. The most

striking example that illustrates the capability of the RBD to mutate without majorly

affecting the ability of the virus to engage host receptor is the variability of the RBD across

the two families of HCoVs: SARS-CoV-2/1 (β-HCoVs) and HCoV-NL63 (α-HCoV) (*23-27,*

50). These HCoVs possess divergent RBDs, but all use the ACE2 receptor for viral entry

suggesting that SARS-CoV-2, and potentially other emerging sarbecoviruses with human

pandemic potential, can tolerate changes in this domain with limited fitness cost.

Therefore, we believe that other sites on the spike protein should be explored as targets

of bnAbs.

We recently isolated a SARS-CoV-1/2 cross-neutralizing antibody from a COVID-19

donor, CC40.8, that exhibits broad cross-reactivity with human β-CoVs (*51*). Here, we

show that the CC40.8 bnAb targets an S2 stem-helix epitope, which is part of the

coronavirus fusion machinery. We first identified a long 25-mer S2 peptide from HCoV-

HKU1 that bound CC40.8 with high affinity and then determined the crystal structure of

CC40.8 with the SARS-CoV-2 S2 peptide. The S2 stem peptide adopts a largely helical

structure that is embedded in a groove between the heavy and light chain

complementarity determining regions (CDRs) of the antibody. Key epitope contact

residues were further validated, by alanine scanning, to be important for peptide binding

and for virus neutralization. These contact residues are largely conserved between β-

CoVs, consistent with the cross reactivity of CC40.8. In SARS-CoV-2 challenge models,

CC40.8 showed in vivo protective efficacy by reducing weight loss and lung tissue viral

titers. Although two recent studies have described S2-stem nAbs isolated from mice and

mice transgenic for human Ig (*52, 53*), CC40.8 represents a human HCoV S2-stem

directed bnAb isolated from natural infection (*51*) and may facilitate development of

antibody-based interventions and prophylactic pan-sarbecovirus and pan-β-coronavirus

vaccine strategies.

Results

CC40.8 binds a conserved peptide from the S2 region of β-coronaviruses.

We recently isolated a bnAb, CC40.8, from a 62-year-old SARS-CoV-2 convalescent

donor from peripheral blood mononuclear cell (PBMC) samples collected 32 days post-

infection (*51*). CC40.8 bnAb neutralizes SARS-CoV-1 and SARS-CoV-2 and exhibits

broad reactivity against β-coronaviruses, notably the endemic coronavirus HCoV-HKU1

(Fig. 1A and B) (*51*). Here, we observed that CC40.8 bnAb can effectively neutralize clade

1b and clade 1a ACE2 receptor-utilizing sarbecoviruses (Fig. 1A, fig. S1A). In addition,

the CC40.8 bnAb was consistently effective against the current SARS-CoV-2 variants of

concern (VOCs) (Fig. 1A, fig. S1A). The effectiveness of CC40.8 bnAb with SARS-CoV-

2 VOCs is consistent with a lack of mutations in the S2 stem helix region in the current

VOCs (*21*). To assess the cell-cell inhibition ability of CC40.8 bnAb, we conducted

experiments in HeLa cells expressing SARS-CoV-2 spike protein or hACE2 receptor. We

observed that CC40.8 bnAb can prevent cell-cell fusion of HeLa cells expressing SARS-

CoV-2 spike protein with HeLa cells expressing the hACE2 receptor (fig. S2).

Using negative-stain electron microscopy (ns-EM), we previously showed that the CC40.8

antibody targets the base of the S2 subunit on HCoV spike proteins, but epitope flexibility

precluded determination of a high-resolution structure (*51*). Here, we pursued epitope

identification, first by peptide mapping. Using HCoV-HKU1 S2 subunit overlapping

biotinylated peptides (15-residue long with a 10-residue overlap) for binding to CC40.8,

we identified that the stem-helix region in the S2 fusion domain contains the epitope (Fig.

1C, fig. S3). Then, through screening with peptides of various lengths that include the

epitope, we identified a 25-residue peptide that showed the strongest binding by biolayer

interferometry (BLI, Fig. 1D). The peptide corresponds to residues 1226-1250 from the

HCoV-HKU1 S2 sequence.

Next, we tested BLI binding of CC40.8 bnAb with peptides encompassing similar S2-

domain regions of other HCoVs. We observed that the antibody binds to the β- but not to

the α-HCoV S2-domain peptides (Fig. 1D). This pattern is consistent with the differential

binding of CC40.8 bnAb to different families of HCoV spike proteins (Fig. 1B) (*51*).

Sequence alignment of the S2 stem-helix domain region showed strong conservation

between SARS-CoV-1 and SARS-CoV-2 with more modest conservation across the

seasonal β-CoVs, consistent with cross-reactive binding patterns (Fig. 1E to G).

To determine whether CC40.8 bnAb affinity maturation was important for cross-reactive

binding or neutralization, we generated an inferred germline (iGL) version of CC40.8 with

corresponding antibody V-D-J germline genes (fig. S1B), as described previously (*54,*

55). Although the CC40.8 bnAb iGL Ab version retained binding to spike proteins and the

stem-helix peptides of β-CoVs, the neutralizing activity was lost against sarbecoviruses

(fig. S1, C to E), suggesting that affinity maturation is critical for neutralization, but cross-

reactive breadth is germline-encoded. Interestingly, the CC40.8 iGL bnAb showed binding

to MERS-CoV spike protein that CC40.8 bnAb fails to bind and exhibited some weak

polyreactivity (fig. S1, fig. S4), suggesting that naive B cells targeting this epitope may

begin with a broader reactivity to CoV spike proteins.

The epitope of CC40.8 bnAb was defined by the crystal structure of a peptide-

antibody complex.

To investigate the molecular nature of the CC40.8 bnAb epitope, we determined the 1.6

Å resolution crystal structure of the antibody Fab fragment with the SARS-CoV-2 25-mer

S2 peptide (Fig. 2A, table S1). The peptide adopts a largely helical structure that traverses

a wide hydrophobic groove formed between the heavy and light chains of the Fab (fig.

S5). The buried surface area on the peptide is about 1150 $Å^2$ (669 $Å^2$ conferred by the

heavy chain and 488 $Å^2$ by the light chain) and is largely contributed by hydrophobic

residue interactions at the paratope-epitope interface, although some hydrogen bonds

and salt bridges are contributed by CDRHs1 to 3, FRH1, and CDRLs1 to 3 (Fig. 2A to C,

fig. S6). Two peptide stretches of ^{1142}QPELD^{1146} and ^{1151}ELDKYF^{1156} and several nearby

residues, F^{1148} N^{1158}, H^{1159}, form the epitope of the bnAb (Fig. 2B and C). Notably,

hydrophobic residues in ^{1151}ELDKYF^{1156} of the stem region, as well as two upstream

residues, L^{1145} and F^{1148}, form the core of the epitope that interacts with a hydrophobic

groove in the antibody lined by heavy chain residues (V33, Y35, W47, Y56, Y58, M96

and V101) and light chain residues (Y32, Y34, L46 and Y49) (Fig. 2C and fig. S6).

Antibody germline and mutated residues both contribute to epitope recognition (fig. S6).

Consistent with our findings, two recent independent studies have shown that

heterologous CoV spike protein immunizations in mice or mice transgenic for the human

Ig locus can induce cross-reactive serum neutralizing antibodies that target the conserved

S2 spike epitope similar to the stem-helix epitope identified in our study, and some

isolated mAbs also show broad reactivity to coronavirus spike proteins (*52, 53*).

The residues important for CC40.8 interaction with virus were also investigated by alanine

scanning mutagenesis of SARS-CoV-2 and HCoV-HKU1 peptides and spike protein by

antibody binding and by neutralization of SARS-CoV-2 spike protein mutants (Fig. 2D and

E, fig. S7). The contact residues determined by crystallography were also found to be

important for peptide binding and neutralization with the S2 helical residues, L^{1145}, E^{1151},

F^{1148} and Y^{1155}, being the most critical (fig. S7). We noted some differences in CC40.8

dependence on S2 residue substitutions for virus neutralization and spike protein binding,

which may reflect differences in conformation or glycosylation between recombinant and

native membrane-associated spike protein (*56, 57*). The conservation of those residues

identified by crystallography and alanine scanning as most critical for interaction of

CC40.8 with virus is high (Fig. 2F) across human β-coronaviruses and related

sarbecoviruses that infect various animal species, thus providing a structural basis for

broad cross-reactivity of the antibody.

The CC40.8 epitope region houses an N-linked glycan (N^{1158}) that is highly conserved

across coronaviruses and may restrict access to this bnAb epitope. To investigate, we

substituted the T^{1160} residue on SARS-CoV-2 virus spike protein with an alanine residue

to eliminate the AsnHisThr (NHT) N-linked glycan attachment site. A modest increase in

neutralization sensitivity of the T1160A variant relative to wild-type virus was observed

(Fig. 2E, fig. S7), suggesting that any steric obstruction of the epitope by the N1158 glycan

is limited.

The CC40.8 bnAb epitope appears to be only partially exposed on the pre-fusion HCoV

spike protein (fig. S8). Previously, a SARS-CoV-2 S2 stem-targeting neutralizing

antibody, B6, was isolated from a mouse immunized with spike protein (*52*) . Here we

compared the structures of antibodies CC40.8 and B6 (fig. S9, A to E). Both antibodies

target a similar epitope on the SARS-CoV-2 spike protein, the conserved S2 stem helix

region, but with different angles of approach; a longer peptide was visualized as the

epitope for CC40.8. The post-fusion spike protein requires a large conformational change

in the S2 stem region, and the superimposed CC40.8 (fig. S8C) and B6 (*52*) would clash

with the post-fusion conformation. On the other hand, if bound to a spike protein in the

pre-fusion state, both antibodies would clash with the adjacent protomer (fig. S9, D and

E), suggesting a possible neutralization mechanism where the antibodies may induce

disruption of the S2 stem 3-helix bundle, and bind to an intermediate state of the spike

protein (fig. S8C). This hypothesis is further supported by comparing the binding of

CC40.8 to S2P and HexaPro (S6P). The two proline mutations (S2P) were introduced to

stabilize the SARS-CoV-2 S trimer in its pre-fusion state (*26, 58*), whereas additional

proline substitutions to the HexaPro or S6P construct further stabilized the SARS-CoV-2

spike trimer (*59*). Here we show that the S6P-stabilized version exhibited much weaker

binding to CC40.8 compared to S2P (fig. S9F), further suggesting that destabilization or

partial disruption of the pre-fusion S trimer is a possible explanation for neutralization by

S2 stem-targeting antibodies, such as CC40.8 or B6. Our previous EM study of the

complex of HCoV-HKU1 S (S2P) and CC40.8 Fab [Fig. 5D in ref (*51*)] showed high

flexibility of the epitope and multiple antibody approach angles, which also suggested

disruption of the 3-helix bundle and induction of flexibility in the S2 stem region.

245
CC40.8 antibody protects against weight loss and reduces viral burden in SARS-

CoV-2 challenge in vivo.

To determine the in vivo efficacy of CC40.8, we conducted passive antibody transfer

followed by SARS-CoV-2 challenge in human ACE2 (hACE2) mice and in Syrian

hamsters. CC40.8 mAb at 4 different doses (300μg, 100μg, 50μg and 10μg per animal)

was intra-peritoneally (i.p.) administered into groups (6 animals per group) of hACE2 mice

(Fig. 3A) (*60*). An RBD nAb (CC12.1; 300 μg/animal) positive control and a Zika-specific

antibody (SMZAb1; 300 μg/animal) negative control were administered i.p. into control

animal groups. All CC40.8- and control mAb-treated animals were challenged with SARS-

CoV-2 (USA-WA1/2020) by intranasal (i.n.) administration of a virus dose of 2×10^4

plaque forming units (PFU), 12 hours post-antibody infusion (Fig. 3A). The animals were

weighed daily to monitor weight changes, as an indicator of disease due to infection and

serum samples were collected to determine the transferred human antibody

concentrations (Fig. S10). Animals were euthanized at day 5 and lung tissues were

collected to determine the SARS-CoV-2 titers by quantitative polymerase chain reaction

(qPCR) and by plaque assays. The CC40.8 bnAb-treated animals showed significantly

reduced weight loss as compared to the SMZAb1-treated control group animals

($P<0.0001$, Fig. 3B, fig. S10), suggesting a protective role for CC40.8. Remarkably, the

animals treated with the lowest dose of CC40.8 bnAb (10 μg/animal) also showed

significant protection against weight loss (P = 0.0005, Fig. 3B, fig. S10). As expected, the

positive control RBD nAb, CC12.1 significantly protected against weight loss (P<0.0001,

Fig. 3B, fig. S10). Consistent with these results, SARS-CoV-2 specific viral RNA copies

and viral titers in day 5 lung tissues were significantly reduced in the CC40.8-treated

animals compared to the SMZAb1 control group animals (P<0.0001, fig. 3C and D).

We also investigated the protective efficacy of CC40.8 mAb by intra-peritoneally (i.p.)

administering into a group of 5 Syrian hamsters (at 2 mg per animal) and subsequently

challenging with SARS-CoV-2 (USA-WA1/2020 dose of 1×10^6 PFU) (fig. S11). SMZAb1

Zika mAb was used as a control. Consistent with hACE2 mouse experiments, the CC40.8

bnAb-treated animals showed substantially reduced weight loss and reduced SARS-CoV-

2 titers in day 5 lung tissues demonstrating its protective role (fig. S11). Altogether, the

findings reveal that CC40.8, despite relatively low in vitro neutralization potency, shows

a substantial degree of protective efficacy against SARS-CoV-2 infection in vivo.

Consistent with these results, a recently isolated S2 stem bnAb, S2P6, has also been

shown to protect against SARS-CoV-2 challenge despite relatively low neutralization

potency (*61*) . Furthermore, this phenomenon of a surprisingly high degree of protection

afforded by antibodies directed to epitopes close to the spike protein membrane and part

of the fusion machinery has been described earlier for HIV (*62*).

The conserved stem-helix epitope defined by bnAb CC40.8 is infrequently targeted

following SARS-CoV-2 infection.

To investigate how frequently the CC40.8 epitope is targeted following SARS-CoV-2

infection, we tested the binding of serum samples from 60 COVID-19 convalescent

donors to 25-mer peptides of HCoVs corresponding to the stem-helix bnAb epitope. We

observed that 6 of 60 (10%) individuals exhibited some degree of cross-reactive binding

with β-HCoV S2 stem peptides (Fig. 4A). We further tested the binding of cross-reactive

serum samples with SARS-CoV-2 S2 stem peptide alanine scan variants spanning the

CC40.8 epitope and observed the presence of CC40.8-like epitope-targeting antibodies

(Fig. 4B). The binding of cross-reactive serum Abs revealed dependence on five common

stem helix residues including a conserved hydrophobic core formed by F^{1148}, L^{1152}, Y^{1155}

and F^{1156} (Fig. 4B). To determine the contribution of S2-stem directed antibodies in overall

SARS-CoV-2 neutralization by serum Abs in cross-reactive COVID-19 donors, we

conducted competition experiments with the SARS-CoV-2 S2 stem-helix peptide. Peptide

competition showed no or minimal effects on the SARS-CoV-2 neutralization (Fig. 4C and

D), suggesting that stem-helix targeting cross-reactive nAbs minimally contribute to the

overall polyclonal serum neutralization in these COVID-19 convalescent donors.

Discussion

The development of effective pan-coronavirus vaccine strategies that can mitigate future outbreaks from new emerging coronaviruses is important (*16, 18*). Two major challenges are the identification of broadly neutralizing antibody (bnAb) targets on CoV spike proteins and the development of vaccine strategies that can reproducibly elicit durable and protective pan-CoV bnAbs. The approach of identifying conserved bnAb surface protein targets by isolating bnAbs from natural infection and utilizing their molecular information in structure-guided immunogen design has greatly contributed to the development of vaccine strategies against a range of complex pathogen surfaces (*63-70*).

The spike S1 subunit shows considerable variation on HCoVs, whereas the S2 subunit is relatively more conserved, especially across the β-HCoVs, and appears to be promising for developing pan-CoV bnAb vaccine strategies. Accordingly, we recently isolated a SARS-CoV-1/2 cross-neutralizing Ab, CC40.8, that exhibits broad reactivity with human β-CoVs (*51*). In this study, using epitope mapping and structural studies, we determined the spike epitope recognized by CC40.8. The epitope is located in the S2 stem-helix region, which is conserved across β-coronaviruses and may thus serve as a promising target for pan-β-coronavirus vaccine strategies. The epitope is highly enriched in hydrophobic residues as well as some charged residues. The bnAbs targeting this region may neutralize by sterically interfering with the fusion machinery (*52, 53*), suggesting a potential target for fusion inhibitors (*71-73*). CC40.8 bnAb represents a human bnAb directed to the HCoV S2 stem helix (*51*). Two more S2 stem human bnAbs, S2P6 and

CV3-25, have also been reported recently (*61, 74*) that target a similar S2 stem epitope

region. Knowledge from these nAbs will be important for developing broad vaccine

strategies for β-coronaviruses.

We noted that cross-reactive antibodies directed to the CC40.8 S2 stem-helix epitope are

much less frequently elicited in human coronavirus natural infections as compared to

strain-specific neutralizing antibody responses (*28*). However, a few recent studies using

more sensitive antibody detection assays have suggested a higher prevalence of

polyclonal stem-helix region-directed antibodies in COVID-19 donors and their possible

association with reduced disease severity (*75-77*). Regardless, the small subset of

individuals in our sample cohort that do make cross-reactive Abs, seem to exhibit broad

reactivity to human β-coronaviruses, which is promising for pan-β-coronavirus vaccine

strategies. In principle, the paucity of these cross-reactive antibodies could be due to poor

accessibility of the S2 stem-helix epitope on the native spike protein relative to other

epitopes, low frequency of bnAb-encoding B cell precursors in humans, or complex

secondary B cell maturation pathways. Low epitope accessibility is clearly a potential

contributor to low immunogenicity. Low precursor frequency seems unlikely, at least for

CC40.8-like antibodies given that this antibody uses a common VH gene segment

(IGHV3-23) and CDRH3 length (10 amino acids) (*78*). Analysis of CC40.8 antibody

variable regions by the Armadillo tool (*79*) revealed the presence of several improbable

somatic mutations that are predicted to contribute to difficulty in elicitation of CC40.8-like

antibodies. Thus, isolation of multiple cross-reactive pan-CoV S2 stem bnAb lineages,

understanding their maturation pathways, and identifying common antibody framework

motifs, are likely to be important for rational vaccine design approaches (*80*).

Encouragingly, two recent studies have described a similar CoV S2 domain bnAb epitope

being targeted by cross-reactive mAbs isolated from heterologous CoV spike protein

immunizations in mice and mice transgenic for the human Ig locus (*52, 53*). These data

suggest that such bnAbs could be induced by both immunization with designed vaccines

as well as coronavirus infection in humans. Nonetheless, it would need to be ascertained

how many sequential immunizations would be needed to broaden the breadth of these

nAb responses.

Interestingly, despite relatively low neutralization potency, CC40.8 showed robust in vivo

protective efficacy against SARS-CoV-2 challenge. The findings illustrate that extra-

neutralizing effector functions of S2 stem bnAbs may be important and need to be

investigated to determine their role in viral suppression or clearance. Indeed, recent

studies have revealed that cross-reactive antibodies to endemic CoVs could serve as a

marker for survival after severe disease (*81, 82*) or protection against COVID-19 (*83*).

Although we show CC40.8 S2 stem-helix bnAb confers in vivo protection against SARS-

CoV-2 infection, our study has limitations. First, our CC40.8 protection studies were

focused on SARS-CoV-2 infection and testing the protective efficacy of CC40.8 against

a broad range of betacoronaviruses will be important. Second, we showed CC40.8

protects against SARS-CoV-2 in two small animal models and investigating its protective

efficacy in non-human primate models is also eventually desirable. Third, CC40.8 bnAb

showed remarkable protection even at very low antibody levels despite its low

neutralization potency, hence warranting future studies to investigate the role of other

factors such as antibody effector function that may contribute to CC40.8-mediated

protection against coronaviruses.

Overall, we describe a cross-neutralizing human bnAb epitope on β-CoVs and provide

molecular details that help explain its broad reactivity. The identification of this conserved

epitope in the coronavirus spike protein should facilitate bnAb-epitope based vaccine

development and antibody-based intervention strategies not only to SARS-CoV-2, but

against existing human coronaviruses and other coronaviruses that could emerge with

pandemic potential.

Materials and Methods

Study Design

The objective of the study was to evaluate a previously discovered SARS-CoV-2 spike

protein stem-helix antibody, CC40.8, for binding to and neutralization of diverse

sarbecoviruses and SARS-CoV-2 Variants of Concern, structurally define its epitope site

and test its protective efficacy. For in vitro binding and neutralization studies, CC40.8 and

control mAbs were tested in duplicate and experiments were repeated independently for

rigor and reproducibility. We did not use any statistical methods to predetermine sample

sizes for the animal studies. All hACE2-trangenic mouse or hamster experiments used 5

or 6 animals per group. A positive and/or a negative control mAb-treated animal group

was included in the in vivo SARS-CoV-2 challenge experiments. Male and female, age

matched (8-week old) animals were randomly assigned in CC40.8 bnAb-treated or control

mAb-treated animal groups for the SARS-CoV-2 challenge studies. All immunological and

virological measurements were performed blinded. Animals were euthanized at day five

post infection to measure weight loss and lung viral load. The serum antibody titers of the

passively transferred antibody were determined daily for the SARS-CoV-2 challenge

experiment in the hACE2 mice. No datapoints were excluded as outliers in any

experiment.

Human cohort information

Plasma from convalescent COVID-19 donors were kindly provided through the "Collection

of Biospecimens from Persons Under Investigation for 2019-Novel Coronavirus Infection

to Understand Viral Shedding and Immune Response Study" UCSD IRB# 200236.

Samples were collected based on COVID-19 diagnosis regardless of gender, race,

ethnicity, disease severity, or other medical conditions. The gender for individuals was

evenly distributed across the human cohort. All human donors were assessed for medical

decision-making capacity using a standardized, approved assessment, and voluntarily

gave informed consent prior to being enrolled in the study. The summary of the

demographic information of the COVID-19 donors is listed in table S2.

Pseudovirus production and generation of mutant spike proteins

Under biosafety level 2 and 3 conditions, MLV-gag/pol (Addgene #14887) and pCMV-

Fluc (Addgene #170575) were co-transfected into HEK293T cells along with plasmids

encoding full-length or variously truncated spike proteins from SARS-CoV-1, WIV1,

SHC014, PANG17, MERS-CoV and SARS-CoV-2 (SARS-CoV-2 variants of concern

(B.1.1.7 (alpha), B.1.351 (beta), P.1 (gamma) and B.1.617.2 (delta)) using Lipofectamine

2000 (Thermo Fisher Scientific cat.# 11668019) to produce single-round of infection

competent pseudo-viruses. The media was changed by fresh Dulbecco's Modified Eagle

Medium (DMEM) with 10% heat-inactivated FBS, 4mM L-Glutamine and 1% P/S 16 hours

post transfection. The supernatant containing MLV-pseudotyped viral particles was

collected 48 hours post transfection, aliquoted and frozen at -80 °C for the neutralization

assay. Amino-acid point mutations in SARS-CoV-2 spike protein-encoding plasmids were

made by using site-directed mutagenesis kit (New England Biolabs cat.# E0554S)

according to the manufacturer's instructions. All the mutations were verified by DNA

sequence analysis (Eton Bioscience).

Neutralization assay

Pseudotyped viral neutralization assay was performed as previously described with

minor modifications (Modified from TZM-bl assay protocol (*84*)). In sterile 96-well half-

area plates (Corning cat.# 3688), 25 µl of virus was immediately mixed with 25 µl of three-

fold serially diluted monoclonal antibodies (mAb) (starting concentration of 300 µg/ml) or

serially diluted plasma from COVID-19 donors and incubated for one hour at 37°C to allow

for antibody neutralization of the pseudotyped virus. Synthesized peptides were optionally

added in the mixture for testing inhibition of neutralization. 10,000 HeLa-hACE2 cells (as

previously generated (*33*)) per well (in 50 µl of media containing 20 µg/ml Dextran) were

directly added to the antibody virus mixture. Plates were incubated at 37°C for 42 to 48

hours. Following the infection, HeLa-hACE2 cells were lysed using 1x luciferase lysis

buffer (25mM Gly-Gly pH 7.8, 15mM MgSO4, 4mM EGTA, 1% Triton X-100). Luciferase

intensity was then read on a Luminometer with luciferase substrate according to the

manufacturer's instructions (Promega cat.# E2620). Percentage of neutralization was

calculated using the following equation: 100 X (1 – (mean fluorescent intensity (MFI) of

sample – average MFI of background) / average of MFI of probe alone – average MFI of

background)). Fifty percent maximal inhibitory concentrations (IC_{50}), the concentrations

required to inhibit infection by 50% compared to the controls, were calculated using the

dose-response-inhibition model with 5-parameter Hill slope equation in GraphPad Prism

7 (GraphPad Software)

Flow cytometry based Cellular-ELISA (CELISA) binding

Binding of monoclonal antibody to various human coronavirus (HCoV) spike proteins

expressed on the surface of HEK293T cells was determined by flow cytometry, as

described previously (*85*). Briefly, HEK293T cells were transfected with different plasmids

encoding full-length HCoV spike proteins and were incubated for 36 to 48 hours at 37°C.

Post incubation cells were trypsinized to prepare a single cell suspension and were

distributed into 96-well plates. Monoclonal antibodies were prepared as 5-fold serial

titrations in FACS buffer (1x phosphate-buffered saline (PBS), 2% fetal bovine serum

(FBS), 1 mM EDTA), starting at 10 µg/ml, 6 dilutions. 50 µl/well of the diluted samples

were added into the cells and incubated on ice for 1 hour. The plates were washed twice

in FACS buffer and stained with 50 µl/well of 1:200 dilution of R-phycoerythrin (PE)-

conjugated mouse anti-human IgG Fc antibody (SouthernBiotech cat.# 9040-09) and

1:1000 dilution of Zombie-NIR viability dye (BioLegend cat.# 423105) on ice in dark for

45 minutes. After another two washes, stained cells were analyzed using flow cytometry

(BD Lyrics cytometers), and the binding data were generated by calculating the percent

(%) PE-positive cells for antigen binding using FlowJo 10 software.

Expression and purification of HCoV spike proteins and SARS-CoV-2 spike protein

mutants

To express the soluble spike ectodomain proteins, the HCoV spike protein encoding

plasmids were transfected into FreeStyle293-F cells (Thermo Fisher Scientific cat.#

R79007). For general production, 350 µg of plasmids were transfected into 1L

FreeStyle293-F cells at the density of 1 million cells per mL. 350 µg plasmids in 16 ml

Opti-MEM™ (Thermo Fisher Scientific cat.# 31985070) were filtered and mixed with 1.8

mL 40K PEI (1mg/mL) in 16 ml Opti-MEM™. After gently mixing the two components, the

combined solution rested at room temperature for 30 minutes and was poured into 1L

FreeStyle293-F cell culture. The cell cultures were centrifuged at 2500xg for 15 minutes

on day 4 after transfection, and the supernatants were filtered through the 0.22 µm

membrane. The His-tagged proteins were purified with the HisPur Ni-NTA Resin (Thermo

Fisher Scientific cat.# 88221). Each column was washed with at least 3 bed volumes of

wash buffer (25 mM Imidazole, pH 7.4), followed by elution with 25 ml of the elution buffer

(250 mM Imidazole, pH 7.4) at slow gravity speed (about 4 seconds per drop). The eluates

were buffer exchanged into PBS by using Amicon tubes, and the proteins were

concentrated afterwards. The proteins were further purified by size-exclusion

chromatography using a Superdex 200 Increase 10/300 GL column (GE Healthcare cat.#

GE28-9909-44). The selected fractions were pooled and concentrated again for further

use.

BioLayer Interferometry (BLI) binding

The determination of monoclonal antibody binding with spike proteins or selected

peptides was conducted in an Octet K2 system (ForteBio). The anti-human IgG Fc

capture (AHC) biosensors (ForteBio cat.# 18-5063) were used to capture IgG first for 60

seconds. After providing baseline in Octet buffer for another 60 seconds, the sensors

were transferred into HCoV spike proteins at various concentrations for 120 seconds for

association, and into Octet buffer for disassociation for 240 seconds. Alternatively, the

hydrated streptavidin biosensors (ForteBio cat.# 18-5020) first captured the N-terminal

biotinylated peptides diluted in Octet buffer (PBS plus 0.1% Tween-20) for 60 seconds,

then transferred into Octet buffer for 60 seconds to remove unbound peptide and provide

the baseline. Then the sensors were immersed in diluted monoclonal antibody IgG for

120 seconds to provide association signal, followed by transferring into Octet buffer to

test for disassociation signal for 240 seconds. The data generated was analyzed using

the ForteBio Data Analysis software for correction, and the kinetic curves were fit to 1:1

binding mode. Note that the IgG: spike protomer binding can be a mixed population of 2:1

and 1:1, such that the term 'apparent affinity' dissociation constants (K_D^{App}) are shown to

reflect the binding affinity between IgGs and spike trimers tested.

HEp2 epithelial cell polyreactive assay

Reactivity to human epithelial type 2 (HEp2) cells was determined by indirect

immunofluorescence on HEp2 slides (Hemagen, cat.# 902360) according to

manufacturer's instructions. Briefly, monoclonal antibody was diluted at 50 μg/mL in PBS

and then incubated onto immobilized HEp2 slides for 30 minutes at room temperature.

After washing 3 times with PBS buffer, one drop of fluorescein isothiocyanate (FITC)-

conjugated goat anti-human IgG was added onto each well and incubated in the dark for

30 minutes at room temperature. After washing, the coverslip was added to HEp2 slide

with glycerol and the slide was photographed on a Nikon fluorescence microscope to

detect FITC signal.

Polyspecificity reagent (PSR) ELISA

Solubilized CHO cell membrane protein (SMP) was coated onto 96-well half-area high-

binding ELISA plates (Corning, cat.# 3690) overnight at 4°C. After washing with PBS plus

0.05% Tween-20 (PBST), plates were blocked with 3% bovine serum albumin (BSA) for

2 hours at 37°C. Antibody samples were diluted at 10 μg/mL in 1% BSA with 3-fold serial

dilution and then added in plates to incubate for 1 hour at 37°C. After 3 thorough washes

with PBST, alkaline phosphatase-conjugated goat anti-human IgG Fc secondary antibody

(Jackson ImmunoResearch, cat.# 109-055-008) was added to the plate and incubated

for 1 hour at 37°C. After a final wash, phosphatase substrate (Sigma-Aldrich, cat.# S0942-

200TAB) was added into each well. Absorption was measured at 405 nm.

Peptide scanning by ELISA binding

N-terminal biotinylated overlapping peptides corresponding to the complete sequence of

HCoV-HKU1 S2 subunit (residue number range: 761-1355 (GenBank: AAT98580.1) were

synthesized at A&A Labs (Synthetic Biomolecules). Each peptide was 15 residue long

with a 10 amino acid overlap. For ELISA binding, 96-well half-area plates (Corning cat. #

3690) were coated overnight at 4°C with 2 µg/ml of streptavidin in PBS. Plates were

washed 3 times with PBST and blocked with 3% (wt/vol) BSA in PBS for 1 hour. After

removal of the blocking buffer, the plates were incubated with peptides in 1% BSA plus

PBST for 1.5 hours at room temperature. After a washing step, monoclonal antibody or

serum samples diluted in 1% BSA/PBST were added into each well and incubated for 1.5

hours. DEN3 human antibody was used as a negative control. After the washes, a

secondary antibody conjugated with alkaline phosphatase-conjugated goat anti-human

IgG Fc secondary antibody (Jackson ImmunoResearch, cat.# 109-055-008) diluted

1:1000 in 1% BSA/PBST, was added to each well and incubated for 1 hour. The plates

were then washed and developed using alkaline phosphatase substrate pNPP tablets

(Sigma-Aldrich, cat.# S0942-200TAB) dissolved in stain buffer. The absorbance was

recorded at an optical density of 405 nm (OD405) using a VersaMax microplate reader

(Molecular Devices), where data were collected using SoftMax software version 5.4.

Cell-cell fusion inhibition assay

HeLa stable cell lines were generated through transduction of lentivirus carrying genes

encoding either human ACE2 (hACE2) and enhanced green fluorescent protein (EGFP)

or nuclear localization signal (NLS)-mCherry and SARS-CoV-2 spike protein. The pBOB

construct carrying these genes was co-transfected into HEK293T cells along with

lentiviral packaging plasmids pMDL, pREV, and pVSV-G (Addgene #12251, #12253,

#8454) by Lipofectamine 2000 (Thermo Fisher Scientific, cat.# 11668019) according to

the manufacturer's instructions. Supernatants were collected 48 hours after transfection,

then were transduced to pre-seeded HeLa cells. 12 hours after transduction, stable cell

lines were collected, scaled up and stored for cell-cell fusion assay. 10,000 NLS-

mCherry+ HeLa cells expressing SARS-COV-2 spike protein were seeded into 96-well

half-well plates on the day before the assay. The culture medium was removed by

aspiration before the assay. 50 µl of 50 µg/ml CC40.8 and DEN3 mAbs diluted in DMEM

with 10% heat-inactivated FBS, 4mM L-Glutamine and 1% P/S were then added to the

pre-seeded cells and incubated for 1 hour in an incubator. 50 µl of 10,000 EGFP+hACE2+

HeLa cells were added to the plates and incubated for 2 hours before taking images under

the microscope.

Sequence alignments of coronavirus spike stem regions

The spike sequences of SARS-CoV-2, SARS-CoV-1, RaTG13, SHC014, Rs4081,

Pang17, RmYN02, Rf1, WIV1, Yun11, BM48-31, BtKY72, HCoV-HKU1, HCoV-OC43,

MERS-CoV, MHV, HCoV-229E and HCoV-NL63 were downloaded from the GenBank

and aligned against the SARS-CoV-2 reference sequence using BioEdit

(http://www.mbio.ncsu.edu/bioedit/bioedit.html).

Expression and purification of CC40.8 Fab

To generate Fab, CC40.8 IgG was digested by Papain (Sigma-Aldrich cat.# P3125) for 4

hours at 37 °C, then was incubated with Protein-A beads at 4 °C for 2 hours to remove

the Fc fragments. CC40.8 Fab was concentrated afterwards and further purified by size-

exclusion chromatography using a Superdex 200 Increase 10/300 GL column (GE

Healthcare cat.# GE28-9909-44). The selected fractions were pooled and concentrated

again for further use.

Crystallization and structural determination

A mixture of 9 mg/ml of CC40.8 Fab and 10× (molar ratio) SARS-CoV-2 stem peptide

was screened for crystallization using the 384 conditions of the JCSG Core Suite (Qiagen)

on our robotic CrystalMation system (Rigaku) at Scripps Research. Crystallization trials

were set-up by the vapor diffusion method in sitting drops containing 0.1 µl of protein and

0.1 µl of reservoir solution. Optimized crystals were then grown in drops containing 0.1 M

sodium acetate buffer at pH 4.26, 0.2 M ammonium sulfate, and 28% (w/v) polyethylene

glycol monomethyl ether 2000 at 20°C. Crystals appeared on day 7, were harvested on

day 15 by soaking in reservoir solution supplemented with 20% (v/v) glycerol, and then

flash cooled and stored in liquid nitrogen until data collection. Diffraction data were

collected at cryogenic temperature (100 K) at Stanford Synchrotron Radiation Lightsource

(SSRL) on the Scripps/Stanford beamline 12-1, with a beam wavelength of 0.97946 Å,

and processed with HKL2000 (*86*). Structures were solved by molecular replacement

using PHASER (*87*). A model of CC40.8 was generated by Repertoire Builder

(https://sysimm.ifrec.osaka-u.ac.jp/rep_builder/) (*88*). Iterative model building and

refinement were carried out in COOT (*89*) and PHENIX (*90*), respectively. Epitope and

paratope residues, as well as their interactions, were identified by accessing PISA at the

European Bioinformatics Institute (http://www.ebi.ac.uk/pdbe/prot_int/pistart.html) (*91*).

Animal Study

8-week old transgenic hACE2 mice were given an i.p. antibody injections 12 hours pre-infection. Mice were infected through intranasal installation of 2×10^4 total plaque-forming units (PFU) per animal of SARS-CoV-2 (USA-WA1/2020) in 25 µL of DMEM. Mice were bled on days 1, 2, 3, and 5 for serum antibody detection and weighed for the duration of the study. At day 5 post-infection, animals were euthanized, and lungs were harvested for quantitative polymerase chain reaction (qPCR) viral titer analysis and plaque live virus analysis. Similar experimental procedures were conducted for the protection study in 8-week old Syrian hamsters except that a higher SARS-CoV-2 (USA-WA1/2020) challenge dose (10^6 total PFU per animal) was used. The research protocol was approved and performed in accordance with Scripps Research IACUC Protocol #20-0003

Antibody detection in hACE2 serum samples by ELISA

Serum samples were obtained at day 1, 2, 3, and 5 to quantify mAb titers. Unconjugated F(ab')$_2$ fragment of goat anti-human F(ab')$_2$ fragment (Jackson ImmuoResearch cat.# 109-006-097) was coated to the ELISA plates overnight, then washed by PBS plus 1% Tween-20 three times. After being blocked by 3% BSA for 2 hours at 37°C, mouse serum dilution series and CC40.8 mAb dilution series for a standard curve were applied to the plates and reacted for 1 hour at 37°C. After three thorough washes with PBS plus 1% Tween-20, alkaline phosphatase-conjugated goat anti-human IgG Fc secondary antibody (Jackson ImmunoResearch, cat.# 109-055-008) was added to the plates before washing

3 times with PBS plus 1% Tween-20 and AP substrate applied for detection. The plates

were read at 405nm and data were analyzed by CurveExpert.

SARS-CoV-2 RNA Quantification

Viral RNA was isolated from lung tissue and subsequently amplified and quantified in a

reverse transcription (RT)-qPCR reaction. Lung tissue was extracted at day 5 post

infection and placed in 1 mL of trIzol reagent (Invitrogen). The samples were then

homogenized using a Bead Ruptor 12 (Omni International). Tissue homogenates were

then spun down and the supernatant was added to an RNA purification column (Qiagen).

Purified RNA was eluted in 60 µL of DNase-, RNase-, endotoxin-free molecular biology

grade water (Millipore). RNA was then subjected to reverse transcription and quantitative

PCR using the CDC's N1 (nucleocapsid) primer sets (Forward 5'-GAC CCC AAA ATC

AGC GAA AT-3'; Reverse 5'-TCT GGT TAC TGC CAG TTG AATCTG-3') and a

fluorescently labeled (FAM) probe (5'-FAM-ACC CCG CAT TAC GTT TGGTGG ACC-

BHQ1-3') (Integrated DNATechnologies) on a BioRad CFX96 Real-Time instrument. For

quantification, a standard curve was generated by diluting 2.5×10^6 PFU RNA equivalents

of SARS-CoV-2. Every run utilized eleven 5-fold serial dilutions of the standard. SARS-

CoV-2-negative mouse lung RNA and no templates were both included as negative

controls for the extraction step as well as the qPCR reaction.

Viral load measurements

SARS-CoV-2 titers were measured by homogenizing organs in DMEM plus 2% fetal calf

serum using 100 µm cell strainers (Myriad cat.# 2825-8367). Homogenized organs were

titrated 1:10 over six steps and layered over Vero-E6 cells. After 1 hour of incubation at

37°C, a 1% methylcellulose in DMEM overlay was added, and the cells were incubated

for 3 days at 37°C. Cells were fixed with 4% paraformaldehyde and plaques were counted

by crystal violet staining.

Statistical Analysis

Statistical analysis was performed using Graph Pad Prism 8 for Mac (Graph Pad

Software). Groups of data were compared using the Kruskal-Wallis non-parametric test.

Dunnett's multiple comparisons test were also performed between experimental groups.

Data were considered statistically significant at $p < 0.05$.

Supplementary Materials

Fig S1 to S11

Table S1 and S2

Data File S1

References

1. C. Huang *et al.*, Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* **395**, 497-506 (2020).
2. Z. Wu, J. M. McGoogan, Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72314 Cases From the Chinese Center for Disease Control and Prevention. *JAMA*, (2020).
3. P. Zhou *et al.*, A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* **579**, 270-273 (2020).
4. J. Cui, F. Li, Z. L. Shi, Origin and evolution of pathogenic coronaviruses. *Nat Rev Microbiol* **17**, 181-192 (2019).
5. T. S. Fung, D. X. Liu, Human Coronavirus: Host-Pathogen Interaction. *Annu Rev Microbiol* **73**, 529-557 (2019).
6. C. Drosten *et al.*, Identification of a novel coronavirus in patients with severe acute respiratory syndrome. *The New England journal of medicine* **348**, 1967-1976 (2003).
7. A. M. Zaki, S. van Boheemen, T. M. Bestebroer, A. D. Osterhaus, R. A. Fouchier, Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia. *The New England journal of medicine* **367**, 1814-1820 (2012).
8. E. Petersen *et al.*, Comparing SARS-CoV-2 with SARS-CoV and influenza pandemics. *Lancet Infect Dis* **20**, e238-e244 (2020).
9. G. Meyerowitz-Katz, L. Merone, A systematic review and meta-analysis of published research data on COVID-19 infection fatality rates. *Int J Infect Dis* **101**, 138-148 (2020).
10. N. Wang *et al.*, Serological Evidence of Bat SARS-Related Coronavirus Infection in Humans, China. *Virol Sin* **33**, 104-107 (2018).
11. V. D. Menachery *et al.*, SARS-like WIV1-CoV poised for human emergence. *Proceedings of the National Academy of Sciences of the United States of America* **113**, 3048-3053 (2016).
12. V. D. Menachery *et al.*, A SARS-like cluster of circulating bat coronaviruses shows potential for human emergence. *Nature medicine* **21**, 1508-1513 (2015).
13. M. Letko, A. Marzi, V. Munster, Functional assessment of cell entry and receptor usage for SARS-CoV-2 and other lineage B betacoronaviruses. *Nat Microbiol* **5**, 562-569 (2020).
14. S. J. Anthony *et al.*, Further Evidence for Bats as the Evolutionary Source of Middle East Respiratory Syndrome Coronavirus. *mBio* **8**, (2017).
15. X. Y. Ge *et al.*, Isolation and characterization of a bat SARS-like coronavirus that uses the ACE2 receptor. *Nature* **503**, 535-538 (2013).
16. D. R. Burton, E. J. Topol, Variant-proof vaccines - invest now for the next pandemic. *Nature* **590**, 386-388 (2021).
17. D. R. Burton, L. M. Walker, Rational Vaccine Design in the Time of COVID-19. *Cell host & microbe* **27**, 695-698 (2020).
18. W. C. Koff, S. F. Berkley, A universal coronavirus vaccine. *Science* **371**, 759 (2021).
19. J. R. Mascola, B. S. Graham, A. S. Fauci, SARS-CoV-2 Viral Variants-Tackling a Moving Target. *JAMA*, (2021).
20. P. Wang *et al.*, Antibody Resistance of SARS-CoV-2 Variants B.1.351 and B.1.1.7. *Nature*, (2021).
21. W. T. Harvey *et al.*, SARS-CoV-2 variants, spike mutations and immune escape. *Nat Rev Microbiol* **19**, 409-424 (2021).
22. C. K. Wibmer *et al.*, SARS-CoV-2 501Y.V2 escapes neutralization by South African COVID-19 donor plasma. *Nature medicine* **27**, 622-625 (2021).
23. F. Li, Structure, Function, and Evolution of Coronavirus Spike Proteins. *Annu Rev Virol* **3**, 237-261 (2016).

713   24.   F. Li, W. Li, M. Farzan, S. C. Harrison, Structure of SARS coronavirus spike receptor-
714   binding domain complexed with receptor. *Science* **309**, 1864-1868 (2005).
715   25.   A. C. Walls *et al.*, Structure, Function, and Antigenicity of the SARS-CoV-2 Spike
716   Glycoprotein. *Cell* **181**, 281-292 e286 (2020).
717   26.   D. Wrapp *et al.*, Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation.
718   *Science* **367**, 1260-1263 (2020).
719   27.   H. Hofmann *et al.*, Human coronavirus NL63 employs the severe acute respiratory
720   syndrome coronavirus receptor for cellular entry. *Proceedings of the National Academy of*
721   *Sciences of the United States of America* **102**, 7988-7993 (2005).
722   28.   L. Dai, G. F. Gao, Viral targets for vaccines against COVID-19. *Nature reviews.*
723   *Immunology*, (2020).
724   29.   F. Amanat, F. Krammer, SARS-CoV-2 Vaccines: Status Report. *Immunity* **52**, 583-589
725   (2020).
726   30.   F. Krammer, SARS-CoV-2 vaccines in development. *Nature* **586**, 516-527 (2020).
727   31.   S. F. Ahmed, A. A. Quadeer, M. R. McKay, Preliminary Identification of Potential Vaccine
728   Targets for the COVID-19 Coronavirus (SARS-CoV-2) Based on SARS-CoV
729   Immunological Studies. *Viruses* **12**, (2020).
730   32.   D. F. Robbiani *et al.*, Convergent antibody responses to SARS-CoV-2 in convalescent
731   individuals. *Nature* **584**, 437-442 (2020).
732   33.   T. F. Rogers *et al.*, Isolation of potent SARS-CoV-2 neutralizing antibodies and protection
733   from disease in a small animal model. *Science* **369**, 956-963 (2020).
734   34.   A. Z. Wec *et al.*, Broad neutralization of SARS-related viruses by human monoclonal
735   antibodies. *Science* **369**, 731-736 (2020).
736   35.   B. Ju *et al.*, Human neutralizing antibodies elicited by SARS-CoV-2 infection. *Nature* **584**,
737   115-119 (2020).
738   36.   R. Shi *et al.*, A human neutralizing antibody targets the receptor-binding site of SARS-
739   CoV-2. *Nature* **584**, 120-124 (2020).
740   37.   S. Du *et al.*, Structurally Resolved SARS-CoV-2 Antibody Shows High Efficacy in Severely
741   Infected Hamsters and Provides a Potent Cocktail Pairing Strategy. *Cell* **183**, 1013-1023
742   e1013 (2020).
743   38.   S. J. Zost *et al.*, Potently neutralizing and protective human antibodies against SARS-
744   CoV-2. *Nature* **584**, 443-449 (2020).
745   39.   Y. Wu *et al.*, A noncompeting pair of human neutralizing antibodies block COVID-19 virus
746   binding to its receptor ACE2. *Science* **368**, 1274-1278 (2020).
747   40.   P. J. M. Brouwer *et al.*, Potent neutralizing antibodies from COVID-19 patients define
748   multiple targets of vulnerability. *Science* **369**, 643-650 (2020).
749   41.   J. Hansen *et al.*, Studies in humanized mice and convalescent humans yield a SARS-
750   CoV-2 antibody cocktail. *Science* **369**, 1010-1014 (2020).
751   42.   C. O. Barnes *et al.*, Structures of Human Antibodies Bound to SARS-CoV-2 Spike Reveal
752   Common Epitopes and Recurrent Features of Antibodies. *Cell* **182**, 828-842 e816 (2020).
753   43.   Y. Cao *et al.*, Potent Neutralizing Antibodies against SARS-CoV-2 Identified by High-
754   Throughput Single-Cell Sequencing of Convalescent Patients' B Cells. *Cell* **182**, 73-84
755   e16 (2020).
756   44.   L. Premkumar *et al.*, The receptor binding domain of the viral spike protein is an
757   immunodominant and highly specific target of antibodies in SARS-CoV-2 patients. *Sci*
758   *Immunol* **5**, (2020).
759   45.   M. Yuan *et al.*, Structural basis of a shared antibody response to SARS-CoV-2. *Science*
760   **369**, 1119-1123 (2020).
761   46.   Z. Wang *et al.*, mRNA vaccine-elicited antibodies to SARS-CoV-2 and circulating variants.
762   *Nature* **592**, 616-622 (2021).

47.   Y. Weisblum *et al.*, Escape from neutralizing antibodies by SARS-CoV-2 spike protein
variants. *Elife* **9**, (2020).
48.   E. Andreano *et al.*, SARS-CoV-2 escape from a highly neutralizing COVID-19
convalescent plasma. *Proceedings of the National Academy of Sciences of the United*
States of America **118**, (2021).
49.   M. Yuan *et al.*, Structural and functional ramifications of antigenic drift in recent SARS-
CoV-2 variants. *Science*, (2021).
50.   K. Wu, W. Li, G. Peng, F. Li, Crystal structure of NL63 respiratory coronavirus receptor-
binding domain complexed with its human receptor. *Proceedings of the National Academy*
of Sciences of the United States of America **106**, 19970-19974 (2009).
51.   G. Song *et al.*, Cross-reactive serum and memory B-cell responses to spike protein in
SARS-CoV-2 and endemic coronavirus infection. *Nature communications* **12**, 2938
(2021).
52.   M. M. Sauer *et al.*, Structural basis for broad coronavirus neutralization. *Nature structural*
& molecular biology **28**, 478-486 (2021).
53.   C. Wang *et al.*, A conserved immunogenic and vulnerable site on the coronavirus spike
protein delineated by cross-reactive monoclonal antibodies. *Nature communications* **12**,
1715 (2021).
54.   R. Andrabi *et al.*, Identification of Common Features in Prototype Broadly Neutralizing
Antibodies to HIV Envelope V2 Apex to Facilitate Vaccine Design. *Immunity* **43**, 959-973
(2015).
55.   J. Jardine *et al.*, Rational HIV immunogen design to target specific germline B cell
receptors. *Science* **340**, 711-716 (2013).
56.   L. Cao *et al.*, Differential processing of HIV envelope glycans on the virus and soluble
recombinant trimer. *Nature communications* **9**, 3693 (2018).
57.   W. B. Struwe *et al.*, Site-Specific Glycosylation of Virion-Derived HIV-1 Env Is Mimicked
by a Soluble Trimeric Immunogen. *Cell reports* **24**, 1958-1966 e1955 (2018).
58.   J. Pallesen *et al.*, Immunogenicity and structures of a rationally designed prefusion MERS-
CoV spike antigen. *Proceedings of the National Academy of Sciences of the United States*
of America **114**, E7348-E7357 (2017).
59.   C. L. Hsieh *et al.*, Structure-based design of prefusion-stabilized SARS-CoV-2 spikes.
Science **369**, 1501-1505 (2020).
60.   L. Bao *et al.*, The pathogenicity of SARS-CoV-2 in hACE2 transgenic mice. *Nature* **583**,
830-833 (2020).
61.   D. Pinto *et al.*, Broad betacoronavirus neutralization by a stem helix-specific human
antibody. *Science* **373**, 1109-1116 (2021).
62.   D. R. Burton, A new lease on life for an HIV-neutralizing antibody class and vaccine target.
Proceedings of the National Academy of Sciences of the United States of America **118**,
(2021).
63.   D. R. Burton, Antibodies, viruses and vaccines. *Nature reviews. Immunology* **2**, 706-713
(2002).
64.   D. R. Burton, What Are the Most Powerful Immunogen Design Vaccine Strategies?
Reverse Vaccinology 2.0 Shows Great Promise. *Cold Spring Harb Perspect Biol*, (2017).
65.   S. F. Andrews, A. B. McDermott, Shaping a universally broad antibody response to
influenza amidst a variable immunoglobulin landscape. *Curr Opin Immunol* **53**, 96-101
(2018).
66.   G. Barba-Spaeth *et al.*, Structural basis of potent Zika-dengue virus antibody cross-
neutralization. *Nature* **536**, 48-53 (2016).
67.   D. Corti, A. Lanzavecchia, Broadly neutralizing antiviral antibodies. *Annual review of*
immunology **31**, 705-742 (2013).

68. C. Dreyfus *et al.*, Highly conserved protective epitopes on influenza B viruses. *Science*
337, 1343-1348 (2012).
69. A. I. Flyak *et al.*, Broadly neutralizing antibodies from human survivors target a conserved
site in the Ebola virus glycoprotein HR2-MPER region. *Nat Microbiol* **3**, 670-677 (2018).
70. B. S. Graham, M. S. A. Gilman, J. S. McLellan, Structure-Based Vaccine Antigen Design.
Annu Rev Med **70**, 91-104 (2019).
71. F. Vigant, N. C. Santos, B. Lee, Broad-spectrum antivirals against viral fusion. *Nat Rev*
Microbiol **13**, 426-437 (2015).
72. R. U. Kadam, I. A. Wilson, Structural basis of influenza virus fusion inhibition by the
antiviral drug Arbidol. *Proceedings of the National Academy of Sciences of the United*
States of America **114**, 206-214 (2017).
73. D. Eggink, B. Berkhout, R. W. Sanders, Inhibition of HIV-1 by fusion inhibitors. *Curr Pharm*
Des **16**, 3716-3728 (2010).
74. N. K. Hurlburt *et al.*, Structural definition of a pan-sarbecovirus neutralizing epitope on the
spike S2 subunit. *bioRxiv*, 2021.2008.2002.454829 (2021).
75. Y. Li *et al.*, Linear epitope landscape of the SARS-CoV-2 Spike protein constructed from
1,051 COVID-19 patients. *Cell reports*, 108915 (2021).
76. Y. Li *et al.*, Linear epitopes of SARS-CoV-2 spike protein elicit neutralizing antibodies in
COVID-19 patients. *Cell Mol Immunol* **17**, 1095-1097 (2020).
77. J. T. Ladner *et al.*, Epitope-resolved profiling of the SARS-CoV-2 antibody response
identifies cross-reactivity with endemic human coronaviruses. *Cell Rep Med* **2**, 100189
(2021).
78. B. Briney, A. Inderbitzin, C. Joyce, D. R. Burton, Commonality despite exceptional
diversity in the baseline human antibody repertoire. *Nature* **566**, 393-397 (2019).
79. K. Wiehe *et al.*, Functional Relevance of Improbable Antibody Mutations for HIV Broadly
Neutralizing Antibody Development. *Cell host & microbe* **23**, 759-765 e756 (2018).
80. R. Andrabi, J. N. Bhiman, D. R. Burton, Strategies for a multi-stage neutralizing antibody-
based HIV vaccine. *Curr Opin Immunol* **53**, 143-151 (2018).
81. T. Zohar *et al.*, Compromised Humoral Functional Evolution Tracks with SARS-CoV-2
Mortality. *Cell* **183**, 1508-1519 e1512 (2020).
82. U. Greenbaum *et al.*, High levels of common cold coronavirus antibodies in convalescent
plasma are associated with improved survival in COVID-19 patients. *medRxiv*, (2021).
83. P. Kaplonek *et al.*, Early cross-coronavirus reactive signatures of humoral immunity
against COVID-19. *Sci Immunol* **6**, eabj2901 (2021).
84. M. Sarzotti-Kelsoe *et al.*, Optimization and validation of the TZM-bl assay for standardized
assessments of neutralizing antibodies against HIV-1. *Journal of immunological methods*
409, 131-146 (2014).
85. L. M. Walker *et al.*, Broad and potent neutralizing antibodies from an African donor reveal
a new HIV-1 vaccine target. *Science* **326**, 285-289 (2009).
86. Z. Otwinowski, W. Minor, Processing of X-ray diffraction data collected in oscillation mode.
Methods Enzymol **276**, 307-326 (1997).
87. Afinogenov, II, [History of the pharmacy of the Main Hospital]. *Voen Med Zh* **318**, 57-60
(1997).
88. D. Schritt *et al.*, Repertoire Builder: high-throughput structural modeling of B and T cell
receptors. *Molecular Systems Design & Engineering* **4**, 761-768 (2019).
89. P. Emsley, B. Lohkamp, W. G. Scott, K. Cowtan, Features and development of Coot. *Acta*
Crystallogr D Biol Crystallogr **66**, 486-501 (2010).
90. P. D. Adams *et al.*, PHENIX: a comprehensive Python-based system for macromolecular
structure solution. *Acta Crystallogr D Biol Crystallogr* **66**, 213-221 (2010).
91. E. Krissinel, K. Henrick, Inference of macromolecular assemblies from crystalline state.
Journal of molecular biology **372**, 774-797 (2007).

92.   S. Henikoff, J. G. Henikoff, Amino acid substitution matrices from protein blocks.
Proceedings of the National Academy of Sciences of the United States of America **89**,
10915-10919 (1992).
93.   E. Jurrus *et al.*, Improvements to the APBS biomolecular solvation software suite. *Protein*
science : a publication of the Protein Society **27**, 112-128 (2018).
94.   T. J. Dolinsky, J. E. Nielsen, J. A. McCammon, N. A. Baker, PDB2PQR: an automated
pipeline for the setup of Poisson-Boltzmann electrostatics calculations. *Nucleic acids*
research **32**, W665-667 (2004).
95.   J. Ye, N. Ma, T. L. Madden, J. M. Ostell, IgBLAST: an immunoglobulin variable domain
sequence analysis tool. *Nucleic acids research* **41**, W34-40 (2013).
96.   Y. Cai *et al.*, Distinct conformational states of SARS-CoV-2 spike protein. *Science* **369**,
1586-1592 (2020).
97.   D. Pinto *et al.*, Cross-neutralization of SARS-CoV-2 by a human monoclonal SARS-CoV
antibody. *Nature* **583**, 290-295 (2020).
98.   V. B. Chen *et al.*, MolProbity: all-atom structure validation for macromolecular
crystallography. *Acta Crystallogr D Biol Crystallogr* **66**, 12-21 (2010).
880

Acknowledgements

We thank all of the human cohort participants for donating samples.

Funding

This work was supported by NIH NIAID CHAVD (UM1 AI44462 to D.R.B.), IAVI

Neutralizing Antibody Center, the Bill and Melinda Gates Foundation (OPP 1170236 and

INV-004923 to I.A.W. and D.R.B.), and the Translational Virology Core of the San Diego

Center for AIDS Research (CFAR) AI036214. S.A.R. was supported by NIH

5T32AI007384). This work was also supported by the John and Mary Tu Foundation and

the James B. Pendleton Charitable Trust (to D.M.S. and D.R.B.). X-ray diffraction data

were collected at Stanford Synchrotron Radiation Lightsource (SSRL). SSRL is a

Directorate of SLAC National Accelerator Laboratory, and an Office of Science User

Facility operated for the U.S. Department of Energy Office of Science by Stanford

University. The SSRL Structural Molecular Biology Program is supported by the DOE

Office of Biological and Environmental Research, and by the National Institutes of Health,

National Institute of General Medical Sciences (including P41GM103393) and the

National Center for Research Resources (P41RR001209).

Author contributions

P.Z., M.Y., G.S., I.A.W., D.R.B., and R.A. conceived and designed the study. N.B., J.R.,

M.P., E.G., S.A.R., D.M.S., and T.F.R. recruited donors and collected and processed

plasma samples. P.Z., G.S., and F.A., performed BLI, ELISA and cell binding and virus

neutralization assays. D.H., and L.P., conducted cell-cell fusion experiment. W.H., S.C.,

and P.Y., generated recombinant protein antigens. M.Y. and X.Z. determined the crystal

structure of the antibody-antigen complex. N.B., N.S., J.R.T., and T.F.R. carried out

animal studies and viral load measurements. P.Z., M.Y., G.S., N.B., N.S., D.H., W.H.,

D.N., J.R.T., T.F.R., I.A.W., D.R.B., and R.A. designed the experiments and analyzed the

data. R.A., P.Z., G.S., M.Y., I.A.W. and D.R.B. wrote the paper, and all authors reviewed

and edited the paper.

Competing interests

R.A., G.S., W.H., T.F.R., and D.R.B. are listed as inventors on pending patent

applications describing the SARS-CoV-2 and HCoV-HKU1 S cross-reactive antibodies.

P.Z., G.S., M.Y., I.A.W., D.R.B. and R.A. are listed as inventors on a pending patent

application describing the S2 stem epitope immunogens identified in this study. All other

authors have no competing interests to declare.

Data Availability: All data associated with this study are in the paper or supplementary

materials.

Figures and legends

Figure 1. Identification of the CC40.8 bnAb epitope on the coronavirus spike protein by epitope mapping.

(A) Neutralization of clade 1b (SARS-CoV-2 and Pang17) and clade 1a (SARS-CoV-1, WIV1 and SHC014) ACE2-utilizing sarbecoviruses by CC40.8 mAb isolated from a COVID-19 donor (left) is shown. CC40.8 neutralizing activity against SARS-CoV-2 (WT-Wuhan) and SARS-CoV-2 variants of concern [B.1.1.7 (alpha), B.1.351 (beta), P.1 (gamma) and B.1.617.2 (delta)] is shown on the right.

(B) Left: Cellular ELISA (CELISA) data show binding of CC40.8 mAb with β-HCoV spikes expressed on 293T cells. Binding to HCoV spikes is recorded as % positive cells using flow cytometry. CC40.8 mAb shows cross-reactive binding with 4 out of 5 human β-HCoV spikes. Right: BioLayer Interferometry (BLI) binding of CC40.8 mAb with human β-HCoV soluble spike proteins. Apparent binding constants (K_D^{App}) for each Ab-antigen interaction

are indicated. K_D^{App} <10^{-12}M indicates that no off-rate could be measured. The raw
experimental curves are shown as dash lines, while the solid lines are the fits.
(C) Epitope mapping of CC40.8 with HCoV-HKU1 S2 subunit overlapping peptides is
shown. A series of HCoV-HKU1 S2 (GenBank: AAT98580.1) overlapping biotinylated
peptides (15-residues long with a 10-residue overlap) were tested for binding to CC40.8
mAb by ELISA. OD405, optical density at 405nm. CC40.8 showed binding to the 95[th] 15-
mer peptide corresponding to the HCoV-HKU1 S2 stem-helix region (residue position
range: 1231-1245). An antibody to dengue virus, DEN3, was used as a control.
(D) BLI data are shown for CC40.8 binding to the HCoV-HKU1 95[th] 15-mer stem peptide
(blue) and HCoV-HKU1 stem peptide variants with 5 additional residues either at the N-
(20-mer: brick red) or C-(20-mer: orange) terminus or added at both termini (25-mer: red).
CC40.8 showed strongest binding to the 25-residue stem peptide corresponding to
HCoV-HKU1 S2 residues 1226-1250. The kinetic curves are fit with a 1:1 binding mode.
(E) BLI data are shown for CC40.8 binding to 25-mer stem peptides derived from different
HCoV spikes. CC40.8 showed binding to the β- but not to the α-HCoV S2 stem peptides.
The HCoV-HKU1 S2 residues 1226-1250 correspond to residues 1140-1164 on SARS-
CoV-2 spike. The kinetic curves are fit with a 1:1 binding mode.
(F) A SARS-CoV-2 spike protein cartoon depicts the S2-stem epitope region in green at
the base of the prefusion spike ectodomain.
(G) Sequence conservation of the CC40.8 stem-helix epitope is shown for SARS-CoV-
1/2, HCoV-HKU1 and HCoV-OC43 human β-CoV spike proteins. Conserved identical
residues are highlighted with blue boxes, and similar residues are in cyan boxes [amino
acids scored greater than or equal to 0 in the BLOSUM62 alignment score matrix (*92*)
were counted as similar here]. An N-linked glycosylation site is indicated with a "#"
symbol.
961

Figure 2. Crystal structure of CC40.8 antibody in complex with the SARS-CoV-2 stem peptide, and S2 stem bnAb epitope residues and conservation across CoVs.
(A) An overall view of the CC40.8-peptide complex structure is shown at 1.6 Å resolution. Heavy and light chains of CC40.8 are shown in orange and yellow semi-transparent surfaces, respectively, with the heavy (H) and light (L) chain complementary determining regions (CDRs) shown as tubes. The SARS-CoV-2 stem-helix peptide is shown as a green tube for the peptide backbone.
(B) An overview of the CC40.8 antibody and S2 stem-peptide interaction is shown. Heavy (H) and light (L) chains of CC40.8 are shown in orange and yellow, respectively, whereas

the SARS-CoV-2 stem peptide is in green. Hydrogen bonds and salt bridges are
represented by black dashed lines.
(C) Details of the interactions between CC40.8 and the SARS-CoV-2 stem peptide are
shown. Residues conserved in SARS-CoV-1, SARS-CoV-2, and other sarbecoviruses as
well as seasonal β-CoVs HCoV-HKU-1, and HCoV-OC43 are labeled with asterisks (*).
(D) BLI data are shown for binding of CC40.8 bnAb to SARS-CoV-2 stem-helix peptide
(top) and soluble spike protein alanine mutants (bottom) spanning the whole epitope. The
stem peptide or spike protein mutants that substantially affect CC40.8 bnAb binding are
shown in assorted colors in comparison to wild-type (WT, red).
(E) Neutralization of SARS-CoV-2 and the stem-helix alanine mutants spanning the whole
epitope by CC40.8 is shown. The WT virus is shown in red and virus mutants that
substantially affect CC40.8 bnAb neutralization are shown in assorted colors. The bold
and dashed color curves indicate substitutions that, respectively, led to a decrease or an
increase in the IC_{50} neutralization titers compared to WT virus.
(F) Sequence conservation is shown for the CC40.8 stem-helix epitope on SARS-CoV-
1/2, sarbecoviruses infecting other animal species, human β-CoVs and mouse hepatitis
virus (MHV). The stem region forming the helix is indicated by black dashes and residues
involved in interaction with CC40.8 antibody are indicated by red dots (cutoff distance =
4 Å). Larger dots indicate residues that are essential for CC40.8 interaction as defined by
alanine scanning mutagenesis where mutation decreased neutralization IC_{50} by at least
10-fold or a complete knock-out (details are shown in fig. S7). Conserved identical
residues are highlighted with blue boxes, and similar residues are in cyan boxes [amino
acids scored greater than or equal to 0 in the BLOSUM62 alignment score matrix (*92*)
were counted as similar here]. An N-linked glycosylation site is indicated with a "#"
symbol. The region that presents a helical secondary structure in the CC40.8/peptide
structure is indicated on top of the panel.
998
999

1000
Figure 3. CC40.8 reduces weight loss, lung viral load, and viral replication following
SARS-CoV-2 challenge in the hACE2 mouse model.
(A) CC40.8 was administered intraperitonially (i.p.) at four different doses (300 μg, 100
μg, 50 μg, and 10 μg) per animal into hACE2 receptor-expressing mice (6 animals per
group). Control animals received CC12.1 RBD nAb (300 μg per animal) or a Zika-specific
mAb, SMZAb1 (300 μg per animal). Each group of animals was challenged intranasally
(i.n.) 12 hours after antibody infusion with 2×10^4 PFU of SARS-CoV-2 (USA-WA1/2020).
Animal weight was monitored daily as an indicator of disease progression and lung tissue
was collected on day 5 for viral load and viral burden assessment.
(B) Percent weight change in CC40.8 or control antibody-treated animals after SARS-
CoV-2 challenge is shown. Percent weight change was calculated from day 0 for all
animals. Data are presented as mean ± SEM.
(C) SARS-CoV-2 viral RNA loads based on the qPCR analysis of lung tissue at day 5
after infection are shown. Data are presented as mean ± SEM.
(D) SARS-CoV-2 infectious virus titers (plaque-forming unit (PFU)) are shown as
determined by plaque assay from lung tissue at day 5 after infection. Data are presented
as mean ± SEM.
Statistical comparisons between groups were performed using a Kruskal-Wallis non-
parametric test followed by Dunnett's multiple comparisons. (*p <0.05, **p <0.01, ***p
<0.001; ****p < 0.0001; ns, p >0.05).
1021

Figure 4. Frequency of CC40.8 S2 epitope-targeting serum antibodies in human COVID-19 donors.

(A) The heatmap shows ELISA binding reactivity profiles of convalescent COVID-19 serum samples with 25-mer peptides corresponding to the CC40.8 bnAb S2 epitope on human β-(SARS-CoV-2, SARS-CoV-1, MERS-CoV, HCoV-HKU1, HCoV-OC43) and α-(HCoV-NL63 and HCoV-229E) coronaviruses. The extent of binding (represented as OD_{405} values) is color coded with red indicating strong reactivity. CC40.8 mAb was the positive control for the binding assay and PBS-BSA solution served as the negative control. Six out of 60 COVID-19 convalescent donors showed cross-reactive binding to various HCoV spike stem-helix peptides.

(B) ELISA-based alanine scan epitope mapping is shown for convalescent COVID-19 serum samples from CC6, CC21, CC40, CC48, CC57 and CC65 donors with SARS-CoV-2 stem peptides (25mer). CC40 serum showed dependence on similar SARS-CoV-2 stem-helix residues as the CC40.8 mAb. SARS-CoV-2 stem-helix residue positions targeted (decrease in ELISA binding compared to WT stem peptide) by multiple cross-reactive COVID-19 serum samples are shown in gray. Five residues, F^{1148}, E^{1151}, L^{1152}, Y^{1155} and F^{1156} were commonly targeted by the cross-reactive COVID-19 serum Abs. These residues form the stem-helix bnAb core epitope.

(C) SARS-CoV-2 neutralization by CC40.8 in the presence of competing SARS-CoV-2 stem peptide is shown. Neutralization data are presented for SARS-CoV-2 by CC40.8 mAb, CC40.8 mAb pre-incubated with SARS-CoV-2 stem peptide (60 µg/ml) and stem peptide-only control. The SARS-CoV-2 stem peptide inhibits the neutralizing activity of CC40.8 mAb.

(D) SARS-CoV-2 neutralization by cross-reactive COVID-19 serum samples was evaluated in the presence of competing SARS-CoV-2 stem peptide. Neutralization of SARS-CoV-2 by serum from COVID-19 convalescent donors, CC6, CC21, CC40, CC48, CC57, CC65, pre-incubated with SARS-CoV-2 stem peptide (60 µg/ml) and stem peptide-only controls was measured. The SARS-CoV-2 stem peptide had minimal effects on neutralization by these COVID-19 convalescent donor serum antibodies.

1054 **SUPPLEMENTARY MATERIALS**
1055

1056

Fig. S1. CC40.8 mature and CC40.8 iGL antibodies bind to spike proteins and stem-helix peptides and mature antibody neutralizes pseudotyped coronaviruses.

(A) IC50 neutralization of CC40.8 broadly neutralizing antibody (bnAb) is shown for sarbecoviruses (SARS-CoV-2, SARS-CoV-1, SHC014, Pang17 and WIV1), MERS-CoV and SARS-CoV-2 variants of concern (alpha (B.1.1.7), beta (B.1.351), gamma (P.1) and delta (B.1.617.2)).

(B) Sequence alignment of CC40.8 heavy and light chains with their corresponding germline V-gene sequences (VH3-23 and VL3-10) is shown with the design of CC40.8 antibody inferred germline (iGL) gene sequences. Dots represent identical residues and dashes represent gaps introduced to preserve the alignment.

(C) BioLayer Interferometry (BLI) binding is shown for CC40.8 iGL antibody with human β-HCoV soluble spike proteins. Apparent binding constants (K_D^{App}) for each antibody-antigen interaction are indicated. The raw experimental curves are shown as dash lines, while the solid lines are the fits.

(D) BLI binding is shown for CC40.8 iGL Ab to 25-mer stem peptides derived from all HCoV spike proteins. The kinetic curves are fit with a 1:1 binding mode.

(E) Binding kinetics (K_D^{App} (spike proteins), K_D (stem-helix peptides) k_{on} and k_{off} constants) of CC40.8 and CC40.8 iGL antibodies with human β-HCoV soluble spike proteins and the 25-mer β-HCoV stem peptides are shown.

(F) IC50 neutralization of CC40.8 iGL is shown for sarbecoviruses (SARS-CoV-2 and SARS-CoV-1) and MERS-CoV.

1078

1079
Fig. S2. CC40.8 antibody inhibits SARS-CoV-2 spike protein- and hACE2-mediated
cell-cell fusion.
(A and B) A schematic diagram of cell-cell fusion assay is shown. SARS-CoV-2 spike-
HeLa cells express nucleus-restricted RFP (Red) and hACE2-HeLa cells express
cytosolic GFP (Green). The interaction of SARS-CoV-2 spike protein and hACE2 can lead
to cell fusion to form syncytia. In the same syncytium, both GFP in the cytoplasm and
RFP in the nucleus can be seen (A). If antibody can block cell-cell fusion, no syncytia can
be seen. Only GFP-expressing hACE2-HeLa cells and RFP-expressing SARS-CoV-2
spike-HeLa cells can be seen (B).
(C) SARS-CoV-2 spike-HeLa cells (red) were pre-incubated with negative control
antibody (DEN3) or CC40.8 S2 stem bnAb for 1 hour, and then mixed with hACE2-HeLa
cells (green). Green syncytia were observed with DEN3, indicating widespread cell-cell
fusion mediated by SARS-CoV-2 spike and hACE2; fusion was inhibited by addition of
CC40.8.

Peptide number	HKU1 S2 residues	Sequence	OD405nm CC40.8	OD405nm DEN3	Peptide number	HKU1 S2 residues	Sequence	OD405nm CC40.8	OD405nm DEN3	Peptide number	HKU1 S2 residues	Sequence	OD405nm CC40.8	OD405nm DEN3
1	761-775	SISASYRFVTFEPFN	0.2	0.2	40	956-970	ESQISGYTTAATVAA	0.1	0.2	79	1151-1165	SYKPISFKTVLVSPG	0.2	0.2
2	766-780	YRFVTFEPFNVSFVN	0.2	0.2	41	961-975	GYTTAATVAAMFPPW	0.2	0.2	80	1156-1170	SFKTVLVSPGLCISG	0.2	0.3
3	771-785	FEPFNVSFVNDSIES	0.1	0.2	42	966-980	ATVAAMFPPWSAAAG	0.1	0.2	81	1161-1175	LVSPGLCISGDVGIA	0.2	0.2
4	776-790	VSFVNDSIESVGGLY	0.1	0.2	43	971-985	MFPPWSAAAGIPFSL	0.2	0.2	82	1166-1180	LCISGDVGIAPKQGY	0.2	0.2
5	781-795	DSIESVGGLYEIKIP	0.2	0.2	44	976-990	SAAAGIPFSLNVQYR	0.2	0.2	83	1171-1185	DVGIAPKQGYFIKHN	0.2	0.2
6	786-800	VGGLYEIKIPTNFTI	0.1	0.2	45	981-995	IPFSLNVQYRINGLG	0.2	0.2	84	1176-1190	PKQGYFIKHNDHWMF	0.2	0.2
7	791-805	EIKIPTNFTIVGQEE	0.1	0.2	46	986-1000	NVQYRINGLGVTMDV	0.2	0.2	85	1181-1195	FIKHNDHWMFTGSSY	0.2	0.3
8	796-810	TNFTIVGQEEFIQTN	0.1	0.2	47	991-1005	INGLGVTMDVLNKNQ	0.1	0.2	86	1186-1200	DHWMFTGSSYYYPEP	0.2	0.2
9	801-815	VGQEEFIQTNSPKVT	0.1	0.2	48	996-1010	VTMDVLNKNQKLIAT	0.2	0.2	87	1191-1205	TGSSYYYPEPISDKN	0.2	0.2
10	806-820	FIQTNSPKVTIDCSL	0.2	0.2	49	1001-1015	LNKNQKLIATAFNNA	0.2	0.2	88	1196-1210	YYPEPISDKNVVFMN	0.2	0.2
11	811-825	SPKVTIDCSLFVCSN	0.2	0.2	50	1006-1020	KLIATAFNNALLSIQ	0.2	0.2	89	1201-1215	ISDKNVVFMNTCSVN	0.2	0.2
12	816-830	IDCSLFVCSNYAACH	0.2	0.2	51	1011-1025	AFNNALLSIQNGFSA	0.2	0.2	90	1206-1220	VVFMNTCSVNFTKAP	0.2	0.2
13	821-835	FVCSNYAACHDLLSE	0.2	0.3	52	1016-1030	LLSIQNGFSATNSAL	0.1	0.2	91	1211-1225	TCSVNFTKAPLVYLN	0.2	0.3
14	826-840	YAACHDLLSEYGTFC	0.2	0.3	53	1021-1035	NGFSATNSALAKIQS	0.2	0.2	92	1216-1230	FTKAPLVYLNHSVPK	0.2	0.2
15	831-845	DLLSEYGTFCDNINS	0.2	0.3	54	1026-1040	TNSALAKIQSVVNSN	0.2	0.2	93	1221-1235	LVYLNHSVPKLSDFE	0.2	0.2
16	836-850	YGTFCDNINSILDEV	0.2	0.2	55	1031-1045	AKIQSVVNSNAQALN	0.2	0.2	94	1226-1240	HSVPKLSDFESELSH	0.2	0.2
17	841-855	DNINSILDEVNGLLD	0.1	0.2	56	1036-1050	VVNSNAQALNSLLQQ	0.2	0.2	95	1231-1245	LSDFESELSHWFKNQ	2.5	0.2
18	846-860	ILDEVNGLLDTTQLH	0.1	0.2	57	1041-1055	AQALNSLLQQLFNKF	0.2	0.3	96	1236-1250	SELSHWFKNQTSIAP	0.2	0.2
19	851-865	NGLLDTTQLHVADTL	0.1	0.2	58	1046-1060	SLLQQLFNKFGAISS	0.2	0.2	97	1241-1255	WFKNQTSIAPNLTLN	0.2	0.2
20	856-870	TTQLHVADTLMQGVT	0.1	0.2	59	1051-1065	LFNKFGAISSSLQEI	0.2	0.2	98	1246-1260	TSIAPNLTLNLHTIN	0.2	0.2
21	861-875	VADTLMQGVTLSSNL	0.1	0.2	60	1056-1070	GAISSSLQEILSRLD	0.2	0.2	99	1251-1265	NLTLNLHTINATFLD	0.2	0.2
22	866-880	MQGVTLSSNLNTNLH	0.1	0.2	61	1061-1075	SLQEILSRLDALEAQ	0.2	0.3	100	1256-1270	LHTINATFLDLYYEM	0.2	0.2
23	871-885	LSSNLNTNLHFDVDN	0.1	0.2	62	1066-1080	LSRLDALEAQVQIDR	0.2	0.2	101	1261-1275	ATFLDLYYEMNLIQE	0.2	0.2
24	876-890	NTNLHFDVDNINFKS	0.1	0.2	63	1071-1085	ALEAQVQIDRLINGR	0.2	0.2	102	1266-1280	LYYEMNLIQESIKSL	0.2	0.2
25	881-895	FDVDNINFKSLVGCL	0.2	0.3	64	1076-1090	VQIDRLINGRLTALN	0.2	0.2	103	1271-1285	NLIQESIKSLNNSYI	0.2	0.2
26	886-900	INFKSLVGCLGPHCG	0.2	0.3	65	1081-1095	LINGRLTALNAYVSQ	0.2	0.2	104	1276-1290	SIKSLNNSYINLKDI	0.2	0.2
27	891-905	LVGCLGPHCGSSSRS	0.2	0.2	66	1086-1100	LTALNAYVSQQLSDI	0.2	0.3	105	1281-1295	NNSYINLKDIGTYEM	0.2	0.2
28	896-910	GPHCGSSSRSFFEDL	0.2	0.2	67	1091-1105	AYVSQQLSDISLVKF	0.2	0.3	106	1286-1300	NLKDIGTYEMYVKWP	0.2	0.2
29	901-915	SSSRSFFEDLLFDKV	0.2	0.2	68	1096-1110	QLSDISLVKFGAALA	0.2	0.2	107	1291-1305	GTYEMYVKWPWYVWL	0.3	0.3
30	906-920	FFEDLLFDKVKLSDV	0.1	0.2	69	1101-1115	SLVKFGAALAMEKVN	0.2	0.2	108	1296-1310	YVKWPWYVWLLISFS	0.3	0.2
31	911-925	LFDKVKLSDVGFVEA	0.2	0.2	70	1106-1120	GAALAMEKVNECVKS	0.2	0.2	109	1301-1315	WYVWLLISFSFIIFL	0.2	0.2
32	916-930	KLSDVGFVEAYNNCT	0.2	0.3	71	1111-1125	MEKVNECVKSQSPRI	0.2	0.3	110	1306-1320	LISFSFIIFLVLLFF	0.1	0.2
33	921-935	GFVEAYNNCTGGSEI	0.2	0.2	72	1116-1130	ECVKSQSPRINFCGN	0.2	0.2	111	1311-1325	FIIFLVLLFFICCCT	0.2	0.2
34	926-940	YNNCTGGSEIRDLLC	0.2	0.2	73	1121-1135	QSPRINFCGNGNHIL	0.2	0.2	112	1316-1330	VLLFFICCCTGCGSA	0.2	0.3
35	931-945	GGSEIRDLLCVQSFN	0.2	0.2	74	1126-1140	NFCGNGNHILSLVQN	0.2	0.2	113	1321-1335	ICCCTGCGSACFSKC	0.2	0.3
36	936-950	RDLLCVQSFNGIKVL	0.2	0.2	75	1131-1145	GNHILSLVQNAPYGL	0.2	0.2	114	1326-1340	GCGSACFSKCHNCCD	0.2	0.3
37	941-955	VQSFNGIKVLPPILS	0.2	0.2	76	1136-1150	SLVQNAPYGLLFMHF	0.2	0.3	115	1331-1345	CFSKCHNCCDEYGGH	0.2	0.2
38	946-960	GIKVLPPILSESQIS	0.2	0.2	77	1141-1155	APYGLLFMHFSYKPI	0.2	0.3	116	1336-1350	HNCCDEYGGHHDFVI	0.2	0.3
39	951-965	PPILSESQISGYTTA	0.2	0.2	78	1146-1160	LFMHFSYKPISFKTV	0.2	0.2	117	1341-1355	EYGGHHDFVIKTSHD	0.2	0.2

Fig. S3. Epitope mapping of CC40.8 antibody with HCoV-HKU1 S2 subunit-derived overlapping peptides.

ELISA binding results are shown for CC40.8 mAb with HCoV-HKU1 S2 subunit overlapping peptides (residue number range: 761-1355). Each HCoV-HKU1 S2 subunit peptide is 15-residues long with a 10-residue overlap. Peptide IDs, S2 subunit residue number ranges of 15-mer peptides, and antibody binding responses are shown. CC40.8 exhibited binding to the 95[th] peptide (residue position range: 1231-1245) corresponding to the HCoV-HKU1 S2 stem-helix region. DEN3, an antibody to dengue virus, was used as a control.

Fig. S4. Polyreactivity analysis of CC40.8 and CC40.8 iGL antibodies.
(A) Immunofluorescence showing binding of antibodies to immobilized HEp2 epithelial cells was detected by FITC-labelled secondary antibody. Positive and negative controls for the Hep2 kit assay are provided by the manufacturer.
(B) Antibodies were tested by enzyme-linked immunosorbent assay (ELISA) for binding to the polyspecificity reagent (PSR) from CHO-cell solubilized membrane protein (SMP). 4E10, an HIV MPER-specific antibody known to display polyreactivity, was used as a positive control.

1115
Fig. S5. CC40.8 bnAb structure with SARS-CoV-2 S2 stem-helix peptide.
(A) Surface area of the SARS-CoV-2 stem peptide is shown. Solvent exposed and buried
areas were calculated with Proteins, Interfaces, Structures and Assemblies (PISA) (*91*).
(B) The SARS-CoV-2 stem peptide inserts into a hydrophobic groove formed by the
heavy and light chains of CC40.8. Surfaces of CC40.8 are color-coded by hydrophobicity
[calculated by Color h (https://pymolwiki.org/index.php/Color_h)].
(C) Electrostatic surface potential of the CC40.8 paratope is shown. Electrostatic potential
was calculated by APBS and PDB2PQR (*93, 94*).
1124
1125

Fig. S6. Contribution of CC40.8 bnAb heavy and light chain germline and somatic mutated V-gene residues in S2 stem epitope recognition.

(A) Alignment of CC40.8 with germline VH3-23 and VL3-10 sequences is shown. Paratope residues [defined as buried surface area (BSA) > 0 Å^2 as calculated by PISA (*91*)] are highlighted with yellow boxes. Somatically mutated residues as calculated by IgBLAST (*95*) are highlighted in red.

B. Detailed interactions between CC40.8 Fab and the SARS-CoV-2 stem peptide are shown. Heavy and light chains of CC40.8 are shown in orange and yellow, while the SARS-CoV-2 stem peptide is in pale green. Hydrogen bonds and salt bridges are represented by black dashed lines. Somatically mutated residues are shown in red. Conserved residues among coronaviruses are indicated by asterisks (*).

Alanine scanning of SARS-CoV-2 Spike

SARS-CoV-2		WT	P1140A	L1141A	Q1142A	P1143A	E1144A	L1145A	D1146A	S1147A	F1148A	K1149A	E1150A	E1151
Neutralization	IC50 (ug/ml, CC40.8)	11.5	1.4	0.6	3.3	238.4	14.6	N/A	84.0	20.5	>300	0.6	24.2	>300
	n-fold	1.0	0.1	0.1	0.3	20.7	1.3	N/A	7.3	1.8	>26.11	0.0	2.1	>26.11
BLI Binding	Response Value (CC40.8)	0.61	0.57	N/A	0.59	0.21	0.62	0.02	0.52	0.55	0.35	0.61	0.51	0.07
	% change with WT	100%	95%	N/A	97%	35%	102%	4%	86%	90%	58%	101%	83%	12%
	Response Value (S309)	0.43	0.43	N/A	0.41	0.45	0.42	0.40	0.43	0.45	0.44	0.42	0.48	0.47
	% change with WT	100%	100%	N/A	94%	103%	97%	91%	100%	103%	101%	98%	110%	108%
SARS-CoV-2		L1152A	D1153A	K1154A	Y1155A	F1156A	K1157A	N1158A	H1159A	T1160A	S1161A	P1162A	D1163A	V1164A
Neutralization	IC50 (ug/ml, CC40.8)	143.1	71.0	4.4	>300	>300	4.7	4.5	6.7	1.9	6.4	16.0	16.0	6.6
	n-fold	12.5	6.2	0.4	>26.11	>26.11	0.4	0.4	0.6	0.2	0.6	1.0	1.4	0.6
BLI Binding	Response Value (CC40.8)	0.65	0.62	0.74	0.34	0.64	0.74	0.65	0.69	0.65	0.49	0.52	0.60	0.75
	% change with WT	108%	102%	122%	55%	106%	122%	107%	114%	107%	81%	86%	99%	124%
	Response Value (S309)	0.48	0.44	0.42	0.46	0.49	0.50	0.43	0.51	0.55	0.47	0.40	0.45	0.47
	% change with WT	110%	101%	98%	107%	113%	114%	99%	117%	127%	108%	93%	103%	110%

Neutralization: n-fold <0.3 ; n-fold >5
BLI Binding: % change <80%

Alanine scanning of SARS-CoV-1/2 S2 stem peptide

SARS-CoV-1/2		WT	P1140A	L1141A	Q1142A	P1143A	E1144A	L1145A	D1146A	S1147A	F1148A	K1149A	E1150A	E1151
BLI Binding	Response Value (CC40.8)	3.8	3.5	3.8	3.8	3.7	3.5	0.9	3.3	3.3	2.3	3.7	3.5	0.6
	% change with WT	100%	92%	99%	99%	97%	92%	22%	86%	87%	61%	96%	91%	16%
SARS-CoV-1/2		L1152A	D1153A	K1154A	Y1155A	F1156A	K1157A	N1158A	H1159A	T1160A	S1161A	P1162A	D1163A	V1164A
BLI Binding	Response Value (CC40.8)	2.9	3.0	3.4	2.1	2.4	3.7	3.0	2.6	3.2	3.3	3.2	2.8	2.5
	% change with WT	76%	80%	89%	54%	62%	97%	79%	68%	84%	87%	83%	74%	66%

Alanine scanning of HCoV-HKU1 S2 stem peptide

HCoV-HKU1		WT	H1140A	S1141A	V1142A	P1143A	K1144A	L1145A	S1146A	D1147A	F1148A	E1149A	S1150A	E1151A
BLI Binding	Response Value (CC40.8)	4.2	4.6	4.3	4.4	4.4	4.4	3.3	4.4	4.0	4.0	3.6	4.0	1.7
	% change with WT	100%	110%	104%	105%	106%	105%	79%	105%	96%	96%	87%	96%	41%
HCoV-HKU1		L1152A	S1153A	H1154A	W1155A	K1156A	F1157A	N1158A	Q1159A	T1160A	S1161A	I1162A	A1163A	P1164A
BLI Binding	Response Value (CC40.8)	3.7	4.0	3.8	1.8	2.7	4.5	4.2	4.2	4.3	4.3	4.3	4.2	3.8
	% change with WT	88%	96%	91%	43%	64%	107%	101%	100%	104%	104%	104%	100%	91%

Fig. S7. Epitope mapping of CC40.8 bnAb by alanine scanning mutagenesis of SARS-CoV-2 spike protein and SARS-CoV-2/HCoV-HKU1 S2 stem peptides using neutralization and BLI binding assays.

The upper panel shows the IC_{50} neutralization of CC40.8 bnAb with wild-type (WT) SARS-CoV-2 and spike mutant pseudoviruses and the BLI binding responses with WT SARS-CoV-2 soluble spike protein and alanine mutants. SARS-CoV-2 receptor binding domain (RBD) antibody S309 was a control for the spike protein binding assays. The IC_{50} fold change (n-fold) was calculated by dividing the mutant value by the WT value. For IC_{50}, n-fold <0.3 are indicated in red, and n-fold >5 in orange. The middle and lower panels show BLI binding responses of CC40.8 antibody to WT and alanine mutants of the SARS-CoV-1/2 and HCoV-HKU1 stem-helix peptides, respectively. Binding response values where the % change in binding (from WT peptide) is below 80% are indicated in yellow. Antibody S309 that recognizes a fairly conserved epitope of the RBD of both SARS-CoV-1 and SARS-CoV-2 was used as control. N/A, not available.

Fig. S8. CC40.8 binds to a buried interface of the 3-helix bundle: predicted mechanism of neutralization.

(A) A SARS-CoV-2 spike protein structure is shown in the pre-fusion state. The three protomers are shown in gray, pale green, and white, respectively, with N-linked glycans represented by sticks. The 3-helix bundle stem region is highlighted in a blue-outlined box. Representative epitope residues of CC40.8 are shown in sticks. The CC40.8 epitope is rich in hydrophobic residues. A cryo-EM structure of SARS-CoV-2 spike protein structure in the pre-fusion state that incorporates the coordinates of the 3-helix bundle stem region (PDB: 6XR8, (*96*)) is shown here.

(B) The SARS-CoV-2 spike protein pre-fusion structure was superimposed on the structure of CC40.8 (orange/yellow) in complex with a SARS-CoV-2 S2 peptide. CC40.8 would clash with the other protomers of the spike protein in the pre-fusion state.

(C) A putative neutralization mechanism of CC40.8 is presented. The S2 3-helix bundle region is shown in green, and heavy and light chains of CC40.8 are shown in orange and yellow, respectively. A model for the mechanism of neutralization is shown and inspired by the interaction of a mouse S2 stem antibody, B6, isolated from a spike protein vaccinated animal that targets a similar stem epitope (*52*) .

Fig. S9. Comparison of bnAbs CC40.8 and B6 that target the S2 stem helix.
(A) A comparison between S2 stem-helix peptides targeted by CC40.8 and B6 is shown. Peptides used for co-crystallization with CC40.8 or B6 are indicated by dashes, with the regions observed in the crystal structures of each study indicated. Residues involved in interactions with CC40.8 and B6 are indicated by black dots (cutoff distance = 4 Å).
(B to E) Structures of CC40.8 and B6 were compared. The heavy and light chains of CC40.8 are colored in orange and yellow, respectively, and those for B6 are in cyan and light cyan. The S2 stem-helix peptides are shown in green. In panels (D) and (E), a SARS-CoV-2 spike protein pre-fusion structure (PDB 6XR8) is superimposed on structures of CC40.8 and B6 in complex with a SARS-CoV-2 S2 peptide in the green protomer (indicated by arrows). Both CC40.8 and B6 would clash with the other protomers of the spike protein in pre-fusion state.
(F) BLI binding kinetics of CC40.8 to S2- and S6- stabilized SARS-CoV-2 spike trimers are shown. An RBD-targeting neutralizing Ab S309 (*97*) was used as a control. Apparent binding constants (K_D^{App}) of antibodies with spike proteins are shown. The raw experimental curves are shown as dash lines, while the solid lines are the fits.

Figure S10. Weight loss, viral titers, and serum antibody titers were measured in hACE2 mice passively administered CC40.8.

(A) Percent day 5 weight change was calculated from day 0 for all animals. Data are presented as mean ± SEM. Significance was calculated with Dunnett's multiple comparisons test between each experimental group and the ZIKV Ab (SMZAb1) control group (***P<0.001, ****P<0.0001).

(B) Serum human IgG concentrations of CC40.8, CC12.1 and SMZAb1 were assessed by ELISA at day 1, 2, 3, and 5 post infection. Data are presented as mean ± SEM.

1199

Fig. S11. CC40.8 reduces weight loss and lung viral load and viral replication following SARS-CoV-2 challenge in Syrian hamsters.

(A) CC40.8 was administered intraperitonially (i.p.) at a 2 mg per animal dose into Syrian hamsters (average: 16.5 mg/kg). Control animals received 2 mg of control SMZAb1. Each group of five animals was challenged intranasally (i.n.) 12 hours after antibody infusion with 1×10^6 PFU of SARS-CoV-2. Animal weight was monitored daily as an indicator of disease progression and lung tissue was collected on day 5 for viral burden assessment. **(B)** Percent weight change is shown for CC40.8 or control antibody-treated animals after SARS-CoV-2 challenge. Percent weight change was calculated from day 0 for all animals. Data are presented as mean ± SEM. **(C)** SARS-CoV-2 titers (PFU) were determined by plaque assay from lung tissue at day 5 after infection. Three out of 5 CC40.8-treated animals had substantially reduced viral titers compared to the SMZAb1 control antibody-treated animals. Data are presented as mean ± SEM.

1214

1215 **Table S1. X-ray data collection and refinement statistics.**

Data collection	CC40.8 Fab + SARS-CoV-2 S2 peptide
Beamline	SSRL12-1
Wavelength (Å)	0.97946 Å
Space group	P 2 2$_1$ 2$_1$
Unit cell parameters	
a, b, c (Å)	54.9, 63.7, 122.0
α, β, γ (°)	90, 90, 90
Resolution (Å) [a]	50.0-1.62 (1.65-1.62)
Unique reflections [a]	54,176 (4,897)
Redundancy [a]	4.3 (3.0)
Completeness (%) [a]	97.0 (89.6)
$<I/\sigma_I>$ [a]	29.9 (1.0)
R_{sym}[b] (%) [a]	7.9 (>100)
R_{pim}[b] (%) [a]	2.8 (47.7)
CC$_{1/2}$[c] (%) [a]	99.4 (56.3)
Refinement statistics	
Resolution (Å)	29.1-1.62
Reflections (work)	54,129
Reflections (test)	1,997
R_{cryst}[d] / R_{free}[e] (%)	17.4/20.6
No. of atoms	3,836
Fab	3,159
Peptide	193
Ligands	35
Solvent	459
Average B-values (Å^2)	28
Fab	26
Peptide	34
Ligands	56
Solvent	39
Wilson B-value (Å^2)	23
RMSD from ideal geometry	
Bond length (Å)	0.006
Bond angle (°)	1.22
Ramachandran statistics (%) [g]	
Favored	98.2
Outliers	0.0
PDB code	7SJS

[a] Numbers in parentheses refer to the highest resolution shell.
[b] $R_{sym} = \Sigma_{hkl} \Sigma_i | I_{hkl,i} - <I_{hkl}> | / \Sigma_{hkl} \Sigma_i I_{hkl,i}$ and $R_{pim} = \Sigma_{hkl} (1/(n-1))^{1/2} \Sigma_i | I_{hkl,i} - <I_{hkl}> | / \Sigma_{hkl} \Sigma_i I_{hkl,i}$, where $I_{hkl,i}$ is the scaled intensity of the ith
measurement of reflection h, k, l, $<I_{hkl}>$ is the average intensity for that reflection, and n is the redundancy.
[c] CC$_{1/2}$ = Pearson correlation coefficient between two random half datasets.
[d] $R_{cryst} = \Sigma_{hkl} | F_o - F_c | / \Sigma_{hkl} | F_o |$ x 100, where F_o and F_c are the observed and calculated structure factors, respectively.
[e] R_{free} was calculated as for R_{cryst}, but on a test set comprising 5% of the data excluded from refinement.
[g] From MolProbity (*98*).
1223

1224 **Table S2. Demographic information of COVID-19 convalescent donors.**

	COVID donor (n = 60)
Age (years)	20 to 72 (median = 46)
Gender	
Male	47% (28/60)
Female	53% (32/60)
Race/Ethnicity	
White, non-Hispanic	80% (48/60)
Hispanic	8.3% (5/60)
Black, non-Hispanic	1.7% (1/60)
Asian, non-Hispanic	3.3% (2/60)
Unknown	6.7% (4/60)
SARS-CoV-2 PCR Positivity	75% (45/60)
Lateral Flow Positivity	60% (36/60)
Disease Severity	
Mild	56.7% (34/60)
Mild to Moderate	6.7% (4/60)
Moderate	25% (15/60)
Moderate to Severe	5% (3/60)
Severe	5% (3/60)
Critical	1.7% (1/60)
Symptoms	
Cough	60% (36/60)
Fever	55% (33/60)
Fatigue	38.3% (23/60)
Anosmia	31.7% (19/60)
Dyspnea	26.7% (16/60)
Diarrhea	16.7% (10/60)
Days Post Symptom Onset at Collection	6 to 90 (median = 35.5)

1225

1226

1227
1228
1229