The long non-coding RNA *MaTAR20* promotes mammary tumor growth by regulating angiogenesis pathways

Sarah D. Diermeier[1,2,*], Kung-Chi Chang[2,3], Ashleigh Frewen[1], Brian A. Benz[2], Suzanne Russo[2], Dawid Nowak[2], Stephen Hearn[2], Allen Yu[2,4], John E. Wilkinson[5], Frank Rigo[6], and David L. Spector[2,3,4*]

[1]Department of Biochemistry, University of Otago, Dunedin 9016, New Zealand

[2]Cold Spring Harbor Laboratory, Cold Spring Harbor, NY 11724, USA

[3]Stony Brook University, Molecular and Cellular Biology Program, Stony Brook, NY 11794, USA

[4]Stony Brook University, Genetics Program, Stony Brook, NY 11794, USA

[5]Department of Pathology, University of Michigan, Ann Arbor, MI 48109, USA

[6]Ionis Pharmaceuticals, Carlsbad, CA 92010, USA

*Correspondence:

Sarah D. Diermeier (sarah.diermeier@otago.ac.nz, +64 20 4221985) or

David L. Spector (spector@cshl.edu, +1 (516) 367-8456)

Running title: *MaTAR20* as new driver of mammary tumor growth

Keywords: long non-coding RNAs; breast cancer; antisense oligonucleotides; Vegf; Tnfsf15

Financial Support:

This research was supported by NCI 5P01CA013106-Project 3 (D.L.S.), CSHL/Northwell Health support (D.L.S), a Susan G. Komen postdoctoral fellowship (S.D.D.) and Rutherford Discovery Fellowship RDF-UOO1802 awarded by the Royal Society Te Apārangi (S.D.D.).

Conflict of interest:

The authors declare no conflict of interest. D.L.S. is a consultant to, and receives research reagents from, Ionis Pharmaceuticals.

Abstract

Long non-coding RNAs (lncRNAs) are an emerging class of regulatory molecules

that have been shown to play important roles in tumorigenesis and cancer progression.

Here, we studied the recently identified lncRNA *Mammary Tumor Associated RNA 20*

(*MaTAR20*) in mammary cancer progression. A CRISPR/Cas9 knockout of *MaTAR20* in

the metastatic 4T1 cell line led to reduced cancer cell proliferation and increased cell

surface adhesion compared to control cells. Consistent with these knockout results

antisense oligonucleotide (ASO) mediated knockdown of *MaTAR20* resulted in reduced

growth and invasion in 4T1 cells, and in primary mammary tumor organoids derived from

the MMTV-PyMT mouse model of breast cancer. Injection of *MaTAR20*-specific ASOs

subcutaneously into tumor bearing MMTV-PyMT mice resulted in smaller and highly

necrotic tumors in comparison to mice injected with a scrambled control ASO. To

investigate the molecular mechanism by which *MaTAR20* acts to advance mammary

tumor progression, we applied a combination of RNA-sequencing and RNA-pulldown

coupled to DNA-sequencing. These analyses demonstrated that the nuclear retained

lncRNA is associated with several essential cancer signaling pathways such as VEGF

signaling. In particular, *MaTAR20* directly binds to and regulates the expression of

Tnfsf15. Our results suggest that *MaTAR20* is likely an important driver of mammary

tumor progression and represents a promising new therapeutic target.

Introduction

Breast cancer (BC) is estimated to account for 30% of new cancer diagnoses and 15% of cancer-related deaths in women in 2021 (1). While currently available chemo- and targeted therapies have led to improved overall survival rates, declines of mortality have slowed over the past decade compared to other types of cancer (1). Metastatic disease in particular is the main cause for BC related mortality (1), indicating the need for innovative approaches to target the metastatic cascade.

Recent studies highlight the potential of long non-coding RNAs (lncRNAs) as new therapeutic targets in cancer (2-13). Many lncRNAs are expressed in a tissue- and cancer-specific manner (14,15) and several previous studies have shown lncRNAs as drivers of tumorigenesis, tumor growth and invasion (reviewed in (16) and (17)). A promising new approach of targeting lncRNAs to reduce mammary tumor growth and metastasis *in vivo* are nucleic acid based therapies (reviewed in (17)). Initial success in targeting oncogenic lncRNAs was shown using locked nucleic acids (LNAs) to target *BCAR4* (5) or antisense oligonucleotides (ASOs) for *Malat1* (6,18).

We previously identified 30 <u>Ma</u>mmary <u>T</u>umor <u>A</u>ssociated <u>R</u>NAs (*MaTARs*) (7). These lncRNAs are over-expressed in mouse models of BC and in human breast tumor tissue compared to normal mammary epithelial cells (7). ASO-mediated knockdown of various *MaTARs* in primary mammary tumor cells resulted in a tumor cell - specific reduction of cell growth (7). Recently, we demonstrated that *MaTAR25* expression contributes to BC progression via regulation of the *Tensin1* gene (13). Here, we investigated the potential of *MaTAR20* (also known as *Gm13387*, *RP23-132N23.1*, *ENSMUSG00000087028*) as a new therapeutic target in BC, as well as the molecular

mechanism by which *MaTAR20* acts. ASO-mediated knockdown or CRISPR/Cas9

knockout of *MaTAR20* results in reduced cell proliferation and invasion. Further, ASO

knockdown of *MaTAR20 in vivo* leads to delayed tumor onset, decreased tumor size,

increased tumor necrosis and reduced metastatic burden. Based on our investigation of

the molecular mechanism of *MaTAR20*, the observed phenotype is likely the result of

reduced tumor vascularization upon *MaTAR20* reduction. *MaTAR20* regulates genes

involved in tumor angiogenesis, such as increasing the expression of the *Vascular*

Endothelial Growth Factor B (*Vegf-B*) gene. *MaTAR20* regulates the balance of the

Tnfsf15/Vegf axis by directly binding to the *Tnfsf15* locus to repress its expression during

tumor progression.

Materials and Methods

Cell culture

Murine 4T1, Py2T, G0771 and D2A cells were cultured in DMEM supplemented

with 10% FBS and 1% penicillin/streptomycin. All cells were grown in a cell culture

incubator at 37 °C and 5% CO2. ASOs were delivered to the cells via free uptake

immediately after seeding the cells. For proliferation assays, cells were seeded in 24-well

plates and trypsinized at individual time points. Cell counts were determined manually

using a hemocytometer. Cell cycle analysis was performed using a BrdU FITC kit (BD)

as described in (13).

Organoid culture

Murine mammary tumor organoids were generated and cultured as previously

described in (7,19). Briefly, organoids were generated from nulliparous MMTV-PyMT

tumors, mixed with growth factor-reduced Matrigel and grown in DMEM/F12 medium

supplemented with 1x ITS (insulin, transferrin, and sodium selenite), 1%

penicillin/streptomycin and murine FGF2 (2.5 nM). For ASO-mediated knockdown

experiments, organoids were seeded at a density of 5 organoids/µl and plated as 80 µl

domes in 24-well dishes. ASOs were added directly (free uptake) to the culture medium

15-20 min after the organoids were plated and in fresh medium at day 3. Organoids were

incubated for a total of 6 days. ASO sequences are provided in Supplementary Table S1.

For visualization purposes and quantification of organoid branching, images were

acquired using an Axio-Observer Live Cell inverted microscope (Zeiss).

Scanning electron microscopy (SEM)

SEM on 2D cells was performed as described previously (20). Briefly, cells cultured

on glass coverslips were fixed in 1.6% glutaraldehyde in PBS for 30 minutes and then

dehydrated in a graded ethanol series. Coverslips were critically point dried (CPD) using

a Samidri 795 CPD device (Tousimis), mounted on aluminum stubs with carbon tabs

(Electron Microscopy Sciences). Samples were coated with gold using an Emitech K550X

Sputter Coater. Samples were imaged with a Hitachi S3500 SEM operated at 5 kV.

For SEM on 3D organoids, Matrigel containing organoid samples were removed

from multi-well tissue culture dishes (using a 3.5 mL disposable transfer pipette and

placed in 45 mL of cold PBS (4°C) in 50 mL capacity polypropylene centrifuge tubes

(VWR Scientific). Matrigel was washed away by 3-4 rounds of suspension in cold PBS

and centrifugation using a refrigerated centrifuge set to 1,000 pm, 20 min, 4°C (Beckman

Coulter Allegra X15r). After the final wash to remove Matrigel, organoids were transferred

to 1.6 mL microcentrifuge tubes and fixed in 1% glutaraldehyde in PBS at 4°C, for 30

minutes, rinsed in distilled water by centrifugation and resuspension and then post-fixed

in 1% aqueous osmium tetroxide (Electron Microscopy Sciences) for 30 minutes at RT.

After osmium fixation, organoids became dense relative to solutions and a centrifuge was

not needed for further handling. Organoids were dehydrated in a graded ethanol series

and then immersed in 50% hexamethyldisilazane (HMDS, Electron Microscopy Sciences)

in 100% ethanol for 5 min and 100% HMDS for 10 min. Organoids in 0.3 mL of HMDS

were aspirated using a transfer pipette and the solution and suspended organoids were

deposited to 12 mm diameter SEM stubs coated with carbon adhesive tabs (Ted Pella

Inc) and allowed to air dry. Samples were coated with gold using an Emitech K550X

Sputter Coater and imaged with a Hitachi S3500 SEM operated at 5 kV.

RNA extraction and qRT-PCR

Total RNA was isolated from cells or organoids using TRIzol. DNase I treatment

was performed for 15 min at RT to remove contaminating DNA. cDNA synthesis was

carried out using the TaqMan Reverse Transcription kit (Life Technologies) and random

hexamers according to the manufacturer's instructions. Quantitative Real time PCR (qRT-

PCR) was performed using the Power SYBR Green Master Mix (Life Technologies).

Cycling conditions were as follows: 15 min at 95 °C followed by 40 cycles of 15 sec at 94

°C, 30 sec at 60°C. *Peptidylprolyl isomerase B* (*cyclophilin B*) was used as an

endogenous control to normalize each sample and relative expression results were

calculated using the $2^{-\Delta\Delta Ct}$ method. A list of primers used is provided in Supplementary

Table S1.

Subcellular localization

For single-molecule FISH, cells were seeded onto acid-washed glass coverslips

and fixed in 4% paraformaldehyde when reaching 50% confluency. RNA-FISH was

carried out using the Affymetrix View ISH Cell Assay Kit and custom probes (Thermo

Fisher) according to the manufacturer's instructions. The slides were imaged on a LSM

710/780 confocal microscope (Zeiss). Cell fractionation assays were performed as

described in (13).

CRISPR/Cas promoter deletion

To generate a genetic knockout, two sgRNAs targeting the promoter region were

combined, creating a deletion. A sgRNA targeting the *Renilla* luciferase gene was used

as non-targeting control. All sgRNAs were cloned into a lentiCRISPR_V2 plasmid

(Addgene #52961) also encoding Cas9 and delivered to the cells using lentiviral

transduction as described in (21). Stable integrands were selected using puromycin

selection and single cell sorted using a FACS Aria (SORP) Cell Sorter. Each single cell

clone was propagated and analyzed by genomic PCR and qRT-PCR to select for

homozygous knockout clones. Sequences for sgRNAs are provided in Supplementary

Table S1.

Invasion assay

Invasion assays were performed as described previously in (7). Briefly, a Cultrex®

96 well BME Cell Invasion Assay (Trevigen) was used. Cells were starved in FBS-free

culture medium, then harvested and seeded at a density of 5×10^4 cells/well into the

invasion chamber. ASOs were added to growth medium containing 10% FBS. The plate

was incubated at 37 °C for 24 h and the assay was performed according to the

manufacturer's instructions. The fluorescence was measured with a SpectraMax i3 Multi-

Mode Detection Platform (Molecular Devices) using the 480/520 nm filter set. Each

sample was measured in triplicate.

170 Rescue Assay

MaTAR20 isoforms were amplified by PCR using Phusion High-Fidelity DNA

Polymerase (NEB) following the manufacturer's instructions. *MaTAR20* isoform 1 was

cloned using XhoI and EcoRI-HF overhangs. The digested PCR product was ligated into

a pCMV6-Entry plasmid (Origene) with T4 DNA ligase. Isoforms 2 and 3 were amplified

by PCR, gel extracted and treated with T4 PNK. The pCMV6-Entry plasmid was digested

using EcoRV-HF and Eco53kI followed by dephosphorylation using Quick CIP. Isoforms

2 and 3 were ligated into the pCMV6-Entry backbone as described above. Ligations were

transformed into competent DH5α cells using heat shock. Transformants were selected

for using Kanamycin at 25 μg/mL and validated using Sanger sequencing.

For ectopic expression of *MaTAR20*, knockout cells were seeded into 96-well

plates at a density of 5,000 cells/well. The plates were incubated overnight (16-18 h) at

37 °C with 5% CO_2. DNA was transfected using Lipofectamine 2000 (Thermo Fisher) in

serum-free DMEM medium at a ratio of 100 ng DNA : 0.2 μL Lipofectamine. After 6 h, the

medium was replaced with 100 μL/well of fresh complete medium. Cells were fixed for

each time point using 100 μL/well of staining solution (150 μg/mL saponin, 4 μg/mL DAPI,

0.5% PFA in PBS) for 30 minutes with gentle shaking at RT, then stored at 4 °C .The cells

were counted using a Cytation5 automated cell counting system.

RNA was isolated 24 h after transfection. The medium was removed and 50 μL of

TRIzol was added to each well of a 96-well plate. RNA was extracted according to the

manufacturer's instructions. For each sample, 1 μg of RNA was treated with ezDNase

and reverse transcribed to cDNA using SuperScript IV VILO following the manufacturer's

instructions (Thermo Fisher). For RT-qPCR testing of the *MaTAR20* rescue, PowerUp

SYBR Green Master Mix (Thermo Fisher) was used. All RT-qPCRs were run on a

QuantStudioTM 3 Real-Time PCR Machine (Applied Biosystems).

Animal experiments

Animal experiments were carried out in the CSHL Animal Shared Resource, in

accordance with IACUC approved procedures. MMTV-PyMT mice (22) were obtained

from Dr. Mikala Egeblad (CSHL). Tumors and normal mammary glands were extracted

immediately after euthanizing the animal and processed to generate primary cells,

organoids or tissue sections. For *in vivo* ASO injections, female MMTV-PyMT

mice were divided into two cohorts, being treated either with a *MaTAR20*-specific

ASO or a scrambled control ASO at ~8-10 weeks of age (after formation of palpable

tumors). MOE ASOs were injected subcutaneously three times per week, cET ASOs

twice per week, both at 50 mg/kg per dose. Tumors were measured twice per week

throughout the treatment course. At the end of the experiment, mice were euthanized,

and primary tumors and lungs were fixed in 4% PFA and embedded in paraffin for histo-

pathological analysis, or snap-frozen for RNA extractions. FFPE blocks were sectioned

and stained with hematoxylin and eosin (H&E). Slides were scanned and analyzed using

the Aperio ImageScope software.

RNA-seq

RNA quality was assessed on an Agilent 2100 Bioanalyzer. Libraries were

prepared on samples with RIN $\geq$9 using the Illumina TruSeq sample prep kit v2 and

sequenced on an Illumina NextSeq instrument. Data was analyzed as previously

described in (7). Briefly, the quality of FASTQ files was assessed using FastQC

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc/). Reads were mapped to

GRCm38 using STAR (23), and the reads per gene record were counted using HTSeq-

count (24). Differential gene expression was performed with DESeq2 (25), and an

adjusted p-value of < 0.05 was set as threshold for statistical significance. KEGG pathway

and GO term enrichment and was carried out using the R/Bioconductor packages GAGE

(26) and Pathview (27).

ChIRP-seq

ChIRP-seq was carried out as previously described (13). Briefly, 20 million cells

were harvested and fixed in 1% glutaraldehyde solution. ChIRP was performed using two

individual tiling pools of biotinylated oligonucleotide probes. A probe pool targeting mouse

Ppib was used as negative control. A list of ChIRP-seq probes is provided in

Supplementary Table S1. ChIRP-Seq libraries were generated using the Illumina TruSeq

ChIP Library Preparation Kit and sequenced on an Illumina NextSeq instrument. ChIRP-

seq   data   quality   was   assessed   using   FastQC

(http://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and paired-end reads were

mapped to GRCm38 using Bowtie2 (28). ChIRP seq analysis was performed using

HOMER (29). For RT-qPCRs testing the ChIRP cDNA, 5 ng cDNA and 5 µM of primers

in 10 µL was used (reagents and cycling conditions as described under "Rescue assay").

Data access

RNA-seq and ChIRP-seq data have been deposited at Gene Expression Omnibus (GEO)

under accession number GSE171085.

Results

MaTAR20 is a tumor-specific, nuclear-retained lncRNA

MaTAR20 is an intergenic transcript located on mouse chromosome 2. According to GENCODE vM18, three isoforms of *MaTAR20* exist, comprising 2,646, 659 and 638 nt (Figure 1A). Our computational analysis of ENCODE expression data revealed that the lncRNA is absent or expressed at low levels in normal mammary glands and other normal tissues, but highly expressed (FPKM >200) in mammary tumors derived from the transgenic MMTV-PyMT mouse model of luminal BC (Figure 1B). Furthermore, *MaTAR20* levels correlate with tumor size in the MMTV-PyMT model (Supplementary Figure S1A).

To examine the localization of *MaTAR20* we performed single molecule RNA-FISH in primary MMTV-PyMT cells (Figure 1C). Our results indicate that *MaTAR20* is indeed an abundant lncRNA in primary mammary tumor cells, and predominantly localizes to the nucleus, averaging 10-40 foci per nucleus. We further confirmed that *MaTAR20* is a nuclear-retained transcript by subcellular fractionation assays, which revealed an enrichment in both the nucleoplasm and the chromatin fraction (Figure 1D), a localization pattern comparable to the lncRNA *Malat1*. Our initial characterization indicates that *MaTAR20* is a tumor-specific, nuclear-enriched lncRNA that may play a role in tumor progression based on its expression profile.

Knockdown of *MaTAR20* leads to reduced organoid branching

To determine if *MaTAR20* is a driver of mammary tumor cell growth, we performed knockdown experiments *ex vivo*, in MMTV-PyMT derived tumor organoids (6,7,30) (Figure 1E-H). Organoids are an excellent model system to study cancer biology and to test new therapeutic treatments (for review, see (31)). Wildtype/untreated organoids

exhibit a branched morphology consistent with collective cell migration (Fig 1G-H) and/or

proliferation (for review, see (32). We achieved *MaTAR20* knockdown efficiencies of 50-

60% using 2'-O-methoxyethyl (MOE) ASOs in 3D mammary tumor organoids by free

uptake (Figure 1E). Treatment of MMTV-PyMT tumor organoids with two independent

MOE ASOs targeting *MaTAR20* lead to a reduction of organoid branching compared to

untreated organoids, or organoids treated with a scrambled control MOE ASO (scASO)

(Figure 1F-H). Based on these and our previous results in primary mammary epithelial

cells (7), we conclude that *MaTAR20* enhances both the proliferative and invasive

potential of mammary tumor cells.

Promoter deletion of *MaTAR20* results in a less aggressive phenotype

To independently validate our ASO-mediated knockdown results, and to develop

a model system that enables us to study the molecular mechanism of *MaTAR20* in detail,

we generated *MaTAR20* loss-of-function cells using the CRISPR/Cas9 system. First, we

tested a panel of mouse mammary tumor cell lines to identify a suitable model system.

Of the tested cell lines, the metastatic 4T1 line (33) showed the highest *MaTAR20*

expression level (Supplementary Figure S2A). Next, we established promoter deletion of

MaTAR20 by stably integrating Cas9 and two guide RNAs (gRNAs): the first gRNA was

designed to bind upstream of the *MaTAR20* transcription start site (at -736) and the

second gRNA binding within the first exon of *MaTAR20* (+12 bp) (Figure 2A), resulting in

promoter deletion of 748 bp. A gRNA targeting the *Renilla* luciferase gene was used to

generate negative control cells. All edited cell populations were single cell sorted to

generate monoclonal cell lines. Two *MaTAR20* knockout clones (KO1, KO2) and two

negative control clones (Ren1, Ren2) were analyzed further. The KO cell lines showed a

~80-85% reduction of *MaTAR20* expression compared to control clones (Figure 2B). We

used four different primer pairs to measure *MaTAR20* levels, one amplifying all three

isoforms of *MaTAR20* (PP1, Figure 2B), and three isoform-specific primer pairs (PP2,

PP3, PP4; Supplementary Figure S2B). Our results indicate that all three isoforms of

MaTAR20 were reduced to comparable levels in the KO cells.

Similar to our ASO-mediated knockdown experiments in primary mammary tumor

cells (7), we observed a 30-40% reduction of cell proliferation in the KO cell lines (Figure

2C). This phenotype was not caused by differences in cell cycle profiles (Figure 2D) and

did not correlate with Cas9 expression (Supplementary Figure S2C). Ectopic expression

of *MaTAR20* in KO cells was able to rescue the cell proliferation phenotype, restoring

growth rates to levels comparable to Ren control cells (Supplementary Figure S3A-B).

This finding also indicates that the *MaTAR20* transcript is responsible for the observed

phenotype, rather than the genomic locus itself playing a regulatory role and that

MaTAR20 plays a role in *trans*.

While culturing the *MaTAR20* KO cells, we noticed a difference in cell adhesion

between the KO clones and the control cell lines. The KO cells were particularly difficult

to dislodge from cell culture flasks, leading to increased (2-4 fold) trypsinization times

required to detach the KO cells. Further examination of the KO and Ren control cells

using scanning electron microscopy revealed that reduced *MaTAR20* levels lead to

stronger surface attachment of the cells, accompanied by more organized, sheet-like

structures (Figure 2E). In summary, promoter deletion of *MaTAR20* leads to reduced

tumor cell growth and a less invasive phenotype compared to control cells, similar to

305 primary cells treated with *MaTAR20* ASOs (Figure 1). Thus, we conclude that the level of

306 *MaTAR20* impacts mammary tumor cell growth and adhesion.

308 **cET ASOs show improved knockdown efficiency *in vitro***

309  To determine the effect of *MaTAR20* reduction on tumor progression, and to

310 investigate whether ASOs are a viable strategy to systemically reduce *MaTAR20*, we

311 performed *in vivo* experiments. First, we designed a constrained ethyl (cET) ASO

312 targeting *MaTAR20*. In comparison to MOE ASOs (Figure 1), cET ASOs show increased

313 potency *in vivo* (34) and are the chemistry of choice for tissues less sensitive to ASO

314 modulation with several clinical trials currently ongoing (35).

315  Before introducing the cET ASO into animals, we tested its knockdown (KD)

316 efficiency *in vitro*. Concentrations as low as 250 nM were sufficient to achieve >60%

317 knockdown of *MaTAR20* within 24 h in 4T1 cells (Figure 3A), and we observed KD

318 efficiencies of >95% using 500 nM after 72 h (Figure 3B). These results compare

319 favorably to concentrations of 4-5 µM of MOE ASOs achieving ~50% KD in organoids

320 (Figure 1) and ~70% KD in primary cells (7). Knockdown of *MaTAR20* using the cET ASO

321 resulted in significantly reduced cell proliferation and invasion (Figure 3C-D). These

322 results agree with our observations using MOE ASOs (Figure 1) and *MaTAR20* promoter

323 deletions (Figure 2).

325 **Knockdown of *MaTAR20 in vivo* leads to a reduction in tumor growth and induces**

326 **tumor necrosis**

To further investigate the potential of cET ASOs targeting *MaTAR20* in mammary

tumors, we injected 100 mg/kg/week of either *MaTAR20* cET or a negative control ASO

(scrambled = scASO cET) into MMTV-PyMT (C57BL/6) mice after palpable tumor

formation (n=7 for scASO, n=8 for *MaTAR20* ASO; Figure 4A). We achieved an 80%

reduction of *MaTAR20* in the tumors on average, further highlighting the potency of the

cET chemistry (Figure 4B). We initiated ASO injections on animals with comparable

average tumor burden in both cohorts and observed a 30% overall reduced tumor burden

in *MaTAR20* cET treated mice over the course of eight weeks (Figure 4C). While mice in

the *MaTAR20* cET group eventually went on to develop tumors, tumor onset was delayed

in comparison to the scASO cET group. Furthermore, when comparing the growth curves

of similarly sized tumors that developed at the same time in both cohorts, tumors in the

MaTAR20 cET group grew slower compared to scASO cET treated mice, resulting in a

difference of 75% after 8 weeks (Figure 4D). After all mice reached the study endpoint,

we carried out hematoxylin and eosin (H&E) staining of the fixed tumors to investigate

potential histo-pathological differences. Notably, tumors in the *MaTAR20* cET treated

group showed severe necrosis compared to the control group (Figure 4E).

Knockdown of *MaTAR20 in vivo* leads to reduced metastatic burden

The MMTV-PyMT model is highly aggressive, with a metastasis incidence rate of

>80% (22). We examined the lungs in our *in vivo* study post-mortem for macro-metastatic

nodules. While 7/7 mice in the scASO group developed at least one macro-metastatic

nodule, only 4/7 mice in the *MaTAR20* ASO cohort presented macro-metastatic lesions

(Figure 4F). We went on to examine the histo-pathology of the four lungs that developed

metastasis by H&E staining and discovered that metastases in the *MaTAR20* ASO

treated group showed cystic, differentiated nodules compared to the solid masses

detected in the scASO cohort (Figure 4G). We suggest that metastatic nodules in animals

that received the *MaTAR20* cET may have developed prior to treatment start, and that

the cystic phenotype is a consequence of systemic *MaTAR20* knockdown.

To independently validate our animal studies, we performed a second *in vivo*

experiment using *MaTAR20* MOE ASO1 and injected it into MMTV-PyMT (FVB) mice

(Supplementary Figure S4). Similar to the cET experiments in C57/Bl6, we observed

reduced tumor growth for the *MaTAR20* ASO group compared to the scASO group from

around week 4 of treatment (Supplementary Figure S4A-B). H&E staining of tumor

sections revealed that animals in the *MaTAR20* MOE group also developed more necrotic

tumors compared to control animals (Supplementary Figure S4C). In addition, we

observed a significant reduction (70%) of macro-metastatic nodules in the lungs of mice

receiving *MaTAR20* MOE ASO1 (Supplementary Figure S4D), confirming the trend of

reduced metastasis upon *MaTAR20* knockdown. In summary, our *in vivo* experiments

show that systemic reduction of *MaTAR20* using ASOs leads to delayed mammary tumor

onset, reduced tumor growth, increased tumor necrosis and reduced metastasis.

MaTAR20 expression level correlates with VEGF expression

We set out to identify the molecular mechanism by which *MaTAR20* impacts tumor

cell growth and invasion. As our RNA-FISH and subcellular fractionation assays indicated

that *MaTAR20* is a nuclear-retained lncRNA (Figure 1), we hypothesized that it may act

by impacting gene expression. First, we performed a computational co-expression

analysis using lncRNA2function (36). Similar approaches have been previously applied

successfully to identify the molecular role of other lncRNAs (37). Notably, we observed a

strong enrichment for pathways involved in angiogenesis, with the top 7 most enriched

pathways being related to vascular endothelial growth factor (VEGF) signaling (Figure

5A), including "Signaling by VEGF", "VEGF hypoxia and angiogenesis" and "Sorafenib

Pharmacodynamics", a multi-kinase inhibitor that has been shown to act on VEGF

receptor (VEGFR) (38). Due to this striking ontology enrichment, we set out to

experimentally assess if *MaTAR20* and VEGF expression correlate. We determined by

qRT-PCR that Vegf-A is reduced by 15% and Vegf-B by 35% in tumors of MMTV-PyMT

mice that received the *MaTAR20* cET ASO compared to the scASO control group (Figure

5B).

We used RNA extracted from whole tumors for the qRT-PCR validations described

above. While MMTV-PyMT tumors generally contain a high percentage of cancer cells

(Figure 4E, Supplementary Figure S4C), qRT-PCR from whole tumors cannot distinguish

between expression changes in tumor or stromal cells. To investigate whether VEGF

expression was reduced in the same cell type as *MaTAR20*, we measured VEGF

expression in 4T1 cells with successful *MaTAR20* KD (Figure 3A/B). Indeed, we observed

a 50% reduction (p-value: 0.05) of Vegf-B (Figure 5C), while Vegf-A was not expressed

at levels sufficient for reliable qPCR detection in these cells (Ct > 30). We also tested

Vegf-B expression in our three knockout cell lines (Figure 2), and observed a statistically

significant, 40-60% down-regulation of Vegf-B compared to control cells (Figure 5D).

Upon ectopic expression of *MaTAR20* in the KO cells, Vegf-B levels were restored (60%

up-regulation, Supplementary Figure S5A). Based on convergent results obtained from

four independent experiments (ASO KD *in vitro*, *in vivo*, CRISPR/Cas9 promoter deletions

and rescue assays), we conclude that down-regulation of *MaTAR20* correlates with

reduced VEGF expression levels.

MaTAR20 loss leads to alterations in cancer signaling and adhesion pathways

To identify how *MaTAR20* impacts Vegf-B signaling, we further investigated the

molecular mechanism by which the lncRNA acts. First, we performed differential RNA-

seq, comparing *MaTAR20* KO to control cells. A total of 223 genes were differentially

expressed ($p < 0.05$), with 129 genes down- and 94 genes up-regulated in the KO cells

(Figure 6A, Supplementary Table S2). Interestingly, one of the most up-regulated genes

was *Vascular endothelial growth inhibitor* (*Vegi*), also known as *Tumor necrosis factor*

(TNF)-like cytokine 1A (TL1A)/TNF superfamily member 15 (*Tnfsf15*). In agreement with

our co-expression analysis (Figure 5A), we observed an overall dysregulation of the

VEGF signaling pathway and of hypoxia-inducible factor 1 (HIF-1) signaling, another

pathway essential for angiogenesis (39) (Supplementary Table S3). In addition, pathways

involved in cancer signaling such as the MAPK, PI3K-Akt and TNF axes were altered, as

were several cell adhesion pathways, including "Focal adhesion" and "Tight junctions"

(Supplementary Table S3). Several of these pathways were previously identified in our

co-expression analysis as well (Figure 5A). Differential expression of important cell

signaling genes likely contributes to the altered phenotypes we observed upon *MaTAR20*

loss, such as reduced organoid branching and tumor cell growth. Changes in cell

adhesion pathways may explain the enhanced adhesion phenotype observed in

MaTAR20 KO cells (Figure 2E), along with reduced invasion and metastatic burden upon

KD of the lncRNA.

MaTAR20* regulates gene expression by direct binding to *Tnfsf15

RNA-seq identifies all differentially expressed genes upon *MaTAR20* loss, both

effects due to direct binding and regulation of the gene by *MaTAR20*, as well as any

secondary or tertiary downstream effects. To hone in on direct targets of *MaTAR20*, we

performed Chromatin Isolation by RNA Purification (ChIRP, (40)) using two separate

pools of nine tiling oligonucleotides each. Both pools were able to enrich *MaTAR20*, with

pool 2 showing overall better pull-down performance (Figure 6B). The difference in

enrichment could be due to one or several oligonucleotides in pool 1 being obstructed

from binding to *MaTAR20*, caused by secondary structures or competition with putative

protein interaction partners. A negative control pull-down reaction for the mRNA

cyclophilin B (*CycloB*) was performed as well to control for unspecific binding of nucleic

acids to biotinylated oligonucleotides and/or streptavidin beads (Figure 6B).

We performed three independent replicates of ChIRP-seq with both probe pools

and *CycloB* negative control probes, and compared genes bound to *MaTAR20* in both

pools. The highest genomic peak score in both pools was to the *MaTAR20* locus, which

we have observed for other lncRNAs such as *MaTAR25* in the past (13). Both pool 1 and

pool 2 were enriched for genes involved in cancer signaling and adhesion pathways that

were also identified in our RNA-seq analysis, such as PI3K-Akt and MAPK signaling,

"Focal adhesion" and "ECM-receptor interactions" (Supplementary Table S4). The more

efficient pool 2 pull-down was also enriched for VEGF and HIF-1 signaling pathways

(Supplementary Table S4). On an individual gene level, we assigned the identified peaks

to the nearest gene and were able to identify 574 genes bound by *MaTAR20* in pool1 and

2,514 genes in pool 2 (excluding *MaTAR20* itself, as well as all genes that were also

found to bind to the negative control *CycloB* probe set) (Supplementary Table S5), with

an overlap of 182 target genes bound by *MaTAR20* in both probe pools (Figure 6C,

Supplementary Table S5).

Comparing RNA-seq and ChIRP-seq results, two genes were identified to be direct

binding partners of *MaTAR20* and also differentially expressed in *MaTAR20* KO cells

(Figure 6D): *Tnfsf15* and *Rab27b*. While *Tnfsf15* was significantly up-regulated upon

MaTAR20 loss, *Rab27b* was repressed (Figure 6E). We performed qRT-PCR at the

Tnfsf15 gene locus to validate direct binding of the lncRNA (Figure 6F). Our *MaTAR20*

ChIRP-seq data identified two putative lncRNA binding regions, one about 5 kb

downstream and the second about 15 kb downstream of the *Tnfsf15* gene. We designed

two primer pairs for each region (p1 and p2 for -15 kb, p3 and p4 for -5 kb), and a primer

pair for a negative control region in between the two putative binding sites, at about -10

kb. Cyclo B probes served as a negative control for all tested regions, representing

background noise. Our results indicate that both *MaTAR20* probe pools were able to

specifically detect the lncRNA at the -5 kb site, while only the more efficient pool2 could

also enrich for MaTAR20 at the -15 kb region. This may indicate higher levels of

MaTAR20 binding to the *Tnfsf15* -5 kb site, which likely represents a *Tnfsf15* enhancer

element. Overall, our results confirm that *MaTAR20* can regulate gene expression by

directly binding to target genes such as *Tnfsf15*.

Based on our findings, we conclude that *MaTAR20* reduces or inhibits the

expression of *Tnfsf15* by directly binding to its genomic locus. Tnfsf15 is a cytokine that

has been described to inhibit VEGF expression (41) and, more generally, VEGF driven

angiogenesis (42). Therefore, we hypothesize that loss of *MaTAR20* leads to a down-

regulation of Vegf and related angiogenesis pathways via up-regulation of Tnfsf15 (Figure

7). As a consequence of the altered Tnfsf15 / Vegf axis, less blood vessels may sprout,

and tumor angiogenesis may be inhibited in tumor bearing mice treated with *MaTAR20*

ASOs. The proposed molecular mechanism agrees with the observed phenotype:

delayed tumor onset, smaller tumors and increased necrosis, combined with lower

metastatic burden in the lungs.

Discussion

Here, we investigated *MaTAR20*, a lncRNA expressed at high levels in mammary

tumors. Loss of *MaTAR20* by either ASO-mediated knockdown or CRISPR/Cas9 genome

editing in mammary tumor cells leads to reduced proliferation, invasion and organoid

branching *in vitro*. In mouse models of BC, treatment with ASOs targeting *MaTAR20*

causes delayed tumor onset, reduced tumor growth and overall reduced metastatic

burden to the lungs. We were able to confirm the observed phenotype using different

ASO sequences, different ASO chemistries (MOE and cET), and in different mouse

background strains (FVB and C57/Bl6). Our results indicate that *MaTAR20* is a new driver

of tumor growth and metastasis and represents a promising new therapeutic target in BC.

We propose that ASOs targeting *MaTAR20* could be a viable new adjunct therapy, as we

obtained highly efficient lncRNA knockdown in tumors upon subcutaneous delivery. We

did not observe any adverse reactions in other mouse tissues or the animal overall,

potentially because *MaTAR20* is expressed at highest levels in the tumor, restricting the

effect of *MaTAR20* KD mostly to tumor tissue. Detailed studies of the human orthologue

hMaTAR20 (7) will be essential to investigate the potential of the lncRNA as new

therapeutic target in breast cancer.

Our data indicate that ASO-mediated reduction of *MaTAR20* may also impact

already existing metastases in the lungs. While systemic delivery of a cancer drug is

essential for clinical use, it creates challenges in distinguishing if the observed phenotype

is due to *MaTAR20* reduction in cancer cells or in the surrounding stroma. *MaTAR20* is

absent or expressed at very low levels in most tissues, however, it is present at moderate

levels in the lungs. Thus, the observed phenotype of cystic, differentiated metastatic

nodules in the lungs may be due to *MaTAR20* reduction in the metastatic cancer cells,

the lung tissue, or both. Future investigation of the impact of *MaTAR20* on already existing

metastatic nodules may be of particular value, as patients with advanced disease would

benefit from a therapeutic approach that is able to attack and/or prevent the formation of

secondary tumors.

Our molecular data indicates that the nuclear-retained lncRNA *MaTAR20* drives

tumor growth and metastasis by impacting the expression of important cancer signaling

pathways and cell adhesion molecules. These findings are reflected in the observed

reduction of cell proliferation, invasive potential and formation of cell protrusions in 2D

and 3D cell culture systems. One particular pathway that stood out across all our analyses

is VEGF signaling and angiogenesis, seemingly centering on Vegf-B in our system.

VEGF-B produced by cancer cells has been described to lead to leaky vascular networks,

which in turn enables the tumor to invade its surrounding tissue with high efficiency (43).

In addition, high VEGF-B levels were associated with poor prognosis (43). Inhibition of

tumor-specific VEGF-B has been suggested as an interesting approach to inhibit cancer

progression and metastasis, with VEGF-B knockdown leading to reduced pulmonary

metastasis in a mouse model of melanoma (43). As *MaTAR20* also leads to reduced lung

metastatic nodules, we hypothesize that it may do so by regulating Vegf-B expression. In

addition, non-angiogenic functions of VEGF-B have been described previously in the

context of invasive BC, which could also play a role here (44).

Our ChIRP assays indicate that *MaTAR20* does not directly bind to the *Vegf-B*

gene, but to a number of other angiogenesis-related genes such as *angiopoietin 2*, among

many others. Of the 182 genes identified by ChIRP-seq analysis, two also show altered

expression in *MaTAR20* KO cells: *Tnfsf15* and *Rab27b*. Tnfsf15 is a cytokine usually

expressed in established vasculature but down-regulated in cancer lesions to allow for

neovascularization (42,45,46). Vegf and Tnfsf15 serve opposing functions, with carefully

balanced expression modulation of both factors playing key roles in vascular and immune

homeostasis. We suggest that *MaTAR20* regulates the balance of the Tnfsf15/Vegf axis

by directly binding to the Tnfsf15 locus and repressing the gene during tumor progression

(Figure 7). Reduced Tnfsf15 levels allow the tumor to recruit blood vessels, which is

essential for sustained tumor growth and enables intravasation of cancer cells into the

circulation, leading to metastasis (Figure 7A). In the absence of *MaTAR20*, we

hypothesize that Tnfsf15 expression will be re-activated, Vegf-B levels decrease, and

neovascularization of the tumor is inhibited (Figure 7B). In BC, high levels of Tnfsf15 in

clinical samples were associated with disease-free survival and overall better prognosis

(47). This further solidifies our hypothesis that tumor-specific down-regulation of Vegf-B

and up-regulation of Tnfsf15 via ASO-mediated *MaTAR20* knockdown could be a

promising therapeutic approach. As the lncRNA is expressed in high levels only in cancer

lesions, we suggest that consequences of *MaTAR20* KD, including reduced

neovascularization, are restricted to tumor tissue as well, representing an attractive new

approach of tumor-specific targeting of Vegf.

In addition to regulating angiogenesis, Tnfsf15 has also been described to be

involved in dendritic cell maturation and T-cell co-activation (46). Future studies are

required to conclude if loss of *MaTAR20* also impacts immune cells in a cancer context,

or if the observed phenotype is in fact driven by reduced vascularization. Further

investigation is also required into the second identified *MaTAR20* target gene, Rab27b.

The small secretory GTPase has been described to control vesicle exocytosis and to

deliver pro-invasive growth regulators into the tumor microenvironment (48), and has

recently been associated with VEGF signaling in cancer as well (49). Rab27b has been

reported to promote proliferation and invasiveness of ER+ BC cells *in vitro* and *in vivo*,

and has been associated with lymph node metastasis and differentiation grade in ER+

human tumors (50). We observed that loss of *MaTAR20* leads to reduced Rab27b

expression, which may further contribute to the decrease in proliferative and invasive

potential we observed here. Future studies will elucidate the detailed regulatory

mechanism of Rab27b by *MaTAR20*.

Acknowledgements

We thank members of the Spector and Diermeier labs for critical discussions and advice

throughout the course of this study. We thank Dr. Mikala Egeblad (CSHL) for providing

MMTV-PyMT mice and Dr. Fred R. Miller (Wayne State University) for providing 4T1

cells. We acknowledge the CSHL Cancer Center Shared Resources (Microscopy, Flow

Cytometry, Animal, Histology, and Next-Gen Sequencing) for services and technical

expertise (NCI 2P3OCA45508). We thank David Jackson (CSHL) and Zachary Lippman

(CSHL) for facilitating access to the SEM instruments. We would also like to thank

Bronwyn Carlisle (Department of Biochemistry, University of Otago) for her assistance

with the *MaTAR20* model illustration (Figure 7).

References

1. Siegel RL, Miller KD, Fuchs HE, Jemal A. Cancer Statistics, 2021. CA: A Cancer Journal for Clinicians. American Cancer Society; 2021;71:7–33.

2. Gupta RA, Shah N, Wang KC, Kim J, Horlings HM, Wong DJ, et al. Long non-coding RNA HOTAIR reprograms chromatin state to promote cancer metastasis. Nature. 2010;464:1071–6.

3. Prensner JR, Iyer MK, Balbin OA, Dhanasekaran SM, Cao Q, Brenner JC, et al. Transcriptome sequencing across a prostate cancer cohort identifies PCAT-1, an unannotated lincRNA implicated in disease progression. Nat Biotechnol. 2011;29:742–9.

4. Tseng Y-Y, Moriarity BS, Gong W, Akiyama R, Tiwari A, Kawakami H, et al. PVT1 dependence in cancer with MYC copy-number increase. Nature. 2014.

5. Xing Z, Lin A, Li C, Liang K, Wang S, Liu Y, et al. lncRNA directs cooperative epigenetic regulation downstream of chemokine signals. Cell. 2014;159:1110–25.

6. Arun G, Diermeier S, Akerman M, Chang K-C, Wilkinson JE, Hearn S, et al. Differentiation of mammary tumors and reduction in metastasis upon Malat1 lncRNA loss. Genes & Development. 2016;30:34–51.

7. Diermeier SD, Chang K-C, Freier SM, Song J, Demerdash El O, Krasnitz A, et al. Mammary Tumor-Associated RNAs Impact Tumor Cell Proliferation, Invasion, and Migration. Cell Reports. 2016;17:261–74.

8. Chakravarty D, Sboner A, Nair SS, Giannopoulou E, Li R, Hennig S, et al. The oestrogen receptor alpha-regulated lncRNA NEAT1 is a critical modulator of prostate cancer. Nature Communications. Nature Publishing Group; 2014;5:5383–16.

9. Sang L-J, Ju H-Q, Liu G-P, Tian T, Ma G-L, Lu Y-X, et al. LncRNA CamK-A Regulates Ca2+-Signaling-Mediated Tumor Microenvironment Remodeling. Molecular Cell. Elsevier Inc; 2018;72:71–7.

10. Hu Q, Ye Y, Chan L-C, Li Y, Liang K, Lin A, et al. Oncogenic lncRNA downregulates cancer cell antigen presentation and intrinsic tumor suppression. Nature Immunology. Springer US; 2019;:1–24.

11. Luo H, Zhu G, Xu J, Lai Q, Yan B, Guo Y, et al. HOTTIP lncRNA Promotes Hematopoietic Stem Cell Self-Renewal Leading to AML-like Disease in Mice. Cancer Cell. Elsevier Inc; 2019;:1–24.

12. Silva-Fisher JM, Dang HX, White NM, Strand MS, Krasnick BA, Rozycki EB, et al. Long non-coding RNA RAMS11 promotes metastatic colorectal cancer progression. Nature Communications. 2020;11:2156.

13.   Chang K-C, Diermeier SD, Yu AT, Brine LD, Russo S, Bhatia S, et al. MaTAR25
lncRNA regulates the Tensin1 gene to impact breast cancer progression. Nature
Communications. Springer US; 2020;:1–19.

14.   Yan X, Hu Z, Feng Y, Hu X, Yuan J, Zhao SD, et al. Comprehensive Genomic
Characterization of Long Non-coding RNAs across Human Cancers. Cancer Cell.
Elsevier Inc; 2015;28:529–40.

15.   Ransohoff JD, Wei Y, Khavari PA. The functions and unique features of long
intergenic non-coding RNA. Nature Reviews Molecular Cell Biology. Nature
Publishing Group; 2018;19:143–57.

16.   Huarte M. The emerging role of lncRNAs in cancer. Nature Medicine.
2015;21:1253–61.

17.   Arun G, Diermeier SD, Spector DL. Therapeutic Targeting of Long Non-Coding
RNAs in Cancer. Trends Mol Med. 2018;24:257–77.

18.   Gutschner T, Hammerle M, Eissmann M, Hsu J, Kim Y, Hung G, et al. The
Noncoding RNA MALAT1 Is a Critical Regulator of the Metastasis Phenotype of
Lung Cancer Cells. Cancer Research. 2013;73:1180–9.

19.   Ewald AJ. Isolation of mouse mammary organoids for long-term time-lapse
imaging. Cold Spring Harb Protoc. 2013;2013:130–3.

20.   Park J, Wysocki RW, Amoozgar Z, Maiorino L, Fein MR, Jorns J, et al. Cancer
cells induce metastasis-supporting neutrophil extracellular DNA traps. Sci Transl
Med. 2016;8:361ra138.

21.   Nowak DG, Katsenelson KC, Watrud KE, Chen M, Mathew G, D'Andrea VD, et al.
The PHLPP2 phosphatase is a druggable driver of prostate cancer progression.
The Journal of Cell Biology. 2019;218:1943–57.

22.   Guy CT, Cardiff RD, Muller WJ. Induction of mammary tumors by expression of
polyomavirus middle T oncogene: a transgenic mouse model for metastatic
disease. Molecular and Cellular Biology. 1992;12:954–61.

23.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR:
ultrafast universal RNA-seq aligner. Bioinformatics. 2012;29:15–21.

24.   Anders S, Pyl PT, Huber W. HTSeq--a Python framework to work with high-
throughput sequencing data. Bioinformatics. 2015;31:166–9.

25.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biol. 2014;15:550.

26. Luo W, Friedman MS, Shedden K, Hankenson KD, Woolf PJ. GAGE: generally
applicable gene set enrichment for pathway analysis. BMC Bioinformatics.
2009;10:161.

27. Luo W, Brouwer C. Pathview: an R/Bioconductor package for pathway-based
data integration and visualization. Bioinformatics. 2013;29:1830–1.

28. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nature
Methods. 2012;9:357–9.

29. Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, et al. Simple
Combinations of Lineage-Determining Transcription Factors Prime cis-Regulatory
Elements Required for Macrophage and B Cell Identities. Molecular Cell. Elsevier
Inc; 2010;38:576–89.

30. Diermeier SD, Spector DL. Antisense Oligonucleotide-mediated Knockdown in
Mammary Tumor Organoids. BIO-PROTOCOL. 2017;7.

31. Tuveson D, Clevers H. Cancer modeling meets human organoid technology.
Science. American Association for the Advancement of Science; 2019;364:952–5.

32. Shamir ER, Ewald AJ. Three-dimensional organotypic culture: experimental
models of mammalian biology and disease. Nature Reviews Molecular Cell
Biology. Nature Publishing Group; 2014;15:647–64.

33. Aslakson CJ, Miller FR. Selective events in the metastatic process defined by
analysis of the sequential dissemination of subpopulations of a mouse mammary
tumor. Cancer Research. 1992;52:1399–405.

34. Pallan PS, Allerson CR, Berdeja A, Seth PP, Swayze EE, Prakash TP, et al.
Structure and nuclease resistance of 2′,4′-constrained 2′-O-methoxyethyl (cMOE)
and 2′-O-ethyl (cEt) modified DNAs. Chem Commun. The Royal Society of
Chemistry; 2012;48:8195–10.

35. Shen X, Corey DR. Chemistry, mechanism and clinical status of antisense
oligonucleotides and duplex RNAs. Nucleic Acids Res. 2018;46:1584–600.

36. Jiang Q, Ma R, Wang J, Wu X, Jin S, Peng J, et al. LncRNA2Function: a
comprehensive resource for functional investigation of human lncRNAs based on
RNA-seq data. BMC Genomics. 2015;16 Suppl 3:S2.

37. Guttman M, Amit I, Garber M, French C, Lin MF, Feldser D, et al. Chromatin
signature reveals over a thousand highly conserved large non-coding RNAs in
mammals. Nature [Internet]. Nature Publishing Group; 2009;457:223–7. Available
from: http://www.nature.com/doifinder/10.1038/nature07672

38. Wilhelm SM, Adnane L, Newell P, Villanueva A, Llovet JM, Lynch M. Preclinical
overview of sorafenib, a multikinase inhibitor that targets both Raf and VEGF and
PDGF receptor tyrosine kinase signaling. Mol Cancer Ther. 2008;7:3129–40.

39. Semenza GL. Targeting HIF-1 for cancer therapy. Nat Rev Cancer. Nature
Publishing Group; 2003;3:721–32.

40. Chu C, Qu K, Zhong FL, Artandi SE, Chang HY. Genomic Maps of Long
Noncoding RNA Occupancy Reveal Principles of RNA-Chromatin Interactions.
Molecular Cell. Elsevier Inc; 2011;44:667–78.

41. Zhang K, Cai H-X, Gao S, Yang G-L, Deng H-T, Xu G-C, et al. TNFSF15
suppresses VEGF production in endothelial cells by stimulating miR-29b
expression via activation of JNK-GATA3 signals. Oncotarget. Impact Journals;
2016;7:69436–49.

42. Yang G-L, Li L-Y. Counterbalance: modulation of VEGF/VEGFR activities by
TNFSF15. Signal Transduction and Targeted Therapy. Springer US; 2018;:1–7.

43. Yang X, Zhang Y, Hosaka K, Andersson P, Wang J, Tholander F, et al. VEGF-B
promotes cancer metastasis through a VEGF-A–independent mechanism and
serves as a marker of poor prognosis for cancer patients. Proceedings of the
National Academy of Sciences. 2015;112:E2900–9.

44. Gunningham SP, Currie MJ, Han C, Robinson BA, Scott PA, Harris AL, et al.
VEGF-B expression in human primary breast cancers is associated with lymph
node metastasis but not angiogenesis. J Pathol. John Wiley & Sons, Ltd;
2001;193:325–32.

45. Deng W, Gu X, Lu Y, Gu C, Zheng Y, Zhang Z, et al. Down-modulation of
TNFSF15 in ovarian cancer by VEGF and MCP-1 is a pre-requisite for tumor
neovascularization. Angiogenesis. Springer Netherlands; 2011;15:71–85.

46. Zhang Z, Li L-Y. TNFSF15 Modulates Neovascularization and Inflammation.
Cancer Microenvironment. 2012;5:237–47.

47. Parr C, Gan CH, Watkins G, Jiang WG. Reduced vascular endothelial growth
inhibitor (VEGI) expression is associated with poor prognosis in breast cancer
patients. Angiogenesis. 2006;9:73–81.

48. Hendrix A, Maynard D, Pauwels P, Braems G, Denys H, Van den Broecke R, et
al. Effect of the Secretory Small GTPase Rab27B on Breast Cancer Growth,
Invasion, and Metastasis. J Natl Cancer Inst. 2010;102:866–80.

49. Tsuruda M, Yoshino H, Okamura S, Kuroshima K, Osako Y, Sakaguchi T, et al.
Oncogenic effects of RAB27B through exosome independent function in renal cell
carcinoma including sunitinib-resistant. Dahiya R, editor. PLoS ONE.
2020;15:e0232545–20.

50.   Li Z, Fang R, Fang J, He S, Liu T. Functional implications of Rab27 GTPases in
708   Cancer. Cell Communication and Signaling; 2018;:1–8.

Figure 1

Figure 1: *MaTAR20* is a tumor-specific, nuclear-retained lncRNA

A: Schematic overview of the *MaTAR20*/*Gm13387* locus (UCSC Browser, mm10) GENCODE vM18 annotation in green.

B: Abundance (FPKM) of *MaTAR20* in ENCODE mouse tissue and HER2+ mammary tumors.

C: Single molecule RNA FISH of MaTAR20 in primary MMTV-PyMT cells. Top panel: merge, middle panel: MaTAR20 (red), bottom panel: DAPI (blue).

D: Cellular fractionation assay followed by qRT-PCR.

E: ASO-mediated knockdown of MaTAR20 in primary MMTV-PyMT organoids. Two specific ASOs were tested for MaTAR20, and a scASO was used as negative control. Bars denote the mean of at least two biological replicates +/- SD.

F: Quantification of organoid branching upon MaTAR20 knockdown. Bars denote the mean of at least three biological replicates +/- SD.

G: Exemplarily images of organoids quantified in F.

H: Exemplarily SEM micrograph of organoids quantified in F.

Figure 2

Figure 2: MaTAR20 KO cells show reduced proliferation and cell adhesion
A: Schematic overview of the knockout strategy. Guide RNAs (gRNAs) are indicated in red, +/-
indicate genomic position relative to the MaTAR20 transcription start site.
B: qRT-PCR to determine relative MaTAR20 expression in MaTAR20 promoter deletion cell lines.
KO = promoter deletion of 748 bp (combination of gRNAs "+12" and "-736"), Ren = negative control,
integration of Cas9 and a gRNA targeting Renilla luciferase. Bars denote the mean of six biological
replicates +/- SD. Statistical significance was determined with a two-tailed Student's t-test; *** p <
0.001.
C: Cell proliferation assay. Number of cells determined after 48 h. Bars denote the mean of three
replicates +/- SD. Statistical significance was determined with a two-tailed Student's t-test; * p <
0.05, ** p < 0.01.
D: Cell cycle profiles of NST-DAPI stained cells, comparing Ren control cells to KO1 and KO2.
E: Scanning electron micrograph of control (Ren) or CRISPR/Cas9-modified cells (KO). Scale as
indicated.

Figure 3

Figure 3: cET ASO-mediated knockdown of MaTAR20 in 4T1 cells lead to reduced cell proliferation and invasion

Bars denote the mean of three biological replicates +/- SD. Statistical significance was determined with a two-tailed Student's t-test; * p < 0.05, ** p < 0.01, *** p < 0.001.

A: qRT-PCR to determine relative MaTAR20 expression. 24 h post treatments with cET ASOs (concentrations as indicated).

B: qRT-PCR to determine relative MaTAR20 expression. 72 h post treatments with 0.5 μM cET ASOs.

C: Proliferation assay. Cells were counted 24 h and 72 h post treatment with 0.5 μM cET ASOs.

D: Invasion assay. Fluorescence units 24 h post treatment with 0.5 μM cET ASOs.

Figure 4

Figure 4: cET ASO-mediated knockdown of *MaTAR20 in vivo* lead to reduced tumor growth and metastatic burden

A: MMTV-PyMT (C57/Bl6) mice were treated for 8 weeks with cET ASOs (100 mg/kg/week), either scASO (cET) or *MaTAR20* cET1. Bars / lines denote the mean of biological replicates +/- SEM. N= 33 tumors from 7 mice for scASO (cET); n = 26 tumors from 8 mice for *MaTAR20* cET1. Statistical significance was determined with a two-tailed Student's t-test; * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

B: qRT-PCR to determine relative *MaTAR20* expression in tumors.

C: Average number of tumors per mouse in each treatment group.

D: Average tumor burden in mm^3 per week in each treatment group.

E: Hematoxylin and Eosin staining of tumor sections. Top left: scASO (cET) treated tumor (whole tumor). Top right: scASO (cET) treated tumor, 10x. Bottom left: *MaTAR20* cET1 treated tumor (whole tumor). Bottom right: *MaTAR20* cET1 treated tumor, 10x.

F: Quantification of macro-metastatic nodules, displayed is the average number per lung.

G: Hematoxylin and Eosin staining of fixed lung sections.

Figure 5

Figure 5: MaTAR20 impacts VEGF expression
A: Co-expression analysis to identify MaTAR20 associated pathways. The 15 most significant pathways are shown.
B: qRT-PCR to determine the relative Vegf-A and Vegf-B expression in tumors. MMTV-PyMT mice were treated for 8 weeks with cET ASOs (100 mg/kg/week), either scASO (cET) or MaTAR20 cET1. Bars denote the mean of biological replicates +/- SEM. N= 23 tumors from 7 mice for scASO (cET); n = 12 tumors from 5 mice for MaTAR20 cET1.
C: qRT-PCR to determine the relative Vegf-B expression in 4T1 cells. 72 h post treatments with 0.5 µM cET ASOs. Bars denote the mean of three biological replicates +/- SD. Statistical significance was determined with a two-tailed Student's t-test; * p < 0.05.
D: qRT-PCR to determine the relative Vegf-B expression in MaTAR20 KO cells. Bars denote the mean of two biological replicates +/- SD. Statistical significance was determined with a two-tailed Student's t-test; * p < 0.05, ** p < 0.01, *** p < 0.001.

Figure 6

Figure 6: RNA-seq and ChIRP-seq identify *Tnfsf15* as a direct *MaTAR20* target gene

A: RNA-seq. Heatmap of significantly differentially expressed genes comparing *MaTAR20* KO and Ren cells. N1-3 indicate three independent biological replicates.

B: ChIRP-seq. qRT-PCR to determine the efficiency of the ChIRP assay. Both *MaTAR20* probe pools enrich *MaTAR20*, with Pool 2 being more efficient. "Cyclo B" indicates a separate pull-down experiment with oligonucleotides specific for the *cyclophilin B* mRNA, which was used as a specificity control. Percentage of MaTAR20 or CycloB pull-down in relation to input is shown. Bars denote the average of at least 3 replicates +/- SD.

C: ChIRP-seq. Venn diagram illustrating the overlap of ChIRP-seq performed with Pool1 vs Pool2 probes. Genes also identified in the negative control (Cyclo B) pull-down were removed from the target lists. 183 genes overlap between both ChIRP probe pools, with Pool1 enriching more genes than Pool 2 overall. All ChIRP experiments were performed in three independent replicates.

D: Venn diagram illustrating the overlap of the 182 target genes identified by ChIRP-seq and the 223 differentially expressed genes according to RNA-seq. 2 genes are both direct genomic binding partners of *MaTAR20* and are differentially expressed upon *MaTAR20* loss.

E: RNA-seq. Expression of Tnfsf15 and Rab27b, comparing the mean log2 fold-change in the two *MaTAR20* KO cell lines ("MaTAR20 KO") to the mean log2 fold-change of the Ren negative control cell lines ("Control").

F: qRT-PCR to validate direct *MaTAR20* binding to the *Tnfsf15* locus. Four tiling primer pairs were designed to two putative *MaTAR20* binding regions in proximity to the *Tnfsf15* gene. A negative control region, located between p1/p2 and p3/p4, shows no *MaTAR20* enrichment. CycloB probes control for unspecific binding at all tested sites. Bars denote the mean of at least two biological replicates +/- SD.

Figure 7

Figure 7: MaTAR20 knockdown leads to smaller, necrotic tumors and reduced metastasis
A: During cancer progression, tumors secret pro-angiogenic factors to recruit blood vessels. Increased vasculature allows for steady tumor growth while also increasing the chances for metastatic spread.
B: In the absence of MaTAR20, tumor growth is slowed down, tumor necrosis increases, and metastasis is reduced. We hypothesize that this could be due to the observed down-regulation of Vegf-B, which may in part be caused by direct binding of MaTAR20 to Tnfsf15. Additional players such as protein binding partners may be of importance and will be elucidated further in future studies.

Figure 7

A

Figure S1:
A: qRT-PCR to determine relative MaTAR20 expression in MMTV-PyMT tumors of different size.
"Early" tumors: <200 mm3, n=7; "middle" tumors: 300 - 1000 mm3, n=4; "late" tumors: >1500 mm3,

n=3.

Figure S2

Figure S2:

A: qRT-PCR to determine the relative *MaTAR20* expression in mouse mammary tumor cell lines. Bars denote the mean of three biological replicates +/- SD.

B: qRT-PCR to determine the relative expression of individual MaTAR20 isoforms in *MaTAR20* promoter deletion cell lines. KO = promoter deletion of 748 bp (combination of gRNAs "+12" and "-736"), Ren = negative control, integration of Cas9 and a gRNA targeting *Renilla* luciferase. Bars denote the mean of two biological replicates +/- SD.

C: Correlation of *MaTAR20* expression (all isoforms, based on six biological replicates, Figure 2B) and Cas9 expression, as determined by qRT-PCR in three biological replicates.

Figure S3

Figure S3:
A: Cell proliferation assay. Number of cells determined after 48 h. Statistical significance was determined comparing KO cells transfected with a MaTAR20 Tx1 (KO+MaTAR20) to KO cells transfected with an empty backbone (KO+bb) and untransfected KO cells. Control Ren cells are shown for reference. Error bars denote SD, and a two-tailed Student's t-test was performed comparing KO+bb to KO+MaTAR20; * p < 0.05.
B: Cell proliferation assay comparing MaTAR20 isoform 1 and isoform 2 after 24 and 48 h. Cell counts normalized to KO cells transfected with empty vector (bb). Bars denote the mean of at least two biological replicates +/- SD. Statistical significance was determined with a two-tailed Student's t-test comparing KO+bb to KO+MaTAR20; * p < 0.05, ** p < 0.01.

Figure S4

Figure S4:
A: MMTV-PyMT (FVB) mice were treated for 6 weeks with MOE ASOs (150 mg/kg/week), either scASO or MaTAR20 ASO1. Bars / lines denote the mean of biological replicates +/- SEM.
B: Average tumor burden in mm3 per week in each treatment group, tumor size 100-200 mm3, n=5.
C: Hematoxylin and Eosin staining of tumor sections. Top: scASO treated tumor. Bottom: MaTAR20 ASO1 treated tumor.
D: Quantification of macro-metastatic nodules, displayed is the average number per lung. Statistical significance was determined with a two-tailed Student's t-test; * p < 0.05. Insets show exemplary images of lungs from scASO and ASO1 treated animals.

Figure S5:
A: qRT-PCR to determine the relative Vegf-B expression in 4T1 MaTAR20 KO cells. Cells were transfected either with MaTAR20 Tx1 (KO+MaTAR20) or an empty backbone (KO+bb). Bars denote the mean of at least three biological replicates +/- SD. Statistical significance was determined with a two-tailed Student's t-test; * $p < 0.05$