Connectivity and Synaptic Physiology in the Mouse and Human Neocortex

Luke Campagnola[*1], Stephanie C Seeman[*1], Thomas Chartrand[1], Lisa Kim[1], Alex Hoggarth[1], Clare Gamlin[1], Shinya Ito[1], Jessica Trinh[1], Pasha Davoudian[1], Cristina Radaelli[1], Mean-Hwan Kim[1], Travis Hage[1], Thomas Braun[2], Lauren Alfiler[1], Rachel Dalley[1], Grace Williams[1], Rachael Larsen[1], Corinne Teeter[1], Tanya L. Daigle[1], Kyla Berry[1], Nadia Dotson[1], Rachel Enstrom[1], Melissa Gorham[1], Madie Hupp[1], Samuel Dingman Lee[1], Kiet Ngo[1], Rusty Nicovich[1], Lydia Potekhina[1], Shea Ransford[1], Amanda Gary[1], Jeff Goldy[1], Delissa McMillen[1], Trangthanh Pham[1], Michael Tieu[1], La'Akea Siverts[1], Miranda Walker[1], Colin Farrell[1], Martin Schroedter[1], Cliff Slaughterbeck[1], Charles Cobb[3], Richard Ellenbogen[4], Ryder P Gwinn[5], C. Dirk Keene[6], Andrew L Ko[4,7], Jeffrey G Ojemann[4,7], Daniel L Silbergeld[4], Daniel Carey[1], Tamara Casper[1], Kirsten Crichton[1], Michael Clark[1], Nick Dee[1], Lauren Ellingwood[1], Jessica Gloe[1], Matthew Kroll[1], Josef Sulc[1], Herman Tung[1], Katherine Wadhwani[1], Krissy Brouner[1], Tom Egdorf[1], Michelle Maxwell[1], Medea McGraw[1], Christina Alice Pom[1], Augustin Ruiz[1], Jasmine Bomben[1], David Feng[1], Nika Hejazinia[1], Shu Shi[1], Aaron Szafer[1], Wayne Wakeman[1], John Phillips[1], Amy Bernard[1], Luke Esposito[1], Florence D D'Orazi[1], Susan Sunkin[1], Kimberly Smith[1], Bosiljka Tasic[1], Anton Arkhipov[1], Staci Sorensen[1], Ed Lein[1], Christof Koch[1], Gabe Murphy[1], Hongkui Zeng[1], Tim Jarsky[♔1]

1. Allen Institute for Brain Science, Seattle, WA, USA
2. Byte Physics (), Berlin, Germany
3. The Ben and Catherine Ivy Center for Advanced Brain Tumor Treatment, Swedish Neuroscience Institute, Seattle, USA
4. Department of Neurological Surgery, University of Washington, Seattle, WA, USA
5. Epilepsy Surgery and Functional Neurosurgery, Swedish Neuroscience Institute, Seattle, United States
6. Department of Pathology, University of Washington, Seattle, WA, USA
7. Regional Epilepsy Center at Harborview Medical Center, Seattle, WA, USA

* These authors contributed equally

♔ Correspondence: timj@alleninstitute.org

Abstract

We present a unique, extensive, public synaptic physiology dataset. The dataset contains over 20,000 neuron pairs probed with multipatch using standardized protocols to capture short-term dynamics. Recordings were made in the human temporal cortex and the adult mouse visual cortex. Our main purpose is to offer data and analyses that provide a more complete picture of the cortical microcircuit to the community. We also make several important findings that relate

connectivity and synaptic properties to the major cell subclasses and cortical layer via the development of novel analysis methods for quantifying connectivity, synapse properties, and synaptic dynamics. We find that excitatory synaptic dynamics depend strongly on the postsynaptic cell subclass, whereas inhibitory synaptic dynamics depend on the presynaptic cell subclass. Despite these associations, short-term synaptic plasticity is heterogeneous in most subclass to subclass connections. We also find that intralaminar connection probability exhibits a strong layer dependence. In human cortex, we find that excitatory synapses are highly reliable, recover rapidly, and are distinct from mouse excitatory synapses.

Introduction

For much of the history of electrophysiological recordings, cells have been categorized based on their most accessible properties--morphology, anatomy (location), intrinsic electrophysiology, and spiking behavior (DeFelipe and Fariñas, 1992; Group et al., 2008). The study of connectivity between these cell classes established "canonical" circuit diagrams describing major excitatory inputs and interlaminar pathways as well as intralaminar recurrent circuits between excitatory and inhibitory cells (Thomson and Bannister, 2003; Douglas and Martin, 2004; Markram et al., 2004). As the field advanced, more attention was given to subdividing cell classes based on their long range connectivity and molecular expression, which offer more tractable and repeatable identification of cell subclasses (Pfeffer et al., 2013; Harris and Shepherd, 2015; Jiang et al., 2015; Karnani et al., 2016; Tremblay et al., 2016; Cardin, 2018). The most commonly used transgenic markers map onto cell subclasses with distinct patterns of connectivity within the cortical circuit: the inhibitory markers parvalbumin (Pvalb), somatostatin (Sst), and vasoactive intestinal peptide (Vip) correspond to interneuron groups with distinct patterns of local connectivity, whereas excitatory markers such as Sim1, Tlx3, and Ntsr1 correspond to pyramidal cell groups with distinct localization and patterns of long range connectivity (Vélez-Fort et al., 2014; Kim et al., 2015). Although these cell subclasses can be further subdivided (Tasic et al., 2016; Hodge et al., 2019; Naka et al., 2019; Gouwens et al., 2020), tools that target these more refined transcriptomic types are not available yet. We describe synaptic physiology at the subclass level as a compromise between cell type resolution and experimental tractability.

The mouse is a compelling model for studying cortical microcircuitry because it affords a high degree of genetic and experimental accessibility. However, if we wish to understand our own cognitive abilities, it is necessary to clarify the human cortical microcircuit's unique features (Lourenço and Bacci, 2017). Access to postoperative tissue specimens has led to the identification of several differences between the mouse and human cortex. For example, it appears that human supragranular cortex has more excitatory cell types, some of which have intrinsic properties that vary with cortical depth (Berg et al., 2020). The synaptic physiology also appears to have some basic differences (Testa-Silva et al., 2014). Foremost among them is that monosynaptic excitatory connections are stronger and in some cases, strong enough to evoke an action potential in connected inhibitory interneurons (Molnár et al., 2008). Is mouse and

human synaptic physiology always different? The limited available synaptic dynamics data suggests not, as recurrent excitatory connections were found to have similar short-term depression in mouse and human cortex (Seeman et al., 2018). We include experiments from human tissue in our survey in order to identify unique and conserved features of the human cortical microcircuit as it relates to cell subclass.

Cortical synapses are dynamic, varying their response strength in ways that are highly stochastic and also modulated by the prior history of activity. Ongoing activity affects both the probability of vesicle release and the quantal response amplitude, causing a combination of short-term facilitation or depression (transient strengthening or weakening of the synaptic "weight"). The dynamical properties of synapses in the cortex are influenced by both the presynaptic and postsynaptic cell types (Reyes et al., 1998; Blackman et al., 2013; Larsen and Sjöström, 2015; Markram et al., 2015; Lefort and Petersen, 2017; Yang et al., 2021)and endow neuronal networks with essential sources of computational diversity (Abbott and Regehr, 2004; Llera-Montero et al., 2019). Using new analyses and models, we have characterized the stochasticity and short term plasticity of over 1000 synapses, offering a more comprehensive view of the diversity of cortical synapse types.

To maximize the value of our dataset as a public resource, we have taken a comprehensive approach including analyses of the strength and probability of chemical and electrical connections (and a subset of polysynaptic connections in human); the latency and kinetic properties of synaptic responses in both voltage- and current-clamp; modeling of quantal release and short-term plasticity; and cell classification features such as morphology, intrinsic physiology, transgenic markers, and cortical layer. In our analysis we have given extra attention to generate measurements that are appropriate for biophysical modeling and that can be more readily compared to future studies, something that has not been done by previous large-scale synaptic physiology studies (Pfeffer et al., 2013; Jiang et al., 2015; Karnani et al., 2016; Lefort and Petersen, 2017). We introduce analyses for making connection probability estimates robust and repeatable, for inferring synaptic quantal and dynamic properties, and for exploring relationships between synaptic properties. We leverage these analyses to compare the intralaminar circuit elements across layers and find that layer 2/3 has several unique features in human and mouse. We also identify previously undescribed circuit motifs that exist in multiple layers. For all circuit motifs, the short-term plasticity appears more heterogeneous than previously understood. Short-term plasticity is also dependent on the cell subclass; excitatory synaptic dynamics depend strongly on the subclass identity of the postsynaptic cell, whereas inhibitory synapses are influenced more by the presynaptic subclass. Additionally, we find that recurrent excitatory connectivity has a common set of features that limit their contribution to excitability. All acquisition software tools, data, analyses, figures, and source code are available for download (https://portal.brain-map.org/explore/connectivity/synaptic-physiology).

Results

We performed 1,853 experiments (Fig 1A) targeting Layer 2 (L2) to L6 of young adult mouse primary visual cortex (VISp; 1,645 experiments) and human fronto-temporal cortex from neurosurgical excised tissue (208 experiments). Each experiment selected up to eight neurons for simultaneous whole-cell patch-clamp (multipatch) recording in acute slices (Fig 1B). Trains of action potentials were elicited in each patched neuron, in turn, while all others were recorded for evidence of a postsynaptic response. We probed 22,833 potential connections (mouse: 20,287; human: 2,546) of which 1,663 were connected by chemical synapses, giving an overall connectivity rate of 7.3% (mouse: 1,463 (7.2%); human: 200 (7.9%)). Over 900,000 individual monosynaptic events were analyzed across all these connections. We observed 160 electrical synaptic connections among inhibitory neurons in mice and 1 in humans. Conversely, we found 69 putative polysynaptic responses among excitatory neurons in humans but none in mice.

Synaptic physiology pipeline

We established a pipeline for conducting large-scale studies in patch clamp electrophysiology focused on connectivity and synaptic properties among cells at the subclass level, delineated by cortical layer (Seeman et al., 2018). In mice we utilized transgenic animals which express unique reporters in two subclasses (Daigle et al., 2018) (Table S1), allowing us to investigate the connectivity across different subclasses in a single experiment. This also enabled us to target rare subclasses and aligns our dataset with related experimental and theoretical projects at the Allen Institute for Brain Science (Harris et al., 2019; Billeh et al., 2020; Gouwens et al., 2020; de Vries et al., 2020) as well as others that utilize similar transgenic animals. Six excitatory subclasses were layer or projection-class specific (Nr5a1 and Rorb, L4; Sim1 or mscRE4-FlpO AAV, L5 ET; Tlx3, L5 IT; Ntsr1, L6 CT) while three inhibitory subclasses (Pvalb, Sst, Vip) were assessed in all targeted layers (Fig 1A). Unlabeled pyramidal cells in human and mouse L2/3 were targeted based on somatic and dendritic trunk shape and confirmed via *post hoc* morphological analysis that verified the presence of dense dendritic spines (Fig 1C, Table S2). Human samples were procured via collaborations with area hospitals performing neurological resections in patients being treated for epilepsy (52 samples) or tumor (20 samples). The majority of human samples were from temporal cortex, with a small fraction from frontal and parietal cortex (Fig 1A).

Multipatch experiments (Fig 1B) were primarily conducted in current-clamp, with a subset of stimuli administered in voltage-clamp as well. Stimulus trains of eight pulses at frequencies ranging from 10 – 200 Hz probed the degree to which individual connections displayed short-term plasticity (STP). After a delay, four additional pulses were delivered to characterize how synapses recovered from STP. To assess how synapses respond to a more realistic firing pattern we also included a "mixed frequency" stimulus. Figure 1B (*Multipatch Experiment*) shows example raw data from two connections in a single experiment. In both connections, postsynaptic potentials (PSPs) are evident on single trials (grey) of the stimulus. The excitatory connection from cell 1 to cell 3 shows evidence of facilitation whereas the inhibitory connection

from cell 5 to cell 1 depresses. Fluorescent images taken during the experiment allowed us to target and verify recorded cells' reporter expression and assign a subclass based on that expression. Subsequent processing of biocytin and DAPI labeled images was used to determine the cortical layer and morphological properties (Fig 1C).

Synapses were detected manually with the aid of custom software tools (see below). Use of a machine classifier supplemented manual synapse detection (Seeman et al., 2018). The classifier results were cross-referenced with the manual results to identify potential false positive and negative results which underwent additional rounds of manual inspection. When a synapse was identified, the data were processed for quality control and grouped according to clamp mode and baseline potential for further analysis of excitatory (EPSP/C) and inhibitory (IPSP/C) connections (Fig 1B *Synapse Processing*). All postsynaptic responses within each group were aligned to the presynaptic spike, allowing us to overlay hundreds of responses and produce a robust average synaptic response (colored trace). From these averages we characterized the strength and kinetics of each synapse (Fig 1B *Synapse Analysis*). Synaptic dynamic properties were characterized from the statistics of individual response amplitudes. Long pulse stimuli were delivered to each recorded cell to characterize intrinsic properties (Fig 1C, *Intrinsic Ephys*) and to assess the presence of electrical coupling between cells. See the Methods for more detail on the pipeline workflow including processing, quality control, and analysis of synaptic responses.

Intralaminar connectivity in mouse VISp

Distance dependence of connectivity

The likelihood that two neurons are locally connected depends on multiple factors such as cell type, cortical region, species, and animal age. A longstanding goal has been to determine the governing principles of local circuit architecture (Markram, 1997; Yoshimura et al., 2005; Ko et al., 2011; Packer and Yuste, 2011; Pfeffer et al., 2013; Jiang et al., 2015; Pala and Petersen, 2015). It is difficult to make direct comparisons among these studies because the observed rate of connectivity depends on the intersomatic distances that are sampled (Perin et al., 2011; Levy and Reyes, 2012) We probed connectivity among cells up to ~200 μm apart (Fig 2A), but could not ensure that intersomatic distances were sampled equally across different connection elements (specific pre-post combinations). In order to make reliable comparisons across elements, we modeled the spatial profile of connectivity versus lateral somatic distance with a Gaussian (Levy and Reyes, 2012) and estimated the peak (p_{max}) and lateral spread of connection probability (sigma: σ) using maximum likelihood estimation (see Methods, Fig S2B). This analysis provides an estimate of the spatial profile of connectivity while being relatively robust to differences in the sampled intersomatic distribution. We initially classified cell pairs among the four combinations of excitatory and inhibitory cell class (E→E, E→I, I→E, I→I). These data were well fit by the Gaussian model (Fig 2A, solid red line) resulting in low peak connectivity among excitatory cells (~5%), moderate connectivity rates among inhibitory cells (~11%), and relatively higher connectivity across E-I cell classes (E→I 13%, I→E 15%). Differences in the lateral spread of connectivity between cell classes is generally understudied

and our results showed a longer σ for within-class connectivity compared to across-class connectivity. By simulating random connectivity experiments with a range of different σ values, we asked what the likelihood was that the σ values for these four groups could have come from the same distribution. This revealed only a 0.25% chance (Bonferroni corrected for multiple comparisons p=0.83%) that I→I and I→E connectivity profiles have identical σ values. No statistical significance was found across other group comparisons. Taken together, we chose to model connectivity for all elements with a fixed σ of 100 µm.

An overabundance of bidirectional connections relative to unidirectional connections can be evidence for connectivity rules that promote the formation of bidirectional connections (Song et al., 2005); however, a more likely explanation is that such measurements come from data that merges subtype-specific connectivity patterns with no preference for bidirectionality (Hoffmann and Triesch, 2017). We quantified the ratio of connected pairs with and without bidirectional connections and observed that reciprocal connections were 3 to 5 times more common than expected for a randomly connected network (red line) in E→E, E→I, I→E, and I→I connections (Fig 2A, bottom row). Further analysis of higher order connectivity motifs in our data and their impact on cortical computation is pursued in a parallel study (Dahmen et al., 2020).

Connection probability measurement; slicing artifacts and detection limits

In addition to the influence of intersomatic distance on connectivity (Holmgren et al., 2003; Perin et al., 2011; Levy and Reyes, 2012), artifacts of the *in vitro* slice preparation may also impact connectivity measurements; cutting of neuronal processes (axons and dendrites) that could have formed synapses draws the most concern (Stepanyants et al., 2009; Levy and Reyes, 2012). Experimentally, we used thick slices (350 µm), targeted cells deep in the slice (Fig S1B, median cell depth = 75 µm), and focused on local (<200 µm apart) intralaminar connections to minimize this effect. Yet, when we subselected our data for cell pairs in which the presynaptic axon was longer than 200 µm (see Methods) we saw that the estimated p_{max} increased by ~20-30% (Fig S1A, D), slightly less than one previous estimate of the effect of slicing artifacts in mouse auditory cortex (Levy and Reyes, 2012). We performed a similar analysis on the depth of the cells from the slice surface and found this had a lesser impact on p_{max} (Fig S1B, ~10-20%). The limited effect of depth on connectivity observed in this study may result from the relatively deep patching achieved, facilitated by software tools to guide the pipette to the selected cell.

Another factor affecting the measurement of the true connection probability is the ability to detect synaptic responses obscured by background electrical activity. Increasing the number of presynaptic action potentials increases the likelihood of detecting small synaptic responses that may be obscured in the noise. We quantified the overall 'detection power' of each cell pair as a function of the amplitude of background noise and the number of recorded presynaptic spikes (see Methods). Unlike slicing artifacts or intersomatic distance, estimates on the number of undetected synapses due to detection limits are seldom reported, although we performed a related sensitivity analysis in our previous study (Seeman et al., 2018). We modeled the relationship between detection power and connection probability in a similar way to intersomatic

distance which resulted in a 50-90% increase in estimated p_{max}, with a larger effect on inhibitory connectivity (Fig S1D), suggesting that the observed rate of connectivity is affected more by detection power than by slicing artifacts.

We observed a two to three and a half fold increase in p_{max} among the four combinations of excitatory and inhibitory cell class connections when simultaneously filtering for axon lengths, pair depths, and detection power values above their respective median (Fig 2A, dotted line). In order to more accurately adjust for the underestimation of connectivity rate, we combined the models for intersomatic distance, presynaptic axon length, cell depth, and detection power discussed above, into a unified connection probability adjustment (see methods, Fig S1D pink curve). We saw that the model-adjusted connectivity rate (Fig 2A dashed line) was similar to the filtered rate across the four connection classes. In some cases (eg. Pvalb→Pvalb, Sst→Vip), p_{max} estimates are near-complete, and are substantially higher than those reported in previous multipatch studies (Jiang et al., 2015; Karnani et al., 2016). This unified model allowed us to extend individualized connectivity adjustments to each pair across matrix elements (Fig S1E). In general we see fairly uniform adjustments for presynaptic axon length and pair depth across the matrix with a larger adjustment for detection power in inhibitory connections as is expected from our grouped results above. It will be necessary to confirm these adjusted functional measurements of connectivity with anatomical methods.

Intralaminar connectivity matrix

We constructed a connectivity matrix in mouse VISp with adjusted p_{max} values in each element scaled in saturation by the 95% confidence interval (more saturated colors have smaller confidence intervals, or CIs; Fig 2B, S2A). Often, we observe connectivity results that are consistent with prior results (Pfeffer et al., 2013; Jiang et al., 2015; Karnani et al., 2016). However, we also saw that intralaminar connection probability varies by layer (Fig 2B and C), long-range projection target of excitatory cells (Fig 2D), and cell subclass more generally (Fig 2 E, F). Our results show that L2/3 connectivity is substantially higher than other layers. This is highlighted by the pyramidal to Pvalb connection in L2/3 (p_{max}=0.76 [0.49, 1.00] CI) compared to L6 (p_{max}=0.41 [0.22, 0.67] CI) (Fig 2C, S2A). Additionally, recurrent connections between excitatory and Vip cells are common in L2/3 (E→Vip 0.47 [0.23, 0.80]; Vip→E 0.14 [0.03, 0.40]) but rare or absent in deeper layers (Fig S2A). Conversely, Pvalb→Vip connections were found in all layers except L2/3.

In addition to differences between layers, we saw some elements that are consistent across layers. For instance, Vip and Sst cells recurrently connect to each other in a similar ratio in all layers. Vip→Sst connections are often discussed in the context of disinhibition of pyramidal cells (Fu et al., 2014; Keller et al., 2020), and yet the role of Sst→Vip connections is comparatively underrepresented in the literature. Our results suggest that reciprocal connectivity between Vip and Sst cells is a common motif across layers that may play a more complex role in visual behavior (Dipoppa et al., 2018).

Sst is thought to avoid connecting with itself (Pfeffer et al., 2013; Naka et al., 2019; Scala et al., 2019) we observed connections between Sst neurons in almost every layer (Fig 2B). It is known that the Sst-IRES-Cre driver can sparsely label fast-spiking interneurons that also express Pvalb (Hu et al., 2013). UMAP clustering of intrinsic properties from inhibitory cre-types showed that slightly more than half of the recurrent Sst connections had both the pre- and postsynaptic cell in a cluster that was spatially distinct from Pvalb cells, suggesting that these connections came from cells that were intrinsically Sst-like (Fig S3A). We confirmed that, in at least one case, both cells in the pair had axons extending into L1 and sparsely spiny dendrites, consistent with Sst neurons (Fig S3B). We performed follow-up experiments utilizing the Patch-seq method (Lee et al., 2020) and further confirmed that cells which transcriptionally mapped to the Sst subclass do form connections with each other (3 connections found out of 47 probed, Fig S3C).

Our previous study focused on recurrent excitatory connectivity in each layer (Seeman et al., 2018). Within L5 we investigated connectivity among and between two different projection classes, intratelencephalic (IT, labeled by Tlx3) and extratelencephalic (ET, labeled by Sim1 and mscRE4). Consistent with our prior work and others (Brown and Hestrin, 2009) connectivity among ET cells is higher than that of IT cells (Fig 2D). Furthermore, connectivity between these projection classes is unidirectional from IT to ET (Brown and Hestrin, 2009) and they receive differential inhibition (Fig 2E,F). Layer 5 Pvalb cells innervate ET and IT at similar rates (Fig 2E) whereas L5 Sst cells preferentially innervate L5 ET cells (Fig 2F). Target specificity of Sst cells based on excitatory projection subclass has also been seen in frontal cortex of the rat, where Sst cells innervate corticopontine cells at a higher rate than corticostriatal (Morishima et al., 2017).

Electrical synapses

In addition to chemical synaptic transmission among cell subclasses, electrical connections, facilitated by gap junctions, were also found between inhibitory subclasses (Fig S4A). The likelihood of electrical synapse connectivity as a function of lateral intersomatic distance could also be approximated by a Gaussian but with a narrower profile, σ = 77 μm (Fig S4B), compared to chemical synapses between inhibitory cells (σ = 131 μm, Fig 2A). To assess the significance of this difference we asked what the likelihood was that the observed σ values came from the same distribution. Over a range of σ values from 65 μm to 140 μm, we found that fewer than 0.1% of simulated experiments had a σ ratio greater than what we observed experimentally (1.7, Fig S4B *lower* dotted black line). This is consistent with previous reports of electrical connections between nearby Pvalb cells which showed that the average distance between electrically coupled cells was short, 40-80 μm (Galarreta and Hestrin, 1999, 2002; Gibson et al., 1999). L2/3 Pvalb cells showed the highest rate of electrical connections (18/106, ~17%). Electrical connections among Vip cells were the most rare with only nine total found across all layers (out of almost 800 probed; six of which were interlaminar between L4 and L5), in contrast to a previous report of Vip electrical connections which were more prevalent (Karnani et al., 2016). A majority of electrical synapses were found between like subclasses (148/154, 96%) and were bidirectional (134/154, 87%). The distance of the gap junction from the soma

coupled with the potential for some rectification of electrical connections (Alcamí and Pereda, 2019) could account for the few cases where reciprocal electrical connections were not observed. There were three cases of cross-subclass electrical synapses, one between an Sst and Pvalb cell and two between an Sst and Vip cell, all in L6. Although rare, gap junctions between cells of different subclass have been reported before (Gibson et al., 1999). The coupling coefficient of electrically coupled pairs was comparable across subclass (Fig S4C) largely due to the lower input resistance of Pvalb cells (Fig S4D). Estimating junctional conductance revealed stronger electrical synapses between Pvalb (0.37 ± 0.38 nS, median ± SD) cells than either Sst (0.21 ± 0.2 nS) or Vip (0.06 ± 0.13 nS) (Fig S4C). Both coupling coefficient and junctional conductance were similar in magnitude between reciprocal pairs (Fig S4C).

Synaptic strength and kinetics

In addition to connectivity, synaptic properties determine the impact of a connection on the postsynaptic neuron and, ultimately, on cortical processing. The strength, latency, and kinetics of local synapses have been described across multiple studies for different cell types (Thomson et al., 2002; Holmgren et al., 2003; Pfeffer et al., 2013; Jouhanneau et al., 2015; Pala and Petersen, 2015; Walker et al., 2016). Still, this study offers the first open dataset in which synapse properties may be directly compared across many cell subclasses and layers. For the purposes of these analyses, transgenic cell subclass is used to identify the pre- and postsynaptic cell type resulting in semi-layer-specificity of excitatory-inhibitory (E-I) and I-E synapses, while pooling I-I synapses across all layers. Pooling was motivated by the observed homogeneity in intralaminar I-I properties across layers (Fig S5) and allowed for more robust comparisons. Nevertheless, pooled results may contain layer-related biases due to differences in the laminar distribution of cell bodies across the inhibitory subclasses. For example, most Vip cells are located in L2/3 (Fig S5A) and thus synapses involving these cells may be biased toward upper layers (Gonchar et al., 2008; Almási et al., 2019). Synaptic properties were extracted from exponential fits to the average postsynaptic response (PSC/P) (see Methods for more detail). Figure 1B shows an example fit to an average EPSP and IPSP from one recording.

We observe patterns in PSP latency at the class and subclass levels. Inhibitory synapses (with generally narrower action potential widths) had shorter latencies (median=1.08 ms) than excitatory synapses (median =1.49 ms, ks=3.17e-32) regardless of the postsynaptic target (I→I 1.1 ms, I→E 1.02 ms, ks=0.02). Pvalb synapses are extremely fast, with sub-millisecond latencies (Fig 3A) highlighted by Pvalb→L6 excitatory synapses (0.97 [0.86, 1.09] ms, median [interquartile range]). Recurrent excitatory synapses had the longest latencies (E→E 1.67 ms, E→I 1.33 ms ks=1.8e-9). It may be valuable to predict PSP latency from presynaptic spiking behavior in network models on a cell type basis and cases where spike properties change during activity. We find that PSP latency is weakly correlated with action potential width for excitatory (r^2 =0.05, p=1.71e^{-05}) synapses and more strongly correlated for inhibitory synapses (r^2 =0.19, p=1.15e^{-49}). The stronger correlation in inhibitory synapses likely reflects that the

calcium entry that drives release occurs during action potential repolarization when the driving force is high and the calcium channels are in the act of closing (Chao and Yang, 2019), whereas at excitatory synapses, there may be appreciable calcium entry during the action potential. As our data was acquired at near physiological temperature, the observed latencies in this study may be considered to reasonably reflect *in-vivo* timing.

Response kinetics, PSP rise time (Fig 3B) and decay time constant (Fig 3C), showed a number of interesting trends at the subclass and class level. Inhibitory→excitatory synapses showed some of the longest rise times (I→E 5.43 ms, I→I 3.6 ms, ks=1.44e-15), while the reverse excitatory→inhibitory synapses displayed the fastest rise times (E→I 2.73 ms, E→E 3.87 ms ks=5.18e-12), likely driven by postsynaptic Pvalb as is highlighted by L4 excitatory synapses onto Pvalb cells (1.45 [1.28, 1.94] ms). Excitatory synapses to Sst cells (see L2/3 Pyr→Sst) had a slower rise time (4.06 [2.99, 4.91] ms) and decay (25.18 [15.37, 38.88] ms), suggesting a contribution of postsynaptic subclass to driving synaptic kinetics. Presynaptic Sst cells, regardless of postsynaptic target, have some of the slowest kinetics (see Sst→Vip rise time: 7.15 [5.7, 10.0] ms, decay: 46.7 [24.41, 235.09] ms; Sst→L5 IT rise time: 6.5 [5.73, 8.55] ms, decay: 32.0 [21.64, 64.36] ms). These examples showcase an organization in synaptic kinetic properties that is driven by the pre- and postsynaptic cell class, and further refined by differences in subclass.

Two measurements were used to represent synaptic strength: the resting-state amplitude (the average response to a single spike after a period of inactivity, Fig 3D) and the 90th percentile amplitude from all responses (Fig 3E), or near-maximal amplitude. As a general rule, excitatory connection strength is related to the postsynaptic cell type. Recurrent excitatory connections showed some of the smallest amplitudes (eg. L5 ET→L5 ET), both in the resting state (0.27 [0.13, 0.5] mV) and in the 90th percentile (0.31 [0.22, 0.57] mV) which is consistent with our previous work showing that these connections are weak and mostly get weaker with activity (Seeman et al., 2018). Excitatory connections onto inhibitory cells are stronger on average. The strongest connections project to Vip cells, although these connections were found predominantly in superficial layers (L2/3 pyr→Vip 0.56 [0.23, 0.84] mV). Excitatory connections onto Sst cells were comparatively weak when measuring the resting state amplitude; however, these synapses are also strongly facilitating, and thus are weakest in the resting state. In contrast to excitatory synapses, inhibitory synaptic strength is most related to the presynaptic cell, with Pvalb cells eliciting the largest PSP amplitudes, in almost every case regardless of postsynaptic cell subclass. One exception is the strength of Pvalb→L5 ET (resting state: -0.33 [-0.31, -0.80] mV, 90th percentile: -0.37 [-0.28, -0.76] mV) which is comparatively weaker than Pvalb→L5 IT (resting state: -0.53 [-0.21, -0.77] mV, 90th percentile: -0.56 [-0.32, -0.72] mV), again highlighting the dichotomy of these two projection classes.

For synapses that facilitate, the resting state PSP amplitude is an underestimate of the potential impact on the postsynaptic cell. To compare synapses in the absence of this bias, we used the 90th percentile of PSP amplitudes as a measure of the near-maximal strength for each synapse. Overall this result follows similar patterns to the resting state strength, with the exception that matrix elements containing the most facilitating synapses are comparatively

stronger, such as E→Sst and Sst→Vip (Fig 3E, left). Facilitation onto inhibitory cells contributes to the longer tails of 90th percentile amplitudes and the rightward shift of E→I (0.74 mV) amplitudes compared to E→E (0.33 mV, ks=1.11e^{-16}) (Fig 3E, histograms). Furthermore, the largest average resting state (2.69 mV), 90th percentile (4.86 mV, L2/3 pyramidal→L2/3 Sst), and single PSP (15.03 mV, L5 IT→L5 Sst) amplitude were all found in an excitatory to Sst connection.

A recent survey of cortical connectivity found that synapse strength positively correlated with connection probability (Jiang et al., 2015). This result suggests an interesting principle of connectivity, but may also result from the reduced detectability of weaker synapses. When we assessed this relationship we found that adjusted connection probability of excitatory synapses was not correlated with synaptic strength (weighted Huber regression r^2 = 0.2, p = 0.5) while inhibitory synapses showed a small correlation (weighted Huber regression r^2 = 0.4, p < 0.01). Recall that we saw a smaller effect of detection power on excitatory compared to inhibitory connections (Fig S1C) suggesting that when detection power is sufficient connection probability is independent of synaptic strength.

Synaptic dynamics

The amplitude of PSPs described above capture the strength of a synapse in response to one presynaptic spike, but during repetitive activity PSP amplitude evolves over a timescale of milliseconds to seconds, a phenomenon known as short-term plasticity (STP; Fig 4, Fig S6). PSP amplitude may increase (facilitate), decrease (depress), or remain stable (pseudo-linear) over the course of a stimulus train as has been described previously (Thomson, 1997; Reyes et al., 1998; Beierlein et al., 2003; Ma et al., 2012; Markram et al., 2015). Our dataset similarly shows examples of depressing, facilitating, and pseudo-linear responses to a 50 Hz stimulus train (Fig 4A). The time course of recovery from STP is an equally important property of synapses, yet one that is not well described. Our 8 + 4 pulse stimulus (Fig 1B, Fig 4A, see Methods) allows us to more fully quantify STP recovery over a broad range of connection types.

STP Induction

Quantifying the magnitude of STP has taken several forms from a paired-pulse ratio (PPR, ratio of the second response to the first), to a ratio of the last pulse in a train to the first. These ratios are sensitive to noise, especially when the signal in the denominator becomes very small; thus we quantified STP using the difference between late (pulses 5-8) and initial response amplitudes, normalized by the 90th percentile response amplitude (Fig 1B, see Methods), termed STP induction. An examination of STP across many matrix elements highlights interesting differences in the average response (Fig 4B). Overall, STP follows a similar pattern to synaptic strength, wherein excitatory synapses are most strongly differentiated by the postsynaptic subclass and inhibitory synapses are differentiated by the presynaptic subclass .

Consistent with previous reports (Beierlein et al., 2003) we observed facilitating synapses from excitatory cells onto Sst in most layers. However, the STP of individual connections spanned a

range of magnitudes from highly facilitating, through pseudo-linear, to moderately depressing (Fig 4E) suggesting a high degree of heterogeneity in STP at L2/3 pyr→Sst synapses. We also observe a difference in the magnitude of facilitation of synapses onto Sst cells from ET and IT cells in L5 where ET to Sst shows stronger facilitation (Fig 4B, ET→Sst 0.16 [0.12, 0.19], IT→Sst 0.09 [0.03, 0.13]). Excitatory connections onto Pvalb (Fig 4E, L4 pyr→Pvalb) were largely depressing on average, except in L6 (L6 pyr→PV; 0.06 [0.0, 0.07]). Notably, a subset of L2/3 excitatory to Pvalb were pseudo-linear, similar to *in vivo* measurements in somatosensory cortex (Pala and Petersen, 2015). Pvalb connections onto other subclasses, both excitatory (Fig 4E, Pvalb→L6 pyr) and inhibitory, were also depressing on average. Connections from Sst cells skewed toward depression (Fig 4E, Sst→L5 IT) but were not as depressed as Pvalb connections (Ma et al., 2012). One notable exception was strong facilitation from Sst to Vip cells (Karnani et al., 2016) (Fig 4B). Our recent study (Seeman et al., 2018) showed that recurrent excitatory connections largely depress which we again see here (Fig 4E, L5ET→L5ET), but that depression is less among L2/3 cells (Fig 4B). All of these results highlight STP at one stimulus frequency, 50 Hz, however it has been shown that depth of facilitation or depression can vary depending on induction frequency (Beierlein et al., 2003; Feldmeyer et al., 2006). We similarly see a range of dependence of STP on stimulus frequency (Fig 4F) and presynaptic or postsynaptic subclass.

STP recovery

We quantified STP recovery using a similar measurement to induction, a normalized ratio of pulse averages (Fig 1B, *Synapse Analysis*) so as to avoid spurious ratios that may result from the inherent variability in individual PSP amplitudes. After a brief interval of 250 ms between the last pulse in the induction train and the first pulse of the recovery train (Fig 1B, *Multipatch Experiment*) we saw that synapses of most connection types had not recovered (Fig 4C). Highly facilitated excitatory to Sst connections were still in a facilitated state and highly depressed Pvalb connections were still depressed. In some cases, particularly recurrent L2/3 connections and L2/3 pyr→Vip connections, we see that the synapse has over-recovered and has gone from a depressed state, on average, to a facilitated state (Fig 4D). STP recovery of connections between Pvalb and Sst cells show that Sst→Pvalb connections recover faster than Pvalb→Sst (Fig 4C), consistent with prior results (Ma et al., 2012).

The variable delays we imposed between the induction and recovery pulses (Fig 1B, *Multipatch Experiment*) of our 50 Hz stimulus allowed us to quantify the time course of synaptic recovery (Fig 4E). From these data we see that at our earliest time point of 125 ms synapses are still in their STP induced state, but that by four seconds they are largely recovered, regardless of cell subclass or whether the synapse started in a facilitated or depressed state. The measurement of STP induction and recovery are based on the average response and don't capture variance across trials. We developed a measure of synaptic variability to quantify the trial to trial dynamics exhibited by different synapses types.

Synaptic variability

Multiple factors contribute to noise in measured PSP amplitudes, including the release probability, quantal variance, and biological and electrical noise sources (Costa et al., 2013). To access the component of the variability driven by synaptic release mechanisms, we used an adjusted coefficient of variation (aCV) that subtracts the experimental noise contribution (see Methods) from the measured PSP variance. Like STP, variability in excitatory synapses changes systematically across postsynaptic subclasses, whereas inhibitory variability is more strongly influenced by the presynaptic subclass (Fig. 4D). Excitatory synapses onto Sst cells exhibited the highest resting state variability, while PV neurons produced some of the most reliable responses. Recurrent excitatory synapses expressed a range of variability, that, as with other dynamics measurements, varied with their layer and projection type. To assess whether variability is affected by STP, we also measured aCV at the end of 50 Hz pulse trains. Synapses with short-term depression generally increased their variability after induction of STP (Fig. 4F; e.g., L4pyr→Pvalb), whereas facilitating synapses tended to decrease their variability (e.g., Sst→Vip).

Human synapses

For the initial characterization of human synaptic connectivity, inhibitory and excitatory cells were distinguished by morphology and the excitatory cells split into putative subclasses by layer. The inhibitory cells, expected to be primarily fast-spiking Pvalb cells due to their prevalence, were pooled across layers based on the consistency of inhibitory synaptic properties through the cortical depth observed in the mouse data (Fig. S6). Due to experimental constraints from tissue health and imaging challenges, sampling was primarily from the supragranular layers of cortex. Without enough data to fully constrain the complete connection probability adjustment model, we applied an adjustment for lateral intersomatic distance only. However, calculations suggest that the remaining adjustments are much smaller than in the mouse dataset (<50%) due to lower noise and more spikes recorded on average. The connection probability was estimated to fall off with distance at a σ of 140 μm (Fig. S7A), moderately larger than the comparable value in mouse (100 μm), reflecting that while cortical expansion is accompanied by the scaling of neuronal morphology, much of this scaling is axial rather than lateral.

Human synaptic properties

Dense sampling of L3, a dramatically expanded layer in anthropoid primate cortex (Balaram and Kaas, 2014), allowed us to split L3 pyramidal cells into L3a and L3b groups for analysis, reflecting known cell type distinctions through this depth (Berg et al., 2020). Examining the connectivity between these subclasses (Fig. 5A), we tested for signatures of functional segregation within supragranular layers, finding a strong bias for recurrent over cross-connections between L3a and L3b and a bias for connections from L2 to L3a over L3b. This descending connection is also more prevalent than the reverse ascending connection, from L3a to L2. Interestingly, the recurrent connectivity within L4 is extremely sparse in the human

data (p_{max}=0.01 [0.0, 0.04] 95% CI, 1/143), significantly lower than that observed in the mouse. Although cross-layer connections from L4 pyramidal cells were not observed in sufficient numbers to estimate connection probabilities, the two observed (38 tested) suggest that the lack of recurrence is not an artifact of a failure to find L4 connections in general, but rather exposes a unique property of the L4 circuit. The excitatory recurrent connection probability has been previously suggested to depend on age (Scala et al., 2019), but could also vary across species or brain area.

The strength, kinetics, and STP properties of the human synapses largely recapitulate observations from the mouse data, with moderate differences across layers and large differences by cell class (Fig. 5B). Recurrent excitatory connections (E-E) had longer latency (1.73 [1.43 2.26] ms, median [quartiles]) than those with a pre- or postsynaptic inhibitory cell (I→E 1.03 [0.86 1.60] ms, ks=2.2e^{-5}; E→I 1.34 [1.14 1.62] ms, ks=8.3e^{-4}), and PSP rise times are faster for E→I than E→E connections (2.48 [1.60 2.95] ms vs. 4.11 [3.15 5.57] ms, ks=8.0e^{-8}), but slower for I→E connections (6.33 [4.03 7.59] ms; ks=0.014, 3.2e^{-6} vs. E→E, E→I). We observed some differences between L2 and L3, including presynaptic L3 cells forming more depressing connections than L2 cells (STP ratios -0.41 [-0.63 -0.14] vs.-0.15 [-0.42 -0.01], ks=7.0e^{-3}). We also note that E→I synapses are uniformly depressing, consistent with the identification of those inhibitory cells as fast-spiking Pvalb cells. In certain properties, we did observe contrasts between human and mouse synaptic properties. The overall amplitudes of L2/3 E→E and E→I connections were significantly larger than the corresponding synapses in mouse, with recurrent excitatory resting state response 0.37 [0.14 0.64] mV (ks=0.039; E→I 0.93 [0.37 1.76] mV, ks=7.0e^{-4}), 90th percentile response 0.51 [0.30 0.96] mV (ks=0.040; E→I 1.17 [0.81 2.24] mV E→I, ks not sig.), and largest single response 12.5 mV (L3 E→I). We also observe dramatically faster recovery from STP in human than mouse excitatory synapses, with most fully recovered at 500 ms (Fig. 5C).This contrast has been previously noted in recurrent L2/3 excitatory synapses (Testa-Silva et al., 2014); our observations suggest that it holds for L5 E→E and L2/3 E→I synapses also.

Human polysynaptic events

Large-amplitude synaptic connections in human, but not mouse cortex, have been reported to trigger polysynaptic, complex events (Molnár et al., 2008). Indeed, we also observed polysynaptic events, primarily short-latency (~3ms) inhibition (Fig 5D) evoked by a confirmed pyramidal cell. Short latency disynaptic inhibition could be driven by an intervening fast-spiking parvalbumin cell, as has been reported in human (Szegedi et al., 2017) and rat (Silberberg and Markram, 2007). This is in contrast to disynaptic inhibition mediated by Sst cells (Kapfer et al., 2007; Silberberg and Markram, 2007; Obermayer et al., 2018) which has a longer latency (~150ms) due to facilitation of excitatory inputs onto Sst cells. Plotting latency versus PSP amplitude of human synapses (Fig 5D) reveals a clear boundary where responses evoked from excitatory cells with a latency of 3 ms or greater are almost exclusively inhibitory (median latency of dIPSPs: 4.04 [3.77, 4.49] ms), compared to monosynaptic EPSPs which had a latency less than 3 ms (median EPSP latency: 1.7 [1.44, 2.13] ms). This potential disynaptic

inhibition (dIPSPs) appeared directional from L2 pyramidal cells to other L2 (n=10) or L3 (n=11) pyramidal cells but just one ascending polysynaptic response originating in L3. This directionality is consistent with the directionality of monosynaptic excitation across layers 2 and 3 as well as supported by higher connection rates from L2 pyramidal cells to inhibitory cells. Median latency of recurrent L2 dIPSPs (4.08 [3.92, 4.56] ms) was similar to that of L2→L3 dIPSPS (3.84 [3.58, 4.24] ms), but recurrent L2 dIPSPs (-0.86 [-0.09, -1.03] mV) were almost three times larger than L2→L3 dIPSPs (-0.28 [-0.11, -0.44] mV). As mentioned above, excitatory inputs from L2 and L3 (Fig 5B) were considerably larger than those in mouse L2/3 (Fig 3E). Very large events in human have been shown to drive spiking in fast-spiking interneurons (Szegedi et al., 2017) and, in fact, in two connections, we observed a pyramidal cell induce an action-potential in a postsynaptic aspiny neuron.

Variation of STP with depth in layers 2 and 3

Another property of supragranular neurons observed in human, but not in mouse, is strong depth-driven variability of intrinsic electrophysiological properties (Kalmbach et al., 2018; Berg et al., 2020). We further examined STP for possible links to this intrinsic variability, expanding on the observed differences in STP between L2 and L3 pyramidal cells. We first refined our criteria for the L2/3 pyramidal subclass using an upper bound on the input resistance of 225 MΩ, excluding L4-type cells that can overlap into L3 based on their smaller size and higher input resistance (Berg et al., 2020). Visualizing the electrophysiology feature space by a UMAP projection of 27 electrophysiology features (Methods), we found that these L4-type cells are situated around the perimeter, indicating distinct properties from L2/3-type cells (Fig 5E). Similarly, the separation of most inhibitory cells into a distinct cluster is consistent with their identification as fast-spiking Pvalb cells. For the L2/3-type cells, projecting a normalized layer depth coordinate (relative to the L2+L3 thickness) onto this space shows a mostly smooth gradient of electrophysiological properties with layer depth, also verified by direct examination of depth correlations with sag and AP upstroke/downstroke ratio (p=2.9×10^{-7}, 6.7×10^{-6}) (Fig 5F). Note that, as previously observed, this structure is not found in mouse L2/3 cells (p>0.09 for both).

Analysis of synaptic STP metrics revealed similarly linear variation of STP with layer depth in synapses from L2/3 pyramidal cells. We identified the depth of a synapse with the presynaptic soma (for postsynaptic, see Fig S7F); since these trends appear consistent across postsynaptic cell classes (Fig 5B STP, Fig S7E), we included both E→E (n=85 human, n=19 mouse) and E→I (n=19 human, n=71 mouse) connections. The paired-pulse STP showed a strong linear relationship with depth in the human data (p=2.6×10^{-5}), varying from weak facilitation for the most superficial cells (0.03 ± 0.03, mean±SEM) to depression for the deepest (-0.37 ± 0.10). A strong correlation was also found with the action potential upstroke/downstroke ratio of the presynaptic cell only (p=2.2×10^{-5} vs p>0.1 for postsynaptic), suggesting that links between spike shape and neurotransmitter release could help explain the variation of STP with depth (Fig 5G). No corresponding trends were found in the mouse data (p>0.1 for all regressions). Although lower sampling of L2/3 pyramidal cells may contribute, regression coefficients for STP against

upstroke/downstroke ratio show a strong contrast (human -0.18 [-0.26, -0.10] CI; mouse 0.06 [-0.04, 0.16]), suggesting that there are real differences between these datasets in the factors contributing to STP variability, whether explained by species, brain area, or other factors.

Modeling short term plasticity

The STP metrics introduced above were chosen for their ease of interpretation, but provide a limited description of the complex behaviors expressed by synapses. For example, we describe the depth of depression and facilitation under a few different conditions, but neglect the rates of depression and recovery and the dependence on stimulus frequency. Ideally, we would like a description that can predict synapse behavior in response to any arbitrary stimulus. Rather than continue to engineer more descriptive statistics, we developed a generative model of stochastic vesicle release and STP with several adjustable parameters (see Methods) and asked which combinations of parameters were best able to explain the responses recorded for each synapse. In this way, we capture and describe more of the dynamic behavior of each synapse with a small number of parameters

Most discussions of short term depression in the cortical literature begin with the assumption that depression is caused by the depletion of vesicles from the readily-releasable pool. However, recent evidence suggests that calcium channel inactivation may be a more prominent mechanism in cortical depression (Nanou and Catterall, 2018). We ran the model on two separate parameter spaces--one that uses vesicle depletion, and another that uses a release-independent depression mechanism. In most cases, the model maximum likelihood value was found in the release-independent parameter space (Fig S11E, left). Release-dependent depression mechanisms should result in negative correlation between consecutive PSP amplitudes; however, we find little evidence for such negative correlations in our data. Likewise, we find little relationship between paired correlation values and the model preference for release-dependent mechanisms (Fig S11E, right). These results are consistent with the proposal that cortical synapses employ release-independent depression mechanisms, and that vesicle depletion plays a relatively minor role in depression.

Model performance was evaluated by using the maximum likelihood parameter set for each synapse to simulate experimental data. This simulated data was then used to generate the same STP metrics that were previously collected from synaptic data. Both resting state PSP amplitude and 90th percentile amplitude are almost perfectly correlated between recorded and simulated data (Fig S12A,B), indicating that the model does exceptionally well at capturing synaptic strength. STP and variability (Fig S12C-F) are also strongly correlated, but with more scatter relative to strength metrics. Notably, STP measured from the second pulse in 50 Hz trains was only half as large as measured in synaptic responses, indicating that the model as parameterized was not able to fully capture STP on this timescale. Resting state variability is fit by the model through the binomial coefficient of variation, which is the product of the resting state release probability and the number of synaptic release sites. Release probability was more

highly correlated with variance than number of release sites, suggesting that synapses may control variability primarily through their release probability.

Organization in synaptic dynamics

With a large dataset describing synapse properties at hand, it becomes possible to ask what patterns emerge from the data. What synaptic features correlate with one another, and do synapses naturally split into clusters based on these features, or do they form a continuum? What aspects of the synaptic feature-space are driven by presynaptic versus postsynaptic cell type? To answer these questions, we used sparse PCA followed by UMAP dimensionality reduction to summarize the output of the stochastic release model described above for 1140 synapses (980 mouse, 160 human) (Fig. 6). This analysis groups synapses based on the similarity of their model results. Therefore, it has access to any synaptic strength and dynamical properties that the model could capture, but does not have access to any information about kinetics, cell subclass, or other cell properties.

Synapses in this analysis form clear excitatory and inhibitory clusters (Fig. 6A), with a continuum of synaptic properties within each cluster. Perhaps the most prominent feature of this organization is that excitatory synapse properties are strongly influenced by the postsynaptic cell subclass (Fig 6B), whereas inhibitory synapse properties are more strongly driven by their presynaptic subclass (Fig 6C), consistent with results described above for strength and STP. The excitatory cluster in Figure 6B (left) is stratified by postsynaptic cell subclass, with the most distinct separations seen between E→Sst, E→Pvalb, mouse E→E, and human E→E. In contrast, inhibitory synapses (right) overall have a more uniform distribution of postsynaptic subclasses, and instead are differentiated by their presynaptic subclass. Many Pvalb synapses in particular occupy an exclusive region of the inhibitory map, while Sst synapses concentrate more strongly at the opposite end and Vip share the central region where all three subclasses overlap. Further subdivisions are present but less prominent. For example, mouse L2/3 E→I synapses concentrate near one edge of the presynaptic subclass map, while human excitatory synapses are separated by their postsynaptic layer.

What synaptic properties determine this organization by subclass? The most clear property that correlates with the UMAP dimensions is synaptic strength, which forms a well defined gradient across the long axis of each cluster (Fig. 6C, left). However, this gradient is mostly orthogonal to the boundaries separating cell classes, indicating that each class occupies a large range of different synaptic strengths, but that the strength differences between postsynaptic subclasses are comparatively small. Orthogonal to the strength axis, we find that response *variability* differentiates synapses and strongly separates subclasses. This axis also correlates with STP, where the most facilitating E→Sst synapses appear at the high-variability end of the axis. At the opposite extreme, human E→E synapses occupy the low-variability edge. Likewise, presynaptic Pvalb cells project the most stable synapses, whereas Sst and Vip project more variable synapses. Thus synaptic strength and variability form two major axes of the reduced synapse-type space. To verify that this is not an artifact of our dimensionality reduction to two

dimensions, we repeated the analysis for three dimensions but found the same 2D structure flattened in 3D space.

Discussion

The mammalian cortex is believed to act as the computational substrate for our highest cognitive abilities, particularly the ability to model the world around us and predict the effects of our actions. It is also of particular interest because many aspects of its structure are repeated across brain regions and conserved across species, suggesting the existence of a general-purpose approach to computation in the cortex. The results we have presented here expand upon a long history of electrophysiological and anatomical experiments exploring the cortical microcircuit's local connectivity. These experiments are technically challenging, making it so far impossible to map complete circuits. Instead, the canonical circuit diagrams established by this research field have evolved piecewise over the years to accommodate new evidence. These efforts offer an excellent depth of insight to isolated regions of the circuit but lack cohesion at a large scale, and furthermore the difficulty of accessing these historical data discourages reuse and reanalysis. We saw an opportunity to run a broader survey than has been attempted in the past, and extend the value of this resource to the community by making our analyses, tools, and data open to the public .

By probing over 20,000 possible connections across 28 mouse lines, we have explored a large fraction of the subtype-specific, intralaminar connectivity in the mouse visual cortex. At the same time, we share the first systematic survey of intralaminar connectivity in the human cortex. Past surveys near this scale have focused on connectivity and strength of synapses; a major advance provided by our study is the depth of characterization and analysis for each synapse, in the context of transgenically identified cell subclasses and species. We find that excitatory dynamics are driven largely by the postsynaptic cell subclass, whereas inhibitory dynamics depend on the presynaptic cell subclass. At the same time, synaptic properties form a continuum within each cluster such that, at the subclass level, short-term plasticity is highly heterogeneous.

Limitations

The methods we use require brain tissue to be dissected and sliced to provide easier visual and physical access to the neurons to be patched. In this process, some neurons are damaged, some connections are severed, and much of the *in vivo* environment is replaced with controlled conditions. At the same time, the synaptic effects we are interested in can be measured only indirectly via their impact on the cell soma. Our ability to resolve these small signals is affected by dendritic filtering, background noise, and the inherent variability of the synapse. Despite these limitations, paired patch clamp recording remains the gold standard in synaptic physiology for its excellent electrical resolution, temporal precision, and control over presynaptic spiking. Past studies have modeled the effect of severed connections (Levy and Reyes, 2012) and

sensitivity (Seeman et al., 2018) on the rate of false negatives in connectivity measurements. We believe this is the first study to quantify these sources of error simultaneously and estimate adjustments, which are substantial, especially for mouse inhibitory connectivity. In addition to providing better estimates of connectivity, reducing experimental bias makes it possible to compare our results between studies with more confidence. For example, recurrent excitatory connections were found more frequently in human, but our analysis of detection power suggests that this difference is due to enhanced signal to noise ratio (arising from a lower noise floor and stronger and more reliable PSPs) in human cortical tissue.

A further confound associated with human neurosurgical tissue is the potential for disease pathology (epilepsy or tumor in this case) to affect the physiological measurements we are making such that we may not be capturing the healthy human brain. While this is important to keep in mind, these tissues are currently our only access to investigating the living human cortex. Additionally, a recent study (Berg et al., 2020) on intrinsic cell physiological features in human found strikingly consistent results across nearly 100 different human tissue donors which varied in age, gender, cortical region, and pathology. This suggests that despite disease state, we can gain useful insight into human cortical function through the use of neurosurgical specimens.

For this study, we focused our efforts on intralaminar connectivity; instances of interlayer connections are present in the dataset but relatively sparsely sampled. For nearby layers, this could be addressed with further experiments. However connections across distant layers are more difficult to interrogate with this method because long axo-dendritic pathways are more likely to be severed (Stepanyants et al., 2009), especially in human cortical tissue that can be several millimeters from pia to white matter. A detailed study of such medium-range connections may require a different methodological approach (Hage et al., 2019).

The transgenic mouse lines used in this study enabled the investigation of connectivity and synaptic physiology between subclasses that are otherwise difficult to target in slice recordings. However, they also merge more refined cell types (Gouwens et al., 2019, 2020) and miss others. The most notable omission is transgenic mice that identify serotonin positive, Vip negative interneurons which make up about 20% of the interneuron population (Tremblay et al., 2016). Merging cell types is a common issue when a single modality is used to assign a cell identity (DeFelipe et al., 2013). To overcome this limitation, we provide a multimodal dataset (physiological and morphological), and these complementary modalities may be used to refine or verify the cell identity. End-users may utilize the various modalities in our dataset in ways that are appropriate for their research question.

Conserved and canonical elements in the intralaminar circuit

As many prior studies have investigated the cortical circuit, a picture has emerged describing the relationships between the excitatory and inhibitory subclasses and their functional relevance. The details of this picture vary somewhat between descriptions, but a few key

elements appear consistently, especially in the systems and theoretical neuroscience literature (Fig 7A). Pvalb interneurons strongly inhibit nearby pyramidal cells and other Pvalb cells, Sst interneurons broadly inhibit nearby cells but avoid other Sst cells, and Vip cells selectively inhibit Sst cells and receive feedback excitation, forming a disinhibitory circuit (Fu et al., 2014; Keller et al., 2020). We have confirmed that each of these motifs is prominent across layers in the intralaminar cortical circuit.

We also find circuit elements that are equally prominent in our data, but are more sparsely acknowledged in the literature. Many recent studies have focused on the importance of the Vip→Sst disinhibitory circuit. In the opposite direction, however, the connection from Sst to Vip has one of the highest connection probabilities, largest IPSP amplitudes, and strongest facilitation in our dataset. Although Sst→Vip connections have been described previously (Karnani et al., 2016), they are often overlooked in consideration of the disinhibitory circuit. The unique synaptic features we observed suggest an important functional ramification on the opposing Vip→Sst disinhibitory pathway. For example, local excitation could drive disinhibition either through the established Vip→Sst pathway or through the reverse Sst→Vip pathway. Furthermore, mutual inhibition between Vip and Sst populations could result in bistability in which either subclass may exclusively drive inhibition of pyramidal cells, but not both simultaneously.

Sst and Vip cells are often described as lacking recurrent connections (Hu et al., 2011; Ma et al., 2012; Naka et al., 2019; Scala et al., 2019) despite some evidence to the contrary (Gibson et al., 1999). We have confirmed that Sst and Vip do have the expected biases in their connectivity across all layers (Sst cells tend to avoid contacting other Sst cells and Vip cells prefer to contact Sst cells). However, we also find sparse, recurrent connections within both interneuron populations about equal to the recurrent connectivity in excitatory populations. Furthermore, we find that the strength of connections from Sst and Vip do not follow the same preferences, having roughly equal strength when connecting to preferred versus non-preferred subclasses. Given their trans-laminar axon projection patterns, recurrent connections within these subclasses may be found more commonly across layer boundaries (Gouwens et al., 2020).

Laminar variations on the cortical circuit

Previous studies that sampled both L2/3 and L5 noted many similarities between the two layers (Pfeffer et al., 2013; Jiang et al., 2015). Although we have confirmed a consistent set of connectivity rules describing the intralaminar circuit, we also find variations on these rules that could support different modes of cortical function. Differences in intralaminar circuitry may contribute to laminar differences in receptive field properties (Niell and Stryker, 2008) or visually mediated behaviors (Fu et al., 2014).

Layer 2/3 has strong interconnections between pyramidal and Vip cells (Fig 7B). In deep layers, these connections are either absent or greatly reduced, and the relative sparsity of Vip cells in deep layers should further enhance this difference (Gonchar et al., 2008; Almási et al., 2019). Direct Vip inhibition of local pyramidal cells further complicates the prevailing view of the

Vip→Sst disinhibitory pathway by, for example, allowing the possibility of feedback inhibition from higher cortical regions. Likewise, local excitatory inputs to Vip cells could be a source of feedforward disinhibition in layer 2/3. In contrast to L2/3, recurrent connections between the Pvalb and Vip subclasses were found most frequently in deep layers (Fig 7C). Ultimately, more targeted experiments and modeling will be needed to explore the functional relevance of these laminar variations in the cortical circuit.

Layer 5 excitatory subclasses differ in their visual responses and long range projections, suggesting different functional roles in the circuit (Kim et al., 2015; Lur et al., 2016). In this context we see several interesting differences in L5 ET and IT neurons. ET pyramidal cells are generally more highly connected, receiving more local excitation and inhibition than IT cells. We confirmed a much higher rate of recurrent connections among L5 ET cells compared to IT as well as the observation that connections between these two subclasses are unidirectional from IT→ET (Fig 7C) (Brown and Hestrin, 2009). Layer 5 ET cells also receive more frequent inhibition from Sst cells, as previously observed in frontal cortex of rat (Morishima et al., 2017).

Two recent studies investigated the correspondence between morphological, electrophysiological, and transcriptomic (MET) features of inhibitory neurons in primary visual cortex and motor cortex (Gouwens et al., 2020; Scala et al., 2020). In visual cortex, different MET types had distinct patterns of local axonal innervation and dendritic morphologies (Gouwens et al., 2020) suggesting that their connectivity will be different. MET types also exhibited layer localization and thus some of the differences in connectivity we observed as a function of layer may reflect differences in connectivity between different MET types. We have a small dataset investigating connectivity of transcriptomically defined Sst cells and our future work exploring the connectivity of MET types more generally may uncover finer circuit details.

Dynamic flexibility in the cortical circuit

Most studies in cortical synaptic physiology describe the circuit in its quiescent state. Ongoing activity *in vivo*, however, dynamically rewires the network by strengthening or weakening synapses. Cortical up- and down-states in particular have the capacity to synchronously facilitate or depress large portions of the network. We find a diversity of dynamic properties even among specific cell subclasses; thus to some extent this dynamic network reshaping occurs at the level of individual synapses. However, we also find that many subclass elements of the cortical circuit show a clear preference for either facilitation or depression (Fig 7D). Overall, most synapses in our dataset were found to express synaptic depression. Pvalb cells in particular receive and project almost exclusively depressing synapses. In contrast, Vip and Sst connections express a mixture of depression and facilitation. Excitatory inputs to these subclasses are often strongly facilitating, as are the interconnections between Vip and Sst. These patterns suggest the ability to dynamically switch between two network modalities where intralaminar activity is either dominated by interactions between pyramidal, Sst, and Vip cells

during sustained activity, or dominated by interactions between pyramidal and Pvalb cells following periods of quiescence.

Excitatory cells receive weak excitation compared to inhibitory cells

Our previous study (Seeman et al., 2018) observed low recurrent excitatory connectivity rates in all layers. We now find that recurrent excitatory synapses have a set of features that distinguish them from excitatory inputs to inhibitory cells and appear to limit their contribution to excitability. In addition to being sparse, they are relatively weak, they get weaker with activity, and they have slower PSP rise times. Slow PSP rise times would limit excitability by raising action potential threshold (Azouz and Gray, 2000). Recurrent excitatory connections also have relatively long latencies. A well-described feature of cortical networks is that inhibition lags behind excitation. This correlation has wide-ranging functional consequences, including creating windows of integration (Pouille and Scanziani, 2001; Pouille et al., 2009). It is interesting to consider the differences between mouse visual cortex and human temporal cortex recurrent excitatory rates in this context. L4 in mouse has the highest recurrent connectivity rate of all layers. L4 also receives potent feedforward inhibition in the visual cortex (Cruikshank et al., 2007) that may coincide with the long latency recurrent excitation to shape the integration window. In human temporal cortex, recurrent excitation is practically absent in L4,and thus is unlikely to shape the integration window.

Unidirectional disynaptic inhibition in human

We observed disynaptic inhibition in human cortex between confirmed spiny pyramidal cells that is unidirectional, originating in L2 and targeting other L2 or L3 pyramidal cells. We did not see disynaptic inhibition in our mouse recordings, which may be due to the stronger excitation we, and others (Molnár et al., 2008), observe in human synapses, particularly onto inhibitory cells. Disynaptic inhibition is often mediated by an interposed Sst cell (Silberberg and Markram, 2007; Obermayer et al., 2018) as they have a low spiking threshold and receive facilitating inputs from excitatory cells. However, the latency of Sst-mediated disynaptic inhibition is often long (> 100 ms) whereas we saw disynaptic IPSPs with a much shorter latency (3 - 6 ms). This suggests the disynaptic inhibition is driven by an intermediate fast-spiking Pvalb cell which, while less common, has been observed in human and can be recruited by very large excitatory events (Szegedi et al., 2017). The unidirectional nature of this disynaptic inhibition from more superficial to deeper cortex further suggests a preferential routing of information by Pvalb cells in human cortex (Szegedi et al., 2017).

Excitatory dynamics depend on target; inhibitory dynamics depend on source

We have taken two complementary approaches to describe the dynamic behavior of synapses. In the first, we take advantage of repeated stimuli of varying temporal structure in order to measure the effects of short-term plasticity and quantal variance. These metrics are generally

easy to interpret but have several drawbacks: they are sensitive to noise, they require successfully repeated stimuli that are not available for all synapses, and they are difficult to use in a biophysical modeling context. In the second approach we address these issues with a new generative model describing quantal release and STP. With this model we are able to use all response data available for each synapse and yield a consistent set of parameters with a clear path to model implementation. Our approach is similar to other recently developed models (Barri et al., 2016; Bird et al., 2016) in that it does not depend on any particular stimulus structure (aside from having a diversity of interspike intervals), and thus frees the experimenter to design stimuli by other criteria and gracefully handles quality issues such as spike failures and early experiment termination.

The model output for each synapse is a multidimensional map of the model likelihood measured across a large parameter space. Each map contains a signature that is unique to the synapse, but shares some features with similar synapses. By reducing the dimensionality of this model output, we were able to ask whether there exists a natural organization among the different types of synapses in our dataset. A few governing principles emerge from this analysis. We find that cortical synapse types can be organized into a two-dimensional feature space, with synaptic strength and variability forming two orthogonal gradients. For the most part, synapses occupy a continuum across these two features. Short-term plasticity features also vary in parallel with variability, but in a discontinuous manner compared to the smoother gradient of variability. In the reduced-dimensionality map, we find that excitatory synapses are strongly differentiated across postsynaptic cell type, whereas inhibitory synapses are differentiated by their presynaptic cell type. This result is consistent with our analyses of the synaptic strength, STP, and variability as they relate to cell subclass, and is similar to results reported by Gupta et al. (2000). In all cases, the subclasses differ most in their synaptic variability, while having largely overlapping distributions of synaptic strength.

Synaptic variability has, in the past, been regarded as an undesirable consequence of signaling via metabolically expensive exocytosis. More recently, advances in machine learning that rely on stochasticity have bolstered the possibility that synaptic variability may offer computational benefits (Llera-Montero et al., 2019). If that is the case, it is further plausible that variability may be modulated by cell type, and that these relationships are crucial features of cortical function. Indeed, our measurements of synaptic variability were found to strongly differentiate cell type in a pattern that is largely, but not entirely, aligned with STP metrics, suggesting the possibility of cell type-specific tuning of variability. More broadly, analysis of our stochastic release model indicates that synaptic strength and variability form the two most important parameters describing synapse behavior. It remains to be explored whether variability itself is an objective parameter directly tuned by the cortex, or if this is a consequence of the relationship between variability and short term plasticity.

Future work

In the near term, we are excited the incorporation of our data into biophysically realistic network models to address questions related to cell-type and synaptic dynamics. With the recent advancements in the segmentation and annotation of large volumes from electron-microscopy (Turner et al., 2020), the opportunity to reconcile connectivity measurements from functional studies like ours with anatomical measurements is on the horizon. The combination of patchSeq (Gouwens et al., 2020; Lee et al., 2020) and multipatch experiments creates the opportunity to gain insights into local connectivity among transcriptomic cell types. A straightforward path is the continuation of the pipeline to explore other brain regions' microcircuitry to reveal the extent to which circuits are either conserved or specialized. A deeper understanding of brain function will require data on the relationships between cell types, learning rules, and modulatory pathways.

Materials and methods

Methods were similar to Seeman, Campagnola et. al, 2018. Information on the Synaptic Physiology pipeline and the dataset are accessible from our website (Synaptic Physiology Coarse Matrix Dataset). Full Standard Operating Procedures can be found: https://www.protocols.io/workspaces/allen-institute-for-brain-science/publications?categories=multipatch

Animals and tissue preparation

All animal procedures were approved by the Institutional Animal Care and Use Committee at the Allen Institute for Brain Science (Seattle, WA), which operates per National Institutes of Health guidelines. Triple (T.L.D., unpublished) and quadruple (Daigle et al., 2018) mouse lines generated using double transgenic mouse lines, were used to target up to two unique cell subclasses in a single animal (see Table S1, http://portal.brain-map.org/explore/toolkit/mice). Each subclass was selectively labeled by fluorescent reporters (tdTomato or EGFP) driven by Cre or FlpO. Layer-specific excitatory cells were targeted using unique transgenic drivers: Nr5a1 and Rorb for layer 4, Sim1, and Tlx3 for layer 5 ET and IT, respectively, and Ntsr1 for layer 6 CT. It was generally not possible to generate crosses of two excitatory drivers, however in L5 we were able to target ET cells via a retroorbital injection of mscRE4-FlpO AAV PHPe.B (Graybuck et al., 2021) into Tlx3-Cre transgenic mice in order to probe interconnections of L5 ET and IT cells. Transgenic lines were not used to target layer 2/3 excitatory cells but were later confirmed through the presence of dendritic spines via post-hoc morphological analysis (see Morphology and Position). Inhibitory cell subclasses, Sst, Pvalb, and Vip, were targeted in all layers.

Female and male adult mice (mean age 46.0 ± 4.6; SD) were anesthetized with 5% isoflurane and transcardially perfused with ice-cold oxygenated slicing aCSF I. All aCSF recipes are in Table S2.

Acute parasagittal slices (350 µm) were produced with a Compresstome (Precisionary Instruments) or VT1200S Vibratome (Leica Biosystems) in ice-cold aCSF I solution. The slicing angle was set to 17° relative to the sagittal plane to preserve pyramidal cells' apical dendrites. Slices were then recovered for 10 min in a holding chamber containing oxygenated aCSF I maintained at 34°C. After recovery, slices were kept in room temperature oxygenated aCSF IV (Table S2).

Human neocortical tissue from Temporal, Frontal, and Parietal lobes was obtained from adult patients undergoing neurosurgery for the treatment of epilepsy or tumor related symptoms (Fig 1A). Tissue obtained from surgery was distal to the core pathological tissue and was deemed not to be of diagnostic value. Surgical specimens were placed in a sterile container filled with pre-chilled (2-4°C), carbogenated aCSF VII containing decreased sodium replaced with NMDG to reduce oxidative damage (Table S2), and delivered from the surgical site to the laboratory within 10-40 min.

In the laboratory, specimens were trimmed to isolate regions of interest and mounted to preserve intact cortical columns (pial surface to white matter) before being sliced in aCSF VII using a Compresstome or Vibratome. Slices were then transferred to oxygenated aCSF VII (34°C) for 10 min, then moved and kept in aCSF VIII at room temperature (Table S2) for a minimum of one hour prior to recording.

Electrophysiological recordings

Slices were placed in custom recording chambers perfused (2-4 mL/min) with aCSF IX which contained one of two external calcium concentrations ($[Ca^{++}]_e$) 1.3 mM or 2.0 mM (Table S2). aCSF IX in the recording chamber was measured at 31-33°C, pH 7.2-7.3, and 30-50% oxygen saturation. In our previous study (Seeman et al., 2018), we conducted experiments in mouse with a $[Ca^{++}]_e$ of 2 mM to be consistent with previous connectivity studies (Markram et al., 1997; Reyes and Sakmann, 1999; Perin et al., 2011; Jiang et al., 2015). However, external calcium concentration *in vivo* has been measured to be closer to 1 mM (Jones and Keep, 1988) and more closely reproduces *in vivo*-like short term plasticity *in vitro* (Borst, 2010). Thus, we reduced $[Ca^{++}]_e$ to 1.3 mM to measure synaptic properties closer to physiological conditions. However, when we compared connection probability, strength, and short-term plasticity for connection elements in which we had data at both $[Ca^{++}]_e$ (21 elements for connectivity and 10 elements for synaptic properties out of 89 targeted intralaminar elements), we found connectivity and synapse characteristics were consistent between the two concentrations (Fig S8). Thus for results reported in this study, data were pooled across conditions. Human experiments continued were conducted with 1.3 mM $[Ca^{++}]_e$ only.

Recording pipettes (Sutter Instruments) were pulled using a DMZ Zeitz-Puller (Zeitz) to a tip resistance of 3-8 MΩ and filled with internal solution containing (in mM): 130 K-gluconate, 10 HEPES, 0 (human) or 0.3 (mouse) ethylene glycol-bis(β-aminoethyl ether)-N,N,N',N'-tetraacetic acid (EGTA), 3 KCl, 0.23 Na2GTP, 6.35 Na2Phosphocreatine, 3.4 Mg-ATP, 13.4 Biocytin, and either 50 µM Cascade Blue dye (excited at 490 nm), or 50 µM Alexa-488 (excited at 565 nm).

Internal solution was measured with osmolarity between 280 and 295 mOsm with pH between 7.2 and 7.3. All electrophysiological values are reported without junction potential correction. We removed EGTA from our internal solution for human recordings to be consistent with previous human electrophysiological studies (Molnár et al., 2008, 2016). A small subset of human recordings were conducted with 0.3 mM EGTA. A comparison of connection elements in which we had both EGTA conditions showed consistent connectivity and synaptic properties and thus the data was pooled (Fig S9).

Eight recording headstages were mounted in a semi-circular arrangement around the recording chamber. The pipette holders were fitted with custom shields to reduce crosstalk artifacts. Each headstage was independently controlled using modified triple-axis motors (Scientifica; PatchStar). Recorded signals were amplified (Multiclamp 700B, Molecular Devices) and digitized (50-200 kHz) using ITC 1600 DAQs (Heka). Pipette pressure was controlled using electro-pneumatic control valves (Proportion-Air; PA2193) or, though manual, mouth applied pressure, available for one pipette at a time. Slices were visualized using oblique (Olympus; WI-OBCD) infrared illumination using 40x or 4x objectives on a custom motorized stage (Scientifica) using a digital sCMOS camera (Hamamatsu; Flash 4.0 V2). Acq4 software (acq4.org; (Campagnola et al., 2014)) was used for pipette positioning, imaging, and subsequent image analysis.

Eight neurons (excitatory or inhibitory) were targeted based on cortical layer, somatic appearance, and depth from the slice's surface in experiments from human and mouse tissue. Neurons in transgenic mice were also targeted based on fluorescent reporter expression. Cells were targeted with a depth of at least 40 μm (Fig S1B) from the surface of the slice with automated pipette control assistance. In order to minimize tissue distortion and damage, pipettes moved through the tissue on a trajectory that was collinear with the long axis of the pipette with minimal positive pressure (10 - 40 mBar).

Whole-cell patch-clamp electrophysiological recordings were performed on neurons that formed a stable seal and had a successful break-in. At least two neurons were measured at the same time per recording, with the mean number of simultaneous recordings being 4 for both mouse and human (see Fig 1A for distributions). Recordings were performed with a holding potential set to either -70 mV (to measure excitatory inputs) or -55 mV (to measure inhibitory inputs) and were maintained within 2 mV using automated bias current injection. Data acquisition was collected using Multi-channel Igor Electrophysiology Suite (MIES; https://github.com/AllenInstitute/MIES), custom software written in Igor Pro (WaveMetrics). A 15-18 second intersweep interval to allow the synapse to recover was used. During a sweep, evoked spikes were distributed in time across recordings such that they were separated by at least 150 ms.

To examine short-term plasticity (STP), cells were stimulated in current and voltage-clamp to drive trains of 12 action potentials (Fig 1B) at different fixed frequencies of 10, 20, 50, 100, and 200 Hz with a delay period between the 8th and 9th pulses (Costa et al., 2013) . The delay period lasted 250 ms for all frequencies with additional delay periods (125, 500, 1000, 2000,

4000 ms) for 50 Hz stimulation. Protocols were repeated five times for each stimulation frequency and delay interval. We also delivered a "mixed frequency" stimulus which was composed of 8 action potentials at 30Hz immediately followed by 30 action potentials, whose intervals were a random resequencing of 29 exponentially increasing intervals between 5 and 100 ms. The intervals were fixed across sweeps and experiments. While in current-clamp, an additional set of stimuli was used to characterize intrinsic properties of each cell (Fig 1C). To estimate input resistance of the cell, a 1-second-long hyperpolarizing square pulse was delivered at an initial amplitude of -20 pA. The voltage response to each current step was measured online and successive current steps were titrated to target response voltages of +2, -2, -5, -10, and -15 mV so as to reliably activate I_h when present. To measure spiking properties, a long (500 ms) depolarizing square pulse stimulus was delivered that started at rheobase and increased 25 pA for 6 intervals. Lastly, we delivered a 15-second sinusoidal chirp that increased in frequency from 0.2 to 40 Hz and evoked a response magnitude that measured ~10 mV from peak to trough.

PatchSeq recordings and processing

PatchSeq recordings were performed in a subset of mouse experiments. To avoid sample contamination, surfaces, equipment, and materials were cleaned using DNA away (Thermo Scientific), RNAse Zap (Sigma-Aldrich), and nuclease-free water (in that order). aCSF V was made daily and filtered before use. Materials used to make and store aCSF V were cleaned thoroughly before use. Recording pipettes were filled with ~1.75 µL of RNAse Inhibitor containing internal solution: 110 mM K-Gluconate, 4 mM KCl, 10 mM HEPES, 1 mM adenosine 5'-triphosphate magnesium salt, 0.3 mM guanosine 5'-triphosphate sodium salt hydrate, 10 mM sodium phosphocreatine, 0.2 mM ethylene glycol-bis (2-aminoehtylether)-N,N,N',N'-tetraacetic acid, 20 µg/mL glycogen, 0.5 U/µL RNase Inhibitor, 0.5 % biocytin, and either 50 µM Cascade Blue dye (excited at 490 nm), or 50 µM Alexa-488 (excited at 565 nm).

In patchSeq experiments, a subset of stimuli were collected to limit progressive cell swelling associated with the addition of RNAse to the internal solution (Lee et al., 2020).

Methods for nuclei extraction and processing are similar to previous patchSeq studies (Gouwens et al., 2020; Lee et al., 2020). At the end of the experiment, pipettes were adjusted to the soma center or placed near the nucleus, if visible. A small amount of negative pressure (~0.5 psi) was applied to all pipettes simultaneously for cytosol extraction. Extraction time varied for each cell; pipettes were slowly (~0.3 µm/s) retracted in the x and z-axis once the soma had visibly shrunk and/or the nucleus was visible at the tip of the pipette. Once pipettes were out of the slice, cytosol and/or nucleus content in each pipette were expelled into individual PCR tubes containing 11.5 µl of lysis buffer (Takara, 634984) and stored in -80 °C.

We used the SMART-Seq v4 Ultra Low Input RNA Kit for Sequencing (Takara, 634894) per the manufacturer's instructions to reverse transcribe RNA and amplify full-length cDNA. To obtain detailed methods, see http://celltypes.brain-map.org, "Transcriptomics Overview Technical

White Paper". We identified transcriptomic types by mapping our Patch-seq transcriptomes data in the same methods mentioned in previous studies (Tasic et al., 2018; Gouwens et al., 2020).

Histology and imaging

After electrophysiological recordings, slices were fixed in solution containing 4% paraformaldehyde and 2.5% glutaraldehyde for at least 40 hours at 4°C. Slices were then transferred and washed in phosphate buffer saline (PBS) solution for 1-7 days before staining.

A 3,3'-diaminobenzidine (DAB) peroxidase substrate kit (Vector Laboratories) generated a brown reaction product in biocytin-filled neurons. Slices were stained with 5 μM 4',6-diamidino-2-phenylindole (DAPI) in PBS for 15 min at room temperature and then triple-washed in PBS (10 min for each wash). Slices were then transferred to 1% hydrogen peroxide (H2O2) in PBS for 30 min and triple washed in PBS. Afterward, slices were mounted onto gelatin-coated slides and cover-slipped with Aqua-Poly/Mount (Polysciences). Slides were dried for approximately 2 days before imaging.

Mounted slides were imaged on an AxioImager Z2 microscope (Zeiss) equipped with an Axiocam 506 monochrome camera. Tiled mosaic images of whole slices were captured with a 20x objective lens (Zeiss Plan-NEOFLUOR 20x/0.5) to generate both biocytin-labeled images and DAPI-labeled images. Biocytin images were used to assess cell morphology and DAPI images were used to identify cortical layer boundaries. To further classify cell morphology, we also used a 63x lens to capture high-resolution z-stacks of biocytin filled cells, which were stitched together using ZEN software and exported as single-plane TIFF files.

Synapse detection

Electrophysiology data was filtered through multiple quality control (QC) steps to ensure it was of good quality to detect a connection (Table S3). Additional QC criteria were applied for characterizing synapses which are discussed in the following section.

Initial data processing consisted of manual synapse detection and curve fitting (Figure 1B *Synapse Processing*) using our [Pair Analysis Tool](#) (Fig S10). During recording, cells were held at -70mV to probe excitatory connections or -55mV to probe inhibitory connections. Within each clamp mode data was binned into two ranges of recorded membrane potentials [-80, -61] (-70mV holding potential) and [-60, -45] (-55mV holding potential). Within each of these four groups, individual postsynaptic responses (PSC/P, Fig S10 white traces) from stimuli <=50 Hz were aligned to the peak rate of rise of the presynaptic spike and averaged (Fig S10A blue traces). Only QC passed PSC/Ps were included in the average. Users visually identified chemical and electrical synapses from the average postsynaptic responses in each of the four quadrants and marked whether the synapse was excitatory or inhibitory (Fig S10). Manual connectivity calls were used to train a machine classifier (Seeman et al., 2018), which was used to reveal possible false positives or false negatives that were later manually corrected.

Synapse characterization

Users manually identified the onset of detected synapses' postsynaptic response (Fig 1B, Fig S10, yellow line). This user-defined latency was used to initialize an automated curve fit of the average response constrained to ± 100 µs of the user-defined latency (Fig S10B red and green traces). The user inspected the automated fit for a good match to the average and could refit as necessary. Once the user was satisfied with the output, the fit was manually passed (Fig S10B green) or failed (Fig S10B red). Contamination by electrical synapses or electrical artifacts, the shape of the fit, and other factors were considered when deciding whether to pass or fail the fit. Users were also able to make notes, which were used to update and test new fitting algorithms.

Passing postsynaptic response fits were used to characterize strength, kinetics, and short-term plasticity of the synapse in a multi-stage process aimed to maximize data inclusion for each characteristic. Kinetics,rise time and decay tau,were measured as a weighted average of curve fit parameters from the two membrane potential ranges. Latency was similarly measured as a weighted average across not only membrane potentials but clamp mode as well as we did not expect PSC/P onset to be influenced by these conditions. Latency values from the individual membrane potential/clamp mode modalities were confirmed to be within 200µs of each other to be included in the weighted average. Thus, each synapse has a singular latency value. PSC/P amplitude was measured at a "resting state" to avoid the influence of short-term plasticity in our estimate of strength. Individual PSC/Ps recorded at the appropriate holding potential for the synapse (-55mV for inhibitory, -70mV for excitatory) and preceded by at least 8 seconds of quiescence were aligned to the presynaptic spike and averaged. This reduced average was fit with kinetic parameters initialized to those from the fit to the average of all responses. For synapse types that show short-term facilitation, the resting state amplitude is often very small and thus, we calculated a second metric to capture the near maximum strength that a synapse can produce. For this, individual PSPs were fit with parameters initialized by results of the average fit. We then calculated the 90th percentile of fit amplitudes to approximate maximum strength (this value was also used to normalize our estimate of short-term plasticity discussed below).

Short-term plasticity (STP) was measured from current-clamp PSPs in response to trains of stimuli at multiple frequencies consisting of eight pulses, followed by a variable delay, and then four more pulses (Fig 1B *Multipatch Experiment*, *Synapse Analysis*). Typically a ratio of pulse amplitudes is calculated to determine depth of STP and recovery from STP. However, for small responses close to the noise threshold or synapses that strongly facilitate or depress, this ratio can cause spurious STP measurements. To correct for this, we normalized STP measurements by the 90th percentile amplitude (see above). The depth of short-term depression or facilitation was measured from the amplitude of individual PSP fits as follows (Fig 1B *Synapse Analysis*):

(Avg(6th, 7th, 8th pulse amplitudes) - 1st pulse amplitude) / 90th percentile amplitude

By this calculation positive values denote facilitating synapses and negative values depressing.

Recovery from STP was similarly calculated from individual PSP fits as:

((Avg(9th-12th pulse amplitudes) - Avg(1st-4th pulse amplitudes)) / 90th percentile.

This yields positive values indicative of recovery beyond the initial state of the synapse and negative values where the synapse has not yet recovered from STP.

PSP/C variability is typically reported as the coefficient of variation (CV) of response amplitudes; however, for weak synapses the CV is dominated by noise and thus does not reflect a property of the synapse. To compensate for this we subtract the amplitude of noise before normalization to yield an "adjusted coefficient of variation" (aCV):

$$aCV(amps,\ noise)\ =\ \frac{\sqrt{\sigma_{amps}^2 - \sigma_{noise}^2}}{\mu_{amps}}$$

Where μ_{amps} and σ_{amps} are the mean and standard deviation of response amplitudes, and σ_{noise} is the standard deviation of background noise, which is measured by performing the same amplitude measurement algorithm on regions of the recording that have no presynaptic stimulus.

The strength of electrical synapses (gap junctions) were quantified as a coupling coefficient and junctional conductance (Galarreta and Hestrin, 1999; Gibson et al., 1999). The voltage change from baseline evoked by a subthreshold long-pulse current injection (Fig 1C) was measured in both the pre- and postsynaptic cell. The coupling coefficient was measured as a least squares linear regression of the voltage change across all sweeps. For cells in which we also measured input resistance the junctional conductance was calculated as G_j = (1/R2) x CC/(1-CC) (Tasic et al., 2016) where CC is the coupling coefficient and R2 is the input resistance of the postsynaptic cell.

Cell characterization

Transgenic Expression

Stack images of the recording site were taken in brightfield, epifluorescence (tdTomato and EGFP), and dye-filled recording pipettes. These images were filtered and overlaid on top of each other (Fig 1B *Multipatch Experiment*) to display the recording site and targeted cells. From the overlap of epifluorescence and pipette dye, we identified each cell's transgenic expression, which was used to define its subclass. If fluorescence overlap could not be confirmed, the expression was marked as unknown. All mouse L2/3 excitatory cells and human cells were fluorescence-negative cells and thus, morphological features were used to identify these cells as discussed below.

Electrophysiology

Intrinsic characterization of individual cells was carried out similarly to that described in (Anon, n.d.). Although we did not collect the full suite of stimuli, the long-pulse sweeps we acquired were sufficient to calculate subthreshold properties such as input resistance, sag, and rheobase; spike train properties such as f-I slope and adaptation index; and single spike properties such as upstroke-downstroke ratio, after-hyperpolarization, and width (Fig1C *Intrinsic Ephys*). For spike upstroke, downstroke, width, threshold, and ISI, 'adaptation ratio' features were calculated as a ratio of the spike features between the first and fifth spike. A subset of cells also had subthreshold frequency response characterized by a logarithmic chirp stimulus (sine wave with exponentially increasing frequency), for which the impedance profile was calculated and characterized by features including the peak frequency and peak ratio. Feature extraction was implemented using the IPFX python package (https://github.com/AllenInstitute/ipfx); custom code used for chirps and some high-level features will be released in a future version of IPFX.

For the human cell dataset, all electrophysiology features were aggregated and visualized using a UMAP projection (McInnes et al., 2018) to gain perspective on the electrophysiological cell type ('e-type') distinctions present. Cells with more than 25% missing features were dropped. The remaining missing features were imputed as a distance-weighted mean of 3 nearest neighbors, and each feature was independently power transformed to a standard Gaussian. Features uninformative for known cell-type distinctions were dropped (assessed by F-score of ANOVA against layer and spininess labels), and the remaining features were visualized by UMAP projection. The e-type distinction made among L2-3 cells, distinguishing 'L4-like' cells with high input resistance, was made via a threshold on the raw feature value but assessed in the full feature space using this visualization.

Morphology and Position

Cell morphology was qualitatively assessed from 63x maximum projection image z-stacks of biocytin filled cells and included features such as dendritic type (spiny-ness), length of truncated axon, and axon origin point of inhibitory cells (Fig 1C *Morphology*). Aspiny or sparsely spiny cells (inhibitory) were defined as such if their dendrites lacked or only had few protrusions. Spiny (excitatory) cells have frequent dendritic protrusions as well as an apical dendrite. (Braitenberg and Schüz, 1998; Gouwens et al., 2019). For the purposes of cell classification throughout our results, "spiny-ness" refers to this analysis. Layer 2/3 pyramidal cells in mouse were largely identified as being "spiny" as we did not have a transgenic driver for this layer. Similarly in human, all cells are identified as excitatory or inhibitory by their dendritic spiny-ness and by the presence/absence of an apical dendrite. A full list of morphological classification can be found in Table S4.

Cortical layer boundaries were determined from DAPI images, with the top of Layer 1 serving as a marker of pia and the bottom of Layer 6 as a marker of white matter. During the experiment, cell position was recorded in the fluorescent images' reference space (Fig 1B *Multipatch Experiment*) and later coregistered with the DAPI image. Image coregistration enabled the cell

soma layer to be established. Other positional metrics such as distance from pia and white matter, fractional cortical depth, and depth within the layer were also calculated from the soma position and layer boundary data using the neuron_morphology python package (https://github.com/AllenInstitute/neuron_morphology). Depth measurements were made using streamlines from the pia to WM boundaries (or the nearest layer boundary in cases where not all layers were complete in the slice). For each cell pair, the pia-WM orientation from the streamlines was averaged and used as a 'vertical' orientation to decompose the soma-soma separation into vertical and lateral distances.

Connection probability estimation

We estimated the connection probability through modeling a probability distribution that depends on the experimental conditions such as the distance between the neuron pair, depth of the neurons in slices, and signal-to-noise ratio. To build a model for the connection probability, we started with a log-likelihood function of the binomial distribution, constructed its probability using multiple experimental variables, and estimated model parameters using maximum-likelihood estimation (MLE).

The log-likelihood function for the binomial distribution is defined as follows.

$$l(\theta|x) = \sum_{i=1}^{n_{conn}} \log\left(p(\theta|x_{conn_i})\right) + \sum_{j=1}^{n_{unc}} \log\left(1 - p(\theta|x_{unc_j})\right),$$

where θ is a set of model parameters, x is a set of experimental variables associated with each pair, p is the model estimate of the connection probability, n_{conn} and n_{unc} are the numbers of connected and unconnected pairs, x_{conn} and x_{unc} are subset of x for connected and unconnected pairs, respectively. The first sum runs over connected pairs; the second sum runs over unconnected pairs. Below, we elaborate how this probability function is constructed.

Gaussian model with maximum likelihood estimation

A full description of how we arrived at a Gaussian model of connection probability can be found in the accompanying notebook and is summarized here.

Connection probability as a function of intersomatic distance of a pair was modeled as a Gaussian function centered at 0 distance:

$$p(\theta|x) = p_{max} e^{-\frac{d^2}{2\sigma^2}},$$

where $\theta = \{p_{max}, \sigma\}$ and $x = \{d\}$, p_{max} is the peak connection probability, σ is the distance constant of connection probability, and *d* is the lateral intersomatic distance of the pair.

The number of samples (pairs of cells) included in the model has a profound impact on our confidence in the model outputs, p_{max} and σ. We used simulated distances drawn from the distribution of measured intersomatic distances and a defined p_{max} and σ to assess the possible error in our fits. We found that while p_{max} was fairly well constrained, σ was not, particularly for lower numbers of samples. We therefore chose to use a fixed σ value, which further constrained the fit of peak connection probability. To determine the fixed σ value, we pooled our data from mouse into four categories based on the cell class of the presynaptic and postsynaptic cells, namely excitatory to excitatory (5,467 cell pairs), excitatory to inhibitory (2,161 pairs), inhibitory to excitatory (1,973 pairs), and inhibitory to inhibitory (6,100 pairs). This allowed us to have several thousand samples in each group to obtain a better estimate of σ (Fig 2A). In order to more fully determine if the σ of these four groups could have been drawn from the same distribution we simulated 10,000 experiments with a true σ that varied between those measured from experimental data. We then did a pair-wise comparison of the σ ratio for each unique comparison among the four groups and calculated the percentile of this distribution where the measured ratio fell. When correcting for multiple comparisons we found that while the I→I and I→E connectivity profiles likely have σ that are not drawn from the same distribution, we could not rule that out for the other comparisons. This analysis was also conducted for a comparison of chemical versus electrical connections among inhibitory cells, with 1,000 simulated trials. From this analysis we chose to fix σ to 100 µm for chemical synapses in order to constrain our estimate of p_{max} for individual matrix elements (Fig 2B). A similar procedure was conducted for human data, resulting in a fixed σ of 140 µm (Fig 5A).

A unified model of connection probability adjustment

We extended the analysis of the connection probability as a function of intersomatic distance to include the effect of tissue slicing and false negative detection of connections due to signal-to-noise ratio. We created a unified model that applies these adjustments to the connection probability, as a function of the pair distance (discussed above), the presynaptic axon length, the depth of the cells, and the detection power.
Using the same log-likelihood function described above, we extended the probability to the following.

$$p(\theta|x) = p_{max} e^{-\frac{d^2}{2\sigma^2}} \prod_k C_k(\theta_k|x_k)$$

where $\theta = \{p_{max}, \sigma, \rho_{ax}, \mu_{depth}, \sigma_{depth}, \mu_{det}, \sigma_{det}\}$, $x = \{d, l_{ax}, z_{depth}, p_{det}\}$, and product over k runs over the following three corrections factors. The correction functions C_k, their parameters θ_k, and their variables x_k are described below.

The model for the presynaptic axon length is a binary step function with a threshold at 200 μm as axons were not measured past this point (Table S4). Also, because the number of neurons with axons measured less than 200 μm were few, we did not have sufficient data to determine the function shape below this threshold. Therefore, we used a single adjustment ratio for the correction.

$$C_{ax}(\rho_{ax}|l_{ax}) = 1 \ (l_{ax} > 200 \ \mu m), \ or \ \rho_{ax} \ (l_{ax} \leq 200 \ \mu m)$$

The model for the average depth of the pair of neurons is as follows.

$$C_{depth}(\mu_{depth}, \sigma_{depth}|z_{depth}) = \frac{1}{2}\left(1 + erf\left(\frac{z_{depth}-\mu_{depth}}{\sqrt{2}\sigma_{depth}}\right)\right),$$

where erf is an error function:

$$erf(x) = \frac{2}{\sqrt{\pi}}\int_0^x e^{-t^2} dt.$$

The detection power (that is proportional to signal-to-noise ratio) is defined as $p_{det} = \sqrt{N_{pre}}/\sigma_{post}$, where N_{pre} is the number of presynaptic test spikes and σ_{post} is the RMS noise-level of the postsynaptic neuron. The model for the detection power p_{det} is also an error function, but in a log space, because the synaptic weight distribution is expected to be a log-normal distribution.

$$C_{det}(\mu_{det}, \sigma_{det}|p_{det}) = \frac{1}{2}\left(1 + erf\left(\frac{\log_{10}(p_{det})-\mu_{det}}{\sqrt{2}\sigma_{det}}\right)\right).$$

We did not see a saturation of the connection probability when the presynaptic cell was inhibitory (Fig S1C), suggesting that there are potentially a large number of undetected synapses. However, we did not want to overestimate the connection probability in the range of detection power where we did not have sufficient data. Therefore, we applied the following constraint to the fit.

$$\mu_{det} + \alpha\sigma_{det} < \beta$$

Where $\alpha \sim 0.6745$, specifying the quartile of the integrand Gaussian of the error function, and $\beta \sim 4.6613$, specifying the quartile of the detection power distribution in our data. This constraint ensures saturation in the high detection power region where data are scarce.

The models and the parameters for these three corrections are determined individually for each variable (Fig S1A-C) and incorporated into the likelihood function used for the distance adjustment. When we estimated final p_{max}, we performed a single-parameter MLE, fixing all the other model parameters to pre-determined values.

Estimating connection probability and confidence intervals

The goal of optimizing the Gaussian MLE was to compare connection probabilities across different cell groups that may have been sampled at different intersomatic distances and have variable presynaptic axon lengths and detection power. Thus, in addition to fitting p_{max} we wanted to calculate a confidence interval of connection probability. Our initial approach was to analyze hundreds of resampled iterations of data from each matrix element. However, this method is computationally expensive and starts to break down when connection probability or the number of samples is very low. We determined that through a derivation of the MLE equation that p_{max} is largely driven by the distribution of probability adjustment factors for all probed connections and the number of connections found, rather than the particular adjustment factor value of each connected pair. This suggests that we can forego the full connection probability model for each element and instead estimate connection probability as a binomial distribution scaled by the same adjustment factors. While the binomial confidence interval is a conservative estimate of confidence in the fit p_{max} it is relatively easy to implement, conceptually simple, and handles cases near the limits of p_{max} well. Thus, p_{max} is calculated as the proportion of connections (number of connections found / number of connections probed) with a binomial confidence interval, both multiplied by the same scale factor. In this case, the scale factor is the mean amplitude of a Gaussian sampled at the intersomatic distances of each element with a σ fixed to 100 µm for mouse and 140 µm for human. Scaled binomial confidence intervals were used to shade connection probability heatmaps in Figure 2B, 5A, and supplemental Figure 2A.

The parameter optimization can result in values for p_{max} (or CI bounds) greater than 1 in cases where the data is not well fit by the fixed-width Gaussian, typically because a class of connections has either low sampling or a true underlying connection probability function with a distinct shape or size. In the resulting figures, p_{max} values and CI upper bounds were clipped at 1, reflecting the fact that such data should be better fit by a flat-topped curve that approaches 1 at its maximum (Fig. S2B). The unclipped values are available in our data and code release for applications like modeling, where estimating the true connection probability vs. distance function may be more important than interpretation of p_{max} as a probability.

Modeling synapse behavior

Stochastic quantal release model

We developed a model of stochastic vesicle release and synaptic dynamics that expands upon standard models (Hennig, 2013) and is similar to some recent models (Barri et al., 2016; Bird et al., 2016; Bykowska et al., 2019). The model was designed to meet several criteria. First, it should give estimates of synaptic quantal parameters (number of release sites, probability of

release, and quantal size) and dynamic parameters (facilitation, docking, etc.). The quantal parameters alone allow the model to predict the overall distribution of response amplitudes (Fig S11A). By allowing quantal parameters to change in response to the history of activity, the model is able to account for changes in the response distribution (Fig S11B,C). Next, the model should operate on individual spike times and response amplitudes rather than requiring structured or repeated stimuli. This ensures that the model can access the complete distribution of response amplitudes and any correlations between adjacent responses. Additionally, operating on individual events rather than averages ensures that we use all available data for each synapse, regardless of the applied stimuli. Finally, the model should fail gracefully in cases with low signal-to-noise ratio by indicating low confidence over its parameters rather than returning unreliable values.

We begin with a standard quantal model with three parameters: N, the number of release sites; q, the amplitude of the postsynaptic response to a single vesicle, and P_r, the probability that each release site will release a vesicle in response to one action potential. We make simplifying assumptions that all release sites in a connection share the same values of P_r and q, and that the response to multiple vesicles released simultaneously is simply the linear sum of individual responses. This component of the model simply predicts that the number of vesicles released per spike is defined by a binomial distribution with parameters N and P_r, and that the distribution of response amplitudes is the same with an additional scaling factor q.

The measured response amplitudes in our dataset only occasionally show binomial characteristics; however in most cases, background recording noise and quantal variability obscure the underlying discrete distribution. We model these sources of variability as independent Gaussian distributions for measurement noise and quantal variance, with parameters σ_m and σ_q, respectively. These combine with the binomial distribution to make a weighted Gaussian mixture model that, on its own, does a decent job of approximating the overall distributions of event amplitudes. The model response amplitude probability distribution $P(X; \theta)$ is calculated as:

$$normpdf(\mu, \sigma, x) = \sqrt{\frac{1}{2\pi\sigma^2}}e^{\frac{-(x-\mu)^2}{2\sigma^2}}$$

$$binompmf(n, p, k) = \frac{n!}{k!(n-k)!}p^k(1-p)^{n-k}$$

$$P(X; \theta) = \sum_{k=0}^{N} binompmf(N, P_r, k)\, normpdf(kq, \sqrt{k\sigma_q^2 + \sigma_m^2}, X)$$

Short term plasticity

Synapses undergo short term plasticity, which we model as changes in the expected distribution of response amplitudes over time (Hennig, 2013). Each incoming action potential causes an instantaneous modification to the quantal parameters N and P_r, which then recover back to their initial values by exponential decay until the time of the next action potential. Although we only seek a phenomenological description of the synapse, this description is designed to mimic three

major classes of synaptic dynamics: vesicle depletion, facilitation, and calcium channel inactivation. Vesicle depletion is implemented by using the response amplitudes and *q* to estimate the most likely number of released vesicles following each spike, which is then subtracted from the releasable vesicle pool. Recovery from vesicle depletion is modeled as an exponential with time constant τ_r. Facilitation and calcium channel inactivation increase or decrease the release probability, respectively, for every incoming spike, and decay back to the initial release probability with time constants τ_f and τ_i. In addition to the decay time constants, these mechanisms introduce an extra two parameters: the amount of facilitation a_f and inactivation a_i per spike.The complete algorithm then looks like:

1. Initialize state variables:
    a. *$N_j = N_r$*
    b. *depression = 0*
    c. *facilitation = 0*
2. For each spike *j* at time *t_j*:
    a. Let *$dt = t_j - t_{j-1}$*
    b. Recover state variables:
        i.   *N_j += (N_r - N_j) * (1 - $e^{-dt/\tau D}$)*
        ii.   *depression *= $e^{-dt/\tau D}$*
        iii. *facilitation *= $e^{-dt/\tau F}$*
    c. Let *P_j = (1 - depression) * (P_r + (1 - P_r) * facilitation)*
    d. Define distribution parameters *θ = {N_j, P_j, q, σ_q, σ_m}*
    e. Estimate likelihood of measured response amplitude *$P(Amp_j | \theta)$* or generate a random sample drawn from *$P(\theta)$*
    f. Apply post-spike modifications to state variables:
        i.   *vesicle_pool -= Amp_j / q*
        ii.   *depression += a_D * (1 - depression)*
        iii. *facilitation += a_F * (1 - facilitation)*

Occasionally a presynaptic stimulus is not followed by a detectable spike, which could be caused by a genuine spike failure or simply by a failure of the spike detection algorithm. In these cases, we assume that a spike did occur for the purpose of updating the model state variables, but we incur a short timeout during which we stop accumulating evidence toward the overall model likelihood.

For any combination of parameters, we estimate a *likelihood* that a recorded set of amplitudes $A_{1..m}$ could be generated by the model by the mean log probability density for each response:

$$L(A) = \frac{1}{m} \sum_{j=1}^{m} ln(P(A_j))$$

If most response amplitudes fall within the modeled regions of high probability density, then the total likelihood for that parameter set will be high. Low-likelihood models result from either bad

parameters (where response amplitudes fall in low density regions of the model probability distribution) or from insufficiently selective parameters (where the probability distribution is spread out over too much area).

Parameter search and optimization

With a measure of model likelihood, we can now attempt to find a set of parameters that maximize this value, yielding a model that best explains the data. Most prior methods use a metric similar to the likelihood defined above along with an optimization method to efficiently find a single point in the parameter space that is most consistent with the recorded data. However, minimization is notoriously difficult in this domain because the model parameters are underconstrained--there exist many solutions that adequately explain the recorded data, and thus large differences in the optimal parameters may simply result from noise or experimental artifacts, rather than physiological differences between synapses (Bykowska et al., 2019; Gontier and Pfister, 2020). To avoid this outcome, we measure the model performance at every point in a large parameter space, thereby identifying the region of the parameter space consistent with the responses recorded from each synapse (Figure S11D). This is similar to some recent methods (Barri et al., 2016; Bird et al., 2016), but differs in that we have implemented a simpler (and thus less computationally expensive) model in order to test many more parameter combinations uniformly.

All combinations of the parameters in a 7-dimensional space were tested, for a total of 6.2 M model tests per synapse. At every point in this parameter space, the quantal amplitude q was estimated and optimized using a minimization algorithm (scikit.optimze). Although this minimization strategy fails to find reliable optima when operating over several parameters, we found it to be reliable in the context of this simpler single-variable optimization. The resulting records for each synapse thus include measured model likelihood as well as the optimal value of q at each point in the parameter space. This yields an 8 dimensional image that serves as a "fingerprint" describing the unique dynamics of each synapse. The size of the parameter space and the number of synapses in our dataset together make this a computationally expensive operation. To reduce this cost, we optimized the core routines used in the model using the python packages numpy and numba.

Synapse typing

To visualize the relationship between cell subclass and synapse properties, we used the UMAP dimensionality reduction method(McInnes et al., 2018) to organize our synapses into a 2 dimensional space for visualization. Features for each synapse, the release model described above is run on a large parameter space, yielding 6.2 million "features" that collectively describe the regions of parameter space that are (and the regions that are not) compatible with the recorded data. We then use sparse PCA to reduce this down to 50 features. This set of features provides a fingerprint of any information available to the release model, including synaptic strength, stochasticity, and short-term plasticity. Next we extend the feature list to include latency and kinetic features (rise time and decay tau) from both current- and voltage-clamp,

since the model does not have access to these. Finally, all features are normalized (scikit-learn), then passed to UMAP for the final dimensionality reduction. The resulting 2D space can then be visualized alongside other synaptic and cellular features to investigate the structure inherent in the data.

References

Abbott LF, Regehr WG (2004) Synaptic computation. Nature 431:796–803.

Alcamí P, Pereda AE (2019) Beyond plasticity: the dynamic impact of electrical synapses on neural circuits. Nat Rev Neurosci 20:253–271.

Almási Z, Dávid C, Witte M, Staiger JF (2019) Distribution Patterns of Three Molecularly Defined Classes of GABAergic Neurons Across Columnar Compartments in Mouse Barrel Cortex. Front Neuroanat 13:45.

Anon (n.d.) Allen Cell Types Database Technical White Paper: Electrophysiology. Available at: http://help.brain-map.org/display/celltypes/Documentation.

Balaram P, Kaas JH (2014) Towards a unified scheme of cortical lamination for primary visual cortex across primates: insights from NeuN and VGLUT2 immunoreactivity. Front Neuroanat 8:81.

Barri A, Wang Y, Hansel D, Mongillo G (2016) Quantifying Repetitive Transmission at Chemical Synapses: A Generative-Model Approach. Eneuro 3:ENEURO.0113-15.2016.

Beierlein M, Gibson JR, Connors BW (2003) Two Dynamically Distinct Inhibitory Networks in Layer 4 of the Neocortex. J Neurophysiol 90:2987–3000.

Berg J, Sorensen SA, Ting JT, Miller JA, Chartrand T, Buchin A, Bakken TE, Budzillo A, Dee N, Ding S-L, Gouwens NW, Hodge RD, Kalmbach B, Lee C, Lee BR, Alfiler L, Baker K, Barkan E, Beller A, Berry K, Bertagnolli D, Bickley K, Bomben J, Braun T, Brouner K, Casper T, Chong P, Crichton K, Dalley R, Frates R de, Desta T, Lee SD, D'Orazi F, Dotson N, Egdorf T, Enstrom R, Farrell C, Feng D, Fong O, Furdan S, Galakhova AA, Gamlin C, Gary A, Glandon A, Goldy J, Gorham M, Goriounova NA, Gratiy S, Graybuck L, Gu H, Hadley K, Hansen N, Heistek TS, Henry AM, Heyer DB, Hill D, Hill C, Hupp M, Jarsky T, Kebede S, Keene L, Kim L, Kim M-H, Kroll M, Latimer C, Levi BP, Link KE, Mallory M, Mann R, Marshall D, Maxwell M, McGraw M, McMillen D, Melief E, Mertens EJ, Mezei L, Mihut N, Mok S, Molnar G, Mukora A, Ng L, Ngo K, Nicovich PR, Nyhus J, Olah G, Oldre A, Omstead V, Ozsvar A, Park D, Peng H, Pham T, Pom CA, Potekhina L, Rajanbabu R, Ransford S, Reid D, Rimorin C, Ruiz A, Sandman D, Sulc J, Sunkin SM, Szafer A, Szemenyei V, Thomsen ER, Tieu M, Torkelson A, Trinh J, Tung H, Wakeman W, Ward K, Wilbers R, Williams G, Yao Z, Yoon J-G, Anastassiou C, Arkhipov A, Barzo P, Bernard A, Cobbs C, Hamer PC de W, Ellenbogen RG, Esposito L, Ferreira M, Gwinn RP, Hawrylycz MJ, Hof PR, Idema S, Jones AR, Keene CD, Ko AL, Murphy GJ, Ng L, Ojemann

JG, Patel AP, Phillips JW, Silbergeld DL, Smith K, Tasic B, Yuste R, Segev I, Kock CPJ de, Mansvelder HD, Tamas G, Zeng H, Koch C, Lein ES (2020) Human cortical expansion involves diversification and specialization of supragranular intratelencephalic-projecting neurons. Biorxiv:2020.03.31.018820.

Billeh YN, Cai B, Gratiy SL, Dai K, Iyer R, Gouwens NW, Abbasi-Asl R, Jia X, Siegle JH, Olsen SR, Koch C, Mihalas S, Arkhipov A (2020) Systematic Integration of Structural and Functional Data into Multi-scale Models of Mouse Primary Visual Cortex. Neuron.

Bird AD, Wall MJ, Richardson MJE (2016) Bayesian Inference of Synaptic Quantal Parameters from Correlated Vesicle Release. Front Comput Neurosc 10:116.

Blackman AV, Abrahamsson T, Costa RP, Lalanne T, Sjöström PJ (2013) Target-cell-specific short-term plasticity in local circuits. Frontiers Synaptic Neurosci 5:11.

Borst GJ (2010) The low synaptic release probability in vivo. Trends Neurosci 33:259–266.

Braitenberg V, Schüz A (1998) Cortex: Statistics and Geometry of Neuronal Connectivity. :103–107.

Brown SP, Hestrin S (2009) Intracortical circuits of pyramidal neurons reflect their long-range axonal targets. Nature 457:1133–1136.

Bykowska O, Gontier C, Sax A-L, Jia DW, Montero ML, Bird AD, Houghton C, Pfister J-P, Costa RP (2019) Model-Based Inference of Synaptic Transmission. Frontiers Synaptic Neurosci 11:21.

Campagnola L, Kratz MB, Manis PB (2014) ACQ4: an open-source software platform for data acquisition and analysis in neurophysiology research. Front Neuroinform 8:3.

Cardin JA (2018) Inhibitory Interneurons Regulate Temporal Precision and Correlations in Cortical Circuits. Trends Neurosci 41:689–700.

Chao OY, Yang Y-M (2019) Timing constraints of action potential evoked Ca2+ current and transmitter release at a central nerve terminal. Sci Rep-uk 9:4448.

Costa RP, Sjöström JP, Rossum MC van (2013) Probabilistic inference of short-term synaptic plasticity in neocortical microcircuits. Front Comput Neurosc 7:75.

Cruikshank SJ, Lewis TJ, Connors BW (2007) Synaptic basis for intense thalamocortical activation of feedforward inhibitory cells in neocortex. Nat Neurosci 10:462–468.

Dahmen D, Recanatesi S, Ocker GK, Jia X, Helias M, Shea-Brown E (2020) Strong coupling and local control of dimensionality across brain areas. Biorxiv:2020.11.02.365072.

Daigle TL, Madisen L, Hage TA, Valley MT, Knoblich U, Larsen RS, Takeno MM, Huang L, Gu H, Larsen R, Mills M, Bosma-Moody A, Siverts LA, Walker M, Graybuck LT, Yao Z, Fong O, Nguyen TN, Garren E, Lenz GH, Chavarha M, Pendergraft J, Harrington J, Hirokawa KE, Harris

JA, Nicovich PR, McGraw MJ, Ollerenshaw DR, Smith KA, Baker CA, Ting JT, Sunkin SM, Lecoq J, Lin MZ, Boyden ES, Murphy GJ, Costa NM da, Waters J, Li L, Tasic B, Zeng H (2018) A Suite of Transgenic Driver and Reporter Mouse Lines with Enhanced Brain-Cell-Type Targeting and Functionality. Cell 174:465-480.e22.

DeFelipe J, Fariñas I (1992) The pyramidal neuron of the cerebral cortex: Morphological and chemical characteristics of the synaptic inputs. Prog Neurobiol 39:563–607.

DeFelipe J, López-Cruz PL, Benavides-Piccione R, Bielza C, Larrañaga P, Anderson S, Burkhalter A, Cauli B, Fairén A, Feldmeyer D, Fishell G, Fitzpatrick D, Freund TF, González-Burgos G, Hestrin S, Hill S, Hof PR, Huang J, Jones EG, Kawaguchi Y, Kisvárday Z, Kubota Y, Lewis DA, Marín O, Markram H, McBain CJ, Meyer HS, Monyer H, Nelson SB, Rockland K, Rossier J, Rubenstein JLR, Rudy B, Scanziani M, Shepherd GM, Sherwood CC, Staiger JF, Tamás G, Thomson A, Wang Y, Yuste R, Ascoli GA (2013) New insights into the classification and nomenclature of cortical GABAergic interneurons. Nat Rev Neurosci 14:202–216.

de Vries SEJ, Lecoq JA, Buice MA, Groblewski PA, Ocker GK, Oliver M, Feng D, Cain N, Ledochowitsch P, Millman D, Roll K, Garrett M, Keenan T, Kuan L, Mihalas S, Olsen S, Thompson C, Wakeman W, Waters J, Williams D, Barber C, Berbesque N, Blanchard B, Bowles N, Caldejon SD, Casal L, Cho A, Cross S, Dang C, Dolbeare T, Edwards M, Galbraith J, Gaudreault N, Gilbert TL, Griffin F, Hargrave P, Howard R, Huang L, Jewell S, Keller N, Knoblich U, Larkin JD, Larsen R, Lau C, Lee E, Lee F, Leon A, Li L, Long F, Luviano J, Mace K, Nguyen T, Perkins J, Robertson M, Seid S, Shea-Brown E, Shi J, Sjoquist N, Slaughterbeck C, Sullivan D, Valenza R, White C, Williford A, Witten DM, Zhuang J, Zeng H, Farrell C, Ng L, Bernard A, Phillips JW, Reid RC, Koch C (2020) A large-scale standardized physiological survey reveals functional organization of the mouse visual cortex. Nat Neurosci 23:138–151.F

Dipoppa M, Ranson A, Krumin M, Pachitariu M, Carandini M, Harris KD (2018) Vision and Locomotion Shape the Interactions between Neuron Types in Mouse Visual Cortex. Neuron 98:602-615.e8.

Douglas RJ, Martin KAC (2004) Neuronal circuits of the neocortex. Neuroscience 27:419–451.

Feldmeyer D, Lübke J, Sakmann B (2006) Efficacy and connectivity of intracolumnar pairs of layer 2/3 pyramidal cells in the barrel cortex of juvenile rats. J Physiology 575:583–602.

Fu Y, Tucciarone JM, Espinosa JS, Sheng N, Darcy DP, Nicoll RA, Huang ZJ, Stryker MP (2014) A Cortical Circuit for Gain Control by Behavioral State. Cell 156:1139–1152.

Galarreta M, Hestrin S (1999) A network of fast-spiking cells in the neocortex connected by electrical synapses. Nature 402:72–75.

Galarreta M, Hestrin S (2002) Electrical and chemical synapses among parvalbumin fast-spiking GABAergic interneurons in adult mouse neocortex. Proc National Acad Sci 99:12438–12443.

Gibson JR, Beierlein M, Connors BW (1999) Two networks of electrically coupled inhibitory neurons in neocortex. Nature 402:75–79.

Gonchar Y, Wang Q, Burkhalter AH (2008) Multiple distinct subtypes of GABAergic neurons in mouse visual cortex identified by triple immunostaining. Front Neuroanat 2:3.

Gontier C, Pfister J-P (2020) Identifiability of a Binomial Synapse. Front Comput Neurosc 14:558477.

Gouwens NW, Sorensen SA, Baftizadeh F, Budzillo A, Lee BR, Jarsky T, Alfiler L, Baker K, Barkan E, Berry K, Bertagnolli D, Bickley K, Bomben J, Braun T, Brouner K, Casper T, Crichton K, Daigle TL, Dalley R, Frates RA de, Dee N, Desta T, Lee SD, Dotson N, Egdorf T, Ellingwood L, Enstrom R, Esposito L, Farrell C, Feng D, Fong O, Gala R, Gamlin C, Gary A, Glandon A, Goldy J, Gorham M, Graybuck L, Gu H, Hadley K, Hawrylycz MJ, Henry AM, Hill D, Hupp M, Kebede S, Kim TK, Kim L, Kroll M, Lee C, Link KE, Mallory M, Mann R, Maxwell M, McGraw M, McMillen D, Mukora A, Ng L, Ng L, Ngo K, Nicovich PR, Oldre A, Park D, Peng H, Penn O, Pham T, Pom A, Popović Z, Potekhina L, Rajanbabu R, Ransford S, Reid D, Rimorin C, Robertson M, Ronellenfitch K, Ruiz A, Sandman D, Smith K, Sulc J, Sunkin SM, Szafer A, Tieu M, Torkelson A, Trinh J, Tung H, Wakeman W, Ward K, Williams G, Zhou Z, Ting JT, Arkhipov A, Sümbül U, Lein ES, Koch C, Yao Z, Tasic B, Berg J, Murphy GJ, Zeng H (2020) Integrated Morphoelectric and Transcriptomic Classification of Cortical GABAergic Cells. Cell 183:935-953.e19.

Gouwens NW, Sorensen SA, Berg J, Lee C, Jarsky T, Ting J, Sunkin SM, Feng D, Anastassiou CA, Barkan E, Bickley K, Blesie N, Braun T, Brouner K, Budzillo A, Caldejon S, Casper T, Castelli D, Chong P, Crichton K, Cuhaciyan C, Daigle TL, Dalley R, Dee N, Desta T, Ding S-L, Dingman S, Doperalski A, Dotson N, Egdorf T, Fisher M, Frates RA de, Garren E, Garwood M, Gary A, Gaudreault N, Godfrey K, Gorham M, Gu H, Habel C, Hadley K, Harrington J, Harris JA, Henry A, Hill D, Josephsen S, Kebede S, Kim L, Kroll M, Lee B, Lemon T, Link KE, Liu X, Long B, Mann R, McGraw M, Mihalas S, Mukora A, Murphy GJ, Ng L, Ngo K, Nguyen TN, Nicovich PR, Oldre A, Park D, Parry S, Perkins J, Potekhina L, Reid D, Robertson M, Sandman D, Schroedter M, Slaughterbeck C, Soler-Llavina G, Sulc J, Szafer A, Tasic B, Taskin N, Teeter C, Thatra N, Tung H, Wakeman W, Williams G, Young R, Zhou Z, Farrell C, Peng H, Hawrylycz MJ, Lein E, Ng L, Arkhipov A, Bernard A, Phillips JW, Zeng H, Koch C (2019) Classification of electrophysiological and morphological neuron types in the mouse visual cortex. Nat Neurosci 22:1182–1195.

Graybuck LT, Daigle TL, Sedeño-Cortés AE, Walker M, Kalmbach B, Lenz GH, Morin E, Nguyen TN, Garren E, Bendrick JL, Kim TK, Zhou T, Mortrud M, Yao S, Siverts LA, Larsen R, Gore BB, Szelenyi ER, Trader C, Balaram P, Velthoven CTJ van, Chiang M, Mich JK, Dee N, Goldy J, Cetin AH, Smith K, Way SW, Esposito L, Yao Z, Gradinaru V, Sunkin SM, Lein E, Levi BP, Ting JT, Zeng H, Tasic B (2021) Enhancer viruses for combinatorial cell-subclass-specific labeling. Neuron.

Group PIN, Ascoli GA, Alonso-Nanclares L, Anderson SA, Barrionuevo G, Benavides-Piccione R, Burkhalter A, Buzsáki G, Cauli B, Defelipe J, Fairén A, Feldmeyer D, Fishell G, Fregnac Y, Freund TF, Gardner D, Gardner EP, Goldberg JH, Helmstaedter M, Hestrin S, Karube F, Kisvárday ZF, Lambolez B, Lewis DA, Marin O, Markram H, Muñoz A, Packer A, Petersen CCH, Rockland KS, Rossier J, Rudy B, Somogyi P, Staiger JF, Tamas G, Thomson AM, Toledo-Rodriguez M, Wang Y, West DC, Yuste R (2008) Petilla terminology: nomenclature of features of GABAergic interneurons of the cerebral cortex. Nat Rev Neurosci 9:557–568.

Gupta A, Wang Y, Markram H (2000) Organizing Principles for a Diversity of GABAergic Interneurons and Synapses in the Neocortex. Science 287:273–278.

Hage TA, Bosma-Moody A, Baker CA, Kratz MB, Campagnola L, Jarsky T, Zeng H, Murphy GJ (2019) Distribution and strength of interlaminar synaptic connectivity in mouse primary visual cortex revealed by two-photon optogenetic stimulation. Biorxiv:2019.12.13.876128.

Harris JA, Mihalas S, Hirokawa KE, Whitesell JD, Choi H, Bernard A, Bohn P, Caldejon S, Casal L, Cho A, Feiner A, Feng D, Gaudreault N, Gerfen CR, Graddis N, Groblewski PA, Henry AM, Ho A, Howard R, Knox JE, Kuan L, Kuang X, Lecoq J, Lesnar P, Li Y, Luviano J, McConoughey S, Mortrud MT, Naeemi M, Ng L, Oh S, Ouellette B, Shen E, Sorensen SA, Wakeman W, Wang Q, Wang Y, Williford A, Phillips JW, Jones AR, Koch C, Zeng H (2019) Hierarchical organization of cortical and thalamic connectivity. Nature 575:195–202.

Harris KD, Shepherd GMG (2015) The neocortical circuit: themes and variations. Nat Neurosci 18:170–181.

Hennig MH (2013) Theoretical models of synaptic short term plasticity. Front Comput Neurosc 7:45.

Hodge RD, Bakken TE, Miller JA, Smith KA, Barkan ER, Graybuck LT, Close JL, Long B, Johansen N, Penn O, Yao Z, Eggermont J, Höllt T, Levi BP, Shehata SI, Aevermann B, Beller A, Bertagnolli D, Brouner K, Casper T, Cobbs C, Dalley R, Dee N, Ding S-L, Ellenbogen RG, Fong O, Garren E, Goldy J, Gwinn RP, Hirschstein D, Keene CD, Keshk M, Ko AL, Lathia K, Mahfouz A, Maltzer Z, McGraw M, Nguyen TN, Nyhus J, Ojemann JG, Oldre A, Parry S, Reynolds S, Rimorin C, Shapovalova NV, Somasundaram S, Szafer A, Thomsen ER, Tieu M, Quon G, Scheuermann RH, Yuste R, Sunkin SM, Lelieveldt B, Feng D, Ng L, Bernard A, Hawrylycz M, Phillips JW, Tasic B, Zeng H, Jones AR, Koch C, Lein ES (2019) Conserved cell types with divergent features in human versus mouse cortex. Nature 573:61–68.

Hoffmann FZ, Triesch J (2017) Nonrandom network connectivity comes in pairs. Netw Neurosci 1:31–41.

Holmgren C, Harkany T, Svennenfors B, Zilberter Y (2003) Pyramidal cell communication within local networks in layer 2/3 of rat neocortex. J Physiology 551:139–153.

Hu H, Cavendish JZ, Agmon A (2013) Not all that glitters is gold: off-target recombination in the somatostatin-IRES-Cre mouse line labels a subset of fast-spiking interneurons. Front Neural Circuit 7:195.

Hu H, Ma Y, Agmon A (2011) Submillisecond Firing Synchrony between Different Subtypes of Cortical Interneurons Connected Chemically But Not Electrically. J Neurosci 31:3351–3361.

Jiang X, Shen S, Cadwell CR, Berens P, Sinz F, Ecker AS, Patel S, Tolias AS (2015) Principles of connectivity among morphologically defined cell types in adult neocortex. Science 350:aac9462.

Jones HC, Keep RF (1988) Brain fluid calcium concentration and response to acute hypercalcaemia during development in the rat. J Physiology 402:579–593.

Jouhanneau J-S, Kremkow J, Dorrn AL, Poulet JFA (2015) In Vivo Monosynaptic Excitatory Transmission between Layer 2 Cortical Pyramidal Neurons. Cell Reports 13:2098–2106.

Kalmbach BE, Buchin A, Long B, Close J, Nandi A, Miller JA, Bakken TE, Hodge RD, Chong P, Frates R de, Dai K, Maltzer Z, Nicovich PR, Keene CD, Silbergeld DL, Gwinn RP, Cobbs C, Ko AL, Ojemann JG, Koch C, Anastassiou CA, Lein ES, Ting JT (2018) h-Channels Contribute to Divergent Intrinsic Membrane Properties of Supragranular Pyramidal Neurons in Human versus Mouse Cerebral Cortex. Neuron 100:1194-1208.e5.

Kapfer C, Glickfeld LL, Atallah BV, Scanziani M (2007) Supralinear increase of recurrent inhibition during sparse activity in the somatosensory cortex. Nat Neurosci 10:743–753.

Karnani MM, Jackson J, Ayzenshtat I, Tucciarone J, Manoocheri K, Snider WG, Yuste R (2016) Cooperative Subnetworks of Molecularly Similar Interneurons in Mouse Neocortex. Neuron 90:86–100.

Keller AJ, Dipoppa M, Roth MM, Caudill MS, Ingrosso A, Miller KD, Scanziani M (2020) A Disinhibitory Circuit for Contextual Modulation in Primary Visual Cortex. Neuron 108:1181-1193.e8.

Kim EJ, Juavinett AL, Kyubwa EM, Jacobs MW, Callaway EM (2015) Three Types of Cortical Layer 5 Neurons That Differ in Brain-wide Connectivity and Function. Neuron 88:1253–1267.

Ko H, Hofer SB, Pichler B, Buchanan KA, Sjöström JP, Mrsic-Flogel TD (2011) Functional specificity of local synaptic connections in neocortical networks. Nature 473:87–91.

Larsen RS, Sjöström PJ (2015) Synapse-type-specific plasticity in local circuits. Curr Opin Neurobiol 35:127–135.

Lee BR, Budzillo A, Hadley K, Miller JA, Jarsky T, Baker K, Hill D, Kim L, Mann R, Ng L, Oldre A, Rajanbabu R, Trinh J, Braun T, Dalley R, Gouwens NW, Kalmbach BE, Kim TK, Smith K, Soler-Llavina GJ, Sorensen SA, Tasic B, Ting JT, Lein ES, Zeng H, Murphy GJ, Berg J (2020)

Scaled, high fidelity electrophysiological, morphological, and transcriptomic cell characterization. Biorxiv:2020.11.04.369082.

Lefort S, Petersen CCH (2017) Layer-Dependent Short-Term Synaptic Plasticity Between Excitatory Neurons in the C2 Barrel Column of Mouse Primary Somatosensory Cortex. Cereb Cortex 27:3869–3878.

Levy RB, Reyes AD (2012) Spatial Profile of Excitatory and Inhibitory Synaptic Connectivity in Mouse Primary Auditory Cortex. J Neurosci 32:5609–5619.

Llera-Montero M, Sacramento J, Costa RP (2019) Computational roles of plastic probabilistic synapses. Curr Opin Neurobiol 54:90–97.

Lourenço J, Bacci A (2017) Human-Specific Cortical Synaptic Connections and Their Plasticity: Is That What Makes Us Human? Plos Biol 15:e2001378.

Lur G, Vinck MA, Tang L, Cardin JA, Higley MJ (2016) Projection-Specific Visual Feature Encoding by Layer 5 Cortical Subnetworks. Cell Reports 14:2538–2545.

Ma Y, Hu H, Agmon A (2012) Short-Term Plasticity of Unitary Inhibitory-to-Inhibitory Synapses Depends on the Presynaptic Interneuron Subtype. J Neurosci 32:983–988.

Markram H (1997) A network of tufted layer 5 pyramidal neurons. Cereb Cortex 7:523--533.

Markram H, Lübke J, Frotscher M, Roth A, Sakmann B (1997) Physiology and anatomy of synaptic connections between thick tufted pyramidal neurones in the developing rat neocortex. J Physiology 500:409–440.

Markram H, Muller E, Ramaswamy S, Reimann MW, Abdellah M, Sanchez CA, Ailamaki A, Alonso-Nanclares L, Antille N, Arsever S, Kahou GAA, Berger TK, Bilgili A, Buncic N, Chalimourda A, Chindemi G, Courcol J-D, Delalondre F, Delattre V, Druckmann S, Dumusc R, Dynes J, Eilemann S, Gal E, Gevaert ME, Ghobril J-P, Gidon A, Graham JW, Gupta A, Haenel V, Hay E, Heinis T, Hernando JB, Hines M, Kanari L, Keller D, Kenyon J, Khazen G, Kim Y, King JG, Kisvarday Z, Kumbhar P, Lasserre S, Le Bé J-V, Magalhães BRC, Merchán-Pérez A, Meystre J, Morrice BR, Muller J, Muñoz-Céspedes A, Muralidhar S, Muthurasa K, Nachbaur D, Newton TH, Nolte M, Ovcharenko A, Palacios J, Pastor L, Perin R, Ranjan R, Riachi I, Rodríguez J-R, Riquelme JL, Rössert C, Sfyrakis K, Shi Y, Shillcock JC, Silberberg G, Silva R, Tauheed F, Telefont M, Toledo-Rodriguez M, Tränkler T, Van Geit W, Díaz JV, Walker R, Wang Y, Zaninetta SM, DeFelipe J, Hill SL, Segev I, Schürmann F (2015) Reconstruction and Simulation of Neocortical Microcircuitry. Cell 163:456–492.

Markram H, Toledo-Rodriguez M, Wang Y, Gupta A, Silberberg G, Wu C (2004) Interneurons of the neocortical inhibitory system. Nat Rev Neurosci 5:793–807.

McInnes L, Healy J, Melville J (2018) UMAP: Uniform Manifold Approximation and Projection for Dimension Reduction. Arxiv.

Molnár G, Oláh S, Komlósi G, Füle M, Szabadics J, Varga C, Barzó P, Tamás G (2008) Complex Events Initiated by Individual Spikes in the Human Cerebral Cortex. Plos Biol 6:e222.

Molnár G, Rózsa M, Baka J, Holderith N, Barzó P, Nusser Z, Tamás G (2016) Human pyramidal to interneuron synapses are mediated by multi-vesicular release and multiple docked vesicles. Elife 5:e18167.

Morishima M, Kobayashi K, Kato S, Kobayashi K, Kawaguchi Y (2017) Segregated Excitatory–Inhibitory Recurrent Subnetworks in Layer 5 of the Rat Frontal Cortex. Cereb Cortex 27:5846–5857.

Naka A, Veit J, Shababo B, Chance RK, Risso D, Stafford D, Snyder B, Egladyous A, Chu D, Sridharan S, Mossing DP, Paninski L, Ngai J, Adesnik H (2019) Complementary networks of cortical somatostatin interneurons enforce layer specific control. Elife 8:e43696.

Nanou E, Catterall WA (2018) Calcium Channels, Synaptic Plasticity, and Neuropsychiatric Disease. Neuron 98:466–481.

Niell CM, Stryker MP (2008) Highly selective receptive fields in mouse visual cortex. J Neurosci Official J Soc Neurosci 28:7520–7536.

Obermayer J, Heistek TS, Kerkhofs A, Goriounova NA, Kroon T, Baayen JC, Idema S, Testa-Silva G, Couey JJ, Mansvelder HD (2018) Lateral inhibition by Martinotti interneurons is facilitated by cholinergic inputs in human and mouse neocortex. Nat Commun 9:4101.

Packer AM, Yuste R (2011) Dense, Unspecific Connectivity of Neocortical Parvalbumin-Positive Interneurons: A Canonical Microcircuit for Inhibition? J Neurosci 31:13260–13271.

Pala A, Petersen CCH (2015) In Vivo Measurement of Cell-Type-Specific Synaptic Connectivity and Synaptic Transmission in Layer 2/3 Mouse Barrel Cortex. Neuron 85:68–75.

Perin R, Berger TK, Markram H (2011) A synaptic organizing principle for cortical neuronal groups. Proc National Acad Sci 108:5419–5424.

Pfeffer CK, Xue M, He M, Huang ZJ, Scanziani M (2013) Inhibition of inhibition in visual cortex: the logic of connections between molecularly distinct interneurons. Nature Neuroscience 16:1068–1076.

Reyes A, Lujan R, Rozov A, Burnashev N, Somogyi P, Sakmann B (1998) Target-cell-specific facilitation and depression in neocortical circuits. Nat Neurosci 1:279–285.

Reyes A, Sakmann B (1999) Developmental Switch in the Short-Term Modification of Unitary EPSPs Evoked in Layer 2/3 and Layer 5 Pyramidal Neurons of Rat Neocortex. J Neurosci 19:3827–3835.

Scala F, Kobak D, Shan S, Bernaerts Y, Laturnus S, Cadwell CR, Hartmanis L, Froudarakis E, Castro JR, Tan ZH, Papadopoulos S, Patel SS, Sandberg R, Berens P, Jiang X, Tolias AS (2019) Layer 4 of mouse neocortex differs in cell types and circuit organization between sensory areas. Nat Commun 10:4174.

Scala F, Kobak D, Bernabucci M, Bernaerts Y, Cadwell CR, Castro JR, Hartmanis L, Jiang X, Laturnus S, Miranda E, Mulherkar S, Tan ZH, Yao Z, Zeng H, Sandberg R, Berens P, Tolias AS (2020) Phenotypic variation of transcriptomic cell types in mouse motor cortex. Nature:1–7.

Seeman SC, Campagnola L, Davoudian PA, Hoggarth A, Hage TA, Bosma-Moody A, Baker CA, Lee JH, Mihalas S, Teeter C, Ko AL, Ojemann JG, Gwinn RP, Silbergeld DL, Cobbs C, Phillips J, Lein E, Murphy G, Koch C, Zeng H, Jarsky T (2018) Sparse recurrent excitatory connectivity in the microcircuit of the adult mouse and human cortex. Elife 7:e37349.

Silberberg G, Markram H (2007) Disynaptic Inhibition between Neocortical Pyramidal Cells Mediated by Martinotti Cells. Neuron 53:735–746.

Song S, Sjöström PJ, Reigl M, Nelson S, Chklovskii DB (2005) Highly Nonrandom Features of Synaptic Connectivity in Local Cortical Circuits. Plos Biol 3:e68.

Stepanyants A, Martinez LM, Ferecskó AS, Kisvárday ZF (2009) The fractions of short- and long-range connections in the visual cortex. Proc National Acad Sci 106:3555–3560.

Szegedi V, Molnár G, Paizs M, Csakvari E, Barzó P, Tamás G, Lamsa K (2017) High-Precision Fast-Spiking Basket Cell Discharges during Complex Events in the Human Neocortex. Eneuro 4:ENEURO.0260-17.2017.

Tasic B, Menon V, Nguyen TN, Kim TK, Jarsky T, Yao Z, Levi B, Gray LT, Sorensen SA, Dolbeare T, Bertagnolli D, Goldy J, Shapovalova N, Parry S, Lee C, Smith K, Bernard A, Madisen L, Sunkin SM, Hawrylycz M, Koch C, Zeng H (2016) Adult mouse cortical cell taxonomy revealed by single cell transcriptomics. Nat Neurosci 19:335–346.

Tasic B, Yao Z, Graybuck LT, Smith KA, Nguyen TN, Bertagnolli D, Goldy J, Garren E, Economo MN, Viswanathan S, Penn O, Bakken T, Menon V, Miller J, Fong O, Hirokawa KE, Lathia K, Rimorin C, Tieu M, Larsen R, Casper T, Barkan E, Kroll M, Parry S, Shapovalova NV, Hirschstein D, Pendergraft J, Sullivan HA, Kim TK, Szafer A, Dee N, Groblewski P, Wickersham I, Cetin A, Harris JA, Levi BP, Sunkin SM, Madisen L, Daigle TL, Looger L, Bernard A, Phillips J, Lein E, Hawrylycz M, Svoboda K, Jones AR, Koch C, Zeng H (2018) Shared and distinct transcriptomic cell types across neocortical areas. Nature 563:72–78.

Testa-Silva G, Verhoog MB, Linaro D, Kock CP de, Baayen JC, Meredith RM, Zeeuw CI, Giugliano M, Mansvelder HD (2014) High Bandwidth Synaptic Communication and Frequency Tracking in Human Neocortex. Plos Biol 12:e1002007.

Thomson AM (1997) Activity-dependent properties of synaptic transmission at two classes of connections made by rat neocortical pyramidal axonsin vitro. J Physiology 502:131–147.

Thomson AM, Bannister AP (2003) Interlaminar Connections in the Neocortex. Cereb Cortex 13:5–14.

Thomson AM, West DC, Wang Y, Bannister AP (2002) Synaptic Connections and Small Circuits Involving Excitatory and Inhibitory Neurons in Layers 2–5 of Adult Rat and Cat Neocortex: Triple Intracellular Recordings and Biocytin Labelling In Vitro. Cereb Cortex 12:936–953.

Tremblay R, Lee S, Rudy B (2016) GABAergic Interneurons in the Neocortex: From Cellular Properties to Circuits. Neuron 91:260–292.

Turner NL, Macrina T, Bae JA, Yang R, Wilson AM, Schneider-Mizell C, Lee K, Lu R, Wu J, Bodor AL, Bleckert AA, Brittain D, Froudarakis E, Dorkenwald S, Collman F, Kemnitz N, Ih D, Silversmith WM, Zung J, Zlateski A, Tartavull I, Yu S, Popovych S, Mu S, Wong W, Jordan CS, Castro M, Buchanan J, Bumbarger DJ, Takeno M, Torres R, Mahalingam G, Elabbady L, Li Y, Cobos E, Zhou P, Suckow S, Becker L, Paninski L, Polleux F, Reimer J, Tolias AS, Reid RC, Costa NM da, Seung HS (2020) Multiscale and multimodal reconstruction of cortical structure and function. Biorxiv:2020.10.14.338681.

Vélez-Fort M, Rousseau CV, Niedworok CJ, Wickersham IR, Rancz EA, Brown APY, Strom M, Margrie TW (2014) The Stimulus Selectivity and Connectivity of Layer Six Principal Cells Reveals Cortical Microcircuits Underlying Visual Processing. Neuron 83:1431–1443.

Walker F, Möck M, Feyerabend M, Guy J, Wagener RJ, Schubert D, Staiger JF, Witte M (2016) Parvalbumin- and vasoactive intestinal polypeptide-expressing neocortical interneurons impose differential inhibition on Martinotti cells. Nat Commun 7:13664.

Yang D, Qi G, Feldmeyer D (2021) Layer 6A pyramidal cells subtypes form synaptic microcircuits with distinct functional and structural properties. Biorxiv:2021.01.13.426506.

Yoshimura Y, Dantzker JL, Callaway EM (2005) Excitatory cortical neurons form fine-scale functional networks. Nature 433:868–873.

Figure 1 - Synaptic Physiology Pipeline: A. Throughput of the pipeline. Each large circle shows statistics for data collected from mouse primary visual cortex (left) and human (right). The center plot shows the age distribution of each organism. Below, the outer small circles show the distributions of recorded cells; in mouse (left) this is denoted by transgenic subclass of excitatory and inhibitory cells and in human this is denoted by cortical region. The inner circles show the number of experiments in which two cells were recorded from, three cells, etc., up to a maximum of eight cells being simultaneously recorded. **B**. *i.Multipatch Experiment*: The top panel shows the set of stimuli used in a multipatch experiment. They consisted of pulse trains at various set frequencies with 8 initial pulses, a delay of 250 ms, and then four more pulses. In the 50 Hz stimulus various delays were interposed between the first 8 and second 4 pulses. The mixed frequency stimulus started with 8 pulses at 30 Hz and the proceeded with 30 pulses at set, but variable instantaneous frequencies ranging from 10-200 Hz. The fluorescent images to the right show cells recorded in an example experiment with the connectivity diagram overlaid. Overall in this experiment, cell 1, an unlabeled spiny cell, received convergent input from both Sst cells (3, 5) and Pvalb cells (7, 8), while cell 8, a Pvalb cell, showed divergent outputs to three different subclasses. Thus, ten synaptic connections were identified in this experiment, characterizing seven of the possible nine interaction types. Below is an example of the electrophysiology recording of cells 1, 3, and 5 during the 50 Hz stimulus highlighted by the orange box. The stimulus and presynaptic spikes were recorded from cell 1 which was presynaptic to cell 3. The response of cell 3 to individual presentations (trials) of the stimulus are shown in grey and the average in blue. A similar example for an inhibitory connection also found in this experiment from cell 5 to cell 1 is shown in green. *ii. Synapse Processing*: A synapse was identified by aligning every presynaptic spike and overlaying the postsynaptic response (black traces) in both voltage clamp (top row) and current clamp (bottom row) with the average in color (colors correspond to the same synapses in the *Multipatch Experiment* panel). Excitatory connections were evaluated at a holding potential between -80 and -61 (left column) mV while inhibitory connections were evaluated at a holding potential between -60 and -45 mV (right column). *iii Synapse Analysis*: When a connection was identified the

average (colored) were fit (black) and metrics such as amplitude and rise time (orange lines) were extracted from the fit output. These fit parameters were used to guide fitting of individual pulses so that we could quantify changes in synapse amplitude over the course of a train. Short term plasticity (STP) was quantified as the amplitude of the first pulse subtracted from the average of pulses 6-8, normalized by the 90th percentile amplitude for the connection overall. Recovery was similarly quantified as the average of pulses 1-4 subtracted from pulses 9-12, normalized by the 90th percentile. **C**. *i Intrinsic Ephys*: Long pulse steps were applied to quantify intrinsic cell properties, and gap junctions (Figure S4). Hyperpolarizing steps (left) delivered to an example cell probed the subthreshold I-V relationship to quantify metrics such as input resistance while depolarizing suprathreshold sweeps (right) were used to measure spiking and firing rate properties. *ii Morphology*: Cells were filled with biocytin during recording and stained. 20x images of the full slice stained with DAPI allowed identification of the cortical layers and 63x z-stack images were used to assess morphological properties such as dendritic type and axon length. Images and cells here are the same as in (B).

Figure 2 - Connectivity: A. Gaussian fit of connection probability as a function of lateral intersomatic distance. All mouse cells were divided into two main classes, excitatory and inhibitory, and pairs classified into the four combinations of those two classes. *Top row*: Connection probability as a function of intersomatic distance was fit with a Gaussian (red line) and output parameters pmax and sigma describe the max connection probability and width of the Gaussian. Adjusted connection probability as a function of intersomatic distance adjusted for presynaptic axon length, depth of the pair from the slice surface, and detection power of connections using a unified model (dashed red line) or via filtering of the data (dotted red line) yield similar results (see Results/Methods). Grey line and area are 40 μm binned average connection probability and 95% confidence interval. Raster below shows distance distribution of connections probed (bottom) and found (top). *Bottom row*: Normalized rate of reciprocal connections. Probed pairs are unordered and number of reciprocal connections counted normalized to expected value of connection probability squared for a randomly connected network (solid red line). **B**. Connection probability matrix for mouse. Connection probability is estimated using the max of a Gaussian fit with a sigma of 100 μm and accounting for all corrections as determined from A (dashed red line, "model"). The shading of each element indicates the 95% CI of the data with higher contrast indicating smaller CI and lower contrast (toward grey) indicating larger CI. The number of connections found out of number of connections probed are printed in each element. Excitatory cell class in layers 4-6 were distinguished by transgenic reporter while layer 2/3 excitatory cells are defined as morphologically spiny. Three inhibitory classes Pvalb, Sst, and Vip are defined by transgenic reporter. **C-E**. Gaussian fit of connection probability vs intersomatic distance (with CI at p_{max}, shaded region) for two contrasting elements with connections found and connections probed raster below. Cross symbol denotes p_{max} with all corrections.

Figure 3 - Synaptic strength and kinetics: A. (left to right) PSP latency matrix; excitatory and inhibitory minimum average latency, median average latency, and maximum average latency representative traces (light to dark colors); latency histograms for the major connection classes (E>E, E>I, I>I, I>E); latency summary plots for a subset of matrix elements. **B.** (left to right) PSP rise time matrix; representative traces for excitatory and inhibitory minimum average rise time, median average rise time, and maximum average rise time (light to dark colors); rise time histograms for the major connection classes; rise time summary plots for a subset of matrix elements. **C.** (left to right) PSP decay tau matrix; representative traces for excitatory and inhibitory minimum average decay tau, median average decay tau, and maximum average decay tau (light to dark colors); decay tau histograms for the major connection classes; decay tau summary plots for a subset of matrix elements. **D.** (left to right) PSP amplitude matrix; representative traces for excitatory and inhibitory minimum average amplitude , median average amplitude , and maximum average amplitude (light to dark colors); amplitude histograms for the major connection classes; amplitude summary plots for a subset of matrix elements. **E.** (left to right) PSP 90th percentile amplitude ; representative traces for excitatory and inhibitory minimum average 90th percentile amplitude, median average 90th percentile amplitude, and maximum average 90th percentile amplitude (light to dark colors); 90th percentile amplitude histograms for the major connection classes; 90th percentile amplitude summary plots for a subset of matrix elements. In all matrices, inhibitory cells are merged across layers.

Figure 4 - Synaptic dynamics: A. Representative depressing, facilitating, and pseudo-linear excitatory synapses (top to bottom); grey dots: individual PSP amplitudes; black lines: average amplitude per pulse in 50 Hz train. **B.** Short-term plasticity matrix. **C.** Recovery (at 250ms) matrix. **D.** Resting state variance (adjusted coefficient of variation) matrix. **E.** Summary plots for paired pulse ratio, STP induction ratio (avg 1st pulse amp : avg of 5th-8th pulse amp) normalized by the 90th percentile, Resting state variance, induced state variance (top to bottom). Each dot corresponds to the average response from one synapse. **F.** Train induced STP (top) at four different frequencies (10, 20, 50, 100 Hz) for each of the elements in E (colors maintained). Each dot is the grand average of all synapses in the element. For L5 ET → L5 ET the blue shading highlights the 95% confidence interval as an example. Lower plot shows recovery from STP at six different delays (125, 250, 500, 1000, 2000, 4000 ms) in a similar manner to the plot above.

Figure 5- Human: A. Connection probability of human synapses. Inhibitory cells are identified by morphology as aspiny or sparsely spiny cells, grouped across layer. **B.** Kinetics, strength, and dynamics of human synapses. All matrices are organized by layer for excitatory cells, with inhibitory cells grouped across layer. Each element is colorized by the median (text in each element) according to the colormap with the saturation scaled to the standard error. Two or more pairs were required to fill in an element. Latency, Rise tau, and Resting state amplitude are quantified from fits of the average PSP response which have passed QC and visual inspection of the fit. **C.**Train induced STP (top) across frequencies for a subset of connection types. Each dot is the median of all synapses in the element, with shading for the 95% CI (bootstrapped) shown for a single example connection type. Lower plot shows recovery from STP at different delays. **D.** Polysynaptic connections from L2/3 pyramidal cells inferred by response latency > 3 ms are plotted with PSP amplitude. All polysynapses from L2 pyramidal cells (left) are inhibitory, while those from L3 pyramidal cells (right) are mixed. Example average inhibitory polysynaptic response shown as inset and highlighted as a larger X symbol. **E.** UMAP projection of intrinsic electrophysiology features from all human cells, colored by cell subclass (left), showing the distinctiveness of inhibitory and L4-type excitatory cells; and by depth (right), showing the structured variability of intrinsic properties with depth in L2/3. **F.** Within L2/3, variation in intrinsic properties is structured and strongly correlated with depth in human but not mouse. Example traces are shown for a superficial and deep human cell (colors as in D), alongside scatter plots of electrophysiology features vs. depth for both species (right). Top shows a phase plane representation of the first spike in a depolarizing step response, quantified as AP upstroke/downstroke ratio. Bottom shows the sag in response to hyperpolarization, quantified as sag ratio. Regression lines shown with bootstrapped 95% CI. **G.** STP of L2/3 excitatory synapses is structured by depth in human and not mouse. Top shows example PSP responses to spike trains for cells from E. The larger response to the first spike is quantified by paired pulse STP, plotted below in relation to postsynaptic depth (left) and AP up/down ratio (right).

Figure 6: Relationships among synapse properties and cell types revealed by dimensionality reduction. **A.** All synapses colored by postsynaptic E/I cell class. The UMAP output generates two clusters: excitatory (left) and inhibitory (right). **B.** Human and mouse synapses colored by postsynaptic subclass. **C.** Mouse synapses colored by presynaptic subclass. **D.** Four synapse properties represented in reduced space, showing 90th percentile PSP amplitude (red=excitatory, blue=inhibitory); STP induced by 50 Hz trains (red=facilitating, blue=depressing), resting state aCV during 50 Hz trains (purple=low variability, yellow=high variability), and the binomial CV derived from model parameters (release probability * number of release sites; purple=high CV, yellow=low CV).

Figure 7: The cortical intralayer circuit differs across layer and with activity. **A)** Some commonly described elements of the intralaminar cortical circuit. Pvalb cells strongly inhibit pyramidal and other Pvalb cells, Sst cells provide broad inhibition, and Vip cells inhibit Sst cells to form a disinhibitory feedback pathway. **B)** Layer 2/3 circuit diagram showing connections between major subclasses. The width of connecting lines roughly represents connection probability and PSP amplitude. Connections that are prominent in mouse L2/3 compared to deeper layers are highlighted in orange, whereas green dashed lines indicate connections that are weak or absent in L2/3. **C)** Circuit diagram of connections found in mouse layers 5 and 6. For simplicity, connections between IT pyramidal and inhibitory cells are omitted. **D)** Two complementary circuits that activate at different times. Red connections are facilitating and will be stronger during sustained activity. Blue connections are depressing and are strongest during quiescent periods.

Figure Supplement 1 Connectivity Adjustments. A. From left: Distribution of presynaptic axon length measured from biocytin fills. If the axon was measured to at least 200 µm the axon it was not measured further except in rare occasions; Connection probability as a function of axon length and 95% confidence interval (grey line/shading) with thresholded fit in the colored line (dotted line denotes fit p_{max}) for all pairs in mouse. Raster below plot shows presynaptic axon lengths of pairs that were probed for connectivity along with pairs that were connected; Same plots as for All Pairs but separated by whether the presynaptic cell was excitatory or inhibitory. **B**. Same plots as A for the average depth of the cell pair from the slice surface. In this case the relationship between connection probability and average pair depth is fit with an error function. **C.** Same plots as A for detection power. Detection power combines the signal to noise ratio of the postsynaptic cell with the number of spikes elicited by the presynaptic cell to probe the connection (see Methods). Detection power as a function of connection probability was also fit with an error function. **D**. Gaussian fit of connection probability as a function of lateral intersomatic distance (similar to Fig 2A) for All, Excitatory, and Inhibitory pairs. The dark red line is the fit only taking into consideration intersomatic distance while each colored line multiplies the effect of the metric fits applied in A-C. The pink curve denotes adjustment for all metrics in A-C. Adjusted p_{max} with a fixed σ of 100 µm are enumerated in the table on the left for each metric and group of pairs. **E.** Adjustment factor of each metric applied to each element in the matrix in Fig 2B.

A

B

Connection probability in sphere intersection model

Figure Supplement 2 Connectivity Matrix. A. The connectivity matrix in Fig 2B with additional details highlighted in the expanded element below the matrix. The center number is the fully adjusted (see Figure S1) p_{max}. The upper and lower numbers on the left of each element are the number of connections found (upper) and number of connections probed (lower). The numbers on the right of each element are the upper and lower 95% confidence interval, respectively. **B**. Probability of connection versus intersomatic distance in a simplified model that assumes a constant density of synapses inside the volume intersection of two spheres. Changing the density of synapses can result in profiles that look qualitatively like an exponential decay, a Gaussian, or a sigmoid.

Figure Supplement 3 Intrinsic uMap and morphologies of Sst recurrent connections. A.UMAP projection of electrophysiology feature space of all mouse inhibitory interneurons. Sst-cre/flp to Sst-cre/flp chemical synaptic connections (colored lines) are overlaid. Umap is color coded by the transgenic cell class. 9 out of 16 Sst to Sst connections had pre- and post-synaptic neurons that mapped with other Sst neurons and apart from Pvalb-cre/flp neurons (note the short green line near the top of the blue line). Only Sst to Sst connections that fall within the Sst island are indicated with a line. Sst-cre/flp cells that are part of a connection are indicated by colors other than blue (Vip-cre/flp), green (Pvalb-cre/flp), and yellow (Sst-cre/flp). **B**. Biocytin image of connected Sst-cre to Sst-cre connected cells with Sst-like morphologies (left). Insets show sparsely spiny dendrites for pre and post synaptic cell. Biocytin image with overlaid reconstructions (right). Insets show sites of close apposition between axon and dendrite of connected neurons. Reconstructions shown in C, show independently (bottom). **C**. UMAP projection of IVSCC patchseq, FACS, and mSeq feature space of mouse inhibitory interneurons. Large dots in the Sst space highlight those from mSeq with Cells from Sst→Sst connections further highlighted in red.

Figure Supplement 4 Electrical Synapses. A. Electrical connectivity matrix among inhibitory cells in each layer from mouse. **B**. (upper) Electrical connection probability as a function of lateral intersomatic distance with 95% confidence interval (grey line/shading) fit with a Gaussian (solid red line). Dotted red line shows Guassian fit of chemical I → I connections (normalized to electrical p_{max}) for reference and to highlight shorter σ of electrical connections. (lower) Cumulative histogram of σ ratio comparing chemical and electrical connections. 1000 experiments were simulated in which the true σ for electrical and chemical connections was set to six evenly distributed values between 65 and 140 µm (light to dark red). We then measured the ratio of the Gaussian fit σ between chemical and electrical connections from those 1000 experiments which are plotted here as a cumulative histogram (a value of 1 indicates that the Gaussian profile of electrical and chemical connections has the same σ). The dotted vertical line denotes the measured σ ratio between chemical and electrical synapses and sits beyond the 99th percentile for all simulations. **C**. Coupling coefficient (top) and junctional conductance (bottom) of recurrent I → I electrical connections. Left plots show a scatter where each dot is the value for a single unidirectional electrical connection and bars denote the median. Right plots show coupling coefficient and junctional conductance of each electrical connection vs it's reciprocal connection (dotted line is unity line). **D**. Junctional conductance as a function of input resistance of the postsynaptic cell.

Figure Supplement 5 Strength and Kinetics. **A**. The distribution of inhibitory cells according to layer. **B**. Latency of inhibitory → inhibitory connections for all of the combinations among Pvalb, Sst, and Vip. For each connection element they are grouped by layer to estimate the variance of I → I latency across layer. The table below shows the p-value from a Kruskal-Wallis (KW) test which suggests that within layer I → I latency does not vary across layers. **C-F**. The same as B for PSP rise time (**C**), PSP decay tau (**D**), PSP resting state amplitude (**E**), and PSP 90th percentile amplitude (**F**). **G**. Average strength or kinetic measurement of each element in the 8 x 8 matrix from Figure 3 with standard deviation sorted from lowest to highest (blue dots, left axis) and number of pairs within each element shown in the grey bars (right axis).

Figure Supplement 6 Dynamics. A. Paired pulse ratio (stp_initial_50hz) of inhibitory → inhibitory connections for all of the combinations among Pvalb, Sst, and Vip. For each connection element they are grouped by layer to estimate the variance of I → I PPR across layer. The table below shows the p-value from a Kruskal-Wallis (KW) test which suggests that within layer I → I PPR does not vary across layers. **B-E**. The same as A for STP induction (**B**), STP recovery (**C**), resting state variability (**D**), and STP induced variability (**E**). **F**. Average dynamics measurement of each element in the 8 x 8 matrix from Figure 4 with standard deviation sorted from lowest to highest (blue dots, left axis) and number of pairs within each element shown in the grey bars (right axis)

Figure Supplement 7 Human Data: A. Gaussian fit of connection probability for all human synapses. Connection probability as a function of lateral intersomatic distance was fit with a Gaussian (red line). Output parameters pmax and size describe the max connection probability and sigma of the Gaussian. Grey line and area are 40um binned average connection probability and 95% confidence interval. Raster below shows distance distribution of connections probed (bottom) and found (top). **B.** Additional STP and variability metrics of human synapse. Matrices are organized by layer for excitatory cells, with inhibitory cells grouped across layer. Each element is colorized by the grand average (text in each element) according to the colormap with the saturation scaled to the standard error. Two or more pairs were required to fill in an element. **C.** Summary plots for a range of synapse strength, timing, STP, and variability measurements. Each dot corresponds to the average response from one synapse. Responses are shown for a subset of matrix elements with sufficient sampling. **D-E.** Additional correlates of STP variability in synapses from L2/3 pyramidal cells. D shows dependence on postsynaptic cell class/subclass in human (left) and mouse (right). E shows decreased correlation with depth (left) and AP upstroke/downstroke ratio (right) when indexed to the postsynaptic cell.

Figure Supplement 8 External Calcium Concentration. A. Connectivity matrices of elements in mouse which were probed with both 2 mM external calcium and 1.3 mM external calcium (top row). The bottom left matrix is a difference of the 2 mM matrix from the 1.3 mM matrix. Red elements are those that showed higher connectivity in 1.3 mM calcium and blue elements those that showed higher connectivity in 2 mM calcium. The bottom right matrix shows uncorrected p-values from a Fisher-exact test for each element. **B.** Violin plot of measured PSP amplitude (resting state and 90th percentile) and induced STP or pairs for each element in 2 mM external calcium (blue) and 1.3 mM external calcium (orange). Measurements from individual pairs denoted by black lines within violin. Pairs/measurements are from different experiments, Kolmogorov-Smirnov p-values are shown in the table to the left for each element.

Figure Supplement 9 Internal EGTA Concentration. A. Connectivity matrices of elements in human which were probed with both 0.3 mM internal EGTA and 0 mM internal EGTA (top row). The bottom left matrix is a difference of the 0.3 mM matrix from the 0 mM matrix. Red elements are those that showed higher connectivity in 0 mM calcium and blue elements those that showed higher connectivity in 0.3 mM calcium. The bottom right matrix shows uncorrected p-values from a Fisher-exact test for each element. **B.** Violin plot of measured PSP amplitude (resting state and 90th percentile) and induced STP or pairs for each element in 0.3 mM internal EGTA (blue) and 0 mM internal EGTA (orange). Measurements from individual pairs denoted by black lines within violin. Pairs/measurements are from different experiments, Kolmogorov-Smirnov p-values are shown in the table to the left for each element.

A

B

Figure Supplement 10 Pair Analysis Tool. A. Pair analysis tool used to analyze synapses. Postsynaptic responses and presynaptic spikes (white traces) from each pair were divided by clamp mode (voltage clamp on the left, current clamp on the right) and then again by baseline potential (depolarized potentials on the top, hyperpolarized potentials on the bottom). Responses from each quadrant were averaged (blue trace). If a synapse was identified the user would select the type (excitatory or inhibitory) from the menu on the left and move the yellow line in any quadrant to the onset of the response (all lines are linked). The user would then select "Fit Response". **B.** The fitting algorithm produces a fit of the PSC/P and plots in red (QC fail, NRMSE too high) or green (QC pass). Fit parameters for each quadrant are printed in the left menu. Users could shift the yellow line and refit to obtain a better fit. When the user was satisfied with the fit result or could not obtain a passing fit the analysis was saved.

Figure Supplement 11: A model of quantal release and short term plasticity. **A.** PSP amplitude histograms for three synapses with the model's average predicted amplitude distribution overlaid. **B.** PSP amplitudes facilitating and reduction of synaptic failures across a 50 Hz spike train for the example L5e→Sst synapse in (A). **C.** Histograms showing PSP amplitudes for the same synapse with model distribution predictions overlaid, showing history-dependent adjustment in model state. **D.** Estimates of model likelihood (bright colors are higher likelihood) across a range of parameters for two synapses. Each image is a maximum projection across all other axes in the model parameter space. **E.** Model results support release-independent depression mechanisms. Left: histogram of the ratios between release-dependent max likelihood and release-independent max likelihood, showing an overall preference for release-independent model parameters. Right: comparison of the same RDD/RID ratio to paired event correlations, with little overall effect. Release-dependent depression should result in negative paired event correlations.

Figure Supplement 12: Comparison between model behavior and measured synapse features. For each synapse, the maximum likelihood model parameters were used to simulate experimental data. Measurements were then performed on the simulated response amplitudes and compared to identical measurements from the recorded data. **A-B.** Measures of synaptic strength correlate strongly with model results. **C-F.** Two measures of STP and variability. **G.** Resting state variability correlates with the binomial CV derived from model parameters. **H-I.** Release probability is more strongly correlated with variability than number of release sites.

Transgenic mouse line	Cre + reporter abbr.	FlpO + reporter abbrev.
Double Transgenic		
Pvalb-IRES-Cre;Ai14	Pvalb-TdTomato	
Sst-IRES-Cre;Ai14	Sst-TdTomato	
Nr5a1-Cre;Ai14	Nr5a1-TdTomato	
Tlx3-Cre_PL56;Ai14	Tlx3-TdTomato	
Sim1-Cre_KJ18;Ai14	Sim1-TdTomato	
Triple Transgenic		
Sst-IRES-Cre;Pvalb-T2A-FlpO;Ai193-hyg	Sst-EGFP	Pvalb-TdTomato
Vip-IRES-Cre;Sst-IRES-FlpO;Ai193-hyg	Vip-EGFP	Sst-TdTomato
Vip-IRES-Cre;Pvalb-T2A-FlpO;Ai193-hyg	Vip-EGFP	Pvalb-TdTomato
Sim1-Cre_KJ18;Sst-IRES-FlpO;Ai193-hyg	Sim1-EGFP	Sst-TdTomato
Sim1-Cre_KJ18;Vip-IRES-FlpO;Ai193-hyg	Sim1-EGFP	Vip-TdTomato
Sim1-Cre_KJ18;Pvalb-T2A-FlpO;Ai193-hyg	Sim1-EGFP	Pvalb-TdTomato
Quadruple Transgenic		
Sst-IRES-Cre;Ai140;Pvalb-T2A-FlpO;Ai65F	Sst-EGFP	Pvalb-TdTomato
Vip-IRES-Cre;Ai140;Sst-IRES-FlpO;Ai65F	Vip-EGFP	Sst-TdTomato
Vip-IRES-Cre;Ai140;Pvalb-T2A-FlpO;Ai65F	Vip-EGFP	Pvalb-TdTomato
Nr5a1-Cre;Ai140;Pvalb-T2A-FlpO;Ai65F	Nr5a1-EGFP	Pvalb-TdTomato
Nr5a1-Cre;Ai140;Sst-IRES-FlpO;Ai65F	Nr5a1-EGFP	Sst-TdTomato
Nr5a1-Cre;Ai140;Vip-IRES-FlpO;Ai65F	Nr5a1-EGFP	Vip-TdTomato
Pvalb-IRES-Cre;Ai140;Rorb-T2A-tTA2;Ai63	Pvalb-EGFP	Rorb-TdTomato
Tlx3-Cre_PL56;Ai140;Pvalb-T2A-FlpO;Ai65F	Tlx3-EGFP	Pvalb-TdTomato
Tlx3-Cre_PL56;Ai140;Sst-IRES-FlpO;Ai65F	Tlx3-EGFP	Sst-TdTomato
Tlx3-Cre_PL56;Ai140;Vip-IRES-FlpO;Ai65F	Tlx3-EGFP	Vip-TdTomato
Sim1-Cre_KJ18;Ai52;Pvalb-T2A-FlpO;Ai65F	Sim1-EGFP	Pvalb-TdTomato
Sim1-Cre_KJ18;Ai139;Sst-IRES-FlpO;Ai65F	Sim1-EGFP	Sst-TdTomato
Sim1-Cre_KJ18;Ai52;Vip-IRES-FlpO;Ai65F	Sim1-EGFP	Vip-TdTomato
Ntsr1-Cre_GN220;Ai140;Pvalb-T2A-FlpO;Ai65F	Ntsr1-EGFP	Pvalb-TdTomato
Ntsr1-Cre_GN220;Ai140;Sst-IRES-FlpO;Ai65F	Ntsr1-EGFP	Sst-TdTomato
Ntsr1-Cre_GN220;Ai140;Vip-IRES-FlpO;Ai65F	Ntsr1-EGFP	Vip-TdTomato
Double Transgenic + retroorbital AAV		
Tlx3-Cre_PL56;Ai193 + mscRE4-FlpO AAV PHPe.B	Tlx3-EGFP	mscRE4-TdTomato targets Layer 5 ET cells

Table Supplement 1 Transgenic Animals. List of transgenic animals used in experiments. Animals were derived from double, triple, or quadruple transgenics. In triple and quadruple transgenics, two subclasses were driven by either Cre or FlpO and expressed either TdTomato or EGFP. In one case we utilized a retroorbital AAV to label L5 ET cells together with L5 IT cells as it was not possible to construct this combination through a triple or quadruple transgenic.

Component	ACSF.I	ACSF.IV	ACSF.V	ACSF.VII	ACSF.IX
Calcium Chloride, dihydrate	0.5 mM	2 mM	1.8 mM	0.5 mM	1.3/2.0 mM
D-Glucose (Dextrose)	25 mM	25 mM	0	25 mM	25 mM
HCl	98 mM	0	0	92 mM	0
HEPES	20 mM	20 mM	5 mM	20 mM	0
Magnesium Chloride	0	0	1.0 mM	0	0
Magnesium Sulfate	10 mM	2 mM	0	10 mM	1.0/2.0 mM
Monosodium Phosphate	1.25 mM	1.25 mM	0	1.2 mM	1.2 mM
Myo inositol	3 mM	3 mM	0	0	0
N-acetylcysteine	12 mM	12.3 mM	0	0	0
N-methyl-d-glucamine	96 mM	0	0	92 mM	0
Potassium Chloride	2.5 mM	2.5 mM	5.4 mM	2.5 mM	3mM
Sodium Bicarbonate	25 mM	25 mM	0	30 mM	18 mM
Sodium Chloride	0	94 mM	135 mM	0	126 mM
Sodium L-Ascorbate	5 mM	5 mM	0	5 mM	0.16 mM
Sodium Pyruvate	3 mM	3 mM	0	3 mM	0
Taurine	0.01 mM	0.01 mM	0	0	0
Thiourea	2 mM	2 mM	0	2 mM	0

Table Supplement 2 ACSF Recipes. Concentrations of each component in ACSF recipes utilized in different stages of our experiments, slice, holding, recording, etc. See Methods for more information on when each ACSF was used.

QC Metric	Analysis level of data being evaluated	Voltage Clamp	Current Clamp
Holding current	Postsynaptic stimulus trial	±800 pA	±800 pA
Holding potential	Postsynaptic stimulus trial		-45 mV to -85 mV
Baseline noise	Postsynaptic stimulus trial	< 200 pA	< 5 mV
Presynaptic spike detected	Presynaptic recording in response to 1 stimulus pulse	✔	✔
Extra spikes	Presynaptic recording in response to 1 stimulus pulse	None within ±8 ms	None within ±8 ms
Standard deviation of postsynaptic response This checks for a relatively stable recording	Postsynaptic recording in response to 1 stimulus pulse	< 15 pA	< 1.5 mV
Max postsynaptic amplitude This checks for artifacts in the response	Postsynaptic recording in response to 1 stimulus pulse	< 500 pA	< 10 mV
Max postsynaptic potential	Postsynaptic recording in response to 1 stimulus pulse		-40 mV
Excitatory connection	Postsynaptic recording in response to one presynaptic spike	Membrane potential between -45 mV and -85 mV	
Inhibitory connection	Postsynaptic recording in response to one presynaptic spike	Membrane potential between -45 mV and -60 mV	

Table Supplement 3 Quality Control. Quality control stages for data processing.

Morphology annotation	Description
Dendrite type	
NEI (Not Enough Information)	Either no biocytin filled cell or poor health precludes definition of dendrite type
Aspiny	No visible spines, non -pyramidal morphology
Sparsely spiny	Visible spines, non -pyramidal morphology
Spiny	Visible spines, pyramidal or stellate morphology
Axon	
Truncation	Obvious cut axon or retraction bulb near surface of the slice that measured ≦ 100 μm
Borderline	Axon measured between 100 -200 μm
Intact	Traceable axon measured > 200 μm
Unclear	Axon untraceable but not obviously truncated, likely due to poor health or biocytin fill
Truncation distance	Measured length of axon from origin to end. Only measured in "truncated" or "borderline" axons
Apical dendrite	
Truncated	Does not reach Layer 1
Intact	Reaches Layer 1
Truncation distance	Measured length of largest apical branch from soma to end of dendrite. Only measured in "truncated" dendrites

Table Supplement 4 Morphology Annotations. Morphological annotations assigned to recorded cells filled with biocytin and imaged at 63x resolution.