Towards a unified theory of the reference frame of the ventriloquism aftereffect

Peter Lokša and Norbert Kopčo

Institute of Computer Science, Faculty of Science, P. J. Šafárik University in Košice, Jesenná

5, 04001 Košice, Slovakia

peter.loksa@upjs.sk, norbert.kopco@upjs.sk

Running Title: Reference Frame of Ventriloquism

ABSTRACT

The ventriloquism aftereffect (VAE), observed as a shift in the perceived locations of sounds

after audio-visual stimulation, requires reference frame alignment since hearing and vision

encode space in different reference frames (head-centered vs. eye-centered). Previous

experimental studies reported inconsistent results, observing either a mixture of head-centered

and eye-centered frames, or a predominantly head-centered frame. Here, a computational

model is introduced to examine these inconsistencies. Based on experimental data, the model

uses the measured size of the ventriloquism effect to predict the VAE adaptation in the

auditory spatial map. Versions of the model examine whether the adaptation is induced by

visual signals in head-centered frame, eye-centered frame, by eye-gaze direction-dependent

signals, or their combination, and whether some biases are induced by the saccade-to-

auditory-target response method used in the experiments. The model is first evaluated on three

separate data sets. It can predict them well even without explicit need for an eye-centered

signals influencing VAE, suggesting that the reference frame of VAE is mainly head-

centered. The model predictions are qualitatively similar but less accurate when all three data

sets are combined, suggesting that interactions between individual neural mechanisms are

more complex than the simple linear combination assumed in the model.

1. Introduction

Auditory spatial perception is highly adaptive and visual signals often guide this adaptation. In the "ventriloquism aftereffect" (VAE), the perceived location of sounds presented alone is shifted after repeated presentations of spatially mismatched visual and auditory stimuli (Bertelson et al., 2006, Recanzone, 1998, Woods and Recanzone, 2004). Complex transformations of spatial representations in the brain are necessary for the visual calibration of auditory space to function correctly, as visual and auditory spatial representations differ in many important ways (van Opstal, 2016). Here, we propose a computational model to examine the visually guided adaptation of auditory spatial representation in the VAE, the related transformations between the reference frames (RFs) of auditory and visual spatial encoding, and the effect that the saccade response method, used in many RF of VAE studies, might have on the results.

We primarily examine the RF in which the VAE occurs. While visual space is initially encoded relative to the direction of eye gaze, the cues for auditory space are computed relative to the head orientation (Groh and Sparks, 1992). A means of aligning these RFs is necessary by the stage at which the visual signals guide auditory spatial adaptation. Several models were developed to describe the ventriloquism aftereffect in humans and birds. The bird models predict the VAE in the barn owls (Haessly et al., 1995, Oess et al., 2020) which cannot move their eyes and therefore their auditory and visual RFs are aligned. The human models mainly focused on spatial and temporal aspects of the ventriloquism aftereffect (Beierholm et al., 2009, Bosen et al., 2018, Kording et al., 2007, Odegaard et al., 2015, Shinn-Cunningham et al., 2005, Watson et al., 2019, Wozny et al., 2010), not considering the different RFs. There are models of the audio-visual reference frame alignment, but those only

consider audio-visual integration (Odegaard et al., 2019, Razavi et al., 2007) and multi-

sensory integration (Pouget et al., 2002) when the auditory and visual stimuli are presented

simultaneously, not the adaptation and transformations underlying the VAE.

Our experimental studies examining RF of VAE in humans and monkeys provided

inconsistent results (described in detail in Section 2). A mixture of eye-centered and head-

centered RFs was identified for recalibration locally induced in the central region of the

audio-visual field (Kopco et al., 2009) while the head-centered RF dominated when was VAE

induced in the audio-visual periphery (Kopco et al., 2019). Additionally, a recent study in

which the VAE was induced over a wide spatial area also concluded that the RFs are mixed

(Watson, 2021). These results imply that the RF used in the VAE is dependent on the region

in which the VAE is induced, possibly due to a non-homogeneity in the auditory spatial

representation. Specifically, recent evidence suggests that, in mammals, auditory space

encoding is based on two or more spatial channels roughly aligned with the left and right

hemifields of the horizontal plane (Groh, 2014, Grothe et al., 2010), implying that the spatial

representation's adaptability might be spatially non-uniform. For example, it might be more

locally adaptable in the subregions of space covered by both hemispheric channels, i.e., in the

center, than in the regions dominated by one channel (periphery). Our current modeling

focuses on a simpler alternative in which the RF transformations are the same across the

audio-visual field, and the observed region-dependence of the RF is due to saccades to

auditory targets, as saccades, which are eye-centered, were used to measure VAE in our

previous studies.

A secondary goal of the current model to propose a mechanism to describe a new

adaptive phenomenon observed in the ventriloquism study of Kopco et al. (2019) (again,

described in detail in Section 2). In that study, adaptation was unexpectedly induced by

spatially aligned audio-visual stimuli, while no such adaptation was observed in Kopco et al.

(2009).

Here, we first summarize the experimental data from Kopco et al. (2009, 2019) to

explain the modeled phenomena (Section 2). Then, the model is introduced (Section 3) and

evaluated (Sections 4 and 5).

2. Summary of data of Kopco et al. (2009, 2019)

------------------------------ Insert Figure 1 around here ------------------------------

In two separate experiments, ventriloquism was induced by audio-visual training trials

either in the central (Kopco et al., 2009) or peripheral (Kopco et al., 2019) subregion of the

horizontal audio-visual field while the eyes fixated one fixation point (FP; red '+' symbol;

upper and middle panels of Figure 1A, respectively). The audio-visual stimuli consisted of a

sound paired with an LED. The aftereffect was evaluated on interleaved auditory-only (A-

only) probe trials using a wide range of target locations while the eyes fixated one of two FPs

(lower panel of Figure 1A). The listener's task in both audio-visual and auditory-only trials

was to perform a saccade to the perceived location of the auditory stimulus from the FP.

The responses to AV stimuli were always very near the visual components of the AV

stimuli, both in the AV-aligned baseline (green dashed and solid lines in Figure 1B

corresponding, respectively, to the peripheral and central experiments) and in the conditions

with the visual component shifted (green lines in Figure 1C). The displaced V component in

the AV-misaligned conditions induced a local ventriloquism aftereffect when measured with

the eyes fixating the training FP (the red solid and dashed lines in Figure 1C show that

maximum ventriloquism was always induced in the trained subregion of the auditory space).

The critical manipulation of these experiments was that a half of probe trials was performed

with eyes fixating a new, non-training FP (blue '+' symbol), located 23.5° to the left of the

training FP (red '+' symbol at +11.75°). If the RF of the VAE is purely head-centered, then

moving the eyes to the new FP location for the testing was expected to have no effect,

resulting in the same pattern of ventriloquism for the non-training and training FPs. On the

other hand, if the RF is purely eye-centered, the observed pattern of induced shifts was

expected to move with the eyes by the 23.5° when the probe trial is performed from the new,

non-trained fixation. The experimental data showed that, in the central experiment, moving

the fixation resulted in a smaller ventriloquism aftereffect with the peak moving in the

direction of eye gaze (blue vs. red dashed line), while in the peripheral experiment only a

negligible effect of the eye fixation position was observed (blue vs. red solid line).

To better visualize these results, the lower panel of Figure 1C shows data expressed as

difference between responses from training vs. non-training FPs from the respective upper

panels, along with the expected patterns of results for the two RFs. The head-centered RF

always predicts that the effect would be identical for the two FPs. Thus, all head-centered

differences (brown lines) are expected at zero. The solid and dashed yellow lines show,

respectively, for the peripheral and central data, the eye-centered RF expected patterns

obtained by subtracting from the red lines the same red lines shifted 23.5° to the left. Finally,

the black solid and dotted lines show the actual differences between the respective red and

blue data from the upper panel. For the central data, the black dashed line falls approximately

in the middle between the head-centered and eye-centered predictions, showing a mixed

nature of the RF of the VAE induced in this region. On the other hand, the black solid line is

always near zero, showing that the RF of the VAE induced in the periphery is predominantly

head-centered. The main goal of the current modeling is to examine this inconsistency.

While the results in Figure 1C are based on the ventriloquism aftereffect induced by

AV-misaligned stimuli, Figure 1B shows the baseline data obtained in runs with auditory and

visual stimuli aligned. In the central-training experiment, the responses from the two FPs were

similar, unbiased at the central locations and with a slight expansive bias in the periphery

(both red and blue dotted lines are near zero in the center, negative in the left-hand portion

and positive in the right-hand portion of the graph). On the other hand, in the peripheral-

training experiment the responses for the middle three targets differed between the two

fixations, where the non-training FP responses fell well below the training-FP responses

(compare the red and blue solid lines). Thus, the peripheral AV-aligned stimuli induced a

fixation-dependent adaptation in the auditory-only responses in the central region, an

adaptation phenomenon that has not been previously described. The black dashed and solid

lines in Figure 1B showing the difference between the corresponding data from the upper

panel, highlight the FP-dependence of the peripheral data in contrast to the FP-independence

in the central data. The secondary goal of the current modeling is to propose a mechanism to

explain this result.

3. Model Description

3.1 Overview

-------------------------------- Insert Figure 2 around here --------------------------------

Figure 2A shows the outline of the model. The model predicts the ventriloquism

aftereffect (i.e., the azimuthal bias in a response to an auditory-only probe; red and blue lines

in Figure 1B-C) as a function of the A-only probe azimuth and the FP location (the

"Response" vs. "Probe stimulus and FP" blocks in Figure 2A). To generate the prediction, it

only uses information about the training AV stimulus locations and the measured AV

response biases for those locations (i.e., the ventriloquism effect; green lines in Figure 1B-C)

for a given type of AV stimuli ("Ventriloquism" block in Figure 2A). The prediction is

computed as an additive combination of two components (the "sum" operation). The main

component is the "Auditory space representation" block (Bosen et al., 2018) which is locally

adapted by the audio-visual ventriloquism signals in different reference frames ("HC" and

"EC" arrows) and which can also be influenced by information about eye position ("FP-

dependent attenuation"). The second component, the "Saccade-related bias" block, represents

the saccade response bias in eye-centered reference frame, which is not directly adapted by

ventriloquism and which is assumed to be present *a priori* when saccades are used as a

response method. Finally, it is assumed that there are interactions between the saccade-related

and ventriloquism effects (the "Saccade-related bias" block is linked to the "Auditory space

representation" block).

The current model assumes that the auditory space representation uses a uniform

population code (Carlile et al., 2001), not a hemispecific space encoding (Lingner, 2018). It

describes the VAE adaptation by only considering the locations from which the auditory

components of audio-visual stimuli were presented during training. Then, it assumes that the

induced VAE at those locations is directly proportional to the measured ventriloquism effect

and those locations, and that it decreases with distance from those locations. Thus, the model

does not require input information about the direction of audio-visual stimulus displacement

during training (whether the visual stimuli were shifted to the left, right, or aligned with the

auditory stimuli). Instead, it only uses the information about where the training occurred and

what the resulting ventriloquism effect was, and it assumes that there is a direct relation

between the observed ventriloquism effect and aftereffect. Supporting this assumption, a

comparison of the VE and VAE data at the trained locations (corresponding green and red

lines in Figure 1C) shows that the VAE is approximately a half of VE in the experimental

data.

There are four versions of the model, differing by whether they include optional

components shown in gray in Figure 2A ("EC" arrow and "FP-dependent attenuation"

subblock), which represent two different hypotheses about how EC signals might influence

the RF of VAE. The basic version of the model, referred to as "HC model", does not include

either of the optional components. Thus, it predicts that the ventriloquism signals influencing

the spatial auditory representation are purely head-centered (the "HC" arrow in Figure 2A)

and all the eye-centered contributions to the observed RF of VAE come from the saccade-

related biases.

In the "HEC model" version, the visual signals adapt the auditory spatial

representation in both head-centered and eye-centered RFs (the optional "EC" arrow) such

that the relative contribution of the HC and EC RFs can be arbitrary. Therefore, the HEC

model reduces to the HC model if the weight of the EC path is set to zero, or it can produce

predictions using only EC RF if the HC path weight is set to zero. Note that a purely EC-

based version of the model was not considered as 1) the peripheral-experiment data only are

consistent with a HC reference frame, so the model would clearly fail, and 2) the HEC model

could behave as such EC-only model by appropriately adjusting the relative weight, if that

were the best fit to the data.

The "dHC model" version assumes that the adaptation of spatial representation

induced by ventriloquism is attenuated when the eyes shift to a new FP away from the

training FP ("FP-dependent" attenuation sub-component), and that this attenuation is

proportional to the distance between the training and new FP. This mechanism might be

related to FP-dependent biases observed in sound localization (Lewald & Ehrenstein, 1998;

Razavi et al., 2007), and it does not require saccades to be used for responses. Since this

attenuation is dependent on the fixation location, it is also in the eye-centered reference frame.

195   Finally, the "dHEC model" incorporates both optional components and thus it assumes

196   that both EC-referenced mechanisms described in the HEC and dHC models affect

197   performance.

198   *3.2 Detailed Specification*

199   The following model specification applies to the most general dHEC model version,

200   with the differences applying to the reduced versions described as needed (all variables in the

201   model use the head-center representation and are in the units of degrees, unless specified

202   otherwise). Panels B-D of Figure 2 provide visualizations of the behavior of the individual

203   model components.

204   Equation 1 describes the predicted bias in responses $\hat{r}$ to a given auditory stimulus at

location s and for eyes fixating the location f, as a weighted sum of a saccade-related bias r_E

and a ventriloquism-related adaptation in auditory spatial representation r_V

$$\hat{r}(s,f) = r_E(s,f) + w \cdot r_V(s,f), \tag{1}$$

Where $w \in [0, \infty]$ is a free parameter specifying the relative weight of the ventriloquism

adaptation. The prediction is determined by considering the observed biases in AV stimulus

responses r_{AV} at the training AV stimulus locations s_{AV} (as described below in Eq. 3).

211   The saccade-related bias at a specific location x for eyes fixating the location f is

modeled as a sigmoidal function

$$r_E(x,f) = \frac{2 \cdot h}{1+e^{-k(x+cf)}} - 1, \tag{2}$$

where $h, k,$ and c are free parameters characterizing the sigmoid. The saccade-related bias

(Figure 2B) is broad and referenced to the FP (i.e., it uses EC RF), exhibiting a combination

of underestimations and overestimations commonly observed in studies of saccades to

auditory targets (Gabriel et al., 2010, Razavi et al., 2007, Yao and Peck, 1997). While similar

functions were previously used to model visually induced spatial auditory adaptation (Zwiers

et al., 2003), the specific shape of the functions used here was chosen to be consistent with

results of Gabriel et al. (2010), which observed both underestimation and overestimation in

saccade responses depending on the target location (or, it can be a result of underestimation of

saccade responses (Yao and Peck, 1997) combined with overestimation in peripheral auditory

localization estimates with a centrally fixed eyes (Bosen et al., 2018)). The resulting

predictions roughly follow the values observed in the peripheral and central no-shift data in

the current study (Figure 1B).

The ventriloquism-driven auditory space adaptation causes bias defined at location x,

for eyes fixating the location f, and for ventriloquism induced at training locations s_{AV}

causing AV response biases r_{AV}, as:

$$r_V(x,f) = d_f \cdot \sum_{i=1}^{N} w_{v,i}(x,f) \cdot \left[r_{AV,i} - r_E\left(s_{AV,i}, f\right) \right], \qquad (3)$$

where d_f is the FP-dependent attenuation of the aftereffect in the dHC and dHEC

models (Eq. 8), N is the number of training locations ($N = 3$ for the current study), i is an

index through these locations, $s_{AV,i}$ is the i-th training location azimuth, and $r_{AV,i}$ is the AV

response bias observed at the i-th training location (the green diamonds in Figure 2B

represent the values of $r_{AV,i}$ in an example simulation in which these values are around 0 and

in which $s_{AV,i}$ are at -7.5, 0, and +7.5°). The differences $r_{AV,i} - r_E\left(s_{AV,i}\right)$ represent the

disparity between the AV response biases (green diamonds) and the saccade-related bias

(red/blue lines in Figure 2B) at the training locations (where the r_E value explicitly

incorporates an interaction between the ventriloquism and the saccade components, in

addition to the ventriloquism effect). The differences are shown in Figure 2C by the red and

blue filled diamonds. $w_{v,i}(x)$ is the strength with which the disparity at the i-th training

location adapts the spatial representation at the location x. In the HEC and dHEC models, this

value is a weighted sum of the adaptation strengths in head-centered and eye-centered

reference frames, defined as:

$$w_{v,i}(x,f) = (1 - w_E) \cdot w_{vH,i}(x) + w_E \cdot w_{vE,i}(x,f), \qquad (4)$$

245   where $w_E \in (0,1)$ is a parameter determining the relative weight of the EC reference frame

246   vs. the HC RF (in the HC and dHC models, $w_E = 0$). Weights $w_{vH,i}$ and $w_{vE,i}$ use normalized

247   Gaussian functions centered at the training locations as a measure of the influence of i-th

248   training location on target at location x:

$$w_{vH,i}(x) = G\left(x, s_{AV,i}, \sigma_H\right), \qquad (5)$$

$$w_{vE,i}(x) = G\left(x, s_{AV,i} + f - f_T, \sigma_E\right), \qquad (6)$$

$$G(x, \mu, \sigma) = \frac{N(x,\mu,\sigma)}{\sum_{i=1}^{N} N(7.5 \cdot (i-2), 0, \sigma)}, \qquad (7)$$

In Eqs. 5 and 6, the parameters σ_H and σ_E represent the width of the influence of the

ventriloquism shift at individual training locations, respectively, for the two reference frames.

$w_{vH,i}$ (Eq. 5) is always centered on the i-th training location in the HC RF, whereas $w_{vE,i}$ (Eq.

6) is centered on the i-th training location in the EC RF. f_T is the training fixation location,

equal to 11.75° in the current study (when $f = f_T$, the two RFs are aligned). Finally, the

Gaussian functions are normalized (Eq. 7) such that the maximum $w_{vH,i}$ or $w_{vE,i}$ after

summing across the three training locations is 1 (the normalization locations $7.5 \cdot (i - 2)$ are

specific for the current training as $7.5°$ is the separation between the loudspeakers used in the

experiments; they need to be modified to model data from experiments using other training

locations).

262   The FP-dependent attenuation of the aftereffect from the non-training FP is defined as

263                                 $$d_f = \exp(-d \cdot |f - f_T|) \tag{8}$$

264   where the free parameter d is the rate of attenuation with distance from the current fixation f

265   to the training fixation f_T (in the HC and HEC models, $d = 0$). Note that the exact form of

266   this dependence cannot be evaluated for the current data as only one training and one non-

267   training fixation were evaluated (i.e., $|f - f_T|$ only had values of 0 or $23.5°$).

268   Figure 2C shows the operation of the ventriloquism adaptation. As mentioned above,

the red and blue filled diamonds are the disparities at the individual training locations driving

the adaptation in HC RF. The blue open diamonds are identical to the blue filled diamonds

except that they are shifted to the left by the difference between the two FPs to illustrate how

the eye gaze shift affects where the adaptation is expected to occur in the EC RF. The red and

blue lines are then the resulting biases r_V for the two fixation locations, each corresponding to

the sum of Gaussians centered at different training locations in the two RFs (and with widths

defined by the σ's). Parameter w_E determines the relative weights of the peaks in the portion

of blue line corresponding to the open diamonds vs. the portion corresponding to the filled

diamonds. Finally, parameter d_f determines how much the blue line deviation from 0 is

reduced compared to the red line (i.e., how much the aftereffect is attenuated at a new fixation

location). So, the blue and red lines show how visually guided adaptation is local and RF-

dependent, decreasing with distance from the AV stimulus locations in HC and EC RFs. It

also shows that since adaptation causes shifts from the saccade-bias response locations

towards AV response locations, if AV responses fall on saccade bias locations, no visually

guided adaptation is predicted to occur. Finally, Figure 2D shows that the model prediction is

a sum of the saccade bias (from Figure 2B) and ventriloquism bias (Figure 2C) weighted by

the parameter w.

4. Simulation Methods

4.1 Stimuli

In the studies of Kopco et al. (2009, 2019), ventriloquism was induced by presenting

AV training stimuli with visual component either shifted to the left of the auditory component

(negative shift), to the right of it (positive shift), or aligned with it (no shift), while the eyes

fixated one location (Figure 1A; upper and middle panels). Thus, nominally, there were 6 AV

conditions (3 shift directions by 2 training regions). For these conditions, predictions could be

compared to data for 9 A-only probe locations and 2 FPs (Figure 1A; lower panel). However,

the main experimental results simulated here were observed after performing two

subtractions: 1) subtracting the training-FP from the non-training-FP data, and 2) subtracting

the positive-shift data from negative-shift data and halving the result, which gave the average

aftereffect magnitude differences (black lines in Figure 1C). These "double differential"

("positive – negative" difference of "training FP – non-training FP" difference) data were the

most stable as they eliminated a lot of between-subject variability related to individual biases

in responses (as will be illustrated later). Therefore, to evaluate the model on these key

differences, the data were also transformed into the difference representation in two steps,

resulting in representation that is equivalent to the difference plots in the bottom panels of

Figure 1B and C (Footnote 1).

The complete data set consisted of 108 data points [9 location x 2 (transformed) FPs x

3 (transformed) shifts x 2 training regions]. Across-subject mean and standard deviation data

were used in the simulations. The model is fitted to the auditory-only trial responses, while

the AV responses are used by the model as parameters for different conditions.

4.2 Model Fitting and Evaluation

In the simulations, the four models' performance was compared on different subsets of

the data. The basic rationale behind these comparisons was that if the minimal HC model

predicts the data well, then there is no evidence that eye-centered signals affect the

ventriloquism aftereffect, and all the observed eye-dependent effects can be ascribed to the

saccade-related bias. On the other hand, if one of the HEC, dHC, and dHEC models performs

best, it is evidence that, even after accounting for the saccade-related effects, the auditory

spatial representation receives some eye-centered signals, causing adaptation that always

depends on the position of the stimuli relative to the eye gaze direction.

-------------------------------- Insert Table 1 around here --------------------------------

Each simulation was performed by fitting the four models to the corresponding subset

of the transformed data using a two-step procedure. First, a systematic search through the

parameter space was performed, using all combinations of 10 values for each parameter, listed

in Table 1 (HC model used 5 parameters, dHC 6, HEC 7, and dHEC 8 parameters). The

ranges were chosen by piloting to cover the expected behavior of the model in different

conditions. Second, the best 100 parameter combinations in terms of weighted MSE were

used as starting positions for non-linear iterative least-squares fitting procedure (Matlab

function lsqnonlin) which, again, minimized the weighted MSE. The parameter values for the

best of these fits were chosen as the optimal values listed in Table 2 and used in the result

figures.

To compare the models' performance while accounting for the number of parameters

used by each model, we computed the Akaike information criterion AICc (Burnham and

Anderson, 2004, Taboga, 2017) for each optimal fit, defined as:

$$\text{AICc} = -2\log(L + 2K + 2K\frac{K+1}{n-K-1}, \qquad (8)$$

$$\log(L) = \frac{-n}{2}\left(\log(2\pi) + \log\frac{\text{SSE}(X)}{n} + 1\right) \qquad (9)$$

where n is the number of experimental data points, K is the number of fitted

parameters, and $\text{SSE}(X)$ is the sum of squares of errors across the data points (i.e., differences

between predictions and across-subject mean data x_i) weighted for each data point by the

inverse of its across-subject standard deviation $\frac{1}{SD(x_i)}$. In general, the model with the lower

AICc is considered to be a better fit for the data. We use the rule that the model with the lower

AICc is substantially better than an alternative model only if the rounded up value ΔAIC is

larger than 2.

5. Results

Four model evaluations were performed, each assessing the four model variants on a

different subset of the Kopco et al. (2009, 2019) data. No Shift evaluation assessed the models

on the AV-aligned no-shift data from both experiments (Figure 1B), while in the Central Data

and Peripheral Data evaluations, the respective data from the AV-misaligned conditions in the

two experiments (dashed and solid lines in Figure 1C) were fitted separately. Finally, in the

Combined Data evaluation the models were fitted on the complete dataset from both

experiments (Figure 1B and C). The results of the 4 evaluations are summarized in Table 2,

which shows for each simulation and model the fitted parameter values and the model's

performance measured using the AICc criterion and the weighted MSE.

------------------------------- Insert Table 2 around here --------------------------------

5.1 No-shift data

------------------------------- Insert Figure 3 around here --------------------------------

This evaluation focused on the AV-aligned data, examining the hypothesis that *the*

saccade-related bias combined with auditory space representation adapted in HC RF are

sufficient to explain the training-region-dependent differences in the AV-aligned data (Figure

1B). I.e., it was predicted that the HC model can accurately describe the baseline data, and

that the optional model components do not need to be considered to explain the different

adaptation effects observed in central vs. peripheral AV-aligned training.

Figure 3 presents the results of the model evaluation and illustrates the function of

individual model components. Panels A and B show the data (circles) and predictions of the

models (lines) with the fitted parameters as listed in Table 2. (Note that only the HC and HEC

models are shown and discussed, as performance was identical for the other two models). The

upper row shows the main results for the difference between the FPs (identical to the black

lines from Figure 1B, now with error bars included), while the middle and bottom rows show

the results separately for the two FPs.

Both models captured very well the difference data, which are near zero for the central

training experiment and have a positive deviation for the peripheral training (upper row of

panels A and B). This conclusion is confirmed by the AICc evaluation which showed no

evidence that either of the models should be preferred even though the HEC model provides a

slightly better fit ($\Delta AIC = 2.4$ for HC). Also, the models captured all the important trends in

the data for the two FPs (middle and bottom row). Specifically, for the central training data,

they captured the slight expansion of the space for the central training data identical for both

FPs (panel A), as well as the FP-dependence of the peripheral training data at the central

locations (panel B).

A comparison of the error bars shows that there was a lot of across-subject variability

when the individual FPs were considered (middle and bottom row), while a large portion of

that variability was eliminated when the differences in biases across the FPs were computed

(upper row). This illustrates why the models were fitted on the transformed data, as those

were much more consistent across subjects, and, with the transformation, the fitting weighed

the difference data (upper row) more, as they were much more reliable (FOOTNOTE 2).

Panels C and D show the predictions of the two model components (columns)

separately for the two FPs (rows) for the central training data, such that the rows are aligned

with the resulting central-data prediction in the same row of panel A (the peripheral

predictions are based on the same model components except that the Auditory space

prediction is shifted by 23.5° to the right from that in Figure 3D; not shown). Each prediction

is, conceptually, a weighted sum of the two components (FOOTNOTE 3). The upper and

lower panels of Figure 3C show that both models predicted similar saccade-related bias,

consisting of expansion (bias toward periphery) at the peripheral target locations (+/-15°, +/-

22.5°, and +/-30°) and bias towards the fixation location for the central 3 locations (-7.5° to

+7.5°). This saccade-related bias was then combined with the auditory space adaptation such

that, at the training locations, the model predictions were shifted towards the AV response

biases, which were near zero for both the central and peripheral training. The HC model

predicts that this ventriloquism shift only occurred in HC RF (beige lines in the upper and

lower panels of Figure 3D), while the HEC model predicts a considerable contribution of the

EC RF (magenta lines at locations -30° to -15° in the bottom panel and at locations -7.5° to

7.5° in the top panel of Figure 3D). However, that contribution only had a small effect on the

overall predictions because the difference between the AV responses and the saccade-related

prediction, which scales the auditory space adaptation from panel D, was small for the no-

shift data, resulting in the small differences between the beige and magenta lines in panel A.

5.2 Central data

401   Central Data simulation only fitted the central-training data from the AV-misaligned

conditions (dashed lines in Figure 1C). For the mixed reference frame observed in these data,

the simulation examined *whether the eye-referenced component is more consistent with the*

eye-referenced shift in adaptation region mechanism (HEC model) or the FP-dependent

attenuation mechanism (dHC model). Figure 4A presents the results of this simulation. The

top panel of Figure 4A shows the FP difference for the AV-misaligned data and the

corresponding model predictions. The HC model's prediction (beige) is fixed at zero, while

the remaining three models fit the central-training data better, confirming that eye-referenced

signals contribute to the ventriloquism adaptation in central region (the improvement vs. HC

model in terms of AICc ranges from 5.9 to 15.6). Further, the HEC model's AICc is worse by

9.6 compared to the dHC model, providing a strong evidence that the mixed reference frame

observed behaviorally is driven by FP-dependent attenuation (dHC), not by ventriloquism

signals in the EC RF (HEC). The HEC model (pink) underestimates the central data for

targets at azimuths around 0° while it predicts a negative deviation at azimuths around -20°,

not observed in the data, which the dHC (green) model does not predict. Finally, the dHC and

dHEC models are comparable in terms of the AICs (difference of 1.4), while the dHEC model

(purple) has the lowest MSE error, indicating that the EC-referenced shift in adaptation region

mechanism might have a minor additional contribution to the adaptation effect.

The middle and bottom panels of Fig 4A show the data and model predictions for the

two FPs separately. All the models capture the basic profile of the adaptation. For the training

FP (middle panel) the dHC model (green) is closer to the data than the HEC model (pink), in

particular for the three left-most azimuths for which the difference prediction had the largest

departure from the data (top panel). Interestingly, for the non-training FP (bottom panel), the

HEC model captures the left-most triplet values better than the other two models. However,

this improved non-training FP prediction results in the non-training FP values being larger

than the corresponding training-FP values (pink line in the bottom panel is above the pink line

in the middle panel), causing the difference prediction (top panel) to have the negative

undershoot, not observed in the data.

The middle and lower panels also illustrate the functioning of the model for the AV-

misaligned data for which the Auditory space representation component dominates the

predictions. The dHC prediction (green line) in the lower panel is simply a scaled down

version of the prediction from the middle panel, while the HEC prediction (pink) has two

Gaussian components that are horizontally aligned and combined in the middle panel, while

one of which is shifted to the left in the bottom panel. The dHEC model (purple) combines

these two mechanisms to obtain predictions that tend to be closest to the data.

-------------------------------- Insert Figure 4 around here --------------------------------

5.3 Peripheral data

Peripheral data simulation fitted only the peripheral-training data from the AV-

misaligned conditions (solid lines in Figure 1C). The upper panel of Figure 4B shows that, in

agreement with the data, all four models produced almost identical predictions of

approximately 0-degree difference, confirming that the reference frame in this experiment

was largely head-centered. Also, Table 2 shows that the HC model was indeed the best in

terms of AICc and the EC-contribution parameters indicate a low contribution in the extended

models ($d_f = 1$ and $w_E = 0.04$). The middle and bottom panels of Figure 4B show that the

models captured the observed ventriloquism from individual FPs well, except for the two left-

most points, for which the effect appears to be negative, which the current models cannot

predict as Gaussians are used to describe the local character of the ventriloquism aftereffect.

6.4 Combined data

In the final evaluation, the four models were fitted to all data, combining the AV-

aligned and AV-misaligned data from central and peripheral experiments (solid and dotted

lines from Figure 1B and 1C). Figure 5 presents the results of this simulation in a layout

similar to Figure 4, with the no-shift data added at the bottom. Overall, the model predictions

in this simulation are less accurate than in the separate data set simulations (sections 6.1-6.3).

The upper row of Figure 5 shows that the models tend to underestimate the FP-difference in

the central data, in particular at azimuths near 0°, while they overestimate the FP-difference in

the peripheral data, especially at the right-most azimuths. This pattern of results is caused by

the largely linear operation of the model, which causes that the peripheral-data and central-

data predictions are approximately identical when aligned with respect to the training region

(i.e., the individual lines in the right-hand panel, when shifted to the left by 23.5°, are almost

identical to the lines in the left-hand panel). Then, to minimize the error for both central and

peripheral data, the model predictions are approximately in the middle of these two data sets.

However, even with this constraint, the dHC and dHEC models perform significantly better

than the HC and HEC model in terms of AICc (Table 2), again supporting the conclusion that

the FP-dependent attenuation mechanism is the most likely mechanism causing the mixed

reference frame observed for the central data, while the HEC mechanism only has a small

contribution.

467   ------------------------------ Insert Figure 5 around here --------------------------------

468   The two middle rows of Figure 5 show predictions separately for the two FPs. Overall,

469   all the models predict the peripheral data well (right-hand column). On the other hand, the

470   predictions for the central data show larger departures, especially for the middle targets and

471   non-training FP, for which the bias in data is reduced more than the models predict. As

472   discussed above, this is a limitation due to linear operation of the models.

473   Finally, the predictions for the AV-aligned data (bottom row) are again worse than

when these data were considered separately (Figure 3), while the models do qualitatively

capture that the biases are larger for the middle targets in the peripheral (panel B) than in the

central (panel A) region. This less accurate prediction is again caused by linear combination

of the saccade-related and auditory-space adaptation components in the model. Notably, all 4

models produce identical predictions of the AV-aligned data, since these predictions are

mostly based on the saccade-related bias model component which is nearly identical across

the 4 models.

6. Summary and Discussion

482   Our study introduced a model that combines ventriloquism-driven auditory space

adaptation with saccade-related biases to predict the reference frame of the ventriloquism

aftereffect. The model considers two forms of eye-centered signals influencing the auditory

space representation: ventriloquism signals in eye-centered reference frame and FP-dependent

attenuation. The main evaluation of the model, performed on the data from Kopco et al.

(2009), found the FP-dependent attenuation mechanism to be the main eye-centered

mechanism that caused the reference frame to be identified as mixed in that study. This result

suggests that the auditory space representation is natively head-centered, and the FP-

dependent signals only modulate the strength of the responses, thus making the result appear

as mixed. Second, the model was also able to explain a new form of auditory space adaptation

induced by AV-aligned stimuli in Kopco et al. (2019) by assuming a specific form of

interaction between saccade-related bias and the ventriloquism adaptation. Finally, when the

AV-aligned and AV-misaligned data from both studies were combined, the model predictions

became less accurate, likely due to the limited linear interactions of the model components

considered here. However, the model evaluation on the combined data still qualitatively

supported the conclusions obtained in the separate evaluations.

The current model only uses the responses on AV training trials to predict the

ventriloquism adaptation, independent of the size of the audio-visual disparity or of whether

the disparity results in hypometric or hypermetric saccades. And it assumes that the ratio of

observed ventriloquism aftereffect to the effect is constant, in our studies at approximately

0.5. With this simple assumption the model can also be applied to predict the results of other

ventriloquism aftereffect studies, even those in which the ventriloquism effect was not

measured, and those that did not use saccades for responding (in the latter case, the saccade-

related bias component of the model can be simply omitted).

The model assumes that the ventriloquism aftereffect measured by saccades is

influenced by some adaptive processes affecting the motor representations guiding the

saccades to AV and/or auditory targets. This mechanism cannot be directly verified as

currently available data are not consistent in terms of whether saccades to auditory targets are

predominantly hypermetric or hypometric (Gabriel et al., 2010, Yao and Peck, 1997), while

even less is known about saccades to misaligned AV targets. Additionally, eye-gaze-

direction-dependent biases in sound localization have been previously observed even when

saccades are not used for responding (Lewald and Ehrenstein, 1998, Razavi et al., 2007), and

these are likely also influence the measured saccade biases. To tease these contributions apart,

future studies need to assess the RF using response methods other than saccades (Kopco et al.,

2015, Lewald and Ehrenstein, 1998).

The neural mechanisms of the ventriloquism aftereffect and its reference frame are not

well understood. Cortical areas involved in ventriloquism aftereffect likely include Heschl's

gyrus, planum temporale, intraparietal sulcus, and inferior parietal lobule (Michalka et al.,

2016, van der Heijden et al., 2019, Zatorre et al., 2002, Zierul et al., 2017). Multiple studies

found some form of hybrid representation or mixed auditory and visual signals in several

areas of the auditory pathway, including the inferior colliculus (Zwiers et al., 2004), primary

auditory cortex (Werner-Reiss et al., 2003), the posterior parietal cortex (Duhamel et al.,

1997, Mullette-Gillman et al., 2005, Mullette-Gillman et al., 2009), as well as in the areas

responsible for planning saccades in the superior colliculus and the frontal eye fields (Schiller

et al., 1979, Wallace and Stein, 1994). In the current model, the saccade-related component

likely corresponds to the saccade-planning areas. The auditory space representation

component likely corresponds to the primary or the higher auditory cortical areas, or the

posterior parietal areas. There is growing evidence that, in mammals, auditory space is

primarily encoded based on two or more spatial channels roughly aligned with the left and

right hemifields of the horizontal plane (Dingle et al., 2012, Groh, 2014, Grothe et al., 2010,

McAlpine et al., 2001, Salminen et al., 2009, Stecker et al., 2005). Considering such an

extension might improve the model's ability to predict the central and peripheral data

simultaneously.

While most recalibration studies examined the aftereffect on the time scales of

minutes (Radeau and Bertelson, 1974, Radeau and Bertelson, 1976, Recanzone, 1998, Woods

and Recanzone, 2004), recent studies demonstrated that it can be elicited very rapidly, e.g., by

a single trial with audio-visual conflict (Wozny and Shams, 2011). If it is the case that the

adaptive processes underlying the ventriloquism aftereffect occur on multiple time scales, as

also suggested in several models of slower ventriloquism aftereffect (Bosen et al., 2018,

Watson et al., 2019), then an open question is whether the reference frame is the same at the

different scales or whether it is different. The current results are mostly applicable to the slow

adaptation on the time scale of minutes, while the RF on the shorter time scales has not been

previously explored.

In summary, our study introduced a model of the reference frame of the ventriloquism

aftereffect, while also considering the effect of saccade response biases on the measured

reference frame. The model assumes simple linear combinations of model components to

predict the data, which was sufficient to predict three data sets separately, but less so when

predicting combined data. Future studies need to expand this model to be able to predict the

combined data simultaneously, e.g., by consider non-linear combinations of the model

components.

Acknowledgments

This work was supported by the Slovak Scientific Grant Agency VEGA 1/0355/20 and

by EU Danube Region Strategy grant ASH (Grant Nos. APVV DS-FR-19-0025, WTZ MULT

07/2020, 45268RE). The authors thank Piotr Majdak for his comments on an early version of

this manuscript.

Footnotes:

Footnote 1: First, the data for the two training FPs were orthogonally transformed

such that instead of using training and non-training FP, a sum and a difference across the two

FPs was used. I.e., instead of having for each shift condition 18 data points corresponding to 9

locations at 2 FPs, we used 18 data points consisting of 9 locations summed across the two

FPs and 9 locations for difference across the 2 FPs. Specifically, if $x_{i,j,k}$ represents a data

point for the i-th target location (1 … 9), j-th AV-shift direction (P - positive, N - negative, 0

- no) and k-th fixation location (T – training, N – non-training), the first transformation

produced a data set y with the same dimensionality as x where each data point $y_{i,j,l}$ is

computed as:

$$y_{i,j,S} = x_{i,j,T} + x_{i,j,N}$$

$$y_{i,j,D} = x_{i,j,T} - x_{i,j,N}.$$

I.e., only the third dimension is changed such that the index l represents the operation

performed across the fixations (S – sum; D – difference).

Second, the positive-shift and negative-shift condition data were transformed in a

similar way, such that instead of positive and negative AV shift we used the aftereffect

magnitude (i.e., a halved difference between the two shifts) and average across the two shifts.

The no-shift data were left unmodified. Specifically, the second transformation produced a

data set z with the same dimensionality as y where each data point $z_{i,m,l}$ is computed as:

$$z_{i,M,l} = 0.5\left(y_{i,P,l} - y_{i,N,l}\right)$$

$$z_{i,A,l} = 0.5\left(y_{i,P,l} + y_{i,N,l}\right)$$

$$z_{i,0,l} = y_{i,0,l}.$$

575   Here, only the second dimension is changed such that the index m represents the operation

576   performed across the AV-shifts (M – magnitude of the aftereffect; A – average across positive

577   & negative shifts; 0 – no shift). Note that the $z_{i,M,l}$ data are plotted as black lines in the bottom

578   right panel of Figure 1C and the $z_{i,0,l}$ data are plotted in the bottom panel of Figure 1B.

579   **Footnote 2:** Note that the second half of the transformed data set, the average across

580   FPs, is not shown, as it can be easily visualized as an average of the individual FP data in the

581   lower two rows of panel B.

582   **Footnote 3:** The prediction is obtained as a combination of the saccade-related bias

583   and the auditory space component scaled 1) by the difference between the AV-training data

584   and the saccade-related bias, and 2) by the weight parameter w (Eqs. 1 and 3). The auditory

585   space graphs in Figure 3B are shown for the values of d_f and $[r_{AV} - r_E]$ fixed to 1 to illustrate

586   the differences between the two models. However, actual values of the parameters need to be

587   considered to generate the predictions in Figure 3A and 3B.

References

BEIERHOLM, U., QUARTZ, S. & SHAMS, L. 2009. Bayesian priors are encoded

independently from likelihoods in human multisensory perception. *Journal of Vision*

ARVO Journal, 9.

BERTELSON, P., FRISSEN, I., VROOMEN, J. & DE GELDER, B. 2006. The aftereffects of

ventriloquism: Patterns of spatial generalization. *Perception and Psychophysics,* 68**,**

428-436.

BOSEN, A. K., FLEMING, J. T., ALLEN, P. D., O'NEIL, W. E. & PAIGE, G. D. 2018.

Multiple time scales of the ventriloquism aftereffect. *PLoS ONE,* 13.

BURNHAM, K. P. & ANDERSON, D. R. 2004. Multimodel Inference Understanding AIC

and BIC in Model Selection. *Sociological Methods & Research,* 33**,** 261-304.

CARLILE, S., HYAMS, S. & DELANEY, S. 2001. Systematic distortions of auditory space

perception following prolonged exposure to broadband noise. *Journal of the*

Acoustical Society of America, 110**,** 416-424.

DINGLE, R. N., HALL, S. E. & PHILLIPS, D. P. 2012. The three-channel model of sound

localization mechanisms: interaural level differences. *J Acoust Soc Am,* 131**,** 4023-9.

DUHAMEL, J.-R., BREMMER, F., BENHAMED, S. & GRAF, W. 1997. Spatial invariance

of visual receptive fields in parietal cortex neurons. *Nature,* 389**,** 845-848.

GABRIEL, D. N., MUNOZ, D. P. & BOEHNKE, S. E. 2010. The eccentricity effect for

auditory saccadic reaction times is independent of target frequency. *Hearing Res,* 262**,**

19-25.

GROH, J. M. 2014. Making space: how the brain knows where things are. *Cambridge, MA:*

Harvard University Press.

GROH, J. M. & SPARKS, D. L. 1992. Two models for transforming auditory signals from

head-centered to eye-centered coordinates. *Biological Cybernetics,* 67**,** 291-302.

GROTHE, B., PECKA, M. & MCALPINE, D. 2010. Mechanisms of sound localization in

mammals. *Physiol Rev* 90**,** 983-1012.

HAESSLY, A., SIROSH, J. & MIIKKULAINEN, R. A model of visually guided plasticity of

the auditory spatial map in the barn owl. Seventeenth Annual Meetings of the

Cognitive Science Society, 1995 Pittsburgh, PA. Erlbaum, 154-158.

KOPCO, N., LIN, I. F., SHINN-CUNNINGHAM, B. G. & GROH, J. M. 2009. Reference

Frame of the Ventriloquism Aftereffect. *Journal of Neuroscience,* 29**,** 13809-13814.

KOPCO, N., LOKSA, P., LIN, I. F., GROH, J. & SHINN-CUNNINGHAM, B. 2019.

Hemisphere-specific properties of the ventriloquism aftereffect. *J Acoust Soc Am,* 146**,**

EL177-183.

KOPCO, N., MARCINEK, L., TOMORIOVA, B. & HLADEK, L. 2015. Contextual

plasticity, top-down, and non-auditory factors in sound localization with a distractor.

Journal of the Acoustical Society of America, 137.

KORDING, K. P., BEACHAM, J. B., MA, W. J., QUARTZ, S., TENENBAUM, J. B. &

SHAMS, L. 2007. Causal Inference in Multisensory Perception. *Plos One,* 2.

LEWALD, J. & EHRENSTEIN, W. H. 1998. Auditory-visual spatial integration: A new

psychophysical approach using laser pointing to acoustic targets. *Journal of the*

Acoustical Society of America, 104**,** 1586-1597.

LINGNER, A., PECKA, M, LEIBOLD, C., GROTHE, B. 2018. A novel concept for dynamic

adjustment of auditory space. *Nature Scientific Reports,* 8.

MCALPINE, D., JIANG, D. & PALMER, A. R. 2001. A neural code for low-frequency

sound localization in mammals. *Nature Neuroscience,* 4**,** 396-401.

MICHALKA, S. W., ROSEN, M. L., KONG, L., SHINN-CUNNINGHAM, B. & SOMERS,

D. C. 2016. Auditory spatial coding flexibly recruits anterior, but not posterior,

visuotopic parietal cortex. *Cerebral Cortex,* 26**,** 1302-1308.

MULLETTE-GILLMAN, O. A., COHEN, Y. E. & GROH, J. M. 2005. Eye-centered, head-

centered, and complex coding of visual and auditory targets in the intraparietal sulcus.

Journal of Neurophysiology, 94**,** 2331-2352.

MULLETTE-GILLMAN, O. A., COHEN, Y. E. & GROH, J. M. 2009. Motor-related signals

in the intraparietal cortex encode locations in a hybrid, rather than eye-centered,

reference frame. *Cerebral Cortex,* 19**,** 1761-75.

ODEGAARD, B., ULRICH, R., CARPENTER, J. & SHAMS, L. 2019. Prior expectation of

objects in space is dependent on the direction of gaze. *Elsevier Cognition,* 182**,** 220-

226.

ODEGAARD, B., WOZNY, D. R. & SHAMS, L. 2015. Biases in Visual, Auditory, and

Audiovisual Perception of Space. *PLOS Computational Biology,* 11**,** e1004649.

OESS, T., ERNST, M. O. & NEUMANN, H. 2020. Computational principles of neural

adaptation for binaural signal integration. *PLOS Comput Biol,* 16.

POUGET, A., DENEVE, S. & DUHAMEL, J. R. 2002. A computational perspective on the

neural basis of multisensory spatial representations. *Nature Reviews Neuroscience,* 3**,**

741-747.

RADEAU, M. & BERTELSON, P. 1974. The after-effects of ventriloquism. *Quarterly*

Journal of Experimental Psychology, 26**,** 63-71.

RADEAU, M. & BERTELSON, P. 1976. The effect of a textured visual field on modality

dominance in a ventriloquism situation. *Perception and Psychophysics,* 20**,** 227-235.

RAZAVI, B., O'NEILL, W. E. & PAIGE, G. D. 2007. Auditory Spatial Perception

661   Dynamically Realigns with Changing Eye Position. *Journal of Neuroscience,* 27**,**

662   10249-10258.

663   RECANZONE, G. H. 1998. Rapidly induced auditory plasticity: The ventriloquism

664   aftereffect. *Proceedings of the National Academy of Sciences of the United States of*

665   *America,* 95**,** 869-875.

666   SALMINEN, N. H., MAY, P. J., ALKU, P. & TIITINEN, H. 2009. A population rate code of

667   auditory space in the human cortex. . *PLoS One* 4:e7600.

668   SHINN-CUNNINGHAM, B. G., KOPCO, N. & MARTIN, T. J. 2005. Localizing nearby

669   sound sources in a classroom: Binaural room impulse resonses. *Journal of the*

670   *Acoustical Society of America,* 117**,** 3100-3115.

671   SCHILLER, P. H., TRUE, S. D. & CONWAY, J. L. 1979. The effects of frontal eye field and

672   superior colliculus ablations on eye movement. *Science,* 206**,** 590-592.

673   STECKER, G. C., HARRINGTON, I. A. & MIDDLEBROOKS, J. C. 2005. Location Coding

674   by Opponent Neural Populations in the Auditory Cortex. *PLoS Biology,* 3**,** e78.

675   TABOGA, M. 2017. Normal distribution - Maximum Likelihood Estimation. *Lectures on*

676   *probability theory and mathematical statistics, Third edition. Kindle Direct*

677   *Publishing.*

678   VAN DER HEIJDEN, K., RAUSCHECKER, J. P., DE GELDER, B. & FORMISANO, E.

679   2019. Cortical mechanisms of spatial hearing. *Nature Reviews Neuroscience,* 20**,** 609-

680   623.

681   VAN OPSTAL, J. 2016. *The Auditory System and Human Sound-Localization Behavior,*

682   Elsevier.

WALLACE, M. T. & STEIN, B. E. 1994. Cross-modal synthesis in midbrain depends on

input from cortex. *Journal of Neurophysiology,* 71**,** 429-432.

WATSON, D. M., AKEROYD, M. A., ROACH, N. W. & S., W. B. 2019. Distinct

mechanisms govern recalibration to audio-visual discrepancies in remote and recent

history. *Sci Rep,* 9.

WATSON, D. M., AKEROYD, M. A., ROACH,, N. W., WEBB B. S. 2021. Multiple spatial

reference frames underpin perceptual recalibration to audio-visual discrepancies. *Plos*

One, 16.

WERNER-REISS, U., KELLY, K. A., TRAUSE, A. S. & UNDERHILL, A. M. 2003. Eye

position affects activity in primary auditory cortex of primates. *Current Biology,* 13**,**

554-562.

WOODS, T. M. & RECANZONE, G. H. 2004. Visually Induced Plasticity of Auditory

Spatial Perception in Macaques. *Current Biology,* 14**,** 1559-1564.

WOZNY, D. R., BEIERHOLM, U. & SHAMS, L. 2010. Probability Matching as a

Computational Strategy Used in Perception. *PLoS Comput Biol,* 6.

WOZNY, D. R. & SHAMS, L. 2011. Recalibration of Auditory Space following Milliseconds

of Cross-Modal Discrepancy. *Journal of Neuroscience,* 31**,** 4607-4612.

YAO, L. & PECK, C. K. 1997. Saccadic eye movements to visual and auditory targets. *Exp*

Brain Res, 115**,** 25-34.

ZATORRE, R. J., BOUFFARD, M., AHAD, P. & BELIN, P. 2002. Where is 'where' in the

human auditory cortex? *Nat Neurosci,* 5**,** 905-9.

ZIERUL, B., RODER, B., TEMPELMANN, C., BRUNS, P. & NOESSELT, T. 2017. The

role of auditory cortex in the spatial ventriloquism aftereffect. *Neuroimage,* 162**,** 257-

268.

ZWIERS, M. P., VAN OPSTAL, A. J. & PAIGE, G. D. 2003. Plasticity in human sound

localization induced by compressed spatial vision. *Nat Neurosci,* 6**,** 175-81.

ZWIERS, M. P., VERSNEL, H. & VAN OPSTAL, A. J. 2004. Involvement of monkey

inferior colliculus in spatial hearing. *J Neurosci,* 24**,** 4145-56.

Table 1: Range and increments of free parameters used in systematic search through the

parameter space during model simulations. Ten values of each parameter were considered with either

linear or quadratic spacing.

Table 2: Fitted model parameters and model performance for each simulation. AICc and MSE

were calculated on the data used in a given simulation. ΔAIC is the increase in AICc for a given model *re*

the model with the lowest AICc. The underlined model names indicate the model version with substantial

evidence of better fit to the data (i.e., rounded up ΔAIC smaller than 2).

Figure Captions

Figure 1: Experimental design and results from Kopco et al. (2009; 2019). A) Experimental

design: nine loudspeakers were evenly distributed at azimuths from -30° to 30°. Two fixation points (FP)

were located 10° below the loudspeakers at ±11.75° from the center. On training trials, audio-visual

stimuli were presented either from the central region (N. Kopco et al., 2009) or peripheral region (N.

Kopco et al., 2019), while the subject fixated the red FP. The audio-visual (AV) stimuli consisted of a

sound paired with an LED offset by -5°, 0°, or -5° (offset direction fixed within a session). On probe trials,

the sound was presented from any of the loudspeakers while the eyes fixated one of the FPs. B) AV-aligned

results: Across-subject mean biases in AV (green) and A-only trials (red and blue lines for respective FPs),

corresponding, respectively, to the ventriloquism effect and aftereffect from the runs with AV-aligned

stimuli. Black lines show the differences between the respective red and blue lines, i.e., differences between

the biases found for the two FPs. C) AV-misaligned results: green, red, blue and black lines as in the AV-

aligned results. Brown and beige lines show the expected results for the difference data if the RF is,

respectively, head-centered and eye-centered. Note: Error bars have been omitted for clarity. N=7 in both

experiments. All horizontal axes are plotted in head-centered reference frame.

Figure 2: Structure of the model in its 4 versions and illustration of its operation. A) Block

diagram of the model in which optional model components that differentiate the 4 model versions are

shown in gray. The model predicts the response bias ("Response") as a function of the "Probe stimulus

location and FP", with additional input parameters such as the training locations and the observed

ventriloquism effect at the training locations ("Ventriloquism"). The prediction is determined as a sum of

two components (square blocks). Component 1: "Auditory space representation" is adapted by AV signals

either only in HC reference frame (used in the HC model version; "HC" arrow) or in a combination of HC

and EC RFs (used in the HEC model; "HC" arrow and the optional "EC" arrow). In the dHC and dHEC

model versions, the size of the auditory space adaptation is reduced depending on the distance of the

current FP from the training FP ("FP-dependent attenuation" optional subblock). Component 2:

"Saccade-related bias" observed when saccades are used for responses. Labels (B), (C), and (D) within

blocks refer to respective panels below that illustrate the function of the blocks by showing the outputs of

the model components. Panels B-D: An illustrative simulation of the HEC model version for AV stimulus

locations and responses shown by green diamonds in panel B (i.e., both near 0). B) Saccade-related bias

predictions (red and blue lines) for the two fixation points (crosses). C) Adaptation of the auditory space

representation resulting from the saccade-related bias and AV response bias from panel B. Diamonds

represent the disparity between AV response bias and saccade-related bias for the training FP (red), and

non-training FP in HC RF (blue filled diamonds) and in EC RF (blue open diamonds). Lines represent

predictions of auditory space adaptation induced by these disparities. (In the HC model version, there

would be no open blue diamonds, and in the dHC and dHEC model versions, the blue line would be scaled

down by the value of d_f.) D) Response bias predicted by the model as a weighted combination of biases

from panels B and C. Model parameters used for the predictions of respective model components are

shown in each panel. All horizontal axes are plotted in HC RF.

Figure 3: Model evaluation on No-Shift data. Model predictions (lines) and experimental data

(symbols) for Central training experiment (A) and Peripheral training experiment (B). Middle and lower

rows: Across-subject mean biases (± standard error) and model predictions for the two FPs separately.

Upper row: Differences between the biases (± 1 standard error) for the two individual FPs and

corresponding differences between the model predictions from the middle and lower rows. C) Predictions

of the saccade-related bias model component of the HC and HEC models (dHC model performed

identically to HC; dHEC identically to HEC). D) Predictions of the auditory space adaptation component

for the central training experiment. Crosses: Fixation points.

Figure 4: Model evaluations preformed separately on the AV-misaligned data from the Central

training (A) and Peripheral training (B) experiments. The layout, color scheme and other aspects as in

Figure 3A and 3B.

Figure 5: Model evaluation performed on the combined AV-aligned and AV-misaligned data

from the Central training (A) and Peripheral training (B) experiments. The layout, color scheme and

other aspects as in Figure 4A and 4B, with the no-shift difference data added to the bottom.

Parameter	Range		Increments
	min	max	
h, w	0	2	linear
k	0.01	20	quadratic
c	0	1.5	quadratic
w_E	0	1	linear
σ_H, σ_E	1	20	linear
d_f	0	1	linear

Simulation	Model	Fitted parameter values								Performance		
		h	k	c	w	w_E	σ_H	σ_E	d_f	AICc	ΔAIC	MSE
No Shift (Figure 3)	HC	1.03	0.31	1.14	1.01	-	12.06	-	-	130.9	2.4	1.59
	HEC	1.13	0.17	0.95	1.24	0.36	12.84	2.98	-	128.5	-	1.26
	dHC	1.03	0.31	1.14	1.01	-	12.06	-	1.00	133.8	5.3	1.59
	dHEC	1.13	0.17	0.95	1.24	0.36	12.84	2.98	1.00	131.9	3.3	1.26
Central (Figure 4A)	HC	1.01	5.64	0.67	0.40	-	18.79	-	-	176.2	15.6	5.48
	HEC	0.96	5.60	0.67	0.48	0.30	18.14	5.01	-	170.2	9.6	3.86
	dHC	1.01	6.84	0.67	0.52	-	15.20	-	0.68	160.6	-	3.22
	dHEC	0.96	13.46	0.67	0.51	0.17	17.99	2.65	0.74	162.0	1.4	2.74
Peripheral (Figure 4B)	HC	0.83	3.40	1.33	0.55	-	12.43	-	-	136.3	-	1.73
	HEC	0.82	5.33	1.33	0.56	0.04	12.12	4.91	-	141.9	5.6	1.68
	dHC	0.83	3.27	1.33	0.55	-	12.43	-	1.00	139.1	2.8	1.73
	dHEC	0.82	2.85	1.33	0.56	0.04	12.12	4.91	1.00	144.2	7.9	1.68
All Data (Figure 5)	HC	0.79	0.82	1.15	0.49	-	14.21	-	-	444.7	10.5	3.25
	HEC	0.77	0.76	1.13	0.53	0.15	13.35	4.83	-	436.9	2.7	2.89
	dHC	0.79	0.91	1.17	0.54	-	13.74	-	0.85	436.4	2.2	2.95
	dHEC	0.77	0.82	1.15	0.55	0.11	13.64	4.36	0.90	434.2	-	2.76

A) Experimental design

Audio-Visual Training Trials

Central

Peripheral

Auditory-only Probe Trials

⬡ Target Speaker

LEDs:

■ Visual Adaptor

✚ Training Fixation Point (TrFP)

✚ Non-Training Fixation Point (Non-TrFP)

← Ventriloquism Shift Direction

B) AV-aligned Baseline

Bias *re.* A-component [°]

Leftward ← → Rightward

Diff. in Response Bias [°]

$Bias_{Tr.\,FP} - Bias_{Non-Tr.\,FP}$

Central   Peripheral

AV Resp
A-only Resps from
Training FP
Non-Training FP
Diff. in A-only Resps

Actual Target Location [°]

C) AV-misaligned

Diff. expected   Central   Peripheral
if Head-Ctrd. RF
if Eye-Ctrd. RF
Observed diff.

Actual Target Location [°]

A) Central **B) Peripheral**

Pos - Neg Shift; Diff between FPs

Pos - Neg Shift; Training FP

Pos - Neg Shift; Non-Training FP

Data

Fixation Position

HC Model

HEC Model

dHC Model

dHEC Model

A) Central   **B) Peripheral**

Pos - Neg Shift; Diff between FPs

Pos - Neg Shift; Training FP

Pos - Neg Shift; Non-Training FP

No-shift; Difference between FPs

Data
Fixation Position
HC Model
HEC Model
dHC Model
dHEC Model