**Title:** *Unraveling Amazon tree community assembly using Maximum Information*

*Entropy*: a quantitative analysis of tropical forest ecology.

**Authors:** Edwin Pos*,1,2 Luiz de Souza Coelho,3 Diogenes de Andrade Lima
Filho,3 Rafael P. Salomão,4,5 Iêda Leão Amaral,3 Francisca Dionízia de Almeida
Matos,3 Carolina V. Castilho,6 Oliver L. Phillips,7 Juan Ernesto Guevara,8,9 Marcelo
de Jesus Veiga Carim,10 Dairon Cárdenas López,11 William E. Magnusson,12
Florian Wittmann,13,14 Mariana Victória Irume,3 Maria Pires Martins,3 Daniel
Sabatier,15 José Renan da Silva Guimarães,10 Jean-François Molino,15 Olaf S.
Bánki,2 Maria Teresa Fernandez Piedade,16 Nigel C.A. Pitman,17 Abel Monteagudo
Mendoza,18 José Ferreira Ramos,3 Joseph E. Hawes,19 Everton José Almeida,20
Luciane Ferreira Barbosa,20 Larissa Cavalheiro,20 Márcia Cléia Vilela dos Santos,20
Bruno Garcia Luize,21 Evlyn Márcia Moraes de Leão Novo,22 Percy Núñez
Vargas,23 Thiago Sanna Freire Silva,24 Eduardo Martins Venticinque,25 Angelo
Gilberto Manzatto,26 Neidiane Farias Costa Reis,27 John Terborgh,28,29 Katia
Regina Casula,27 Euridice N. Honorio Coronado,30,7 Juan Carlos Montero,31,3
Beatriz S. Marimon,32 Ben Hur Marimon-Junior,32 Ted R. Feldpausch,33,7 Alvaro
Duque,34 Chris Baraloto,35 Nicolás Castaño Arboleda,11 Julien Engel,15,35 Pascal
Petronelli,36 Charles Eugene Zartman,3 Timothy J. Killeen,37 Rodolfo Vasquez,18
Bonifacio Mostacedo,38 Rafael L. Assis,39 Jochen Schöngart,16 Hernán
Castellanos,40 Marcelo Brilhante de Medeiros,41 Marcelo Fragomeni Simon,41 Ana
Andrade,42 José Luís Camargo,42 Layon O. Demarchi,16 William F. Laurance,29
Susan G.W. Laurance,29 Emanuelle de Sousa Farias,43,44 Maria Aparecida Lopes,45
José Leonardo Lima Magalhães,46,47 Henrique Eduardo Mendonça Nascimento,3
Helder Lima de Queiroz,48 Gerardo A. Aymard C.,49 Roel Brienen,7 Juan David
Cardenas Revilla,3 Flávia R.C. Costa,3 Adriano Quaresma,16 Ima Célia Guimarães
Vieira,5 Bruno Barçante Ladvocat Cintra,50 Pablo R. Stevenson,51 Yuri Oliveira
Feitosa,52 Joost F. Duivenvoorden,53 Hugo F. Mogollón,54 Leandro Valle Ferreira,5
James A. Comiskey,55,56 Freddie Draper,57,35 José Julio de Toledo,58 Gabriel
Damasco,59 Nállarett Dávila,60 Roosevelt García-Villacorta,61,62 Aline Lopes,16,63
Alberto Vicentini,12 Janaína Costa Noronha,64 Flávia Rodrigues Barbosa,64
Rainiellen de Sá Carpanedo,64 Thaise Emilio,65,12 Carolina Levis,66,67 Domingos
de Jesus Rodrigues,64 Juliana Schietti,3 Priscila Souza,3 Alfonso Alonso,56
Francisco Dallmeier,56 Vitor H.F. Gomes,68,69 Jon Lloyd,70 David Neill,71 Daniel
Praia Portela de Aguiar,16 Alejandro Araujo-Murakami,72 Luzmila Arroyo,72
Fernanda Antunes Carvalho,12,73 Fernanda Coelho de Souza,12,7 Dário Dantas do
Amaral,5 Kenneth J. Feeley,74,75 Rogerio Gribel,3 Marcelo Petratti Pansonato,3,76
Jos Barlow,77 Erika Berenguer,78,77 Joice Ferreira,47 Paul V.A. Fine,59 Marcelino
Carneiro Guedes,79 Eliana M. Jimenez,80 Juan Carlos Licona,31 Maria Cristina
Peñuela Mora,81 Carlos A. Peres,82 Boris Eduardo Villa Zegarra,83 Carlos Cerón,84
Terry W. Henkel,85 Paul Maas,2 Marcos Silveira,86 Juliana Stropp,87 Raquel
Thomas-Caesar,88 Tim R. Baker,7 Doug Daly,89 Kyle G. Dexter,90,91 John Ethan
Householder,13 Isau Huamantupa-Chuquimaco,23 Toby Pennington,33,91 Marcos
Ríos Paredes,92 Alfredo Fuentes,93,94 José Luis Marcelo Pena,95 Miles R.
Silman,96 J. Sebastián Tello,94 Jerome Chave,97 Fernando Cornejo Valverde,98
Anthony Di Fiore,99 Renato Richard Hilário,58 Juan Fernando Phillips,100 Gonzalo
Rivas-Torres,101,102 Tinde R. van Andel,2,103 Patricio von Hildebrand,104
Edelcilio Marques Barbosa,3 Luiz Carlos de Matos Bonates,3 Hilda Paulette Dávila

Doza,92 Émile Fonty,105,15 Ricardo Zárate Gómez,106 Therany Gonzales,107
George Pepe Gallardo Gonzales,92 Jean-Louis Guillaumet†,108 Bruce Hoffman,109
André Braga Junqueira,110 Yadvinder Malhi,111 Ires Paula de Andrade Miranda,3
Linder Felipe Mozombite Pinto,92 Adriana Prieto,112 Agustín Rudas,112 Ademir R.
Ruschel,47 Natalino Silva,113 César I.A. Vela,114 Vincent Antoine Vos,115 Egleé L.
Zent,116 Stanford Zent,116 Bianca Weiss Albuquerque,16 Angela Cano,51,117
Diego F. Correa,51,118 Janaina Barbosa Pedrosa Costa,79 Bernardo Monteiro
Flores,119 Milena Holmgren,120 Marcelo Trindade Nascimento,121 Alexandre A.
Oliveira,76 Hirma Ramirez-Angulo,122 Maira Rocha,16 Veridiana Vizoni
Scudeller,123 Rodrigo Sierra,124 Milton Tirado,124 Maria Natalia Umaña,125
Geertje van der Heijden,126 Emilio Vilanova Torre,122,127 Corine Vriesendorp,17
Ophelia Wang,128 Kenneth R. Young,129 Manuel Augusto Ahuite Reategui,130
Cláudia Baider,131,76 Henrik Balslev,132 Sasha Cárdenas,51 Luisa Fernanda
Casas,51 William Farfan-Rios,23,133,94 Cid Ferreira,3 Reynaldo Linares-
Palomino,56 Casimiro Mendoza,134,135 Italo Mesones,59 Armando Torres-
Lezama,122 Ligia Estela Urrego Giraldo,34 Daniel Villarroel,72 Roderick Zagt,136
Miguel N. Alexiades,137 Karina Garcia-Cabrera,96 Lionel Hernandez,40 William
Milliken,65 Walter Palacios Cuenca,138 Susamar Pansini,27 Daniela Pauletto,139
Freddy Ramirez Arevalo,140 Adeilza Felipe Sampaio,27 Elvis H. Valderrama
Sandoval,141,140 Luis Valenzuela Gamarra,18 Gerhard Boenisch,142 Jens
Kattge,143 Nathan Kraft,144 Aurora Levesley,7 Karina Melgaço,7 Georgia
Pickavance,7 Lourens Poorter,67 Hans ter Steege,2
71
72    * Corresponding author: e.t.pos@uu.nl
73    † Deceased 01-2018
74
**Affiliations**
1 Quantitative Biodiversity Dynamics, Ecology & Biodiversity & Utrecht University
Botanic Gardens, Utrecht University, Padualaan 8, Utrecht, 3584 CH, The
Netherlands
2Naturalis Biodiversity Center, PO Box 9517, Leiden, 2300 RA, The Netherlands
3Coordenação de Biodiversidade, Instituto Nacional de Pesquisas da Amazônia -
INPA, Av. André Araújo, 2936, Petrópolis, Manaus, AM, 69067-375, Brazil
4Programa Professor Visitante Nacional Sênior na Amazônia - CAPES, Universidade
Federal Rural da Amazônia, Av. Perimetral, s/n, Belém, PA, Brazil
5Coordenação de Botânica, Museu Paraense Emílio Goeldi, Av. Magalhães Barata
376, C.P. 399, Belém, PA, 66040-170, Brazil
6EMBRAPA – Centro de Pesquisa Agroflorestal de Roraima, BR 174, km 8 –
Distrito Industrial, Boa Vista, RR, 69301-970, Brazil
7School of Geography, University of Leeds, Woodhouse Lane, Leeds, LS2 9JT, UK
8Grupo de Investigación en Biodiversidad, Medio Ambiente y Salud-BIOMAS,
Universidad de las Américas, Campus Queri, Quito, Ecuador
9Keller Science Action Center, The Field Museum, 1400 S. Lake Shore Drive,
Chicago, IL, 60605-2496, USA
10Departamento de Botânica, Instituto de Pesquisas Científicas e Tecnológicas do
Amapá - IEPA, Rodovia JK, Km 10, Campus do IEPA da Fazendinha, Amapá,
68901-025, Brazil
11Herbario Amazónico Colombiano, Instituto SINCHI, Calle 20 No 5-44, Bogotá,
DC, Colombia

12Coordenação de Pesquisas em Ecologia, Instituto Nacional de Pesquisas da
Amazônia - INPA, Av. André Araújo, 2936, Petrópolis, Manaus, AM, 69067-375,
Brazil
13Dep. of Wetland Ecology, Institute of Geography and Geoecology, Karlsruhe
Institute of Technology - KIT, Josefstr.1, Rastatt, D-76437, Germany
14Biogeochemistry, Max Planck Institute for Chemistry, Hahn-Meitner Weg 1,
Mainz, 55128, Germany
15AMAP, IRD, Cirad, CNRS, INRA, Université de Montpellier, Montpellier, F-
34398, France
16Coordenação de Dinâmica Ambiental, Instituto Nacional de Pesquisas da
Amazônia - INPA, Av. André Araújo, 2936, Petrópolis, Manaus, AM, 69067-375,
Brazil
17Science and Education, The Field Museum, 1400 S. Lake Shore Drive, Chicago,
IL, 60605-2496, USA
18Jardín Botánico de Missouri, Oxapampa, Pasco, Peru
19Applied Ecology Research Group, School of Life Sciences, Anglia Ruskin
University, East Road, Cambridge, CB1 1PT, UK
20ICNHS, Universidade Federal de Mato Grosso, Av. Alexandre Ferronato, 1200,
Sinop, MT, 78557-267, Brazil
21Departamento de Ecologia, Universidade Estadual Paulista - UNESP – Instituto de
Biociências – IB, Av. 24 A, 1515, Bela Vista, Rio Claro, SP, 13506-900, Brazil
22Divisao de Sensoriamento Remoto – DSR, Instituto Nacional de Pesquisas
Espaciais – INPE, Av. dos Astronautas, 1758, Jardim da Granja, São José dos
Campos, SP, 12227-010, Brazil
23Herbario Vargas, Universidad Nacional de San Antonio Abad del Cusco, Avenida
de la Cultura, Nro 733, Cusco, Cuzco, Peru
24Biological and Environmental Sciences, University of Stirling, Stirling, FK9 4LA,
UK
25Centro de Biociências, Departamento de Ecologia, Universidade Federal do Rio
Grande do Norte, Av. Senador Salgado Filho, 3000 , Natal, RN, 59072-970, Brazil
26Departamento de Biologia, Universidade Federal de Rondônia, Rodovia BR 364 s/n
Km 9,5 - Sentido Acre, Unir, Porto Velho, RO, 76.824-027, Brazil
27Programa de Pós- Graduação em Biodiversidade e Biotecnologia PPG- Bionorte,
Universidade Federal de Rondônia, Campus Porto Velho Km 9,5 bairro Rural, Porto
Velho, RO, 76.824-027, Brazil
28Department of Biology and Florida Museum of Natural History, University of
Florida, Gainesville, FL, 32611, USA
29Centre for Tropical Environmental and Sustainability Science and College of
Science and Engineering, James Cook University, Cairns, Queensland, 4870,
Australia
30Instituto de Investigaciones de la Amazonía Peruana (IIAP), Av. A. Quiñones km
2,5, Iquitos, Loreto, 784, Peru
31Instituto Boliviano de Investigacion Forestal, Av. 6 de agosto #28, Km. 14, Doble
via La Guardia, Casilla 6204, Santa Cruz, Santa Cruz, Bolivia
32Programa de Pós-Graduação em Ecologia e Conservação, Universidade do Estado
de Mato Grosso, Nova Xavantina, MT, Brazil
33Geography, College of Life and Environmental Sciences, University of Exeter,
Rennes Drive, Exeter, EX4 4RJ, UK
34Departamento de Ciencias Forestales, Universidad Nacional de Colombia, Calle 64
x Cra 65, Medellín, Antioquia, 1027, Colombia

35International Center for Tropical Botany (ICTB) Department of Biological
Sciences, Florida International University, 11200 SW 8th Street, OE 243, Miami, FL,
33199, USA
36Cirad UMR Ecofog, AgrosParisTech,CNRS,INRA,Univ Guyane, Campus
agronomique, Kourou Cedex, 97379, France
37Agteca-Amazonica, Santa Cruz, Bolivia
38Facultad de Ciencias Agrícolas, Universidad Autónoma Gabriel René Moreno,
Santa Cruz, Santa Cruz, Bolivia
39Natural History Museum, University of Oslo, Postboks 1172, Oslo, 0318, Norway
40Centro de Investigaciones Ecológicas de Guayana, Universidad Nacional
Experimental de Guayana, Calle Chile, urbaniz Chilemex, Puerto Ordaz, Bolivar,
Venezuela
41Prédio da Botânica e Ecologia, Embrapa Recursos Genéticos e Biotecnologia,
Parque Estação Biológica, Av. W5 Norte, Brasilia, DF, 70770-917, Brazil
42Projeto Dinâmica Biológica de Fragmentos Florestais, Instituto Nacional de
Pesquisas da Amazônia - INPA, Av. André Araújo, 2936, Petrópolis, Manaus, AM,
69067-375, Brazil
43Laboratório de Ecologia de Doenças Transmissíveis da Amazônia (EDTA),
Instituto Leônidas e Maria Deane, Fiocruz, Rua Terezina, 476, Adrianópolis, Manaus,
AM, 69060-001, Brazil
44Programa de Pós-graduação em Biodiversidade e Saúde, Instituto Oswaldo Cruz -
IOC/FIOCRUZ, Pav. Arthur Neiva – Térreo, Av. Brasil, 4365 – Manguinhos, Rio de
Janeiro, RJ, 21040-360, Brazil
45Instituto de Ciências Biológicas, Universidade Federal do Pará, Av. Augusto
Corrêa 01, Belém, PA, 66075-110, Brazil
46Programa de Pós-Graduação em Ecologia, Universidade Federal do Pará, Av.
Augusto Corrêa 01, Belém, PA, 66075-110, Brazil
47Embrapa Amazônia Oriental, Trav. Dr. Enéas Pinheiro s/nº, Belém, PA, 66095-
100, Brazil
48Diretoria Técnico-Científica, Instituto de Desenvolvimento Sustentável Mamirauá,
Estrada do Bexiga, 2584, Tefé, AM, 69470-000, Brazil
49Programa de Ciencias del Agro y el Mar, Herbario Universitario (PORT),
UNELLEZ-Guanare, Guanare, Portuguesa, 3350, Venezuela
50Instituto de Biociências - Dept. Botanica, Universidade de Sao Paulo - USP, Rua do
Matão 277, Cidade Universitária, São Paulo, SP, 05508-090, Brazil
51Laboratorio de Ecología de Bosques Tropicales y Primatología, Universidad de los
Andes, Carrera 1 # 18a- 10, Bogotá, DC, 111711, Colombia
52Programa de Pós-Graduação em Biologia (Botânica), Instituto Nacional de
Pesquisas da Amazônia - INPA, Av. André Araújo, 2936, Petrópolis, Manaus, AM,
69067-375, Brazil
53Institute of Biodiversity and Ecosystem Dynamics, University of Amsterdam,
Sciencepark 904, Amsterdam, 1098 XH, The Netherlands
54Endangered Species Coalition, 8530 Geren Rd., Silver Spring, MD, 20901, USA
55Inventory and Monitoring Program, National Park Service, 120 Chatham Lane,
Fredericksburg, VA, 22405, USA
56Center for Conservation and Sustainability, Smithsonian Conservation Biology
Institute, 1100 Jefferson Dr. SW, Suite 3123, Washington, DC, 20560-0705, USA
57Department of Global Ecology, Carnegie Institution for Science, 260 Panama St.,
Stanford, CA, 94305, USA

58Universidade Federal do Amapá, Ciências Ambientais, Rod. Juscelino Kubitschek
km2, Macapá, AP, 68902-280, Brazil
59Department of Integrative Biology, University of California, Berkeley, CA, 94720-
3140, USA
60Biologia Vegetal, Universidade Estadual de Campinas, Caixa Postal 6109,
Campinas, SP, 13.083-970, Brazil
61Department of Ecology and Evolutionary Biology, Cornell University, Corson Hall,
215 Tower Road, Ithaca, NY, 14850, USA
62Peruvian Center for Biodiversity and Conservation (PCBC), Iquitos, Peru
63Department of Ecology, University of Brasilia, Brasilia, DF, 70904-970, Brazil
64ICNHS, Federal University of Mato Grosso, Av. Alexandre Ferronato 1200, Setor
Industrial, Sinop, MT, 78.557-267, Brazil
65Natural Capital and Plant Health, Royal Botanic Gardens, Kew, Richmond, Surrey,
TW9 3AB, UK
66Programa de Pós-Graduação em Ecologia, Instituto Nacional de Pesquisas da
Amazônia - INPA, Av. André Araújo, 2936, Petrópolis, Manaus, AM, 69067-375,
Brazil
67Forest Ecology and Forest Management Group, Wageningen University &
Research, Droevendaalsesteeg 3, Wageningen, P.O. Box 47, 6700 AA, The
Netherlands
68Escola de Negócios Tecnologia e Inovação, Centro Universitário do Pará, Belém,
PA, Brazil
69Universidade Federal do Pará, Rua Augusto Corrêa 01, Belém, PA, 66075-110,
Brazil
70Faculty of Natural Sciences, Department of Life Sciences, Imperial College
London, Silwood Park, South Kensington Campus, London, SW7 2AZ, UK
71Ecosistemas, Biodiversidad y Conservación de Especies, Universidad Estatal
Amazónica, Km. 2 1/2 vía a Tena (Paso Lateral), Puyo, Pastaza, Ecuador
72Museo de Historia Natural Noel Kempff Mercado, Universidad Autónoma Gabriel
Rene Moreno, Avenida Irala 565 Casilla Post al 2489, Santa Cruz, Santa Cruz,
Bolivia
73Universidade Federal de Minas Gerais, Instituto de Ciências Biológicas,
Departamento de Genética, Ecologia e Evolução, Av. Antônio Carlos, 6627
Pampulha, Belo Horizonte, MG, 31270-901, Brazil
74Department of Biology, University of Miami, Coral Gables, FL, 33146, USA
75Fairchild Tropical Botanic Garden, Coral Gables, FL, 33156, USA
76Instituto de Biociências - Dept. Ecologia, Universidade de Sao Paulo - USP, Rua do
Matão, Trav. 14, no. 321, Cidade Universitária, São Paulo, SP, 05508-090, Brazil
77Lancaster Environment Centre, Lancaster University, Lancaster, Lancashire, LA1
4YQ, UK
78Environmental Change Institute, University of Oxford, Oxford, Oxfordshire, OX1
3QY, UK
79Empresa Brasileira de Pesquisa Agropecuária, Embrapa Amapá, Rod. Juscelino
Kubitschek km 5, Macapá, Amapá, 68903-419, Brazil
80Grupo de Ecología y Conservación de Fauna y Flora Silvestre, Instituto Amazónico
de Investigaciones Imani, Universidad Nacional de Colombia sede Amazonia, Leticia,
Amazonas, Colombia
81Universidad Regional Amazónica IKIAM, Km 7 via Muyuna, Tena, Napo, Ecuador
82School of Environmental Sciences, University of East Anglia, Norwich, NR4 7TJ,
UK

83Direccíon de Evaluación Forestal y de Fauna Silvestre, Av. Javier Praod Oeste 693,
Magdalena del Mar, Peru
84Escuela de Biología Herbario Alfredo Paredes, Universidad Central, Ap. Postal
17.01.2177, Quito, Pichincha, Ecuador
85Department of Biological Sciences, Humboldt State University, 1 Harpst Street,
Arcata, CA, 95521, USA
86Museu Universitário / Centro de Ciências Biológicas e da Natureza / Laboratório de
Botânica e Ecologia Vegetal, Universidade Federal do Acre, Rio Branco, AC, 69915-
559, Brazil
87Institute of Biological and Health Sciences, Federal University of Alagoas, Av.
Lourival Melo Mota, s/n, Tabuleiro do Martins, Maceio, AL, 57072-970, Brazil
88Iwokrama International Centre for Rain Forest Conservation and Development,
Georgetown, Guyana
89New York Botanical Garden, 2900 Southern Blvd, Bronx, New York, NY, 10458-
5126, USA
90School of Geosciences, University of Edinburgh, 201 Crew Building, King's
Buildings, Edinburgh, EH9 3JN, UK
91Tropical Diversity Section, Royal Botanic Garden Edinburgh, 20a Inverleith Row,
Edinburgh, Scotland, EH3 5LR, UK
92Servicios de Biodiversidad EIRL, Jr. Independencia 405, Iquitos, Loreto, 784, Peru
93Herbario Nacional de Bolivia, Universitario UMSA, Casilla 10077 Correo Central,
La Paz, La Paz, Bolivia
94Center for Conservation and Sustainable Development, Missouri Botanical Garden,
P.O. Box 299, St. Louis, MO, 63166-0299, USA
95Universidad Nacional de Jaén, Carretera Jaén San Ignacio Km 23, Jaén, Cajamarca,
06801, Peru
96Biology Department and Center for Energy, Environment and Sustainability, Wake
Forest University, 1834 Wake Forest Rd, Winston Salem, NC, 27106, USA
97Laboratoire Evolution et Diversité Biologique, CNRS and Université Paul Sabatier,
UMR 5174 EDB, Toulouse, 31000, France
98Andes to Amazon Biodiversity Program, Madre de Dios, Madre de Dios, Peru
99Department of Anthropology, University of Texas at Austin, SAC 5.150, 2201
Speedway Stop C3200, Austin, TX, 78712, USA
100Fundación Puerto Rastrojo, Cra 10 No. 24-76 Oficina 1201, Bogotá, DC,
Colombia
101Colegio de Ciencias Biológicas y Ambientales-COCIBA & Galapagos Institute
for the Arts and Sciences-GAIAS, Universidad San Francisco de Quito-USFQ, Quito,
Pichincha, Ecuador
102Department of Wildlife Ecology and Conservation, University of Florida, 110
Newins-Ziegler Hall, Gainesville, FL, 32611, USA
103Biosystematics group, Wageningen University, Droevendaalsesteeg 1,
Wageningen, 6708 PB, The Netherlands
104Fundación Estación de Biología, Cra 10 No. 24-76 Oficina 1201, Bogotá, DC,
Colombia
105Direction régionale de la Guyane, ONF, Cayenne, F-97300, French Guiana
106PROTERRA, Instituto de Investigaciones de la Amazonía Peruana (IIAP), Av. A.
Quiñones km 2,5, Iquitos, Loreto, 784, Peru
107ACEER Foundation, Jirón Cusco N° 370, Puerto Maldonado, Madre de Dios,
Peru

108Departement EV, Muséum national d'histoire naturelle de Paris, 16 rue Buffon,
Paris, 75005, France
109Amazon Conservation Team, Doekhieweg Oost #24, Paramaribo, Suriname
110Institut de Ciència i Tecnologia Ambientals, Universitat Autònoma de Barcelona,
08193 Bellaterra, Barcelona, Spain
111Environmental Change Institute, Oxford University Centre for the Environment,
Dyson Perrins Building, South Parks Road, Oxford, England, OX1 3QY, UK
112Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Apartado
7945, Bogotá, DC, Colombia
113Instituto de Ciência Agrárias, Universidade Federal Rural da Amazônia, Av.
Presidente Tancredo Neves 2501, Belém, PA, 66.077-830, Brazil
114Escuela Profesional de Ingeniería Forestal, Universidad Nacional de San Antonio
Abad del Cusco, Jirón San Martín 451, Puerto Maldonado, Madre de Dios, Peru
115Universidad Autónoma del Beni José Ballivián, Campus Universitario Final, Av.
Ejercito, Riberalta, Beni, Bolivia
116Laboratory of Human Ecology, Instituto Venezolano de Investigaciones
Científicas - IVIC, Ado 20632, Caracas, DC, 1020A, Venezuela
117Cambridge University Botanic Garden, 1 Brookside., Cambridge, CB2 1JE, UK
118School of Agriculture and Food Sciences - ARC Centre of Excellence for
Environmental Decisions CEED, The University of Queensland, St. Lucia, QLD
4072, Australia
119University of Campinas, Plant Biology Department, Rua Monteiro Lobato, 255,
Cidade Universitária Zeferino Vaz, Barão Geraldo, Campinas, São Paulo, CEP
13083-862, Brazil
120Resource Ecology Group, Wageningen University & Research,
Droevendaalsesteeg 3a, Lumen, building number 100, Wageningen, Gelderland, 6708
PB, The Netherlands
121Laboratório de Ciências Ambientais, Universidade Estadual do Norte Fluminense,
Av. Alberto Lamego 2000, Campos dos Goyatacazes, RJ, 28013-620, Brazil
122Instituto de Investigaciones para el Desarrollo Forestal (INDEFOR), Universidad
de los Andes, Conjunto Forestal, 5101, Mérida, Mérida, Venezuela
123Departamento de Biologia, Universidade Federal do Amazonas - UFAM –
Instituto de Ciências Biológicas – ICB1, Av General Rodrigo Octavio 6200, Manaus,
AM, 69080-900, Brazil
124GeoIS, El Día 369 y El Telégrafo, 3° Piso, Quito, Pichincha, Ecuador
125Department of Ecology and Evolutionary Biology, University of Michigan, Ann
Arbor, MI, 48109, USA
126University of Nottingham, University Park, Nottingham, NG7 2RD, UK
127School of Environmental and Forest Sciences, University of Washington, Seattle,
WA, 98195-2100, USA
128Environmental Science and Policy, Northern Arizona University, Flagstaff, AZ,
86011, USA
129Geography and the Environment, University of Texas at Austin, 305 E. 23rd
Street, CLA building, Austin, TX, 78712, USA
130Medio Ambiente, PLUSPRETOL, Iquitos, Loreto, Peru
131The Mauritius Herbarium, Agricultural Services, Ministry of Agro-Industry and
Food Security, Reduit, 80835, Mauritius
132Department of Bioscience, Aarhus University, Building 1540 Ny Munkegade,
Aarhus C, Aarhus, DK-8000, Denmark

133Living Earth Collaborative, Washington University in Saint Louis, St. Louis, MO,
63130, USA
134Escuela de Ciencias Forestales (ESFOR), Universidad Mayor de San Simon
(UMSS), Sacta, Cochabamba, Bolivia
135FOMABO, Manejo Forestal en las Tierras Tropicales de Bolivia, Sacta,
Cochabamba, Bolivia
136Tropenbos International, Lawickse Allee 11 PO Box 232, Wageningen, 6700 AE,
The Netherlands
137School of Anthropology and Conservation, University of Kent, Marlowe Building,
Canterbury, Kent, CT2 7NR, UK
138Herbario Nacional del Ecuador, Universidad Técnica del Norte, Quito, Pichincha,
Ecuador
139Instituto de Biodiversidade e Floresta, Universidade Federal do Oeste do Pará,
Rua Vera Paz, Campus Tapajós, Santarém, PA, 68015-110, Brazil
140Facultad de Biologia, Universidad Nacional de la Amazonia Peruana, Pevas 5ta
cdra, Iquitos, Loreto, Peru
141Department of Biology, University of Missouri, St. Louis, MO, 63121, USA
142Department of Biogeochemical Integration, Max-Planck-Institute for
Biogeochemistry, P.O. Box 10 01 64, Jena, 07701, Germany
143Functional Biogeography, Max-Planck-Institute for Biogeochemistry, P.O. Box
10 01 64, Jena, 07701, Germany
144Department of Ecology and Evolutionary Biology, UCLA, 621 Charles E. Young
Drive South, Box 951606, Los Angeles, CA, 90095, USA
368

**Abstract** In a time of rapid global change, the question of what determines patterns in species abundance distribution remains a priority for understanding the complex dynamics of ecosystems. The constrained maximization of information entropy provides a framework for the understanding of such complex systems dynamics by a quantitative analysis of important constraints via predictions using least biased probability distributions. We apply it to over two thousand hectares of Amazonian tree inventories across seven forest types and thirteen functional traits, representing major global axes of plant strategies. Results show that constraints formed by regional relative abundances of genera explain eight times more of local relative abundances than constraints based on directional selection for specific functional traits, although the latter does show clear signals of environmental dependency. These results provide a quantitative insight by inference from large-scale data using cross-disciplinary methods, furthering our understanding of ecological dynamics.

**Introduction**

Drivers of species distributions and their predictions have been a long-standing search in ecology, with approaches varying from deterministic to neutral (i.e. stochastic) and almost everything in between (e.g. near-neutral, continuum or emergent-neutral [1,2]). Most models are based on prior assumptions of processes that drive community dynamics. The Maximum Entropy Formalism (hereafter called MEF) makes no such, potentially unjustified, a-priori assumptions in generating predictions of species abundance distributions, as such it is a useful construct to infer processes driving community dynamics given the constraints imposed by prior knowledge (e.g. functional traits or summed regional abundances) [3]. Quantifying the relative importance of these distinct constraints can thus provide additional answers to understand the complexity of community dynamics (see Supporting Materials SM: boxes S1-S3). This is especially so because, although many different tests are available that link variation in taxon abundances to 1) trait variation, 2) taxon turnover between habitats or environments and 3) the distance decay of similarities between samples, none quantify the importance of these relative to each other. The MEF as applied here, however, is capable of and designed to do exactly this by decomposing variation to separate information explained by each of these aspects in a four-step model (Fig. 1 and Box S2). Its application to an unprecedented large tree inventory database on genus level taxonomy consisting of > 2,000 1-ha plots distributed over Amazonia [4] and a genus trait database of 13 key functional traits representing global axes of plant strategies [5] allows us to advance the study of Amazonian tree community dynamics from a new cross-disciplinary perspective.

**Results**

Principles from information theory [6–8] can be used in an ecological setting to predict the most likely abundance state for each taxon while simultaneously maximizing entropy based on constraints. Maximization of entropy allows quantifying the information yield for each constraint and therefor identifies which constraints reduce entropy the most. Here we specifically use Shipley's mathematical framework (CATS) for the MEF calculations, similar to earlier studies [9–11].

*Predictive power of the four-step model*

Using a uniform prior and CWM values (Community Weighted Means) as constraints accounted for 23% on average of total deviance between observed and predicted relative abundances (measured by $R^2_{KL}$ values, see Box S2 equation 5). Filtered by forest type this was 34% for podzol forests, *várzea* 25%, *igapó* 23%, swamp forests 34%, 21% and 24% for Guyana Shield and Pebas *terra firme* respectively and 20% for Brazilian Shield *terra firme* forests (see Table S1 for detailed decomposition). Using observed metacommunity relative abundances as prior regardless of functional traits accounted for on average 56% for the combined dataset with for all forest types between 51 and 55%, except for the Guyana Shield *terra firme* with 62%. The hybrid model (including both traits as constraints and the metacommunity prior) performed slightly better for the combined dataset (average 60%) with a minimum of 57% for swamp and *várzea* forests and a maximum of 66% for Guyana Shield *terra firme* forests. To compensate for spurious relationships between regional abundances and local trait constraints, regardless of selection, explanatory power was regarded relative to model bias yielding the pure trait and metacommunity effects (Box S3, Fig. 2 and Table S1). This lowered the proportion of information accounted for and yielded

average pure metacommunity effects of 40% for the overall dataset ranging between

26 and 45% for each forest type separately with pure trait effects explaining only 5%

of information for the combined dataset on average with for each forest type between

3 and 9%. Although the latter was lowered substantially, the explanatory power did

appear to be strongly dependent on forest type. The online supplementary material

provides additional results relating to the predictive power of each model as well as

the spatial gradient of the pure trait and metacommunity effects (Figs. S2-3).

*Direction and strength of selection of trait-based constraints*

Each trait showed significant differences in lambda when compared between forest

types (Fig. S1, see methods for a definition of lambda). Scatterplots of CWM trait

values versus lambda show that, in general, higher lambda values correspond with

higher CWM trait values (Figure S7). A number of functional traits associated with

low nutrient conditions (e.g. ectomycorrhiza) and life history strategies suited for

protection against herbivores (e.g. latex, resin and high leaf C content) were clearly

positively associated with abundance in nutrient poor environments (podzols),

indicated by the positive lambda values. In contrast, having fleshy fruits and high leaf

N and P content were clearly negatively associated with abundance on these soils.

Nodulation was also negatively associated with abundance on poor soils. The ability

to accumulate aluminium was positively associated with abundance on those soils

commonly associated with higher aluminium content such as *igapó* (strong positive

effects). In contrast, it was strongly negatively associated with abundance on other

soils, with negative lambda values for podzol and *várzea* forests. Traits such as SLA

or winged fruits also showed patterns dependent on forest type, although less

pronounced.

*Effect of regional metacommunity prior*

There was a remarkable similar mean 22% decrease of the information explained

purely by the metacommunity prior for each forest type (Fig. S3). For the separate

forest types, although the initial pure metacommunity effect varied, the decline

appeared remarkably similar with a mean 25% decrease in pure metacommunity

effect for podzol, 23% for *várzea*, 26% for *igapó* and 27% for swamp forests with

*terra firme* forests having a smaller decline of approximately 21%, averaged over the

three subregions. It should be noted there is an obvious risk that when sampling size is

increased, this also includes more environmental heterogeneity as samples are coming

from a variety of localities potentially leading to changing composition. If this were

the case, however, the regional prior ($q_i$ from Fig. 1 and Box S2) would also change,

as taxa might be abundant in some places but rare or absent in others. As the

metacommunity effect is the explained information that remains relative to any trait

effects (i.e. information unique to the neutral prior) and the pure trait effects are the

explained information remaining after correcting for pure metacommunity effects

(Box S3) this effect should then be accompanied by an increase in pure trait effect for

each sample. This was not observed, not even within the different forest types.

Instead, the trait effect gradually went up and then remained constant (Fig. S4).

**Discussion**

The MEF emerges from a well-founded theoretical and empirical body of ecology and evolutionary biology, regarding natural selection, migration and population dynamics. From an ecological point of view, it can be used to quantify the relative association between directional or stabilizing selection for functional traits versus the importance of relative regional abundance regardless of these traits by imposing these as constraints. Our results show that pure trait effects, on average, explained only 5% of the information when all forest types were taken together whereas the pure metacommunity effect, however, explained eight times more taken all forest types together (40%). Greater trait dissimilarity was positively associated with higher pure trait effects, indicating trait-based selection, although the assumed influence of dispersal regardless of these traits appeared to confer more information explaining tree genus composition of the Amazon rainforest. The strength and direction of selection indicated clear directional selective pressure for life history strategies of either growth or protection, depending on forest type (see supplementary online material S-A for a more detailed exploration of ecological interpretation). Including community weighted variance as reflective of potential stabilizing selection did not provide additional information. Although this could be interpreted as indicative of weak or absent stabilizing selection, it is more likely to be an artefact of many genera not being shared among localities due to the sheer geographical scale resulting in a strong mismatch between observed, predicted and uniform relative abundances resulting in a model bias higher than information yielded by including these constraints (see also box S2).

Despite showing clear patterns in environmental selection and dispersal effects, there

was a large proportion of information left unexplained (44% on average). Potentially,

local demographic stochasticity could weaken any link between functional traits

measured and regional abundances of genera. This would, however, mean that almost

half of the information contained in relative abundances are the result of random

population dynamics and are not structurally governed. Alternatively, this could be

due to functional traits reflective of processes not taken into account in this study,

such as traits reflective of interactions between trophic levels (e.g. traits influencing

specific plant-pathogen interactions). Another and at least equally likely hypothesis

for (local) unexplained information is that when scaling up, the ratio of genus richness

to total abundance decreases rapidly initially but stabilizes again as relatively non-

overlapping habitats are included in the regional abundance distributions and more

genera are included again due to the different habitats. This would result in a change

of the regional abundance distribution (i.e. the prior) to which each local community

is compared, resulting in higher local unexplained information. Further study into

these aspects could provide additional insight, though the data necessary for these

scales is lacking for Amazonian trees.

*Metacommunity importance*

Although the initial explanatory power of the metacommunity prior differed between

forest types, the decay pattern was very similar. As the effects of either traits or the

metacommunity are measured in the goodness-of-fit predictions on local relative

abundances, this implies that at small spatial scales the surrounding regional

abundances provide better estimators than functional traits, while at larger spatial

scales this shifts to the traits. The ecological translation would be that on small spatial

scales, local communities share similar environmental conditions leaving dispersal

and drift acting predominantly in changing community composition, at least for genus

level taxonomy. As the potential regional pool is increased, more and more

environmental heterogeneity and non-overlapping regions are likely to be introduced.

The more gradual decline of *terra firme* forests can then arguably be attributed to

these forests having the largest relative surface area of Amazonia (even for the

separate subregions), potentially giving these forests an almost continuous

metacommunity without gaps, resulting in a more gradual transition from

metacommunity to trait relative importance. The fact that metacommunity effects do

not change anymore after certain distances would indicate the effect of dispersal

potentially occurs over very large distances. It should be noted that as these

calculations are done at community and genus level, they do not measure single

dispersal events but rather the effect of dispersal on community composition much

deeper in time. In other words, this effect suggests more than a dispersal event every

now and then. Instead, it argues for prolonged mixing of forests on large geographical

and temporal scales, supported by recent findings demonstrating a lack of

geographical phylogenetic structure of lineages for Amazonian tree genera [12].

*Conclusion*

Using an unprecedented scale of data and applying the Maximum Entropy Formalism

from information theory we show that constraints formed by regional relative

abundances of genera explain eight times more of local relative abundances then

constraints based on directional selection for specific functional traits, although the

latter does show clear signals of environmental dependency. There is, however, still

much to be explored due to the large unexplained effects and analyses on finer

taxonomic (i.e. species level) and environmental (e.g. microhabitat) scales could

resolve these issues. The relatively large effects of the regional pool of genera over

great distances does suggest an important role for long term dispersal and mixing of

Amazonian trees, especially for the Amazonian interior.

**Methods**

*Empirical data*

The ATDN [4,13,14] consists of over 2000 tree inventory plots distributed over the

Amazon basin and the Guiana Shield, collectively referred to as Amazonia (a map of

all current plots can be found at https://atdn.myspecies.info/). Only those plots with

trees ≥ 10 cm diameter at breast height were used, leaving 2011 plots with a mean of

558 individuals per plot identified to at least genus level. Most plots used are 1 ha in

size (1414) with 492 being smaller (minimum size of .1 ha) and 105 larger (maximum

size of 80 ha). Genera have been standardized to the W3 Tropicos database [15] using

the Taxonomic Name Resolution Service (TNRS, see [16]). After filtering based on

above criteria and solving nomenclature issues, 1,121,935 individuals belonging to

over 828 genera remained. Plots were distributed over seven abiotically different

forest types: Podzol forests (PZ), *Igapó* (IG, black water flood forests), *Várzea* (VA

white water flood forests), Swamp (SW) and *Terra firme* forests (TF) with subregions

BS (Brazilian Shield), GS (Guyana Shield) and PB (Pebas) (see also [17] for details

regarding these forest types). Trait data were extracted from several sources. Wood

density was mostly derived from [18]. Traits related to leaf characteristics mostly

came from four large datasets [19–24] , including additional data from other

sources [25–27] as well as unpublished data (J. Lloyd, A.A. de Oliveira, L. Poorter,

M. van de Sande & Mazzei, M. van de Sande & L. Poorter). Data on seed mass came

from [28–30] as well as different flora's and tree guides. As this particular trait can

vary over several orders of magnitude, this was included on a log-scale [29,31].

Ectomycorrhizal aspects were derived from literature [32], the same applies to

nodulation  [33,34]. Traits involved in aluminum accumulation were based

on  [35,36] and references therein. For binary traits (yes/no), a genus was considered

having a certain trait only when >50% of the genus was positive for that specific trait.

*Functional traits and trait imputation*

Constraints were formed by Community Weighted genus Means (CWM) of functional

traits (Table 1), related to key ecological life history aspects. According to principles

of natural selection, CWM values will likely be biased towards favourable trait values

for that particular environment in the case of directional selection, as taxa with these

traits will be more abundant due to environmental selection. Previous studies included

community weighted variance (CWV) as well as indicative of potential stabilizing

selection  [11,37]. In our case, however, including CWV as constraints resulted in a

model bias that was consistently higher than information including trait or

metacommunity aspects, CWV was therefore not included as constraints in the final

analysis. As for many traits it has been shown earlier that the interspecific variability

was larger than the intraspecific variability, this allowed the use of data from different

sources to at least calculate a mean species trait value. Genus trait values were

subsequently computed as genus-level means of species values if known within the

genus and considered constant for each genus. Genus level of taxonomy was used as

the available trait database had the most information on this taxonomic level (see

Table 1). Unknown values for traits were estimated by Multiple Imputation with

Chained Equations (MICE, see [15]) by delta adjustment, subtracting a fixed amount

(delta), with sensitivity of this adjustment to the imputations of the observed versus

imputed data analysed using density plots (Fig. S8) and a linear regression model.

This procedure was done using the *mice* package [38], available on the R repository,

under predictive mean matching (*pmm* setting, 50 iterations). Results showed

imputations were stable and showed near identical patterns with each imputation

scenario (see Figs S5-6 and Table S2). After imputation, all trait values were

transformed to Community Weighted Means (CWM) of each trait ($J$) for each plot

($K$) ($\bar{T}_{JK}$) as $\bar{T}_{JK} = \sum_{i=1}^{S} t_{ij} ra_{ik}$ with *ra* the relative abundance of the $i^{th}$ genus in the

$k^{th}$ plot following earlier uses [37].

*MEF procedure predictions and ecological inference*

Figure 1 provides a schematic procedure overview, box S1 provides an overview of

important terms and Boxes S2-3 further mathematical details. Initially, a maximally

uninformative prior is specified, where $q_i$ (Box S1 equation 1) equals 1/S, indicative

of each species having equal abundances, and trait constraints are randomly permuted

multiple times (n=50) among genera to test whether inclusion of specified constraints

significantly changes derived probability distributions (see also [39]). Subsequently,

the same prior is used but now observed trait CWM values belonging to specific

genera are used as constraints (following earlier applications using simulated

communities [11]). Third, observed regional abundances are used as prior with

permutated trait constraints and finally both observed regional abundances and

observed trait CWM are used as prior and constraints. *Maxent2*, an updated version of

the *maxent* function currently in the FD library of R provided the computational

platform. Proportions of uncertainty explained by each model are given by the

Kullback-Leibler divergence $R^2_{KL}$, a generalization of the classic $R^2$ goodness of fit.

In contrast with standard linear regression models having squared goodness-of-fits

measurements, the $R^2_{KL}$ is much more related to the concept of relative entropy,

quantifying the information lost when one distribution is compared to another by

means of quantifying the statistical distance between two distributions [40]. Pure trait,

pure metacommunity, joint metacommunity-trait and unexplained effects are

calculated as proportions of total biologically relevant information (Box S1 and Box

S2). Data was rarefied to smallest sample size (swamp forests; 28) and calculations

bootstrapped 25 times. Results indicated no significant change compared to using all

data, hence the total dataset was used for all analyses.

*Strength and direction of selection*

Predictions of genus relative abundances are computed as a function of traits reflected

in the CWM values and a series of constants ($\lambda_{jk}$: the Lagrange Multipliers). Each

multiplier quantifies the association between a unit of change for a particular trait *j*

and a proportional change in predicted relative abundance $p_{ik}$ (the i$^{th}$ genus in the k$^{th}$

community) considering all other traits are constant, formally described as:

$\partial p_{ik} / \partial t_{ij} = \lambda_{jk} p_{ik} (1 - p_{ik})$ (see appendix 1 from [41]). Positive values indicate

larger trait values associated with higher abundances (positive selection), negative

values indicate the opposite (negative selection) with changes proportional to lambda.

Values approximating zero indicate no association between specific traits and relative

abundances of species. Decomposing $\lambda_{jk}$ and comparing by means of a One-Way

Analysis of Variance for each trait separately between forest types allows studying

both the strength and direction of selection in different habitats. Note that this is done

for the same constraint between forest types, as lambda values for each constraint do

not scale linearly between different constraints.

*Estimation of metacommunity size*

Iteratively increasing the regional species pool considered as prior in concentric circles of a fixed radius of 50 km allows estimating the spatial effect of metacommunity size. Due to computational limits, the number of permutations for the MEF calculations (see above) was reduced to two, shuffling the combinations of genera and traits at least once. Comparison of results from the analyses using all plots indicated small effects of a smaller perturbations (average of 5% difference for metacommunity effect between 5 and 50 permutations). The relationship between pure metacommunity effect and radius of metacommunity size was fitted using a smoothing loess regression (function *loess* and *predict;* R-package *stats* with span set at 0.1). Fits subsequently were used to predict values of metacommunity effect based on geographical distance to visualize general patterns for each forest type. Exponential decay of pure metacommunity effect was described using a self-start asymptotic regression function (*SSasymp*) of the form $y(t) \sim y_f + (y_0 - y_f)e^{-exp(log(\alpha))t}$ (*nls* from *stats,* [42]. A list of all packages used in R in addition to those preloaded can be found in the supplementary online material (SA2).

**References and notes:**

[1] R. A. Chisholm and S. W. Pacala, *Niche and Neutral Models Predict Asymptotically Equivalent Species Abundance Distributions in High-Diversity Ecological Communities.*, Proc. Natl. Acad. Sci. U. S. A. **107**, 15821 (2010).

[2] D. Tilman, *Niche Tradeoffs, Neutrality, and Community Structure: A Stochastic Theory of Resource Competition, Invasion, and Community Assembly.*, Proc. Natl. Acad. Sci. U. S. A. **101**, 10854 (2004).

[3] J. Harte, *Maximum Entropy and Ecology: A Theory of Abundance, Distribution, and Energetics* (OUP Oxford, 2011).

[4] *The Amazon Tree Diversity Network*, http://atdn.myspecies.info/.

[5] S. Díaz, J. Kattge, J. H. C. Cornelissen, I. J. Wright, S. Lavorel, S. Dray, B. Reu, M. Kleyer, C. Wirth, I. Colin Prentice, E. Garnier, G. Bönisch, M. Westoby, H. Poorter, P. B. Reich, A. T. Moles, J. Dickie, A. N. Gillison, A. E. Zanne, J. Chave, S. Joseph Wright, S. N. Sheremet Ev, H. Jactel, C. Baraloto, B. Cerabolini, S. Pierce, B. Shipley, D. Kirkup, F. Casanoves, J. S. Joswig, A. Günther, V. Falczuk, N. Rüger, M. D. Mahecha, and L. D. Gorné, *The Global Spectrum of Plant Form and Function*, Nature **529**, 167 (2016).

[6] E. T. Jaynes, *Information Theory and Statistical Mechanics I*, Phys. Rev. **106**, 620 (1957).

[7] E. T. Jaynes, *Information Theory and Statistical Mechanics. II*, Phys. Rev. **108**, 171 (1957).

[8] C. E. Shannon, *A Mathematical Theory of Communication*, Bell Syst. Tech. J. **27**, 379 (1948).

[9] B. Shipley, *Measuring and Interpreting Trait-Based Selection versus Meta-Community Effects during Local Community Assembly*, J. Veg. Sci. **25**, 55

699     (2014).

700     [10]    B. Shipley, C. E. T. Paine, and C. Baraloto, *Quantifying the Importance of*

701             *Local Niche-Based and Stochastic Processes to Tropical Tree Community*

702             *Assembly*, Ecology **93**, 760 (2012).

703     [11]    J. Loranger, F. Munoz, B. Shipley, and C. Violle, *What Makes Trait–*

704             *Abundance Relationships When Both Environmental Filtering and Stochastic*

705             *Neutral Dynamics Are at Play?*, Oikos **127**, 1735 (2018).

706     [12]    K. G. Dexter, M. Lavin, B. M. Torke, A. D. Twyford, T. A. Kursar, P. D.

707             Coley, C. Drake, R. Hollands, and R. T. Pennington, *Dispersal Assembly of*

708             *Rain Forest Tree Communities across the Amazon Basin*, Proc. Natl. Acad. Sci.

709             **114**, 2645 (2017).

710     [13]    H. ter Steege, N. C. A. Pitman, D. Sabatier, C. Baraloto, R. P. Salomão, J. E.

711             Guevara, O. L. Phillips, C. V Castilho, W. E. Magnusson, J.-F. Molino, A.

712             Monteagudo, P. Núñez Vargas, J. C. Montero, T. R. Feldpausch, E. N. H.

713             Coronado, T. J. Killeen, B. Mostacedo, R. Vasquez, R. L. Assis, J. Terborgh, F.

714             Wittmann, A. Andrade, W. F. Laurance, S. G. W. Laurance, B. S. Marimon,

715             B.-H. Marimon, I. C. Guimarães Vieira, I. L. Amaral, R. Brienen, H.

716             Castellanos, D. Cárdenas López, J. F. Duivenvoorden, H. F. Mogollón, F. D. de

717             A. Matos, N. Dávila, R. García-Villacorta, P. R. Stevenson Diaz, F. Costa, T.

718             Emilio, C. Levis, J. Schietti, P. Souza, A. Alonso, F. Dallmeier, A. J. D.

719             Montoya, M. T. Fernandez Piedade, A. Araujo-Murakami, L. Arroyo, R.

720             Gribel, P. V. A. Fine, C. A. Peres, M. Toledo, G. A. Aymard C, T. R. Baker, C.

721             Cerón, J. Engel, T. W. Henkel, P. Maas, P. Petronelli, J. Stropp, C. E. Zartman,

722             D. Daly, D. Neill, M. Silveira, M. R. Paredes, J. Chave, D. de A. Lima Filho, P.

723             M. Jørgensen, A. Fuentes, J. Schöngart, F. Cornejo Valverde, A. Di Fiore, E.

M. Jimenez, M. C. Peñuela Mora, J. F. Phillips, G. Rivas, T. R. van Andel, P.

von Hildebrand, B. Hoffman, E. L. Zent, Y. Malhi, A. Prieto, A. Rudas, A. R.

Ruschell, N. Silva, V. Vos, S. Zent, A. A. Oliveira, A. C. Schutz, T. Gonzales,

M. Trindade Nascimento, H. Ramirez-Angulo, R. Sierra, M. Tirado, M. N.

Umaña Medina, G. van der Heijden, C. I. A. Vela, E. Vilanova Torre, C.

Vriesendorp, O. Wang, K. R. Young, C. Baider, H. Balslev, C. Ferreira, I.

Mesones, A. Torres-Lezama, L. E. Urrego Giraldo, R. Zagt, M. N. Alexiades,

L. Hernandez, I. Huamantupa-Chuquimaco, W. Milliken, W. Palacios Cuenca,

D. Pauletto, E. Valderrama Sandoval, L. Valenzuela Gamarra, K. G. Dexter, K.

Feeley, G. Lopez-Gonzalez, M. R. Silman, S. P. Hubbell, F. He, R. Condit, L.

Borda-de-Agua, J. Kellner, H. Ter Steege, G. A. Black, T. H. Dobzhansky, C.

Pavan, J. M. Pires, T. Dobzhansky, G. A. Black, M. J. G. Hopkins, M. J.

Costello, R. M. May, N. E. Stork, P. Haripersaud, H. ter Steege, J.-J. de

Granville, H. Chevillotte, M. Hoff, D. P. Bebber, M. A. Carine, J. R. Wood, A.

H. Wortley, D. J. Harris, G. T. Prance, G. Davidse, J. Paige, T. D. Pennington,

N. K. Robson, R. W. Scotland, B. J. McGill, R. S. Etienne, J. S. Gray, D.

Alonso, M. J. Anderson, H. K. Benecha, M. Dornelas, B. J. Enquist, J. L.

Green, F. He, A. H. Hurlbert, A. E. Magurran, P. A. Marquet, B. A. Maurer, A.

Ostling, C. U. Soykan, K. I. Ugland, E. P. White, R. J. Warren, D. K. Skelly,

O. J. Schmitz, M. A. Bradford, N. C. A. Pitman, J. W. Terborgh, M. R. Silman,

P. N. V, D. A. Neill, C. E. Cerón, W. A. Palacios, M. Aulestia, N. C. A.

Pitman, M. R. Silman, J. W. Terborgh, F. D. Lozano, M. W. Schwartz, M. W.

Schwartz, D. Simberloff, J. E. Richardson, R. T. Pennington, T. D. Pennington,

P. M. Hollingsworth, T. L. Couvreur, F. Forest, W. J. Baker, S. Cavers, C. W.

Dick, D. H. Janzen, S. A. Mangan, S. A. Schnitzer, E. A. Herre, K. M. Mack,

M. C. Valencia, E. I. Sanchez, J. D. Bever, W. Balée, D. G. Campbell, C.

Levis, P. F. de Souza, J. Schietti, T. Emilio, J. L. P. V. Pinto, C. R. Clement, F.

R. C. Costa, D. A. Posey, E. Montoya, V. Rull, N. D. Stansell, M. B. Abbott, S.

Nogué, B. W. Bird, W. A. Díaz, C. Gomez-Navarro, C. Jaramillo, F. Herrera,

S. L. Wing, R. Callejas, C. H. McMichael, D. R. Piperno, M. B. Bush, M. R.

Silman, A. R. Zimmerman, M. F. Raczka, L. C. Lobato, H. ter Steege, P. P.

Haripersaud, O. S. Bánki, F. Schieving, S. J. Phillips, R. P. Anderson, R. E.

Schapire, S. J. Phillips, M. Dudik, C. A. Quesada, J. Lloyd, L. O. Anderson, N.

M. Fyllas, M. Schwarz, C. I. Czimczik, B. Rollet, H. ter Steege, N. C. Pitman,

O. L. Phillips, J. Chave, D. Sabatier, A. Duque, J. F. Molino, M. F. Prévost, R.

Spichiger, H. Castellanos, P. von Hildebrand, R. Vásquez, P. M. Fearnside, D.

Mouillot, D. R. Bellwood, C. Baraloto, J. Chave, R. Galzin, M. Harmelin-

Vivien, M. Kulbicki, S. Lavergne, S. Lavorel, N. Mouquet, C. E. Paine, J.

Renaud, W. Thuiller, G. Lopez-Gonzalez, S. L. Lewis, M. Burkitt, O. L.

Phillips, P. J. M. Maas, L. Y. T. Westra, H. Rainer, A. Q. Lobão, R. H. J.

Erkens, P. V. A. Fine, D. C. Daly, G. V. Muñoz, I. Mesones, K. M. Cameron,

K. J. Feeley, M. R. Silman, M. Dufrene, P. Legendre, J. Bunge, M. Fitzpatrick,

J. Bunge, L. Woodard, D. Böhning, J. A. Foster, S. Connolly, H. K. Allen, J.-P.

Z. Wang, B. G. Lindsay, A. Chao, R. K. Colwell, C.-W. Lin, N. J. Gotelli, C.

X. Mao, R. K. Colwell, U. Brose, N. D. Martinez, R. J. Williams, I. J. Good, A.

Chao, S.-M. Lee, I. Rocchetti, J. Bunge, D. Bohning, R. A. Fisher, A. S.

Corbet, C. B. Williams, F. W. Preston, J.-P. Wang, P. V. A. Fine, and R. H.

Ree, *Hyperdominance in the Amazonian Tree Flora*, Science **342**, 1243092

(2013).

[14]  H. ter Steege, P. I. Prado, R. A. F. d. Lima, E. Pos, L. de Souza Coelho, D. de Andrade Lima Filho, R. P. Salomão, I. L. Amaral, F. D. de Almeida Matos, C. V. Castilho, O. L. Phillips, J. E. Guevara, M. de Jesus Veiga Carim, D. Cárdenas López, W. E. Magnusson, F. Wittmann, M. P. Martins, D. Sabatier, M. V. Irume, J. R. da Silva Guimarães, J. F. Molino, O. S. Bánki, M. T. F. Piedade, N. C. A. Pitman, J. F. Ramos, A. Monteagudo Mendoza, E. M. Venticinque, B. G. Luize, P. Núñez Vargas, T. S. F. Silva, E. M. M. de Leão Novo, N. F. C. Reis, J. Terborgh, A. G. Manzatto, K. R. Casula, E. N. Honorio Coronado, J. C. Montero, A. Duque, F. R. C. Costa, N. Castaño Arboleda, J. Schöngart, C. E. Zartman, T. J. Killeen, B. S. Marimon, B. H. Marimon-Junior, R. Vasquez, B. Mostacedo, L. O. Demarchi, T. R. Feldpausch, J. Engel, P. Petronelli, C. Baraloto, R. L. Assis, H. Castellanos, M. F. Simon, M. B. de Medeiros, A. Quaresma, S. G. W. Laurance, L. M. Rincón, A. Andrade, T. R. Sousa, J. L. Camargo, J. Schietti, W. F. Laurance, H. L. de Queiroz, H. E. M. Nascimento, M. A. Lopes, E. de Sousa Farias, J. L. L. Magalhães, R. Brienen, G. A. Aymard C, J. D. C. Revilla, I. C. G. Vieira, B. B. L. Cintra, P. R. Stevenson, Y. O. Feitosa, J. F. Duivenvoorden, H. F. Mogollón, A. Araujo-Murakami, L. V. Ferreira, J. R. Lozada, J. A. Comiskey, J. J. de Toledo, G. Damasco, N. Dávila, A. Lopes, R. García-Villacorta, F. Draper, A. Vicentini, F. Cornejo Valverde, J. Lloyd, V. H. F. Gomes, D. Neill, A. Alonso, F. Dallmeier, F. C. de Souza, R. Gribel, L. Arroyo, F. A. Carvalho, D. P. P. de Aguiar, D. D. do Amaral, M. P. Pansonato, K. J. Feeley, E. Berenguer, P. V. A. Fine, M. C. Guedes, J. Barlow, J. Ferreira, B. Villa, M. C. Peñuela Mora, E. M. Jimenez, J. C. Licona, C. Cerón, R. Thomas, P. Maas, M. Silveira, T. W.

Henkel, J. Stropp, M. R. Paredes, K. G. Dexter, D. Daly, T. R. Baker, I.

Huamantupa-Chuquimaco, W. Milliken, T. Pennington, J. S. Tello, J. L. M.

Pena, C. A. Peres, B. Klitgaard, A. Fuentes, M. R. Silman, A. Di Fiore, P. von

Hildebrand, J. Chave, T. R. van Andel, R. R. Hilário, J. F. Phillips, G. Rivas-

Torres, J. C. Noronha, A. Prieto, T. Gonzales, R. de Sá Carpanedo, G. P. G.

Gonzales, R. Z. Gómez, D. de Jesus Rodrigues, E. L. Zent, A. R. Ruschel, V.

A. Vos, É. Fonty, A. B. Junqueira, H. P. D. Doza, B. Hoffman, S. Zent, E. M.

Barbosa, Y. Malhi, L. C. de Matos Bonates, I. P. de Andrade Miranda, N.

Silva, F. R. Barbosa, C. I. A. Vela, L. F. M. Pinto, A. Rudas, B. W.

Albuquerque, M. N. Umaña, Y. A. Carrero Márquez, G. van der Heijden, K. R.

Young, M. Tirado, D. F. Correa, R. Sierra, J. B. P. Costa, M. Rocha, E.

Vilanova Torre, O. Wang, A. A. Oliveira, M. Kalamandeen, C. Vriesendorp, H.

Ramirez-Angulo, M. Holmgren, M. T. Nascimento, D. Galbraith, B. M. Flores,

V. V. Scudeller, A. Cano, M. A. Ahuite Reategui, I. Mesones, C. Baider, C.

Mendoza, R. Zagt, L. E. Urrego Giraldo, C. Ferreira, D. Villarroel, R. Linares-

Palomino, W. Farfan-Rios, W. Farfan-Rios, L. F. Casas, S. Cárdenas, H.

Balslev, A. Torres-Lezama, M. N. Alexiades, K. Garcia-Cabrera, L. Valenzuela

Gamarra, E. H. Valderrama Sandoval, F. Ramirez Arevalo, L. Hernandez, A. F.

Sampaio, S. Pansini, W. Palacios Cuenca, E. A. de Oliveira, D. Pauletto, A.

Levesley, K. Melgaço, and G. Pickavance, *Biased-Corrected Richness*

*Estimates for the Amazonian Tree Flora*, Sci. Rep. **10**, 1 (2020).

[15]  *Tropicos, Tropicos.Org. Missouri Botanical Garden. Accessed 2018, Version*

*Number Unknown Due to Lack of Release-History.*, (unpublished).

[16] B. Boyle, N. Hopkins, Z. Lu, J. A. Raygoza Garay, D. Mozzherin, T. Rees, N.

Matasci, M. L. Narro, W. H. Piel, S. J. McKay, S. Lowry, C. Freeland, R. K.

Peet, and B. J. Enquist, *The Taxonomic Name Resolution Service: An Online*

*Tool for Automated Standardization of Plant Names.*, BMC Bioinformatics **14**,

16 (2013).

[17] T. Haugaasen and C. Peres, *Floristic, Edaphic and Structural Characteristics*

*of Flooded and Unflooded Forests in the Lower Rio Purús Region of Central*

*Amazonia, Brazil*, Acta Amaz. **36**, 25 (2006).

[18] J. Chave, D. Coomes, and S. Jansen, *Towards a Worldwide Wood Economics*

*Spectrum*, Ecol. Lett. 351 (2009).

[19] N. J. B. Kraft, R. Valencia, and D. D. Ackerly, *Functional Traits and Niche-*

*Based Tree Community Assembly in an Amazonian Forest.*, Science **322**, 580

(2008).

[20] N. M. Fyllas, S. Patino, T. R. Baker, G. Bielefeld Nardoto, L. A. Martinelli, C.

A. Quesada, R. Paiva, M. Schwarz, V. Horna, L. M. Mercado, A. Santos, L.

Arroyo, E. M. Jiměnez, F. J. Luizao, D. A. Neill, N. Silva, A. Prieto, A. Rudas,

M. Silviera, I. C. G. Vieira, G. Lopez-Gonzalez, Y. Malhi, O. L. Phillips, and J.

Lloyd, *Basin-Wide Variations in Foliar Properties of Amazonian Forest:*

*Phylogeny, Soils and Climate*, Biogeosciences **6**, 2677 (2009).

[21] C. Baraloto, C. E. T. Paine, L. Poorter, J. Beauchene, D. Bonal, A. M.

Domenach, B. Hérault, S. Patiño, J. C. Roggy, and J. Chave, *Decoupled Leaf*

*and Stem Economics in Rain Forest Trees*, Ecol. Lett. **13**, 1338 (2010).

[22] C. Fortunel, P. V. A. Fine, and C. Baraloto, *Leaf, Stem and Root Tissue*

*Strategies across 758 Neotropical Tree Species*, Funct. Ecol. **26**, 1153 (2012).

[23] C. E. T. Paine, C. Baraloto, J. Chave, and B. Hérault, *Functional Traits of*

*Individual Trees Reveal Ecological Constraints on Community Assembly in*

*Tropical Rain Forests*, Oikos **120**, 720 (2011).

[24] S. Patiño, N. M. Fyllas, T. R. Baker, R. Paiva, C. A. Quesada, A. J. B. Santos,

M. Schwarz, H. ter Steege, O. L. Phillips, and J. Lloyd., *Coordination of*

*Physiological and Structural Traits in Amazon Forest Trees.*, Biogeosciences

**9**, 775 (2012).

[25] F. M. J. Ohler, *Phytomass and Mineral Content in Untouched Forest*, (1980).

[26] J. Thompson, J. Proctor, V. Viana, W. Milliken, J. A. Ratter, and D. A. Scott,

*Ecological Studies on a Lowland Evergreen Rain Forest on Maracá Island,*

*Roraima, Brazil. I. Physical Environment, Forest Structure and Leaf*

*Chemistry*, J. Ecol. 689 (1992).

[27] T. L. Pons, K. Perreijn, C. Van Kessel, and M. J. A. Werger, *Symbiotic*

*Nitrogen Fixation in a Tropical Rainforest: 15N Natural Abundance*

*Measurements Supported by Experimental Isotopic Enrichment*, New Phytol.

**173**, 154 (2006).

[28] S. Foster and C. H. Janson, *The Relationship between Seed Size and*

*Establishment Conditions in Tropical Woody Plants*, Ecology **66**, 773 (1985).

[29] D. S. Hammond and V. K. Brown, *Hammond and Brown 1995 Seed Size.*

*Ecology.Pdf*, Ecology **76**, 2544 (1995).

[30] Royal Botanic Gardens Kew, *Seed Information Database (SID)*, (unpublished).

[31] H. ter Steege and D. Hammond, *Character Convergence, Diversity, and*

*Disturbance in Tropical Rain Forest in Guyana*, Ecology **82**, 3197 (2001).

[32]   L. Tedersoo and M. C. Brundrett, *Evolution of Ectomycorrhizal Symbiosis in*

*Plants.*, Biogeogr. Mycorrhizal Symbiosis. Springer Int. Publ. Basel 407

(2017).

[33]   J. I. Sprent, *Nodulation in Legumes*, (unpublished).

[34]   P. S. Soltis, D. E. Soltis, and M. W. Chase, *Angiosperm Phylogeny Inferred*

*from Multiple Genes as a Tool for Comparative Biology.*, Nature **402**, 402

(1999).

[35]   S. Jansen, T. Watanabe, S. Dessein, E. Smets, and E. Robbrecht, *A*

*Comparative Study of Metal Levels in Leaves of Some Al-Accumulating*

*Rubiaceae*, Ann. Bot. **91**, 657 (2003).

[36]   S. Jansen, T. Watanabe, and E. Smets., *Aluminium Accumulation in Leaves of*

*127 Species in Melastomataceae, with Comments on the Order Myrtales*, Ann.

Bot. **90**, 53 (2002).

[37]   G. Sonnier, B. Shipley, and M. L. Navas, *Quantifying Relationships between*

*Traits and Explicitly Measured Gradients of Stress and Disturbance in Early*

*Successional Plant Communities*, J. Veg. Sci. **21**, 1014 (2010).

[38]   S. van Buuren and K. Groothuis-Oudshoorn, *Mice: Multivariate Imputation by*

*Chained Equations in R*, J. Stat. Softw. **45**, 1 (2011).

[39]   S. H. Roxburgh and K. Mokany, *On Testing Predictions of Species Relative*

*Abundance from Maximum Entropy Optimisation*, Oikos **119**, 583 (2010).

[40]   A. Colin Cameron and F. A. G. Windmeijer, *An R-Squared Measure of*

*Goodness of Fit for Some Common Nonlinear Regression Models*, J. Econom.

**77**, 329 (1997).

[41]    G. Sonnier, M. L. Navas, A. Fayolle, and B. Shipley, *Quantifying Trait*

897           *Selection Driving Community Assembly: A Test in Herbaceous Plant*

898           *Communities under Contrasted Land Use Regimes*, Oikos **121**, 1103 (2012).

899    [42]    R Core Team, *R: A Language and Environment for Statistical Computing*.

900    [43]    E. Weiher, A. van der Werf, K. Thompson, M. Roderick, E. Garnier, and O.

901           Eriksson, *Challenging Theophrastus: A Common Core List of Plant Traits for*

902           *Functional Ecology*, J. Veg. Sci. **10**, 609 (1999).

903    [44]    K. G. Raghothama, *Phosphate Acquisition*, Annu. Rev. Plant Physiol. Plant

904           Mol. Biol. **50**, 665 (1999).

905    [45]    L. Poorter, M. Van De Plassche, S. Willems, and R. G. A. Boot, *Leaf Traits*

906           *and Herbivory Rates of Tropical Tree Species Differing in Successional Status*,

907           Plant Biol. **6**, 746 (2004).

908    [46]    A. A. Agrawal, *Macroevolution of Plant Defense Strategies*, Trends Ecol. Evol.

909           **22**, 103 (2007).

910    [47]    J. I. Sprent and R. Parsons, *Nitrogen Fixation in Legume and Non-Legume*

911           *Trees*, F. Crop. Res. **65**, 183 (2000).

912    [48]    J. V. Colpaert, J. H. L. Wevers, E. Krznaric, and K. Adriaensen, *How Metal-*

913           *Tolerant Ecotypes of Ectomycorrhizal Fungi Protect Plants from Heavy Metal*

914           *Pollution*, Ann. For. Sci. **68**, 17 (2011).

915    [49]    C. Poschenrieder, B. Gunsé, I. Corrales, and J. Barceló, *A Glance into*

916           *Aluminum Toxicity and Resistance in Plants*, Sci. Total Environ. **400**, 356

917           (2008).

918    [50]    H. F. Howe and J. Smallwood, *Ecology of Seed Dispersal Ecology of Seed*

919           *Dispersal*, **13**, 201 (2013).

**Author Contributions:**

E.T. Pos and H. ter Steege designed the study. E.T.Pos performed analyses and took the lead in writing the manuscript, H. ter Steege supervised the writing and provided regular feedback both for the manuscript and the interpretation of the results. All other authors provided feedback on the manuscript and provided their data from the Amazon Tree Diversity Network or trait data. Authors E.T. Pos to L.V. Gamarra provided tree inventory data, authors G. Boenisch, J. Kattge, N. Kraft, A. Levesley, K. Melgaço, G. Pickavance, L. Poorter provided data on functional traits, C. Baraloto, J. Lloyd, A. A. Oliveira and H. ter Steege provided both tree inventory and functional trait data. **Competing interests:** Authors declare no competing interests. **Data and materials availability:** R scripts are available on the github repository of E.T. Pos (EdwinTPos). The data that support the findings of this study are available from The Amazon Tree Diversity Network (ATDN) upon reasonable request.

**Acknowledgements:**

This paper is the result of the work of hundreds of different scientists and research

institutions in the Amazon over the past 80 years. Without their hard work this

analysis would have been impossible. The 25-ha Long-Term Ecological Research

Project of Amacayacu is a collaborative project of the Instituto Amazónico de

Investigaciones Científicas Sinchi and the Universidad Nacional de Colombia Sede

Medellín, in partnership with the Unidad de Manejo Especial de Parques Naturales

Nacionales and the Center for Tropical Forest Science of the Smithsonian Tropical

Research Institute (CTFS). The Amacayacu Forest Dynamics Plot is part of the

Center for Tropical Forest Science, a global network of large-scale demographic tree

plots. We acknowledge the Director and staff of the Amacayacu National Park for

supporting and maintaining the project in this National Park. We would also like to

thank Prof. John Harte, Prof. Bill Shipley and the anonymous reviewers for their

constructive feedback during the review process.

**Funding:** HtS and RS were supported by grant 407232/2013-3 - PVE -

MEC/MCTI/CAPES/CNPq/FAPs; CB was supported by grant FAPESP 95/3058-0 -

CRS 068/96 WWF Brasil - The Body Shop; DS, JFM, JE, PP and JC benefited from

an "Investissement d'Avenir" grant managed by the Agence Nationale de la

Recherche (CEBA: ANR-10-LABX-25-01); HLQ/MAP/JLLM received financial

supported by MCT/CNPq/CT-INFRA/GEOMA #550373/2010-1 and # 457515/2012-

0, and JLLM were supported by grant CAPES/PDSE # 88881.135761/2016-01 and

CAPES/Fapespa #1530801; The Brazilian National Research Council (CNPq)

provided a productivity grant to EMV (Grant 308040/2017-1); Floristic identification

in plots in the RAINFOR forest monitoring network have been supported by the

Natural Environment Research Council (grants NE/B503384/1, NE/ D01025X/1,

NE/I02982X/1, NE/F005806/1, NE/D005590/1 and NE/I028122/1) and the Gordon

and Betty Moore Foundation; B.M.F. is funded by FAPESP grant 2016/25086-3.

BSM, BHMJ and OLP were supported by grants CNPq/CAPES/FAPS/BC-Newton

Fund #441244/2016-5 and FAPEMAT/0589267/2016; TWH was funded by National

Science Foundation grant DEB-1556338.

TABLE 1

| Functional trait | Units | Mean | SD | Est % | Associated challenge |
|---|---|---|---|---|---|
| Wood density (*WD*) | g/cm$^3$ | 0.63 | 0.17 | 30 | Longevity [43] |
| Seed Mass Class (*SMC*) | categorical (1-8) | 4.3 | 1.4 | 31 | Dispersal, Fecundity, Establishment [43] |
| Specific Leaf Area (*SLA*) | mm$^2$/mg | 15 | 5.9 | 41 | Establishment, Plasticity, Disturbance [43] |
| Leaf nitrogen content (*N*) | mg/g | 22.3 | 7.30 | 41 | Photosynthetic capacity [43] |
| Leaf phosphorus content (*P*) | mg/g | 1 | 0.77 | 50 | Limited available P for metabolism [44] |
| Leaf carbon content (*C*) | mg/g | 468 | 38.1 | 54 | Herbivore resistance (C:N) [45] |
| Latex | 1=no, 2 =yes | 1.2 | 0.43 | 46 | Herbivore resistance [46] |
| Resin | 1=no, 2 =yes | 1.1 | 0.35 | 58 | Herbivore resistance [46] |
| Root Nodules (*Nodules*) | 1=no, 2 =yes | 1.1 | 0.28 | 0 | Nitrogen fixation [47] |
| Ectomycorrhiza (*EctoMyco*) | 1=no, 2 =yes | 1.01 | 0.11 | 0 | Organic N fixation , heavy metal pollution [48] |
| Aluminum accumulation (*AlAcc*) | 1=no, 2 =yes | 1.1 | 0.21 | 3 | Heavy metal pollution [49] |
| Fleshy Fruits (*Fleshy*) | 1=no, 2 =yes | 1.6 | 0.50 | 7 | Dispersal (*specificity*) [50] |
| Winged seeds (*Wings*) | 1=no, 2 =yes | 1.2 | 0.42 | 39 | Dispersal (*limitation*) [50] |

**Table 1. Overview of used functional traits.** Mean and standard deviation (SD) are calculated after predictive mean matching (percentage of estimated values is given by Est (%)). Associated challenge indicates different aspects of life history and selective environment related to specific functional traits, sources are given in the footnote. For specific methodology of measurement protocols and calculation for each trait we refer to the original sources of the data (see main text).

FIGURE 1

**Fig. 1. Schematic depiction of the MEF procedure.** Left panel shows a genus abundances per site and a functional trait matrix per genus, bottom half outlines calculations. Middle and right panel show different scenarios of neutral and deterministic dynamics under infinite or limited migration. Figure was custom made using Adobe Illustrator (Adobe Inc., 2019. Adobe Illustrator).

FIGURE 2

**Fig. 2. Visual representation of pure trait, pure metacommunity, hybrid model and the remaining unexplained information for each separate forest type.** Abbreviations indicate different types: igapó (IG), podzol (PZ), swamp (SW), Brazilian shield terra firme (TFBS), Guiana Shield terra firme (TFGS), Pebas terra firme (TFPB) and várzea (VA). Boxplots show median value of pure effects over all samples, with lower and upper hinges corresponding to 25th and 75th percentiles. Whiskers extends from hinge to largest or smallest value no further than 1.5 * IQR from hinge. Points beyond this range are plotted individually and only positive values were plotted.

**Entities:**
*The basic unit of the MEF model which can exist in different states. Here, this constitutes a collection of genera existing at a site, hence each entity can be considered a single genus.*

**States:**
*Classification of different ways any entity can exist. Within the system, states of each entity describe their specific abundance at that site. Microstates constitute the spatial and temporal composition for the states of the entities in the system. Macrostates depict the entities of a system independent of the spatial or temporal composition, e.g., the overall relative abundance distribution but not including processes leading to this distribution (such as dispersal).*

**Traits or properties:**
*The measurable attributes for each entity, of which the values can be different for each entity. For example, genera differ in average wood density, seed mass, height etcetera. Here defined by the functional traits as described in the main text.*

**Maximally uninformative prior:**
*All the information regarding the states before specific constraints are introduced. Described as maximally uninformative as all empirical information should be introduced in the form of constraints quantify the maximal gain of information regarding the different constraints (e.g. traits or prior distribution in this case).*

**Prior distribution:**
*The expected states for the entities, here either constituted by the observed relative abundance of each entity in the summed sample (i.e. summed abundances describing the metacommunity) or by a maximally uninformed (uniform) distribution (see above). The former would be a neutral prior (expected local abundance is equal to the abundance in the larger metacommunity).*

**Community-weighted mean or variance:**
*The mean or variance of genus-level trait value over all constituent present species (for each entity) weighted by the relative abundance of each entity at a specific site.*

**Box S1. Different ingredients necessary for analyses using MEF.** Definitions of the most important terms used in the MEF analyses and throughout the main text to provide the necessary framework of understanding, adapted from [1].

BOX S2

The Maximum Entropy Formalism as applied here works based on a conceptual model called CATS (*Community Assembly by Trait Selection* [1–3]) and makes use of three inputs:

i) A **trait matrix** containing the measured functional traits of each of the S total genera in the total regional pool, these can be of either discrete or continuous form.

ii) A **vector of $n$ community weighted trait values**, estimating the average trait value over all individuals in the local community for each of the traits

iii) A **prior probability distribution** specifying the regional abundance distribution, quantifying potential contributions of the regional pool of recruits to the structure of local communities. Using these three sources of information, the model predicts relative abundances ($p_i$) in the form of Bayesian probabilities for each genus in each local community without assuming any a priori relations or processes. This is achieved by finding the vector of relative abundances maximizing entropy:

$$1)\ RE = -\sum_{i=1}^{S} p_i \ln\left(\frac{p_i}{q_i}\right)$$

with $q_i$ the regional species pool abundance of species $i$ and $RE$ (Relative Entropy) subject to the known constraints for $j$ traits and $i$ species.:

$$2)\ \bar{t}_j = \sum_{i=1}^{S} o_i t_{ij} \quad \text{and} \quad 3)\ \sum_{i=1}^{S} p_i = 1$$

The solution is a generalized exponential distribution where the λ values measure the importance of each trait when all other traits are constant:

$$4)\ p_i = \frac{q_i e^{\sum_{j=1}^{n} \lambda_j t_{ij}}}{\sum_{i=1}^{S} q_i e^{\sum_{j=1}^{T} \lambda_j t_{ij}}}$$

Note that when all λ values are zero, i.e. there is no trait based selection, $p_i = q_i$

The final step is to measure the proportion of total deviance accounted for between observed and predicted relative abundances for each of the four-step solution. These are the $R^2_{KL}$ values, a generalization of the classic $R^2$ index of maximum likelihood estimation using the Kullback-Leibler index [4,5]:

i) $\overline{R}^2_{KL}(u)$: fit of model bias, the model null hypotheses given a uniform prior (i.e. equal distribution in the regional pool of recruits).

ii) $R^2_{KL}(u, t)$: fit using again a uniform prior but including traits as constraints.

iii) $\overline{R}^2_{KL}(m)$: fit using the metacommunity prior but excluding traits as constraints

iv) $R^2_{KL}(m, t)$: fit using the metacommunity prior and including traits as constraints

The general form of the $R^2_{KL}$ divergence is calculated by:

$$5)\ R^2_{KL} = 1 - \frac{\sum_{k=1}^{c} \sum_{i=1}^{S} O_{ik} \ln\left(\frac{O_{ik}}{P_{ik}}\right)}{\sum_{j=k}^{c} \sum_{i=1}^{S} O_{ik} \ln\left(\frac{O_{ij}}{Q_{i,0}}\right)}$$

With the following parameters:

$O_{ik}$ as the observed relative abundances of the $i^{th}$ genus in the $k^{th}$ community,

$P_{ik}$ the accompanying predicted values for the specific model of the four solution step as described in the main text and,

$Q_{i,0}$ the predicted relative abundances given only the maximum uninformative prior.

Further details on the calculation of all separate $R^2_{KL}$ values and accompanying pure trait, pure metacommunity, joint information and biologically unexplained information can be found Box S3.

**Box S2. Mathematical description of the Maximum Entropy Formalism for the four-step solution.** Left panel shows necessary ingredients and formulation of the Maximum Entropy Formalism. Right side panel shows decomposition of the proportion of total deviance accounted for between observed and predicted relative abundances for each of the four-step solution, adapted from [5].

BOX S3

*The purpose of using MEF is to decompose the deviance between observed and predicted relative abundances using the four-step solution as described in the main text. The values generated are described below. The $R^2_{KL}$ value is a generalization of the classic $R^2$ index of maximum likelihood estimation using the Kullback-Leibler index for a non-linear regression including a multinomial error structure [2,4,5]. In essence, it is a way of measuring the proportion of total deviance accounted for by that specific model from one of the four steps:*

$\overline{R}^2_{KL}(u)$: fit of model bias, the model null hypotheses given a uniform prior and permuted traits
$R^2_{KL}(u, t)$: fit using a uniform prior but including observed traits as constraints
$\overline{R}^2_{KL}(m)$: fit using the metacommunity prior but excluding observed traits as constraints
$R^2_{KL}(m, t)$: fit using the metacommunity prior and including observed traits as constraints

*1) The increase in the explained deviance due to traits can be calculated either by*

$\Lambda R^2_{KL}(t|\varphi) = R^2_{KL}(u, t) - \overline{R}^2_{KL}(u)$
Increase in explained deviance due to traits beyond that due solely to model bias

or $\Lambda R^2_{KL}(t|m) = R^2_{KL}(m, t) - \overline{R}^2_{KL}(m)$
Increase in explained deviance due to traits beyond contributions made by the meta-community

*2) The increase in explained deviance due dispersal mass effects via the metacommunity can be calculated by either:*

$\Lambda R^2_{KL}(m|\varphi) = \overline{R}^2_{KL}(m) - \overline{R}^2_{KL}(u)$
Increase in explained deviance (if any) due to the metacommunity beyond that due to model bias

or $\Lambda R^2_{KL}(m|t) = R^2_{KL}(m, t) - R^2_{KL}(u, t)$
Increase in explained deviance due to the meta-community given traits, relative to the explained deviance due only to the traits: i.e. information unique to neutral prior

*3) And finally the joint information and the biologically unexplained information:*

$\Lambda R^2_{KL}(m+t) = \Lambda R^2_{KL}(m|\varphi) - \Lambda R^2_{KL}(m|t) = \Lambda R^2_{KL}(t|\varphi) - \Lambda R^2_{KL}(t|m)$
Joint information gain, or increase in explained deviance due to both the metacommunity prior and the constraints based on the traits

$1 - \Lambda R^2_{KL}(m,t)$
Biologically unexplained variation

*From these values the pure trait, pure metacommunity, joint effect and biologically unexplained variation can be calculated by the following calculations:*

Pure trait effects: $\Lambda R^2_{KL}(t|m) / (1 - \overline{R}^2_{KL}(u))$
Pure metacommunity effects: $\Lambda R^2_{KL}(m|t) / (1 - \overline{R}^2_{KL}(u))$
Joint metacommunity and trait effects: $\Lambda R^2_{KL}(m+t) / (1 - \overline{R}^2_{KL}(u))$
Unexplained effects: $1 - \Lambda R^2_{KL}(m,t) / (1 - \overline{R}^2_{KL}(u))$

**Box S3. Detailed decomposition of the four-step solution from the MEF.** Mathematical description of the decomposition based on the constraints and prior distributions (both uniform and neutral) for each of the steps from the four-step solution to measure the proportion of total deviance accounted for by each specific model from one of the four steps, adapted from [5].

**Fig. S1. Mean lambda values with standard error bars for each functional trait and compared between forest types.** Forest type abbreviations are *igapó* (IG), podzol (PZ), swamp (SW), Brazilian shield *terra firme* (TFBS), Guiana Shield *terra* firme (TFGS), Pebas *terra* firme (TFPB) and *várzea* (VA). Positive values indicate positive selection, reflective of a strong association between higher trait values and higher abundances, negative values reflect the opposite with high trait values associated with lower abundances. Differences between forest types were tested with a one-way analysis of variance with significance levels corresponding to: ns non-significant, * p < .05, ** p < .01 and *** p < .001. Abbreviations indicate functional traits: wood density (WD), seed mass class (SMC), specific leaf area (SLA), nitrogen (N), phosphorus (P) and carbon (C) leaf content. Further traits include the presence/absence of Latex, Resin, Nodules, Ectomycorrhiza (EctoMyco), the ability to accumulate aluminium (AlAcc), and the presence/absence of fleshy fruits (Fleshy) and winged seeds (Wings).

**Fig. S2. Observed relative abundances for each genus in all plots plotted against predicted relative abundance per plot (left) and summed (right) using only the traits as constraints in combination with a uniform prior (top) or the hybrid model using both traits and the metacommunity relative abundance as prior (bottom) on a log-log scale.** Top figures show predictions using only a uniform prior, left separate for all plots and right for all genera summed over all plots. Bottom figures show predictions using the regional prior, again separate for all plots and genera (left) and summed over all plots for each genus (right). Red points indicate taxa with observed relative abundances over 1e-1. Lines show the x=y prediction and $R^2$ values correspond to the Pearson's correlation coefficient.

**Fig. S3. Distance decay of pure metacommunity effect.** X-axis represents radius of

metacommunity prior; i.e. first 100 km consists of just a few plots and at 3800 km all

plots are taken into account with dashed line indicating the mean number of plots for

that distance included as metacommunity prior. Y-axis represents the pure

metacommunity effect, i.e. the increase in explained deviance due to the

metacommunity (given traits), but relative to the explained deviance due only to the

traits. It is the information unique to neutral prior taken relative to the model bias.

Solid lines indicate predictions from loess regression based on all points with different

colors indicating the forest types with abbreviations as in main text. Blue vertical lines

indicate 1000 and 2500 km boundary points. Blue shading reflects minimum and

maximum loess regression predicted values.

FIGURE S4

**Fig. S4. Distance decay of pure trait effect for each forest type separately and the overall dataset.** X-axis represents the radius of the metacommunity prior; i.e. the first 100 km consists of just a few plots and at 3800 km all plots are taken into account. Y-axis represent the pure trait effect, i.e. the increase in explained deviance due to traits beyond contributions made by the meta-community and relative to the model bias (see also Box S2). Colors indicate the different forest types with abbreviations as in main text. Lines indicate the predictions following from the loess regression based on all points. Blue vertical lines indicate the 1000 and 2500 km boundary points. Blue shading reflects maximum values for that distance of the whole dataset.

# FIGURE S5

**Fig. S5. Density plots for each functional trait showing density curves for observed (blue) versus imputed data (red; 15 iterations) under the different delta adjustment scenarios.** Only Nitrogen showed substantial deviation for the larger delta adjustment.

FIGURE S6

**Fig. S6. Scatterplot of observed vs. imputed leaf nitrogen and phosphorus content under different delta adjustment scenarios.** As Leaf Nitrogen content showed substantial deviation in the larger delta adjustment scenario it was here plotted versus Leaf Phosphorus content, which showed hardly any deviation.

**Fig. S7 Scatterplots of the CWM values against lambda values, colored according to forest type**. Titles are abbreviations for functional traits as used in the main text. Plots show in some cases these are clearly correlated (e.g. wood density, C) but for many others not (e.g. SMC and AlAcc).

FIGURE S8

**Fig. S8 Density plot of the per-plot-Pearson correlation coefficient between predicted relative abundances of each genus.** Models either used a uniform prior and functional traits (blue) or the actual observed prior and functional traits (red), results show a large increase in accuracy for the latter.

TABLE S1

**Table S1. Decomposition of results from the various maximum entropy models, combined and separated by forest type (PZ podzol, IG i*gapó*, VA *várzea*, SW Swamp, TF *terra firme* with subregions BS (Brazilian Shield), GS (Guyana Shield) and PB (Pebas formation).** Top rows indicate estimated proportions ($R^2_{KL}$) of total information reflective of variation in local relative abundance explained for by the various models. Middle rows indicate the specific information gain from any one of the used models relative to the model bias. Bottom rows show the actual effects of traits, the metacommunity and the joint information relative to the model bias.

| | Forest types | | | | | | | |
|---|---|---|---|---|---|---|---|---|
| **Explained proportions** | PZ | VA | IG | SW | TFBS | TFGS | TFPB | Combined |
| $\overline{R}^2_{KL}(u)$ *model bias fit* | 0.17 | 0.11 | 0.12 | 0.19 | 0.08 | 0.08 | 0.06 | 0.09 |
| $\overline{R}^2_{KL}(m)$ *pure neutral model fit* | 0.51 | 0.51 | 0.52 | 0.55 | 0.55 | 0.62 | 0.55 | 0.56 |
| $R^2_{KL}(u,t)$ *pure trait model fit* | 0.34 | 0.25 | 0.23 | 0.34 | 0.21 | 0.24 | 0.20 | 0.23 |
| $R^2_{KL}(m,t)$ *hybrid model fit* | 0.59 | 0.57 | 0.55 | 0.57 | 0.59 | 0.66 | 0.59 | 0.60 |
| **Increase in explained deviance** | | | | | | | | |
| $\Lambda R^2_{KL}(m|\varphi)$ *metacommunity effect beyond model bias* | 0.35 | 0.40 | 0.40 | 0.36 | 0.47 | 0.54 | 0.48 | 0.47 |
| $\Lambda R^2_{KL}(t|\varphi)$ *trait effect beyond model bias* | 0.17 | 0.14 | 0.11 | 0.15 | 0.13 | 0.16 | 0.14 | 0.14 |
| $\Lambda R^2_{KL}(t|m)$ *trait effect beyond metacommunity effect* | 0.07 | 0.05 | 0.03 | 0.02 | 0.04 | 0.04 | 0.04 | 0.04 |
| $\Lambda R^2_{KL}(m|t)$ *metacommunity effect relative to trait effects* | 0.25 | 0.32 | 0.32 | 0.23 | 0.38 | 0.42 | 0.39 | 0.37 |
| $\Lambda R^2_{KL}(m+t)$ *joint effect of metacommunity and traits* | 0.10 | 0.08 | 0.08 | 0.13 | 0.09 | 0.12 | 0.09 | 0.10 |
| $1- \Lambda R^2_{KL}(m,t)$ *unexplained effects* | 0.41 | 0.43 | 0.45 | 0.43 | 0.41 | 0.34 | 0.41 | 0.40 |
| **Biologically relevant information** | | | | | | | | |
| Pure trait effect *Information from traits, relative to bias* | 0.09 | 0.06 | 0.04 | 0.03 | 0.05 | 0.04 | 0.05 | 0.05 |
| Pure metacommunity effect *Information from metacommunity, relative to bias* | 0.30 | 0.36 | 0.37 | 0.26 | 0.42 | 0.45 | 0.41 | 0.40 |
| Joint effect *Information from joint effect, relative to bias* | 0.11 | 0.09 | 0.09 | 0.17 | 0.09 | 0.13 | 0.10 | 0.11 |
| Unexplained information *Left over information not explained, relative to bias* | 0.49 | 0.49 | 0.51 | 0.54 | 0.44 | 0.38 | 0.44 | 0.44 |

TABLE S2

**Table S2. Summary statistics overview for the linear models of the various scenarios under the delta adjustment technique as described in the main text.** Rows indicate the different delta adjustments used with the columns representing the standard summary statistics of the linear model comparing the imputed versus observed trait values. Results showed similar patterns with each imputation scenario, indicating a robust imputation procedure.

| | *Summary statistics linear model lm(imputed ~ observed)* | | | | | | |
|---|---|---|---|---|---|---|---|
| **Scenario** | Intercept | Std. error | T value | Pr. | $R^2$ | Adj. $R^2$ | Sigma |
| *Delta 0* | -.33 | .07 | -4.85 | 1.42e-06 | .32 | .33 | .63 |
| *Delta -2.5* | -.34 | .06 | -5.93 | 4.31e-09 | .37 | .37 | .58 |
| *Delta -5* | -.16 | .04 | -3.57 | 3.77e-04 | .40 | .40 | .53 |
| *Delta -7.5* | .64 | .09 | -7.33 | 5.18e-13 | .42 | .42 | .88 |
| *Delta -10* | 0.09 | .04 | 3.17 | 1.57e-03 | .47 | .47 | .48 |

966
967
968

**S-A Ecological interpretation of the MEF results**

Signals of quantitative environmental selection were found to be highest for podzol forests, whereas its counterpart in the form of the dispersal mass effect from the regional pool of genera had the second lowest value. Podzol forests, having extremely nutrient poor soils could reflect a much stronger selective environment than any of the other forest types. *Terra firme* forests, presumably reflective of a less strong selective environment in terms of resource availability, showed the opposite, with approximately half of the pure trait effect in comparison with podzol forests (even when rarefied to accommodate for different sample sizes). Traits associated with protection against herbivores such as latex [7] and high leaf carbon content showed higher values associated with greater abundance on podzol soils, whereas traits indicative of investment in growth and photosynthetic ability such as high foliar concentrations of P and N [8] showed strong negative associations on nutrient poor soils. The ability to accumulate aluminium was also strongly positively associated with relative abundance on *igapó* forests, which can potentially be richer in aluminium. Lambda values also showed strong negative lambda values for wood density in swamp and forests, fitting high tree mortality and many individuals belonging to pioneer species in especially the western Amazonian swamp forests. *Várzea* and Pebas *terra firme* forests showed a similar response. As the Pebas consists mainly of Andean sediments it has higher nutrient content, promoting lower wood density, supported by our results whereas *várzea* forests are also often flooded. There were also traits that showed no specific (strong) signal of selection on certain forest types (either positive or negative), such as latex on *igapó* and ectomycorrhiza on terra firme forests (see Fig. S1 for all lambda values). Interestingly, terra firme forests in general showed the smallest lambda values overall (positive or negative). This may be indicative of either more pronounced demographic stochasticity or ecological drift eliminating the association between traits and relative abundance. Lower effects of selection in general or more (random) variation due to the larger species pool in comparison with other forest types, however, could also be the result of mixing heterogeneous microenvironments into a single environmental class. Support for such heterogeneity within terra firme forests having influence on distribution of functional traits on valleys or plateaus has recently been found [9]. In addition, natural but also anthropogenic [10] disturbance history affects biotic community composition and can lead to changes in tree community through time, blurring relationships between traits and relative abundances. It should further be noted that, although for terra firme forests we were able to make a distinction by subregion, true within forest type heterogeneity was not taken into account. This might cause an underestimation of the deterministic effect but as of yet cannot be corrected for on this scale and is worth to be investigated in future studies. In addition, podzol forests have a smaller connected surface area and accompanying smaller number of genera in comparison with terra firme forests, adding to the calculated stronger trait effects [11,12]. When more detailed understanding and knowledge of these functional traits would be provided, this would most likely increase the explanatory power of the MEF. The fact, however, that we do not have a very specific knowledge of these interactions and specific traits is precisely the reason why the MEF can provide additional insight.

It should be noted that for species level analyses any micro environmental gradients
might prove to also show (stronger) selection at local scales [13,14], as it has been
shown that most variation in community composition, due to selection in regard to
habitat filtering and niche conservatism, is found at lower taxonomic levels, such as
between species within genera [15,16]. In contrast, theoretically it has been shown
and tested that immigration numbers are actually very robust across taxonomic
scales [17], validating our results of the metacommunity importance using genus level
taxonomy. Spatial patterns of metacommunity effects showed shallowest declines in
the centre, supporting the suggestion that high diversity of the Amazonian interior
could be explained by influx of recruits due to large (overlapping) ranges. This mid-
domain effect [18], however, would also predict lower species richness for the edges
due to lower range overlap, assuming a closed community. This is not the case, as
there is a strong species richness gradient from West (rich) to Eastern Amazonian
forests (poor) [19]. The lower metacommunity effect for the edges then is most likely
not due to less absolute influx of genera, but rather less influx from the Amazonian
tree community. Influx from the species-rich Andes could account for the high
diversity [20], yet low Amazonian metacommunity effect for Western Amazonian
forests. In contrast, South-eastern parts of Amazonia receive influx from tree species-
poor biomes (i.e. the Cerrado) resulting in lower diversity but also low
metacommunity effect for Amazonian trees in this region.

**S-A2 List of packages used in addition to standard R [1] preloaded packages**

*Vegan* Jari Oksanen, F. Guillaume Blanchet, Michael Friendly, Roeland Kindt, Pierre Legendre, Dan McGlinn, Peter R. Minchin, R. B. O'Hara, Gavin L. Simpson, Peter Solymos, M. Henry H. Stevens, Eduard Szoecs and Helene Wagner (2020). vegan: Community Ecology Package. R package version 2.5-7.   https://CRAN.R-project.org/package=vegan

*FD* Laliberté, E., and P. Legendre (2010) A distance-based framework for measuring functional diversity from multiple traits. Ecology 91:299-305.

*FD* Laliberté, E., Legendre, P., and B. Shipley. (2014). FD: measuring functional diversity from multiple traits, and other tools for functional ecology. R package version 1.0-12.

*Fields* Douglas Nychka, Reinhard Furrer, John Paige, Stephan Sain (2021). "fields: Tools for spatial data." R package version 13.3, https://github.com/dnychka/fieldsRPackage

*Binr* Sergei Izrailev (2015). binr: Cut Numeric Values into Evenly Distributed Groups. R package version 1.1. https://CRAN.R-project.org/package=binr

*doParallel* Microsoft Corporation and Steve Weston (2020). doParallel: Foreach Parallel Adaptor for the 'parallel' Package. R package version 1.0.16. https://CRAN.R-project.org/package=doParallel

*Itertools* Steve Weston and Hadley Wickham (2014). itertools: Iterator Tools. R package version 0.1-3. https://CRAN.R-project.org/package=itertools

*Reshape* Hadley Wickham (2007). Reshaping Data with the reshape Package. Journal of Statistical Software, 21(12), 1-20. URL http://www.jstatsoft.org/v21/i12/.

*Ggplot2* H. Wickham. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York, 2016.

*Plotrix* Lemon, J. (2006) Plotrix: a package in the red light district of R. R-News, 6(4): 8-12.

*Mice* Stef van Buuren, Karin Groothuis-Oudshoorn (2011). mice: Multivariate Imputation by Chained Equations in R. Journal of Statistical Software, 45(3), 1-67. DOI 10.18637/jss.v045.i03.

## REFERENCES SUPPLEMENTARY MATERIAL

[1]   B. Shipley, *From Plant Traits to Vegetation Structure. Chance and Selection in the Assembly of Ecological Communities* (Cambridge University Press, 2010).

[2]   B. Shipley, C. E. T. Paine, and C. Baraloto, *Quantifying the Importance of Local Niche-Based and Stochastic Processes to Tropical Tree Community Assembly*, Ecology **93**, 760 (2012).

[3]   J. Loranger, F. Munoz, B. Shipley, and C. Violle, *What Makes Trait–Abundance Relationships When Both Environmental Filtering and Stochastic Neutral Dynamics Are at Play?*, Oikos **127**, 1735 (2018).

[4]   A. Colin Cameron and F. A. G. Windmeijer, *An R-Squared Measure of Goodness of Fit for Some Common Nonlinear Regression Models*, J. Econom. **77**, 329 (1997).

[5]   B. Shipley, *Measuring and Interpreting Trait-Based Selection versus Meta-Community Effects during Local Community Assembly*, J. Veg. Sci. **25**, 55 (2014).

[6]   H. ter Steege, N. C. a Pitman, O. L. Phillips, J. Chave, D. Sabatier, A. Duque, J.-F. Molino, M.-F. Prévost, R. Spichiger, H. Castellanos, P. von Hildebrand, and R. Vásquez, *Continental-Scale Patterns of Canopy Tree Composition and Function across Amazonia.*, Nature **443**, 444 (2006).

[7]   A. A. Agrawal, *Macroevolution of Plant Defense Strategies*, Trends Ecol. Evol. **22**, 103 (2007).

[8]   J. C. Ordoñez, P. M. Van Bodegom, J. P. M. Witte, I. J. Wright, P. B. Reich, and R. Aerts, *A Global Study of Relationships between Leaf Traits, Climate and Soil Measures of Nutrient Fertility*, Glob. Ecol. Biogeogr. **18**, 137 (2009).

[9]   R. S. Oliveira, F. R. C. Costa, E. van Baalen, A. de Jonge, P. R. Bittencourt, Y. Almanza, F. de V. Barros, E. C. Cordoba, M. V. Fagundes, S. Garcia, Z. T. T. M. Guimaraes, M. Hertel, J. Schietti, J. Rodrigues-Souza, and L. Poorter, *Embolism Resistance Drives the Distribution of Amazonian Rainforest Tree Species along Hydro-Topographic Gradients*, New Phytol. (2018).

[10]  C. Levis, B. M. Flores, P. A. Moreira, B. G. Luize, R. P. Alves, J. Franco-Moraes, J. Lins, E. Konings, M. Peña-Claros, F. Bongers, F. R. C. Costa, and C. R. Clement, *How People Domesticated Amazonian Forests*, Front. Ecol. Evol. **5**, (2018).

[11]  J. E. Guevara, G. Damasco, C. Baraloto, P. V. A. Fine, M. C. Peñuela, C. Castilho, A. Vincentini, D. Cárdenas, F. Wittmann, N. Targhetta, O. Phillips, J. Stropp, I. Amaral, P. Maas, A. Monteagudo, E. M. Jimenez, R. Thomas, R. Brienen, Á. Duque, W. Magnusson, C. Ferreira, E. Honorio, F. de Almeida Matos, F. R. Arevalo, J. Engel, P. Petronelli, R. Vasquez, and H. ter Steege, *Low Phylogenetic Beta Diversity and Geographic Neo-Endemism in Amazonian White-Sand Forests*, Biotropica **48**, 34 (2016).

[12] H. ter Steege, N. C. A. Pitman, D. Sabatier, C. Baraloto, R. P. Salomão, J. E. Guevara, O. L. Phillips, C. V Castilho, W. E. Magnusson, J.-F. Molino, A. Monteagudo, P. Núñez Vargas, J. C. Montero, T. R. Feldpausch, E. N. H. Coronado, T. J. Killeen, B. Mostacedo, R. Vasquez, R. L. Assis, J. Terborgh, F. Wittmann, A. Andrade, W. F. Laurance, S. G. W. Laurance, B. S. Marimon, B.-H. Marimon, I. C. Guimarães Vieira, I. L. Amaral, R. Brienen, H. Castellanos, D. Cárdenas López, J. F. Duivenvoorden, H. F. Mogollón, F. D. de A. Matos, N. Dávila, R. García-Villacorta, P. R. Stevenson Diaz, F. Costa, T. Emilio, C. Levis, J. Schietti, P. Souza, A. Alonso, F. Dallmeier, A. J. D. Montoya, M. T. Fernandez Piedade, A. Araujo-Murakami, L. Arroyo, R. Gribel, P. V. A. Fine, C. A. Peres, M. Toledo, G. A. Aymard C, T. R. Baker, C. Cerón, J. Engel, T. W. Henkel, P. Maas, P. Petronelli, J. Stropp, C. E. Zartman, D. Daly, D. Neill, M. Silveira, M. R. Paredes, J. Chave, D. de A. Lima Filho, P. M. Jørgensen, A. Fuentes, J. Schöngart, F. Cornejo Valverde, A. Di Fiore, E. M. Jimenez, M. C. Peñuela Mora, J. F. Phillips, G. Rivas, T. R. van Andel, P. von Hildebrand, B. Hoffman, E. L. Zent, Y. Malhi, A. Prieto, A. Rudas, A. R. Ruschell, N. Silva, V. Vos, S. Zent, A. A. Oliveira, A. C. Schutz, T. Gonzales, M. Trindade Nascimento, H. Ramirez-Angulo, R. Sierra, M. Tirado, M. N. Umaña Medina, G. van der Heijden, C. I. A. Vela, E. Vilanova Torre, C. Vriesendorp, O. Wang, K. R. Young, C. Baider, H. Balslev, C. Ferreira, I. Mesones, A. Torres-Lezama, L. E. Urrego Giraldo, R. Zagt, M. N. Alexiades, L. Hernandez, I. Huamantupa-Chuquimaco, W. Milliken, W. Palacios Cuenca, D. Pauletto, E. Valderrama Sandoval, L. Valenzuela Gamarra, K. G. Dexter, K. Feeley, G. Lopez-Gonzalez, M. R. Silman, S. P. Hubbell, F. He, R. Condit, L. Borda-de-Agua, J. Kellner, H. Ter Steege, G. A. Black, T. H. Dobzhansky, C. Pavan, J. M. Pires, T. Dobzhansky, G. A. Black, M. J. G. Hopkins, M. J. Costello, R. M. May, N. E. Stork, P. Haripersaud, H. ter Steege, J.-J. de Granville, H. Chevillotte, M. Hoff, D. P. Bebber, M. A. Carine, J. R. Wood, A. H. Wortley, D. J. Harris, G. T. Prance, G. Davidse, J. Paige, T. D. Pennington, N. K. Robson, R. W. Scotland, B. J. McGill, R. S. Etienne, J. S. Gray, D. Alonso, M. J. Anderson, H. K. Benecha, M. Dornelas, B. J. Enquist, J. L. Green, F. He, A. H. Hurlbert, A. E. Magurran, P. A. Marquet, B. A. Maurer, A. Ostling, C. U. Soykan, K. I. Ugland, E. P. White, R. J. Warren, D. K. Skelly, O. J. Schmitz, M. A. Bradford, N. C. A. Pitman, J. W. Terborgh, M. R. Silman, P. N. V, D. A. Neill, C. E. Cerón, W. A. Palacios, M. Aulestia, N. C. A. Pitman, M. R. Silman, J. W. Terborgh, F. D. Lozano, M. W. Schwartz, M. W. Schwartz, D. Simberloff, J. E. Richardson, R. T. Pennington, T. D. Pennington, P. M. Hollingsworth, T. L. Couvreur, F. Forest, W. J. Baker, S. Cavers, C. W. Dick, D. H. Janzen, S. A. Mangan, S. A. Schnitzer, E. A. Herre, K. M. Mack, M. C. Valencia, E. I. Sanchez, J. D. Bever, W. Balée, D. G. Campbell, C. Levis, P. F. de Souza, J. Schietti, T. Emilio, J. L. P. V. Pinto, C. R. Clement, F. R. C. Costa, D. A. Posey, E. Montoya, V. Rull, N. D. Stansell, M. B. Abbott, S. Nogué, B. W. Bird, W. A. Díaz, C. Gomez-Navarro, C. Jaramillo, F. Herrera, S. L. Wing, R. Callejas, C. H. McMichael, D. R. Piperno, M. B. Bush, M. R. Silman, A. R. Zimmerman, M. F. Raczka, L. C. Lobato, H. ter Steege, P. P. Haripersaud, O. S. Bánki, F. Schieving, S. J. Phillips, R. P. Anderson, R. E. Schapire, S. J. Phillips, M. Dudik, C. A. Quesada, J. Lloyd, L. O. Anderson, N. M. Fyllas, M. Schwarz, C. I. Czimczik, B. Rollet, H. ter Steege, N. C. Pitman, O. L. Phillips, J. Chave, D. Sabatier, A. Duque, J. F. Molino, M. F. Prévost, R. Spichiger, H. Castellanos, P. von Hildebrand, R. Vásquez, P. M. Fearnside, D.

Mouillot, D. R. Bellwood, C. Baraloto, J. Chave, R. Galzin, M. Harmelin-
Vivien, M. Kulbicki, S. Lavergne, S. Lavorel, N. Mouquet, C. E. Paine, J.
Renaud, W. Thuiller, G. Lopez-Gonzalez, S. L. Lewis, M. Burkitt, O. L.
Phillips, P. J. M. Maas, L. Y. T. Westra, H. Rainer, A. Q. Lobão, R. H. J.
Erkens, P. V. A. Fine, D. C. Daly, G. V. Muñoz, I. Mesones, K. M. Cameron,
K. J. Feeley, M. R. Silman, M. Dufrene, P. Legendre, J. Bunge, M. Fitzpatrick,
J. Bunge, L. Woodard, D. Böhning, J. A. Foster, S. Connolly, H. K. Allen, J.-P.
Z. Wang, B. G. Lindsay, A. Chao, R. K. Colwell, C.-W. Lin, N. J. Gotelli, C.
X. Mao, R. K. Colwell, U. Brose, N. D. Martinez, R. J. Williams, I. J. Good, A.
Chao, S.-M. Lee, I. Rocchetti, J. Bunge, D. Bohning, R. A. Fisher, A. S.
Corbet, C. B. Williams, F. W. Preston, J.-P. Wang, P. V. A. Fine, and R. H.
Ree, *Hyperdominance in the Amazonian Tree Flora*, Science **342**, 1243092
(2013).
[13]  L. H. M. Cosme, J. Schietti, F. R. C. Costa, and R. S. Oliveira, *The Importance*
*of Hydraulic Architecture to the Distribution Patterns of Trees in a Central*
*Amazonian Forest*, New Phytol. **215**, 113 (2017).
[14]  S. E. Russo, S. J. Davies, D. A. King, and S. Tan, *Soil-Related Performance*
*Variation and Distributions of Tree Species in a Bornean Rain Forest*, J. Ecol.
**93**, 879 (2005).
[15]  K. J. Gaston, *Species-Range Size Distributions: Products of Speciation,*
*Extinction and Transformation: Philosophical Transactions of the Royal*
*Society of London Series B*, **353**, 219 (1998).
[16]  R. E. Ricklefs, H. Qian, and P. S. White, *The Region Effect on Mesoscale Plant*
*Species Richness between Eastern Asia and Eastern North America*, **2**, 129
(2004).
[17]  F. Munoz, B. R. Ramesh, and P. Couteron, *How Do Habitat Filtering and*
*Niche Conservatism Affect Community Composition at Different Taxonomic*
*Resolutions?*, Ecology **95**, 2179 (2014).
[18]  R. K. Colwell, C. Rahbek, and N. J. Gotelli, *The Mid-Domain Effect and*
*Species Richness Patterns:What Have We Learned so Far?*, Am. Nat. **163**, E1
(2004).
[19]  H. ter Steege, N. Pitman, D. Sabatier, H. Castellanos, P. Van der Hout, D. C.
Daly, M. Silveira, O. Phillips, R. Thomas, J. V. A. N. Essen, H. Mogollon, and
W. Morawetz, *A Spatial Model of Tree a -Diversity and Tree Density for the*
*Amazon*, Biodivers. Conserv. **12**, 2255 (2003).
[20]  T. F. Rangel, N. R. Edwards, P. B. Holden, J. A. F. Diniz-Filho, W. D.
Gosling, M. T. P. Coelho, F. A. S. Cassemiro, C. Rahbek, and R. K. Colwell,
*Modeling the Ecology and Evolution of Biodiversity: Biogeographical Cradles,*
*Museums, and Graves. Science*, Science. **361**, (2018).
1218