1 **Virulence shift in a sexual clade of Type X *Toxoplasma* infecting Southern Sea Otters**
2

KENNARD, A[1], MILLER, MA[2], KHAN, A[1*], QUINONES, M[3], MILLER, N[4], SUNDAR, N[1], JAMES,
ER[5], ROOS, DR[4], CONRAD, PA[6] AND GRIGG, ME[1,5].
[1]Laboratory of Parasitic Diseases, NIAID, National Institutes of Health, Bethesda, MD, USA. [2]California
Department of Fish and Game, Santa Cruz, CA, USA. [3]Bioinformatics and Computational Biosciences Branch,
NIAID, National Institutes of Health, Bethesda, MD, USA. [4]Department of Biology and Penn Genome Frontiers
Institute, University of Pennsylvania, Philadelphia, Pennsylvania, USA. [5]Department of Medicine, University of
British Columbia, Vancouver, BC, Canada. [6]School of Veterinary Medicine, University of California, Davis, CA,
USA.
11
*Current Address: Agricultural Research Service, USDA, Beltsville, MD, USA.

Abstract

How virulent protozoal pathogens capable of causing overt disease are maintained in nature
is an important paradigm of eukaryotic pathogenesis. Here we used population genetics and
molecular methods to study the evolution and emergence of a marine invasion of new
genetic variants of *Toxoplasma gondii,* referred collectively as Type X (HG12). 53 *Toxoplasma*
isolates were obtained from mustelids that stranded between 1998-2004 with
toxoplasmosis (ranging from chronic infection to fatal encephalitis). Over 74% of the sea
otters collected throughout their geographic range were infected with Type X as determined
by multi-locus PCR-DNA sequencing. Depending on the locus investigated, Type X strains
possessed one of three allelic types that had independently assorted across the strains
examined; either genetically distinct alleles, referred to as "γ" or "δ", or a Type II allele.
Phylogenetic incongruence among locus-specific trees, genome-wide CGH array and WGS
analyses confirmed that Type X is a sexual clade of natural recombinants that resemble F1
progeny from a genetic cross between Type II and a mosaic of two distinct "γ" or "δ"
ancestries. A single Type X genotype (19/53; 36%) had expanded in sea otters largely as
subclinical chronic infections, but it was highly pathogenic to mice (LD_{100}= 1 parasite). To
determine whether murine virulence genes could be mapped within this naturally occurring
population, we performed a genome scan and identified four QTLs with LOD scores greater
than 4.0. Targeted disruption of ROP33, the strongest candidate from among 16 genes within
the highest QTL on Chromosome VIIa established ROP33 as a murine virulence locus. The
ability of this highly pathogenic clone to expand and cause the majority of sea otter infections
supports a virulence shift model whereby generalist pathogens like *Toxoplasma* utilize their
sexual cycles to produce new strains with an expanded biological potential. Such a trait
enables pathogens to extend their host range or be naturally selected within their vast
intermediate host range to maximize their transmission. Our work thus establishes a
rationale for how virulent strains can be maintained cryptically in nature across a pathogen's
broad host range, and act as reservoirs for epidemic disease.

Importance
Waterborne outbreaks of protozoal parasites are increasingly causing fatal disease in a wide
range of animals, including humans. Population expansion of felids near marine estuarine
environments has led to increased exposure of marine wildlife to highly infectious
Toxoplasma gondii oocysts shed in the feces of cats that are dispersed by storm events. In
North America *Toxoplasma* is thought to possess a highly clonal population structure

dominated by 4 clonal lineages (I, II, III, and X). Population genetic analysis of 53 *Toxoplasma*
isolates collected longitudinally from mustelids infected with *Toxoplasma* that stranded
between 1998-2004 identified a majority of otters (74%) to be infected with Type X
Toxoplasma, and that Type X is not a clonal lineage, but rather a recombinant clade of strains
consistent with a recent genetic cross that produced at least 12 distinct haplotypes.
Importantly, one Type X haplotype expanded in 36% of otters across their geographic range
and caused relatively benign infections, however it was highly pathogenic to mice. A genome
scan was performed to identify a new virulence locus, a secreted serine threonine kinase
(ROP33), that is pathogenic in mice, but not sea otters. Our data support a virulence shift
model whereby generalist pathogens like *Toxoplasma* utilize their sexual cycles to produce
virulent strains that are maintained cryptically in nature, according to their differential
capacity to cause disease within the pathogen's broad intermediate host range. This type of
"zoonotic selection" has important public health implications. Strains capable of causing fatal
infections can persist in nature by circulating as chronic infections in intermediate host
species that act as reservoirs for epidemic disease.
63
64   **Introduction**
65   How successful microbes maintain reservoirs of virulent strains in nature is
understudied and is an important paradigm of infectious diseases. Whereas many studies
document how microbial pathogens acquire or admix genetic material to influence their
pathogenicity, relatively less is known how emergent strains are selectively maintained in
nature. Specifically, it is well established that viruses utilize reassortment to rapidly produce
admixture lines that possess a range of altered biological potential, including virulence [1].
Bacteria transfer mobile plasmid elements that encode pathogenicity islands via conjugation
to confer an altered virulence potential on those strains that receive them [2-4]. Among both
fungi and protozoa, genetic hybridization by sexual recombination is a potent way to
produce genomic variation or new admixture lines that possess increased virulence and are
capable of causing disease outbreaks or altering host range [5-7]. What is less clear are the
mechanisms at play allowing highly prevalent pathogens to maintain virulent strains
cryptically in nature. We sought to test whether generalist pathogens, such as the protozoan
parasite *Toxoplasma gondii*, recovered from natural infections in one animal host, leverages
its broad range of intermediate hosts to selectively partition parasite genotypic diversity and
virulence potential.
81   *Toxoplasma gondii* is a highly successful and prevalent protozoan pathogen that
infects a wide range of wildlife, livestock and 20-80% of humans worldwide [8-10]. Its
success in nature is largely attributed to its highly flexible life cycle. It is propagated asexually
by carnivory among essentially all warm-blooded vertebrates (via orally infectious tissue
cysts) or sexually within its definitive felid host by self-mating (when a single parasite
undergoes fertilization and sexually clones itself) or by out-crossing (producing as many as
10^8 genetic hybrids that are transmissible as environmentally stable, highly infectious
oocysts). Genetic hybridization has been previously shown to produce genotypes that
possess a wide spectrum of biological potential, including altered virulence or a capacity to
expand as successful epidemic clones [7, 11, 12]. Importantly, parasite clones possess a
broad spectrum of disease states that are highly dependent on both parasite genetics and the
animal species infected [8, 13-15]. In rodents, low dose inoculums of Type I *Toxoplasma*
strains are uniformly lethal to laboratory mice [16] whereas Type II strains are considerably

less virulent and routinely establish chronic, transmissible infections [14]. In contrast, wild-
derived CIM mice that encode resistant alleles of immunity-related GTPases (IRGs) confer
resistance to virulent Type I strains by inactivating parasite-specific secreted kinases
(ROP18/ROP5) that control mouse susceptibility to infection [17, 18]. The end result of this
host genetic, parasite genetic interplay determines the potential for a disease-producing
virulent clone to expand or be maintained cryptically within a strain-specific, or host species-
specific, manner, but this has not so far been demonstrated to occur in nature and has only
been inferred by laboratory studies. Likewise, laboratory rats are resistant to Type I
infections and fail to transmit parasites, but again this is entirely rat species-dependent [19].
The molecular basis for this resistance is an allele-dependent activation of the NLRP1
inflammasome resulting in macrophage pyroptosis and the inhibition of parasite growth in
Lewis (LEW) and Sprague Dawley (SD) rats, but not in Brown Norway (BN) or Fischer CDF
rats, which go on to establish chronic, transmissible infections of mouse-virulent Type I
strains [20-22]. While these laboratory studies suggest the possibility that virulent strains
can be maintained cryptically across the pathogen's host range [23], the phenomenon has
not been systematically investigated nor demonstrated to occur in a natural setting.
Between 1998 and 2004, *Toxoplasma gondii* emerged as a significant disease in
California sea otters (*Enhydra lutris nereis*), a federally listed threatened species [24, 25]. Full
necropsies performed on each stranded sea otter identified a wide spectrum of disease
sequelae among *Toxoplasma*-infected otters, ranging from chronic asymptomatic infection
to fatal meningoencephalitis at time of death [25, 26]. Importantly, the majority of sea otters
were found to be infected by a single outbreak clone, referred to as Type X (or HG12) [24].
Based on multi-locus PCR-DNA sequencing (MLST) using a limited set of genotyping
markers, Type X was defined as the fourth clonal lineage in the US [27] and it is highly
prevalent in sylvatic niches as chronic, subclinical or mild infections in 47% of US wildlife
[28, 29]. The range of disease sequelae detected in sea otters, however, was not consistent
with infection by a single clone. Other possibilities proposed to explain variation in disease
susceptibility have included co-infection [30], exposure to environmental pollutants, or
toxins that cause immunosuppression [31]. More recently, WGS performed on *Toxoplasma*
isolates has called into question whether MLST analyses are sufficiently resolved to predict
clonotypes because they fail to capture extant genetic heterogeneity [15]. Indeed, a previous
study concluded that Type X was comprised of two genotypes [32], but these two genotypes
differed by only a few SNPs, so it was not clear whether strain variation was likewise playing
a contributing role in sea otter disease in this novel host-parasite interaction.
To test for this and to determine the true extent of genetic diversity among naturally
circulating Type X isolates, we sequenced a wide selection of linked and unlinked markers
across the nuclear and organellar genomes for 53 isolates collected longitudinally from
infected sea otters across a seven year period. Based on these results, we selected 16 isolates
for WGS. Analysis of our data support a model whereby Type X exists as a recombinant clade
of strains that resemble F1 progeny from a natural cross between a Type II strain and a novel
genotype that has not been identified previously in nature. We show that one sea otter
genotype is widely distributed and has expanded to cause the majority of infections, which
were largely subclinical and consistent with a strain that has been naturally selected in sea
otters. This genotype, however, was highly pathogenic to laboratory mice and a population
level, forward genetic approach using WGS mapped a mouse virulence gene to a novel
parasite-specific secreted kinase ROP33, also referred to as WNG3 [33]. Our natural

population dataset argues that *Toxoplasma* leverages its broad range of intermediate hosts to partition genetic diversity. Further, that intermediate hosts play a central role in the natural selection, expansion and maintenance of cryptically virulent strains in a host species-specific manner and that these infected hosts act as reservoirs for pathogenic or epidemic disease in another host species.

Results

Genetic characterization of *Toxoplasma gondii* strains isolated from sea otters

In the late 1990s, an increasing number of sea otters stranded along the California coast with protozoal encephalitis due to *Toxoplasma gondii* [25, 29, 32, 34]. An initial genetic analysis performed on a subset of 35 *T. gondii* isolates established that 21 were a new genetic type, referred to as Type X (also known as Haplogroup 12, or HG12), but this conclusion was limited to sequence typing at a single locus, GRA6 [25]. We developed a multi-locus (MLST) PCR-DNA sequence-based typing scheme using an additional 4 unlinked markers (BSR4, BAG1, SAG3, and ROP1) that possess a wide-range of phylogenetic strength, including markers under neutral (BSR4), diversifying (GRA6, SAG3, ROP1), and purifying (BAG1) selection to determine the molecular genotypes of 53 *T. gondii* isolates collected longitudinally over a seven-year period between 1998-2004 from fresh, beachcast California sea otters. Isolates were obtained throughout the California sea otter range (Supplemental Figure 1). Antibody titers against *Toxoplasma* (when available), stranding location, otter sex, and co-infection status with other protozoal agents was available for 52 of the sea otters (Supplemental Figure 1).

Point-source outbreak of Type II strains infecting sea otters

To determine the *T. gondii* genotypes among the sea otter isolates, the 5 locus MLST was applied. Typing at just the GRA6 marker identified three sequence types, either a Type II allele in 16 (30%) isolates, or two non-archetypal alleles, previously identified as X or A [25, 32], in 25 (47%) and 12 (23%) isolates, respectively. Expanding the genetic analysis across five markers showed that 2 of 16 isolates with a Type II allele at GRA6 were predicted to be non Type II, recombinant strains, because they possessed non-archetypal alleles at ROP1 (3675) or BSR4 and BAG1 (3671) (Figure 1A). Of the remaining 14 otters infected with Type II strains, 12 (86%) were identified within a restricted geographic range, a 40 mile stretch off the California coast line containing Moss Landing and Monterey Bay, which is consistent with a point source outbreak of a *Toxoplasma* strain that is found most frequently infecting human and domestic livestock in the USA. The other two Type II infected otters were recovered from Pismo Beach in Central California. Further, the majority of Type II infected otters (9/14; 64%) were collected during the first half of the study (Supplemental Figure 1). Males were predominantly infected with this genotype, and it was the only genotype that had a male-specific infection sex bias. [Fisher's two-sided exact test II vs. all; p=0.008]

Type X is a recombinant clade of strains

Of the remaining 39 (74%) *Toxoplasma* isolates, collectively referred to as Type X or A by the allele present at GRA6, a maximum of only three alleles were identified at the additional sequenced loci, either a canonical Type II allele or one of two genetically distinct alleles,

referred to as "γ" or "δ", that appeared to segregate independently across the isolates (Figure
1A). In total 8 distinct haplotypes designated A-H were resolved based on their differential
inheritance of the limited alleles, consistent with recombination between a Type II strain and
a mosaic of two distinct ancestries as the most plausible explanation for the genetic
relationship among the non-Type II strains. The Type X haplotype designated A was
dominant (19/39; 49%). It was widely distributed across the entire geographic range of the
Southern sea otter (Supplemental Figure 1), and it was isolated in every year except 2000.
Importantly, this genotype had expanded during the time frame of the study period, and it
was responsible for the majority of infections by 2003 (Figure 1B).
195
196 **Acute virulence in mice is dependent on Type X haplotype**
197   Mouse virulence is highly dependent on *Toxoplasma* genotype [7, 16, 35]. To further
characterize the otter isolates, 18 strains, including two Type II and at least one isolate
representing each of the 8 distinct Type X haplotypes, were tested for their mouse virulence
phenotype. Isolates were selected based largely on their ability to expand *in vitro* in human
foreskin fibroblasts. Virulence was assayed in female CD-1 outbred mice using a low-dose,
50 tachyzoite, intra-peritoneal injection model. At this inoculum, mouse avirulent strains
(*e.g.* Type II) establish chronic, transmissible infections whereas mouse virulent strains (*e.g.*
Type I) die acutely within 10-14 days.
205   Among the otter isolates, three mouse virulence phenotypes were identified: virulent
(red; all mice died acutely), intermediate virulent (blue; some mice survived acute disease),
and avirulent (green; all mice survived acute infection and produced chronic disease). The
two otter isolates that possessed a Type II MLST were avirulent (Figure 1C; green) and
phenocopied the infection kinetics of two well studied Type II lines Me49 and 76K. Among
the Type X haplotypes, E and G were also avirulent. Three otter isolates from haplotype A,
the genotype that had expanded to cause the majority of sea otter infections, were highly
pathogenic to mice; all infected mice died acutely (Figure 1C; red). Haplotypes D and H were
likewise highly virulent in mice. The three haplotype F otter isolates all possessed an
intermediate virulence phenotype (Figure 1C; blue). Of particular note, the 5 isolates from
haplotypes B and C had a mix of two mouse virulence phenotypes, they were either avirulent
(green) or they were intermediate virulent (blue) (Figure 1C). Because the Type X isolates
displayed a range of virulence phenotypes, the data collectively support a model whereby
Type X represents a clade of recombinants from a genetic admixture of a mouse avirulent
Type II strain with at least one other parent that is mouse virulent, rather than expansion of
a single clonal lineage. Furthermore, not all isolates within each subgroup displayed
equivalent virulence kinetics, in particular isolates from the B and C haplotypes were either
avirulent or possessed intermediate virulence, suggesting that the 5 marker MLST was
insufficiently resolved to fully capture total genetic and phenotypic diversity among the
isolates recovered from infected Southern sea otters.
225   To explore this possibility, the 18 isolates assayed through mice plus 3 additional
isolates, one each from haplotypes A, C and D, were further genotyped at 18 markers (17
nuclear-encoded single copy gene loci plus one microsatellite marker) representing an
expanded set of linked and unlinked loci across the genome. Additionally, 2 markers, one
each from the organellar genomes of the apicoplast and mitochondria, were included to
determine the ancestry of each maternally inherited genome. This represents a significant
increase in the number of markers that were previously used to conclude that Type X (HG12)

is a clonal lineage [10, 27, 36]. Altogether, the 20 marker MLST method provided 15,404 bp
of sequence information to determine the genetic relationship among *Toxoplasma* strains
isolated from Southern sea otters (Supplemental Table 1).
235
Type X is comprised of 12 distinct haplotypes by an expanded MLST analysis
To establish the number of haplotypes among the 21 isolates, sequence data for the
17 nuclear-encoded single copy gene loci was concatenated and analysed using eBURST
(Figure 1D). eBURST resolves strains into clonal complexes (CC) that are in linkage
disequilibrium. In this case, coded MLST data was used to identify strains that differ at one
locus (or share 16 out of 17 markers) and isolates that differed at the same genetic locus
were grouped together and connected by a line within each clonal complex. As expected,
eBURST clustered the two Type II isolates (2987, 3131) into a single clonal complex with
Me49 (CC1). Type X subgroup G isolate 3675, originally identified by its allele at the
microsatellite marker ROP1, was included within CC1 because it differed from Type II strains
at only SAG4. For the remaining 18 Type X isolates, 15 clustered into 3 clonal complexes
(designated CC2-CC4), and 3 unique haplotypes were resolved. The eBURST analysis failed
to support a clonal lineage designation for Type X. Further, the increased resolution provided
by the 17 markers subdivided the 8 distinct haplotypes defined using 5 loci in Figure 1A into
12 haplotypes (Figure 1D) which was consistent with another study that resolved Type X
Toxoplasma genotypes isolated from southern sea otters into 8 genotypes [26]. One isolate
each from subgroups A (n=4; 3168), C (n=4; 3160), D (n=3; 4166) and F (n=3; 3387) was
resolved further as a unique haplotype based on their inheritance of alleles at PK1, SAG4 and
GRA7 (Supplemental Table 1).
255
Type X resembles a recombinant clade of F1 progeny from a natural cross
eBURST is insufficiently resolved to distinguish among isolates that exist either as
variants within a clonal lineage, because they possess alleles that differ by only minor
mutational drift, or as genetic admixtures, that possess genetically distinct alleles derived
from different parental types. Previous studies suggested that Type X (or HG12) is a clonal
lineage infecting sea otters and wildlife in North America, and that HG12 represents an
expanded clone from a sexual cross between a Type II strain and a new genetic lineage, that
was referred to as "γ", but this was based largely on the alleles found at just two loci, GRA6
and GRA7 [24, 25, 27]. To investigate whether the genetic relationship among the 12 distinct
haplotypes is supported by a model of genetic drift, or that of recombination, individual
maximum likelihood trees were created for each of the 17 nuclear-encoded single gene copy
loci, the microsatellite locus, and the two organellar loci (Supplemental Figure 1). To
differentiate between alleles undergoing minor mutational drift, from those that have
evolved independently as distinct genetic outgroups, 1000 bootstrap replicates were run for
each tree, and supported nodes above 60% are indicated at each marker. For each tree, the
19 isolates identified as Type X were labeled in purple and isolates belonging to the clonal
lineage I (n=1), II (n=3), and III (n=1) were labeled in red, green, and blue respectively
(Figure 2).
At 5 genetic markers located on chromosomes Ib, II, III, XI, and XII, all 12 distinct Type
X haplotypes possessed a canonical Type II allele (green), with no minor mutational drift
detected, consistent with Type II being one of the parental genetic backgrounds
(Supplemental Table 1). At the L358 marker on chromosome V, however, all Type X isolates

except 3675 possessed a Type I or "α" allele (red), indicating the presence of mixed ancestry.
Further, at the BSR4 marker on chromosome IV, all Type X isolates except 3675 possessed
an entirely novel allele that was readily differentiated from Type I, II and III strains with
strong bootstrap support (Figure 2B). At all remaining loci, Type X isolates possessed either
a Type II or one of 2 genetically distinct alleles, which could not be readily explained by minor
mutational drift. Hence, across the 17 nuclear-encoded markers, each isolate possessed
either a Type II allele, or an allele of distinct ancestry that was referred to as belonging to the
"α" (red), "γ" (purple) or "δ" (orange) lineages.
The maternally inherited organellar genome markers were likewise mixed. At the
mitochondrial marker CO1, two alleles were identified, either an allele common to Type I, II
and III strains or a novel allele designated "γ" (Figure 2A). Further, 3 distinct alleles were
identified at the apicolast marker APICO: a Type I or "α" allele (red), a Type II allele (green)
or a novel "γ" allele (purple), supporting the mixed ancestry designation. Each of the 12 Type
X haplotypes were therefore a mosaic of mixed ancestry that possessed some combination
of Type II, Type I, "γ" or "δ" alleles that had segregated independently across the loci
investigated. The data are consistent with Type X existing as a recombinant clade resembling
F1 progeny that was recently derived, without sufficient time to develop minor mutational
drift (Supplemental Table 1).
In support of the sexual cross model, incongruity between phylogenies for each
haplotype was readily observed between markers located in different portions of the
genome. In Figure 2C, 8 isolates shared the same ancestry between the two genetic loci
located on separate chromosomes VIIa and VIIb (GRA7 and BAG1 respectively) and were
shaded either green (Type II alleles at both loci) or purple (γ lineage alleles at both loci). In
contrast, 13 isolates had independently segregated chromosomes of mixed ancestry, colored
orange and pink (to depict isolates that were discordant and possessed different ancestral
alleles at each locus), and the crossing of orange and pink lines highlighted this lack of
congruence (Figure 2C). For example, isolate 3671, had a γ lineage allele at GRA7 but a δ
lineage allele at BAG1. Furthermore, recombination within a chromosome was likewise
observed at linked markers for 18 strains (Figure 2D). For example, isolate 3387, had a γ
lineage allele at GRA7 but a δ lineage allele at SAG4, whereas 3675 had a II lineage allele
GRA7 but a γ lineage allele SAG4, formally establishing that recombination had occurred and
supporting genetic hybridization as the most plausible explanation for the origin of the 12
Type X haplotypes.
311
Type X is an admixture cross between Type II and a novel genetic background by CGH-
array hybridization
A subset of Type X isolates, 2 each from haplotypes A (3142, 3265), C (3026, 3045), F
(3503, 3429) and one each from D (3178) and H (3671) were next hybridized against a
photolithographic microarray that possessed 1517 polymorphic Type I, II and III strain-
specific genotyping probes distributed genome-wide [37]. This was done to test whether the
MLST genetic markers accurately predicted the presence of large genome-wide haplotype
specific blocks, consistent with genetic hybridization. Each Type-specific SNP on the CGH
array is represented as a dot colored either red, green or blue respectively for a hybridizing
strain that possesses a Type I, II or III SNP at the probe position. Dots colored grey only
identify probes that failed to hybridize, consistent with a strain that does not possess a Type

I, II or III specific SNP at the probe position. Hybridization with DNA from a Type I (GT1), Type II (Me49), and Type III (CTG) strain identified hybridization patterns consistent with genotype, indicating that all genotype-specific probes were functioning as expected (Figure 3A). Hybridization with the Type X isolates identified large, contiguous Type II haploblocks, as evidenced by hybridization of all Type II-specific probes present on Chr III, VI, VIIb and XII, for example. However, in other portions of the genome, the SNP diversity pattern was novel. The hybridization pattern in these regions either shared patch-work similarity with Type I (i.e., at Chr Ib, VI) or was highly divergent with large contiguous blocks of SNP probes colored grey that failed to hybridize, which was consistent with introgression of a non-Type I, II or III genetic ancestry, rather than minor mutational drift as the explanation for lack of reactivity (i.e., Chr IV, XI, right end of Chr VII, XII, left end of Chr VIII).

The pattern of hybridizing SNPs was unique for each of the seven Type X isolates examined, but it was only minimally different; either 3, 5, or 15 SNP differences were detected in pairwise comparisons within haplotype A (3142, 3265), C (3026, 3045), and F (3503, 3429), respectively (Figure 3A). At this resolution, it was not possible to distinguish between CGH array hybridization efficiency, from that of minor mutational drift. Indeed, the CGH arrays each had approximately the same number of SNP differences (3-15) between haplotypes as they did within a haplotype, which is more consistent with hybridization efficiency as the probable explanation for these hybridization differences. It did, however, establish that 3671 is a distinct genetic admixture because it possessed a different Type II hybridization pattern at Chr Ia, left hand of Chr Ib, VI, XI and right side of Chr XII, which was distinct from all other Type X strains analyzed (Figure 3A).

Type X is a recombinant clade at WGS resolution

The lack of resolution using the CGH array approach indicated that WGS was required to infer an accurate genetic history model for the Type X strains. Using genome-wide polymorphism data derived by identifying all variant SNP positions after reference mapping a Type I, II, III, 16 Type X, and three previously WGS sequenced HG12 isolates (WTD-1, RAY, ARI) against the published Me49 genome, we generated an unrooted Neighbor-Net tree (Figure 3B). The Neighbor-Net analysis established that all Type X isolates, in similar to Type I and III strains, shared blocks of genome-wide ancestry with Type II, represented by the reticulated pattern of edge blocks that depict recombination events between Type II and the different genetic backgrounds within X, I and III. It did not support a clonal lineage designation as the majority of Type X isolates appeared on separate branches (Figure 3B, Inset). Only clonal complexes 1 and 4 (CC1, Type II; CC4, 3045, 3160, 4167) were supported at WGS resolution, and 3 previously sequenced HG12 isolates from 2 people (ARI, RAY) and a deer (WTD-1) formed well-supported clades with Type X isolates 4166 (WTD-1; RAY) and 3168 (ARI) indicating that Type X has a wider distribution beyond sea otter infection. Outside of the reticulated network of edge blocks, the branch length for each Type X isolate was significantly less than that observed for Type I and III, synonymous with a more recent origin, without sufficient time to accumulate private SNPs by mutational drift. When murine virulence data for each genotype was mapped onto the Neighbor-Net tree (see inset, Figure 3B), a clear partitioning of the different virulence phenotypes was resolved along branches within the network, consistent with Type X isolates existing as sister progeny with only minor mutational drift detected at each supported branch.

To identify the size, genome distribution, and total number of Type II admixed haploblocks introgressed into each Type X isolate, pairwise SNP diversity plots were generated for all Type X isolates, as well as reference Type I and III strains that have been previously shown to have recombined with Type II. The Type I GT1 strain possessed an Me49 Chr Ia and IV, and an Me49 admixture block at the left side of Chr VIIa, and right end of Chr XI (Figure 4). As expected, the Type III VEG strain possessed many more Me49 admixture blocks distributed genome-wide than GT1, comprising ~40% of its genome. Only Type X haplotype H (3671) possessed haploblocks that were highly similar in sequence to Me49, on Chr Ia, II, III, VI, VIII, IX, XI, and XII (Figure 4). In other regions, however, 3671 shared regions clearly Me49-like, but that appeared to have diverged somewhat by mutational drift (average 3-5 SNPs per 10kb block) on Chr II or introgressed into Chr IV, V, VIIa, VIIb, VIII, IX, and XII. Intermixed within the Type II regions of the 3671 genome were a limited number of large haploblocks containing divergent SNP density synonymous with hybridization by a strain possessing distinct γ or δ genetic ancestry (average of 50-100 SNPs per 10kb block). When the pairwise SNP density analysis was expanded to different Type X haplotypes, isolates within each haplotype possessed highly similar patchwork mosaics of Type II-like or divergent (Type X) haploblocks that were specific to each haplotype. For example, haplotype A isolates 3142 and 3168 SNP density plots were highly similar to each other, and to haplotype C isolate 3045, but were readily distinguishable from isolates 4166 and 3503, strains within haplotypes D and F, respectively (Figure 4). Specifically, isolate 3142 had a Type X haploblock inheritance pattern at the right end of Chr V, and in the middle of Chr Ib, VIIa, VIII, IX and XII whereas 4166, which was indistinguishable from a previously sequenced HG12 strain (RAY) recovered from a human patient, possessed either Type II or 3671 haploblocks in these regions (Figure 4). Although the haplotype F isolate 3503 was highly similar in genomic organization to 3142, it possessed a Type II haploblock at the right end of Chr V that readily distinguished it from haplotype A. The pairwise SNP analysis established that the majority of Type X haplotypes possess a genomic architecture that is strikingly similar, but different at a limited number of admixture blocks. Coupled with the low allelic diversity, the data support a genetic history model whereby Type X resembles a sexual clade of recently-derived natural recombinants from a relatively limited number of crosses.

PopNet analysis identifies only limited admixture blocks among Type X isolates

While the Neighbor-Net analysis established that Type X exists as a recombinant clade of strains, it failed to predict the precise number of genetic ancestries, or how they had admixed positionally across the chromosomes. PopNet was used to paint chromosomes according to their local inheritance patterns [38]. Included in the analysis were all sequenced Type X and II strains, as well as reference Type I and III strains, known to have admixed with Type II. PopNet identified 4 statistically supported ancestries and showed that each genome was a mosaic of these distinct ancestries, supporting the admixture model.

Within the circle depicting the Type X clade, 5 distinct subgroupings were identified based on the number and position of shared ancestral blocks; these were grouped together based on line thickness (Figure 5A). These same groupings were supported both by Pairwise SNP plots and Neighbor-Net analysis, but the PopNet analysis showed which of the 5 ancestries had introgressed positionally across the mosaic genomes. Clear recombination blocks were readily resolved, and the genome architecture was remarkably similar between the groupings. A custom script designed to identify only major crossover points between the

two parents (Type II and the mosaic ancestry of the γ/δ parent) identified either a limited
number of single and double recombination events, or the inheritance of whole
chromosomes of either Type II (Chr II) or γ/δ (Chr Ia, IV, XI) parental ancestry, consistent
with this group of strains representing related sister progeny (Figure 5B). Because murine
virulence plotted on the Neighbor-Net tree identified clear pathogenicity differences in mice
that clustered based on parasite genotype, and that the Type X strains resembled a natural
clade of recombinant progeny (akin to F_1) that possessed only a restricted number of
crossover points, a population-based QTL was performed to see if genes could be mapped
that contribute to murine virulence within the set of 18 natural isolates for which virulence
data existed (Figure 5C),
424
Natural Population-based QTL identifies ROP33 (WNG-3) as a new murine Vir locus

All Type X isolates recovered from sea otters resembled offspring from one (or a few
crosses) between a Type II strain, which is avirulent in mice, and unknown parent(s) that
are mosaic for two distinct ancestries. Because these natural isolates possessed a range of
virulence phenotypes when infected into mice (Figure 1C), a genome scan was performed to
determine the log-likelihood for association of discrete genome haploblocks with the acute
virulence phenotype. Four quantitative trait loci (QTL) peaks were identified with logarithm
of odds (LOD) scores 4.0 or greater on chromosomes V, VIIa, VIII, and X (Figure 6A). The
average size of the genomic regions spanned by the QTLs were in the range of 100-200kb,
except for the one on chromosome V, which was >700 kb (Figure 6B). To identify candidate
genes within the four peaks, the following inclusion criteria were assessed: presence of a
signal peptide and/or transmembrane domain, gene expression and polymorphism
differences, as well as every genes CRISPR genome-wide mutagenesis score for essentiality
[39](Supplemental Table 2). *ROP33* on chromosome VIIa stood out as the best candidate
gene to target for reverse genetics as it was abundantly expressed during acute infection, it
was predicted to be a functional serine-threonine protein kinase, it was highly polymorphic,
and is part of a family of divergent WNG (with-no-Gly-loop) kinases that regulate tubular
membrane biogenesis in the parasite [33]. Further, one allele of *ROP33* was strongly
correlated with acute virulence (p=0.00031; fishers two-sided exact test) (Figure 6C).

To determine whether *ROP33* was a novel murine virulence gene, the *ROP33* locus was
disrupted by targeted deletion using CRISPR-Cas9 facilitated double crossover homologous
recombination in the mouse virulent RH∆*ku80*∆*rop18*∆*hxgprt* strain [40]. This strain is
virulent in mice (LD_{100}=500 tachyzoites) and was engineered to accept targeted replacement
of the *rop33* gene using an HXGPRT gene flanked by 30bp of homology just outside of the
rop33 promoter and 3'UTR region. Following selection in mycophenolic acid (MPA) and
xanthine (to select for the *HXGPRT* gene), the population was screened for disruption of the
rop33 gene by PCR. Mice were next infected with either the parent RH∆*ku80*∆*rop18*∆*hxgprt*
strain or the Rop33- mutant to assess murine virulence. Groups of 5 outbred CD-1 mice were
injected intraperitoneally with 500 tachyzoites and the results shown are for one of two
independent experiments. All mice infected with the parent RH line succumbed to infection
within 20 days (Figure 6D). In contrast, targeted deletion of the *rop33* gene was protective,
the majority of mice survived acute infection, and our results suggest that ROP33 is a novel
Toxoplasma acute virulence gene. Importantly, the lack of virulence was not the result of a
failure of the mutant parasite to proliferate *in vivo*, as no difference in parasite load was

detected through acute infection by bioluminescence imaging, indicating that deletion of
ROP33 does not appear to affect parasite proliferation *in vivo* (Figure 6E). All surviving mice
were confirmed to be positive by serology, indicating that they had been productively
infected. These data strongly suggest that ROP33 is a new virulence factor for murine
infection that was identified using natural isolates derived from a population genetic cross
performed in nature.
465
466   **Discussion**
467   Here, we investigated the emergence of a presumed clonal lineage of *Toxoplasma*
gondii (Type X; also referred to as HG12) recovered from a single marine host, infected
southern sea otters in California, that were collected longitudinally over a 7 year period in
order to determine the genetic basis for this epizootic in a federally listed, threatened
species. Although Type X has previously been identified as the 4th clonal lineage in North
American wildlife, work done here established that Type X is not a clonal lineage, but rather
a recombinant clade of strains that resemble F_1 progeny from at least one natural cross. DNA
sequence analysis using a wide selection of linked and unlinked markers across the nuclear
and organellar genomes of Type X isolates identified Type X to be a composite of Type II and
an unknown mosaic of two distinct ancestries, referred to as γ and δ, that have recombined
to produce a highly invasive clade of strains causing both subclinical infection as well as
mortality in threatened marine mammals, including the southern sea otter. Our data suggest
that Type X was derived from one or a limited number of crosses from two highly related
parental strains. One sea otter genotype was widely distributed and found to have expanded
to cause the majority of subclinical infections, but it was highly virulent to mice. Our natural
population dataset supports a model whereby *Toxoplasma*'s sexual life cycle is facilitating
the evolution and expansion of cryptically virulent strains that are capable of causing
pathogenic disease in a host-species specific manner. Hence, for generalist parasites that
have broad host ranges, natural selection among intermediate hosts appears to maximize
transmission and establishes how virulent strains can be maintained cryptically in nature.
The expansion of a mouse virulent clone into a new ecological niche, such as near-shore
marine mammals of the Eastern Pacific coastline, demonstrates how *Toxoplasma gondii* is
leveraging its intermediate host range to selectively partition parasite genetic diversity and
epidemic disease potential.
491   In previous studies, unknown strains having a distinct ancestry, referred to as α and
β, were found to have crossed with Type II to create the Type I and Type III clonal strains
respectively [11]. The frequency and pattern of recombined blocks inherited within the Type
X strains is parsimonious with previously described models for the genetic history of Types
I and III in which an unidentified ancestor (respectively α and β) sexually recombined with
an ancestral Type II to create the new lineages (Figure 5B) [11, 41]. Additionally, it has been
suggested previously that Type X is a product of genetic hybridization between a Type II
lineage and an unknown γ lineage [25, 27, 28, 32, 42]. Here, we have further refined this
perspective and concluded that Type X is a mosaic of a cross between Type II strains and two
distinct γ and δ ancestries . The γ/δ lineage has likely recombined at least once with Type II
to create the Type X recombinant clade of strains. Currently, no single isolate has been found
which could be defined as either γ or δ, in similar to the α and β lines thought to have admixed
with Type II to produce the clonal lines I and III, respectively [11, 41].

Previous work on Type X circulating in wildlife samples using a limited set of
genotyping markers identified Type X as the 4th clonal lineage in North America [10, 27].
However, this designation was based on the discovery of genetically distinct alleles at only
two loci, GRA6 and GRA7, which was the original basis for distinguishing the Type X lineage
as a new genotype from that of Type II, and that it had undergone a hybridization event with
a novel genotype [25]. It is now well established that Type X commonly infects wildlife across
North America [10, 14, 29, 30, 32, 36]. This study examined *Toxoplasma* strains collected
longitudinally across a 7 year period and large geographic range from sea otter infections,
and it was comprised of infections ranging from asymptomatic to subclinical or considered
the primary cause of death. As previous work had classified Type X as clonal, finding such a
myriad of different disease states among infected sea otters was not expected [10, 27, 32,
36]. To ascertain whether parasite genotype was a critical parameter influencing the
spectrum of disease, genotyping studies were carried out using an expanded set of markers
to ascertain the true population genetic structure of the isolates belonging to Type X. All
genotyping markers used in this study were sequenced to increase resolution rather than
evaluated strictly for their PCR-RFLP genotype, as was the standard for many of the previous
studies. While the markers used herein only surveyed a small part of the genome, they
identified at least 12 distinct haplotypes within Type X, confirming that they do not exist as
a clonal population, as hypothesized in another study [26]. Furthermore, at any given locus,
only a Type II allele or one or another of two distinct alleles was identified indicating limited
allelic diversity. Hence, each of the 12 haplotypes possessed one of three allelic types at any
one locus that had independently segregated across the markers examined. This result is
parsimonious with a recombinant clade of strains that reflect genetic hybrids from a limited,
but distinct set of ancestries. Our results were further confirmed both by CGH array analyses
as well as whole genome sequencing. The previous misclassification as a clonal population
appears to be the result of the low resolution genotyping analyses performed, and the
placement of the markers in predominantly Type II regions of the genome [10, 14, 43]. For
this reason, future studies should use WGS to not only discriminate between haplogroups
within a population but also to determine the true genetic ancestry of each haplogroup.
Toxoplasma population genetics would also greatly benefit from an increase in the number
of isolates that are whole genome sequenced. WGS should be expanded to include both a
greater diversity of strains as well as multiple isolates from within a canonical clonal lineage.
The Type I and III clonal lineages are thought to be derived from a limited number of
sexual crosses, and all strains within these clonotypes share large haploblocks of Type II-like
sequence [11, 42, 44, 45]. This finding also holds true for the Type X lineage. At all loci
surveyed, Type X displayed at least one isolate with a Type II inherited allele, and the
introgression of large haploblocks of Type II ancestry across the genomes was confirmed in
the 19 strains that were resolved at WGS resolution. In regions that did not clade with Type
II, two different allelic types were identified. We concluded that these represented distinct
ancestries that we referred to as γ and δ. This designation was supported by the inherent
reticulation of the clade, seen in the NeighborNet tree (Figure 2B), which possessed short
branch lengths radiating from a reticulated network with multiple strains on a single branch,
rather than producing a star-like phylogeny with multiple alleles present at each major
branch, the result of private SNPs that accumulated through genetic drift. In fact, the lack of
genetic drift between these loci supported genetic hybridization as the most parsimonious
explanation for the relationship between the Type X haplotypes.

Sexual replication with genetic recombination is known to occur in South America at high
frequency [41, 44, 46]. Additionally, recent studies on the population genetic structure of
Toxoplasma has established that genetic hybridization, at whole genome resolution, is extant
for the vast majority of sequenced strains that represent the breadth of genetic diversity
within *Toxoplasma* [15, 38]. Evidence of sexual recombination across the population, as well
as the evidence of sexual replication within the Type X clade shown here, may indicate that
sexual recombination is occurring more frequently than previously suspected within North
America.
No barriers appear to exist, or have been described, to limit sexual recombination from
occurring in the laboratory setting, although there is a lack of understanding for the genetic
factors that regulate sexual replication in the felid host [7, 45, 47-49]. It is not clear why
sexual recombination does not appear to be occurring in North American wildlife, to the
degree that it does in South America, although it is possible that self-mating within the
population masks the detection of genetic outcrossing [12, 42, 50]. Especially among closely
related isolates, unisexual mating cannot be resolved using current genotyping methods that
rely on low-resolution analyses and thus cannot readily distinguish asexual expansion from
unisexual mating [12].
Toxoplasma has been shown previously to utilize its sexual cycle to expand its biological
potential and alter its pathogenesis [7, 47, 49]. Sexual recombination, by the ability to
reshuffle parental alleles into new combinations, can produce progeny with a vast range of
altered biological potentials [7]. Our work herein has identified a single *Toxoplasma*
genotype that was derived from a recent genetic cross that has been naturally selected in
southern sea otters across a space of 7 years. Importantly, this same genotype was highly
pathogenic to outbred laboratory mice, with all mice dying in 10-15 days with an inoculum
of just 50 tachyzoites. In contrast, the two Type X B haplotype isolates that caused subclinical
infection in outbred laboratory mice were highly pathogenic to sea otters that stranded with
a fatal meningoencephalitis [25]. Our data thus support a virulence shift model for the
expansion and propagation of *Toxoplasma* in nature, whereby *Toxoplasma* utilizes its sexual
cycle to generate a wide spectrum of novel genotypes that possess altered biological
potentials that can be naturally selected across *Toxoplasma*'s vast host range for optimal
transmission. Hence, natural selection among intermediate hosts allows *Toxoplasma* to
maintain highly pathogenic strains cryptically that can cause serious disease in some animal
hosts, but not in others. Examples of this type of selection have been observed previously in
laboratory mice that express variable IRG (Immunity Related GTPase) gene arrays. IRGs are
host proteins primarily responsible for combatting *Toxoplasma* lysis of murine cells and
Toxoplasma encodes a suite of highly polymorphic Rhoptry kinase genes (ROPKs) whose
primary function is to inactivate host IRGs [51-53]. While laboratory mice are highly clonal,
wild mice and their subsequent IRGs are more diverse, as are the ROPKs expressed by
different *Toxoplasma* strains that are capable of infecting wild mice [17, 54]. Similarly, TLR
11/12 is one of the primary rodent innate immune response sensors (otherwise referred to
as PAMPs) that detects infection by *Toxoplasma*. However, not all intermediate hosts share
a functional combination of TLR11/12, thus, *Toxoplasma* strains that are selected for
infection in mice need to bypass TLR11/12 recognition, whereas this is not a barrier to
infection in a human cell, for example [54, 55]. Thus, intermediate hosts are capable of
naturally selecting *Toxoplasma* strain genotypes that establish chronic, transmissible
infections in a host species, parasite strain-dependent manner. In this way, the parasite finds

a balance between its virulence, fitness and infectious transmissibility across its wide host range. It is this capacity to maintain cryptically virulent strains that may in part explain *Toxoplasma's* success in nature; it has the ability to cause disease outbreaks, expand its host range into new ecological niches, or alter its pathogenicity in both a parasite strain and host genetic manner to maximize its transmission.

In other protozoan parasites, when host partitioning occurs, it is synonymous with speciation. For example, *Plasmodium* parasites often partition by the mosquito host species they co-evolve with to maximize parasite transmission [56-60]. Likewise, the apicomplexan parasite *Sarcocystis* largely maintains separate species for each intermediate-definitive host species combination to maintain its life cycle [12, 61]. In contrast, *Toxoplasma* forms transmissible cysts in virtually all warm-blooded vertebrates, and these cysts are infectious for its definitive felid host, as well as a large array of intermediate hosts. What is different, is that across the genetic diversity of *Toxoplasma*, specific strain genotypes are being selectively expanded among its vast array of different intermediate hosts, based primarily on parasite genotype and the suite of polymorphic effector proteins encoded by a particular strain. It is this fact that allows *Toxoplasma* to maintain cryptically virulent strains across its broad intermediate host range. Hence, the intermediate host is playing a central role in the natural selection, expansion, and maintenance of virulent strains across the broad host range of this generalist parasite. In effect, infected hosts act as reservoirs for the pathogenic or epidemic potential of the species. Our study showed that one sub-type of *Toxoplasma* had expanded in the majority of sea otters because it caused primarily subclinical, chronic infections in sea otters, but that it was uniformly lethal to outbred laboratory mice. This is not unlike the expansion of strains bearing Type I alleles in birds that harbor asymptomatic infections but prove highly pathogenic to laboratory mice [62]. Additionally, while Type II isolates are commonly found infecting domestic livestock in North America, Type X is more common in sylvatic hosts, leading others to propose that separate cycles exist within the *Toxoplasma* population that overlap solely within the feline definitive host [10, 29, 63].

Finally, only a limited number of crossovers were detected among strains WGS sequenced within the recombinant clade of Type X haplotypes. Because the pedigree of one of the parents was known, we concluded that the isolates resembled F_1 progeny from a natural cross, which prompted us to perform a population-based QTL analysis on an unmanipulated, natural population of genetic hybrids. Taking advantage of a genome-wide, high resolution SNP map we had generated for all sequenced haplotypes, our analysis identified multiple punctate QTL peaks containing a limited number of candidate loci associated with differences in mouse virulence for reverse genetic follow-up. The serine-threonine protein kinase ROP33 stood out as the best candidate to influence pathogenicity, it was identified previously to be an active protein kinase that is both abundantly expressed and highly polymorphic. Further, ROP33 is a divergent WNG kinase (WNG-3) that is related to ROP35 (WNG-1), a critical regulator of tubular membrane biogenesis, the formation of the parasite intravesicular network (IVN) and the phosphorylation of dense granule proteins associated with IVN biogenesis [33]. In fact dense granule proteins GRA2 and GRA6 are required for IVN biogenesis, and IVN deficient parasites grow normally *in vitro*, but are attenuated in mouse virulence assays [64, 65]. Other related ROP proteins, including ROP 5, 16, 17, and 18 are known to hijack host immune signaling pathways to alter *Toxoplasma* virulence during rodent infection. However, all Type X isolates investigated in our study, whether mouse virulent or avirulent, only expressed avirulent allele combinations of the

known ROP 5/16/18 virulence factors, which is why we chose to knock-out the gene in a
Ku80 deficient strain of *Toxoplasma* that was both mouse virulent, and deficient in ROP18.
This is because no change in mouse virulence was reported when ROP33 was deleted in a
wild-type RH strain background [66]. In addition, previous genetic mapping studies failed to
identify the QTL for this locus, even though it is polymorphic between Type I, II and III strains
for which genetic crosses have been performed. It is likely that the ROP5 and ROP18
virulence factors were dominant, and ROP33 is analogous to ROP17, another virulence factor
that was only identified after the virulence enhancing capacity of ROP18 was removed [40].
Our results show that parasites deficient in ROP33 in the highly permissive RHΔku80Δ*rop18*
parent, were avirulent, whereas an infectious dose of 500 tachyzoites of the RHΔku80Δ*rop18*
parent genotype was uniformly virulent. Specifically how ROP33 contributes to murine
virulence, what dense granule proteins it phosphorylates, the integrity of the IVN and what
host immune signaling pathways are modified are being actively investigated. Future work
will dissect the host or parasite factors that ROP33 targets to reduce or influence parasite
transmission and pathogenicity. Our study establishes that it is possible to use natural
isolates derived from a population genetic cross in nature to identify genetic loci associated
with a specific quantitative phenotype.
659
660   **Materials and Methods**
661
662   **Parasite culture**
663   Fifty-three previously published *T. gondii* strains isolated from brain and/or tissue
samples from stranded Pacific coast southern sea otters were provided by Dr. Patricia
Conrad [25, 26]. *Toxoplasma* strains were isolated at the Marine Wildlife Veterinary Care and
Research Centre in Santa Cruz, CA, or at the University of California, Davis School of
Veterinary Medicine. Cause of death (COD) was determined for each sea otter based on
primary pathology reports from marine veterinarians at date of necropsy, as previously
published [25, 26, 31, 67]. *Toxoplasma* isolates each had an ATOS ("as the otter swims")
number assigned to them based on the stranding location where each sea otter was
discovered. Isolates were named sequentially based on the date the sea otter host stranded.
Parasites were cultured and maintained in human foreskin fibroblast (HFF) monolayers as
described previously [68].
674
675   **DNA extraction and genetic typing markers**
676   Parasites were syringe lysed using a 27-gauge needle and filtered through a 3.0
micron polycarbonate filter to remove cellular debris. DNA was extracted from cell pellets
using the Qiagen DNeasy Blood and Tissue kit (Qiagen). All 53 isolates were typed using 5
previously described PCR-RFLP and one DNA sequencing marker [7, 25, 69-71]. BSR4, BAG1,
and ROP1 were PCR amplified and typed based on known RFLP identity. GRA6 and SAG3
were typed according to previously identified representative SNPs based on Sanger
sequence data done by the National Institute of Allergy and Infectious Disease's Rocky
Mountain Laboratory (NIAID RML).
684   Alleles observed via sequencing were designated as canonical I, II, or III. When
sequences differed from canonical alleles, previously described methods of characterization
were used [29]. Briefly, when genetic variation differed from references by two mutations
the allele was categorized as unique, while only one mutation difference was labeled a drifted

allele from the closest reference sequence. When more than two nucleotide mutations were
present, or the sequence was as distant from the reference strains as references are from
each other, alleles were classified as novel alleles.
21 strains representing the 8 unique genotypes identified by the original 5 markers
were selected to characterize further. These 21 strains were PCR amplified, sequenced, and
genotyped using 20 markers: 18 at both linked and unlinked genomic loci, encompassing 13
of the 14 chromosomes, and 2 loci on organellar genomes (apicoplast and mitochondria).
15,430 bp with 335 SNPs (~0.024% of the genome). Sequences were examined and
nucleotides verified using SeqMan Pro alignment software (Lasergene). Sequences for
reference strains (ME49, GT1, and VEG) were downloaded from ToxoDB [72].
698
699 Generation of the *ROP33* knockout strain in *RHΔku80Δhxgprt* parasites
Deletion of the *ROP33* gene was generated using the Type I parasite RH that was
rendered deficient in the expression of *KU80* and the drug selectable marker *HXGPRT*.
Briefly, the CRISPR/Cas9 targeting plasmid pSAG1:CAS9,U6:sgUPRT (received from Prof. L.
David Sibley, Department of Molecular Microbiology, Washington University School of
Medicine) was modified with a guide RNA (GTCGGACGCGAAACTCGCTT) to target the 5'
region of the *ROP33* gene (sgROP33) using Q5 mutagenesis (New England Biolabs, MA). Then
a CRISPR/Cas9 replacement construct was created using Gibson assembly (New England
Biolabs) to stitch together a 1kb flanking region of the 5' UTR region of ROP33
(TGGT1_chrVIIa, 3921773 to 3922773 bp) with the selectable marker HXGPRT and a 1kb
flanking region of the 3'UTR region of ROP33 (TGGT1_chrVIIa, 3926189 to 3927189 bp) to
replace the *ROP33* gene using the selectable drug cassette. To delete the *ROP33* gene, a total
of 50 μg of the sgROP33 and replacement construct (5:1 ratio) were co-transfected into a
ROP18 deficient strain of RH (RH *Δku80Δhxgprtrop18*), which was cultured in human
foreskin fibroblasts (HFF) cells and maintained in Dulbecco's modified Eagle's medium
(DMEM) supplemented with 10% fetal bovine serum, 2 mM glutamine and 25 μg / ml
gentamicin. After transfection, parasites were selected for stable integration of the targeting
construct using mycophenolic acid (12.5 mg / mL in MeOH; MPA) / xanthine (20 mg / mL in
1M KOH; XAN) as described previously [73]. After 15 days, the resistant population was
cloned by limiting dilution and single clones were screened by PCR for targeted deletion of
the *ROP33* gene.
720
721 Mouse virulence assay
To determine the virulence of the sea otter isolates in a mouse model, groups of five
or more, 6-8 week-old, female CD1 outbred mice were intraperitoneally injected with 50 *T.*
gondii tachyzoites resuspended in 500 μl of PBS that had been expanded in HFF cells. Mice
were weighed daily to measure infection-induced cachexia and mouse survival was assayed
over 42 days [7]. At 14 days, mice were bled, and serum extracted to test for seroconversion
via indirect fluorescent antibody test (IFAT) against ME49 tachyzoites [74]. Strain virulence
was classified as follows: avirulent strains killed no seropositive mice within 42 days of
infection, intermediate virulent strains killed some but not all seropositive mice within 42
days post-infection, and virulent strains killed all seropositive mice. To assess the
contribution of ROP33 to mouse virulence, groups of five 6-8 week old female outbred CD-1
mice were injected intraperitoneally with 500 tachyzoites of either the parent

RHΔ*ku80*Δ*rop18*Δ*hxgprt* strain or the *Rop33-* mutant. Infectivity and virulence were assessed
as above.
735
Phylogenetic Tree and DNA Marker Analysis
The default settings of Clustal X were used to align DNA sequences for all markers
individually and as a concatenated set of markers [75]. MSF files of aligned marker sequences
were imported into Molecular Evolutionary Genetic Analysis (MEGA) Version 7 to create a
maximum likelihood tree using Tamura-Nei model distance analysis with uniform rates of
substitution across all sites, and 1000 bootstrap support for all branch points [76].
Consensus trees were rooted on ME49, as Type II is an inherent parent in all strains analyzed.
All scales are by 0.001 nucleotide difference unit. Distinct parental lineages with bootstrap
support over 60% are indicated on each tree to distinguish II alleles from novel γ and δ
alleles. These novel alleles are indicated in purple and orange respectively on the 17 loci
MLST in Supplementary Table 2. The Nexus file of the concatenated marker alignment was
imported into SplitsTree 4 Version 4.13.1 where NetworkNet analysis was run using default
settings [77].
749
eBURST Clonal Complex Analysis
Alleles determined from the 17 nuclear encoded markers (excluding ROP1 due to its
classification as a microsatellite marker) were given numerical designations to create a
multilocus sequence-typing scheme for the eBURST program [78]. Default settings were
used to evaluate the isolated strains' alleles. Strains were displayed as dots and colors of dots
were depicted accordingly: Type I, II, III, and X being red, green, blue, and purple
respectively. Dot size was representative of the number of strains contained within each
genotype, larger dots representing more strains bearing the same genotype. Strains sharing
16 of the 17 nuclear encoded markers were designated as clonal complexes and represented
by connecting lines in the diagram shown.
760
Genomic hybridization to Affymetrix arrays
Genomic DNA was sheared and biotin labeled before hybridizing to a custom *T. gondii*
Affymetrix microarray as previously published [37]. High-fidelity SNPs were characterized
via a custom R script to identify SNPs belonging to each of the three reference strains (I, II,
and III). Three reference strains (GT1, ME49, and CTG) are shown to demonstrate ideal
hybridization within canonical lineages. Sea otter Type X DNA isolate hybridizations are
shown below the reference strains.
768
Whole-Genome Sequence Analysis of Type X Recombination
DNA from 16 isolates that grew best in tissue culture as well as 3 representative Type
X strains (ARI, RAY, and WTD1), for a total of 19 Type X strains was isolated. 3 µg of DNA
isolated from each of these 19 strains was sent to NIAID RML for whole-genome sequencing
using Illumina HiSeq technology. Fastq reads were reference mapped to the *T. gondii* ME49
assemblage Version 8.2 [72] using BWA 0.7.5a to align the reads to the reference genome
and GATK 3.7 in coordination with Picard 1.131 following best practices to quality control
the mapped reads [80]. Following mapping, the gVCF method of GATK was used to combine
SNP calls using stand_call_conf of 30.0, nct of 10, and ploidy of 1 (for haploid genomes) to

call 568,592 single nucleotide polymorphism positions across the whole genomes of these
strains [80]. The derived VCF formatted SNP file was curated using GATK and VCFTools to
produce a tabular file containing only biallelic SNPs with no large insertions or deletions
[81]. A custom script was then utilized to convert this SNP file into a fasta file of strain
polymorphic positions across the Type X and reference genomes. These SNP fasta files were
run in SplitsTree4 using default parameters for BioNJ with 1000 bootstrap support to create
a NeighborNet tree [72]. Interconnected reticulation between strains is indicative of
recombination between strains while mitotic drift was visualized by divergent branching.
786
SNP Density Fingerprint Analysis of Recombinant Progeny
The same tabular, biallelic SNP file used in the NeighborNet analysis was modified
using a custom R script based on location mapping to isolate strain-specific polymorphic
locations from the tabular VCF. Further R scripts group these SNPs into 100 kb windows
which are mapped across the genomes of these strains to display where diversity from the
reference genome (ME49) is most apparent [44, 82]. The larger the number of SNPs in a
particular window, the more divergent the strain is from the ME49 reference. Distinct
haploblocks where genomic recombination has occurred are apparent in areas where
haploblock diversity significantly varies from the surrounding regions on the same
chromosome.
797
PopNet Characterization of Strain Interrelatedness
The tabular, biallelic SNP file created from WGS of Type X and used to derive the SNP
density plots and NeighborNet tree was uploaded into PopNet using default parameters to
assess the diversity and interrelatedness of strains [38]. Cytoscape was used to visualize the
recombination and Markov clustering outputs. Genomes are displayed in circularized format
with chromosomes concatenated into a circular genome display. The background of each
strain is painted to match the group that shares the most common ancestry over the entire
genome with the strain shown. Chromosome painting was done in 10 kb increments and the
sequence haploblock was painted based on its shared ancestry. For instance, a Type X isolate
haploblock that was most closely related to the ancestral Type II strain was painted green to
indicate its inheritance. Strains that are more closely related have thicker connecting lines
between the circles of the strain isolates.
810
Virulence Gene Identification by QTL
To find novel virulence alleles, the tabular, biallelic SNP file was utilized to run a
quantitative trait locus (QTL) analysis on the Type X and Type II isolates of *Toxoplasma*
sequenced at WGS resolution [83]. SNP calls were down-selected by a custom java script to
include one SNP every 5 kb. This allows the QTL software to analyze the breadth of the
WGS data in QTL, which was built to handle marker typing data without the depth inherent
in WGS data. Custom scripts code the SNPs into reference (ME49) versus alternative (Type
X's γ/δ lineage which substitute as the secondary parent) alleles. This down-selected
marker dataset was combined with previously determined low-dose murine virulence for
these strains. This data was then input into J/qtl and a one QTL genome scan was run using
the default settings for the EM algorithm (maximum likelihood) with 1000 permutations to

identify genomic locations significantly associated with the percentage of murine death due to Type X infection [84].

The QTL calculations identified four significantly associated regions based on a LOD score of greater than 4.0. Within these genomic regions, 450 genes were predicted based on ToxoDB documentation. These potential genes were down-selected based on presence of a signal peptide and/or transmembrane domain, gene expression, polymorphism, and genome-wide CRISPR score for essentiality to identify 32 virulence candidate genes within the Type X strains. Of these, ROP33 was selected for further interrogation based on its high LOD score and similarity to previously identified virulence effector proteins.

Funding statement:

This work was supported in part by the Intramural Research Program of the National Institute of Allergy and Infectious Diseases (NIAID) at the National Institutes of Health and by the National Science Foundation, Ecology of Infectious Diseases Grant 052576.

Ethics statements:

The animal study protocol LPD 22E was reviewed and approved by the Animal Care and Use Committee of the Intramural Research Program of the National Institute of Allergy and Infectious Diseases, National Institutes of Health. No human studies or identifiable human data are presented in this study. All datasets generated for this study are available upon request to the corresponding author.

Acknowledgements

We thank Andrea Packham and Ann Melli for their assistance in culturing and isolation of the *Toxoplasma* isolates, and Javi Zhang for his help with PopNet analysis. We also thank all volunteers and staff at the California Department of Fish and Wildlife, the Monterey Bay Aquarium, the Marine Mammal Center, and the United States Geological Survey for their efforts to recover stranded sea otters that facilitated the isolation of *Toxoplasma gondii* strains utilized in this paper.

References

1. Vijaykrishna D, Smith GJ, Pybus OG, Zhu H, Bhatt S, Poon LL, et al. Long-term evolution and transmission dynamics of swine influenza A virus. Nature. 2011;473(7348):519-22. doi: 10.1038/nature10004. PubMed PMID: 21614079.

2. Bellanger X, Payot S, Leblond-Bourget N, Guedon G. Conjugative and mobilizable genomic islands in bacteria: evolution and diversity. FEMS Microbiol Rev. 2014;38(4):720-60. doi: 10.1111/1574-6976.12058. PubMed PMID: 24372381.

3. Schneider G, Dobrindt U, Middendorf B, Hochhut B, Szijarto V, Emody L, et al. Mobilisation and remobilisation of a large archetypal pathogenicity island of uropathogenic *Escherichia coli* in vitro support the role of conjugation for horizontal transfer of genomic islands. BMC Microbiol. 2011;11:210. doi: 10.1186/1471-2180-11-210. PubMed PMID: 21943043; PubMed Central PMCID: PMC3202238.

4. Wardal E, Markowska K, Zabicka D, Wroblewska M, Giemza M, Mik E, et al. Molecular analysis of vanA outbreak of *Enterococcus faecium* in two Warsaw hospitals: the importance of mobile genetic elements. Biomed Res Int. 2014;2014:575367. doi:

10.1155/2014/575367. PubMed PMID: 25003118; PubMed Central PMCID:
PMC4070583.

5. English ED, Adomako-Ankomah Y, Boyle JP. Secreted effectors in *Toxoplasma gondii* and
related species: determinants of host range and pathogenesis? Parasite immunology.
2015;37(3):127-40. doi: 10.1111/pim.12166. PubMed PMID: 25655311.

6. Fraser JA, Giles SS, Wenink EC, Geunes-Boyer SG, Wright JR, Diezmann S, et al. Same-sex
mating and the origin of the Vancouver Island *Cryptococcus gattii* outbreak. Nature.
2005;437(7063):1360-4. Epub 2005/10/14. doi: 10.1038/nature04220. PubMed
PMID: 16222245.

7. Grigg ME, Bonnefoy S, Hehl AB, Suzuki Y, Boothroyd JC. Success and virulence in
Toxoplasma as the result of sexual recombination between two distinct ancestries.
Science. 2001;294(5540):161-5. doi: DOI 10.1126/science.1061888. PubMed PMID:
WOS:000171448800052.

8. Boothroyd JC, Grigg ME. Population biology of *Toxoplasma gondii* and its relevance to
human infection: do different strains cause different disease? Curr Opin Microbiol. 5.
England2002. p. 438-42.

9. Boyer K, Hill D, Mui E, Wroblewski K, Karrison T, Dubey JP, et al. Unrecognized ingestion
of *Toxoplasma gondii* oocysts leads to congenital toxoplasmosis and causes epidemics in
North America. Clin Infect Dis. 2011;53(11):1081-9. Epub 2011/10/25. doi:
10.1093/cid/cir667. PubMed PMID: 22021924; PubMed Central PMCID:
PMCPMC3246875.

10. Behnke MS, Khan A, Wootton JC, Dubey JP, Tang K, Sibley LD. Virulence differences in
Toxoplasma mediated by amplification of a family of polymorphic pseudokinases. Proc
Natl Acad Sci U S A. 2011;108(23):9631-6. doi: 10.1073/pnas.1015338108. PubMed
PMID: 21586633; PubMed Central PMCID: PMC3111276.

11. Boyle JP, Rajasekar B, Saeij JP, Ajioka JW, Berriman M, Paulsen I, et al. Just one cross
appears capable of dramatically altering the population biology of a eukaryotic
pathogen like *Toxoplasma gondii*. Proc Natl Acad Sci U S A. 2006;103(27):10514-9. doi:
10.1073/pnas.0510319103. PubMed PMID: 16801557; PubMed Central PMCID:
PMC1502489.

12. Wendte JM, Miller MA, Lambourn DM, Magargal SL, Jessup DA, Grigg ME. Self-mating in
the definitive host potentiates clonal outbreaks of the apicomplexan parasites
Sarcocystis neurona and *Toxoplasma gondii*. PLoS Genetics. 2010;6(12):e1001261. doi:
10.1371/journal.pgen.1001261. PubMed PMID: 21203443; PubMed Central PMCID:
PMC3009688.

13. Boothroyd JC. Expansion of host range as a driving force in the evolution of *Toxoplasma*.
Mem I Oswaldo Cruz. 2009;104(2):179-84. PubMed PMID: WOS:000267051500009.

14. Howe DK, Sibley LD. *Toxoplasma gondii* comprises three clonal lineages: correlation of
parasite genotype with human disease. J Infect Dis. 1995;172(6):1561-6. PubMed PMID:
7594717.

15. Lorenzi H, Khan A, Behnke MS, Namasivayam S, Swapna LS, Hadjithomas M, et al. Local
admixture of amplified and diversified secreted pathogenesis determinants shapes
mosaic *Toxoplasma gondii* genomes. Nat Commun. 2016;7:10147. doi:
10.1038/ncomms10147. PubMed PMID: 26738725; PubMed Central PMCID:
PMCPMC4729833.

16. Sibley LD, Boothroyd JC. Virulent strains of *Toxoplasma gondii* comprise a single clonal
lineage. Nature. 1992;359(6390):82-5. doi: 10.1038/359082a0. PubMed PMID:
1355855.
17. Lilue J, Muller UB, Steinfeldt T, Howard JC. Reciprocal virulence and resistance
polymorphism in the relationship between *Toxoplasma gondii* and the house mouse.
Elife. 2013;2:e01298. doi: 10.7554/eLife.01298. PubMed PMID: 24175088; PubMed
Central PMCID: PMC3810784.
18. Steinfeldt T, Konen-Waisman S, Tong L, Pawlowski N, Lamkemeyer T, Sibley LD, et al.
Phosphorylation of mouse immunity-related GTPase (IRG) resistance proteins is an
evasion strategy for virulent *Toxoplasma gondii*. PLoS Biology. 2010;8(12):e1000576.
Epub 2011/01/05. doi: 10.1371/journal.pbio.1000576. PubMed PMID: 21203588;
PubMed Central PMCID: PMCPMC3006384.
19. Cavailles P, Sergent V, Bisanz C, Papapietro O, Colacios C, Mas M, et al. The rat *Toxo1*
locus directs toxoplasmosis outcome and controls parasite proliferation and spreading
by macrophage-dependent mechanisms. Proc Natl Acad Sci U S A. 2006;103(3):744-9.
Epub 2006/01/13. doi: 10.1073/pnas.0506643103. PubMed PMID: 16407112; PubMed
Central PMCID: PMCPMC1334643.
20. Cirelli KM, Gorfu G, Hassan MA, Printz M, Crown D, Leppla SH, et al. Inflammasome
sensor NLRP1 controls rat macrophage susceptibility to *Toxoplasma gondii.* PLoS
Pathog. 2014;10(3):e1003927. Epub 2014/03/15. doi: 10.1371/journal.ppat.1003927.
PubMed PMID: 24626226; PubMed Central PMCID: PMCPMC3953412.
21. Ewald SE, Chavarria-Smith J, Boothroyd JC. NLRP1 is an inflammasome sensor for
Toxoplasma gondii. Infection and Immunity. 2014;82(1):460-8. Epub 2013/11/13. doi:
10.1128/IAI.01170-13. PubMed PMID: 24218483; PubMed Central PMCID:
PMCPMC3911858.
22. Gorfu G, Cirelli KM, Melo MB, Mayer-Barber K, Crown D, Koller BH, et al. Dual role for
inflammasome sensors NLRP1 and NLRP3 in murine resistance to *Toxoplasma gondii*.
mBio. 2014;5(1). Epub 2014/02/20. doi: 10.1128/mBio.01117-13. PubMed PMID:
24549849; PubMed Central PMCID: PMCPMC3944820.
23. Dubey JP, Shen SK, Kwok OC, Frenkel JK. Infection and immunity with the RH strain of
Toxoplasma gondii in rats and mice. The Journal of Parasitology. 1999;85(4):657-62.
PubMed PMID: 10461945.
24. Conrad PA, Miller MA, Kreuder C, James ER, Mazet J, Dabritz H, et al. Transmission of
Toxoplasma: clues from the study of sea otters as sentinels of *Toxoplasma gondii* flow
into the marine environment. Int J Parasitol. 2005;35(11-12):1155-68. Epub
2005/09/15. doi: 10.1016/j.ijpara.2005.07.002. PubMed PMID: 16157341.
25. Miller MA, Grigg ME, Kreuder C, James ER, Melli AC, Crosbie PR, et al. An unusual
genotype of *Toxoplasma gondii* is common in California sea otters (*Enhydra lutris nereis*)
and is a cause of mortality. Int J Parasitol. 2004;34(3):275-84. doi:
10.1016/j.ijpara.2003.12.008. PubMed PMID: 15003489.
26. Shapiro K, VanWormer E, Packham A, Dodd E, Conrad PA, Miller M. Type X strains of
Toxoplasma gondii are virulent for southern sea otters (*Enhydra lutris nereis*) and
present in felids from nearby watersheds. Proc Biol Sci. 2019;286(1909):20191334.
Epub 2019/08/23. doi: 10.1098/rspb.2019.1334. PubMed PMID: 31431162; PubMed
Central PMCID: PMCPMC6732395.

27. Khan A, Dubey JP, Su C, Ajioka JW, Rosenthal BM, Sibley LD. Genetic analyses of atypical
Toxoplasma gondii strains reveal a fourth clonal lineage in North America. Int J Parasitol.
2011;41(6):645-55. doi: 10.1016/j.ijpara.2011.01.005. PubMed PMID: 21320505;
PubMed Central PMCID: PMC3081397.

28. Dubey JP, Velmurugan GV, Rajendran C, Yabsley MJ, Thomas NJ, Beckmen KB, et al.
Genetic characterisation of *Toxoplasma gondii* in wildlife from North America revealed
widespread and high prevalence of the fourth clonal type. Int J Parasitol.
2011;41(11):1139-47. doi: 10.1016/j.ijpara.2011.06.005. PubMed PMID: 21802422.

29. VanWormer E, Miller MA, Conrad PA, Grigg ME, Rejmanek D, Carpenter TE, et al. Using
molecular epidemiology to track *Toxoplasma gondii* from terrestrial carnivores to
marine hosts: implications for public health and conservation. PLoS Neglected Tropical
Diseases. 2014;8(5):e2852. doi: 10.1371/journal.pntd.0002852. PubMed PMID:
24874796; PubMed Central PMCID: PMC4038486.

30. Gibson AK, Raverty S, Lambourn DM, Huggins J, Magargal SL, Grigg ME. Polyparasitism
is associated with increased disease severity in *Toxoplasma gondii*-infected marine
sentinel species. PLoS Neglected Tropical Diseases. 2011;5(5):e1142. doi:
10.1371/journal.pntd.0001142. PubMed PMID: 21629726; PubMed Central PMCID:
PMC3101184.

31. Kreuder C, Miller MA, Jessup DA, Lowenstine LJ, Harris MD, Ames JA, et al. Patterns of
mortality in southern sea otters (*Enhydra lutris nereis*) from 1998-2001. J Wildl Dis.
2003;39(3):495-509. doi: 10.7589/0090-3558-39.3.495. PubMed PMID: 14567210.

32. Sundar N, Cole RA, Thomas NJ, Majumdar D, Dubey JP, Su C. Genetic diversity among sea
otter isolates of *Toxoplasma gondii.* Veterinary Parasitology. 2008;151(2-4):125-32. doi:
10.1016/j.vetpar.2007.11.012. PubMed PMID: 18155841.

33. Beraki T, Hu X, Broncel M, Young JC, O'Shaughnessy WJ, Borek D, et al. Divergent kinase
regulates membrane ultrastructure of the *Toxoplasma* parasitophorous vacuole. Proc
Natl Acad Sci U S A. 2019;116(13):6361-70. Epub 2019/03/10. doi:
10.1073/pnas.1816161116. PubMed PMID: 30850550; PubMed Central PMCID:
PMCPMC6442604.

34. Miller MA, Miller WA, Conrad PA, James ER, Melli AC, Leutenegger CM, et al. Type X
Toxoplasma gondii in a wild mussel and terrestrial carnivores from coastal California:
new linkages between terrestrial mammals, runoff and toxoplasmosis of sea otters. Int
J Parasitol. 2008;38(11):1319-28. doi: 10.1016/j.ijpara.2008.02.005. PubMed PMID:
18452923.

35. Saeij JP, Boyle JP, Boothroyd JC. Differences among the three major strains of
Toxoplasma gondii and their specific interactions with the infected host. Trends in
Parasitology. 2005;21(10):476-81. doi: 10.1016/j.pt.2005.08.001. PubMed PMID:
16098810.

36. Su C, Khan A, Zhou P, Majumdar D, Ajzenberg D, Darde ML, et al. Globally diverse
Toxoplasma gondii isolates comprise six major clades originating from a small number
of distinct ancestral lineages. Proc Natl Acad Sci U S A. 2012;109(15):5844-9. doi:
10.1073/pnas.1203190109. PubMed PMID: 22431627; PubMed Central PMCID:
PMCPMC3326454.

37. Khan A, Miller N, Roos DS, Dubey JP, Ajzenberg D, Darde ML, et al. A monomorphic
haplotype of chromosome Ia is associated with widespread success in clonal and
nonclonal populations of *Toxoplasma gondii*. mBio. 2011;2(6):e00228-11. doi:

10.1128/mBio.00228-11. PubMed PMID: 22068979; PubMed Central PMCID:
PMC3215432.

38. Zhang J, Khan A, Kennard A, Grigg ME, Parkinson J. PopNet: A Markov Clustering
Approach to Study Population Genetic Structure. Mol Biol Evol. 2017;34(7):1799-811.
Epub 2017/04/07. doi: 10.1093/molbev/msx110. PubMed PMID: 28383661; PubMed
Central PMCID: PMCPMC5850731.

39. Sidik SM, Huet D, Ganesan SM, Huynh MH, Wang T, Nasamu AS, et al. A Genome-wide
CRISPR Screen in *Toxoplasma* Identifies Essential Apicomplexan Genes. Cell.
2016;166(6):1423-35 e12. Epub 2016/09/07. doi: 10.1016/j.cell.2016.08.019. PubMed
PMID: 27594426; PubMed Central PMCID: PMCPMC5017925.

40. Etheridge RD, Alaganan A, Tang K, Lou HJ, Turk BE, Sibley LD. The *Toxoplasma*
pseudokinase ROP5 forms complexes with ROP18 and ROP17 kinases that synergize to
control acute virulence in mice. Cell Host & Microbe. 2014;15(5):537-50. Epub
2014/05/17. doi: 10.1016/j.chom.2014.04.002. PubMed PMID: 24832449; PubMed
Central PMCID: PMCPMC4086214.

41. Khan A, Fux B, Su C, Dubey JP, Darde ML, Ajioka JW, et al. Recent transcontinental sweep
of *Toxoplasma gondii* driven by a single monomorphic chromosome. Proc Natl Acad Sci
U S A. 2007;104(37):14872-7. Epub 2007/09/07. doi: 10.1073/pnas.0702356104.
PubMed PMID: 17804804; PubMed Central PMCID: PMCPMC1965483.

42. Grigg ME, Sundar N. Sexual recombination punctuated by outbreaks and clonal
expansions predicts *Toxoplasma gondii* population genetics. Int J Parasitol.
2009;39(8):925-33. doi: 10.1016/j.ijpara.2009.02.005. PubMed PMID: 19217909;
PubMed Central PMCID: PMC2713429.

43. Ajzenberg D, Cogne N, Paris L, Bessieres MH, Thulliez P, Filisetti D, et al. Genotype of 86
Toxoplasma gondii isolates associated with human congenital toxoplasmosis, and
correlation with clinical findings. J Infect Dis. 2002;186(5):684-9. Epub 2002/08/27.
doi: 10.1086/342663. PubMed PMID: 12195356.

44. Minot S, Melo MB, Li F, Lu D, Niedelman W, Levine SS, et al. Admixture and
recombination among *Toxoplasma gondii* lineages explain global genome diversity. P
Natl Acad Sci USA. 2012;109(33):13458-63. doi: DOI 10.1073/pnas.1117047109.
PubMed PMID: WOS:000307807000067; PubMed Central PMCID: PMCPMC3421188.

45. Khan A, Shaik JS, Behnke M, Wang QL, Dubey JP, Lorenzi HA, et al. NextGen sequencing
reveals short double crossovers contribute disproportionately to genetic diversity in
Toxoplasma gondii. Bmc Genomics. 2014;15. doi: Artn 1168
10.1186/1471-2164-15-1168. PubMed PMID: WOS:000349049000003.

46. Rajendran C, Su C, Dubey JP. Molecular genotyping of *Toxoplasma gondii* from Central
and South America revealed high diversity within and between populations. Infection,
Genetics and Evolution : journal of molecular epidemiology and evolutionary genetics in
infectious diseases. 2012;12(2):359-68. Epub 2012/01/10. doi:
10.1016/j.meegid.2011.12.010. PubMed PMID: 22226702.

47. Saeij JP, Boyle JP, Coller S, Taylor S, Sibley LD, Brooke-Powell ET, et al. Polymorphic
secreted kinases are key virulence factors in toxoplasmosis. Science.
2006;314(5806):1780-3. doi: 10.1126/science.1133690. PubMed PMID: 17170306;
PubMed Central PMCID: PMC2646183.

48. Saeij JP, Coller S, Boyle JP, Jerome ME, White MW, Boothroyd JC. *Toxoplasma* co-opts host
gene expression by injection of a polymorphic kinase homologue. Nature.

2007;445(7125):324-7. doi: 10.1038/nature05395. PubMed PMID: 17183270; PubMed
Central PMCID: PMC2637441.

49. Taylor S, Barragan A, Su C, Fux B, Fentress SJ, Tang K, et al. A secreted serine-threonine
kinase determines virulence in the eukaryotic pathogen *Toxoplasma gondii.* Science.
2006;314(5806):1776-80. doi: 10.1126/science.1133643. PubMed PMID: 17170305.

50. Ferguson D. *Toxoplasma gondii* and sex: essential or optional extra? Trends in
Parasitology. 2002;18(8):351. Epub 2002/10/16. PubMed PMID: 12377284.

51. Fleckenstein MC, Reese ML, Konen-Waisman S, Boothroyd JC, Howard JC, Steinfeldt T. A
Toxoplasma gondii pseudokinase inhibits host IRG resistance proteins. PLoS Biology.
2012;10(7):e1001358. doi: 10.1371/journal.pbio.1001358. PubMed PMID: 22802726;
PubMed Central PMCID: PMC3393671.

52. Hunn JP, Feng CG, Sher A, Howard JC. The immunity-related GTPases in mammals: a fast-
evolving cell-autonomous resistance system against intracellular pathogens. Mamm
Genome. 2011;22(1-2):43-54. doi: 10.1007/s00335-010-9293-3. PubMed PMID:
21052678; PubMed Central PMCID: PMC3438224.

53. Yamamoto M, Okuyama M, Ma JS, Kimura T, Kamiyama N, Saiga H, et al. A cluster of
interferon-gamma-inducible p65 GTPases plays a critical role in host defense against
Toxoplasma gondii. Immunity. 2012;37(2):302-13. doi: 10.1016/j.immuni.2012.06.009.
PubMed PMID: 22795875.

54. Gazzinelli RT, Mendonca-Neto R, Lilue J, Howard J, Sher A. Innate Resistance against
Toxoplasma gondii: An Evolutionary Tale of Mice, Cats, and Men. Cell Host & Microbe.
2014;15(2):132-8. Epub 2014/02/18. doi: 10.1016/j.chom.2014.01.004. PubMed
PMID: 24528860.

55. Koblansky AA, Jankovic D, Oh H, Hieny S, Sungnak W, Mathur R, et al. Recognition of
profilin by Toll-like receptor 12 is critical for host resistance to *Toxoplasma gondii.*
Immunity. 2013;38(1):119-30. doi: 10.1016/j.immuni.2012.09.016. PubMed PMID:
23246311; PubMed Central PMCID: PMC3601573.

56. Mackinnon MJ, Read AF. Virulence in malaria: an evolutionary viewpoint. Philosophical
transactions of the Royal Society of London Series B, Biological sciences.
2004;359(1446):965-86. doi: 10.1098/rstb.2003.1414. PubMed PMID: 15306410;
PubMed Central PMCID: PMC1693375.

57. Ariey F, Witkowski B, Amaratunga C, Beghain J, Langlois AC, Khim N, et al. A molecular
marker of artemisinin-resistant *Plasmodium falciparum* malaria. Nature.
2014;505(7481):50-5. Epub 2013/12/20. doi: 10.1038/nature12876. PubMed PMID:
24352242.

58. Bopp SE, Manary MJ, Bright AT, Johnston GL, Dharia NV, Luna FL, et al. Mitotic evolution
of *Plasmodium falciparum* shows a stable core genome but recombination in antigen
families. PLoS Genetics. 2013;9(2):e1003293. doi: 10.1371/journal.pgen.1003293.
PubMed PMID: 23408914; PubMed Central PMCID: PMC3567157.

59. Miotto O, Almagro-Garcia J, Manske M, Macinnis B, Campino S, Rockett KA, et al. Multiple
populations of artemisinin-resistant *Plasmodium falciparum* in Cambodia. Nature
Genetics. 2013;45(6):648-55. Epub 2013/04/30. doi: 10.1038/ng.2624. PubMed PMID:
23624527; PubMed Central PMCID: PMC3807790.

60. Arisue N, Hashimoto T. Phylogeny and evolution of apicoplasts and apicomplexan
parasites. Parasitology International. 2014. doi: 10.1016/j.parint.2014.10.005. PubMed
PMID: 25451217.

61. Barbosa L, Johnson CK, Lambourn DM, Gibson AK, Haman KH, Huggins JL, et al. A novel
Sarcocystis neurona genotype XIII is associated with severe encephalitis in an
unexpectedly broad range of marine mammals from the northeastern Pacific Ocean. Int
J Parasitol. 2015. doi: 10.1016/j.ijpara.2015.02.013. PubMed PMID: 25997588.
62. Dubey JP, Graham DH, da Silva DS, Lehmann T, Bahia-Oliveira LM. *Toxoplasma gondii*
isolates of free-ranging chickens from Rio de Janeiro, Brazil: mouse mortality, genotype,
and oocyst shedding by cats. The Journal of Parasitology. 2003;89(4):851-3. Epub
2003/10/10. doi: 10.1645/GE-60R. PubMed PMID: 14533703.
63. Wendte JM, Gibson AK, Grigg ME. Population genetics of *Toxoplasma gondii*: new
perspectives from parasite genotypes in wildlife. Veterinary Parasitology.
2011;182(1):96-111. doi: 10.1016/j.vetpar.2011.07.018. PubMed PMID: 21824730.
64. Labruyere E, Lingnau M, Mercier C, Sibley LD. Differential membrane targeting of the
secretory proteins GRA4 and GRA6 within the parasitophorous vacuole formed by
Toxoplasma gondii. Mol Biochem Parasitol. 1999;102(2):311-24. Epub 1999/09/25. doi:
10.1016/s0166-6851(99)00092-4. PubMed PMID: 10498186.
65. Mercier C, Howe DK, Mordue D, Lingnau M, Sibley LD. Targeted disruption of the GRA2
locus in *Toxoplasma gondii* decreases acute virulence in mice. Infection and Immunity.
1998;66(9):4176-82. Epub 1998/08/26. PubMed PMID: 9712765.
66. Wang JL, Li TT, Elsheikha HM, Chen K, Zhu WN, Yue DM, et al. Functional
Characterization of Rhoptry Kinome in the Virulent *Toxoplasma gondii* RH Strain. Front
Microbiol. 2017;8:84. Epub 2017/02/09. doi: 10.3389/fmicb.2017.00084. PubMed
PMID: 28174572; PubMed Central PMCID: PMCPMC5258691.
67. Miller MA, Gardner IA, Packham A, Mazet JK, Hanni KD, Jessup D, et al. Evaluation of an
Indirect Fluorescent Antibody Test (Ifat) for Demonstration of Antibodies to
Toxoplasma Gondii in the Sea Otter (*Enhydra Lutris*). Journal of Parasitology.
2002;88(3):594-9. doi: 10.1645/0022-3395(2002)088[0594:eoaifa]2.0.co;2.
68. Pszenny V, Angel SO, Duschak VG, Paulino M, Ledesma B, Yabo MI, et al. Molecular
cloning, sequencing and expression of a serine proteinase inhibitor gene from
Toxoplasma gondii. Mol Biochem Parasitol. 2000;107(2):241-9. PubMed PMID:
10779600.
69. Su C, Zhang X, Dubey JP. Genotyping of *Toxoplasma gondii* by multilocus PCR-RFLP
markers: a high resolution and simple method for identification of parasites. Int J
Parasitol. 2006;36(7):841-8. doi: 10.1016/j.ijpara.2006.03.003. PubMed PMID:
16643922.
70. Fazaeli A, Carter PE, Darde ML, Pennington TH. Molecular typing of *Toxoplasma gondii*
strains by GRA6 gene sequence analysis. Int J Parasitol. 2000;30(5):637-42. PubMed
PMID: 10779578.
71. Howe DK, Sibley LD. *Toxoplasma gondii*: analysis of different laboratory stocks of the RH
strain reveals genetic heterogeneity. Exp Parasitol. 1994;78(2):242-5. doi:
10.1006/expr.1994.1024. PubMed PMID: 7907030.
72. Gajria B, Bahl A, Brestelli J, Dommer J, Fischer S, Gao X, et al. ToxoDB: an integrated
Toxoplasma gondii database resource. Nucleic Acids Res. 2008;36(Database
issue):D553-6. doi: 10.1093/nar/gkm981. PubMed PMID: 18003657.
73. Donald RG, Carter D, Ullman B, Roos DS. Insertional tagging, cloning, and expression of
the *Toxoplasma gondii* hypoxanthine-xanthine-guanine phosphoribosyltransferase
gene. Use as a selectable marker for stable transformation. J Biol Chem.

1996;271(24):14010-9. Epub 1996/06/14. doi: 10.1074/jbc.271.24.14010. PubMed
PMID: 8662859.
74. Fletcher S. Indirect Fluorescent Antibody Technique in the Serology of *Toxoplasma*
gondii. J Clin Pathol. 1965;18:193-9. Epub 1965/03/01. PubMed PMID: 14276154.
75. Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, McWilliam H, et al.
Clustal W and Clustal X version 2.0. Bioinformatics. 2007;23(21):2947-8. doi:
10.1093/bioinformatics/btm404. PubMed PMID: 17846036.
76. Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. MEGA6: Molecular Evolutionary
Genetics Analysis version 6.0. Mol Biol Evol. 2013;30(12):2725-9. doi:
10.1093/molbev/mst197. PubMed PMID: 24132122.
77. Huson DH, Bryant D. Application of phylogenetic networks in evolutionary studies. Mol
Biol Evol. 2006;23(2):254-67. doi: 10.1093/molbev/msj030. PubMed PMID: 16221896.
78. Feil EJ, Li BC, Aanensen DM, Hanage WP, Spratt BG. eBURST: Inferring Patterns of
Evolutionary Descent among Clusters of Related Bacterial Genotypes from Multilocus
Sequence Typing Data. Journal of Bacteriology. 2004;186(5):1518-30. doi:
10.1128/jb.186.5.1518-1530.2004.
79. Rozas J, Sanchez-DelBarrio JC, Messeguer X, Rozas R. DnaSP, DNA polymorphism
analyses by the coalescent and other methods. Bioinformatics. 2003;19(18):2496-7. doi:
10.1093/bioinformatics/btg359.
80. Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, Del Angel G, Levy-Moonshine A, et
al. From FastQ data to high confidence variant calls: the Genome Analysis Toolkit best
practices pipeline. Curr Protoc Bioinformatics. 2013;11(1110):11 0 1- 0 33. doi:
10.1002/0471250953.bi1110s43. PubMed PMID: 25431634.
81. Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., Genomes
Project Analysis. (2011). The variant call format and VCFtools. *Bioinformatics, 27*(15),
2156-2158. doi:10.1093/bioinformatics/btr330.
82. Yin, T., Cook, D., & Lawrence, M. (2012). ggbio: an R package for extending the grammar
of graphics for genomic data. *Genome Biol, 13*(8), R77. doi:10.1186/gb-2012-13-8-r77
83. Behnke, M. S., Khan, A., Lauron, E. J., Jimah, J. R., Wang, Q., Tolia, N. H., & Sibley, L. D.
(2015). Rhoptry Proteins ROP5 and ROP18 Are Major Murine Virulence Factors in
Genetically Divergent South American Strains of Toxoplasma gondii. *PLoS Genet, 11*(8),
e1005434. doi:10.1371/journal.pgen.1005434.
84. Smith, R., Sheppard, K., DiPetrillo, K., & Churchill, G. (2009). Quantitative trait locus
analysis using J/qtl. *Methods Mol Biol, 573*, 175-188. doi:10.1007/978-1-60761-247-
6_10.

1176
1177   **Competing interests**
1178   The authors declare no competing financial interests.
1179
Figure Legends
<u>**Figure 1**</u>**: Parasite diversity and mouse virulence within the Type X clade**
A)   53 isolates genotyped using 5 unlinked markers identified 9 distinct *T. gondii*
genotypes infecting southern sea otters. Type I, II, and III alleles are represented by red,
green, and blue, respectively. Sea otters were infected with *T. gondii* strains that possessed
either a non-archetypal allele (purple or orange) or a Type II allele (green) at the loci
examined. The number of sea otter infections (#SO) and percentage of total infections (%SO)

is shown for each genotype group. Each group was comprised of the following isolates: **A**
(3097, 3142, 3168, 3265, 3483, 3488, 3520, 3637, 3659, 3744, 3786, 3821, 3865, 3947, 3950,
4003, 4045, 4071, 4151), **B** (3458, 3523, 3728, 3897), **C** (3026, 3045, 3077, 3160, 4167), **D**
(3133, 3178, 3183, 3451, 4166), **E** (3819), **F** (3387, 3429, 3503), **G** (3675), **H** (3671), **II**
(2987, 2994, 3005, 3009, 3087, 3131, 3208, 3396, 3521, 3576, 3587, 3636, 3739, 4181).
B)   Total Sea Otter Strandings: number of sea otter strandings between 1998-2004 are
charted according to their 5 marker type designation of Type II (green) or sub-types of Type
X (range of purples). Type II infections not only clustered geographically (not shown) but
also temporally while Type X infections are steady over time. Additionally, the Type X, sub-
type A infections which were predominantly avirulent in sea otters show increased sea otter
infections over time.
C)   CD1 mice were infected with 50 tachyzoites intraperitoneally with one of 18 isolates
representing all singletons (**E, G, H**) and at least two isolates from each of the other 6 distinct
genotypes (**II, A, B, C, D, F**) recovered from sea otters. Mouse seroconversion and survival
was monitored for 30 days. At least two independent infection experiments were performed
using 5 mice each for every strain tested. Results shown are for those mice that
seroconverted or died acutely during infection. Strains are grouped and colored based on
their genotype and virulence in mice. **Red**, mouse virulent, **Blue**, intermediate mouse
virulent (some mice survived acute infection); **Green**, mouse avirulent (all mice survived
acute infection). Number of mice infected by each genotype and their relative virulence is
indicated.
D)   eBURST analysis to identify linkage disequilibrium across 17 nuclear-encoded linked
and unlinked markers identified 4 clonal complexes and 3 unique Type X genotypes among
21 isolates selected from the 9 distinct clades in Figure 1A. Isolate colors represent the
lineage Type from A: red (Type I), green (Type II), blue (Type III), and purple (Type X; all
subgroups (A-H)). Dot size is proportional to the number of isolates with that genotype;
larger dots are multi-isolate genotypes. Clonal complexes (CC) are indicated by lines
connecting isolates and are highlighted with ovals corresponding to their allelic identity (X-
purple, II-green)
1216
Figure 2: **Phylogenetic analyses support the designation of Type X as a recombinant**
clade of related strains
Maximum likelihood trees of sequenced markers are depicted. Allele designations were
dependent on a bootstrap support greater than 60%. Isolates are colored based on their
genotype from Figure 1A (I: red, II: green, III: blue, X: purple). A) Mitochondrial marker, CO1.
B) Nuclear genome marker BSR4 on Chromosome IV. At BSR4, all Type X strains possessed
a single allele, referred to as γ that was readily distinguished from Type I, II and III alleles C)
Comparison of two unlinked nuclear-encoded markers GRA7 and BAG1. At both markers,
Type X strains possessed one of two distinct alleles, referred to as γ or δ. Isolates possessing
δ alleles at both markers were highlighted in purple, those that possessed Type II alleles at
both markers were highlighted in green. Isolates that were recombinant and possessed a γ
allele at GRA7 but a δ allele at BAG1 were highlighted in pink, and conversely, a δ allele at
GRA7 and a γ allele at BAG1 were highlighted in orange. D) Comparison of linked genomic
markers GRA7 and SAG4. Genetic recombination is highlighted within this chromosome.
Isolates that are linked and have Type II lineage alleles at both markers are highlighted in

green. Isolates that show recombination between the γ/δ lineage alleles and the Type II
lineage alleles are highlighted in pink and orange depending on recombination
directionality.
1235

Figure 3: Genome-wide SNP typing of Type X displays haploblock recombination
across the genome
A) CGH-array hybridization of Type X strains against Type I, II and III lineage-specific probes
identified 8 distinct Type II and non-Type II hybridization patterns across the 8 Type X
strains examined. SNPs are represented by a dot in one of three rows indicating where the
isolate has hybridized to the microarray with a hybridization characteristic of lineage Type
I (red), II (green), or III (blue). Grey indicates no hybridization at this location. Chromosomes
are represented as alternating grey and white bars.
B) Linkage disequilibrium and network reticulation demonstrates the existence of recombination
across the genomes of Type X. A NeighborNet tree based on whole genome sequence identified
568,592 SNP variant positions across the 19 Type X strains reference mapped to ME49. Type I
(red), II (green), III (blue), and X (purple) strains were annotated in the box beside the group based
on their previous MLST designation. Two clusters of Type X strains were identified. Strains within
the inset were colored based on their murine virulence: virulent (red), intermediate virulent (blue),
and avirulent (green).
1251

Figure 4: SNP density plots identify distinct and different recombination crossover
points between the Type II and γ/δ lineage among the Type X strains examined
WGS reads for all strains were mapped to the ME49 reference genome. Each row represents
the SNPs of one strain mapped in a sliding 10 kb window across the genome, separated into
chromosomes columns. Vertical bars across the row represent the number of SNPs, the
height of the bar (from y-axis 0-200 bp) represents the amount of divergence from the ME49
genome. Type X strains were grouped based on their MLST designation.
1259

Figure 5: Whole genome sequencing establishes Type X as a recombinant clade of
related strains
A) PopNet analysis for all strains sequenced at WGS resolution. The genome of each strain is
represented by a circle of concatenated chromosomes starting at the top with Chromosome
Ia and rotating clockwise to Chromosome XII. All strains were clustered into four distinct
color groups based on their degree of shared ancestry (Type II-green, Type X-pink, Type I-
cyan, Type III-blue) and each isolate has an inner circle that is arrayed in color-group
haploblocks based on shared ancestry. Line thickness between strains indicates their
interrelatedness, boldest lines indicate 90-100% of the genome is in linkage disequilibrium,
as shown for strains 4167, 3819, 3045 and 3160.
B) Genetic model showing cross-over recombination points for isolate 3142. Type II ancestry
is shown in green whereas the mosaic of γ/δ ancestry is shown in black.
C) 21 WGS isolates are shown with their group designation for the 5 loci MLST (Figure 1A),
eBURST clonal complex identity (Figure 1D), 17 loci MLST (Table 1), and murine virulence
phenotype (Figure 1C).
1275

1276 **Figure 6: Population level QTL analysis identifies multiple chromosomal locations**
1277 **associated with acute murine virulence**

A) WGS data from Type X strains was down-selected to one SNP every 5 kb and analyzed using standard QTL methods with 1000 bootstrap support. LOD scores are shown across the chromosomes for these strains based on genetic association with acute murine virulence as shown in Table 1.

B) Significantly associated peaks from the QTL were identified based on clusters of SNP peaks. The LOD scores for the tallest portion of the associated genomic region are listed.

C) Distance-based Neighbor-joining ROP33 tree. Alignment of *ROP33* DNA sequences were used to construct a phylogenetic tree representing the majority of ROP33 alleles encoded within the species *Toxoplasma gondii.* Mouse virulence for each isolate is depicted by a color-code, per Figure 1C. **Red**, mouse virulent, **Blue**, intermediate mouse virulent (some CD-1 mice survived acute infection); **Green**, mouse avirulent (all CD-1 mice survived acute infection). Inoculum size was 50 tachyzoites injected intraperitoneally in CD-1 outbred mice.

D) Type I RH parasites deficient in ROP18 and ROP33 are less virulent than Type I RH parasites deficient in ROP18 alone, indicating that ROP33 is a virulence locus in *Toxoplasma.* CD-1 mice were infected intraperitoneally with 500 tachyzoites of the RH*Δrop18* or RH*Δrop18Δrop33* DKO, only the results for seropositive mice are shown. Data are combined from two independent experiments, each with five mice per group.

E) Bioluminescent detection of parasite burden *in vivo* at 9 days post-infection of CD-1 mice with 50 tachyzoites injected intraperitoneally. A representative image with photon output in photons/second/cm2/surface radiance (*sr*) is shown.

Supplemental Figure 1: **Stranding location, necropsy date, sex, *Toxoplasma* titer (IFAT), and co-infection status of every California Sea otter that yielded a *Toxoplasma* isolate for genetic characterization**

 A) Geographical map for all *Toxoplasma*-infected sea otter stranding locations along the California coastline. Each Sea otter stranding location is represented by a black dot.

B) Isolates of *Toxoplasma gondii* used in this paper. Isolates in yellow were genotyped previously using PCR-RFLP analyses at 4 loci (*B1, SAG1, SAG2, SAG3*) and isolates 3131, 3133, 3160, 3265 were DNA sequenced at the *GRA6 locus* [25]. ATOS numbers and stranding locations identify the location of each stranded Sea otter. Isolates were grouped together based on their 5 locus DNA sequence genotype and then by their ATOS number. "Tg" indicates *Toxoplasma gondii,* "Sn" indicates co-infection with *Sarcocystis neurona.*

Supplemental Table 1: **Expanded sequencing marker genotyping of Type X isolates reveals chromosomal segregation and recombination within chromosomes**

21 isolates characterized at 17 nuclear markers (ROP1 was excluded as it is a microsatellite marker, and prone to elevated mutation rates) clade into 12 distinct Type X genotypes (X1-X12). Isolate numbers in red were sequenced at whole-genome (WGS) resolution. Types I, II, III, γ and δ lineage alleles, as determined by phylogenetic comparisons, are colored red, green, blue, purple, and orange, respectively. Shades of colors represent genetic drift below the 60% bootstrap delineation from the canonical allele. White represents uninformative or incomplete DNA sequencing results.

Supplemental Table 2: **Potential virulence candidate genes derived from the QTL analysis**

1323 Chromosome peaks with LOD scores over 3 as shown in Figure 3B were interrogated via
1324 ToxoDB to select genes in these regions which have predicted signal sequences (Signal) and
1325 transmembrane (TM) domains in *Toxoplasma.* Genes are listed with gene ID, genomic
1326 location, and predicted protein function.

A

	Ia	*Ib*	*II*	*III*	*IV*	*V*	*VI*	*VIIa*	*VIIb*	*VIII*	*IX*	*X*	*XI*	*XII*
I (GT1)														
II (ME49)														
III (CTG)														
3142														
3265														
3026														
3045														
3178														
3503														
3429														
3671														

B

3675
ME49 2987
3131
[II]

RAY 4166 3671

WTD1

0.01

[III]

3387
3458
3168
ARI 3819
3133 3045 3429
3142 3160
4167 3503

VEG

[X]

[I]

RH-88

0.001

3387

3458

3168

ARI

3133

3142

3160
3045
4167   3819

3503
3429

A

Type X Strains Mapped to ME49, 10 kb window

A

B

γ/δ X II

Ia
Ib
II
III
IV
V
VI
VIIa
VIIb
VIII
IX
X
XI
XII

C

Isolate	5 loci	CC	17 loci	Mouse VIR
2987	II	CC1	II	-
3131	II	CC1	II	-
3142	A	CC2	X1	+
3265	A	CC2	X1	+
3947	A	CC2	X1	n.d.
3168	A	CC3	X2	+
3458	B	CC3	X3	+/-
3523	B	CC3	X3	-
3026	C	CC4	X4	-
3045	C	CC4	X4	-
4167	C	CC4	X4	+/-
3160	C	CC4	X5	n.d.
3133	D	CC3	X6	+
3178	D	CC3	X6	n.d.
4166	D	-	X7	+
3819	E	-	X8	-
3503	F	CC2	X9	+/-
3429	F	CC2	X9	+/-
3387	F	CC3	X10	+/-
3675	G	CC1	X11	-
3671	H	-	X12	+

Chrm	Start (bp)	End (bp)	LOD
V	1430036	2165042	4.091
VIIa	3815011	4085005	4.400
VIIb	3035265	3160030	3.680
VIII	6745008	6905157	4.194
X	7314	290066	4.190

A — Percent.Death LOD Scores vs Genetic Markers. α Threshold: 0.100; α Threshold: 0.050

B

C

D

E

RH   Rop33-

3.02E+09   4.06E+09

A CALIFORNIA

B

Isolate	ATOS #	5 loci	17 loci	Stranding Location	Necropsy Date	Sex	IFAT	Infection
2994	249	II		Santa Cruz	05/15/98	M	no serum	Tg
3739	264	II		Santa Cruz 26th Ave Beach	06/12/02	M	2560	Tg
3576	274	II		New Brighton State Beach	09/05/01	M	20480	Tg
3587	281	II		Rio Del Mar Beach	09/22/01	M	1280	Tg
2987	291	II	II	Mannressa State Beach	05/11/98	F	1280	Tg
4181	303	II		Sunset State Beach	04/14/04	F	40960	Sn/Tg
3009	321	II		Elkhorn Slough	06/04/98	M	640	Tg
3521	325	II		Salinas River State Beach	05/19/01	M	2560	Tg
3208	357	II		Monterey Bay	07/12/99	M	5120	Tg
3636	376	II		Del Monte Beach	12/27/01	M	80	Tg
3087	377	II		Monterey Harbor	10/28/98	F	640	Tg
3131	379	II	II	Monterey Bay	02/05/99	M	320	Tg
3005	928	II		Pismo Beach	06/01/98	M	160	Tg
3396	928	II		Oceano Dunes	11/11/00	M	20480	Tg
3097	84	A		Dunes Beach, San Mateo	11/16/98	M	5120	Tg
3142	180	A	X1	Ano Nuevo Island	02/24/99	F	320	Tg
3786	180	A		Ano Nuevo Island	09/20/02	M	2560	Sn
4045	260	A		Seabright Beach	10/21/03	M	10240	Tg
3950	263	A		Santa Cruz, 21st Street Beach	07/16/03	M	10240	Sn/Tg
3520	306	A		Sunset Beach	05/16/01	M	320	Tg
3659	374	A		Del Monte Beach	02/28/02	F	40	Tg
3265	426	A	X1	Carmel Beach	09/28/99	F	10240	Tg
3168	450	A	X2	Monterey Bay	04/09/99	F	2560	Tg
3865	480	A		Carmel River State Beach	04/03/03	F	10240	Tg
3947	523	A	X1	Pfieffer Burns Beach	06/26/03	F	20480	Tg
4003	750	A		Cambria	09/17/03	F	10240	Tg
4071	807	A		Cayucos, North of E Steet	12/03/03	F	2560	Tg
3483	820	A		Morro Strand	03/30/01	F	320	Sn/Tg
4151	825	A		Morro Bay	04/04/04	F	>40960	Sn/Tg
3488	919	A		Pismo Beach, ~400m North of Grand Ave	04/04/01	F	640	Tg
3821	922	A		Pismo Beach	01/06/03	M	2560	Tg
3744	924	A		Pismo Beach	07/02/02	F	5120	Tg
3637	926	A		Oceana Dunes	02/14/03	F	no serum	Tg
3523	141	B	X3	Pescadero Point	05/31/01	F	5120	Sn/Tg
3458	422	B	X3	Stillwater Cove	01/16/01	F	<80	Tg
3897	435	B		Pt Lobos State Reserve	04/22/03	F	10240	Tg
3728	1,445	B		Ventura County (Pt Mugu St Pk)	05/29/02	M	320	Tg
3077	145	C		Bean Hollow, Half Moon Bay	09/27/98	M	1280	Tg
3045	808	C	X4	Cayucas	07/09/98	F	no serum	Tg
3026	840	C	X4	Morro Bay Sandspit	06/24/98	M	320	Tg
4167	852	C	X4	Morro Creek	04/09/04	M	10240	Sn/Tg
3160	933	C	X5	Pismo Beach	03/29/99	M	320	Tg
3178	321	D	X6	Moss Landing Harbor	03/26/02	F	320	Tg
3183	816	D		Morro Strand	05/04/99	F	640	Tg
3133	818	D	X6	Morro Strand Campground	02/04/99	M	20480	Tg
3451	836	D		Morro Bay, Sandspit between port 3 & 4	11/29/00	M	<80	Tg
4166	827	D	X7	Morro Bay	04/09/04	M	40960	Sn/Tg
3819	732	E	X8	San Simeon Bay	12/28/02	F	2560	Tg
3429	827	F	X9	Morro Bay	10/30/00	F	2560	Tg
3387	922	F	X10	Pismo Beach	07/19/00	F	1280	Tg
3503	1,319	F	X9	Rincon Pt., Ventura County	08/09/01	M	2560	Tg
3675	435	G	X11	Point Lobos	03/20/02	F	1280	Tg
3671	321	H	X12	Elkhorn Slough	03/13/02	M	640	Sn/Tg
		n=53	n=21					

Marker	5 Loci	17 Loci	C228	Chr_II_10	C292	BSR4	L358	PK1	SAG4	ROP18	GRA7	BAG1	SRS2	SAG1	SAG2	BTB	GRA6	ROP1	UPRT	SAG3	APICO	COX1
Chromosome			Ib	II	III	IV	V	VI	VIIa	VIIa	VIIa	VIIb	VIII	VIII	VIII	IX	X	XI	XI	XII	API	MITO
Length			421	639	397	1138	354	1011	174	1617	460	1917	983	700	1243	331	758	960	1214	253	442	418
Type I		I																				
Type II		II																				
Type III		III																				
2987	II	II																				
3131	II	II																				
3142	A	X1						44			44	44					44					
3265	A	X1						44			44	44					44					
3947	A	X1						44			44	44					44	44				
3168	A	X2															44	44				
3458	B	X3										44										
3523	B	X3										44										
4167	C	X4						44		44	44											
3026	C	X4						44		44	44											
3045	C	X4						44		44	44											
3160	C	X5						44		44												
3133	D	X6										44					44					
3178	D	X6															44					
4166	D	X7					44	44									44					
3819	E	X8						44			44											
3503	F	X9						44		44	44											
3429	F	X9						44		44	44											
3387	F	X10									44	44						44				
3675	G	X11																				
3671	H	X12									44											

Type II		green
α		red
β		blue
γ	32	purple
δ	44	orange

Chr	Start (bp)	End (bp)	LOD	TM	Signal	Gene ID	Genomic Location(s)	Product Description
5	220001	350021	3.25	Y	Y	TGME49_220240	TGME49_chrV:235,019..240,463(-)	hypothetical protein
				Y	Y	TGME49_220300	TGME49_chrV:275,453..283,005(-)	ribosomal protein L15 protein
				Y	Y	TGME49_220330	TGME49_chrV:295,153..299,369(-)	hypothetical protein
				Y	Y	TGME49_220350	TGME49_chrV:310,715..318,211(+)	tRNA ligases class II (D, K and N) domain-containing protein
5	1430036	2165042	4.09	Y	Y	TGME49_285870	TGME49_chrV:2,158,544..2,162,541(-)	SAG-related sequence SRS20A
				Y	Y	TGME49_285940	TGME49_chrV:2,123,872..2,135,543(-)	hypothetical protein
				Y	Y	TGME49_286180	TGME49_chrV:1,933,873..1,942,288(-)	tRNA ligases class I (M) protein
				Y	Y	TGME49_286450	TGME49_chrV:1,824,587..1,826,598(+)	dense granule protein GRA5
				Y	Y	TGME49_286530	TGME49_chrV:1,790,919..1,793,716(+)	hypothetical protein
				Y	Y	TGME49_286620	TGME49_chrV:1,754,511..1,762,820(-)	S1 RNA binding domain-containing protein
				Y	Y	TGME49_286630	TGME49_chrV:1,750,888..1,752,970(+)	redoxin domain-containing protein
				Y	Y	TGME49_286770	TGME49_chrV:1,682,413..1,683,569(+)	hypothetical protein
				Y	Y	TGME49_213920	TGME49_chrV:1,428,849..1,440,483(-)	hypothetical protein
7a	3815011	4085005	4.40	Y	Y	TGME49_201130	TGME49_chrVIIa:4,019,644..4,026,659(+)	rhoptry kinase family protein ROP33
				Y	Y	TGME49_201180	TGME49_chrVIIa:3,982,746..3,988,716(-)	hypothetical protein
				Y	Y	TGME49_201390	TGME49_chrVIIa:3,908,769..3,911,287(-)	hypothetical protein
				Y	Y	TGME49_201750	TGME49_chrVIIa: 3,822,835 – 3,823,206(+)	Tctex-1 family protein
				Y	Y	TGME49_201760	TGME49_chrVIIa: 3,816,484 – 3,820,369(+)	hypothetical protein
7b	3035265	3160030	3.68	Y	Y	TGME49_258828	TGME49_chrVIIb:3,159,970..3,160,907(+)	hypothetical protein
				Y	Y	TGME49_258840	TGME49_chrVIIb:3,137,682..3,141,698(+)	hypothetical protein
				Y	Y	TGME49_258860	TGME49_chrVIIb:3,122,191..3,126,503(+)	hypothetical protein
				Y	Y	TGME49_258890	TGME49_chrVIIb:3,103,220..3,104,003(-)	hypothetical protein
				Y	Y	TGME49_258910	TGME49_chrVIIb:3,092,653..3,098,081(+)	hypothetical protein
				Y	Y	TGME49_258950	TGME49_chrVIIb:3,079,180..3,083,732(+)	lectin family protein
				Y	Y	TGME49_258990	TGME49_chrVIIb:3,038,025..3,049,513(+)	Bromodomain-containing protein
				Y	Y	TGME49_259000	TGME49_chrVIIb:3,035,953..3,049,513(+)	hypothetical protein
8	5435051	6125010	3.8	Y	Y	TGME49_269150	TGME49_chrVIII:6,039,099..6,042,614(-)	DHHC zinc finger domain-containing protein
				Y	Y	TGME49_269190	TGME49_chrVIII:6,002,854..6,007,557(-)	glyceraldehyde-3-phosphate dehydrogenase GAPDH2
				Y	Y	TGME49_269400	TGME49_chrVIII:5,836,645..5,840,235(+)	oxidoreductase, short chain dehydrogenase/reductase family protein
				Y	Y	TGME49_269750	TGME49_chrVIII:5,625,999..5,628,567(+)	CrcB family protein
				Y	Y	TGME49_269885	TGME49_chrVIII:5,564,529..5,572,853(+)	rhoptry metalloprotease toxolysin TLN1
				Y	Y	TGME49_269920	TGME49_chrVIII:5,535,689..5,545,827(-)	phosphatidylserine decarboxylase
				Y	Y	TGME49_269980	TGME49_chrVIII:5,483,536..5,489,390(-)	preprotein translocase Sec61, putative
				Y	Y	TGME49_269690	TGME49_chrVIII:5,680,572..5,684,829(+)	hypothetical protein
				Y	Y	TGME49_270030	TGME49_chrVIII:5,448,751..5,449,486(+)	hypothetical protein
8	6745008	6905157	4.19	Y	Y	TGME49_200230	TGME49_chrVIII:6,753,611..6,755,422(-)	microneme protein MIC17C
				Y	Y	TGME49_200440	TGME49_chrVIII:6,883,172..6,887,049(+)	hypothetical protein
10	280002	290066	4.19	N/A	N/A	N/A	N/A	N/A