Title: *Scn1a*-GFP transgenic mouse revealed Nav1.1 expression in neocortical pyramidal

tract projection neurons

Authors: Tetsushi Yamagata[1, 2, ¶], Ikuo Ogiwara[2, 3, ¶], Tetsuya Tatsukawa[2, ¶], Toshimitsu

Suzuki[1, 2], Yuka Otsuka[1], Nao Imaeda[1], Emi Mazaki[2, †], Ikuyo Inoue[2, ‡], Natsuko Tokonami[2, §],

Yurina Hibi[1], Shigeyoshi Itohara[4], and Kazuhiro Yamakawa[1, 2, *]

Affiliations:

[1]Department of Neurodevelopmental Disorder Genetics, Institute of Brain Science, Nagoya

City University Graduate School of Medical Sciences, Nagoya, Aichi 467-8601, Japan.

[2]Laboratory for Neurogenetics, RIKEN Center for Brain Science, Wako, Saitama 351-0198,

Japan.

[3]Department of Physiology, Nippon Medical School, Tokyo 113-8602, Japan.

[4]Laboratory for Behavioral Genetics, RIKEN Center for Brain Science, Wako, Saitama

351-0198, Japan.

[¶]These authors contributed equally to this work.

[†]Present address; International Research Center for Neurointelligence (IRCN), The University

of Tokyo, Institutes for Advanced Study, Tokyo 113-0033, Japan.

[‡]Present address; Medical-risk Avoidance based on iPS Cells Team, RIKEN Center for

Advanced Intelligence Project, Kyoto 606-8507, Japan.

[§]Present address; Institute of Physiology, University of Zürich, CH-8057 Zürich, Switzerland.

[*]Correspondence and requests for materials should be addressed to K.Y. (E-mail:

yamakawa@med.nagoya-cu.ac.jp)

Abstract

Expressions of voltage-gated sodium channels Nav1.1 and Nav1.2, encoded by *SCN1A* and

SCN2A genes respectively, have been reported to be mutually exclusive in most brain regions.

In juvenile and adult neocortex, Nav1.1 is predominantly expressed in inhibitory neurons

while Nav1.2 is in excitatory neurons. Although a distinct subpopulation of layer V (L5)

neocortical excitatory neurons were also reported to express Nav1.1, their nature has been

uncharacterized. In hippocampus, Nav1.1 has been proposed to be expressed only in

inhibitory neurons. By using newly-generated transgenic mouse lines expressing *Scn1a*

promoter-driven green fluorescent protein (GFP), here we confirm the mutually-exclusive

expressions of Nav1.1 and Nav1.2 and the absence of Nav1.1 in hippocampal excitatory

neurons. We also show that Nav1.1 is expressed in inhibitory and a subpopulation of

excitatory neurons not only in L5 but all layers of neocortex. By using neocortical excitatory

projection neuron markers including FEZF2 for L5 pyramidal tract (PT) and TBR1 for layer

VI (L6) cortico-thalamic (CT) projection neurons, we further show that most L5 PT neurons

and a minor subpopulation of layer II/III (L2/3) cortico-cortical (CC) neurons express Nav1.1

while the majority of L6 CT, L5/6 cortico-striatal (CS) and L2/3 CC neurons express Nav1.2.

These observations now contribute to the elucidation of pathological neural circuits for

diseases such as epilepsies and neurodevelopmental disorders caused by *SCN1A* and *SCN2A*

mutations.

Introduction

Voltage-gated sodium channels (VGSCs) play crucial roles in the generation and propagation of action potentials, contributing to excitability and information processing (Catterall, 2012). They consist of one main pore-forming alpha- and one or two subsidiary beta-subunits that regulate kinetics or subcellular trafficking of the alpha subunits. Human has nine alpha (Nav1.1~Nav1.9) and four beta (beta-1~beta-4) subunits. Among alphas, Nav1.1, Nav1.2, Nav1.3 and Nav1.6, encoded by *SCN1A*, *SCN2A*, *SCN3A* and *SCN8A*, respectively, are expressed in central nervous system. *SCN3A* is mainly expressed embryonically (Brysch et al., 1991), and *SCN1A*, *SCN2A* and *SCN8A* are major alphas after birth. Although these three genes show mutations in a wide spectrum of neurological diseases such as epilepsy, autism spectrum disorder (ASD) and intellectual disability, two of those, *SCN1A* and *SCN2A,* are major ones (reviewed in Yamakawa et al., 2016; Meisler et al., 2021). To understand the circuit basis of these diseases, it's indispensable to know the detailed distributions of these molecules in the brain.

We previously reported that expressions of Nav1.1 and Nav1.2 seem to be mutually-exclusive in many brain regions (Yamagata et al., 2017). In adult neocortex and hippocampus, Nav1.1 is dominantly expressed in medial ganglionic eminence-derived parvalbumin-positive (PV-IN) and somatostatin-positive (SST-IN) inhibitory neurons (Ogiwara et al., 2007; Lorincz and Nusser, 2008; Ogiwara et al., 2013; Li et al., 2014; Tai et al., 2014; Tian et al., 2014; Yamagata et al, 2017). In the neocortex, some amount of Nav1.1 is also expressed in a distinct subset of layer V (L5) excitatory neurons (Ogiwara et al., 2013), but their natures were unknown. In the hippocampus, Nav1.1 seems to be expressed in inhibitory but not in excitatory neurons (Ogiwara et al., 2007; Ogiwara et al., 2013). In contrast, a major amount of Nav1.2 (~95%) is expressed in excitatory neurons including the most of neocortical and all of hippocampal ones, and a minor amount is expressed in caudal ganglionic eminence-derived inhibitory neurons such as vasoactive intestinal polypeptide (VIP)-positive ones (Lorincz and Nusser 2010; Yamagata et al., 2017; Ogiwara et al., 2018). However, a recent study reported that a subpopulation (more than half) of VIP-positive inhibitory neurons is Nav1.1-positive (Goff and Goldberg, 2019).

VGSCs are mainly localized at axons and therefore it is not always easy to identify their origins, soma. To overcome this, here in this study we generated bacterial artificial chromosome (BAC) transgenic mouse lines that express GFP under the control of *Scn1a* promoters, and we carefully investigated the GFP/Nav1.1 distribution in mouse brain. Our analysis confirmed that expressions of Nav1.1 and Nav1.2 are mutually-exclusive and that in

neocortex Nav1.1 is expressed in both inhibitory and excitatory neurons while in

hippocampus only in inhibitory but totally absent in excitatory neurons. Furthermore by using

a transcription factor FEZF2 (FEZ family zinc finger protein 2 transcriptional factor), also

referred to as Fezl, Fez1, Zfp312 and Fez, as a marker for L5 PT neurons (Inoue et al., 2004;

Chen et al., 2005; Chen et al., 2008; Molyneaux et al., 2005; Lodato et al., 2014 ; Matho et al.,

2021), and a transcription factor TBR1 which suppresses FEZF2 expression and therefore

does not overlap with FEZF2 (Han et al., 2011; McKenna et al., 2011, Matho et al., 2021), we

found that most of L5 FEZF2-positive neurons are GFP-positive while L5/6 TBR1-positive

neurons are largely GFP-negative and Nav1.2-positive. These results proposed that Nav1.1 is

expressed in L5 PT while Nav1.2 in L5/6 non-PT neurons such as L5/6 CS and L6 CT

projection neurons. A majority of L2/3 excitatory neurons express Nav1.2 but a minor

subpopulation are GFP-positive, suggesting that most of CC projection neurons express

Nav1.2 but the distinct minor population express Nav1.1. These results refine the expression

loci of Nav1.1 and Nav1.2 in the brain and should contribute to the understanding of circuit

mechanisms for diseases caused by *SCN1A* and *SCN2A* mutations.

Results

Generation and verification of *Scn1a*-GFP transgenic mouse lines

Scn1a-GFP founder mice were generated from C57BL/6J zygotes microinjected with a

modified *Scn1a*-GFP BAC construct harboring all, upstream and downstream,

Scn1a-promoters (Nakayama et al., 2010) (Figure 1A) (see Materials and Methods for details).

Western blot analysis (Figure 1B) and immunohistochemistry (Supplementary figure S1)

showed robust GFP expression and mostly normal expression levels of Nav1.1 in *Scn1a*-GFP

mouse lines #184 and #233. Both lines showed a similar distribution of chromogenic GFP

immunosignals across the entire brain (Figures 2A-H), and a similar distribution was also

obtained in fluorescence detection of GFP (Figures 2I-L and Supplementary figure S2). In

neocortex (Figures 2B, F, J and Supplementary figures S2B, F), GFP-positive cells were

distributed throughout all cortical layers. In hippocampus (Figures 2C, G, K and

Supplementary figures S2C, G), cells with intense GFP signals, which are assumed to be

PV-IN and SST-IN (Ogiwara et al, 2007; Tai et al., 2014) (see also Figure 8), were scattered

in stratum oriens, pyramidale, radiatum, lucidum and lacunosum-moleculare of the CA (cornu

ammonis) fields, hilus and molecular layer of dentate gyrus. Of note, somata of dentate

granule cells were apparently GFP-negative. CA1~3 pyramidal cells were twined around with

fibrous GFP immunosignals. We previously reported that the fibrous Nav1.1-signals clinging

to somata of hippocampal CA1~3 pyramidal cells were disappeared by conditional

elimination of Nav1.1 in PV-INs but not in excitatory neurons, and therefore concluded that

these Nav1.1-immunopositive fibers are axon terminals of PV-INs (Ogiwara et al., 2013). As

such, GFP signals are fibrous but do not form cell shapes in the CA pyramidal cell layer

(Figures 2C, G, K and Supplementary figures S2C, G), and therefore these CA pyramidal

cells themselves are assumed to be GFP-negative. These observations further confirmed our

previous proposal that hippocampal excitatory neurons are negative for Nav1.1 (Ogiwara et

al., 2007; Ogiwara et al., 2013). In cerebellum (Figures 2D, H, L and Supplementary figures

S2D, H), GFP signals appeared in Purkinje, basket, and deep cerebellar nuclei cells, again

consistent to the previous reports (Ogiwara et al., 2007; Ogiwara et al., 2013). In the

following analyses, we used the line #233 which shows stronger GFP signals than #184.

Quantification of Nav1.1 signals in western blot analyses of brain lysates from the

Scn1a-GFP mice and their wild-type littermates (N = 5 animals per each genotype) showed no

difference between genotypes, while that of GFP somehow deviated among individual

Scn1a-GFP mice (Supplementary figure S3). Fluorescence imaging of the *Scn1a*-GFP sagittal

brain sections at postnatal day 15 (P15), 4-week-old (4W) and 8W showed that GFP-signals

continue to be intense in caudal region such as thalamus, mid brain, and brainstem (Figure 3),

which is well consistent with our previous report of Nav1.1 protein and *Scn1a* mRNA

distributions in wild-type mouse brain (Ogiwara et al, 2007).

Nav1.1 is expressed in both excitatory and inhibitory neurons in neocortex but only in

inhibitory neurons in hippocampus

In the neocortex of *Scn1a*-GFP mouse, a large number of cells with GFP-positive somata

(GFP-positive cells) were broadly distributed across all cortical layers (Figure 3 and

Supplementary figure S4). Intensities of GFP signals in primary somatosensory cortex (S1) at

L2/3 are much higher than other areas such as primary motor cortex (M1) (Supplementary

figure S4), however the cell population (density) of GFP-positive cells did not differ in these

areas indicating that GFP-signals for GFP-positive cells are stronger in S1 at L2/3. Although

GFP signals are strong in PV-INs (see Figure 8), cell density of PV-INs is not specifically

high at S1 area and therefore most cells with strong GFP signals in S1 at L2/3 may not be

PV-INs but excitatory neurons.

In order to know the ratio of GFP-positive cells among all neurons, we further performed

immunohistochemical staining using NeuN-antibody on *Scn1a*-GFP mouse at P15 and cells

were counted at M1 and S1 (Supplementary figure S5 and Supplementary table S1). The

NeuN staining showed that GFP-positive cells occupy 30% (L2/3), 32% (L5) and 22% (L6)

of NeuN- or GFP-positive cells at P15 (Supplementary figure S5B and Supplementary table

S1). However, we noticed that sparsely-distributed cells with intense GFP-signals, which are

assumed to be PV-INs (see Figure 8), were often NeuN-negative (Supplementary figure S5A -

arrowheads), reminiscent of a previous report that NeuN expression is absent in cerebellar

inhibitory neurons such as Golgi, basket and satellite cells in cerebellum (Weyer et al., 2003).

Therefore, NeuN-positive cells do not represent all neurons in neocortex as well.

NeuN/GFP-double negative neurons could even exist and therefore above figure

(Supplementary figure S5B) may deviate from the real ratios of GFP-positive cells among all

neurons.

Next, we performed triple-immunostaining of Nav1.1, GFP and ankyrinG on brains of

Scn1a-GFP mouse at P15. In the neocortex (Figure 4), axon initial segments (AISs) of cells

with Nav1.1-positive somata were always Nav1.1-positive but somata of cells with

Nav1.1-positive AISs were occasionally Nav1.1-negative (Figures 4A-C). Cell counting

revealed that 17% (L2/3), 21% (L5) and 8% (L6) of neurons (cells with ankyrinG-positive

AISs) were GFP-positive (Figure 4D-left panel and Supplementary table S2). Of note, all

cells with Nav1.1-positive AISs or somata were GFP-positive, but AISs or somata for only

half of GFP-positive cells were Nav1.1-positive (Figure 4D and Supplementary tables S3, S4),

possibly due to undetectably low levels of Nav1.1 immunosignals in a subpopulation of

GFP-positive cells. The above ratios of GFP-positive cells among neurons (cells with

ankyrinG-positive AISs) obtained in the triple-immunostaining of Nav1.1, GFP and ankyrinG

are rather discordant to those obtained in the later experiment of triple-immunostaining of

Nav1.2, GFP and ankyrinG, 23% (L2/3), 30% (L5) and 21% (L6) (see Figure 11). Therefore,

we additionally performed double-immunostaining of GFP and ankyrinG on brains of

Scn1a-GFP mouse at P15, and the ratios of of GFP-positive cells among neurons were 30%

(L2/3), 26% (L5) and 9% (L6) (Supplementary figure S6 and Supplementary table S5).

Averaged ratios of GFP-positive cells among neurons of Figures 4D, 11B and Supplementary

figure S6B are 23% (L2/3), 26% (L5) and 13% (L6) (Supplementary figure S7 and

Supplementary table S6), which are actually significantly lower than those obtained in the

NeuN-staining (Supplementary figure S5 and Supplementary table S1).

In contrast to the neocortex where only half of GFP-positive cells were Nav1.1-positive, in

the hippocampus all GFP-positive cells were Nav1.1-positive and all Nav1.1-positive cells

were GFP-positive (Figure 5). Actually, most of excitatory neurons such as CA1~3 pyramidal

cells and dentate granule cells were GFP-negative. As described above (Figures 2C, G),

fibrous GFP and Nav1.1 signals twining around CA1~3 pyramidal cells' somata which are

assumed to be axon terminals of PV-INs were again observed (Figures 5A, B, D). Cell

counting in the hippocampal CA1 region showed that 98% of cells with GFP-positive somata

were Nav1.1-positive at their AISs and 100% of cells with Nav1.1-positive AISs were

GFP-positive (Figure 5F and Supplementary tables S7, S8).

Double *in situ* hybridization of *Scn1a* and GFP mRNAs showed that these signals well

overlap in both neocortex and hippocampus of *Scn1a*-GFP mice (Figure 6), further supporting

that the GFP signals well represent endogenous *Scn1a*/Nav1.1 expression. Again, in

neocortex *Scn1a* and GFP mRNAs seem to be expressed in a number of neurons including

some of excitatory pyramidal cells, while in hippocampus they are absent in excitatory

neurons such as CA1~3 pyramidal cells and dentate granule cells. All of these distributions of

Scn1a and GFP mRNAs in *Scn1a*-GFP transgenic mouse brain are consistent to our previous

report of regional distributions of *Scn1a* mRNA in wild-type mouse (Ogiwara et al, 2007).

To investigate the ratio of inhibitory neurons in GFP-positive cells, we generated and

examined *Scn1a*-GFP and vesicular GABA transporter (*Vgat*)-Cre (Ogiwara et al., 2013)

double transgenic mice in which *Vgat*-Cre is expressed in all GABAergic inhibitory neurons

and visualized by floxed tdTomato transgene (Figure 7). In the neocortex at 4W, 23% (L2/3),

28% (L5) and 27% (L6) of GFP-positive cells were Tomato-positive inhibitory neurons and

73% (L2/3), 77% (L5) and 83% (L6) of Tomato-positive cells were GFP-positive (Figure 7C

and Supplementary tables S9, S10). These results suggest that a significant subpopulation of

neocortical excitatory neurons also express Nav1.1. Our previous observation that Nav1.1 is

expressed in callosal axons of neocortical excitatory neurons (Ogiwara et al., 2013) supports

that a subpopulation of L2/3 CC neurons express Nav1.1. Unlike in neocortex, in the

hippocampus most of GFP-positive cells were Tomato-positive, 98% (CA1) and 94% (DG),

and majorities of Tomato-positive GABAergic neurons are GFP-positive, 93% (CA1) and

77% (DG). These results further confirmed that in hippocampus Nav1.1 is expressed in

inhibitory neurons but not in excitatory neurons. Although somata of pyramidal cells in

CA2/3 region are weakly GFP-positive in this and some other experiments (Figure 7B,

Supplementary figure S2G), those were GFP-negative in other experiments (Figures 2K, 5A,

D) and therefore the Nav1.1 expression in CA2/3 pyramidal cells would be minimal if any.

We further performed immunohistochemical staining of PV and SST in neocortex and

hippocampus of *Scn1a*-GFP mice at 4W (Figure 8). PV and SST do not co-express in cells

and do not overlap. PV-INs and SST-INs were both GFP-positive, and especially GFP signals

in PV-INs were intense (Figure 8A). Cell counting revealed that 21% (L2/3), 37% (L5), 37%

(L6), 58% (CA1), 42% (CA2/3), and 41% (DG) of GFP-positive cells were PV or

SST-positive depending on regions in neocortex and hippocampus (Figure 8B and

Supplementary tables S11-S13). All PV-INs were GFP-positive (Figure 8B - middle), and

most of SST-INs were GFP-positive (Figure 8B - right). Comparison of these results with

those of *Vgat*-Cre mouse (Figure 7) suggests that GFP-positive GABAergic neurons in

neocortex are mostly PV- or SST-positive, while in hippocampus a half of those are

PV/SST-negative GABAergic neurons. Higher ratios of PV- or SST-positive cells (Figure 8B)

compared with those of *Vgat*-Cre-positive cells (Figure 7C) among GFP-positive cells would

be explained by that we counted PV-positive cells even if their PV-immunosignals are

moderate and a significant subpopulation of such cells are known to be excitatory neurons

(Jinno et al., 2004; Tanahira et al., 2009; Matho et al., 2021). Quantitative analysis of GFP

signal intensity and area size of cells revealed that GFP signal intensities in PV-positive cells

were significantly higher than those in PV-negative cells and GFP signal intensities in

SST-positive cells were lower than those in PV-positive cells but similar to PV/SST-double

negative cells (Figure 9 and Supplementary tables S14-S16). These results indicate that

Nav1.1 expression level in PV-INs is significantly higher than those in excitatory neurons and

PV-negative GABAergic neurons including SST-INs.

Nav1.1 and Nav1.2 expressions are mutually exclusive in mouse brain

We previously reported that expressions of Nav1.1 and Nav1.2 seem to be mutually-exclusive

in multiple brain regions including neocortex, hippocampal CA1, dentate gyrus, striatum,

globus pallidus, and cerebellum in wild-type mice (Yamagata et al., 2017). To further confirm

it, here we performed triple immunostaining for Nav1.1, Nav1.2 and ankyrinG, and counted

Nav1.1- or Nav1.2-immunopositive AISs in the neocortex of *Scn1a*-GFP mice (Figure 10).

The staining again showed that Nav1.1 and Nav1.2 expressions are mutually exclusive in

brain regions including neocortex (Figure 10A). The counting revealed that 5% (L2/3), 6%

(L5) and 3% (L6) of AISs at P15 were Nav1.1-positive and 78% (L2/3), 69% (L5) and 69%

(L6) of AISs at P15 were Nav1.2-positive in the neocortex (Figure 10B and Supplementary

table S17). Of note, less than 0.5% of AISs are Nav1.1/Nav1.2-double positive confirming

that Nav1.1 and Nav1.2 do not co-express. These results are consistent with our previous

study (Yamagata et al., 2017) and further support that expressions of Nav1.1 and Nav1.2 are

mutually exclusive in mouse neocortex at least at immunohistochemical level.

As mentioned, GFP signals in *Scn1a*-GFP mouse can represent even moderate or low

Nav1.1 expressions which cannot be detected by immunohistochemical staining, so some of

GFP-positive cells may still express Nav1.2. To investigate whether and if so how much of

GFP-positive cells have Nav1.2-positive AISs in *Scn1a*-GFP mouse neocortex, we performed

triple immunohistochemical staining for Nav1.2, GFP and ankyrinG (Figure 11 and

Supplementary tables S18-S20). The staining showed that AISs of GFP-positive cells are

largely negative for Nav1.2, and cells with Nav1.2-positive AISs are mostly GFP-negative

(Figure 11A). Cell counting revealed that 88% (L2/3), 90% (L5) and 95% (L6) of cells with

Nav1.2-positive AISs at P15 were GFP-negative (Figure 11B-middle panel and

Supplementary table S19), and 69% (L2/3), 83% (L5), and 86% (L6) of AISs of GFP-positive

cells at P15 were Nav1.2-negative (Figure 11B-right panel and Supplementary table S20).

These results indicate that the co-expression of GFP and Nav1.2 would be minimal if any.

Neocortical pyramidal tract and a subpopulation of cortico-cortical projection neurons

express Nav1.1

Neocortical excitatory neurons can be divided into functionally distinct subpopulations, a

majority of those are pyramidal cells which have axons of long-range projections such as L2/3

CC, L5 PT, L5/6 CS and L6 CT projection neurons (Shepherd, 2013). Although PT neurons

also project their axon collaterals to ipsilateral striatum, CS neurons project bilaterally to ipsi-

and contralateral striata. A transcription factor FEZF2 is expressed in L5 PT neurons, forms

their axonal projections and defines their targets (Inoue et al., 2004; Chen et al., 2005; Chen et

al., 2008; Lodato et al., 2014). Most of PT neurons are FEZF2-positive (Molyneaux et al.,

2005; Matho et al., 2021). We previously reported that a subpopulation of neocortical L5

pyramidal neurons is Nav1.1-positive (Ogiwara et al., 2013), but their natures were unclear.

To investigate those, here we performed immunohistochemical staining of FEZF2 and GFP

on *Scn1a*-GFP mouse brains (Figures 12 and 13, Supplementary tables S21-S23). In L5 where

a major population of FEZF2-positive cells locate (Figure 12A), a majority of FEZF2-positive

neurons were GFP-positive (83% and 96% of FEZF2-positive neurons were GFP-positive at

P15 and 4W, respectively) (Figure 12B - middle panels, Supplementary table S22). In L2/3,

FEZF2-positive cells were scarce (Figure 12A). In L6, a certain number of FEZF2-posiive

cells exist but overlaps of FEZF2 and GFP signals are much less compared to those in L5

(Figure 12 and Supplementary tables S21-S23). Quantitative analyses revealed that

FEZF2/GFP-double positive cells in L5 showed significantly lower GFP immunosignal

intensities and larger signal areas (soma sizes) compared to those of

FEZF2-negative/GFP-positive cells (Figure 13 and Supplementary table S24), indicating that

L5 PT neurons showed lower Nav1.1 expression compared to other neurons such as PV-INs

(see also Figure 9). Soma sizes of GFP-positive cells in L6 were overall smaller than those of

FEZF2/GFP-double positive cells in L5, and there was no statistically significant difference in

size between the FEZF2-positive/negative subpopulations. However, FEZF2/GFP-double

positive cells still showed lower intensity of GFP signals compared to

FEZF2-negative/GFP-positive cells (Figure 13 and Supplementary table S24).

We further performed triple immunostaining of FEZF2, Nav1.1 and ankyrinG on

Scn1a-GFP mice at P15, and found that 11% of FEZF2-positive cells have Nav1.1-positive

AIS in L5 of *Scn1a*-GFP mouse neocortex (Supplementary figure S8 and Supplementary

tables S25-S27). The low ratios of FEZF2/Nav1.1-double positive cells are most possibly due

to immunohistochemically-undetectable low levels of Nav1.1 expression in these excitatory

neurons.

We also performed triple immunostaining of FEZF2, Nav1.2 and ankyrinG on *Scn1a*-GFP

mice at P15 (Supplementary figure S9 and Supplementary tables S28-S31). The staining

showed that 20% of neurons (cells with ankyrinG-positive AISs) in L5 are FEZF2-positive

and a half of L5 FEZF2-positive cells have Nav1.2-positive AISs. Together with the

observation that most of FEZF2-positive cells are GFP-positive (Figure 12), these results

indicate that a subpopulation of FEZF2-positive PT neurons may express both Nav1.1 and

Nav1.2.

We additionally performed triple immunostaining of FEZF2, GFP, and Nav1.2 on

Scn1a-GFP mice at P15 (Supplementary figure S10 and Supplementary table S32), showing

that in L5 74% of FEZF2-positive cells are GFP-positive but a majority of their AISs are

Nav1.2-negative. The ratios of Nav1.2-positive cells among FEZF2-positive cells obtained in

the triple-immunostaining of FEZF2, GFP and Nav1.2 (Supplementary figure S10) are 27%

(L5) and 40% (L6). These results further support the above notion that a subpopulation of

FEZF2-positive PT neurons may express both Nav1.1 and Nav1.2. Although further studies

such retrograde tracking analyses are required to confirm and figure out the detailed circuits,

all these results propose that the majority of L5 PT neurons express Nav1.1.

The majority of cortico-thalamic, cortico-striatal and cortico-cortical projection neurons

express Nav1.2

TBR1 (T-box brain 1 transcription factor) is a negative regulator of FEZF2 and therefore not

expressed in the PT neurons, all of which are known to be FEZF2-positive (Chen et al., 2008;

Han et al., 2011; McKenna et al., 2011). TBR1 is predominantly expressed in L6 CT neurons

and subpopulations of L2/3 and L5 non-PT excitatory neurons instead (Han et al., 2011;

McKenna et al., 2011; Matho et al., 2021). To further elucidate the distributions of GFP

(Nav1.1)-expressing neurons in neocortex, we performed immunohistochemical staining of

TBR1 on *Scn1a*-GFP mice (Figure 14 and Supplementary tables S33-S35) and quantitated

GFP signal intensities and area sizes of cells (Figure 15 and Supplementary table S36).

TBR1-positive cells were predominant in neocortical L6, some in L5 and a few in L2/3. In L5,

contrary to the high ratios of GFP-positive cells among FEZF2-positive cells (83% at P15,

and 96% at 4W) (Figure 12B), the ratios of GFP-positive cells among TBR1-positive cells are

quite low (11% at P15, and 5% at 4W) (Figure 14B-middle panels and Supplementary table

S34). Soma sizes of TBR1/GFP-double positive cells were smaller than those of

TBR1-negative/GFP-positive cells (Figure 15-middle panel and Supplementary table S36). In

L6 where a major population of TBR1-positive neurons locate, the ratios of GFP-positive

cells among TBR1-positive cells are still low (15% at P15, and 26% at 4W) (Figure

14B-middle panels and Supplementary table S34). Soma sizes of TBR1/GFP-double positive

cells were also smaller than those of TBR1-negative/GFP-positive cells, and

TBR1/GFP-double positive cells showed lower intensity of GFP immunosignals compared to

TBR1-negative/GFP-positive cells (Figure 15-right panel and Supplementary table S36). We

additionally performed triple immunostaining for TBR1, Nav1.1, and ankyrinG on

Scn1a-GFP mice (Supplementary figure S11 and Supplementary tables S37-S39). Notably,

the ratios of Nav1.1-positive cells among TBR1-positive cells are 0% in all layers

(Supplementary figure S11B-right panel and Supplementary table S39). These results indicate

that the major population of TBR1-positive cells do not express Nav1.1.

To investigate whether TBR1-positive cells express Nav1.2, we performed triple

immunostaining of TBR1, Nav1.2 and ankyrinG on *Scn1a*-GFP mice (Supplementary figure

S12 and Supplementary tables S40-S43). In L5, contrary to the low ratios of GFP-positive

cells among TBR1-positive cells (11% at P15 and 5% at 4W) (Figure 14B-middle panels and

Supplementary table S34), the ratios of Nav1.2-positive cells among TBR1-positeve cells are

high (69% (L2/3), 69%(L5) and 69% (L6) at P15) (Supplementary figure S12B-right-upper

panel and Supplementary table S42). The ratios of TBR1-positive cells among

Nav1.2-positive cells are 29% (L2/3), 53% (L5) and 62% (L6) at P15 (Supplementary figure

S12B-middle upper panel and Supplementary table S41).

We further performed triple immunostaining for TBR1, Nav1.2 and GFP on *Scn1a*-GFP

mice (Supplementary figure S13 and Supplementary table S44) and found that most (88%) of

L6 TBR1-positive cells are GFP-negative.

Taken together, these results indicate that most TBR1-positive neurons including L6 CT

neurons do not express Nav1.1 but expresses Nav1.2.

As a whole, above results showed that a minor subpopulation of L2/3 CC and L5 PT

neurons express Nav1.1 while the majority of L2/3 CC, L5/6 CS and L6 CT neurons express

Nav1.2. A breakdown of L5 neuron species containing PT neurons is specifically described

(Supplementary figure S14).

Discussion

In our present study, we developed *Scn1a* promoter-driven GFP mice in which the expression

of GFP mimics that of Nav1.1. All PV-INs and most of SST-INs were GFP-positive in the

neocortex and hippocampus of the *Scn1a*-GFP mouse, being consistent with the previous

reports of Nav1.1 expression in those inhibitory neurons (Ogiwara et al., 2007; Lorincz and

Nusser, 2008; Ogiwara et al., 2013; Li et al., 2014; Tai et al., 2014; Tian et al., 2014;

Yamagata et al, 2017). All Nav1.1-positive cells were GFP-positive. Reversely all

GFP-positive cells were also Nav1.1-positive in the hippocampus, but in the neocortex only a

half of GFP-positive cells were Nav1.1-positive. This is largely because in the hippocampus

Nav1.1 expression is restricted to inhibitory neurons, but in the neocortex Nav1.1 is expressed

not only in inhibitory but also in a subpopulation of excitatory neurons in which Nav1.1

expression is low and not easily detected immunohistochemically by anti-Nav1.1 antibodies.

In neocortex, one third of GFP-positive cells were GABAergic cells such as PV-INs and

SST-INs, and remained two-third were excitatory neurons. GFP signals were especially

intense in PV-positive cells indicating strong Nav1.1 expression in those cells, while

GFP-signals in SST-positive cells were similar to those in excitatory neurons. In addition,

extensive immunostaining analyses using projection neuron markers FEZF2 and TBR1

together with anti-Nav1.2 antibody also revealed that most L5 PT neurons and a minor

subpopulation of L2/3 CC neurons express Nav1.1 while the majority of L6 CT, L5/6 CS and

L2/3 CC neurons express Nav1.2.

The above observations should contribute to understanding of neural circuits responsible

for diseases such as epilepsy and neurodevelopmental disorders caused by *SCN1A* and *SCN2A*

mutations. Dravet syndrome is a sporadic intractable epileptic encephalopathy characterized

by early onset (6 months ~ 1 year after birth) epileptic seizures, intellectual disability, autistic

features, ataxia and increased risk of sudden unexpected death in epilepsy (SUDEP). De novo

loss-of-function mutations of *SCN1A* are found in more than 80% of the patients (Claes et al.,

2001; Sugawara et al., 2002; Fujiwara et al., 2003; Dravet et al., 2005; Depienne et al., 2009;

Meng et al., 2015). In mice, loss-of-function *Scn1a* mutations caused clinical features

reminiscent of Dravet syndrome, including early-onset epileptic seizures, hyperactivity,

learning and memory deficits, reduced sociability and ataxic gaits and premature sudden

death (Yu et al., 2006; Ogiwara et al., 2007; Oakley et al., 2009; Cao et al., 2012; Han et al.,

2012; Kalume et al., 2013; Ito et al., 2013). As also shown in the present study, Nav1.1 is

densely localized at AISs of inhibitory cells such PV-IN (Ogiwara et al., 2007; Ogiwara et al.,

2013; Li et al., 2014; Tai et al., 2014) and selective elimination of Nav1.1 in PV-IN in mice

leads to epileptic seizures, sudden death and deficits in social behavior and spatial memory

(Ogiwara et al., 2013; Tatsukawa et al., 2018). It is thus plausible that Nav1.1

haplo-deficiency in PV-IN plays a pivotal role in the pathophysiology of many clinical

aspects of Dravet syndrome. Notably, mice with selective Nav1.1 haplo-elimination in global

inhibitory neurons were at a greater risk of lethal seizure than systemic Nav1.1

haplo-deficient mice, and the mortality risk of mice with Nav1.1 haplo-deficiency in

inhibitory neurons was significantly decreased or improved with additional Nav1.1

haplo-elimination in dorsal telencephalic excitatory neurons (Ogiwara et al., 2013), which

indicates beneficial effects of Nav1.1 deficiency in excitatory neurons for epileptic seizures

and sudden death. Because of the absence of Nav1.1 in hippocampal excitatory neurons

(Ogiwara et al., 2007; Ogiwara et al., 2013; Yamagata et al., 2017, and the present study), the

ameliorating effect was most possibly caused by Nav1.1 haploinsufficiency in neocortical

excitatory neurons. Kalume and colleagues (2013) reported that parasympathetic

hyperactivity is observed in Nav1.1 haplo-deficient mice and it causes ictal bradycardia and

finally result in seizure-associated sudden death.

Our present finding of Nav1.1 expression in L5 pyramidal tract projection neurons which

innervate the vagus nerve may possibly elucidate the ameliorating effects of Nav1.1

haploinsufficiency in neocortical excitatory neurons for sudden death of Nav1.1

haplo-deficient mice and may contribute to the understanding of the neural circuit for SUDEP

in patients with Dravet syndrome. Further studies including retrograde tracing and

electrophysiological analyses are awaited.

We previously proposed that impaired cortico-striatal excitatory neurotransmission causes

epilepsies in *Scn2a* haplodeficient mouse (Miyamoto et al., 2019). Our present finding of

Nav1.2 expression in CS neurons is consistent and further support the proposal. Because

SCN2A has been well established as one of top genes which show de novo loss-of-function

mutations in patients with ASD (Hoischen et al., 2014; Johnson et al., 2016) and because

impaired striatal function was suggested in multiple ASD animal models (Fuccillo et al.,

2016), our finding of Nav1.2 expression in CS neurons may also contribute to the

understanding of neural circuit for ASD caused by *SCN2A* mutations.

In summary, the present investigations using a newly-developed *Scn1a* promoter-driven

GFP mice together with anti-Nav1.1/Nav1.2 antibodies and neocortical neuron markers

revealed the expression loci of Nav1.1 and Nav1.2 in more detail. Further developments of

transgenic mice for other sodium channel genes' promoter-driven reporter molecules and

combinatorial analyses of those mice are awaited to segregate and redefine their unique

functional roles in each of highly diverse neuronal species and complexed neural circuits.

Materials and Methods

Animal work statement

All animal experimental protocols were approved by the Animal Experiment Committee of

Nagoya City University and RIKEN Center for Brain Science. Mice were handled in

accordance with the guidelines of the Animal Experiment Committee.

Mice

Scn1a-GFP BAC transgenic mice were generated as follows. A murine BAC clone

RP23-232A20 containing the *Scn1a* locus was obtained from the BACPAC Resource Center

(https://bacpacresources.org). A GFP reporter cassette, comprising a red-shifted variant GFP

cDNA and a downstream polyadenylation signal derived from pIRES2-EGFP (Takara Bio),

was inserted in-frame into the initiation codon of the *Scn1a* coding exon 1 using the Red/ET

Recombineering kit (Gene Bridges), according to the manufacturer's instructions. A correctly

modified BAC clone verified using PCR and restriction mapping was digested with *Sac*II,

purified using CL-4B sepharose (GE Healthcare), and injected into pronuclei of C57BL/6J

zygotes. Mice carrying the BAC transgene were identified using PCR with primers:

m*Scn1a*_TG_check_F1, 5'-TGTTCTCCACGTTTCTGGTT-3', m*Scn1a*_TG_check_R1,

5'-TTAGCCTTCTCTTCTGCAATG-3' and EGFP_R1, 5'-GCTCCTGGACGTAGCCTTC-3'

that detect the wild-type *Scn1a* allele as an internal control (186 bp) and the inserted

transgene (371 bp). Of fifteen independent founder lines that were crossed with C57BL/6J

mice, twelve lines successfully transmitted the transgene to their progeny. Of twelve founders,

two lines (#184 and 233) that display much stronger green fluorescent intensity compared

with other lines were selected, and maintained on a congenic C57BL/6J background. The

mouse lines had normal growth and development. The line #233 has been deposited to the

RIKEN BioResource Research Center (https://web.brc.riken.jp/en/) for distribution under the

registration number RBRC10241. *Vgat*-Cre BAC transgenic mice and loxP flanked

transcription terminator cassette CAG promotor driven tdTomato transgenic

(*Rosa26*-tdTomato; B6.Cg- Gt(ROSA)26Sortm14(CAG-tdTomato)Hze/J, Stock No: 007914,

The Jackson Laboratory, USA) mice were maintained on a C57BL/6J background. To

generate triple mutant mice (*Scn1a*-GFP$^{gfp/-}$, *Vgat*-Cre$^{cre/-}$, *Rosa26*-tdTomamto$^{tomato/-}$),

heterozygous *Scn1a*-GFP and *Vgat*-Cre mice were mated with homozygous *Rosa26*-tdTomato

transgenic mice.

Western blot analysis

Mouse brains at 5W were isolated and homogenized in homogenization buffer [(0.32 M

sucrose, 10 mM HEPES, 2 mM EDTA and 1× complete protease inhibitor cocktail (Roche

Diagnostics), pH 7.4)], and centrifuged for 15 min at 1,000 *g*. The supernatants were next

centrifuged for 30 min at 30,000 *g*. The resulting supernatants were designated as the cytosol

fraction. The pellets were subsequently resuspended in lysis buffer (50 mM HEPES and 2

mM EDTA, pH 7.4) and centrifuged for 30 min at 30,000 *g*. The resulting pellets, designated

as the membrane fraction, were dissolved in 2 M Urea, 1× NuPAGE reducing agent (Thermo

Fisher Scientific) and 1× NuPAGE LDS sample buffer (Thermo Fisher Scientific). The

cytosol and membrane fractions were separated on the NuPAGE Novex Tris-acetate 3–8% gel

(Thermo Fisher Scientific) or the PAG mini SuperSep Ace Tris-glycine 5–20% gel

(FUJIFILM Wako Pure Chemical), and transferred to nitrocellulose membranes (Bio-Rad).

Membranes were probed with the rabbit anti-Nav1.1 (250 ng/ml; IO1, Ogiwara et al., 2007),

chicken anti-GFP (1:5,000; ab13970, Abcam) and mouse anti-β tubulin (1:10,000; T0198,

Sigma-Aldrich) antibodies, and incubated with the horseradish peroxidase-conjugated goat

anti-rabbit IgG (1:2,000; sc-2004, Santa Cruz Biotechnology), rabbit anti-chicken IgY

(1:1,000; G1351, Promega) and goat anti-mouse IgG (1:5,000; W4011, Promega) antibodies.

Blots were detected using the enhanced chemiluminescence reagent (PerkinElmer). The

intensity of the Nav1.1 immunosignals were quantified using the Image Studio Lite software

(LI-COR, Lincoln, Nebraska USA) and normalized to the level of β tubulin or

Glyceraldehyde 3-phosphate dehydrogenase.

Histochemistry

Mice were deeply anesthetized, perfused transcardially with 4% paraformaldehyde (PFA) in

PBS (10 mM phosphate buffer, 2.7 mM KCl, and 137 mM NaCl, pH 7.4) or

periodate-lysine-4% PFA (PLP). Brains were removed from the skull and post-fixed. For

fluorescent imaging, PFA-fixed brains were cryoprotected with 30% sucrose in PBS, cut in 30

µm parasagittal sections, and mounted on glass slides. The sections on glass slides were

treated with TrueBlack Lipofuscin Autofluorescence Quencher (Biotium) to reduce

background fluorescence. For immunostaining, frozen parasagittal sections (30 µm) were

blocked with 4% BlockAce (DS Pharma Biomedical) in PBS for 1 hour at room temperature

(RT), and incubated with rat anti-GFP (1:500; GF090R, Nacalai Tesque). The sections were

then incubated with the secondary antibodies conjugated with biotin (1:200; Vector

Laboratories). The antibody-antigen complexes were visualized using the Vectastain Elite

ABC kit (Vector Laboratories) with Metal enhanced DAB substrate (34065, PIERCE). For

immunofluorescent staining, we prepared 6 µm parasagittal sections from paraffin embedded

PLP-fixed brains of mice. The sections were processed as previously described (Yamagata et

al., 2017). Following antibodies were used to detect GFP, Nav1.1, Nav1.2, TBR1, FEZF2,

ankyrinG, NeuN, parvalbumin and somatostatin; mouse anti-GFP antibodies (1:500;

11814460001, Roche Diagnostics), anti-Nav1.1 antibodies (1:10,000; rabbit IO1, 1:500; goat

SC-16031, Santa Cruz Biotechnology), anti-Nav1.2 antibodies (1:1,000; rabbit ASC-002,

Alomone Labs; goat SC-31371, Santa Cruz Biotechnology), rabbit anti-TBR1 antibody

(1:1,000; ab31940, Abcam or 1:500; SC-376258, Santa Cruz Biotechnology), rabbit

anti-FEZF2 antibody (1:500; #18997, IBL), ankyrinG antibodies (1:500; mouse SC-12719,

rabbit SC-28561; goat, SC-31778, Santa Cruz Biotechnology), mouse anti-NeuN biotin

conjugated antibody (1:2,000; MAB377B, Millipore), rabbit anti-parvalbumin (1:5,000;

PC255L, Merck) and rabbit anti-somatostatin (1:5,000; T-4103, Peninsula Laboratories,

1:1,000; SC-7819, Santa Cruz Biotechnology) antibodies. As secondary antibodies, Alexa

Fluor Plus 488, 555, 594 and 647 conjugated antibodies (1:1,000; A32723, A32766, A32794,

A32754, A32849, A32795, A32787, Thermo Fisher Scientific) were used. To detect NeuN,

Alexa-647 conjugated streptavidin (1:1,000; S21374, Thermo Fisher Science) was used.

Images were captured using fluorescence microscopes (BZ-8100 and BZ-X710, Keyence),

and processed with Adobe Photoshop Elements 10 (Adobe Systems) and BZ-X analyzer

(Keyence).

Fluorescence and Immunofluorescence quantification

For quantification of inhibitory neurons in GFP-positive cells, we used

Scn1a-GFP/*Vgat*-Cre/*Rosa26*-tdTomamto mice at 4W. We acquired multiple color images of

primary motor cortex and hippocampus from three parasagittal sections per animal. Six

images per region of interest were manually counted and summarized using Adobe Photoshop

Elements 10 and Excel (Microsoft). On immunofluorescence quantification, we used

Scn1a-GFP mice at P15 and/or 4W for the quantification of immunosignals. For

quantification of GFP, NeuN, PV, SST, Nav1.1, Nav1.2, FEZF2 or TBR1-positive cells, we

acquired multiple color images of primary motor cortex and hippocampus from three

parasagittal sections per animal. Six ~ nine images per region of interest were manually

quantified and summarized. For quantification of PV, SST, FEZF2 or TBR1-positive cells,

intensity and area size of GFP fluorescent signals were measured by Fiji software. Statistical

analyses were performed by one-way ANOVA followed by Tukey–Kramer post-hoc multiple

comparison test using Kyplot 6.0 (KyensLab Inc.). P-value smaller than 0.05 was considered

statistically significant. Data are presented as the mean ± standard error of the mean (SEM).

***In situ* hybridization**

Frozen sections (30 µm) of PFA-fixed mouse brains at 4W were incubated in 0.3% H_2O_2

in PBS for 30 min at RT to quench endogenous peroxidases, and mounted on glass slides. The

sections on slides were UV irradiated with 1,250 mJ/cm^2 (Bio-Rad), permeabilized with 0.3%

TritonX-100 in PBS for 15 min at RT, and digested with 1 µg/ml proteinase K (Nacalai

Tesque) in 10mM Tris-HCl and 1mM EDTA, pH 8.0, for 30 min at 37°C, washed twice with

100 mM glycine in PBS for 5 min at RT, fixed with 4% formaldehyde in PBS for 5 min at RT,

and acetylated with 0.25% acetic anhydride in 100 mM triethanolamine, pH8.0. After

acetylation, the sections were washed twice with 0.1 M phosphate buffer, pH 8.0, incubated in

a hybridization buffer [(50% formamide, 5× SSPE, 0.1% SDS, and 1 mg/ml Yeast tRNA

(Roche Diagnostics)] containing the Avidin solution (Vector Laboratories) for 2 hr at 60°C,

and hybridized with 2 µg/ml digoxigenin (DIG)- and dinitrophenol (DNP)-labeled probes in a

hybridization buffer containing the Biotin solution (Vector Laboratories) overnight at 60°C in

a humidified chamber. The hybridized sections were washed with 50% formamide in 2× SSC

for 15 min at 50°C twice, incubated in TNE (1 mM EDTA, 500 mM NaCl, 10 mM Tris-HCl,

pH 8.0) for 10 min at 37°C, treated with 20 µg/ml RNase A (Nacalai Tesque) in TNE for 15

min at 37°C, washed 2× SSC twice for 15 min each at 37°C twice and 0.2× SSC twice for 15

min each at 37°C. After washing twice in a high stringency buffer (10 mM Tris, 10 mM

EDTA and 500 mM NaCl, pH 8.0) for 10 min each at RT, the sections were blocked with a

blocking buffer [20 mM Tris and 150 mM NaCl, pH 7.5 containing 0.05% Tween-20, 4%

BlockAce (DS Pharma Biomedical) and 0.5× Blocking reagent (Roche Diagnostics)] for 1 hr

at RT, and incubated with the alkaline phosphatase-conjugated sheep anti-DIG (1:500;

11093274910, Roche Diagnostics) and biotinylated rabbit anti-DNP (1:100; BA-0603, Vector

laboratories) antibodies in a blocking buffer overnight at 4°C, followed by incubation with the

biotinylated goat anti-rabbit antibody (1:200; BA-1000, Vector laboratories) in a blocking

buffer at RT for 1 to 2hr. The probes were visualized using the NBT/BCIP kit (Roche

Diagnostics), VECTASTAIN Elite ABC kit (Vector laboratories) and ImmPACT DAB

substrate (Vector laboratories).

The DIG-labeled RNA probes for *Scn1a* designed to target the 3'-untranlated region

(nucleotides 6,488–7,102 from accession number NM_001313997.1) were described

previously (Ogiwara et al., 2007), and synthesized using the MEGAscript transcription kits

(Thermo Fisher Scientific) with DIG-11-UTP (Roche Diagnostics). The DNP-labeled RNA

probes for GFP were derived from the fragment corresponding to nucleotides 1,256–1,983 in

pIRES2-EGFP (Takara Bio), and prepared using the MEGAscript transcription kits (Thermo

Fisher Scientific) with DNP-11-UTP (PerkinElmer).

Acknowledgment

We thank Dr. Yaguchi (Laboratory for Behavioral Genetics) and the staff members at the

Research Resources Division of RIKEN Center for Brain Science for technical assistance in

generating *Scn1a*-GFP BAC Tg mice and Dr. Kaneda (Nippon Medical School) for his

support. This study was supported in part by MEXT/JSPS KAKENHI JP20H03566,

JP17H01564, JP16H06276, AMED JP18dm0107092, RIKEN Center for Brain Science

(K.Y.); Takeda Science Foundation, Kiyokun Foundation (I.O. and K.Y.); MEXT/JSPS

KAKENHI JP19790747, JP21791020, JP16K15564, JP19K08284 (I.O.); and Japan Epilepsy

Research Foundation (I.O. and T.T.).

Competing interests

No competing interests declared.

588
589

References

1. Brysch W, Creutzfeldt OD, Lüno K, Schlingensiepen R, Schlingensiepen KH. Regional and temporal expression of sodium channel messenger RNAs in the rat brain during development. *Exp Brain Res.* **86**(3): 562-567 (1991).

2. Cao D, Ohtani H, Ogiwara I, Ohtani S, Takahashi Y, Yamakawa K, Inoue Y. Efficacy of stiripentol in hyperthermia-induced seizures in a mouse model of Dravet syndrome. *Epilepsia.* **53**, 1140-1145 (2012).

3. Catterall WA. Voltage-gated sodium channels at 60: structure, function and pathophysiology. *J. Physiol.* **590**(11):2577-2589 (2012).

4. Chen B, Schaevitz LR., McConnell SK. Fezl regulates the differentiation and axon targeting of layer 5 subcortical projection neurons in cerebral cortex. *Proc. Natl. Acad. Sci. USA.* **102**:17184–17189 (2005).

5. Chen B, Wang SS, Hattox AM, Rayburn H, Nelson SB, McConnell SK. The Fezf2-Ctip2 genetic pathway regulates the fate choice of subcortical projection neurons in the developing cerebral cortex. *Proc. Natl. Acad. Sci. USA.* **105**(32):11382-11387 (2008).

6. Claes L, Del-Favero J, Ceulemans B, Lagae L, Van Broeckhoven C, De Jonghe P. De novo mutations in the sodium-channel gene *SCN1A* cause severe myoclonic epilepsy of infancy. *Am. J. Hum. Genet.* **68,** 1327–1332 (2001).

7. Depienne C, Trouillard O, Saint-Martin C, Gourfinkel-An I, Bouteiller D, Carpentier W, Keren B, Abert B, Gautier A, Baulac S, Arzimanoglou A, Cazeneuve C, Nabbout R, LeGuern E. Spectrum of *SCN1A* gene mutations associated with Dravet syndrome: analysis of 333 patients. *J. Med. Genet.* **46**, 183–191 (2009).

8. Dravet C, Bureau M, Oguni H, Fukuyama Y, Cokar O. Severe myoclonic epilepsy in infancy (Dravet syndrome) in *Epileptic Syndromes in Infancy, Childhood and Adolescence, 4th edn* (eds. Roger, J. *et al.*) 89–113 (John Libbey Eurotext Ltd, Montrouge, France, 2005).

9. Fuccillo MV. Striatal Circuits as a Common Node for Autism Pathophysiology. *Front. Neurosci.* **10**:27 (2016).

10. Fujiwara T, Sugawara T, Mazaki-Miyazaki E, Takahashi Y, Fukushima K, Watanabe M, Hara K, Morikawa T, Yagi K, Yamakawa K, Inoue Y. Mutations of sodium channel α subunit type 1 (*SCN1A*) in intractable childhood epilepsies with frequent generalized tonic-clonic seizures. *Brain.* **126**, 531–546 (2003).

11. Goff KM, Goldberg EM. Vasoactive intestinal peptide-expressing interneurons are impaired in a mouse model of Dravet syndrome. *Elife.* **8,** e46846 (2019).

12. Han S, Tai C, Westenbroek RE, Yu FH, Cheah CS, Potter GB, Rubenstein JL, Scheuer T,
de la Iglesia HO, Catterall WA. Autistic-like behaviour in *Scn1a*$^{+/-}$ mice and rescue by
enhanced GABA-mediated neurotransmission. *Nature*. **489**, 385–390 (2012).

13. Han W, Kwan KY, Shim S, Lam MMS, Shin Y, Xu X, Zhu Y, Li M, Sestan N. TBR1
directly represses Fezf2 to control the laminar origin and development of the corticospinal
tract. *Proc. Natl. Acad. Sci. USA* **108**(7), 3041-3046 (2011)

14. Hoischen A, Krumm N, Eichler EE. Prioritization of neurodevelopmental disease genes
by discovery of new mutations. *Nat. Neurosci*. **17**(6):764-772 (2014).

15. Ito S, Ogiwara I, Yamada K, Miyamoto H, Hensch TK, Osawa M, Yamakawa K. Mouse
with Nav1.1 haploinsufficiency, a model for Dravet syndrome, exhibits lowered
sociability and learning impairment. *Neurobiol. Dis*. **49**, 29–40 (2013).

16. Inoue K, Terashima T, Nishikawa T, Takumi T. Fez1 is layer-specifically expressed in the
adult mouse neocortex. *Eur J Neurosci*. **20**(11):2909-16 (2004).

17. Jinno S, Kosaka T. Parvalbumin is expressed in glutamatergic and GABAergic
corticostriatal pathway in mice. *J Comp Neurol*. **477**(2):188-201 (2004).

18. Johnson MR, Shkura K, Langley SR, Delahaye-Duriez A, Srivastava P, Hill WD,
Rackham OJ, Davies G, Harris SE, Moreno-Moral A, Rotival M, Speed D, Petrovski S,
Katz A, Hayward C, Porteous DJ, Smith BH, Padmanabhan S, Hocking LJ, Starr JM,
Liewald DC, Visconti A, Falchi M, Bottolo L, Rossetti T, Danis B, Mazzuferi M, Foerch
P, Grote A, Helmstaedter C, Becker AJ, Kaminski RM, Deary IJ, Petretto E. Systems
genetics identifies a convergent gene network for cognition and neurodevelopmental
disease. *Nat. Neurosci*. **19**(2):223-232 (2016).

19. Kalume F, Westenbroek RE, Cheah CS, Yu FH, Oakley JC, Scheuer T, Catterall WA.
Sudden unexpected death in a mouse model of Dravet syndrome. *J. Clin. Invest*. **123**,
1798–1808 (2013).

20. Li T, Tian C, Scalmani P, Frassoni C, Mantegazza M, Wang Y, Yang M, Wu S, Shu Y.
Action potential initiation in neocortical inhibitory interneurons. *PLoS Biol*. **12,** e1001944:
(2014).

21. Lodato S, Molyneaux BJ, Zuccaro E, Goff LA, Chen HH, Yuan W, Meleski A, Takahashi
E, Mahony S, Rinn JL, Gifford DK, Arlotta P. Gene co-regulation by Fezf2 selects
neurotransmitter identity and connectivity of corticospinal neurons. *Nat Neurosci*.
17(8):1046-54 (2014).

22. Lorincz A, Nusser Z. Cell-type-dependent molecular composition of the axon initial
segment. *J. Neurosci*. **28**(53), 14329-14340 (2008).

23. Lorincz A, Nusser Z. Molecular identity of dendritic voltage-gated sodium channels.
Science **328**(5980):906-909 (2010).

24. Matho KS, Huilgol D, Galbavy W, He M, Kim G, An X, Lu J, Wu P, Di Bella DJ, Shetty
AS, Palaniswamy R, Hatfield J, Raudales R, Narasimhan A, Gamache E, Levine JM,
Tucciarone J, Szelenyi E, Harris JA, Mitra PP, Osten P, Arlotta P, Huang ZJ. Genetic
dissection of the glutamatergic neuron system in cerebral cortex. *Nature*.
598(7879):182-187 (2021)

25. McKenna WL, Betancourt J, Larkin KA, Abrams B, Guo C, Rubenstein JL, Chen B. Tbr1
and Fezf2 regulate alternate corticofugal neuronal identities during neocortical
development. *J. Neurosci*. **31**(2):549-564 (2011).

26. Meisler MH, Hill SF, Yu W. Sodium channelopathies in neurodevelopmental disorders.
Nat. Rev. Neurosci. **22**(3):152-166 (2021).

27. Meng H, Xu HQ, Yu L, Lin GW, He N, Su T, Shi YW, Li B, Wang J, Liu XR, Tang B,
Long YS, Yi YH, Liao WP. The *SCN1A* mutation database: updating information and
analysis of the relationships among genotype, functional alteration, and phenotype. *Hum.*
Mutat. 36, 573–580 (2015).

28. Miyamoto H, Tatsukawa T, Shimohata A, Yamagata T, Suzuki T, Amano K, Mazaki E,
Raveau M, Ogiwara I, Oba-Asaka A, Hensch TK, Itohara S, Sakimura K, Kobayashi K,
Kobayashi K, Yamakawa K. Impaired cortico-striatal excitatory transmission triggers
epilepsy. *Nat. Commun*. **10**(1):1917 (2019).

29. Molyneaux BJ, Arlotta P, Hirata T, Hibi M, Macklis JD. Fezl is required for the birth and
specification of corticospinal motor neurons. *Neuron.* **47**:817–831 (2005).

30. Nakayama T, Ogiwara I, Ito K, Kaneda M, Mazaki E, Osaka H, Ohtani H, Inoue Y,
Fujiwara T, Uematsu M, Haginoya K, Tsuchiya S, Yamakawa K. Deletions of SCN1A 5'
genomic region with promoter activity in Dravet syndrome. *Hum. Mutat*. **31**(7):820-829
(2010).

31. Oakley JC, Kalume F, Yu FH, Scheuer T, Catterall WA. Temperature- and age-dependent
seizures in a mouse model of severe myoclonic epilepsy in infancy. *Proc. Natl. Acad. Sci.*
USA. **106**, 3994–3999 (2009).

32. Ogiwara I, Miyamoto H, Morita N, Atapour N, Mazaki E, Inoue I, Takeuchi T, Itohara S,
Yanagawa Y, Obata K, Furuichi T, Hensch TK, Yamakawa K. Nav1.1 Localizes to axons
of parvalbumin-positive inhibitory interneurons: A circuit basis for epileptic seizures in
mice carrying an *Scn1a* gene mutation. *J. Neurosci*. **27**, 5903–5914 (2007)

33. Ogiwara I, Iwasato T, Miyamoto H, Iwata R, Yamagata T, Mazaki E, Yanagawa Y,
Tamamaki N, Hensch TK, Itohara S, Yamakawa K. Nav1.1 haploinsufficiency in
excitatory neurons ameliorates seizure-associated sudden death in a mouse model of
Dravet syndrome. *Hum. Mol. Genet.* **22**, 4784–4804 (2013).
34. Ogiwara I, Miyamoto H, Tatsukawa T, Yamagata T, Nakayama T, Atapour N, Miura E,
Mazaki E, Ernst SJ, Cao D, Ohtani H, Itohara S, Yanagawa Y, Montal M, Yuzaki M,
Inoue Y, Hensch TK, Noebels JL, Yamakawa K. Nav1.2 haplodeficiency in excitatory
neurons causes absence-like seizures in mice. *Commun. Biol.* **1**:96 (2018).
35. Paxinos G, Franklin BJ. The mouse brain in stereotaxic coordinates. Academic Press
(2001).
36. Shepherd GM. Corticostriatal connectivity and its role in disease. *Nat. Rev. Neurosci.* **14**,
278-291 (2013).
37. Sugawara T, Mazaki-Miyazaki E, Fukushima K, Shimomura J, Fujiwara T, Hamano S,
Inoue Y, Yamakawa K. Frequent mutations of *SCN1A* in severe myoclonic epilepsy in
infancy. *Neurology* **58**, 1122–1124 (2002).
38. Tai T, Abe Y, Westenbroek RE, Scheuer T, Catterall WA. Impaired excitability of
somatostatin- and parvalbumin-expressing cortical interneurons in a mouse model of
Dravet syndrome. *Proc. Natl. Acad. Sci. USA.* **111**, E3139–3148 (2014).
39. Tanahira C, Higo S, Watanabe K, Tomioka R, Ebihara S, Kaneko T, Tamamaki N.
Parvalbumin neurons in the forebrain as revealed by parvalbumin-Cre transgenic mice.
Neurosci Res. **63**(3):213-23 (2009).
40. Tatsukawa T, Ogiwara I, Mazaki E, Shimohata A, Yamakawa K. Impairments in social
novelty recognition and spatial memory in mice with conditional deletion of *Scn1a* in
parvalbumin-expressing cells. *Neurobiol. Dis.* **112**, 24–34 (2018).
41. Tian C, Wang K, Ke W, Guo H, Shu Y. Molecular identity of axonal sodium channels in
human cortical pyramidal cells. *Front Cell Neurosci.* **8**: 297 (2014).
42. Weyer A, Schilling K. Developmental and cell type-specific expression of the neuronal
marker NeuN in the murine cerebellum. *J Neurosci Res.* **73**, 400-409 (2003).
43. Yamagata T, Ogiwara I, Mazaki E, Yanagawa Y, Yamakawa K. Nav1.2 is expressed in
caudal ganglionic eminence-derived disinhibitory interneurons: Mutually exclusive
distributions of Nav1.1 and Nav1.2. *Biochem. Biophys. Res. Commun.* **491**, 1070–1076
(2017).
44. Yamakawa, K. Mutations of voltage-gated sodium channel genes *SCN1A* and *SCN2A* in
epilepsy, intellectual disability, and autism in *Neuronal and synaptic dysfunction in*

autism spectrum disorder and intellectual disability (eds. Sara, C. & Verpelli, C.)

233–251 (Elsevier, Amsterdam, Netherlands, 2016).

45. Yu FH, Mantegazza M, Westenbroek RE, Robbins CA, Kalume F, Burton KA, Spain WJ,

McKnight GS, Scheuer T, Catterall WA. Reduced sodium current in GABAergic

interneurons in a mouse model of severe myoclonic epilepsy in infancy. *Nat. Neurosci.* **9**,

1142–1149 (2006).

Figure legends

Figure 1. Generation of *Scn1a*-GFP mice.

(**A**) Schematic representation of the modified BAC construct containing the *Scn1a*-GFP

transgene. A GFP reporter cassette consisting of GFP cDNA and a polyadenylation signal was

inserted at the ATG initiation codon in the coding exon 1 of *Scn1a*. Filled and hatched boxes

indicate the coding and non-coding exons of *Scn9a* and *Scn1a*. Arrows indicate the start sites

and orientation of transcription of *Scn1a*. (**B**) Western blot analysis for *Scn1a*-GFP and

endogenous Nav1.1. The whole cytosolic fractions from 5W *Scn1a*-GFP brains (lines #184

and #233) were probed with anti-GFP and their membrane fractions were probed with

anti-Nav1.1 antibodies. β-Tubulin was used as an internal control. pA, polyadenylation signal;

Tg, hemizygous *Scn1a*-GFP transgenic mice; Wt, wild-type littermates.

Figure 2. Distributions of GFP signals in brains are similar among *Scn1a*-GFP mouse

lines.

Chromogenic immunostaining of GFP (brown) with Nissl counterstaining (violet) of lines

#184 and #233 (A-H) and GFP fluorescence images of line #233 (I-L) on parasagittal sections

from 5–6W *Scn1a*-GFP brains. Boxed areas in (A, E, I, B, F, J, C, G, K, D, H, L) are

magnified in (B-D, F-H, J-L, B1, F1, J1, C1-3, G1-3, K1-3, D1, H1, L1). The two lines (lines

#184 and #233) showed a similar distribution pattern of GFP-expressing cells across all brain

regions (A-H), but the signals in the line #233 are more intense than the line #184. In

neocortex (B, F, J), GFP-expressing cells were scattered throughout the entire region. In the

hippocampus (C, G, K), GFP-positive inhibitory neurons were sparsely distributed (see also

Figure 8), while excitatory neurons in stratum pyramidale and stratum granulosum are

GFP-negative. In cerebellum (D, H, L), Purkinje, basket, and deep cerebellar nuclei cells were

GFP-positive. IHC, immunohistochemistry; CA, cornu ammonis; DG, dentate gyrus; o,

stratum oriens; p, stratum pyramidale; r, stratum radiatum; lm, stratum

lacunosum-moleculare; l, stratum lucidum; m, stratum moleculare; g, stratum granulosum; h,

hilus; DCN, deep cerebellar nuclei; M, molecular layer; P, Purkinje cell layer; G, granular cell

layer. Scale bars; 1 mm (A, E, I), 500 μm (C, D, G, H, K, L), 100 μm (B, F, J), 50 μm (B1,

C1-3, D1, F1, G1-3, H1, J1, K1-3, L1).

Figure 3. Distribution of GFP signals in *Scn1a*-GFP mouse brain are largely maintained

through development.

Fluorescent images of parasagittal sections from P15 (**A**), 4W (**B**) and 8W (**C**) *Scn1a*-GFP

mouse brains (line #233). GFP signals were observed in multiple brain regions. APT, anterior

pretectal nucleus; CPu, caudate putamen; Cx, cerebral cortex; DCN, deep cerebellar nuclei;

HP, hippocampus; IC, inferior colliculus; Mo, medulla oblongata; Ob, olfactory bulb; P, pons;

RT, reticular thalamic nucleus; SC, superior colliculus; STN, subthalamic nucleus; VPM,

ventral posteromedial thalamic nucleus; ZI, zona incerta. Scale bars; 1 mm.

Figure 4. All cells with Nav1.1-positive AISs are GFP-positive but only half or less

population of AISs for GFP-positive cells are Nav1.1-positive in *Scn1a*-GFP mouse

neocortex.

(**A**) Triple immunofluorescent staining of parasagittal section from P15 *Scn1a*-GFP mouse

brain (line #233) by mouse anti-GFP (green), rabbit anti-Nav1.1 (magenta), and goat

anti-ankyrinG (cyan) antibodies. Regions at primary motor cortex are shown. (**B**, **C**)

Magnified images outlined in (A) are shown in (B) and (C). Arrows indicate AISs of cells

with GFP-positive somata in which both somata and AISs are positive for Nav1.1.

Arrowheads indicate AISs of cells with GFP-positive somata in which AISs but not somata

are positive for Nav1.1. All images are oriented from pial surface (top) to callosal (bottom).

Scale bars; 100 μm (A), 50 μm (B and C). (**D**) Cell counting of three *Scn1a*-GFP mice. Bar

graphs indicating the percentage of cells with GFP- and Nav1.1-positive/negative somata and

AISs per cells with ankyrinG-positive AISs (left panel), the percentage of cells with

GFP-positive/negative somata per cells with ankyrinG-positive AISs and Nav1.1-positive

somata and/or AISs (middle panel), and the percentage of cells with Nav1.1-positive/negative

somata and/or AISs per cells with ankyrinG-positive AISs and GFP-positive somata (right

panel) in L2/3, L5, and L6 (see also Supplementary tables S2-S4). Only cells with

ankyrinG-positive AISs were counted. Nav1.1 immunosignals were occasionally observed in

somata, but in such cases Nav1.1 signals were always observed in their AISs if visible by

ankyrinG staining. Note that 99% (L2/3), 99% (L5) and 97% (L6) of cells with

Nav1.1-positive AISs have GFP-positive somata (middle panel), but only half or less of cells

with GFP-positive somata have Nav1.1-positive AISs (right panel). L2/3, L5: neocortical

layer II/III and V. AnkG, ankyrinG; +, positive; -, negative.

Figure 5. All cells with Nav1.1-positive AISs are GFP-positive and all AISs of

GFP-positive cells are Nav1.1-positive in *Scn1a*-GFP mouse hippocampus.

(**A**-**D**) Triple immunofluorescent staining of parasagittal section from P15 *Scn1a*-GFP mouse

brain (line #233) by mouse anti-GFP (green), rabbit anti-Nav1.1 (magenta), and goat

anti-ankyrinG (cyan) antibodies. Regions at hippocampus were shown. Note that GFP and

Nav1.1 immunosignals mostly overlap at somata. CA1, cornu ammonis 1; CA2, cornu

ammonis 2; CA3, cornu ammonis 3; DG, dentate gyrus. Images are oriented from pial surface

(top) to callosal (bottom). Scale bars; 100 μm. (**E**) Magnified images for co-expression of

GFP and Nav1.1 in cells at CA1 region. Arrowheads indicate Nav1.1-positive AISs of

GFP-expression cells. Scale bar; 50 μm. (**F**) Bar graphs indicate the percentage of cells in

hippocampal CA1 region with Nav1.1-positive/negative AISs per cells with GFP-positive

somata and ankyrinG-positive AISs (left panel), and the percentage of cells with

GFP-positive/negative somata per cells with Nav1.1/ankyrinG-double positive AISs (right

panel) (see also Supplementary tables S7, S8). Only cells with ankyrinG-positive AISs were

counted. GFP/Nav1.1-double negative cells, most of which are pyramidal cells, were not

counted because of the accumulated nature of their ankyrinG-positive AISs. AnkG, ankyrinG;

+, positive; -, negative.

Figure 6. GFP and *Scn1a* mRNAs expressions goodly overlap in *Scn1a*-GFP mouse

brain.

Double *in situ* hybridization for *Scn1a*-GFP transgene mRNA and endogenous *Scn1a* mRNA

on parasagittal sections from 4W *Scn1a*-GFP brains (line #233). (**A**) Sections were hybridized

with antisense (left) and sense (right) RNA probes for GFP transgene (brown) and

endogenous *Scn1a* (blue) mRNA species and chromogenically stained. Magnified images

outlined in (A) are shown in (**B-D**) for antisense probes, and (**E-G**) for sense probes. o,

stratum oriens; p, stratum pyramidale; r, stratum radiatum; lm, stratum

lacunosum-moleculare; m, stratum moleculare; g, stratum granulosum, CA1, cornu ammonis

1; DG, dentate gyrus. Scale bars; 500 μm (A), 50 μm (B-G).

Figure 7. One third of GFP-positive cells in neocortex are inhibitory neurons, but most

of GFP-positive cells in hippocampus are inhibitory neurons.

(**A, B**) GFP (green) and Tomato (magenta) fluorescent images of parasagittal sections from

4W *Scn1a*-GFP/*Vgat*-cre/*Rosa26*-tdTomato mouse. Regions at primary motor cortex (A) and

hippocampus (B) are shown. Scale bar; 100 μm. (**C**) Bar graphs indicate the percentage of

cells with Tomato-positive/negative somata per cells with GFP-positive somata (left panel)

(see also Supplementary table S9) and the percentage of cells with GFP-positive/negative

somata per cells with Tomato-positive somata (right panel) (see also Supplementary table

S10) in L2/3, L5, L6, CA1 and DG. Cells in primary motor cortex and hippocampus of

Scn1a-GFP mouse at 4W were counted. L2/3, L5, L6, CA1 and DG: neocortical layer II/III, V,

VI, cornu ammonis 1, dentate gyrus. +, positive; -, negative.

Figure 8. Parvalbumin or somatostatin-positive inhibitory neurons are GFP-positive in

***Scn1a*-GFP mouse neocortex and hippocampus.**

(**A**) Triple immunofluorescent staining of parasagittal sections from 4W *Scn1a*-GFP mouse

(line #233) by mouse anti-GFP (green), rabbit anti-parvalbumin (PV) (magenta) and goat

anti-somatostatin (SST) (Cyan) antibodies. Regions at neocortex and hippocampus are shown.

Merged images were shown in the right columns. Arrows indicate SST/GFP-double positive

cells. Arrowheads indicate PV/GFP-double positive. o, stratum oriens; p, stratum pyramidale;

r, stratum radiatum; h, hilus; g, stratum granulosum; m, stratum moleculare. All images are

oriented from pial surface (top) to callosal (bottom). Scale bars; 50 μm. (**B**) Bar graphs

indicate the percentage of cells with PV- and SST-positive/negative somata per cells with

GFP-positive somata (left panel) (see also Supplementary table S11), the percentage of cells

with GFP-positive/negative somata per cells with PV-positive somata (middle panel) (see also

Supplementary table S12), and the percentage of cells with GFP-positive/negative somata per

cells with SST-positive somata (right panel) (see also Supplementary table S13) in L2/3, L5,

L6, CA1, CA2/3 and DG. Cells in neocortex and hippocampus of *Scn1a*-GFP mouse at 4W

were counted. L2/3, L5, L6, CA1, CA2/3 and DG: neocortical layer II/III, V, VI, cornu

ammonis 1, 2 plus 3, dentate gyrus. +, positive; -, negative.

Figure 9. GFP-signals in parvalbumin-positive inhibitory neurons are higher than

PV-negative/GFP-positive cells in *Scn1a*-GFP mouse neocortex.

(**A**) Scatter plots of intensities and area sizes of GFP immunosignals in GFP-positive cells

with PV or SST-positive or negative somata. Cells at primary motor cortex (upper panels) and

hippocampus (lower panels) in parasagittal sections from 4W *Scn1a*-GFP mouse (line #233)

were analyzed. PV-positive (magenta circles) or SST-positive (black circles) and negative

(green circles) cells in neocortical L2/3, L5, and L6 or hippocampal CA1, CA2/3, and DG are

plotted (see also Supplementary table S14). (**B**) Box-plots represent values for the intensity

and area size in each cell type (see also Supplementary tables S15, S16). Cross marks indicate

mean values in each cell type. Statistical significance was assessed using one-way ANOVA

followed by Tukey–Kramer post-hoc multiple comparison test. *; $p < 0.05$, **; $p < 0.01$, ***;

p < 0.001. Note that GFP signal intensities of PV/GFP-double positive cells were significantly

higher than that of SST/GFP-double positive cells and PV/SST-negative/GFP-positive cells

(all layers), while GFP signal intensities of SST/GFP-double positive cells were similar to

PV/SST-negative/GFP-positive cells in neocortex. In hippocampus, GFP signal intensities of

SST/GFP-double positive cells were significantly lower than that of

SST-negative/GFP-positive cells at CA2/3 and DG. CA1, CA2/3 and DG: cornu ammonis 1,

2 plus 3, dentate gyrus. a.u., arbitrary unit; +, positive; -, negative.

Figure 10. Nav1.1 and Nav1.2 are mutually exclusive at AISs in mouse brain.

(**A**) Triple immunofluorescent staining on parasagittal sections from *Scn1a*-GFP mouse at

P15 by rabbit anti-Nav1.1 (green), goat anti-Nav1.2 (magenta), and mouse anti-ankyrinG

(cyan) antibodies. Merged images are shown in the right panels. Arrows indicate

Nav1.2-positive AISs. Arrowheads indicate Nav1.1-positive AISs. Note that there are no

Nav1.1/Nav1.2-double positive AISs. Images are oriented from pial surface (top) to callosal

(bottom). Scale bars; 50 μm. (**B**) Bar graphs indicating the percentage of Nav1.1 and

Nav1.2-positive/negative AISs per AISs detected by ankyrinG-staining in L2/3, L5, and L6 of

Scn1a-GFP mice. Note that Nav1.1/Nav1.2-double positive AISs were less than 0.5% of all

AISs in these layers (see Supplementary table S17). L2/3, L5, L6: neocortical layer II/III, V,

VI. AnkG, ankyrinG; +, positive; -, negative.

Figure 11. Cells with Nav1.2-positive AISs are mostly GFP-negative in *Scn1a*-GFP

mouse neocortex.

(**A**) Triple immunofluorescent staining of parasagittal sections from P15 *Scn1a*-GFP mouse

brain (line #233) by mouse anti-GFP (green), goat anti-Nav1.2 (magenta) and rabbit

anti-ankyrinG (cyan) antibodies. Merged images of the signals are shown in the right panels.

Arrows indicate Nav1.2-positive AISs of cells with GFP-negative somata. Arrowheads

indicate Nav1.2-negative AISs of cells with GFP-positive somata. All images are oriented

from pial surface (top) to callosal (bottom). Scale bars; 50μm. (**B**) Bar graphs indicating the

percentage of cells with GFP- and Nav1.2-positive/negative somata and AISs per cells with

ankyrinG-positive AISs (left panel) (see also Supplementary table S18), the percentage of

cells with GFP-positive/negative somata per cells with ankyrinG/Nav1.2-double positive AISs

(middle panel) (see also Supplementary table S19), and the percentage of cells with

Nav1.2-positive/negative AISs per cells with ankyrinG-positive AISs and GFP-positive

somata (right panel) (see also Supplementary table S20) in L2/3, L5, and L6. Note that 88%

(L2/3), 90% (L5) and 95% (L6) of cells with Nav1.2-positive AISs have GFP-negative

somata (middle panel), and 68% (L2/3), 83% (L5) and 86% (L6) of cells with GFP-positive

somata have Nav1.2-negative AISs (right panel). L2/3, L5, L6: neocortical layer II/III, V, VI.

AnkG, ankyrinG; +, positive; -, negative.

Figure 12. Cells positive for FEZF2 are mostly GFP-positive at L5 of *Scn1a*-GFP mouse

neocortex.

(**A**) Double immunostaining of FEZF2 and GFP in neocortex of 4W *Scn1a*-GFP mouse (line

#233) by rabbit anti-FEZF2 (magenta) and mouse anti-GFP (green) antibodies. Arrows

indicate FEZF2/GFP-double positive cells. Magnified images outlined in (A) are shown in

(A1-A3). Note that FEZF2 signals mostly overlap with GFP signals in L5. Many of the

remained GFP-positive/FEZF2-negative cells have intense GFP signals and are assumed to be

inhibitory neurons (see Figure 8). Scale bars; 100 μm (A), 50 μm (A1-A3). (**B**) Bar graphs

indicating the percentage of cells with FEZF2- and GFP-positive/negative nuclei and somata

per cells with GFP-positive somata and/or FEZF2-positive nuclei (left panels) (see also

Supplementary table S21), the percentage of cells with GFP-positive/negative somata per

cells with FEZF2-positive nuclei (middle panels) (see also Supplementary table S22), and the

percentage of cells with FEZF2-positive/negative nuclei per cells with GFP-positive somata

(right panels) (see also Supplementary table S23) in L2/3, L5, and L6. Cells at primary motor

cortex of *Scn1a*-GFP mouse at P15 and 4W were counted. Note that 83% (P15) and 96%

(4W) of cells with FEZF2-positive cells are GFP-positive in L5 (middle panels), but a half of

cells with GFP-positive cells are FEZF2-positive in L5 (right panel). L2/3, L5, L6: neocortical

layer II/III, V, VI. +, positive; -, negative.

Figure 13. FEZF2-positive cells have lower GFP signal intensities in *Scn1a*-GFP mouse

neocortex.

Scatter plots of intensities and area sizes of GFP immunosignals in GFP-positive cells with

FEZF2-positive/negative nuclei. Cells at primary motor cortex in parasagittal sections from

4W *Scn1a*-GFP mouse (line #233) were analyzed. FEZF2-positive (orange squares) and

negative (black circles) cells in neocortical L2/3, L5, and L6 are plotted (see also

Supplementary table S24). Note that GFP signal intensities of FEZF2/GFP-double positive

cells were significantly lower than that of FEZF2-negative/GFP-positive cells (L5, L6), and

signal area size of FEZF2/GFP-double positive cells were significantly larger than that of

FEZF2-negative/GFP-positive cells (L2/3, L5). Statistical significance was assessed using

t-test. a.u., arbitrary unit; +, positive; -, negative.

Figure 14. GFP-positive cells were mostly negative for TBR1 at L5 of *Scn1a*-GFP mouse

neocortex.

(**A**) Double immunostaining of TBR1 and GFP in neocortex of 4W *Scn1a*-GFP mouse (line

#233) detected by mouse rabbit anti-TBR1 (magenta) and anti-GFP (green) antibodies.

Arrows indicate TBR1-negative/GFP-positive cells. Magnified images outlined in (A) are

shown in (A1-A6). Note that at L5 GFP-positive cells were mostly TBR1-negative but at L2/3

more than half of GFP-positive cells were TBR1-positive. Scale bars; 100 μm (A), 50 μm

(A1-A6). (**B**) Bar graphs indicating the percentage of cells with TBR1- and

GFP-positive/negative nuclei and somata per GFP-positive cells and/or TBR1-positive nuclei

(left panels) (see also Supplementary table S33), the percentage of cells with

GFP-positive/negative somata per cells with TBR1-positive nuclei (middle panels) (see also

Supplementary table S34), and the percentage of cells with TBR1-positive/negative nuclei per

cells with GFP-positive somata (right panels) (see also Supplementary table S35) in L2/3, L5,

and L6. Cells in primary motor cortex of *Scn1a*-GFP mouse at P15 and 4W were counted.

Note that 86% (P15) and 95% (4W) of cells with TBR1-positive cells are GFP-negative in L5

(middle panels), and 90% (P15) and 96% (4W) of cells with GFP-positive cells are

TBR1-negative in L5 (right panel). L2/3, L5, L6: neocortical layer II/III, V, VI. +, positive; -,

negative.

Figure 15. TBR1-positive cells have lower GFP signal intensities in *Scn1a*-GFP mouse

neocortex.

Scatter plots of intensities and area sizes of GFP immunosignals in GFP-positive cells with

TBR1-positive/negative nuclei. Cells at primary motor cortex in parasagittal sections from

4W *Scn1a*-GFP mouse (line #233) were analyzed. TBR1-positive (orange squares) and

negative (black circles) cells in neocortical L2/3, L5, and L6 are plotted (see also

Supplementary table S36). Note that GFP signal intensities of TBR1/GFP-double positive

cells were significantly lower than that of TBR1-negative/GFP-positive cells (L6), and signal

area size of TBR1/GFP-double positive cells was significantly smaller than that of

TBR1-negative/GFP-positive cells (L5, L6). Statistical significance was assessed using t-test.

a.u., arbitrary unit; +, positive; -, negative.

Supplementary Information

Supplementary figure S1. GFP expressions in brains of two *Scn1a*-GFP transgenic

mouse lines.

Chromogenic immunostaining of GFP in parasagittal sections from brains of 5W *Scn1a*-GFP

mouse lines (#184, #233) and wild-type controls by anti-GFP antibody (brown). Sections

were counterstained with Nissl for labeling of neurons (violet). IHC, immunohistochemistry.

Scale bars; 1 mm.

Supplementary figure S2. Similar GFP distribution in brains of two *Scn1a*-GFP mouse

lines (#184, #233).

Fluorescence images of parasagittal sections from brains of P15-17 *Scn1a*-GFP mouse lines

#184 (**A-D**) and #233 (**E-H**). Magnified images outlined in (A, E) are shown in (B-D, F-H)

and further magnified in (B1, C1~3, D1, F1, G1~3, H1). Although both lines #184 and #233

show a similar distribution of GFP-signals across all brain regions, the GFP signals are more

intense in #233 than that in #184 in caudal part of the brain such as mid brain and brainstem.

CA, cornu ammonis; DG, dentate gyrus; o, stratum oriens; p, stratum pyramidale; r, stratum

radiatum; l, stratum lucidum; g, stratum granulosum; h, hilus; DCN, deep cerebellar nuclei; M,

molecular layer; P, Purkinje cell layer; G, granular cell layer. Scale bars; 1 mm (A, E), 500

µm (B-D, F-H), 50 µm (B1, C1-3, D1, F1, G1-3, H1).

Supplementary figure S3. Expression levels of Nav1.1 are stable but that of GFP are

rather variable among *Scn1a*-GFP individual mice.

(**A**) Western blot analysis for Nav1.1 in *Scn1a*-GFP and wild-type mice. The whole

membrane fractions from 5W *Scn1a*-GFP brains (line #233) were probed with anti-Nav1.1

antibody (left panel). Nav1.1 values quantified from the blots were normalized by

beta-tubulin (TUBB), and the mean of Nav1.1 amount in wild-type was assigned as a value of

100 (right panel). The amount of Nav1.1 was not significantly changed in the *Scn1a*-GFP

mice compared to wild-type mice (t-test; $p > 0.05$). The boxes show median, 25th and 75th

percentiles, and whiskers represent minimum and maximum values. Cross marks indicate

mean values for each genotype. (**B**) Western blot analysis for GFP in *Scn1a*-GFP and

wild-type mice. The whole cytosolic fractions from the same series of mice used in (A) were

probed with anti-GFP antibody (left panel). GFP values were normalized by glyceraldehyde

3-phosphate dehydrogenase (GAPDH), and the lowest GFP amount in the #2 mouse was

assigned as a value of 1 (right panel). Statistical significance was assessed using t-test.

1004

Supplementary figure S4. Distribution of GFP fluorescent signals in neocortical layers of

Scn1a-**GFP mouse brain.**

(**A**) Fluorescent images of parasagittal section spanning whole neocortical area of P15

Scn1a-GFP brain (line #233). (**B-D**) Outlined areas in (A) are magnified in (B-D) and further

in (B1-B3, C1-C2, D1-D3). GFP signals in L2/3 of primary somatosensory cortex (C) are

brighter than other regions such as primary motor cortex (B) and secondary visual cortex (D).

Brain regions were defined using mouse brain atlas (Paxinos and Franklin, 2001) as reference.

Dashed lines indicate the borders between M1, S1 and V2 areas. M1, primary motor cortex;

S1, primary somatosensory cortex; V2, secondary visual cortex. Scale bars; 1 mm (A), 100

μm (B-D), 50 μm (B1-B3, C1-C3, D1-D3).

1015

Supplementary figure S5. Immunostaining of NeuN and GFP in the neocortex of

Scn1a-**GFP mouse.**

(**A**) Double immunostaining of parasagittal sections from P15 *Scn1a*-GFP mouse brain (line

#233) by mouse anti-GFP (green) and mouse anti-NeuN (magenta) antibodies with DAPI

(blue). Merged images are shown in the right panels. Arrows indicate GFP/NeuN-double

positive cells. Arrowheads indicate GFP-positive but NeuN-negative cells. Note that cells

with intense GFP-signals, which are assumed to be inhibitory neurons (see Figure 8), were

NeuN-negative. All images are oriented from pial surface (top) to callosal (bottom). Scale

bars; 50 μm. (**B**) Bar graphs indicating the percentage of cells with GFP-positive somata

among cells with GFP- and/or NeuN-positive somata in neocortical layers. Cells in primary

motor cortex (M1) and primary somatosensory cortex (S1) of P15 *Scn1a*-GFP mice were

counted (see also Supplementary table S1). L2/3, L5, L6: neocortical layer II/III, V, VI. +,

positive; -, negative.

1029

Supplementary figure S6. Immunostaining of GFP and ankyrinG in the neocortex of

Scn1a-**GFP mouse.**

(**A**) Double immunostaining of parasagittal sections from P15 *Scn1a*-GFP mouse brain (line

#233) by mouse anti-GFP (green) and goat anti-ankyrinG (cyan) antibodies. Merged images

are shown in the right panels. Arrows indicate ankyrinG-positive AISs with GFP-positive

soma. All images are oriented from pial surface (top) to callosal (bottom). Scale bars; 50μm.

(B) Bar graphs indicating the percentage of cells with GFP-positive/negative somata per cells

with ankyrinG-positive AISs in L2/3, L5, and L6 (see also Supplementary table S5). Note that

30% (L2/3), 26% (L5) and 9% (L6) of cells with ankyrinG-positive AISs have GFP-positive

somata. L2/3, L5, L6: neocortical layer II/III, V, VI. AnkG, ankyrinG; +, positive; -, negative.

1040

Supplementary figure S7. GFP-positive cells are more abundant in layers II/III and V

than in layer VI of the neocortex.

Bar graphs indicating the average percentage of cells with GFP-positive/negative somata per

cells with ankyrinG-positive AISs in three different assessments (see also Supplementary

table S6). Note that 23% (L2/3), 26% (L5) and 13% (L6) of cells with ankyrinG positive AISs

have GFP-positive somata. L2/3, L5, L6: neocortical layer II/III, V, VI. AnkG, ankyrinG; +,

positive; -, negative.

1048

Supplementary figure S8. A part of FEZF2-positive cells at L5 have Nav1.1-positive AIS

of *Scn1a*-GFP mouse neocortex.

(**A**) Triple immunostaining of parasagittal section from P15 *Scn1a*-GFP mouse brain (line

#233) by rabbit anti-FEZF2 (magenta), goat anti-Nav1.1 (green), and mouse anti-ankyrinG

(cyan) antibodies. Merged images are shown in the right panels. An arrow indicates

Nav1.1-positive AIS of FEZF2-positive cell. All images are oriented from pial surface (top)

to callosal (bottom). Scale bars; 50μm. (**B**) Bar graphs indicating the percentage of cells with

FEZF2- and Nav1.1-positive/negative nuclei or AISs per cells with ankyrinG-positive AISs

(left panel) (see also Supplementary table S25), the percentage of cells with

FEZF2-positive/negative nuclei per cells with ankyrinG and Nav1.1-positive AISs (middle

panel) (see also Supplementary table S26), and the percentage of cells with

Nav1.1-positive/negative AISs per cells with ankyrinG-positive AISs and FEZF2-positive

nuclei (right panel) (see also Supplementary table S27). Cells in primary motor cortex of

Scn1a-GFP mouse at P15 were counted. L5, L6: neocortical layer V, VI. AnkG, ankyrinG; +,

positive; -, negative.

1064

Supplementary figure S9. A half of FEZF2-positive cells at L5 have Nav1.2-positive AIS

of *Scn1a*-GFP mouse neocortex.

(**A**) Triple immunostaining of parasagittal section from P15 *Scn1a*-GFP mouse brain (line

#233) by rabbit anti-FEZF2 (green), goat anti-Nav1.2 (magenta), and mouse anti-ankyrinG

(cyan) antibodies. Merged images are shown in the right panels. An arrow indicates

FEZF2/Nav1.2-double positive cell. An arrowhead indicates Nav1.2-negative AIS of

FEZF2-positive cell. All images are oriented from pial surface (top) to callosal (bottom).

Scale bars; 50 μm. (**B**) Bar graphs indicating the percentage of cells with FEZF2 and

Nav1.2-positive/negative nuclei or AISs per cells with ankyrinG-positive AISs (left-upper

panel) (see also Supplementary table S28), the percentage of cells with

FEZF2-positive/negative nuclei per cells with ankyrinG and Nav1.2-positive AISs

(middle-upper panel) (see also Supplementary table S29), and the percentage of cells with

Nav1.2-positive/negative AISs per cells with ankyrinG-positive AISs and FEZF2-positive

nuclei (right-upper panel) (see also Supplementary table S30). Cells with

ankyrinG-positive/negative AISs per cells with FEZF2-positive nuclei (lower panel) (see also

Supplementary table S31). Cells in primary motor cortex of *Scn1a*-GFP mouse at P15 were

counted. L5, L6: neocortical layer V, VI. +; positive. -; negative.

1082

Supplementary figure S10. Most of the FEZF2/GFP-double positive cells are

Nav1.2-negative at L5 in *Scn1a*-GFP mouse neocortex.

(**A**) Triple immunostaining of parasagittal sections from P15 *Scn1a*-GFP mouse brain (line

#233) by rabbit anti-FEZF2 (cyan), mouse anti-GFP (green), and goat anti-Nav1.2 (magenta)

antibodies. Merged images are shown in the right panels. Arrows indicate FEZF2/GFP-double

positive cells (cells with GFP-positive somata and FEZF2-positive nuclei). Note that

Nav1.2-positive AISs do not belong to GFP/FEZF2-double positive cells. All images are

oriented from pial surface (top) to callosal (bottom). Scale bars; 25 μm. (**B**) Bar graphs

indicating the percentage of cells with GFP- and Nav1.2-positive/negative somata and AISs

per cells with FEZF2-positive nuclei in L5 and L6 (see also Supplementary table S32). For

this graph, to obtain correct cell population for Nav1.2-negative cells, virtual cell numbers

were estimated using percentage of ankyrinG/FEZF2-double positive cells in Supplementary

figure S9B-lower panel (see also Supplementary table S28). Cells in primary motor cortex of

Scn1a-GFP mouse at P15 were counted. L5, L6: neocortical layer V, VI. +, positive; -,

negative.

1098

Supplementary figure S11. AISs of TBR1-positive cells are Nav1.1-negative in

***Scn1a*-GFP mouse neocortex.**

(**A**) Triple immunostaining of parasagittal section from P15 *Scn1a*-GFP mouse brain (line

#233) by rabbit anti-TBR1 (magenta), goat anti-Nav1.1 (green), and mouse anti-ankyrinG

(cyan) antibodies. The regions of motor cortex were shown. Merged images are shown in the

right panels. Arrowheads indicate Nav1.1-positive cells. All images are oriented from pial

surface (top) to callosal (bottom). Scale bars; 50 μm. (**B**) Bar graphs indicating the percentage

of cells with TBR1- and Nav1.1-positive/negative nuclei or AISs per cells with

ankyrinG-positive AISs (left panel) (see also Supplementary table S37), the percentage of

cells with TBR1-positive/negative nuclei per cells with ankyrinG- and Nav1.1-positive AISs

(middle panel) (see also Supplementary table S38), and the percentage of cells with

Nav1.1-positive/negative AISs per cells with ankyrinG-positive AISs and TBR1-positive

nuclei (right panel) (see also Supplementary table S39). Cells in primary motor cortex of

Scn1a-GFP mouse at P15 were counted. L2/3, L5, L6: neocortical layer II/III, V, VI. AnkG,

ankyrinG; +, positive; -, negative.

1114

Supplementary figure S12. 70% of TBR1-positive cells have Nav1.2-positive AIS in

***Scn1a*-GFP mouse neocortex.**

(**A**) Triple immunostaining of parasagittal section from P15 *Scn1a*-GFP mouse brain (line

#233) by rabbit anti-TBR1 (green), goat anti-Nav1.2 (magenta), and mouse anti-ankyrinG

(cyan) antibodies. Merged images are shown in the right panels. Arrows indicate

TBR1/ankyrinG-double positive cells. Arrowheads indicate Nav1.2-positive AIS of

TBR1-positive cells. All images are oriented from pial surface (top) to callosal (bottom).

Scale bars; 25 μm. (**B**) Bar graphs indicating the percentage of cells with TBR1- and

Nav1.2-positive/negative nuclei or AISs per cells with ankyrinG-positive AISs (left-upper

panel) (see also Supplementary table S40), the percentage of cells with

TBR1-positive/negative nuclei per cells with ankyrinG- and Nav1.2-positive AISs

(middle-upper panel) (see also Supplementary table S41), and the percentage of cells with

Nav1.2-positive/negative AISs per cells with ankyrinG-positive AISs and TBR1-positive

nuclei (right-upper panel) (see also Supplementary table S42). Cells with

ankyrinG-positive/negative AISs per cells with TBR1-positive nuclei (lower panel) (see also

Supplementary table S43). Cells in primary motor cortex of *Scn1a*-GFP mouse at P15 were

counted. L2/3, L5, L6: neocortical layer II/III, V, VI. AnkG, ankyrinG; +, positive; -,

negative.

1133

Supplementary figure S13. TBR1/Nav1.2-double positive cells are GFP-negative at L5

and L6 in *Scn1a*-GFP mouse neocortex.

(**A**) Triple immunostaining of parasagittal section from P15 *Scn1a*-GFP mouse brain (line

#233) by rabbit anti-TBR1 (cyan), mouse anti-GFP (green), and goat anti-Nav1.2 (magenta)

antibodies. Merged images are shown in the right panels. Arrows indicate TBR1/GFP-double

positive cells. Arrowheads indicate Nav1.2-positive AIS of TBR1-positive cells. All images

are oriented from pial surface (top) to callosal (bottom). Scale bar; 50 µm. (**B**) Bar graphs

indicating the percentage of cells with GFP- and Nav1.2-positive/negative somata and AISs

per cells with TBR1-positive nuclei in L2/3, L5, and L6 (see also Supplementary table S44).

For this graph, to obtain correct cell population for Nav1.2-negative cells, virtual cell numbers

were estimated using percentage of ankyrinG/TBR1-double positive cells in Supplementary

figure S12B-lower panel (see also Supplementary table S40). Cells in primary motor cortex of

Scn1a-GFP mouse at P15 were counted. L2/3, L5, L6: neocortical layer II/III, V, VI. +,

positive; -, negative.

1148

1149   **Supplementary figure S14. Distributions of Nav1.1 (GFP) and Nav1.2 in neocortical**

1150   **layer V revealed by the analysis of *Scn1a*-GFP mouse.**

1151

1152

	Region [a]	per cells with GFP+ and/or NeuN+ somata		
		% GFP+, NeuN- [b]	% GFP+, NeuN+ [b]	% GFP-, NeuN+ [b]
M1	L2/3	6.8 ± 0.5	23.3 ± 0.5	70.2 ± 0.5
	L5	8.4 ± 0.6	23.8 ± 0.1	67.7 ± 1.3
	L6	6.5 ± 0.6	15.2 ± 2.2	78.3 ± 2.3
S1	L2/3	6.7 ± 0.1	21.3 ± 2.0	72.0 ± 2.1
	L5	10.2 ± 0.9	22.6 ± 2.1	67.1 ± 1.3
	L6	5.9 ± 0.7	11.7 ± 1.8	82.4 ± 1.8

1153

Supplementary table S1. Percentage of cells with GFP- and/or NeuN-positive somata

among cells with GFP- and/or NeuN-positive somata in *Scn1a*-GFP mouse neocortex.

Values for Supplementary figure S5B. [a]M1, S1, L2/3, L5, L6: primary motor cortex, primary

somatosensory cortex, neocortical layer II/III, V, VI. Cells in three *Scn1a*-GFP mice (line

#233) at P15 were counted; [b]L2/3 (M1; N = 2599 cells, S1; N=2932), L5 (M1; N = 1734 cells,

S1; N=1827) and L6 (M1; N = 2465 cells, S1; N=2335). Values are presented as mean ± SEM.

+, positive; -, negative.

1161

1162

Region [a]	per cells with AnkG+ AISs			
	% GFP+ [b]		% GFP- [b]	
L2/3	16.6 ± 1.9		83.4 ± 1.9	
L5	21.4 ± 3.3		78.6 ± 3.3	
L6	7.9 ± 1.7		92.1 ± 1.7	
	% Nav1.1+	% Nav1.1-	% Nav1.1+	% Nav1.1-
L2/3	5.5 ± 1.1	11.1 ± 0.8	0.2 ± 0.1	83.4 ± 2.0
L5	8.0 ± 0.2	13.4 ± 3.3	0.2 ± 0.1	78.7 ± 3.1
L6	3.7 ± 0.5	4.2 ± 1.3	0.1 ± 0.1	92.0 ± 1.6

1163

Supplementary table S2. Percentage of cells with GFP- and Nav1.1-positive/negative

somata and AISs per cells with ankyrinG-positive AISs in *Scn1a*-GFP mouse neocortex.

Values for Figure 4D-left panel. [a]L2/3, L5, L6: neocortical layer II/III, V, VI. Cells in three

Scn1a-GFP mice (line #233) at P15 were counted; [b]L2/3 (N = 1412 cells), L5 (N = 1054 cells),

and L6 (N = 1481 cells). Values are presented as mean ± SEM. AnkG, ankyrinG; +, positive;

-, negative.

1170

1171

| Region [a] | per cells with AnkG+ AISs and Nav1.1+ somata and/or AISs | |
	% GFP+ [b]	% GFP- [b]
L2/3	98.5 ± 0.8	1.55 ± 0.8
L5	98.7 ± 0.6	1.27 ± 0.6
L6	97.2 ± 2.8	2.82 ± 2.8

1172

Supplementary table S3. Percentage of cells with GFP-positive/negative somata per cells

with ankyrinG-positive AISs and Nav1.1-positive somata and/or AISs in *Scn1a*-GFP

mouse neocortex. Values for Figure 4D-middle panel. [a]L2/3, L5, L6: neocortical layer II/III,

V, VI. Cells in three *Scn1a*-GFP mice (line #233) at P15 were counted; [b]L2/3 (N = 108 cells),

L5 (N = 124 cells), and L6 (N = 82 cells). Values are presented as mean ± SEM. AnkG,

ankyrinG; +, positive; -, negative.

1179

1180

| Region [a] | per cells with AnkG+ AISs and GFP+ somata | |
	% Nav1.1+ [b]	% Nav1.1- [b]
L2/3	32.8 ± 3.5	67.2 ± 3.5
L5	39.9 ± 6.4	60.1 ± 6.4
L6	49.5 ± 8.4	50.5 ± 8.4

1181

Supplementary table S4. Percentage of cells with GFP-positive/negative somata per cells

with ankyrinG-positive AISs and Nav1.1+ somata and/or AISs in *Scn1a*-GFP mouse

neocortex. Values for Figure 4D-right panel. [a]L2/3, L5, L6: neocortical layer II/III, V, VI.

Cells in three *Scn1a*-GFP mice (line #233) at P15 were counted; [b]L2/3 (N = 229 cells), L5 (N

= 220 cells), and L6 (N = 111 cells). Values are presented as mean ± SEM. AnkG, ankyrinG;

+, positive; -, negative.

1188

1189

1190

| Region [a] | per cells with AnkG+ AISs | |
	% GFP+ [b]	% GFP- [b]
L2/3	30.0 ± 1.1	70.0 ± 1.1
L5	26.4 ± 1.6	73.6 ± 1.6
L6	9.0 ± 1.0	91.0 ± 1.0

1191

1192 **Supplementary table S5. Percentage of cells with GFP-positive/negative somata per cells**

1193 **with ankyrinG-positive AISs in *Scn1a*-GFP mouse neocortex.** Values for Supplementary

1194 figure S6B. [a]L2/3, L5, L6: neocortical layer II/III, V, VI. Cells in three *Scn1a*-GFP mice (line

1195 #233) at P15 were counted; [b]L2/3 (N = 1787 cells), L5 (N = 1450 cells), and L6 (N = 2330

1196 cells). Values are presented as mean ± SEM. AnkG, ankyrinG; +, positive; -, negative.

1197

1198

| Region [a] | % GFP+/AnkG+ AIS [b] | Nav1.1/GFP/AnkG | Nav1.2/GFP/AnkG | Average | SEM |
		% GFP+/AnkG+ AIS [b]	% GFP+/AnkG+ AIS [b]		
L2/3	30.0	16.6	22.6	23.1	3.9
L5	26.4	21.4	30.2	26.0	2.5
L6	9.0	7.9	20.7	12.5	4.1

1199

1200 **Supplementary table S6. Percentage of cells with GFP-positive somata per cells with**

1201 **ankyrinG-positive AISs in the three different assessments.** Values for Supplementary

1202 figure S7. [a]L2/3, L5, L6: neocortical layer II/III, V, VI. [b]Values indicate that percentage of

1203 GFP positive cells in Supplementary figure S6B, Figure 4D-left panel and Figure 11B-left

1204 panel, respectively. AnkG, ankyrinG; +, positive; -, negative.

1205

1206

1207

| Region [a] | per cells with GFP+ somata and AnkG+ AISs | |
	% Nav1.1+ [b]	% Nav1.1- [b]
CA1	98.3 ± 0.9	1.7 ± 0.9

1208

1209 **Supplementary table S7. Percentage of cells with Nav1.1-positive/negative AISs per cells**

1210 **with GFP-positive somata and ankyrinG-positive AISs in *Scn1a*-GFP mouse**

1211 **hippocampus.** Values for Figure 5F-left panel. [a]CA1: Cornu ammonis 1. [b]Cells in three

1212 *Scn1a*-GFP mice (line #233) at P15 were counted (N = 117 cells). Values are presented as

1213 mean ± SEM. AnkG, ankyrinG; +, positive; -, negative.

1214

1215

| Region [a] | per cells with Nav1.1+ and AnkG+ AISs | |
	% GFP+ [b]	% GFP- [b]
CA1	100.0 ± 0.0	0.0 ± 0.0

1216

1217 **Supplementary table S8. Percentage of cells with GFP-positive/negative somata per cells**

1218 **with Nav1.1/ankyrinG-double positive AISs in *Scn1a*-GFP mouse hippocampus.** Values

1219 for Figure 5F-right panel. [a]CA1: Cornu ammonis 1. [b]Cells in three *Scn1a*-GFP mice (line

1220 #233) at P15 were counted (N = 114 cells). Values are presented as mean ± SEM. AnkG,

1221 ankyrinG; +, positive; -, negative.

1222

	per cells with GFP+ somata	
Region [a]	%Tomato+ [b]	% Tomato- [b]
L2/3	23.2 ± 2.4	76.8 ± 2.4
L5	28.1 ± 0.8	71.9 ± 0.8
L6	27.4 ± 3.3	72.6 ± 3.3
CA1	97.9 ± 0.1	2.1 ± 0.1
DG	93.5 ± 4.5	6.5 ± 4.5

1223

Supplementary table S9. Percentage of cells with Tomato-positive/negative somata per

cells with GFP-positive somata in *Scn1a*-GFP/*Vgat*-Cre/*Rosa26*-tdTomato mouse

neocortex and hippocampus. Values for Figure 7C-left panels. [a]L2/3, L5, L6, CA1, DG:

neocortical layer II/III, V, VI, cornu ammonis 1, dentate gyrus. Cells in two

Scn1a-GFP/*Vgat*-Cre/*Rosa26*-tdTomato mice (line #233) at 4W were counted; [b]L2/3 (N =

882 cells), L5 (N = 693 cells), L6 (N = 590 cells), CA1 (N = 270 cells) and DG (N = 191

cells). Values are presented as mean ± SEM. +, positive; -, negative.

1231

1232

	per cells with Tomato+ somata	
Region [a]	%GFP+ [b]	% GFP- [b]
L2/3	72.7 ± 5.0	27.3 ± 5.0
L5	76.9 ± 0.0	23.1 ± 0.0
L6	83.0 ± 5.5	17.0 ± 5.5
CA1	93.2 ± 0.2	6.8 ± 0.2
DG	77.2 ± 5.2	22.8 ± 5.2

1233

Supplementary table S10. Percentage of cells with GFP-positive/negative somata per

cells with Tomato-positive somata in *Scn1a*-GFP/*Vgat*-Cre/*Rosa26*-tdTomato mouse

neocortex and hippocampus. Values for Figure 7C-right panels. [a]L2/3, L5, L6, CA1, DG:

neocortical layer II/III, V, VI, cornu ammonis 1, dentate gyrus. Cells in two

Scn1a-GFP/*Vgat*-Cre/*Rosa26*-tdTomato mice (line #233) at 4W were counted; [b]L2/3 (N =

263 cells), L5 (N = 224 cells), L6 (N = 300 cells), CA1 (N = 276 cells) and DG (N = 364

cells). Values are presented as mean ± SEM. +, positive; -, negative.

1241

Region [a]	per cells with GFP+ somata			
	% PV+, SST+ [b]	% PV+ [b]	% SST+ [b]	% PV-, SST- [b]
L2/3	0.1 ± 0.1	14.7 ± 1.4	6.4 ± 2.5	78.8 ± 3.8
L5	0.1 ± 0.1	26.3 ± 1.5	11.1 ± 2.1	62.4 ± 3.7
L6	0.4 ± 0.4	21.8 ± 2.2	14.8 ± 1.6	63.0 ± 3.7
CA1	2.0 ± 0.3	33.6 ± 3.2	22.7 ± 3.5	41.7 ± 7.4
CA2/3	1.4 ± 0.3	24.6 ± 5.1	16.0 ± 1.5	58.1 ± 6.5
DG	0.0 ± 0.0	19.3 ± 4.5	21.4 ± 1.4	59.3 ± 3.4

1242

Supplementary table S11. Percentage of cells with PV-positive/negative somata per cells
with GFP-positive somata in *Scn1a*-GFP mouse neocortex and hippocampus. Values for
Figure 8B-left panels. [a]L2/3, L5, L6, CA1, CA2/3, DG: neocortical layer II/III, V, VI, cornu
ammonis 1, 2 plus 3, dentate gyrus. Cells in three *Scn1a*-GFP mice (line #233) at 4W were
counted; [b]L2/3 (N = 631 cells), L5 (N = 533 cells), L6 (N = 390 cells), CA1 (N = 231 cells),
CA2/3 (N = 311 cells) and DG (N = 258 cells). Values are presented as mean ± SEM. +,
positive; -, negative.

1250

Region [a]	per cells with PV+ somata	
	%GFP+ [b]	% GFP- [b]
L2/3	100.0 ± 0.0	0.0 ± 0.0
L5	98.4 ± 1.5	1.6 ± 1.5
L6	100.0 ± 0.0	0.0 ± 0.0
CA1	100.0 ± 0.0	0.0 ± 0.0
CA2/3	100.0 ± 0.0	0.0 ± 0.0
DG	100.0 ± 0.0	0.0 ± 0.0

1251

Supplementary table S12. Percentage of cells with GFP-positive/negative somata per
cells with PV-positive somata. Values for Figure 8B-middle panels. [a]L2/3, L5, L6, CA1,
CA2/3, DG: neocortical layer II/III, V, VI, cornu ammonis 1, 2 plus 3, dentate gyrus. Cells in
three *Scn1a*-GFP mice (line #233) at 4W were counted; [b]L2/3 (N = 95 cells), L5 (N = 144
cells), L6 (N = 84 cells), CA1 (N = 83 cells), CA2/3 (N = 78 cells) and DG (N = 37 cells).
Values are presented as mean ± SEM. +, positive; -, negative.

1258

Region [a]	per cells with SST+ somata	
	%GFP+ [b]	% GFP- [b]
L2/3	89.8 ± 7.6	10.2 ± 7.6
L5	83.3 ± 4.1	16.7 ± 4.1
L6	95.4 ± 2.8	4.6 ± 2.8
CA1	100.0 ± 0.0	0.0 ± 0.0
CA2/3	96.4 ± 1.9	3.4 ± 1.9
DG	99.3 ± 0.7	0.7 ± 0.7

1260

Supplementary table S13. Percentage of cells with GFP-positive/negative somata per

cells with SST-positive somata. Values for Figure 8B-right panels. [a]L2/3, L5, L6, CA1,

CA2/3, DG: neocortical layer II/III, V, VI, cornu ammonis 1, 2 plus 3, dentate gyrus. Cells in

three *Scn1a*-GFP mice (line #233) at 4W were counted; [b]L2/3 (N = 46 cells), L5 (N = 76

cells), L6 (N = 58 cells), CA1 (N = 58 cells), CA2/3 (N = 56 cells) and DG (N = 55 cells).

Values are presented as mean ± SEM. +, positive; -, negative.

1267

Region [a]	GFP+, PV-, SST-		GFP+, PV+		GFP+, SST+	
	Area size [b]	Intensity [b]	Area size [b]	Intensity [b]	Area size [b]	Intensity [b]
L2/3	1037.1 ± 19.1	79.0 ± 1.2	1090.0 ± 51.8	112.0 ± 4.9	986.5 ± 55.7	76.4 ± 4.8
L5	1401.3 ± 63.6	62.9 ± 0.8	1051.4 ± 36.3	77.9 ± 2.0	969.1 ± 51.5	58.5 ± 1.4
L6	1003.2 ± 36.6	52.0 ± 1.1	1333.4 ± 61.0	65.7 ± 2.1	1130.2 ± 57.7	57.0 ± 2.9
CA1	1290.6 ± 72.6	76.5 ± 3.9	1544.1 ± 106.6	84.1 ± 5.6	1508.0 ± 99.5	74.9 ± 3.0
CA2/3	988.0 ± 49.5	74.9 ± 2.5	1494.8 ± 134.4	83.5 ± 3.5	1146.9 ± 86.9	61.2 ± 2.6
DG	1242.1 ± 61.5	73.6 ± 2.4	1721.0 ± 159.1	82.7 ± 5.5	1382.1 ± 123.2	51.5 ± 1.4

1268

Supplementary table S14. Area size and intensity of GFP immunosignals in

GFP-positive cells with PV/SST-positive or negative somata in *Scn1a*-GFP mouse

neocortex and hippocampus. Values for Figure 9A. [a]L2/3, L5, L6, CA1, CA2/3, DG:

neocortical layer II/III, V, VI, cornu ammonis 1, 2 plus 3, dentate gyrus. Cells in two

Scn1a-GFP mice (line #233) at 4W were counted; [b]L2/3 (N = 157 cells), L5 (N = 122 cells)

and L6 (N = 109 cells), CA1 (N = 52 cells), CA2/3 (N = 85 cells) and DG (N = 60 cells).

Values are presented as mean ± SEM. +, positive; -, negative.

1276

1277

| Region [a] | Cell type [b] | p-value for intensity [c] | |
		PV-, SST-, GFP+	PV+, GFP+
L2/3	PV+, GFP+	1.4×10^{-6}	NA
	SST+, GFP+	8.4×10^{-1}	1.4×10^{-6}
L5	PV+, GFP+	1.5×10^{-6}	NA
	SST+, GFP+	1.3×10^{-1}	1.5×10^{-6}
L6	PV+, GFP+	1.7×10^{-6}	NA
	SST+, GFP+,	1.4×10^{-1}	9.6×10^{-3}
CA1	PV+ GFP+	4.1×10^{-1}	NA
	SST+, GFP+	9.6×10^{-1}	3.6×10^{-1}
CA2/3	PV+, GFP+	8.4×10^{-2}	NA
	SST+, GFP+	3.7×10^{-2}	9.0×10^{-4}
DG	PV+, GFP+	1.8×10^{-1}	NA
	SST+, GFP+	5.9×10^{-6}	6.2×10^{-1}

1278

Supplementary table S15. Statistical significance for intensity of GFP immunosignals in

GFP-positive cells with PV/SST-positive or negative somata. Values for Figure 9B-left

panels. [a]L2/3, L5, L6, CA1, CA2/3, DG: neocortical layer II/III, V, VI, cornu ammonis 1, 2

plus 3, dentate gyrus. Cells in two *Scn1a*-GFP mice (line #233) at 4W were counted; [b]L2/3 (N

= 157 cells), L5 (N = 122 cells) and L6 (N = 109 cells), CA1 (N = 52 cells), CA2/3 (N = 85

cells) and DG (N = 60 cells). [c]Statistical significance was assessed using one-way ANOVA

followed by Tukey–Kramer post-hoc multiple comparison test. Values are presented as mean

± SEM. +, positive; -, negative; NA, not available.

1287

Region [a]	Cell type [b]	p-value for area size [c]	
		PV-, SST-, GFP+	**PV+, GFP+**
L2/3	PV+, GFP+	4.9×10^{-1}	NA
	SST+, GFP+	7.2×10^{-1}	3.6×10^{-1}
L5	PV+, GFP+	1.2×10^{-4}	NA
	SST+, GFP+	1.5×10^{-4}	7.3×10^{-1}
L6	PV+, GFP+	8.0×10^{-6}	NA
	SST+, GFP+	1.9×10^{-1}	3.8×10^{-2}
CA1	PV+, GFP+	1.2×10^{-1}	NA
	SST+, GFP+	2.0×10^{-1}	9.6×10^{-1}
CA2/3	PV+, GFP+	8.2×10^{-5}	NA
	SST+, GFP+	5.6×10^{-1}	9.8×10^{-2}
DG	PV+, GFP+	9.7×10^{-3}	NA
	SST+, GFP+	5.1×10^{-1}	1.5×10^{-1}

1288

Supplementary table S16. Statistical significance for area size of GFP immunosignals in

GFP-positive cells with PV/SST-positive or negative somata. Values for Figure 9B-right

panels. [a]L2/3, L5, L6, CA1, CA2/3, DG: neocortical layer II/III, V, VI, cornu ammonis 1, 2

plus 3, dentate gyrus. Cells in two *Scn1a*-GFP mice (line #233) at 4W were counted; [b]L2/3 (N

= 157 cells), L5 (N = 122 cells) and L6 (N = 109 cells), CA1 (N = 52 cells), CA2/3 (N = 85

cells) and DG (N = 60 cells). [c]Statistical significance was assessed using one-way ANOVA

followed by Tukey–Kramer post-hoc multiple comparison test. Values are presented as mean

± SEM. +, positive; -, negative; NA, not available.

1297

Region [a]	per cells with AnkG+ AISs			
	% Nav1.1+ [b]		% Nav1.1- [b]	
L2/3	4.7 ± 0.1		95.3 ± 0.1	
L5	6.0 ± 1.5		94.0 ± 1.5	
L6	2.5 ± 0.3		97.5 ± 0.3	
	% Nav1.2+	% Nav1.2-	% Nav1.2+	% Nav1.2-
L2/3	0.3 ± 0.2	4.4 ± 0.1	78.1 ± 3.6	17.2 ± 3.7
L5	0.1 ± 0.1	5.9 ± 1.4	69.3 ± 4.5	24.7 ± 5.7
L6	0.1 ± 0.1	2.3 ± 0.1	69.3 ± 3.4	28.3 ± 3.2

1298

Supplementary table S17. Percentage of cells with Nav1.1- and Nav1.2-positive/negative cells per cells with ankyrinG-positive AISs in *Scn1a*-GFP mouse neocortex. Values for Figure 10B. [a]L2/3, L5, L6: neocortical layer II/III, V, VI. Cells in three *Scn1a*-GFP mice (line #233) at P15 were counted; [b]L2/3 (N = 1300 cells), L5 (N = 895 cells), and L6 (N = 1295 cells). Values are presented as mean ± SEM. AnkG, ankyrinG; +, positive; -, negative.

Region [a]	per cells with AnkG+ AISs			
	% GFP+ [b]		% GFP- [b]	
L2/3	22.6 ± 2.9		77.4 ± 2.9	
L5	30.2 ± 0.3		69.8 ± 0.3	
L6	20.7 ± 1.4		79.3 ± 1.4	
	% Nav1.2+	% Nav1.2-	% Nav1.2+	% Nav1.2-
L2/3	7.0 ± 1.4	15.6 ± 1.6	53.3 ± 2.5	24.1 ± 0.9
L5	5.2 ± 1.2	25.0 ± 1.5	47.0 ± 2.2	22.8 ± 1.9
L6	3.2 ± 0.8	17.6 ± 0.9	59.3 ± 2.4	20.0 ± 3.7

1306

Supplementary table S18. Percentage of cells with GFP- and Nav1.2-positive/negative somata and AISs per cells with ankyrinG-positive AISs in *Scn1a*-GFP mouse neocortex. Values for Figure 11B-left panel. [a]L2/3, L5, L6: neocortical layer II/III, V, VI. Cells in three *Scn1a*-GFP mice (line #233) at P15 were counted; [b]L2/3 (N = 877 cells), L5 (N = 724 cells) and L6 (N = 682 cells). Values are presented as mean ± SEM. AnkG, ankyrinG; +, positive; -, negative.

1313

1314

| Region [a] | per cells with AnkG+ and Nav1.2+ AISs | |
	% GFP+ [b]	% GFP- [b]
L2/3	11.8 ± 2.7	88.2 ± 2.7
L5	10.0 ± 1.6	90.0 ± 1.6
L6	5.0 ± 1.0	95.0 ± 1.0

1315

Supplementary table S19. Percentage of cells with GFP-positive/negative somata per

cells with Nav1.2/ankyrinG-double positive AISs in *Scn1a*-GFP mouse neocortex. Values

for Figure 11B-middle panel. [a]L2/3, L5, L6: neocortical layer II/III, V, VI. Cells in three

Scn1a-GFP mice (line #233) at P15 were counted; [b]L2/3 (N = 527 cells), L5 (N = 378 cells)

and L6 (N = 422 cells). Values are presented as mean ± SEM. AnkG, ankyrinG; +, positive; -,

negative.

1322

1323

| Region [a] | per cells with AnkG+ AISs and GFP+ somata | |
	% Nav1.2+ [b]	% Nav1.2- [b]
L2/3	31.5 ± 2.3	68.5 ± 2.3
L5	17.1 ± 4.4	82.9 ± 4.4
L6	14.4 ± 2.6	85.6 ± 2.6

1324

Supplementary table S20. Percentage of cells with Nav1.2-positive/negative AISs per

cells with ankyrinG-positive AISs and GFP+ somata in *Scn1a*-GFP mouse neocortex.

Values for Figure 11B-right panel. [a]L2/3, L5, L6: neocortical layer II/III, V, VI. Cells in three

Scn1a-GFP mice (line #233) at P15 were counted; [b]L2/3 (N = 197 cells), L5 (N = 215 cells)

and L6 (N = 139 cells). Values are presented as mean ± SEM. AnkG, ankyrinG; +, positive; -,

negative.

1331

1332

Age [a]	Region [b]	per cells with GFP+ somata and/or FEZF2+ nuclei		
		% GFP+, FEZF2- [c]	% GFP+, FEZF2+ [c]	% GFP-, FEZF2+ [c]
P15	L2/3	72.9 ± 3.1	13.1 ± 4.4	14.0 ± 2.2
	L5	40.4 ± 1.5	49.8 ± 2.4	9.9 ± 1.7
	L6	38.0 ± 9.0	5.0 ± 3.6	56.9 ± 8.5
4W	L2/3	88.6 ± 2.7	6.6 ± 0.8	4.7 ± 1.9
	L5	51.1 ± 0.1	47.2 ± 0.3	1.7 ± 0.3
	L6	59.3 ± 8.5	15.7 ± 1.3	25.0 ± 9.8

Supplementary table S21. Percentage of cells with FEZF2- and GFP-positive/negative nuclei and somata per cells with GFP-positive somata and/or FEZF2-positive nuclei in *Scn1a*-GFP mouse neocortex. Values for Figure 12B-left panels. [a]Two animals were counted for each age. [b]L2/3, L5, L6: neocortical layer II/III, V, VI. Cells in *Scn1a*-GFP mice (line #233) were counted; [c]L2/3 (P15, N = 670 cells; 4W, N = 325 cells), L5 (P15, N = 733 cells; 4W, N = 431 cells) and L6 (P15, N = 466 cells; 4W, N = 386 cells). Values are presented as mean ± SEM. +, positive; -, negative.

Age [a]	Region [b]	per cells with FEZF2+ nuclei	
		% GFP+ [c]	% GFP- [c]
P15	L2/3	43.2 ± 13.9	56.8 ± 13.9
	L5	83.4 ± 2.9	16.6 ± 2.9
	L6	8.0 ± 5.0	92.0 ± 5.0
4W	L2/3	48.2 ± 18.5	51.8 ± 18.5
	L5	95.7 ± 0.4	4.3 ± 0.4
	L6	36.2 ± 0.4	63.8 ± 0.4

Supplementary table S22. Percentage of cells with GFP-positive/negative somata per cells with FEZF2-positive nuclei in *Scn1a*-GFP mouse neocortex. Values for Figure 12B-middle panels. [a]Two animals were counted for each age. [b]L2/3, L5, L6: neocortical layer II/III, V, VI. Cells in *Scn1a*-GFP mice (line #233) were counted; [c]L2/3 (P15, N = 198 cells; 4W, N = 25 cells), L5 (P15, N = 431 cells; 4W, N = 214 cells) and L6 (P15, N = 283 cells; 4W, N = 202 cells). Values are presented as mean ± SEM. +, positive; -, negative.

1351

		per cells with GFP+ somata	
Age [a]	Region [b]	% FEZF2+ [c]	% FEZF2- [c]
P15	L2/3	16.1 ± 4.9	83.9 ± 4.9
	L5	54.3 ± 2.2	45.7 ± 2.2
	L6	12.0 ± 9.6	88.0 ± 9.6
4W	L2/3	5.8 ± 0.3	94.2 ± 0.3
	L5	49.0 ± 1.2	51.0 ± 1.2
	L6	17.0 ± 3.7	83.0 ± 3.7

1352

Supplementary table S23. Percentage of cells with FEZF2-positive/negative nuclei per

cells with GFP-positive somata in *Scn1a*-GFP mouse neocortex. Values for Figure

12B-right panels. [a]Two animals were counted for each age. [b]L2/3, L5, L6: neocortical layer

II/III, V, VI. Cells in *Scn1a*-GFP mice (line #233) at P15 were counted; [c]L2/3 (P15, N = 567

cells; 4W, N = 318 cells), L5 (P15, N = 479 cells; 4W, N = 363 cells) and L6 (P15, N = 202

cells; 4W, N = 228 cells). Values are presented as mean ± SEM. +, positive; -, negative.

1359

1360

	FEZF2+, GFP+		FEZF2-, GFP+		p-value	
Region [a]	Area size [b]	Intensity [b]	Area size [b]	Intensity [b]	Area size [c]	Intensity [c]
L2/3	510.8 ± 9.0	50.4 ± 0.8	377.3 ± 9.0	48.0 ± 0.8	1.0×10^{-2}	5.8×10^{-1}
L5	577.4 ± 21.4	47.6 ± 1.6	461.6 ± 18.1	25.9 ± 3.1	7.3×10^{-5}	3.8×10^{-8}
L6	389.3 ± 27.9	43.0 ± 3.2	372.1 ± 19.0	52.4 ± 2.3	6.4×10^{-1}	3.8×10^{-2}

1361

Supplementary table S24. Area size and intensity of GFP immunosignals in

GFP-positive cells with FEZF2-positive or negative nuclei in *Scn1a*-GFP mouse

neocortex. Values for Figure 13. [a]L2/3, L5, L6: neocortical layer II/III, V, VI. Cells in one

Scn1a-GFP mouse (line #233) at 4W were counted; [b]L2/3 (N = 159 cells), L5 (N = 147 cells)

and L6 (N = 80 cells). [c]Statistical significance was assessed using t-test. Values are presented

as mean ± SEM. +, positive; -, negative.

1368

1369

1370

Region [a]	per cells with AnkG+ AISs			
	% FEZF2+ [b]		% FEZF2- [b]	
L5	52.8 ± 0.7		47.2 ± 0.7	
L6	66.3 ± 5.5		33.7 ± 5.5	
	% Nav1.1+	% Nav1.1-	% Nav1.1+	% Nav1.1-
L5	2.2 ± 0.4	18.5 ± 0.8	4.9 ± 1.6	74.4 ± 1.3
L6	0.4 ± 0.3	18.7 ± 1.4	2.3 ± 0.3	78.6 ± 1.7

1371

1372 **Supplementary table S25. Percentage of cells with FEZF2- and Nav1.1-positive/negative**
1373 **nuclei or AISs per cells with ankyrinG-positive AISs in *Scn1a*-GFP mouse neocortex.**
1374 Values for Supplementary figure S8B-left panel. [a]L5, L6: neocortical layer II/III, V, VI. Cells
1375 in three *Scn1a*-GFP mice (line #233) at P15 were counted; [b]L5 (N = 588 cells) and L6 (N =
1376 580 cells). Values are presented as mean ± SEM. AnkG, ankyrinG; +, positive; -, negative.

1377

1378

Region [a]	per cells with AnkG+ and Nav1.1+ AISs	
	% FEZF2+ [b]	% FEZF2- [b]
L5	35.9 ± 9.7	64.1 ± 9.7
L6	12.4 ± 8.6	87.6 ± 8.6

1379

1380 **Supplementary table S26. Percentage of cells with FEZF2-positive/negative nuclei per**
1381 **cells with ankyrinG/Nav1.1-double positive AISs in *Scn1a*-GFP mouse neocortex.** Values
1382 for Supplementary figure S8B-middle panel. [a]L5, L6: neocortical layer V, VI. Cells in three
1383 *Scn1a*-GFP mice (line #233) at P15 were counted; [b]L5 (N = 115 cells) and L6 (N = 45 cells).
1384 Values are presented as mean ± SEM. AnkG, ankyrinG; +, positive; -, negative.

1385

1386

	per cells with AnkG+ AISs and FEZF2+ nuclei	
Region [a]	% Nav1.1+ [b]	% Nav1.1- [b]
L5	11.1 ± 1.9	88.9 ± 1.9
L6	1.7 ± 1.3	98.3 ± 1.3

1387

Supplementary table S27. Percentage of cells with Nav1.1-positive/negative AISs per

cells with ankyrinG-positive AISs and FEZF2-positive nuclei in *Scn1a*-GFP mouse

neocortex. Values for Supplementary figure S8B-right panel. [a]L5, L6: neocortical layer V, VI.

Cells in three *Scn1a*-GFP mice (line #233) at P15 were counted; [b]L5 (N = 326 cells) and L6

(N = 324 cells). Values are presented as mean ± SEM. AnkG, ankyrinG; +, positive; -,

negative.

1394

1395

	per cells with AnkG+ AISs			
Region [a]	% FEZF2+ [b]		% FEZF2- [b]	
L5	20.4 ± 5.7		79.6 ± 5.7	
L6	11.6 ± 3.4		88.4 ± 3.4	
	% Nav1.2+	% Nav1.2-	% Nav1.2+	% Nav1.2-
L5	8.9 ± 1.7	11.5 ± 1.0	50.1 ± 0.9	29.5 ± 2.6
L6	6.0 ± 1.3	5.6 ± 3.3	49.3 ± 4.8	39.1 ± 2.2

1396

1397 **Supplementary table S28. Percentage of cells with FEZF2- and Nav1.2-positive/negative**

1398 **nuclei or AISs per cells with ankyrinG-positive AISs in *Scn1a*-GFP mouse neocortex.**

1399 Values for Supplementary figure S9B-left-upper panel. [a]L5, L6: neocortical layer V, VI. Cells

1400 in three *Scn1a*-GFP mice (line #233) at P15 were counted; [b]L5 (N = 644 cells) and L6 (N =

1401 758 cells). Values are presented as mean ± SEM. AnkG, ankyrinG; +, positive; -, negative.

1402

1403

1404

| | per cells with AnkG+ and Nav1.2+ AISs | |
Region [a]	% FEZF2+ [b]	% FEZF2- [b]
L5	15.1 ± 2.9	84.9 ± 2.9
L6	10.5 ± 4.3	89.5 ± 4.3

1405

1406 **Supplementary table S29. Percentage of cells with FEZF2-positive/negative nuclei per**
1407 **cells with ankyrinG/Nav1.2-double positive AISs in *Scn1a*-GFP mouse neocortex.** Values
1408 for Supplementary figure S9B-middle-upper panel. [a]L5, L6: neocortical layer V, VI. Cells in
1409 three *Scn1a*-GFP mice (line #233) at P15 were counted; [b]L5 (N = 376 cells) and L6 (N = 415
1410 cells). Values are presented as mean ± SEM. AnkG, ankyrinG; +, positive; -, negative.
1411

| | per cells with AnkG+ AISs and FEZF2+ nuclei | |
Region [a]	% Nav1.2+ [b]	% Nav1.2- [b]
L5	51.2 ± 3.7	48.8 ± 3.7
L6	41.6 ± 9.5	58.4 ± 9.5

1412

Supplementary table S30. Percentage of cells with Nav1.2-positive/negative AISs per
cells with ankyrinG-positive AISs and FEZF2-positive nuclei in *Scn1a*-GFP mouse
neocortex. Values for Supplementary figure S9B-right-upper panel. [a]L5, L6: neocortical layer
V, VI. Cells in three *Scn1a*-GFP mice (line #233) at P15 were counted; [b]L5 (N = 135 cells)
and L6 (N = 94 cells). Values are presented as mean ± SEM. AnkG, ankyrinG; +, positive; -,
negative.
1419

1420

1421

Region [a]	per cells with FEZF2+ nuclei	
	% AnkG+ [b]	% AnkG- [b]
L5	38.6 ± 7.1	61.4 ± 7.1
L6	39.4 ± 6.3	60.6 ± 6.3

1422

1423 **Supplementary table S31. Percentage of cells with ankyrinG-positive/negative AISs per**

1424 **cells with FEZF2-positive nuclei in *Scn1a*-GFP mouse neocortex.** Values for

1425 Supplementary figure S9B-lower panel. [a]L5, L6: neocortical layer V, VI. Cells in three

1426 *Scn1a*-GFP mice (line #233) at P15 were counted; [b]L5 (N = 301 cells), and L6 (N = 257 cells).

1427 Values are presented as mean ± SEM. AnkG, ankyrinG; +, positive; -, negative.

1428

1429

Region [a]	per cells with FEZF2+ nuclei			
	% GFP+ [b]		% GFP- [b]	
L5	74.4 ± 3.4		25.6 ± 3.4	
L6	16.5 ± 4.7		83.5 ± 4.7	
	% Nav1.2+	% Nav1.2- [c]	% Nav1.2+	% Nav1.2- [c]
L5	13.9 ± 2.7	56.4 ± 6.8	13.2 ± 5.7	16.5 ± 3.2
L6	5.3 ± 3.7	9.6 ± 3.3	34.7 ± 3.0	50.3 ± 8.7

1430

Supplementary table S32. Percentage of cells with GFP- and Nav1.2-positive/negative

somata and AISs per cells with FEZF2-positive nuclei in *Scn1a*-GFP mouse neocortex.

Values for Supplementary figure S10B. [a]L5, L6: neocortical layer V, VI. Cells in three

Scn1a-GFP mice (line #233) at P15 were counted; [b]L5 (N = 267 cells) and L6 (N = 188 cells).

[c]In these values, because they were including ankyrinG-negative cells, to obtain correct cell

population for Nav1.2-negative/ankyrinG-positive cells, virtual cell number were estimated

using ratio of ankyrinG/FEZF2 double positive cells in Supplementary figure S9B.Values are

presented as mean ± SEM. +, positive; -, negative.

1439

1440

1441

| Age [a] | Region [b] | per cells with GFP+ somata and/or TBR1+ nuclei | | |
		% GFP+, TBR1- [c]	% GFP+, TBR1+ [c]	% GFP-, TBR1+ [c]
P15	L2/3	46.5 ± 0.8	38.7 ± 0.4	14.8 ± 0.3
	L5	49.3 ± 1.2	5.5 ± 2.0	45.1 ± 0.8
	L6	14.8 ± 2.3	12.9 ± 7.4	72.3 ± 9.8
4W	L2/3	35.3 ± 0.2	42.1 ± 1.5	22.6 ± 1.3
	L5	54.7 ± 0.1	2.4 ± 1.8	42.9 ± 0.1
	L6	13.3 ± 1.8	22.1 ± 4.5	64.6 ± 6.3

1442

Supplementary table S33. Percentage of cells with TBR1- and GFP-positive/negative nuclei and somata per GFP-positive cells and/or TBR1-positive nuclei in *Scn1a*-GFP mouse neocortex. Values for Figure 14B-left panels. [a]Two animals were counted for each age. [b]L2/3, L5, L6: neocortical layer II/III, V, VI. Cells in *Scn1a*-GFP mice (line #233) were counted; [c]L2/3 (P15, N = 704 cells; 4W, N = 549 cells), L5 (P15, N = 639 cells; 4W, N = 400 cells) and L6 (P15, N = 990 cells; 4W, N = 598 cells). Values are presented as mean ± SEM. +, positive; -, negative.

1450

1451

| Age [a] | Region [b] | per cells with TBR1+ nuclei | |
		% GFP+ [c]	% GFP- [c]
P15	L2/3	71.9 ± 0.4	28.1 ± 0.4
	L5	10.7 ± 3.9	89.3 ± 3.9
	L6	15.4 ± 9.1	84.6 ± 9.1
4W	L2/3	64.7 ± 1.6	35.3 ± 1.6
	L5	5.1 ± 3.9	94.9 ± 3.9
	L6	25.6 ± 5.7	74.4 ± 5.7

1452

Supplementary table S34. Percentage of cells with GFP-positive/negative somata per cells with TBR1-positive nuclei in *Scn1a*-GFP mouse neocortex. Values for Figure 14B-middle panels. [a]Two animals were counted for each age. [b]L2/3, L5, L6: neocortical layer II/III, V, VI. Cells in *Scn1a*-GFP mice (line #233) were counted; [c]L2/3 (P15, N = 367 cells;

1457   4W, N = 358 cells), L5 (P15, N = 322 cells; 4W, N = 180 cells) and L6 (P15, N = 848 cells;

1458   4W, N = 518 cells). Values are presented as mean ± SEM. +, positive; -, negative.

1459

1460

| Age [a] | Region [b] | per cells with GFP+ somata | |
		% TBR1+ [c]	% TBR1- [c]
	L2/3	45.1 ± 0.7	54.9 ± 0.7
P15	L5	10.0 ± 3.5	90.0 ± 3.5
	L6	41.5 ± 12.9	58.5 ± 12.9
	L2/3	54.4 ± 0.7	45.6 ± 0.7
4W	L5	4.1 ± 3.2	95.9 ± 3.2
	L6	61.2 ± 2.9	38.8 ± 2.9

1461

Supplementary table S35. Percentage of cells with TBR1-positive/negative nuclei per

cells with GFP-positive somata in *Scn1a*-GFP mouse neocortex. Values for Figure

14B-right panels. [a]Two animals were counted for each age. [b]L2/3, L5, L6: neocortical layer

II/III, V, VI. Cells in *Scn1a*-GFP mice (line #233) were counted; [c]L2/3 (P15, N = 601 cells;

4W, N = 422 cells), L5 (P15, N = 350 cells; 4W, N = 228 cells) and L6 (P15, N = 235 cells;

4W, N = 222 cells). Values are presented as mean ± SEM. +, positive; -, negative.

1468

1469

| | TBR1+, GFP+ | | TBR1-, GFP+ | | p-value | |
Region [a]	Area size [b]	Intensity [b]	Area size [b]	Intensity [b]	Area size [c]	Intensity [c]
L2/3	372.9 ± 7.5	41.8 ± 1.1	377.0 ± 11.0	43.8 ± 1.8	7.5×10^{-1}	2.9×10^{-1}
L5	337.5 ± 21.4	36.0 ± 1.7	481.6 ± 16.2	42.0 ± 1.5	8.6×10^{-5}	5.9×10^{-2}
L6	318.2 ± 10.3	35.9 ± 0.9	475.8 ± 24.3	50.6 ± 2.3	8.8×10^{-10}	1.1×10^{-9}

1470

Supplementary table S36. Area size and intensity of GFP immunosignals in

GFP-positive cells with TBR1-positive or negative nuclei in *Scn1a*-GFP mouse neocortex.

Values for Figure 15. [a]L2/3, L5, L6: neocortical layer II/III, V, VI. Cells in one *Scn1a*-GFP

mouse (line #233) at 4W were counted; [b]L2/3 (N = 189 cells), L5 (N = 118 cells) and L6 (N =

105 cells). [c]Statistical significance was assessed using t-test. Values are presented as mean ±

SEM. +, positive; -, negative.

1477

1478

1479

1480

Region[a]	per cells with AnkG+ AISs			
	% TBR1+[b]		% TBR1-[b]	
L2/3	34.5 ± 0.7		65.5 ± 0.7	
L5	52.8 ± 0.7		47.2 ± 0.7	
L6	66.3 ± 5.5		33.7 ± 5.5	
	% Nav1.1+	% Nav1.1-	% Nav1.1+	% Nav1.1-
L2/3	0.0 ± 0.0	34.5 ± 0.7	6.9 ± 0.6	58.6 ± 0.5
L5	0.0 ± 0.0	52.8 ± 0.7	10.2 ± 1.1	37.0 ± 1.7
L6	0.0 ± 0.0	66.3 ± 5.5	3.3 ± 0.7	30.4 ± 5.7

1481

Supplementary table S37. Percentage of cells with TBR1- and Nav1.1-positive/negative

nuclei or AISs per cells with ankyrinG-positive AISs in *Scn1a*-GFP mouse neocortex.

Values for Supplementary figure S11B-left panel. [a]L2/3, L5, L6: neocortical layer II/III, V, VI.

Cells in three *Scn1a*-GFP mice (line #233) at P15 were counted; [b]L2/3 (N = 879 cells), L5 (N

= 614 cells) and L6 (N = 816 cells). Values are presented as mean ± SEM. AnkG, ankyrinG;

+, positive; -, negative.

1488

1489

Region[a]	per cells with AnkG+ and Nav1.1+ AISs	
	% TBR1+[b]	% TBR1-[b]
L2/3	0.0 ± 0.0	100.0 ± 0.0
L5	0.0 ± 0.0	100.0 ± 0.0
L6	0.0 ± 0.0	100.0 ± 0.0

Supplementary table S38. Percentage of cells with TBR1-positive/negative nuclei per

cells with ankyrinG/Nav1.1-double positive AISs in *Scn1a*-GFP mouse neocortex. Values

for Supplementary figure S11B-middle panel. [a]L2/3, L5, L6: neocortical layer II/III, V, VI.

Cells in three *Scn1a*-GFP mice (line #233) at P15 were counted; [b]L2/3 (N = 59 cells), L5 (N =

63 cells) and L6 (N = 28 cells). Values are presented as mean ± SEM. AnkG, ankyrinG; +,

positive; -, negative.

1496

1497

| Region [a] | per cells with AnkG+ AISs and TBR1+ nuclei | |
	% Nav1.1+ [b]	% Nav1.1- [b]
L2/3	0.0 ± 0.0	100.0 ± 0.0
L5	0.0 ± 0.0	100.0 ± 0.0
L6	0.0 ± 0.0	100.0 ± 0.0

 Supplementary table S39. Percentage of cells with Nav1.1-positive/negative AISs per

 cells with ankyrinG-positive AISs and TBR1-positive nuclei in *Scn1a*-GFP mouse

 neocortex. Values for Supplementary figure S11B-right panel. [a]L2/3, L5, L6: neocortical

 layer II/III, V, VI. Cells in three *Scn1a*-GFP mice (line #233) at P15 were counted; [b]L2/3 (N =

 296 cells), L5 (N = 319 cells) and L6 (N = 552 cells). Values are presented as mean ± SEM.

 AnkG, ankyrinG; +, positive; -, negative.

| Region [a] | per cells with AnkG+ AISs | | | |
	% TBR1+ [b]		% TBR1- [b]	
L2/3	33.3 ± 5.0		66.7 ± 5.0	
L5	48.9 ± 2.5		51.1 ± 2.5	
L6	62.7 ± 3.5		37.3 ± 3.5	
	% Nav1.2+	% Nav1.2-	% Nav1.2+	% Nav1.2-
L2/3	22.6 ± 2.2	10.7 ± 2.9	57.4 ± 3.5	9.3 ± 1.6
L5	34.6 ± 1.6	14.3 ± 1.0	31.3 ± 0.9	19.8 ± 2.6
L6	42.4 ± 1.2	20.3 ± 3.3	26.5 ± 4.8	10.8 ± 2.2

 Supplementary table S40. Percentage of cells with TBR1- and Nav1.2-positive/negative

 nuclei or AISs per cells with ankyrinG-positive AISs in *Scn1a*-GFP mouse neocortex.

 Values for Supplementary figure S12B-left-upper panel. [a]L2/3, L5, L6: neocortical layer II/III,

 V, VI. Cells in three *Scn1a*-GFP mice (line #233) at P15 were counted; [b]L2/3 (N = 967 cells),

 L5 (N = 667 cells), and L6 (N = 833 cells). Values are presented as mean ± SEM. AnkG,

 ankyrinG; +, positive; -, negative.

1517

	per cells with AnkG+ and Nav1.2+ AISs	
Region [a]	% TBR1+ [b]	% TBR1- [b]
L2/3	28.5 ± 3.2	71.5 ± 3.2
L5	52.8 ± 1.4	47.2 ± 1.4
L6	62.4 ± 4.5	37.6 ± 4.5

1518

Supplementary table S41. Percentage of cells with TBR1-positive/negative nuclei per

cells with ankyrinG/Nav1.2-double positive AISs in *Scn1a*-GFP mouse neocortex. Values

for Supplementary figure S12B-middle-upper panel. [a]L2/3, L5, L6: neocortical layer II/III, V,

VI. Cells in three *Scn1a*-GFP mice (line #233) at P15 were counted; [b]L2/3 (N = 995 cells), L5

(N = 667 cells) and L6 (N = 923 cells). Values are presented as mean ± SEM. AnkG,

ankyrinG; +, positive; -, negative.

1525

1526

	per cells with AnkG+ AISs and TBR1+ nuclei	
Region [a]	% Nav1.2+ [b]	% Nav1.2- [b]
L2/3	69.3 ± 3.5	30.8 ± 3.5
L5	68.9 ± 2.7	31.1 ± 2.7
L6	68.7 ± 3.8	31.3 ± 3.8

1527

Supplementary table S42. Percentage of cells with Nav1.2-positive/negative AISs per

cells with ankyrinG-positive AISs and TBR1-positive nuclei in *Scn1a*-GFP mouse

neocortex. Values for Supplementary figure S12B-right-upper panel. [a]L2/3, L5, L6:

neocortical layer II/III, V, VI. Cells in three *Scn1a*-GFP mice (line #233) at P15 were

counted; [b]L2/3 (N = 317 cells), L5 (N = 309 cells) and L6 (N = 523 cells). Values are

presented as mean ± SEM. AnkG, ankyrinG; +, positive; -, negative.

1534

1535

1536

	per cells with TBR1+ nuclei	
Region [a]	% AnkG+ [b]	% AnkG- [b]
L2/3	62.9 ± 4.9	37.1 ± 4.9
L5	52.2 ± 5.5	47.8 ± 5.5
L6	52.3 ± 3.4	47.7 ± 3.4

1537

Supplementary table S43. Percentage of cells with AnkG-positive/negative AISs per cells
with TBR1-positive nuclei in *Scn1a*-GFP mouse neocortex. Values for Supplementary
figure S12B-lower panel. [a]L2/3, L5, L6: neocortical layer II/III, V, VI. Cells in three
Scn1a-GFP mice (line #233) at P15 were counted; [b]L2/3 (N = 509 cells), L5 (N = 582 cells)
and L6 (N = 998 cells). Values are presented as mean ± SEM. AnkG, ankyrinG; +, positive; -,
negative.

1544

1545

	per cells with TBR1+ nuclei			
Region [a]	% GFP+ [b]		% GFP- [b]	
L2/3	57.7 ± 5.0		42.3 ± 5.0	
L5	29.7 ± 6.0		70.3 ± 6.0	
L6	11.6 ± 2.5		88.4 ± 2.5	
	% Nav1.2+	% Nav1.2- [c]	% Nav1.2+	% Nav1.2- [c]
L2/3	30.2 ± 6.8	27.4 ± 2.5	20.4 ± 3.6	22.0 ± 2.7
L5	12.8 ± 5.6	16.3 ± 2.6	36.8 ± 5.4	34.1 ± 1.8
L6	5.0 ± 1.3	7.0 ± 2.3	36.6 ± 5.2	51.5 ± 2.8

1546

Supplementary table S44. Percentage of cells with GFP- and Nav1.2-positive/negative
somata and AISs per cells with TBR1-positive nuclei in *Scn1a*-GFP mouse neocortex.
Values for Supplementary figure S13B. [a]L2/3, L5, L6: neocortical layer II/III, V, VI. Cells in
three *Scn1a*-GFP mice (line #233) at P15 were counted; [b]L2/3 (N = 461 cells), L5 (N = 464
cells) and L6 (N = 913 cells). [c]In these values, because they were including AnkG-negative
cells, to obtain correct cell population for Nav1.2-/AnkG+ cells, virtual cell numbers were
estimated using ratio of AnkG+/TBR1+ cells in Supplementary figure S12B. Values are
presented as mean ± SEM. AnkG, ankyrinG; +, positive; -, negative.

1555

Yamagata et al., Figure 1

GFP IHC + Nissl stain (5W)

Tg #184   Tg #233

Wild-type   Wild-type

Yamagata et al., Supplementary figure S1

Yamagata et al., Figure 2

GFP (fluorescence)

#184 (P17)

#233 (P15)

Yamagata et al., Supplementary figure S2

Yamagata et al., Supplementary figure S3

GFP (fluorescence) #233

Yamagata et al., Figure 3

Yamagata et al., Supplementary figure S4

A

	GFP	NeuN	DAPI	Merge

Neocortex M1 (P15) — Layer II/III, Layer V, Layer VI

B

per cells with GFP+ and/or NeuN+ somata

M1 — L2/3, L5, L6 — 0, 50, 100 (%)

S1 — L2/3, L5, L6 — 0, 50, 100 (%)

■ GFP+, NeuN−   ■ GFP+, NeuN+   ■ GFP−, NeuN+

Yamagata et al., Supplementary figure S5

Yamagata et al., Figure 4

A

	GFP	Ankyrin G	Merge

Neocortex (P15)
- Layer II/III
- Layer V
- Layer VI

B

per cells with AnkG+ AISs

GFP+　GFP-

Yamagata et al., Supplementary figure S6

Yamagata et al., Supplementary figure S7

A GFP

B Nav1.1

C Ankyrin G

D GFP+Nav1.1

Hippocampus (P15)

CA1

CA2

CA3

DG

E GFP   Nav1.1   Ankyrin G   Merge

F

per cells with GFP+
somata and AnkG+ AISs

0   50   100 (%)

■ Nav1.1+   ■ Nav1.1-

per cells with Nav1.1+
and AnkG+ AISs

0   50   100 (%)

■ GFP+   ■ GFP-

Yamagata et al., Figure 5

Yamagata et al., Figure 6

Yamagata et al., Figure 7

Yamagata et al., Figure 8

Yamagata et al., Figure 9

A

B

Yamagata et al., Figure 10

Yamagata et al., Figure 11

Yamagata et al., Figure 12

Yamagata et al., Figure 13

Yamagata et al., Supplementary figure S8

Yamagata et al., Supplementary figure S9

A

Neocortex (P15)

	FEZF2	GFP	Nav1.2	Merge
Layer II/III				
Layer V				
Layer VI				

B

per cells with FEZF2+ nuclei

- GFP+, Nav1.2+
- GFP+, Nav1.2-
- GFP-, Nav1.2+
- GFP-, Nav1.2-

Yamagata et al., Supplementary figure S10

Yamagata et al., Figure 14

Yamagata et al., Figure 15

A

| | TBR1 | Nav1.1 | Ankyrin G | Merge |

Neocortex (P15)
Layer II/III
Layer V
Layer VI

B

per cells with AnkG+ AISs

per cells with AnkG+ and Nav1.1+ AISs

per cells with AnkG+ AISs and TBR1+ nuclei

0   50   100 (%)

L2/3
L5
L6

■ TBR1+, Nav1.1+   ■ TBR1+, Nav1.1-   ■ TBR1-, Nav1.1+   ■ TBR1-, Nav1.1-

Yamagata et al., Supplementary figure S11

Yamagata et al., Supplementary figure S12

Yamagata et al., Supplementary figure S13

Neocortical Layer V (L5) neurons

Yamagata et al., Supplementary figure S14