Temporal perturbations cause movement-context independent but modality-specific sensorimotor adaptation

Nadine Schlichting*, Tatiana Kartashova, Michael Wiesing, Eckart Zimmermann

Institute for Experimental Psychology

Heinrich-Heine-University Düsseldorf, Germany

Address for correspondence:

Nadine Schlichting

Universitätsstraße 1

40225 Düsseldorf, Germany

nadine.schlichting@hhu.de

Abstract

Complex, goal-directed and time-critical movements require the processing of temporal features in sensory information as well as the fine-tuned temporal interplay of several effectors. Temporal estimates used to produce such behavior may thus be obtained through perceptual or motor processes. To disentangle the two options, we tested whether adaptation to a temporal perturbation in an interval reproduction task transfers to interval reproduction tasks with varying sensory information (visual appearance of targets, modality, virtual reality (VR) environment or real-world) or varying movement types (continuous arm movements or ballistic clicking movements). Halfway through the experiments we introduced a temporal perturbation, such that continuous pointing movements were artificially slowed down in VR, causing participants to adapt their behavior to sustain performance. In four experiments, we found that sensorimotor adaptation to temporal perturbations is independent of environment context and movement type, but modality specific. Our findings suggest that motor errors induced by temporal sensorimotor adaptation affect the modality specific perceptual processing of temporal estimates.

Keywords: interval timing, visuomotor adaptation, motor planning, VR

Graphical Abstract

New & Noteworthy

Without calibration through error signals from the external world, internal sensory estimates could only be interpreted arbitrarily. In the present work, we investigated whether actions calibrate temporal estimates using an adaptation paradigm. We found that adaptation transfers between different types of movement and movement-goals, but only if the modality of the adapted task and the transfer-task was the same. Our results suggest that motor errors calibrate time upstream of motor areas, probably in sensory areas.

Introduction

In a 100 m sprint race, a good start can make the difference in winning the gold or silver medal. One crucial aspect to nail the start is to measure the interval between the *ready-* and *set-* signal in order to predict when the *go*-signal will occur, so that the athlete can immediately leave the starting block and, ideally, save precious milliseconds. To generate a prediction about the time of the go-signal, the sprinter has to measure the duration of the first interval and reproduce it by initiating the sprint.

The measurement of the first interval duration might be accomplished in perceptual areas, in the sprinter example by the auditory system. The temporal estimate is then handed to motor areas in order to generate a movement as soon as the interval exceeds. However, a more economical solution to the problem would be if these estimates are directly entailed in the motor planning of the sprint start movements, since transfer between representations might induce noise and delays (see also 1). Indeed, Jazayeri and Shadlen (2) demonstrated in macaques that in intra-parietal cortex temporal intervals are measured prospectively in relation to the desired motor plan to reproduce these intervals. If temporal estimates are entailed in motor planning and actions, then motor planning and actions may alter our ability to measure time conversely. Indeed, evidence for the influence of motor actions on duration estimates is accumulating: The frequency of finger tapping (3–5); the length (6) or type (7) of movements; or the mere preparation of a ballistic action (8) bias temporal estimates. In these studies, the motor action is task-irrelevant and experimental manipulations depend on the participant's ability to consciously alter specific movement parameters. Hence, it is impossible to say whether there is a clock-system informing the motor system and vice versa, or whether they are essentially the same. Interval production and reproduction paradigms in which, just like in the sprint example, a time-critical motor action (e.g., a saccade, button press, or arm movement) is an essential part the task offer additional insights into the entanglement of the motor system and time. One such insight is that the well-known temporal context effect in interval reproduction tasks, in which duration estimates are biased towards previously perceived intervals (for a review, see 9), has been shown to originate in motor areas of the brain (2, 10) and occur already during the encoding of the to-be-reproduced interval during which no motor action is executed (11, 12), and during which decision biases can be ruled out (13).

We approached the question of whether temporal estimates are obtained through perceptual or motor processes by inducing temporal adaptation in a motor reproduction task and by then measuring transfer of adaptation. We tested transfer of adaptation to a motor task that was not

trained during adaptation or we kept the motor reproduction task identical but varied the sensory stimuli that defined the temporal intervals and measured adaptation transfer. If adaptation takes place in perceptual areas, one would expect to observe adapted interval reproductions irrespective of the motor tasks that is used to respond. By contrast, if motor areas generate temporal estimates, only the motor task that was adapted should produce reproduction estimates that differ from baseline behavior. We asked participants to reproduce intervals by performing different time-critical actions. The temporal estimate was directly contingent on the movement dynamics, and the error following a movement would inform about its temporal accuracy. The study was conducted in Virtual Reality (VR), enabling us to provide systematically distorted feedback about the movement and its temporal accuracy that leads to temporal adaptation of the movement. In a ready-set-go paradigm, the interval between *ready* and *set* had to be reproduced by performing either a ballistic finger movement (clicking a button on the controller, *clicking reproduction*) or by a continuous movement of the arm to hit a target at the time of the *go*-signal (*pointing reproduction*; note that contrary to the sprint example, the movement had to be completed by the time of the *go*-signal, see Figure 1A). During the adaptation phase, feedback about pointing performance was manipulated such that movements were artificially slowed down, forcing participants to adapt their movement speed in order to sustain performance. In four experiments, we tested whether adaptation to a temporal perturbation is specific to the movement or task (transfer to ballistic finger movement; Experiment 1 and 2), interval range (Experiment 1), target location (Experiment 2), the VR setting itself (Experiment 3), or modality (Experiment 4).

Experiment 1: Task independency of temporal sensorimotor adaptation

Materials & Methods

Apparatus. All experiments were conducted on a Windows 10 based desktop computer (Alienware Aurora R8, Intel(R) Core™ i7-8700 CPU @ 3.20 GHz, 16 GB RAM, NVIDIA GeForce GTX 1080TI graphics card) connected to an HTC Vive Pro Eye Head Mounted Display (HMD) (HTC Corporation, Taoyuan, Taiwan). The HMD presents stimuli on two low-persistence organic light-emitting diode (OLED) displays with a resolution of 1,440 x 1,600 pixels per eye and a refresh rate of 90 Hz. Additionally, participants used a Vive motion-controller for their right hand. The virtual environment (VE) was rendered using SteamVR and a custom-made program created in Unity game engine, version 2019.1.13f1 (Unity Technologies, San Francisco, U.S.). Head and hand movements were tracked via the HMD and controller using the SteamVR 1.0 tracking system.

According to previous research, this system provides a robust tracking of head and hand motion with a 360° coverage, provided tracking loss is prevented (14). Tests of Verdelet et al. (15) demonstrated a submillimeter precision (0.237 mm) and an accuracy of 8.7 mm for static and 8.5 mm for dynamic objects. While the system can update the user's pose (position and orientation) at a higher rate (up to 1000 Hz for the HMD and 250 Hz the controllers), in this study the sampling rate for both HMD and controller was limited by the HMD's refresh rate of 90 Hz. Because participants always responded to stimuli in front of them, we did not need the full coverage around participants in the present study. Hence, in order to minimize the chance of occlusions of the HMD or the controller and thereby avoiding tracking loss, our setup had both base stations facing the participant. Throughout the experiment, participants held the controller with an outstretched index finger placed on top of the controller with the fingertip matching the tracking origin of the controller as close as possible (see Figure 1A). Participants' hands were presented as gloves instead of bare hands. Previous research has shown that the appearance of self-avatars can influence the self-perception and behavior of participants. Most notably the so-called Proteus Effect, which describes the tendency of participants to infer their expected behaviors and attitudes from their self-avatar's appearance (16, 17). By presenting gloves we were able to cover several characteristics that might otherwise be incongruent to the participants real hands, such as the skin color or gender of the hands.

Participants remained seated during the entire experiment. The apparatus is the same for all reported experiments.

Participants. 12 participants (8 female, 4 authors, age range 19-42 years, all right-handed) were tested in exchange for a monetary reward (€10 per hour) or course credits. Sample size in all experiments was based on previous similar research (e.g., 5-8 participants per experiment in Anobile et al. (2020). All participants had normal or corrected-to-normal vision. Participants gave informed consent prior to participation. The experiments were carried out along the principles laid down in the Declaration of Helsinki. All experiments were approved by the local ethics committee of the psychological department of the Heinrich-Heine-University Düsseldorf. This holds for all reported experiments.

Pointing reproduction task. In the pointing task participants had to measure and immediately reproduce a 1s interval by reaching a target with the controller (ready-set-go paradigm, see Figure 1A, left column). At the beginning of the trial participants had to place the controller behind the start line in a sphere (diameter = 10 cm), which was located slightly to their

right bottom at x = 20 cm, y = -40 cm, and z = 30 cm, with respect to their head position (x = 0, y = 0, z = 0 cm). To their left they saw a small black sphere, the target (diameter = 3 cm, x = -15, y = -40, z = 30 cm, distance between start line and target was 30 cm). The target changed its color to red for 0.1 s, first to mark the start of the interval presentation (*ready*) and again after 1s to mark the end of the interval presentation and the start of the reproduction (*set*). Participants had to virtually touch the sphere to end their reproduction (*go*). As soon as participants crossed the start line with the controller, they saw a VR-hand following the movement of their physical hand. We will refer to the time between crossing the start line and reaching the target as *movement duration*. Participants received immediate feedback on their performance: The deviation of participants' reproduction from the target interval (a negative number corresponded to under-reproductions) was displayed above the target, additionally color-coded in red (deviation > 0.3 s), yellow (0.1 > deviation < 0.3 s), or green (deviation < 0.1 s). The next trial started once participants moved their hand back to the start position and pressed the trigger-button of the controller with their middle finger. We used three different kinds of trials: Training-trials, adaptation-trials, and test-trials (see Figure 1B for a temporal outline of the experiment). Training-trials were as described above and used to accustom participants to the VE and for de-adaptation at the end of the experiment. In adaptation-trials participants saw the VR-hand move at half the speed of their actual movement, that is, participants received delayed visual feedback by means of the VR hand. The reproduction of the 1 s interval depended on the VR-hand reaching the sphere with the index finger, so that for an accurate reproduction, participants had to adapt their movement by speeding up the pointing action. An alternative strategy is to start the hand movement earlier without changing movement speed, or to use a combination of both faster movement and earlier movement start. Changes in movement speed are thought to reflect implicit adaptation, while changes in movement onset are thought to be more cognitively controlled (18). Feedback was given on every trial. In test-trials participants did not see the VR-hand movement and received no feedback.

Figure 1. Experiment 1: Task independency of temporal sensorimotor adaptation. A) In a pointing trial, participants had to reproduce the interval marked by the ready- and set-signal by reaching the target (black sphere) in time for the go-signal. Visual feedback was provided by means of a VR-hand, appearing as soon as the movement was initiated. In clicking trials, the interval between the ready- and set-signal had to be reproduced by clicking the trigger button. No additional movement feedback was provided. Participants wore the VR headset at all times. *B)* Temporal outline and trial order of a single session. *C)* reproductions in the pointing (top panel) and clicking reproduction task (bottom panel). Data was averaged over all three sessions. Error bars represent 95% within-subject CIs (19, 20).

Clicking reproduction task. For the clicking task, we used the same ready-set-go paradigm as in the adaptation task. Participants ended their reproductions (i.e., indicate the go-signal) by pressing the trigger button of the controller (see Figure 1A, right column). Tested intervals ranged between 0.85 s and 1.15 s in steps of 0.05 s. In training-trials participants received feedback as described above. In test-trials participants received no feedback on their performance.

Procedure. Figure 1B depicts the temporal outline of the experiment. Each participant completed the experiment three times in three sessions, separated by at least 4 hours. In each session participants first got accustomed to the VE and the tasks by completing 10 training-trials of both tasks. In the pre-adaptation phase, each participant completed 10 test-trials of the pointing

task and 70 test-trials of the clicking task (i.e., each interval was presented 10 times). This first test phase was followed by 40 adaptation-trials of the pointing task. The 70 post-adaptation clicking reproduction task trials were interleaved with adaptation trials of the pointing task (five trials each). This was followed by 10 test-trials of the pointing task, again interleaved with adaptation-trials. At the end of the experiment, participants completed 10 more training-trials of the pointing task to de-adapt.

Statistical analysis. As an individual measure of the magnitude of motor adaptation we calculated the proportional change in the averaged movement onset in pre- versus post-adaptation pointing test-trials (($onset_{post}$ - $onset_{pre}$) / $onset_{post}$) for each participant and session separately, reflecting the amount of implicit motor learning in each participant and session. Movement onset was identified as the most likely candidate to quantify adaptation. This was tested by fitting linear and exponential models and comparing their performance in predicting development of movement onset and movement duration over the course of the adaptation phase. Additionally, we compared the average movement duration and movement onset of five trials in the beginning (omitting the very first five trials of the adaptation phase, so that fast, strategic recalibration does not obscure this measure) and at the end of the adaptation phase by means of two-sided t-tests. A difference between early and late adaptation phase would hint at gradual changes in behavior over the course of trials.

Participants' behavior in the pointing reproduction task was analyzed by means of Linear-Mixed-Models (LMMs) using the lme4 (21) and lmerTest (22) packages in R version 4.0.3 (23). Models were constructed to predict the reproduced 1 s interval with predictors including adaptation (coded as 0 for all pre-adaptation trials and as the difference in movement onset in pre- and post-adaption trials in the pointing task, dependent on participant and session, see above) and session (1-3, as factor). In all models, intercepts varied by participant. All models with different combinations of these predictors, with or without interactions between main effects, were compared by means of likelihood ratio tests, BIC values (we consider a reduction of 10 as evidence to include a given factor) and Bayes Factors (BF$_{10}$; throughout the manuscript we report the BF as the evidence of the alternative hypothesis over the null hypothesis; we consider BF$_{10}$ > 3 as evidence for the alternative hypothesis) calculated using the BayesFactor package (24). We report the resulting best model and statistical evidence for or against effects.

Models that were constructed to predict participants' behavior in the clicking reproduction task incorporated the following predictors: Interval length (i.e., the to-be-reproduced interval, zero-

centered), previously presented durations (i.e., interval duration of trial N-1, N-2, etc.), adaptation (see above), and session (see above). In all models, intercepts varied by participant. We proceeded with model selection as described above.

To rule out that effects of time-in-experiment (e.g., fatigue) are driving differences in pre- and post-adaptation performance, we compared root mean squared errors (RMSEs), calculated from an estimated linear slope and scaled by mean reproduction for each participant (25), as a measure of the variable error in pre- and post-adaptation performance in both tasks. This test was based on the rationale that with increasing fatigue, reproductions in post-adaptation trials should become more variable than in pre-adaptation trials. For the analysis, we constructed models with predictors including: Interval length (for the clicking task only, coded as described above), session, and adaptation (coded as described above). In all models, intercepts varied by participant.

Finally, to quantify the amount of transfer from the adapted pointing to the clicking reproduction task, we compared the difference between pointing reproductions in test-trials before and after adaptation with the change in clicking reproductions induced by the adaptation procedure:

$$transfer = \frac{(pointing\ reproduction_{post} - pointing\ reproduction_{pre})}{(clicking\ reproduction_{post} - clicking\ reproduction_{pre})} \times 100 \qquad (1)$$

Trials in which the reproduced duration was shorter or longer than 3 median absolute deviations were excluded from the analysis (26). This led to the exclusion of 2.2% of trials in the pointing task and 2.6% of trials in the clicking task.

Results

Adaptation phase. The adaptation time course is shown in Figure 2, with separate curves for movement duration (bottom, dark grey), movement onset (middle, medium grey), and interval reproductions (top, light grey). During the adaptation phase movement onset exhibits clear signs of adaptation: Initiation of the pointing movement adapted slowly and gradually over the course of adaptation-trials. Movement duration, in contrast, adapted within a few trials and was retained at this level during the remaining adaptation phase. For both movement onset and duration, a linear model fitted the data best and reflected the gradual decrease in movement duration (slope = -0.0004 $\pm$ 0.0001 SE, $X^2(1) = 9.53$, $p < .001$, ΔBIC = 6.26, $BF_{10} = 49.86$) and movement onset (slope = -0.002 $\pm$ 0.0003 SE, $X^2(1) = 36.36$, $p < .001$, ΔBIC = 30.19, $BF_{10} = 50.29$) over the course of the adaptation phase. Comparing measures during early (trial 6-10) and late phases of adaptation (trial 36-40), we found differences for movement onset ($t(11) = 3.38$, $p = .006$, $d = 0.98$, $BF_{10} = 8.76$),

but not for movement duration ($t(11) = -0.64$, $p = .53$, $d = -0.19$, $BF_{10} = 0.34$). Adaptation strength was calculated based on pre- and post-movement onset durations ($M = -0.16$, 95% CI [-0.23, -0.10]). The smaller this value, the more sensorimotor adaptation was observed. Because more strongly adapted participants presumably show more adaptation after- or transfer-effects, we incorporated this parameter in the subsequent analyses.

Figure 2. Experiment 1: Adaptation time course. Reproduction (top curve, light grey) and *movement start* (middle curve, medium grey) were measured relative to the onset of the *set*-signal. *Movement duration* (bottom curve, dark grey) was measured from the time the controller passed the start line. Note that the *movement duration* and *movement onset* lines do not add up to replicate the *reproduction* line because of averaging, on a single subject/single session basis they do. Shaded error bars represent 95% CIs, red lines represent linear fits. Data points on the right represent movement onset (top) and -duration (bottom), averaged over five trials in the early and late phase of adaptation (see colored bars in A), as a measure of adaptation.

Pointing reproductions. For the analysis of reproduction performance in test trials, depicted in Figure 1C, top panel, we constructed linear mixed models (LMMs) to predict 1 s pointing reproductions. The final model to predict reproductions included the factor adaptation strength (coded as 0 for all pre-adaptation trials, and as the above described adaptation strength based on movement onset for all post-adaptation trials; $X^2(1) = 294.72$, $p < .001$, $\Delta BIC = 288.17$, $BF_{10} > 1000$). In other words, 1 s pointing reproductions in post-adaptation trials were shorter than in pre-adaption trials, and this difference was greater for participants who adapted their movement onset more strongly. Adding the factor session was not warranted ($X^2(2) = 3.16$, $p = .21$, $\Delta BIC = -9.95$, $BF_{10} = 0.04$), meaning there was no difference in performance between sessions.

To rule out that effects of time-in-experiment (e.g., fatigue) are driving differences in pre- and post-adaptation performance, we compared RMSEs as a measure of the variable error in pre- and post-adaptation performance in both tasks. RMSEs were not affected by adaptation ($X^2(1) = 0.78$, $p = .377$, $\Delta\text{BIC} = -3.50$, $\text{BF}_{10} = 0.35$). This means that it is unlikely that differences in performance in pre- and post-adaptation trials were driven by time-in-experiment effects. Including session as a predictor was warranted following the likelihood ratio test, however, BIC-values and Bayes Factor analysis did not lead to the same conclusion ($X^2(2) = 7.18$, $p = .028$, $\Delta\text{BIC} = -1.37$, $\text{BF}_{10} = 1.92$). Thus, there is mixed evidence concerning the stability of precision over sessions.

Additional information on pointing trajectories and velocity profiles can be found online at https://osf.io/zbgy9/.

Clicking reproductions. Figure 1C, bottom panel, depicts reproduction performance for the seven different intervals, split on pre- and post-adaption trials. In the LMM analysis, the general trend that longer intervals were reproduced as longer was captured by including the factor interval length ($X^2(1) = 204.48$, $p < .001$, $\Delta\text{BIC} = 195.98$, $\text{BF}_{10} > 1000$). Regression-towards-the-mean effects were captured by including the duration of the previous trial (N-1, $X^2(1) = 18.24$, $p < .001$, $\Delta\text{BIC} = 9.74$, $\text{BF}_{10} = 564.51$) and of the trial before the previous (N-2, $X^2(1) = 6.87$, $p = .009$, $\Delta\text{BIC} = -1.62$, $\text{BF}_{10} = 1.9$) in the final model. Note that evidence for the inclusion of the factor N-2 is ambiguous. Apart from this general regression towards the mean, it appears that intervals were systematically underestimated following adaptation-trials of the pointing task. This is reflected in the final model by including the factor adaptation strength ($X^2(1) = 154.59$, $p < .001$, $\Delta\text{BIC} = 146.10$, $\text{BF}_{10} > 1000$), showing that more strongly adapted participants in the pointing task show larger transfer to the clicking task. Together, these results suggest that 1) adaptation effects transferred to another interval reproduction task in which the movement required to produce the go-signal differed substantially from the one that was adapted (clicking instead of pointing); 2) adaptation generalizes to a broader range of intervals; and 3) participants who adapted their motor behavior more strongly in the pointing task also showed larger differences in pre- and post-adaptation clicking reproductions. The parallel existence of adaptation and temporal context effects suggests that sensorimotor adaptation and context effect do not interact or cancel each other out, but affect reproductions independently. While temporal context has been shown to already affect the initial temporal estimate (11, 12), its neural origins have been found in motor areas (2). In neuroimaging studies investigating the locus of sensorimotor adaptation, the cerebellum is, apart

from cortical motor areas, thought to play a critical role (18). Thus, these two effects may originate in different neural substrates (e.g., LIP/SMA vs. cerebellum) or during different processing stages (e.g., perception vs. motor prediction or planning).

We did not find evidence for effects of adaptation ($X^2(1) = 0.76$, $p = .384$, ΔBIC = -5.46, $BF_{10} = 0.14$) or session ($X^2(2) = 5.06$, $p = .080$, ΔBIC = -7.39, $BF_{10} = 0.24$) on RMSE. Errors adhered to Weber's law (27) and varied for different intervals ($X^2(1) = 12.78$, $p < .001$, ΔBIC = 6.56, $BF_{10} = 49.71$). As for the analysis of RMSEs in the pointing task, these results speak against an interpretation of pre-post-adaptation performance differences being driven by, for example, increased fatigue over the course of the experiment.

Transfer. To quantify the amount of adaptation transfer from the pointing to the clicking reproduction task, we calculated transfer as the difference between pointing reproductions in test-trials before and after adaptation with the change in clicking reproductions induced by the adaptation procedure. In this experiment, 41.00 % (95% CI [18.59, 63.41]) of the adaptation effect in the pointing task transferred to the clicking task, which, tested with a one-sample t-test, differed significantly from zero ($t(11) = 4.03$, $p = .002$, $d = 1.16$, $BF_{10} = 22.41$).

Discussion

Results of Experiment 1 revealed that sensorimotor adaptation affected participants' timing abilities and transferred to another type of movement. As already touched upon in the introduction, sensorimotor adaptation effects are often reported to exhibit spatial selectivity, that is, adaptation effects are observed only in the region in which the adapter was shown or the adaptation movement was executed. In Experiment 1, the coordinates of the target (the black sphere) are the same in both tasks and in all kinds of trials. To test in how far adaptation effects in our VR setup are location specific, we changed the target position in test-trials of the pointing and/or the clicking task in Experiment 2, meaning that in some cases the target in test trials appeared mirrored compared to adaptation trials.

Experiment 2: Location independency of temporal sensorimotor adaptation

Materials & Methods

Participants. 7 participants (4 female, 3 authors, age range 23-42 years, all right-handed) who already participated in Experiment 1 and 2 were re-tested in Experiment 2.

Pointing reproduction task. The pointing task was essentially the same as in Experiment 1 (Figure 3A, left column), with the only differences that the visual scene for test trials was mirrored in two out of four sessions (i.e., in mirrored sessions participants had to point from left to right).

Figure 3. Experiment 2: Location independency of temporal sensorimotor adaptation. A) In a pointing trial, participants had to reproduce the interval marked by the ready- and set-signal by reaching the target (black sphere) in time for the go-signal. Visual feedback was provided by means of a VR-hand, appearing as soon as the movement was initiated. In clicking trials, the interval between the ready- and set-signal had to be reproduced by clicking the trigger button. No additional movement feedback was provided. The location and pointing direction in test trials varied (original and mirrored). Participants wore the VR headset at all times. *B)* Temporal outline and trial order of the four different sessions. White errors inform about the target location. *C)* Reproductions in the pointing (top panel) and clicking reproduction task (bottom panel). Data was pooled together depending on whether the target changed location in the pointing task for pointing reproductions, or whether the target changed location in the clicking task for clicking reproductions. Error bars represent 95% within-subject CIs (19, 20).

Clicking reproduction task. The clicking task was essentially the same as in Experiment 1 (Figure 4A, right column), with the only differences that we used only the 1s interval; and that the visual scene for test trials was mirrored in two out of four sessions (Figure 3B).

Procedure. Each participant completed four sessions, separated by at least two hours. Sessions differed in the way mirrored conditions were combined: Test trials were mirrored 1) for both tasks, 2) for the pointing reproduction task only, 3) for the clicking reproduction task only, or 4) for none of the tasks. Trial structure was identical to Experiment 1, with the only exception that only 30 pre- and post-adaptation clicking test-trials were performed. The temporal outline of the experiment is depicted in Figure 3C.

Statistical analyses. Analysis for the location dependent motor adaptation experiment was identical to Experiment 1, with the only exception that 'interval' was not included as a predictor in any model, and we added the predictor location (coding for whether the target in test-trials appeared at the same or different location than in adaptation-trials). 0.9% of trials in the pointing task and 6.7% of trials in the clicking task were excluded from the analysis (see Experiment 1 for exclusion criteria).

Results

Pointing reproductions. Analogue to the model analysis performed for pointing reproductions in Experiment 1, including the factor adaptation strength ($M_{same\ location}$ = -0.13, 95% CI [-0.26, 0]; $M_{different\ location}$ = -0.01, 95% CI [-0.14, 0.1]) to predict interval reproductions in the pointing task improved the model fit ($X^2(1) = 17.32$, p < .001, $\Delta BIC = 11.01$, $BF_{10} = 466.87$). To test whether target location in test trials affected reproductions additionally, we included the factor location (coded as "same" or "different" compared to adaptation trials). Location did, however, not improve the model fit ($X^2(1) = 1.27$, p = .26, $\Delta BIC = -5.05$, $BF_{10} = 0.18$). This shows that, regardless of whether the target in test-trials appeared at the same location as in adaptation-trials, reproductions were influenced by sensorimotor adaptation in that reproductions were systematically shorter in post-adaptation compared to pre-adaptation test-trials (Figure 3C, top panel).

There was no evidence for effects of adaptation ($X^2(1) < .01$, p = .995, $\Delta BIC = -3.33$, $BF_{10} = 0.34$) or location ($X^2(1) = 0.10$, p = .755, $\Delta BIC = -3.23$, $BF_{10} = 0.36$) on RMSEs, ruling out time-in-experiment effects or location dependent training effects.

Clicking reproductions. Figure 3C, bottom panel, depicts reproduction performance, split on pre- and post-adaption trials and on the location of the target. The final model included the predictor adaptation strength (($M_{same\ location}$ = -0.08, 95% CI [-0.23, 0.06]; $M_{different\ location}$ = -0.06, 95% CI [-0.21, 0.09]), $X^2(1) = 17.32$, $p < .001$, $\Delta BIC = 11.01$, $BF_{10} = 488.11$), while the location of the target did not improve the model fit ($X^2(1) = 1.25$, $p = .26$, $\Delta BIC = -5.07$, $BF_{10} = 0.18$). Thus,

intervals in the clicking task were systematically under-reproduced after adapting to the altered temporal properties in the pointing task. As for the pointing task, effects of adaptation on clicking reproductions are independent of location.

We did not find evidence for effects of adaptation ($X^2(1) = 1.83$, $p = .176$, ΔBIC $= -1.50$, $BF_{10} = 0.69$) or location ($X^2(1) = 1.76$, $p = .185$, ΔBIC $= -1.57$, $BF_{10} = 0.66$) on RMSEs.

Transfer. Because we did not find evidence for an effect of location on pointing reproductions, nor on clicking reproductions, we averaged over all location conditions for the calculation of transfer from adaptation in the pointing to the clicking task. The amount of adaptation transfer was 59.62 % (95% CI [23.99, 95.49], $t(6) = 4.09$, $p = .006$, $d = 1.55$, $BF_{10} = 9.49$). Adapting to altered temporal aspects of timed actions seems to affect all motor planning regarding direction or type of movement.

Discussion

In the previous experiments we found that sensorimotor adaptation to temporal perturbations generalizes to all motor actions aimed to reproduce a given interval. These results, however, could be caused by a general correction of movements to overcome the temporal lag associated to the VE, and not by adaptation of the motor system. To rule out that the above described findings apply to VR only, we tested whether adaptation effects also transfer to a pointing task outside of VR.

Experiment 3: Environment independency of temporal sensorimotor adaptation

Materials & Methods

Apparatus. In the real-world task, participants had to point to a green LED, attached to an Arduino microcontroller and controlled by Unity, which was positioned at the same location as the target in the VE (see schematic outline in Figure 4A).

Participants. 5 participants (3 female, 2 authors, age range 23-36 years, all right-handed) who already participated in Experiment 1, 2 and 3 were re-tested in Experiment 3.

Figure 4. Experiment 3: Environment independency of temporal sensorimotor adaptation. A) Participants had to reproduce the interval marked by the ready- and set-signal by reaching the target (black sphere in VR, green LED in real-world) in time for the go-signal. In VR, visual feedback was provided by means of a VR-hand, appearing as soon as the movement was initiated. In real-world, participants could see their hand. Participants wore the VR headset dependent on task requirements. *B)* Temporal outline and trial order. *C)* Averaged reproductions in the pointing task separated for VR and real-world task (top panel). Influence of adaptation in single subject data separated for VR and real-world task (connected dots represent one subject), plotted as pre-adaptation reproductions against post-adaptation reproductions. Dots below the dashed line reflect effects of adaptation (under-reproduction in post-adaptation trials compared to pre-adaptation trials), and the distance to the dashed line reflects the strength of adaptation (larger distance ~ larger adaptation effect). Error bars represent 95% within-subject CIs (19, 20).

Pointing reproduction task. The pointing task was the same as in Experiment 1, with the only differences that an additional set of 10 test-trials were performed outside of the VE (see Figure 4A and B). For the real-world pointing task, a green LED, attached to an Arduino microcontroller controlled by Unity, was positioned in the physical location directly below the virtual target. To ensure that participants were able to find the start position in the non-VR condition, a blue LED was attached to the Arduino and lit up when the hand was overlapping with the virtual start position.

Procedure. This experiment comprised one session only. Participants first got accustomed to the VE and the task by completing 10 training-trials. In the pre-adaptation phase, each participant completed 10 test-trials of the pointing task within and outside of the VE. The pre-adaptation test phase was followed by 40 adaptation-trials. In the post-adaptation phase participants again performed 10 test trials of the pointing task within and outside of the VE, interleaved with adaptation-trials. At the end of the experiment, participants underwent the de-adaptation procedure as in the other experiments.

Statistical analyses. Analysis for the VR-dependent motor adaptation experiment was identical to Experiment 2, with the exception that instead of the predictor 'location' we included the binary predictor 'VR'. The transfer from adaptation within VR to adaptation outside VR was calculated following formula (1). No trials had to be excluded from the analysis (see Experiment 1 for exclusion criteria).

Results

Pointing reproductions. While including the factor adaptation strength to predict reproductions in the pointing task did not improved the model fit (M_{VR} = -0.01 (95% CI [-0.21, 0.18]), $M_{real\text{-}world}$ = -0.13 (95% CI [-0.32, 0.07]), $X^2(1) = 3.63$, $p = .06$, ΔBIC = -1.64, $BF_{10} = 0.96$), including a binary factor adaptation (coded as "pre" and "post") did significantly improve model fits ($X^2(1) = 58.70$, $p < .001$, ΔBIC = 53.43, $BF_{10} > 1000$). Including a factor encoding environment (coded as "VR" or "real-world") was not warranted ($X^2(1) = 0.02$, $p = .90$, ΔBIC = -5.26, $BF_{10} = 0.14$). This shows that, regardless of whether the pointing task was performed within the VR environment or not, reproductions were influenced by sensorimotor adaptation, irrespective of how strongly participants were adapted (Figure 4C).

There was no evidence for effects of adaptation ($X^2(1) = 0.95$, $p = .33$, ΔBIC = -2.05, $BF_{10} = 0.55$) or VR ($X^2(1) = 0.21$, $p = .65$, ΔBIC = -2.79, $BF_{10} = 0.43$) on RMSEs, ruling out time-in-experiment effects or environment dependent effects.

Transfer from VR to the real world. The amount of adaptation transfer from VR to non-VR was 85.16% (95% CI [41.7, 128.62], $t(4) = 5.44$, $p = .006$, $d = 2.43$, $BF_{10} = 10.81$). Adaptation was not contextually cued by being in the VE or by wearing VR goggles. Instead, the crucial component driving the transfer of adaptation effects from VR to the real world may be the movement goal – reaching the target at the go-signal – which was the same within VR and the real world. The finding of substantial transfer further confirms the presence of sensorimotor adaptation,

because, unlike in VR test-trials, participants can see their hand in the analogue task and thus have immediate visual feedback on their movement time course.

Experiment 4: Modality dependence of temporal sensorimotor adaptation

Materials & Methods

Apparatus. The experimental hardware was identical to the previous experiments. During the experiment, participants remained seated in front of a table. A rubber mat (20 x 20 x 1 cm) served as the home position on which participants placed their right hand between trials. Additionally, two speakers (Genuine Altec Lansing Rev A00) where placed 30 cm to the left/right and 25 cm in front of the participant. The speakers were not rendered in the VE.

Participants. 17 participants (12 female, age range 19-30 years, one left-handed), two of which already participated in one or more of the previously reported experiments, were tested in Experiment 4.

Virtual Environment. In contrast to previous experiments, Experiment 4 was created in Unreal Engine 4.25 (UE) and a new VE was developed for the experiment (Figure 5A). The VE consisted of a customized version of UE's "Archviz Interior" sample project (28). The original sample makes use of real-time raytracing, a modern rendering technique that is currently too performance-intensive to meet the high performance demands of VR rendering. Instead, we replaced all ray-tracing effects by precalculated lighting and reflections. Additionally, all 3D models and textures were optimized (i.e., lowered in resolution) until we were able to hit stable 90 frames per second (FPS).

Within the VE, participants were sitting in front of a virtual table (160 x 60 x 70 cm), which was co-located with a physical table. The starting line of the previous experiments was replaced by the rubber mat (20 x 20 cm), which was also co-located with its physical counterpart. The center of the mat was placed about 50 cm in front and 24 cm to the right of the participants.

Pointing reproduction task. The pointing reproduction task (depicted in Figure 6A) was essentially the same as in Experiment 1. Again, participants were required to reproduce a 1 s interval by touching a target with their right index-finger. To start a trial, participants had to place their right hand on the start location and press the thumbpad. Afterwards, either a mid-gray target sphere (diameter 7 cm) or a mid-gray buzzer (diameter 9 cm) appeared about 50 cm in front and 27 cm to the left of the participants. After a randomized interval of 0.5 – 1.0 s, the target lit up in bright gray for 0.1 s to mark the start of the interval presentation (*ready*) and again after 1 s to mark

Figure 5. Virtual Environment of Experiment 4. A) Screenshots of the Virtual Environment *B)* Participants held the controller with the thumb being on the thumbpad and an outstretched index finger. The controller was not visible in the experiment.

again after 1 s to mark the end of the interval presentation and the start of the reproduction (*set*). As in the previous experiment, participants had to hit the target with their index-finger of the right hand to end the reproduction (*go*). While the target sphere had only to be touched, the buzzer had to pressed down by 1 cm with the right index-finger before a response was registered. The buzzer and sphere conditions were tested in separate sessions. In both conditions, as soon as participants hit the target sphere or fully pressed the buzzer, they received vibro-tactile feedback in form of a controller vibration for 0.2 s.

In training and adaptation trials, participants received immediate feedback on their performance in form of a color change of the target as well as textual feedback in the same color presented on the display: If the reproduction was in a range of 1 ± 0.15 s, color changed to green and the message "Gut!" (good!) appeared on the display. Following reproductions faster than 0.85 s resulted in a color change to red and the message "Zu schnell" (too fast!) appeared on the display. If reproductions exceeded the target interval by more than 0.15 s, the color to blue and the message "Zu langsam!" (too slow!) appeared on the display.

Figure 6. Experiment 4: Modality dependence of temporal sensorimotor adaptation. A) Participants had to reproduce the interval marked by the ready- and set-signal by either reaching a target (pointing to a sphere or buzzer, left columns) or clicking a button on the controller (clicking task, right columns) in time for the go-signal. For the clicking task, the ready-set signal was auditory, and the sound came either from the direction of the sphere used in training trials, or was not localized at all (i.e., ambient). Note that during test trials participants did not see the glove representing their hand. *B)* Temporal outline and trial order. *C)* Averaged reproductions in test trials of each condition. Error bars represent 95% within-subject CIs (19, 20).

In trainings trials and during the first half of a session, participants received veridical feedback about their performance. In the second half, participants received perturbated feedback about their performance, indicating that the participants were too slow. The manipulation was achieved by multiplying the actual reproduction time with a specific factor before displaying it as feedback. After the first half of the experiment the factor was increased stepwise over the course of five trials from 1.06 to finally 1.30, artificially indicating that participants are slower than they are. In test trials, no feedback about the performance was given and the glove was not visible.

Clicking task. For the clicking task, we used the same ready-set-go paradigm as in the pointing reproduction task (see Figure 6A). However, instead of a color change, participants heard two tones (sine wave; duration: 0.1 s; frequency: 880 Hz) indicating the *ready* and *set* signal. The task for participants was to indicate the *go* signal via button press of the thumbpad. The clicking task was tested in two separate sessions. In one session, the tone was presented as an ambient sound, without a specific direction. In the other clicking session, the tones were spatialized, such that the tone came from the black sphere. At trial start, the target sphere of the pointing task above the table and both tones were emitted. To achieve spatialized sound, Unreal's build-in spatialization plugin was used. Furthermore, the tone was played through the left speaker only, which roughly matched the location of the target.

Procedure. Figure 6B depicts the temporal outline of the experiment. Each participant completed four sessions, separated by at least 30 minutes. In all sessions, participants were adapted using the pointing reproduction task. In contrast to the previous experiments, here we used an n-1 adaptation procedure, that is, a test trial followed immediately after an adaptation trial (i.e., no block of adaptation trials was presented).

In one session, test trials consisted of the same pointing reproduction task as used for the adaptation (condition *visual sphere*). In another session we tested if adaptation effects transfer to different objects (condition *visual buzzer*). Instead of pointing at the sphere, participants had to press down a buzzer in the test trials.

To examine if adaptation effects transfer to another sensory modality, participants had to reproduce an interval indicated by auditory instead of visual stimuli in two of four sessions. In one auditory session, the tones were spatialized, as if they were coming from the target sphere (condition *auditory sphere*). In the other auditory condition, an ambient sound was presented (condition *auditory ambient*).

Participants started with 20 trainings trials of the pointing reproduction task. Session including the buzzer or clicking conditions, contained 10 additional trials of the respective task. Following training, participants completed 40 test trials with veridical feedback, each interleaved with a pointing adaptation trial. Afterwards, we introduced the feedback perturbation, stepwise over the course of five trials, followed by 39 additional test trials with perturbated feedback. At the end of the experiment, participants completed 10 more training-trials of the pointing task to de-adapt. The session *visual sphere* contained 196 trials in total. During the other sessions participants completed ten additional trainings trials of the respective test-condition, resulting in 216 trials in total.

Statistical analyses. For the analysis of reproduction performance in test trials, depicted in Figure 6C, we constructed LMMs to predict reproductions. Models included the predictors adaptation (coded as 0 for all no-adaptation trials and as 1 for all adaptation trials), condition (*visual sphere, visual buzzer, auditory sphere, and auditory ambient*) and the feedback of the previous (adaptation) trial as a measure of adaptation transfer (n-1 feedback, coded as -1 if the feedback was "too slow", 1 if the feedback was "too fast", and 0 if the feedback was "good"). In all models, intercepts varied by participant. Models with different combinations of these predictors, with or without interactions between main effects, were again, if applicable, compared by means of likelihood ratio tests, BIC values and Bayes Factors. We report the resulting best model and statistical evidence for or against effects. Post-hoc comparisons were calculated using the emmeans package in R (Searle, Speed, & Milliken, 1980) with Tukey-adjusted *p*-values.

In condition *visual sphere* 8.09% of all trials had to be excluded from the analysis, 13.94% in condition *visual buzzer*, 8.77% in condition *auditory sphere*, and 8.32% in condition *auditory ambient* (see Experiment 1 for exclusion criteria).

Results

Reproduction performance. The final model to predict reproductions included the factor adaptation, condition, their interaction ($X^2(3) = 241.4$, $p < .001$, $\Delta BIC = 215.97$, $BF_{10} > 1000$), the feedback of the previous (adaptation) trial and its interaction with condition ($X^2(3) = 151.19$, $p < 001$, $\Delta BIC = 125.75$, $BF_{10} > 1000$). Post-hoc comparisons showed that during the no-adaptation half of the experiment, reproductions did not differ from each other ($ps > .1$), while in the adaptation half of the experiment auditory conditions differed from visual conditions ($ps < .001$), but visual and auditory conditions did not differ from each other ($ps > .2$). In all conditions reproductions during the adaptation phase were lower than during the no-adaptation phase ($ps < .05$).

Discussion

These results suggest that adaptation affected the visual pointing tasks much stronger than the auditory clicking tasks. Since we already established that spatiotemporal adaptation does transfer from pointing to clicking movements in Experiment 1, the lack of substantial adaptation effects in auditory clicking tasks must be caused by the modality manipulation. Further post hoc tests to decompose the interaction between condition and feedback presented to participants in the previous trial revealed that the effect of n-1 feedback differed between auditory and visual conditions (ps < .001), and between condition *visual sphere* and *visual buzzer* (p = .04). This shows that the more similar the test-task was to adaptation trials (auditory-visual vs. visual-visual) and the more similar the target object (buzzer-sphere vs. sphere-sphere), the more participants incorporated the feedback received in the previous trial, or in other words, the more transfer from adaptation to test trials was observable.

General Discussion

In a set of four experiments we explored whether temporal estimates are obtained through perceptual or motor processes. To this end, we tested whether adaptation to a temporal perturbation in a motor reproduction task transfers to movements other than the adapted movement, to contexts other than the adapted context, and to modalities other than the adapted modality. Participants had to adjust their pointing movement required to reproduce an interval to cope with slowed down hand movements in a virtual environment. Over the course of this adaptation phase participants incorporated performance feedback and gradually varied movement onset (i.e., started the movement earlier) and speeded up their movements (i.e., decreased movement duration). This adaptation affected subsequent pointing reproductions, causing systematic under-reproductions once the temporal perturbation was removed. Looking at trials without direct performance feedback, we tested whether adaptation transfers when varying the motor reproduction task and when varying the sensory stimuli that marked the temporal interval in appearance or modality.

We found that adaptation transfers from continuous arm reaching movements to ballistic finger movements. Continuous and ballistic movements differ drastically in the requirements of their control. Continuous movements are monitored online to regulate their speed and duration in order to steer the hand to the desired goal location. For ballistic movements, like clicking, only the onset can be variably controlled whereas the movement itself, once initiated, underlies an automatic routine. Since timing adaptation affected the planning of movements with such different dynamics,

it seems as if the motor system as a whole (or for the right arm) adapted to the temporal perturbation. We also found that adaptation effects transferred to a pointing movement in the opposite direction, ruling out that adaptation is specific to location. Typically, transfer of adaptation is highly specific to the particular movement (i.e., movement type and direction, context, effector, or task) that is trained in the adaptation trials. This is true for adaptation to spatial (18) and spatio-temporal distortions (30) (e.g., intercepting moving targets with delayed movement feedback), as well as for motor-aftereffects on temporal judgements (e.g., the speed of finger tapping influences perceived duration in a subsequent visual interval discrimination task: 3, 31–34). In the current study, temporal sensorimotor adaptation effects were not spatially selective, and adaptation effects persisted regardless of the type of movement (continuous pointing or ballistic clicking). Adaptation effects even persisted outside of the VE, ruling out that participants altered their behavior because of the motion-to-photon latency of the VR system and that adaptation effects were object-based (i.e., tied to the target in the VE). In other words, the locus of adaptation effects was neither (fully) extrinsic, nor (fully) object-centered. Instead, observed effects of adaptation were intrinsic and affected temporal predictions and temporal planning of all goal directed motor actions (i.e., motor actions performed to reproduce an interval) tested in this study.

Temporal adaptation did not transfer across sensory modalities of the stimuli that marked the temporal interval. Here, adaptation was induced with visual stimuli. When subjects were asked to reproduce intervals marked by auditory stimuli adaptation was significantly weaker than for visually defined intervals. Stimulus changes within the visual modality did not affect adaptation. Movements executed toward a sphere or a buzzer yielded comparable adaptation strengths, as did movements executed towards a sphere in VR or to a LED in the real world.

These motor transfer results suggest that temporal adaptation in the current experiments occurred upstream of motor processing. If adaptation takes place before temporal estimates are used for motor reproduction, all movements should be modified as was found in the present study. The absence of transfer for movements executed to auditory stimuli is in agreement with this interpretation. If adaptation affects the visual processing of the temporal interval, aftereffects should be restricted to visually marked intervals. Adaptation of visual time would explain all transfer findings. This interpretation is also consistent with previous studies that found changes in purely visual time estimation tasks after motor adaptation (3).

How does motor information adapt visual time? In a recent review, De Kock et al. (35) discuss two general models how movements might shape the perception of time. In the first, which

they call *Feedforward Enhancement*, temporal estimates are generated within motor areas directly. The second, termed *Active Sensing*, assumes that motor signals influence processing in sensory regions. Similar to the Active Sensing framework, we recently showed that post-saccadic errors calibrate visual localization, that is, motor induced errors affect visual perception (36). In the current study, motor errors induced temporal adaptation and - as argued above - transfer findings showed that adaptation most likely resulted from processing in visual areas. Our data extends previous findings (36): Motor errors not only recalibrate our perception of space, but also of time, allowing us to smoothly interact with our dynamic environment.

In conclusion, generalized adaptation transfer between different movement types and specificity of adaptation to the visual modality suggest that motor errors induce adaptation of visual temporal processing.

Grants

This research was supported by the European Research Council (project moreSense, grant agreement 757184) and by the Deutsche Forschungsgemeinschaft (DFG, ZI 1456).

Disclosure

The authors declare no competing interests.

Additional information

An earlier version of this manuscript was published as a bioRxiv-preprint, available at https://doi.org/10.1101/2021.03.31.437803. All data and analysis scripts are available online at https://osf.io/zbgy9/.

References

1.   **Remington ED**, **Parks TV**, **Jazayeri M**. Late Bayesian inference in mental transformations. *Nat Commun* 9: 4419, 2018. doi: 10.1038/s41467-018-06726-9.

2.   **Jazayeri M**, **Shadlen MN**. A neural mechanism for sensing and reproducing a time interval. *Current Biology* 25: 2599–2609, 2015. doi: 10.1016/j.cub.2015.08.038.

3.   **Anobile G**, **Domenici N**, **Togoli I**, **Burr D**, **Arrighi R**. Distortions of visual time induced by motor adaptation. *Journal of Experimental Psychology: General* 149: 1333–1343, 2020. doi: 10.1037/xge0000709.

4.   **Tomassini A**, **Vercillo T**, **Torricelli F**, **Morrone MC**. Rhythmic motor behaviour influences perception of visual time. *Proc R Soc B* 285: 20181597, 2018. doi: 10.1098/rspb.2018.1597.

5.   **Yokosaka T**, **Kuroki S**, **Nishida S**, **Watanabe J**. Apparent time interval of visual stimuli is compressed during fast hand movement. *PLoS ONE* 10: e0124901, 2015. doi: 10.1371/journal.pone.0124901.

6.   **Yon D**, **Edey R**, **Ivry RB**, **Press C**. Time on your hands: Perceived duration of sensory events is biased toward concurrent actions. *Journal of Experimental Psychology: General* 146: 182–193, 2017. doi: 10.1037/xge0000254.

7.   **Ueda S**, **Shimoda S**. Subjective time compression induced by continuous action. *Sci Rep* 11: 13488, 2021. doi: 10.1038/s41598-021-92946-x.

8.   **Hagura N**, **Kanai R**, **Orgs G**, **Haggard P**. Ready steady slow: Action preparation slows the subjective passage of time. *Proc R Soc B* 279: 4399–4406, 2012. doi: 10.1098/rspb.2012.1339.

9.   **Van Rijn H**. Accounting for memory mechanisms in interval timing: A review. *Current Opinion in Behavioral Sciences* 8: 245–249, 2016. doi: 10.1016/j.cobeha.2016.02.016.

10.   **Sohn H**, **Narain D**, **Meirhaeghe N**, **Jazayeri M**. Bayesian computation through cortical latent dynamics. *Neuron* 103: 934-947.e5, 2019. doi: 10.1016/j.neuron.2019.06.012.

11.   **Damsma A**, **Schlichting N**, **Van Rijn H**. Temporal context actively shapes EEG signatures of time perception. *Journal of Neuroscience* 41: 4514–4523, 2021. doi: 10.1523/JNEUROSCI.0628-20.2021.

12.   **Zimmermann E**, **Cicchini GM**. Temporal context affects interval timing at the perceptual level. *Sci Rep* 10: 8767, 2020. doi: 10.1038/s41598-020-65609-6.

13.   **De Kock R**, **Zhou W**, **Joiner WM**, **Wiener M**. Slowing the body slows down time perception. *eLife* 10: e63607, 2021. doi: 10.7554/eLife.63607.

14.   **Niehorster DC**, **Li L**, **Lappe M**. The accuracy and precision of position and orientation tracking in the HTC Vive Virtual Reality system for scientific research. *i-Perception* 8: 204166951770820, 2017. doi: 10.1177/2041669517708205.

15. **Verdelet G**, **Salemme R**, **Desoche C**, **Volland F**, **Farne A**, **Coudert A**, **Hermann R**, **Truy E**, **Gaveau V**, **Pavani F**. Assessing spatial and temporal reliability of the Vive System as a tool for naturalistic behavioural research. In: *2019 International Conference on 3D Immersion (IC3D)*. 2019 International Conference on 3D Immersion (IC3D). IEEE, p. 1–8.

16. **Yee N**, **Bailenson JN**, **Ducheneaut N**. The Proteus Effect: Implications of Transformed Digital Self-Representation on Online and Offline Behavior. *Communication Research* 36: 285–312, 2009. doi: 10.1177/0093650208330254.

17. **Yee N**, **Bailenson J**. The Proteus Effect: The Effect of Transformed Self-Representation on Behavior. *Human Comm Res* 33: 271–290, 2007. doi: 10.1111/j.1468-2958.2007.00299.x.

18. **Krakauer JW**, **Hadjiosif AM**, **Xu J**, **Wong AL**, **Haith AM**. Motor learning. *Comprehensive Physiology* 9: 613–663, 2019. doi: 10.1002/cphy.c170043.

19. **Cousineau D**. Varieties of confidence intervals. *ACP* 13: 140–155, 2017. doi: 10.5709/acp-0214-z.

20. **Morey RD**. Confidence intervals from normalized data: A correction to Cousineau (2005). *TQMP* 4: 61–64, 2008. doi: 10.20982/tqmp.04.2.p061.

21. **Bates D**, **Mächler M**, **Bolker B**, **Walker S**. Fitting Linear Mixed-Effects Models using lme4. *J Stat Soft* 67, 2015. doi: 10.18637/jss.v067.i01.

22. **Kuznetsova A**, **Brockhoff PB**, **Christensen RHB**. lmerTest Package: Tests in Linear Mixed Effects Models. *J Stat Soft* 82, 2017. doi: 10.18637/jss.v082.i13.

23. **R Core Team**. R: A language environment for statistical computing [Online]. R Foundation for Statistical Computing. https://www.R-project.org/.

24. **Morey RD**, **Rouder JN**. BayesFactor: Computation of Bayes factors for common designs [Online]. https://cran.r-project.org/web/packages/BayesFactor/index.html.

25. **Maaß SC**, **Van Rijn H**. 1-s productions: A validation of an efficient measure of clock variability. *Front Hum Neurosci* 12: 519, 2018. doi: 10.3389/fnhum.2018.00519.

26. **Leys C**, **Ley C**, **Klein O**, **Bernard P**, **Licata L**. Detecting outliers: Do not use standard deviation around the mean, use absolute deviation around the median. *Journal of Experimental Social Psychology* 49: 764–766, 2013. doi: 10.1016/j.jesp.2013.03.013.

27. **Grondin S**. About the (non)scalar property for time perception. In: *Neurobiology of Interval Timing*, edited by Merchant H, de Lafuente V. Springer, p. 17–32.

28. **EpicGames**. New Archviz Interior Rendering sample project now available! [Online]. *Unreal Engine*: 2019. https://www.unrealengine.com/en-US/blog/new-archviz-interior-rendering-sample-project-now-available [12 Aug. 2021].

29. **Searle SR**, **Speed FM**, **Milliken GA**. Population marginal means in the linear model: An alternative to least squares means. *The American Statistician* 34: 216–221, 1980. doi: 10.1080/00031305.1980.10483031.

30. **de la Malla C**, **Lopez-Moliner J**, **Brenner E**. Dealing with delays does not transfer across sensorimotor tasks. *Journal of Vision* 14: 8–8, 2014. doi: 10.1167/14.12.8.

31. **Burr D**, **Tozzi A**, **Morrone MC**. Neural mechanisms for timing visual events are spatially selective in real-world coordinates. *Nat Neurosci* 10: 423–425, 2007. doi: 10.1038/nn1874.

32. **Fornaciai M**, **Arrighi R**, **Burr DC**. Adaptation-induced compression of event time occurs only for translational motion. *Sci Rep* 6: 23341, 2016. doi: 10.1038/srep23341.

33. **Frassinetti F**, **Magnani B**, **Oliveri M**. Prismatic Lenses Shift Time Perception. *Psychol Sci* 20: 949–954, 2009. doi: 10.1111/j.1467-9280.2009.02390.x.

34. **Johnston A**, **Arnold DH**, **Nishida S**. Spatially localized distortions of event time. *Current Biology* 16: 472–479, 2006. doi: 10.1016/j.cub.2006.01.032.

35. **De Kock R**, **Gladhill KA**, **Ali MN**, **Joiner WM**, **Wiener M**. How movements shape the perception of time. *Trends in Cognitive Sciences* 25: 950–963, 2021. doi: 10.1016/j.tics.2021.08.002.

36. **Cont C**, **Zimmermann E**. The motor representation of sensory experience. *Current Biology* 31: 1–8, 2020. doi: 10.1016/j.cub.2020.11.032.