Title: A heterochromatic knob reducing the flowering time in maize

Authors: Renata Flávia Carvalho[1], Margarida Lopes Rodrigues Aguiar-Perecin[1],

Wellington Ronildo Clarindo[2], Roberto Fristche-Neto[1,3], Mateus Mondin[1*]

Affiliations:

[1]University of São Paulo, "Luiz de Queiroz" College of Agriculture - ESALQ,

Piracicaba, SP 13418-900, Brazil.

[2]Department of General Biology, Federal University of Viçosa, Viçosa, MG 36570-900,

Brazil.

[3]International Rice Research Institute (IRRI) - Breeding Analytics and Data

Management Unit, Laguna 4031, Philippines

*Correspondence to: mmondin@usp.br

Abstract: Maize flowering time is an important agronomic trait, which is associated

with variations in the genome size and heterochromatic knobs content. We integrated

three steps to show this association. Firstly, we selected inbred lines varying for

heterochromatic knob composition at specific sites in the homozygous state. Then, we

produced heterozygous hybrids for knobs, which allow us to carry out genetic mapping.

Second, we measured the genome size and flowering time for all materials. Knob

composition did not affect the genome size. Finally, we developed an association study

and identified a knob marker on chromosome 9 showing the strongest association with

flowering time. Indeed, modeling allele substitution and dominance effects could offer

only one heterochromatic knob locus that could affect flowering time, making it earlier

rather than the knob composition.

One Sentence Summary: Flowering time in maize is associated with specific knob

position rather than genome size and knob composition.

Main Text: The relationship between heterochromatic knobs, genome size, and

flowering time (FT) in maize is a long debate. Knobs have been intriguing geneticists

for more than 100 years, and since their discovery, their functions remain under

investigation. These maize genome regions are constitutive heterochromatins with late

replication during the cell cycle, extensively composed of two highly repetitive satellite

DNA families, K180 and TR-1 (*1*, *2*). Due to their monotonous DNA sequence

organization, no direct function has been attributed to the knobs. However, the first

eminent observation was their wide variability among landraces, inbred lines, and

hybrids, following a biogeographic distribution, suggesting a possible role affecting the

expression of some phenotypic features FT (*3–5*).

Moreover, other exciting aspects of the heterochromatic knobs in the maize genome

have been elucidated by several studies. Knobs have been shown to affect local

recombination (*6*, *7*) and genes adjacent to these regions reduce their expression level

(*8*). The meiotic drive mechanism influences the number of knobs in different species of

Zea (*9–11*). Such a system would favour knob transmissions preferentially during

female meiosis by a neocentromere activity, which might have contributed to maize

genome remodelling throughout its evolutionary history. Response to abiotic stress was

observed in maize plants from the transcriptional activation of the knob repetitive

sequences, being this activation selective, temporary, and accompanied by epigenetic

changes (*12*). Lastly, associations between heterochromatic knobs and agronomic traits

have also been reported (*13*).

There is growing interest in the role of variation of genomic content in creating

phenotypic modifications within a species. These changes are due to copy number

variation (CNV), which has been used to describe duplications, deletions, and insertions

between individuals of a species; and presence/absence variation (PAV) that describes

the presence or not of sequences on the genome of different individuals of the same

species. Together, they form the pan-genome of species (*14*). Maize is also a model

species for studies of pan-genome. Differences in chromosomal structure between

maize landraces were firstly identified through cytogenetic studies. Barbara McClintock

and colleagues analysed the content and size of heterochromatic knobs to characterize

this variation in the genome (*4*).

Moreover, quantification of DNA content through flow cytometry has

documented significant variability in the genome size (GS) between landraces and

inbred lines (*15*, *16*). Currently, modern cytogenomic techniques have sampled the wide

variation in the copy numbers resulting from repetitive sequences, which make up most

of the maize genome (*17–19*). New combinations of alleles arise from the variation of

genomic content within species, contributing to phenotypic variation (*20*). Therefore,

this can have an influence on several important characteristics, including flowering time

(*17*). Flowering time is a quantitative trait of extreme relevance for cultivated plants

since it controls plant adaptation to the environment. Breeding programs through this

feature outline strategies to make earlier or later varieties, allowing their expansion to

other regions and consequently increasing yield (*21*).

Recent studies about maize genome present the triangle formed between genome

size, knob content, and flowering time in the natural populations. These surveys have

considered just the knob numbers or estimated the knob abundances by low coverage

sequencing (*16*, *17*, *22*). However, few studies have evaluated the knob effect

depending on its homozygous or heterozygous condition (*23*). Different researchers

have attributed variation in maize genome size to heterochromatin content differences,

especially in knobs (*16*, *22–24*). Moreover, it is proposed that there is a relationship

between decreased genome sizes in high altitudes with reduced flowering time (*17*, *25*).

Besides, the same association is observed for knobs abundance occurring along

altitudinal clines. However, no experimental design has isolated the effects of

homozygous or heterozygous knobs and test whether they are correlated genetically. In

this case, the relationship between knobs and flowering time remains unclear. Here, we

derived a panel of inbred lines and their hybrids to verify the association between

genome size, knob constitution, and flowering time for male (MF) and female (FF)

inflorescence. Firstly, in chronological order, inbred lines with specific knob

constitutions for chromosomes K3L, K5L, K7S, and K9S were derived until S9

generation from a segregating single seed of a flint variety (*26*). Then we created the

hybrids between inbred lines, and finally, we started to estimate the flowering time.

These near-isogenic inbred lines and their hybrids, varying for knob positions at K3L,

K5L, K7S, and K9S (Fig. 1 and 2 and Data S2), were used to assay flowering time

under controlled environmental conditions and their genome size measured by flow

cytometry. All data were used to analyse the genome association study (for more

information, see Supplementary Materials and Data S1).

Knob composition does not correlate with maize genome size. There is a

wide variability of genome size in natural maize populations, and these differences

often are associated with heterochromatin percentage (*27*). Here, we measured the

genome size for the inbred lines and hybrids to test whether knob composition

contributes to the DNA content variation. In our analyses, genome size was considered

a quantitative trait, and its heritability was estimated at 26%. The hybrids presented a

higher mean genome size than inbred lines, showing 5477 Mbp (2C = 5.6 pg) and 5281

Mbp (2C = 5.4 pg), respectively (Fig. 3). However, no significant differences were

found. Even when the presence and absence of the knob were considered, no differences

were observed (Fig. S1). Each inbred line and hybrid can have different knob

configurations at the chromosomes K3L, K5L, K7S, and K9S. These combinations can

be homozygous for presence or absence in the lines or heterozygous in the hybrids (Fig.

1 and 2). These knob combinations were used to create dosage classes and test their

effects on genome size and flowering time (Fig. 4). Our results show knob dosages are

not enough to explain the genome size increase or decrease in the inbred lines and

hybrids. However, we observed that genome size varies broadly despite the same knob

composition. For instance, the hybrids presenting the highest (441311/2 x 441324/1 –

2C = 6.31 pg) and the smallest genome size (442213/1 x 441311/2 – 2C = 4.72 pg)

share the same knob constitution (Data S1). For inbred lines, the largest genome size

(131311/1-04 - 2C = 5.69 pg) has a lesser number of knobs than the shortest genome

(442213/1 - 2C = 4.65 pg), where we have homozygous knob presence for all the

chromosomes (Data S1). Therefore, no pattern was observed, which indicates that there

is probably no linear relationship between the number of knobs present in the maize

genome with the increase or decrease in the DNA content in our study. The genome size

associated with the heterochromatin percentage and knob numbers with flowering time

for the maize genome is given in some studies. Most of these studies have shown

positive correlations between genome size and knob content (*5*, *15*, *22*, *28*). A recent

study provided evidence that natural selection plays a substantial role in reducing the

maize genome size at high altitudes (*17*). The authors also showed that the abundance of

transposable elements (TEs) and heterochromatic knobs are significantly correlated with

altitude, and the knobs act as significant effect loci in the genome size. Unlike what they

found, our data does not show this same relationship. We performed an adapted

genome-wide association study (GWAS) with the heterochromatic knob full panel to

identify associations with genome size and flowering time, where the knobs were used

as our genetic markers (see Supplementary Materials). Our null hypothesis was that

there were significant differences in genome size and days for male and female

flowering due to the knob conditions in the K3L, K5L, K7S, and K9S positions. To test

our hypothesis, we used the inbred lines and hybrids panel, to which all knobs were

mapped (Fig. 2, Data S1). The panel has a matrix-*like* structure with different knob

combinations for presence (++ or 1) or absence (00 or -1) when homozygous and as

heterozygous when just one of the homologous has a knob (+0 or 0). Firstly, only the

allele substitution effect model was performed, and no significant association was found

between knobs and FT (Fig. S2). Moreover, this information was supported by a null

correlation between genome size and knob dosage classes (Table 1 and Fig. S3). Our

results also corroborate a survey showing no significant correlation between maize

population genome size and heterochromatin percentage. Even though larger genomes

exhibited the highest heterochromatin content (*16*); hence, all these observations lead us

to infer that maize genome size is independent of knob composition and might be

controlled genetically. Another hypothesis is that as our materials differ in the

composition of only four knob positions, genome size measurements by flow cytometry

would not detect slight differences in genome size due to knobs. Furthermore, we might

infer differences in the genome size among our inbred lines and hybrids result from

transposable element (TE) variability once the loss of TE during selfing has been

demonstrated (*29*).

Genome size does not affect maize flowering time. Negative values for

Pearson's correlation were found comparing genome size with female and male

flowering time (Table 1). Our results show a negative correlation between genome size

and flowering time in maize for the first time. Previously, a positive relationship

between genome size and flowering time has been reported (*5*, *28*, *30*). Also, significant

positive correlations between genome size, knobs abundance, and flowering time were

found in maize inbred lines (*22*). However, these correlations were lost when a kinship

matrix was introduced in the analyses. In this study, the authors performed an

association study where genome size was also correlated with flowering time. Three

genomic regions associated with genome size were found, mapped close to the knob

region on chromosome 8 (*22*). In contrast, another study analysing maize landraces in

northern Argentina found no correlation between genome size and flowering time (*16*).

Nevertheless, a positive relationship between the vegetative cycle and heterochromatin

percentage was observed (*16*).

Relationship between flowering time and knob constitution. We evaluated

days to MF and FF individually (Data S1) and estimated heritability to be 51% for MF

and 41% for FF. The mean values for days to MF and FF hybrids were 63 and 64 days,

respectively, while the flowering for inbred lines was 70 days to MF and 72 days to FF

(Fig. 3). Heterochromatic knob configurations as heterozygotes exhibited shorter

flowering times than those in homozygous states (Fig S1). The flowering time data also

were plotted, showing its amplitude inside the inbred line families and hybrids (Fig. 4).

We observed synchronicity for both traits within each group analysed, and hybrids had

shorter flowering time than inbred line families.

Further, it is interesting to notice that the flowering time amplitude of hybrids is

much smaller than the other groups, ranging mainly between 60 and 70 days, in

agreement with the observed mean values. A broader variation in flowering time is

observed for inbred line families (58 - 114 days), with distribution densities mainly

varying between 60 and 80 days. The JD 4-4 family had a smaller flowering time

distribution for FF and FM among families. The knob dosage class's analyses cannot

explain the flowering behaviour of all maize materials. It was possible to observe that

the mean values of dosage classes were very similar for MF and FF (Fig. 5). It was also

interesting to note that the inbred lines with only two knobs (dosage class = 2) flowered

later than those with other dosage classes (dosage class = 3-8). A previous study argued

that repetitive sequences would have indirect effects on flowering time due to their

impact on genome size and might depend on the environment (*17*). However, this effect

was not found in this study since there was no relationship between genome size and

knob composition (Table 1 and Fig S2).

The genetic architecture of flowering time in maize has been widely studied,

given that this trait reflects the plant adaptation to the environment. Since maize is

distributed throughout America and is adapted to a wide range of environments,

understanding how flowering time is regulated is of paramount importance and

generates valuable information for breeding programs (*31*). Besides, these studies, in

the last two decades, dissected the maize flowering time using different approaches such

as linkage and association mapping (*32–35*), population genetics (*36, 37*),

archaeological DNA studies (*38*), genome-wide association studies (*22, 39–41*), and

gene analysis (*42–45*). In inbred maize lines, the genetic architecture of flowering time

has been attributed to the cumulative effect of numerous quantitative trait loci (QTL),

each with a small impact on this trait (*31*). In the research described here, we used two

GWAS models to illustrate the interactions between flowering time and knob

conditions. The first model took into account only the allele substitution effect of the

markers (Fig. 6A, 6B, and S4), and the second considered the dominance effect (Fig.

6C, 6D, and S5). Our analysis showed only one significant marker-trait association for

both allele substitution and dominance effect models regarding flowering time. This

significant association was observed just for the knob on the short arm of chromosome

9 (K9S) (Fig. 6). Regardless of the GWAS model, the knob marker on chromosome 9

displayed the same performance concerning the flowering time, showing the *p*-value

highly significant (Table 2 and Table S1).

Regarding the allele substitution effect model, the presence of the K9S

decreased the MF and FF of the inbred lines and hybrids by -0.45 and -0.51 days,

respectively. While on the dominance effect model, this effect was even more

significant, decreasing flowering time in one and a half days (MF = -1.40 and FF = -

1.53) (Fig. 7). Only the knob on chromosome 9 significantly reduced the flowering time

in the inbred lines and hybrids. Decreasing in flowering time is more affected by the

presence of knobs in the homozygous or heterozygous configuration. Our analysis also

showed slight effects for each of the knobs on flowering time, albeit these were not

significant (Table S1). These results showed that the knob composition was not

essential to model the genotype effects on the trait. The heritability of the markers for

both models varied from 0.02 to 0.15 (Table 2). The flowering time violin plot also

supported the GWAS analyses (Fig. 5). Since all hybrids are homozygous or

heterozygous for the K9S, corroborating the specific contribution of this knob to

decrease maize flowering time.

Especially in maize, the literature describes the hypothesis of a putative

contribution of the heterochromatic knobs to the flowering time variation (*23*).

Previously, studies had already indicated that gene groups responsible for the plant

morphological and physiological traits would be correlated with the presence of knobs

(*11, 13*). According to the authors, once close to these knobs, recombination in adjacent

regions was suppressed, influencing such traits. More recently, through fluorescent *in*

situ hybridization (FISH), it was demonstrated that knobs are located in areas dense in genes where large knobs can reduce recombination locally (*6*). Comparisons between European maize genomes and US Corn Belt revealed variation in their repetitive sequences and gene content. The germplasms were separated by the intensity and position of knob regions. However, additional sites with small arrangements of knobs conserved in flint and dent lines were observed. This study also showed that the knob sequences could affect genes surrounded by them, decreasing their expression level (*8*). Besides, the functional role of the knobs has also been traced by the relationship between their abundance, the genome size, and the flowering time, as shown above (*17*, *22*).

Flowering time-related QTLs have been found across the maize genome (*31*, *35*, *38*). In chromosome 9, flowering time-related QTLs were found, and some genes were identified (*40*, *46*). The main gene located in chromosome 9 was ZmCCT9 (photoperiod sensitive) and mapped at the long arm, opposite the knob position. Other candidate genes were found along the chromosome, but the precise location was not defined yet (*33*, *47*). In our study, the heterochromatic knob mapped at the short arm of chromosome 9 is strongly associated with flowering time. Constitutive heterochromatins have well-defined roles in the genomes, such as centromeres and telomeres (*48*, *49*). That leads us to infer that heterochromatic knobs may also have a role in the maize genome, affecting certain phenotypic traits. However, further investigation in these regions will be needed to find possible candidate genes associated with knobs due to the inhibition of crossing over on their neighbourhood.

Concluding remarks. Despite more than 100 years of studies on maize genetics, the discussion on the role of knobs remains a current issue and has shown that this region is an essential fraction of the maize genome. Elimination dynamics of

components of the maize genome over successive self-fertilizations were reported (*29*).

Among the five components analysed in the genome (gene content, ribosomal DNA, B

chromosomes, TEs, and knobs), TEs were the most significantly lost components. The

results provide insights into the constitutive role played by knobs in the maize genome.

Since, like TEs, knobs are repetitive sequences, which make up about 10% of the maize

genome and, for unknown reasons, are not eliminated from it after generations of self-

fertilization. That strengthens the hypothesis that heterochromatic knobs may have a

functional role within the maize genome, even composed almost exclusively by

repetitive sequences.

Throughout the maize genome evolution, it was proposed that flowering time

was a trait influenced by changes in the genome size. At high altitudes, maize flowering

time was shorter than at low altitudes, followed by a smaller genome (*5*, *16*, *25*). At the

same time, most studies have also indicated a positive correlation between the genome

size and the knob abundance and that the heterochromatin content decreases along the

altitude clines. We found no significant association between knobs and genome size

from our data. The knob present in the short arm of chromosome 9 is associated with

reducing the flowering time when homozygous and heterozygous. Our results suggest a

role of the knobs in the flowering time different from those previously described.

Unravel the mystery behind the repetitive sequences has been hard work all over

the years. It becomes even more difficult for maize since its genome comprises more

than 85% of these sequences. For the first time, maize inbred lines were selected to have

knobs in specific locations, and their hybrids developed to carry heterozygous and

homozygous knobs. After that, the analyses of adapted GWAS were carried out,

showing the contribution of a single locus to the early maize flowering time. The maize

flowering time is a complex trait, and several studies have provided insights into its

genetic architecture. To this complexity, we add our data that suggest that gene-*like* components, such as specific heterochromatic knobs, are capable of affecting the maize flowering time, probably by interaction with particular genes playing a central role in the regulation of the feature expression. Furthermore, as knobs suppress local recombination (*6*), some genes would be associated with them.

References and Notes:

1. W. J. Peacock, E. S. Dennis, M. M. Rhoades, a J. Pryor, Highly repeated DNA sequence limited to knob heterochromatin in maize. *Proc. Natl. Acad. Sci. U. S. A.* **78**, 4490–4 (1981).

2. E. V. Ananiev, R. L. Phillips, H. W. Rines, A knob-associated tandem repeat in maize capable of forming fold-back DNA segments: are chromosome knobs megatransposons? *Proc. Natl. Acad. Sci. U. S. A.* **95**, 10785–90 (1998).

3. Y. Jian, C. Xu, Z. Guo, S. Wang, Y. Xu, C. Zou, Maize (Zea mays L.) genome size indicated by 180-bp knob abundance is associated with flowering time. *Sci. Rep.* **7**, 1–9 (2017).

4. B. McClintock, A. K. Takeo, A. Blumenschein, *Chromosome constitution of races of maize. Its significance in the interpretation of relationships between races and varieties in the Americas* (Colegio de Postgraduados, Chapingo, 1981).

5. A. L. Rayburn, H. J. Price, J. D. Smith, J. R. Gold, C-Band Heterochromatin and DNA Content in Zea mays. *Am. J. Bot.* **72**, 1610–1617 (1985).

6. R. Ghaffari, E. K. S. Cannon, L. B. Kanizay, C. J. Lawrence, R. K. Dawe, Maize chromosomal knobs are located in gene-dense areas and suppress local recombination. *Chromosoma.* **122**, 67–75 (2013).

7. S. M. Stack, L. A. Shearer, L. Lohmiller, L. K. Anderson, Meiotic crossing over

in maize knob heterochromatin. *Genetics*. **205**, 1101–1112 (2017).

8.   G. Haberer, N. Kamal, E. Bauer, H. Gundlach, I. Fischer, M. A. Seidel, M.

Spannagl, C. Marcon, A. Ruban, C. Urbany, A. Nemri, F. Hochholdinger, M.

Ouzunova, A. Houben, C.-C. Schön, K. F. X. Mayer, European maize genomes

highlight intraspecies variation in repeat and gene content. *Nat. Genet.* **52**, 950–

957 (2020).

9.   E. S. Buckler, T. L. Phelps-Durr, C. S. Keith Buckler, R. K. Dawe, J. F. Doebley,

T. P. Holtsford, Meiotic Drive of Chromosomal Knobs Reshaped the Maize

Genome. *Genetics*. **153**, 415–426 (1999).

10.   D. M. Higgins, E. G. Lowry, L. B. Kanizay, P. W. Becraft, D. W. Hall, R. K.

Dawe, Fitness costs and variation in transmission distortion associated with the

abnormal chromosome 10 meiotic drive system in maize. *Genetics*. **208**, 297–305

(2018).

11.   M. M. Rhoades, E. Dempsey, The Effect of Abnormal Chromosome 10 on

Preferential Segregation and Crossing over in Maize. *Genetics*. **53**, 989–1020

(1966).

12.   Y. Hu, L. Zhang, S. He, M. Huang, J. Tan, L. Zhao, S. Yan, H. Li, K. Zhou, Y.

Liang, L. Li, Cold stress selectively unsilences tandem repeats in

heterochromatin associated with accumulation of H3K9ac. *Plant, Cell Environ.*

35, 2130–2142 (2012).

13.   A. Blumenshein, Identification of chromosome segments that contribute to the

specificity of racial characteristics in maize. PhD Thesis, University of North

Carolina (1964).

14.   N. M. Springer, K. Ying, Y. Fu, T. Ji, C. T. Yeh, Y. Jia, W. Wu, T. Richmond, J.

Kitzman, H. Rosenbaum, A. L. Iniguez, W. B. Barbazuk, J. A. Jeddeloh, D.

Nettleton, P. S. Schnable, Maize inbreds exhibit high levels of copy number

variation (CNV) and presence/absence variation (PAV) in genome content. *PLoS*

Genet. **5** (2009), doi:10.1371/journal.pgen.1000734.

15.   D. A. Laurie, M. D. Bennett, Nuclear DNA content in the genera Zea and

Sorghum. Intergeneric, interspecific and intraspecific variation. *Heredity (Edinb).*

55, 307–313 (1985).

16.   M. F. Realini, L. Poggio, J. Cámara-Hernández, G. E. González, Intra-specific

variation in genome size in maize: cytological and phenotypic correlates. *AoB*

Plants. **8**, 1–12 (2015).

17.   P. Bilinski, P. S. Albert, J. J. Berg, J. A. Birchler, M. N. Grote, A. Lorant, J.

Quezada, K. Swarts, J. Yang, J. Ross-Ibarra, Parallel altitudinal clines reveal

trends in adaptive evolution of genome size in Zea mays. *PLoS Genet.* **14**,

e1007162 (2018).

18.   A. Kato, J. C. Lamb, J. a Birchler, Chromosome painting using repetitive DNA

sequences as probes for somatic chromosome identification in maize. *Proc. Natl.*

Acad. Sci. U. S. A. **101**, 13554–13559 (2004).

19.   P. S. Albert, Z. Gao, T. V. Danilova, J. A. Birchler, Diversity of chromosomal

karyotypes in maize and its relatives. *Cytogenet. Genome Res.* **129**, 6–16 (2010).

20.   A. B. Brohammer, T. J. Y. Kono, C. N. Hirsch, *The Maize Genome* (Springer

International Publishing, Cham, 2018; http://dx.doi.org/10.1007/978-3-319-

97427-9_2), *Compendium of Plant Genomes*.

21.   C. Jung, A. E. Müller, Flowering time control and applications in plant breeding.

Trends Plant Sci. **14**, 563–573 (2009).

22. Y. Jian, C. Xu, Z. Guo, S. Wang, Y. Xu, C. Zou, Maize (Zea mays L.) genome

size indicated by 180-bp knob abundance is associated with flowering time. *Sci.*

Rep. **7**, 5954 (2017).

23. S. Chughtai, D. Steffensen, Some insights into the genetics of flowering time.

Maize Genet. Coop. Newsl. **61**, 98–99 (1987).

24. S. R. Chughtai, D. M. Steffensen, J. Crossa, M. Aslam, H. I. Javed, M. Hussain,

in *The Genetics and Exploitation of Heterosis in Crops* (CIMMYT, Mexico), pp.

387–388.

25. L. Poggio, M. Rosato, A. M. Chiavarino, C. A. Naranjo, Genome size and

environmental correlations in maize (Zea mays ssp. mays, Poaceae). *Ann. Bot.*

82A, 107–115 (1998).

26. M. J. U. Decico, Análise da segregação de knobs em progênies F2 e de

retrocruzamento derivadas de uma variedade de milho (1991), pp. 1–147.

27. M. F. Fourastié, A. M. Gottlieb, L. Poggio, G. E. González, Are cytological

parameters of maize landraces (Zea mays ssp. mays) adapted along an altitudinal

cline? *J. Plant Res.* **131**, 285–296 (2018).

28. C. M. Tito, L. Poggio, C. A. Naranjo, Cytogenetic studies in the genus Zea. *TAG*

Theor. Appl. Genet. **83**, 58–64 (1991).

29. K. Roessler, A. Muyle, C. M. Diez, G. R. J. J. Gaut, A. Bousios, M. C. Stitzer, D.

K. Seymour, J. F. Doebley, Q. Liu, B. S. Gaut, The genome-wide dynamics of

purging during selfing in maize. *Nat. Plants*. **5**, 980–990 (2019).

30. A. L. Rayburn, J. W. Dudley, D. P. Biradar, Selection for Early Flowering

Results in Simultaneous Selection for Reduced Nuclear DNA Content in Maize.

Plant Breed. **112**, 318–322 (1994).

31.   E. S. Buckler, J. B. Holland, P. J. Bradbury, C. B. Acharya, P. J. Brown, C.

Browne, E. Ersoz, S. Flint-Garcia, A. Garcia, J. C. Glaubitz, M. M. Goodman, C.

Harjes, K. Guill, D. E. Kroon, S. Larsson, The Genetic Architecture of Maize.

Science (80-.). **325**, 714–718 (2009).

32.   J. Alberto, R. Navarro, M. Willcox, J. Burgueño, C. Romay, K. Swarts, S.

Trachsel, E. Preciado, A. Terron, H. V. Delgado, V. Vidal, A. Ortega, A.

Espinoza Banda, N. Orlando Gómez Montiel, I. Ortiz-Monasterio, F. San

Vicente, A. Guadarrama Espinoza, G. Atlin, P. Wenzl, S. Hearne, E. S. Buckler,

A study of allelic diversity underlying flowering-time adaptation in maize

landraces. *Nat. Publ. Gr.* **49** (2017), doi:10.1038/ng.3784.

33.   Y. Y. X. Li, C. Li, P. J. Bradbury, X. Liu, F. Lu, C. M. Romay, J. C. Glaubitz, X.

Wu, B. Peng, Y. Shi, Y. Song, D. Zhang, E. S. Buckler, Z. Zhang, Y. Y. X. Li, T.

Wang, Identification of genetic variants associated with maize flowering time

using an extremely large multi-genetic background population. *Plant J.* **86**, 391–

402 (2016).

34.   S. Ducrocq, C. Giauffret, D. Madur, V. Combes, F. Dumas, S. Jouanne, D.

Coubriche, P. Jamin, L. Moreau, A. Charcosset, Fine Mapping and Haplotype

Structure Analysis of a Major Flowering Time Quantitative Trait Locus on Maize

Chromosome 10. *Genetics*. **183**, 1555–1563 (2009).

35.   F. Chardon, B. Virlon, L. Moreau, M. Falque, J. Joets, L. Decousset, A.

Murigneux, A. Charcosset, Genetic architecture of flowering time in maize as

inferred from quantitative trait loci meta-analysis and synteny conservation with

the rice genome. *Genetics*. **168**, 2169–2185 (2004).

36.   L. Guo, X. Wang, M. Zhao, C. Huang, C. Li, D. Li, C. J. Yang, A. M. York, W.

Xue, G. Xu, Y. Liang, Q. Chen, J. F. Doebley, F. Tian, Stepwise cis-Regulatory

Changes in ZCN8 Contribute to Maize Flowering-Time Adaptation. *Curr. Biol.*

28, 3005-3015.e4 (2018).

37.   X. Wang, Y. Xu, S. Zhang, L. Cao, Y. Huang, J. Cheng, G. Wu, S. Tian, C.

Chen, Y. Liu, H. Yu, X. Yang, H. Lan, N. Wang, L. Wang, J. Xu, X. Jiang, Z.

Xie, M. Tan, R. M. Larkin, L.-L. Chen, B.-G. Ma, Y. Ruan, X. Deng, Q. Xu,

Genomic analyses of primitive, wild and cultivated citrus provide insights into

asexual reproduction. *Nat. Genet.* **49**, 765–772 (2017).

38.   C. J. Yang, L. F. Samayoa, P. J. Bradbury, B. A. Olukolu, W. Xue, A. M. York,

M. R. Tuholski, W. Wang, L. L. Daskalska, M. A. Neumeyer, J. de Jesus

Sanchez-Gonzalez, M. C. Romay, J. C. Glaubitz, Q. Sun, E. S. Buckler, J. B.

Holland, J. F. Doebley, The genetic architecture of teosinte catalyzed and

constrained maize domestication. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 5643–5652

(2019).

39.   Y. Liang, Q. Liu, X. Wang, C. Huang, G. Xu, S. Hey, H. Y. Lin, C. Li, D. Xu, L.

Wu, C. Wang, W. Wu, J. Xia, X. Han, S. Lu, J. Lai, W. Song, P. S. Schnable, F.

Tian, ZmMADS69 functions as a flowering activator through the ZmRap2.7-

ZCN8 regulatory module and contributes to maize flowering time adaptation.

New Phytol. **221**, 2335–2347 (2019).

40.   H. Y. Hung, L. M. Shannon, F. Tian, P. J. Bradbury, C. Chen, S. A. Flint-Garcia,

M. D. McMullen, D. Ware, E. S. Buckler, J. F. Doebley, J. B. Holland, ZmCCT

and the genetic basis of day-length adaptation underlying the postdomestication

spread of maize. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 11068 (2012).

41.   Q. Yang, Z. Li, W. Li, L. Ku, C. Wang, J. Ye, K. Li, N. Yang, Y. Li, T. Zhong, J.

Li, Y. Chen, J. Yan, X. Yang, M. Xu, CACTA-like transposable element in
ZmCCT attenuated photoperiod sensitivity and accelerated the postdomestication
spread of maize. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 16969–16974 (2013).

42.   E. Stephenson, S. Estrada, X. Meng, J. Ourada, M. G. Muszynski, J. E. Habben,
O. N. Danilevskaya, Over-expression of the photoperiod response regulator
ZmCCT10 modifies plant architecture, flowering time and inflorescence
morphology in maize. *PLoS One*. **14**, e0203728 (2019).

43.   P. Alter, S. Bircheneder, L.-Z. Zhou, U. Schlüter, M. Gahrtz, U. Sonnewald, T.
Dresselhaus, Flowering Time-Regulated Genes in Maize Include the
Transcription Factor ZmMADS1. *Plant Physiol.* **172**, 389–404 (2016).

44.   M. A. A. Minow, L. M. Ávila, K. Turner, E. Ponzoni, I. Mascheretti, F. M.
Dussault, L. Lukens, V. Rossi, J. Colasanti, Distinct gene networks modulate
floral induction of autonomous maize and photoperiod-dependent teosinte. *J.*
Exp. Bot. **69**, 2937–2952 (2018).

45.   C. M. Lazakis, V. Coneva, J. Colasanti, ZCN8 encodes a potential orthologue of
Arabidopsis FT florigen that integrates both endogenous and photoperiod
flowering signals in maize. *J. Exp. Bot.* **62**, 4833–4842 (2011).

46.   C. Huang, H. Sun, D. Xu, Q. Chen, Y. Liang, X. Wang, G. Xu, J. Tian, C. Wang,
D. Li, L. Wu, X. Yang, W. Jin, J. F. Doebley, F. Tian, ZmCCT9 enhances maize
adaptation to higher latitudes. *Proc. Natl. Acad. Sci. U. S. A.* **115**, E334–E341
(2017).

47.   T. A. Miller, E. H. Muslin, J. E. Dorweiler, A maize CONSTANS-like gene,
conz1, exhibits distinct diurnal expression patterns in varied photoperiods.
Planta. **227**, 1377–1388 (2008).

48.   E. M. C. Tacconi, M. Tarsounas, How homologous recombination maintains

telomere integrity. *Chromosoma*. **124** (2015), pp. 119–130.

49.   S. Salvi, G. Sponza, M. Morgante, D. Tomes, X. Niu, K. A. Fengler, R. Meeley,

E. V Ananiev, S. Svitashev, E. Bruggemann, B. Li, C. F. Hainey, S. Radovic, G.

Zaina, J.-A. Rafalski, S. V Tingey, G.-H. Miao, R. L. Phillips, R. Tuberosa,

Conserved noncoding genomic sequences associated with a flowering-time

quantitative trait locus in maize. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 11376–81

(2007).

50.   M. Mondin, J. a Santos-Serejo, M. R. Bertão, P. Laborda, D. Pizzaia, M. L. R.

Aguiar-Perecin, Karyotype variability in tropical maize sister inbred lines and

hybrids compared with KYS standard line. *Front. Plant Sci.* **5**, 544 (2014).

51.   M. M. Praça-Fontes, C. R. Carvalho, W. R. Clarindo, C. D. Cruz, Revisiting the

DNA C-values of the genome size-standards used in plant flow cytometry to

choose the "best primary standards." *Plant Cell Rep.* **30**, 1183–1191 (2011).

52.   F. Otto, in *Methods in Cell Biology* (1990), vol. 33, pp. 105–110.

53.   J. Dolezel, J. Greilhuber, J. Suda, Estimation of nuclear DNA content in plants

using flow cytometry. *Nat. Protoc.* **2**, 2233–44 (2007).

54.   X. Liu, M. Huang, B. Fan, E. S. Buckler, Z. Zhang, Iterative Usage of Fixed and

Random Effect Models for Powerful and Efficient Genome-Wide Association

Studies. *PLOS Genet.* **12**, 1–24 (2016).

55.   M. L. R. Aguiar-Perecin, J. U. Decico, in *Maize Genetics Cooperation*

Newsletter (1988), vol. 62, p. 100.

56.   Y. A. Tsepilov, S. Y. Shin, N. Soranzo, T. D. Spector, C. Prehn, J. Adamski, G.

Kastenmüller, R. Wang-Sattler, K. Strauch, C. Gieger, Y. S. Aulchenko, J. S.

Ried, Nonadditive effects of genes in human metabolomics. *Genetics*. **200**, 707–

718 (2015).

Acknowledgments Coordenação de Aperfeiçoamento de Pessoal de Nível Superior—

Brasil (CAPES) and Conselho Nacional de Desenvolvimento Científico e Tecnológico

(CNPq) for the financial support. **Funding:** RFC was supported by schoolarship from

Coordenação de Aperfeiçoamento de Pessoal de Nível Superior—Brasil (CAPES) and

Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq); MM is PET-

MEC fellowship. **Author contributions:** MLRAP and MM designed the idea; MLRAP,

MM and RFC designed the experiments, RFN designed and carried out statistical

analysis; RFC and WRC performed flow cytometry analysis; RFC carried out

cytogenetic analysis, field experiments and managed the results; RFC, MM and

MLRAP wrote the paper. **Competing interests:** Authors declare no competing

interests. **Data and materials availability:** All data is available in the main text or the

supplementary materials.

Supplementary Materials:

Materials and Methods

Supplementary text

Figures S1-S5

Table S1

External Databases S1-S2

References

Fig. 1. Representation of the knob conditions for chromosome 9 (K9S). (A) Absence
knob; **(B)** heterozygous knob and **(C)** homozygous knob.

Fig. 2. A sampling of genotypes used in the experiments on flowering time and genome size. (A) – **(G)** Somatic karyotypes of JD lines; and **(H)** – **(L)** hybrids, labeled by FISH using a probe for the knob 180-bp repeat (green).

493
Fig. 3. Flowering time and Genome Size of the hybrid and inbred lines. Boxplots

compare mean values between male and female flowering time and genome size for

inbred lines and hybrids. The flowering time showed in days to flowering and the

genome size in picograms. The boxes indicate the first quartile (lower line), the second

quartile or mean (central line), and the third quartile (upper line). Additionally, the

whiskers represent the standard deviation with the dots as the outliers.

Fig. 4. Flowering time among inbred line families and hybrids. The violin plot

shows the FT distribution in days for all analyzed materials. The violin shape represents

the estimated value density of the trait within each group. Hybrid, n = 175; JD 1-3, n =

36; JD 2-1, n = 25; JD 4-4, n = 63.

Fig. 5. Comparison between dosage class of knobs for male and female flowering time, and genome size. The flowering time is expressed in days to flowering and genome size in picograms. The class means all possible combinations of presence and absence homozygous or heterozygous of the knobs considering only chromosomes 3, 5, 7, and 9.

Fig. 6. Manhattan plots of GWAS for male (mf) and female (ff) flowering time

using the knob positions (K3L, K5L, K7S, and K9S). (A) and **(B)** are plots showing

the allele substitution effect model, **(C)** and **(D)** the dominance effect model, both with

p-values < 0.05. The green lines are the significance threshold, a Bonferroni-corrected

significance threshold used to identify significant associations.

Fig. 7. Boxplots show male (MF) and female (FF) flowering time for each knob condition in the K3L, K5L, K7S, and K9S chromosomes, considering both allele substitution and dominance effects. The x-axis shows the classification for each condition 00 = knob absence; +0 = heterozygous knob presence and ++ = homozygous knob presence. The y-axis shows the flowering days of female and male plants.

523 **Table 1. Pearson's correlation between traits**

	Male flowering	Genome size	Knob dosage classes
Female flowering	0.95	-0.14	-0.20
Male flowering		-0.17	-0.24
Genome size			-0.08

525 **Table 2. Significant marker-trait association for flowering time.** Male flowering

526 (MF), female flowering (FF), knob in the short arm of chromosome 9 (K9S), *p*-value

527 Bonferroni test, minor allele frequency (MAF), allele substitution model (A),

528 dominance model (D), and heritability (h^2).

Trait	Marker	Chr.	Position	*P-value*	MAF	Effect (A or D)	h^2
MF	K9S	9	Short Arm	1.32×10^{-2}	0.17	-0.51 (A)	0.05
	K9S	9	Short Arm	4.21×10^{-3}	0.46	-1.41 (D)	0.15
FF	K9S	9	Short Arm	2.12×10^{-2}	0.17	-0.45 (A)	0.05
	K9S	9	Short Arm	3.75×10^{-3}	0.46	-1.53 (D)	0.02