Glycosylated Antibiotics: New Promising Bacterial Efflux Pumps Inhibitors

Zahraa Ali Kamaz[1*], Haruna Isiyaku Umar[2], Parth Doshi[3], Praveenya Suri[4]

[1,2,3,4] Ioncure Tech Pvt. Ltd. 560065, Karnataka, India

110085 Rohini sector 13, Delhi, India.

* Corresponding author.
E-mail: biologyme1983@gmail.com

Running title: Zahraa K / *Glycosylated Antibiotics: New Promising Bacterial Efflux Pumps Inhibitors.*

Total counts of words 3580, total figures 7, total tables 3

Abstract

Antimicrobial resistance is considered a major concern; bacteria have evolved mechanisms to

overcome antibiotics' action through evolutionary process. One major resistance mechanism that

bacteria developed is the pumping of the antibiotics out of bacterial cells by transmembrane

transporter proteins known as efflux pumps. To overcome such resistance, small molecules known

as efflux pumps inhibitors (EPIs), are created to inhibit the efflux pumps functions. The new

inhibitors are glycosylated antibiotics which have been created by adding N-acetyl glucose amine

moiety to acceptor hydroxyl group of five common antibiotics are Ampicillin, Amoxicillin,

Chloramphenicol, Ciprofloxacin, and Tetracycline. To test the effectiveness of the new EPIs in

inhibiting AcrB-TolC and MexA-OprM efflux pumps functions, docking and simulation studies

have been applied. Docked glycosylated tetracycline has given the highest binding energy in the

active sites of both pumps, with -9.4 against AcrB and -8.8 against MexA. The simulation study

has confirmed the binding of the glycosylated tetracycline in the active sites of both pumps, as

well as its stability during the biological dynamicity of both pumps (opening and closing channels).

However, for the cost purpose, this study included short simulation time of 2ns that prevent the

proteins-ligands complexes from reaching the plateau.

Key words

Antibiotics; Antimicrobial resistance; Molecular docking; Efflux pumps inhibitors; Glycosylated

antibiotics

Introduction:

Bacterial antibiotics resistance is a globally concern problem, bacteria have evolved mechanisms to counter antibiotics' actions, main one involving the pumping out of antibiotics from bacterial inside into the external environment by specialized membrane proteins known as efflux pumps, which confer bacterial resistance to a wide range of antibiotics [1,2]. Efflux pumps are bacterial transport proteins that extrude toxins and antibiotics out of bacterial cells by energy dependent mechanisms [3], thereby classified into ABC active transporters, derived their energy from ATP molecules while other transporters' families derived their energy by chemical gradients which are MFS (Major Facilitator Superfamily), MATE (Multidrug and toxic compounds extrusion), SMR (The small multidrug resistance), RND (Resistance Nodulation-division), and DMT (Drug metabolite transporter) [4,5]. Well studied efflux pumps are AcrB-TolC in *E.coli* and Mex-OPrM that mostly found in *Pseudomonas aeruginosa* [6,7], molecular based structures for both transporters have been elucidated extensively, TolC and OprM are tunnel structures that span the outer membrane of Gram-negative bacteria, It's a homotrimer composed of α- helical barrel, β-barrel and α/β mixed fold [1], AcrB and Mex are fusion or linker proteins that are directly in contact with TolC and OprM respectively, whereas the inner cytoplasmic domain is the transporter protein, some of its parts spanning the cytoplasmic membrane to come in contact with fusion parts, for better descriptions, both pumps structures are illustrated in **(Figure 1)**, [1,8].

As the important role of bacterial efflux pumps in bacterial treatments' failures and concurrent bacterial infections, numerous natural and synthetic compounds have been repurposed as efflux pumps inhibitors (EPIs), such compounds either cause conformational change in the structure of efflux pumps to inhibit its functional assembly or block the substrates binding site of the efflux pumps [9]. PAβN (Phenylalanine-arginine β-naphthylamide) is the first discovered inhibitor for Mex-OprM efflux pump that blocks the binding site of the pump resulting in an increase of the efflux sensitive antibiotic inside bacterial cells, Pyridopyrimidine derivative D13-9001 restore the antibiotic activity by competitive binding with the substrate binding site of AcrB-TolC and Mex-OprM efflux pumps [10]. Several other natural EPIs are plants alkaloids, polyphenols, quinolones derivatives and others, majority of them with unknown exact mechanisms of action [4,11].

EPIs are considered as potential antibiotics adjuncts to overcome multi-drugs resistance, working by combining EPIs with antibiotics course of treatments, or simply just modifying the antibiotics structure to create new EPIs or so called antibiotics analogues such as tetracycline analogue that recently known to restore tetracycline effectiveness in resistant bacteria [12], glycosylation of antibiotics is mostly common among macrolides, bacterial glycosyltransferase transfers UDP-glucose moiety into antibiotic's OH receptor and circumvents the antibiotics activity [13,14]. Glycosylated antibiotic can be synthesized through modifying the chemical structure of the specified antibiotic, such new class of antibiotics might restore the antibiotic activity in the resistant bacteria by working as EPIs or antibiotics analogues, to test the hypothesis, sweet antibiotics have been docked against the active binding site of AcrB and MexA efflux pumps and molecular dynamic simulation has been applied to figure out the conformational changes of the mentioned efflux pumps upon sweet antibiotics binding.

152 **Results**

1. Pharmacokinetic properties

Lead likeness properties for all five glycosylated antibiotics were measured, Lipinski's role of 5 to
predict drugs likeness properties for all new glycosylated antibiotics were followed, the drugs
criteria are MW of the lead ≤ 500, less than 5 hydrogen donors, less than 10 hydrogen bond
acceptor, Log P or lipophilicity not greater than 5 [15], all lead-likeness properties along with
Lipinski's number of violations are listed in (**Table 1**).

2. **Docking results**

The binding energy of all ligands "glycosylated and non-glycosylated antibiotics" against AcrB and
MexA efflux pumps are illustrated in (**Tables 2, 3**). Glycosylated tetracycline has given the highest
binding energy against both efflux pumps, with -9.7 against AcrB efflux pump and -8.8 against
MexA, the interacted amino acids residues in the binding pockets for both pumps against
glycosylated tetracycline are explained in **(Figures 2, 3)**. Glycosylated tetracycline has shown
some similarity in the binding pocket of MexA efflux pump when compare it to the binding pocket
pyridopyrimidine in the same pump, shared amino acids residues are GLN110, LYS 108 and SER
107 **(Figure 3b)**. of Two EPIs were included in the docking against studied efflux pumps for
comparison.

3. Molecular dynamic simulation

RMSD value for the 6TA6- glycosylated tetracycline complex was plotted as in **(Figure 4)**, protein-
ligand complex was not reaching the plateau since short time of simulation was devoted.
However, the efflux pump dynamicity has been captured during the opening and the closing state
of the pump **(Figure 5)**. The md simulation of AcrB efflux pump was done in 2 ns scale, RMSD
value for AcrB-glycosylated tetracycline complex is explained in **(Figure 6)**, whereas **(Figure 7)**
shows the complex conformational change during the opening and the closing process.

Discussion

In order for the newly drugs candidates to pass clinical trials, there are set of ADME properties that the drugs should meet. ADME stands for Absorption, Distribution, Metabolism and excretion, most of drugs are administered orally, so the oral bioavailability is such an important feature [16].

Three out of the five glycosylated antibiotics in the current study were seen to violate Lipinski's rule for molecular weight determinant. As the importance of low molecular weight for orally administered drugs [17], however, there are other important features that all the understudied glycosylated antibiotics have such as acceptable oral bioavailability, no toxicity and no blood brain barrier permeability. A study has confirmed the presence of many FDA approved antibiotics and antifungals that have violated the rule of 5, the same study had explained that such drugs are still orally bioavailable because it has a specific group that works as transporters' substrate [18]. The rule of 5 appears as important data set that the oral rout drugs should meet, in a study has examined about 111 non-oral rout drugs, concluded that Lipinski's criteria for such drugs are not so important to meet [19].

The drugs' journey starts by its absorption through gastrointestinal system, reaching blood stream and binding with the specified target to exert its action. The targeted proteins are two bacterial efflux pumps known as AcrB and MexA, the glycosylated tetracycline has shown the highest binding energy against both target proteins. Tetracycline resistance mediated by efflux pumps has increased recently, various factors are responsible for such resistance rise including the pumps genes are DNA mobilized that can transfer from one bacterium to another. Another factor is the long-term use of tetracycline causing the bacteria to co- evolve itself to extrude such toxins outside of the bacterial cells [20]. Developing molecules that target bacterial efflux pumps has been considered as a new treatment approach, such molecules are known as efflux pumps inhibitors (EPIs). One of the big issues that hurdle the use of EPIs is their toxicity to human cells [20]. Finding safe and effective EPIs is on the increase, of which one potential way is through modifying the structure of the effluxed antibiotics, such modification might help in blocking the binding capability of the used antibiotic and successfully reverse the antibiotic resistance mechanism [21]. New tetracycline derivative known as glycylcycline has been developed as a combination therapy along with tetracycline to overcome Tet efflux pump resistance [22]. Glycylcycline which sometimes known as a tigecycline has similar structure to tetracycline with substitution of an N-alkylglycylamido group at D-9 position [23]. Another modification can be achieved through adding various glycosidic chains at C, O, N atoms on antibiotics, such glycosylation is a naturally occurring process that have been observed among number of

antibiotics including macrolides and streptomycin [24]. Understudied glycosylated antibiotics
have shown promising binding energy against the target proteins in compare to synthetically
available EPIs. Because the targets are efflux pumps transporter proteins which are a dynamic
structure, molecular dynamic simulation has been applied to test the stability of the newly drugs
in the binding pockets of its targets.

The molecular dynamic simulation studies of bacterial efflux pumps are a new era of interesting,
investigating ligands stability among different conformational changes that efflux pumps
undergone such as the transition from substrate binding state into opening state is facing issues
at large scale proteins level [25]. The first md simulation studies for AcrB-TolC and MexA-OprM
have been started since 1999 and still ongoing progresses till now, different simulation
approaches have been used including mutable basin simulation, steered md simulation and un
biased equilibrium md simulation [26]. Identifying ligand binding site or what's so called the
hotspots is crucial in drug discovery, mixed solvent md simulation is successfully used for such a
purpose, one of its big limitation is the cost of running such kind of simulation [27].
239
240
241 **Conclusion**: Glycosylated tetracycline is showing promising results to work as EPIs without
242 worrying about safety issues, however, it's clinical approving require applying longer simulation
243 time of 50 ns or more, as well as further in vitro studies are required to measure its activity
244 against resistant bacteria. Glycosylation of other antibiotics with lower molecular weight and less
245 Lipinski's role of 5 violations are recommended.

Materials and Methods

1. Glycosylation of antibiotics

Three dimensional structures of amoxicillin, ampicillin, chloramphenicol, ciprofloxacin and

tetracycline were retrieved from pubchem web server (https://pubchem.ncbi.nlm.nih.gov/) in

simple document format (SDF). The Pubchem CIDs for the studied antibiotics are 33613, 6249,

5959, 2764, and 54675776 respectively.

N-acetyl glucose amine moiety has been added to the hydroxyl acceptor group for each selected

antibiotic, using Marvin sketch software 20.20, then the ligands were energetically optimized to

the most stable structures using merk molecular force field 94 (MMFF94).

2. Pharmacophore kinetics properties

The drug-likeness properties have been measured for all glycosylated antibiotics using the

website http://www.swissadme.ch/. The role of 5 for Lipinski as well as others features have

been measured including the oral bioavailability, GI absorption, Blood brain barrier permeability,

and liver cytotoxicity.

3. Molecular docking

3D x-ray crystallographic structures of AcrB and MexA were retrieved from the protein data bank

(PDB) with IDs 2I6W and 6TA6 respectively. After which the proteins were prepared for docking

and protein-ligand complexes minimized using the relevant tools in Cresset Flare© software,

version 4.0 (https://www.cresset-group.com/flare/). The protein minimization was done based on

the GAFF force field, with gradient cutoff of 0.200 kcal/mol/A and iterations was set to 2000

iterations [28]. Furthermore, the docking was achieved using flexible docking protocol [29]. In

brief; Python Prescription 0.8, a suite housing Auto Dock Vina, was employed for the molecular

docking study of the selected ligands with the two efflux pumps. The specific target site for AcrB

was set using the grid box with dimensions (29.1800 x 22.3491 x 31.3636) Å, while that of MexA

was (68.1632 x 57.6364 x 103.9827) Å and their centres were adjusted based on the active site of

the enzymes. At the end of the molecular docking, 10 binding poses with least binding affinity for

each protein-ligand complex were generated for all the ligands, scoring results were also created

after retrieving from text files which were employed for comparative analysis. The protein-ligand

complexes as well the molecular interactions were all visualized in 3D and 2D using PyMOL©

Molecular Graphics (version 2.4, 2016, Shrodinger LLC) and Protein- Ligand interaction Profiler

PLIP server (https://plip-tool.biotec.tu-dresden.de/plip-web/plip/index).

4. Molecular dynamic simulation

For glycosylated tetracycline which give the lowest binding affinity against both efflux pumps'
binding sites, molecular dynamic simulation was performed using NAMD 2.8, the topology file
for the efflux pumps proteins were generated using automatic PSF builder in VMD program,
whereas, the topology file for the ligand was generated using CHARMM GUI server
(http://www.charmm-gui.org/), 2ns simulation was performed with time step of 2 fs, periodic
boundary conditions were used, the temperature kept at 310 K and the pressure was stable at
1.013 bar, 100 steps of equilibration were completed before starting the solvate simulation, the
whole md simulation protocol can be found [30]. CDC trajectory file for both efflux pumps- ligand
complexes was analyzed in the form of measuring RMSD values using VMD graphical visualization
program 1.9 version.

317
318

Authors' contributions
ZK carried out the study experimental design, molecular dynamic simulation work and writing
the first draft of the research paper. HU carried out the modification of antibiotics' structures
work, did part of the docking work and reviewed the paper. PD did most of the docking work,
and analyzed the docking results. PS confirmed the docking results by replicated it and
reviewed the paper.

325
326

327 **Competing interests**
328 The authors have declared no competing interests

345
346
347
348
349 **References**
350
[1]. Jamshidi S, Sutton JM, Rahman KM. An overview of bacterial efflux pumps and
computational approaches to study efflux pump inhibitors. Future Medicinal Chemistry.
2016.
[2]. Soto SM. Role of efflux pumps in the antibiotic resistance of bacteria embedded in a
biofilm. Virulence. 2013;
[3]. Blanco P, Hernando-Amado S, Reales-Calderon J, Corona F, Lira F, Alcalde-Rico M, et al.
Bacterial Multidrug Efflux Pumps: Much More Than Antibiotic Resistance Determinants.
Microorganisms. 2016;
[4]. Sharma A, Gupta VK, Pathania R. Efflux pump inhibitors for bacterial pathogens: From
bench to bedside. Indian Journal of Medical Research. 2019.
[5]. Soto SM. Soto 2013. 2013;223–9.
[6]. Tsutsumi K, Yonehara R, Ishizaka-Ikeda E, Miyazaki N, Maeda S, Iwasaki K, et al.
Structures of the wild-type MexAB–OprM tripartite pump reveal its complex formation
and drug efflux mechanism. Nat Commun. 2019;
[7]. Pos KM. Drug transport mechanism of the AcrB efflux pump. Biochimica et Biophysica
Acta - Proteins and Proteomics. 2009.
[8]. Yu EW, Aires JR, Nikaido H. AcrB multidrug efflux pump of Escherichia coli: Composite
substrate-binding cavity of exceptional flexibility generates its extremely wide substrate
specificity. J Bacteriol. 2003;185(19):5657–64.
[9]. P. Tegos G, Haynes M, Jacob Strouse J, Md. T. Khan M, G. Bologa C, I. Oprea T, et al.
Microbial Efflux Pump Inhibition: Tactics and Strategies. Curr Pharm Des. 2012;
[10]. Y. Mahmood H, Jamshidi S, Mark Sutton J, M. Rahman K. Current Advances in Developing
Inhibitors of Bacterial Multidrug Efflux Pumps. Curr Med Chem. 2016;
[11]. Pagès JM, Masi M, Barbe J. Inhibitors of efflux pumps in Gram-negative bacteria. Trends
Mol Med. 2005;11(8):382–9.
[12]. Rodrigues L, Cravo P, Viveiros M. Efflux pump inhibitors as a promising adjunct therapy
against drug resistant tuberculosis: a new strategy to revisit mycobacterial targets and
repurpose old drugs. Expert Review of Anti-Infective Therapy. 2020.
[13]. Quirós LM, Carbajo RJ, Braña AF, Salas JA. Glycosylation of macrolide antibiotics.
Purification and kinetic studies of a macrolide glycosyltransferase from Streptomyces
antibioticus. J Biol Chem. 2000;275(16):11713–20.
[14]. Bambeke F, Pages J-M, Lee V. Inhibitors of Bacterial Efflux Pumps as Adjuvants in
Antibiotic Treatments and Diagnostic Tools for Detection of Resistance by Efflux. Recent
Pat Antiinfect Drug Discov. 2008;1(2):157–75.
[15]. Neidle S. Design Principles for Quadruplex-binding Small Molecules. In: Therapeutic
Applications of Quadruplex Nucleic Acids. 2012.
[16]. Schneider G. Prediction of Drug-Like Properties. In: Adaptive Systems in Drug Design.
2020.

[17]. Ursu O, Rayan A, Goldblum A, Oprea TI. Understanding drug-likeness. Wiley
Interdisciplinary Reviews: Computational Molecular Science. 2011.
[18]. Kadam R, Roy N. Recent trends in drug-likeness prediction: A comprehensive review of in
silico methods. Indian Journal of Pharmaceutical Sciences. 2007.
[19]. Choy Y Bin, Prausnitz MR. The rule of five for non-oral routes of drug delivery:
Ophthalmic, inhalation and transdermal. Pharmaceutical Research. 2011.
[20]. Tuckman M, Petersen PJ, Howe AYM, Orlowski M, Mullen S, Chan K, et al. Occurrence of
tetracycline resistance genes among Escherichia coli isolates from the phase 3 clinical
trials for tigecycline. Antimicrob Agents Chemother. 2007;
[21]. Marquez B. Bacterial efflux systems and efflux pumps inhibitors. Biochimie. 2005.
[22]. Chopra I. New developments in tetracycline antibiotics: Glycylcyclines and tetracycline
efflux pump inhibitors. Drug Resist Updat. 2002;
[23]. Zhanel GG, Homenuik K, Nichol K, Noreddin A, Vercaigne L, Embil J, et al. The
Glycylcyclines: A Comparative Review with the Tetracyclines. Drugs. 2004.
[24]. Piepersberg W. Glycosylation of Antibiotics and Other Agents from Actinomycetes. In
2001.
[25]. Wong K, Ma J, Rothnie A, Biggin PC, Kerr ID. Towards understanding promiscuity in
multidrug efflux pumps. Trends in Biochemical Sciences. 2014.
[26]. Fischer N, Raunest M, Schmidt TH, Koch DC, Kandt C. Efflux pump-mediated antibiotics
resistance: Insights from computational structural biology. Interdiscip Sci Comput Life Sci.
2014;
[27]. Bernardi RC, Melo MCR, Schulten K. Enhanced sampling techniques in molecular
dynamics simulations of biological systems. Biochimica et Biophysica Acta - General
Subjects. 2015.
[28]. Stroganov O V., Novikov FN, Zeifman AA, Stroylov VS, Chilov GG. TSAR, a new graph-
theoretical approach to computational modeling of protein side-chain flexibility:
Modeling of ionization properties of proteins. Proteins Struct Funct Bioinforma. 2011;
[29]. Trott O, Olson AJ. AutoDock Vina: Improving the speed and accuracy of docking with a
new scoring function, efficient optimization, and multithreading. J Comput Chem. 2009;
[30]. Nair VV, Varghese V, Paul RR, Jose A, Sinu CR, Menon RS, et al. Supporting Information. J
Am Chem Soc. 2012;

Figure legends

Figure 1 The structural assembly of MexA efflux pump
The structural components of MexA efflux pump on the left panel, that its similar to the structure of AcrB efflux pump on the right side, the Image has been taken from [1].

Figure 2 The binding site of the glycosylated tetracycline in Acrb efflux pump binding pocket
 A. The left side is the binding pose of the glycosylated tetracycline
 B. The right side represents the interaction between amino acids residues of AcrB chain A and the glycosylated tetracycline, the new antibiotic showed hydrophobic interaction with PRO 40A, GLU 673A and Hydrogen bonds with TYR 77A, SER 79A, THR 91A, SER 134A, GLU 683A, GLU 826A.

Figure 3 The binding sites of all of the glycosylated antibiotics in the binding pocket of MexA efflux pump
A. The left side, shows the binding pose of amoxicillin (cyan), ampicillin (pink), chloramphenicol (yellow), ciprofloxacin (red) and tetracycline (blue) in MexA. The drugs occupy similar region in MexA except amoxicillin.
B. The right side is the interaction between residue of MexAB – OprM efflux pump and Glycosylated tetracycline, that showed hydrophobic interaction ASP 230D, THR 233D, GLU 796J and hydrogen bonds with PRO 32D, ASN 36D, ARG 147E, ASP 176D, THR 178D, ASN 194J, SER 235D, VAL 236D, THR 237D, MET 797.

Figure 4 The binding site of pyridopyrimidine derivative D13-9001
Interacted amino acids residues in the binding pocket of MexA upon Pyridopyrimidine binding, shared residues with the glycosylated tetracycline binding pocket are highlighted in yellow.

Figure 5 RMSD analysis for MexA-OprM- glycosylated tetracycline complex
CDC trajectory file for the solvate complex was used to analyze the RMSD value, 1150 frames were used, the protein-ligand complex didn't reach the plateau because of short simulation time.

 Figure 6 MexA efflux pump opening and closing states upon glycosylated tetracycline binding
 the dynamic movement of MexA efflux pump upon binding with the glycosylated tetracycline, it shows the stability of the ligand during the closing of the channel **A.** and the opening of the channel **B.**

Figure 7 RMSD analysis for AcrB- glycosylated tetracycline complex

RMSD value for AcrB- glycosylated tetracycline was calculated using VMD program, 4250 frames were being used in the analysis of the CDC trajectory file. Again protein-ligand complex didn't reach the plateau.

Figure 8 AcrB efflux pump opening and closing state upon glycosylated tetracycline binding

Shows the closing state **A.** and the opening of AcrB efflux pump **B.**, the glycosylated tetracycline is stable in the binding pocket of the target protein.

Tables

Table 1 Pharmacokinetics properties of the newly glycosylated antibiotics.

Table 2 Docking results of N-acetyl glucose amine antibiotics against AcrB efflux pump in *E. coli* and MexA efflux pump in *Pseudomonas aeruginosa*.

Table 3 Binding Energy of Non- glycosylated antibiotics docked against ArcB and MexA.

Supplementary material

Figure 9 The binding sites of all glycosylated antibiotics against AcrB efflux pump

Binding pose of amoxicillin (yellow), ampicillin (red), chloramphenicol (white), ciprofloxacin (blue) and tetracycline (orange) in AcrB. The drugs occupy similar region in AcrB.

Figure 10 The binding sites of all glycosylated antibiotics against MexA efflux pump

Binding pose of amoxicillin (cyan), ampicillin (pink), chloramphenicol (yellow), ciprofloxacin (red) and tetracycline (blue) in MexA. The drugs occupy similar region in MexA except amoxicillin.

Table 4 The binding affinities of the glycosylated tetracycline against AcrB and MexA.

Figure 1 The structural assembly of MexA efflux pump
The structural components of MexA efflux pump on the left panel, that its similar to the structure of AcrB efflux pump on the right side, the Image has been taken from [1].

A

B

Figure 2 The binding site of the glycosylated tetracycline in Acrb efflux pump binding pocket

A. The left side is the binding pose of the glycosylated tetracycline

B. The right side represents the interaction between amino acids residues of AcrB chain A and the glycosylated tetracycline, the new antibiotic showed hydrophobic interaction with PRO 40A, GLU 673A and Hydrogen bonds with TYR 77A, SER 79A, THR 91A, SER 134A, GLU 683A, GLU 826A.

A

B

Figure 3 The binding sites of all of the glycosylated antibiotics in the binding pocket of MexA efflux pump

A. The left side, shows the binding pose of amoxicillin (cyan), ampicillin (pink), chloramphenicol (yellow), ciprofloxacin (red) and tetracycline (blue) in MexA. The drugs occupy similar region in MexA except amoxicillin.

B. The right side is the interaction between residue of MexAB – OprM efflux pump and Glycosylated tetracycline, that showed hydrophobic interaction ASP 230D, THR 233D, GLU 796J and hydrogen bonds with PRO 32D, ASN 36D, ARG 147E, ASP 176D, THR 178D, ASN 194J, SER 235D, VAL 236D, THR 237D, MET 797.

Figure 4 The binding site of pyridopyrimidine derivative D13-9001

Interacted amino acids residues in the binding pocket of MexA upon Pyridopyrimidine binding, shared residues with the glycosylated tetracycline binding pocket are highlighted in yellow.

Figure 5 RMSD analysis for MexA-OprM- glycosylated tetracycline complex
CDC trajectory file for the solvate complex was used to analyze the RMSD value, 1150 frames were used, the protein-ligand complex didn't reach the plateau because of short simulation time.

A

B

Figure 6 MexA efflux pump opening and closing states upon glycosylated tetracycline binding
 the dynamic movement of MexA efflux pump upon binding with the glycosylated tetracycline, it shows the stability of the ligand during the closing of the channel **A.** and the opening of the channel **B**.

Figure 7 RMSD analysis for AcrB- glycosylated tetracycline complex

RMSD value for AcrB- glycosylated tetracycline was calculated using VMD program, 4250 frames were being used in the analysis of the CDC trajectory file. Again protein-ligand complex didn't reach the plateau.

A

B

Figure 8 AcrB efflux pump opening and closing state upon glycosylated tetracycline binding
Shows the closing state **A.** and the opening of AcrB efflux pump **B.**, the glycosylated tetracycline is stable in the binding pocket of the target protein.

Table 1: Pharmacokinetics properties of the newly glycosylated antibiotics.

Glycosylated antibiotics	No.H bond donors	No.H bond acceptors	MW	Log p	Blood brain barrier permeability	GI absorption	Liver cytotoxicity	Lipinski's No. violations
Glycosylated Ampicillin	6	10	552.6	1.88	No	Low	No	3
Glycosylated Amoxicillin	7	11	568.6	-0.48	No	Low	No	3
Glycosylated Chloramphenicol	6	10	484.29	0.42	No	Low	No	2
Glycosylated Ciprofloxacin	5	10	492.5	2.04	No	Low	No	1
Glycosylated Tetracycline	9	14	605.59	1.34	No	Low	No	3

Table 2: Docking results of N-acetyl glucose amine antibiotics against AcrB efflux pump in *E. coli* and MexA efflux pump in *Pseudomonas aeruginosa*

Ligand	Binding Energy (kcal/mol)	
	ArcB	MexA
Glycosylated Amoxicillin	-8.7	-7.5
Glycosylated Ampicillin	-8.3	-7.9
Glycosylated Chloramphenicol	-8.0	-7.8
Glycosylated Ciprofloxacin	-9.0	-7.8
Glycosylated Tetracycline	-9.7	-8.8

Table 3. Binding Energy of Non- glycosylated antibiotics docked against ArcB and MexA.

Ligand	Binding Energy (kcal/mol)	
	ArcB	MexA
Amoxicillin	-7.8	-6.3
Ampicillin	-7.7	-6.7
Chloramphenicol	-6.1	-5.8
Ciprofloxacin	-7.5	-6.4
Tetracycline	-8.3	-6.8
PAβN	-9.3	-9.1
Pyridopyrimidine	-8.1	-9.6