Short-interval reburns in the boreal forest alter soil bacterial communities, reflecting increased pH and poor conifer seedling establishment

Jamie Woolet[a,b], Ellen Whitman[c,d], Marc-André Parisien[c], Dan K. Thompson[c], Mike D. Flannigan[d], and Thea Whitman[a*]

a. Department of Soil Science, University of Wisconsin-Madison, 1525 Observatory Dr., Madison, WI, 53706, USA

b. Department of Forest and Rangeland Stewardship, Colorado State University, 1001 Amy Van Dyken Way, Fort Collins, CO, 80521, USA

c. Northern Forestry Centre, Canadian Forest Service, Natural Resources Canada, 5320 122 Street, Edmonton, AB, T6H 3S5, Canada

d. Department of Renewable Resources, University of Alberta, 751 General Services Building, Edmonton, AB, T6G 2H1, Canada

* Corresponding author: twhitman@wisc.edu; 608.263.4947

Abstract: Increasing burn rates (percentage area burned annually) in some biomes are leading to fires burning in close succession, triggering rapid vegetation change as well as altering soil properties. Despite the importance of soil microbes for nutrient cycling and as plant symbionts, the effects of increased fire frequency on belowground microbial communities remain largely unknown. We present a study of the effects of short interval reburns (defined here as <20 years between fires) on soil bacterial communities in the boreal forest of northwestern Canada, using a paired site design that spans wetlands and uplands, with 50 sites total. We asked whether short interval reburns significantly alter soil bacterial community composition and richness, and which bacterial taxa are associated with greater or lower fire frequency. We found that, while short interval reburns had no significant effect on bacterial richness, there were significant changes in overall community composition. We did not find correlations between understory vegetation community dissimilarities and bacterial community dissimilarities, suggesting the primary drivers of changes induced by short interval reburns may differ between plants and microbes. We identified an abundant *Blastococcus sp.* that was consistently enriched in short interval reburns, in both wetlands and uplands, indicating its role as a strongly "pyrophilous" bacterium. We also identified an abundant *Callaberonia sordidicola* taxon as being consistently depleted in short interval reburns. This endophytic diazotrophic organism is a robust colonizer of pine and spruce seedlings and has the ability to increase seedling growth, due in part to large contributions of fixed nitrogen. Its depletion in short-interval reburn sites raises questions about whether this is contributing to – or merely reflects – poor conifer seedling recolonization post-fire at short-interval reburns.

Introduction

The boreal zone is one the world's largest biomes, spanning 1.89 billion ha across the northern hemisphere (Brandt *et al.,* 2013). This zone consists of forests of cold-tolerant tree species, lakes, rivers, wetlands, and naturally treeless areas, such as shrublands and grasslands (Brandt, 2009). The Canadian boreal forest represents 28% - about 552 million ha - of the world's boreal zone (Brandt *et al.* 2013) and provides habitat for thousands of species, supplies numerous ecosystem services including timber, timber products, and water filtration, is home to 12% of Canada's population, and offers many other economic and cultural resources (Bogdanski, 2008). Furthermore, this system stores about 10 - 30% of the global terrestrial carbon stocks, mostly belowground in peatlands and soils (Bradshaw and Warkentin, 2015; Kasischke, 2000), which may be threatened by changing fire regimes (Ribeiro-Kumara *et al.*, 2020, Walker *et al.*, 2019).

Fire is a common and widespread disturbance throughout much of the western Canadian boreal zone, where the average fire-free interval has been observed to range anywhere between 30 – 100s of years (Larsen 1997; Stocks 2002). Fire is a critical event for maintaining healthy boreal ecosystems by shaping vegetation composition, soil chemical properties, and animal communities (Rowe and Scotter, 1973). Over the past 50 years, there has been a shift in the forest fire regime for many areas of the North American boreal forest, including lengthened burn season, increased lightening ignitions, and increased area burned (Hanes *et al*., 2019; Wotton and Flannigan, 1993; Jain *et al*., 2017; Veraverbeke *et al.,* 2017). These shifting disturbance regimes can have adverse effects on ecosystems and may degrade forest resilience to fire (Johnstone *et al*., 2016). Forest resilience, broadly, is the ability for a forest to return to pre-disturbance conditions, often determined by ecological memory of past states (*e.g.*, via seed banks) (Johnstone *et al*., 2016) and the regeneration of plant communities (Gill *et al*., 2017). Under normal fire regimes, forests often have self-regulatory processes that limit disturbance frequency (Peterson, 2002). Under drought conditions and as the forest ages, however, these self-regulatory processes can weaken, allowing for increased fire frequency (Parks *et al.,* 2018). In the relatively uncommon case when young forests (<20 years) reburn, increased fire frequency (short interval reburns) in boreal forests can alter vegetation composition (Whitman *et al.,* 2019b), change above-ground plant production (Johnstone and Chapin, 2006), and potentially induce forest-type

conversions (*i.e.*, *Picea* spp.-dominated to *Populus tremuloides*- or *Pinus banksiana*-dominated)

(Johnstone and Chapin, 2006; Gill *et al*, 2017). Short interval reburns also reduce soil organic

horizon thickness (Johnstone and Chapin, 2006; Hoy *et al.,* 2016), change soil chemical

properties by depleting total C and N (Pellegrini and Jackson, 2020; Pellegrini *et al.*, 2020), and

can potentially decrease microbial decomposition rates (Köster *et* al., 2015; Pellegrini *et al.*,

2020). Furthermore, novel wildfire regimes have been found to have long-term effects on

biogeochemical soil processes, decreasing mineral soil organic carbon (SOC), soil extracellular

enzyme activity, and soil microbial respiration (Dove *et al.,* 2020), which may interact with

vegetation responses to fire (Knelman *et al*., 2015). However, it is less clear how short interval

reburns may affect soil microbial communities.

Soil microbial communities provide numerous critical ecosystem functions, including cycling

carbon and nitrogen, supporting plant growth and diversity, preventing erosion, and maintaining

soil structure via biofilms and fungal hyphae (Van Der Heijden *et al.,* 2007; Saleem *et al.,* 2019).

Fire can affect the soil microbial communities directly, via heating and oxidation of the soil

environment, and indirectly, by increased exposure to climatic variation and from changes to the

physicochemical environment (Hart *et al.*, 2005). Immediately post-fire, microbial biomass may

decrease due to direct killing of microbes or the loss of nutrient resources (Dooley and Treseder,

2011; Holden and Treseder, 2013; Pressler *et al*., 2019). Microbial community structures may

take decades to recover to previous states, often requiring plant community reestablishment first

(Dooley and Treseder, 2011; Ferrenberg *et al.,* 2013). The extent to which microbial community

composition is affected by fire is influenced by fire severity and changes to vegetation, moisture,

pH, and soil carbon after fires (Whitman *et al*., 2019a; Hart *et al*., 2005; Holden and Treseder,

2013; Sáenz de Miera *et al*., 2020).

Microbial resistance and resilience may inform our understanding of forest resilience following

fire. Microbes have innate traits that differ from those of some of their larger-organism

counterparts (high abundances, widespread dispersal potential, comparatively rapid growth

potential, and comparatively rapid evolutionary adaptations (Shade, 2012)).The stability of their

populations over time can be influenced by an individual's stress tolerance and phenotypic

plasticity, a population's growth rate and adaptability, and a community's richness, evenness,

and microbial interactions (Shade, 2012). Microbial communities may show resistance by

recovering as a compositionally different community, yet remaining functionally similar, in that

the ecosystem process rates of interest remain unchanged (Allison and Martiny, 2008). Short

interval reburns have the potential to change soil properties and vegetation community

composition, both of which are factors that shape microbial communities (Chandra *et al.,* 2016;

Woolet and Whitman, 2020; Van Der Heijden *et al.*, 2007; Bardgett and van der Putten, 2014).

However, there are few studies that examine how microbial communities respond to these

potentially transformative short interval reburns. Because microbes play a critical role in

ecosystem functioning and structure, it is important to understand how they respond to fire

regime changes.

Using a paired sample study design, Whitman *et al*. (2019b) showed that changes in fire-return

interval caused changes to conifer and broadleaf recruitment, soil organic horizon depth, and

herbaceous vegetation cover. Here, we set out to investigate soil bacterial community responses

to paired short interval (SI) vs. long interval (LI) reburns at the same sites (Whitman *et al.*

2019b). Specifically, we asked 1) Do microbial communities have a different response to short

interval reburns compared to normal fire intervals?, 2) Do short interval reburns reduce bacterial

community richness?, and 3) Which bacterial taxa respond positively to increased fire

frequency? We hypothesized that:

1) There would be a significant effect of SI vs. LI reburns on bacterial community

119   composition, where the (dis)similarity of bacterial communities between paired SI and LI

120   sites would be dependent on:

121   a) Time since last fire (TSLF): paired sites that had a longer time to recover at the

122   sampling date would be more similar than sites that had less time to recover

123   b) Vegetation transition: paired sites that show a change in leading tree species

124   would be less similar than pairs that did not see a change in leading tree species

125   due to changes in rhizosphere interactions and differences in litter inputs

126   c) Drainage class: paired wetland sites would be more similar than paired upland

127   sites due to wetland sites generally having lower severity fires and thus retaining

128   more of the organic horizon

129   d) Difference in fire-free interval (FFI): paired sites that had greater differences in

130   fire-free interval would be less similar due to larger pre-fire differences in

131   successional trajectory

2) Bacterial richness would not be consistently different between paired SI and LI sites,

but would be predicted by pair-specific characteristics (TSLF, difference in FFI)

3) Bacteria that have higher relative abundance in SI sites than LI sites would include

taxa that have previously been identified as being responsive to fires and associated with

increasing burn severity in this region, such as certain taxa within the genera

Blastococcus, *Arthrobacter*, and *Massilia* (Whitman *et al.*, 2019a).

Methods

Study region description

Our study area is located in the Canadian boreal forest in the Northwest Territories and Alberta

around Tu Nedhé (Great Slave Lake) (Figure 1).

Figure 1. Location and fire history of the study area (Figure adapted from Whitman *et al*., 2019b). (**a**) Recent (1984–2016) fire perimeters (dark grey) within the boreal forest of North America (green). (**b**) Fire history within the study region around Tu Nedhé (Great Slave Lake). Recent wildfires (1980–2015) are coloured grey, while sampled wildfires are coloured by year of occurrence. Shapes represent sampling locations. The boundary of Wood Buffalo National Park (WBNP) is outlined in grey and major roads are shown in black. (**c**) Detail showing an example of the sampling design of paired sites with short (triangles) and long (squares) interval sites. E.g., consider the triangle (SI) and square (LI) in SW corner – both were burned in the 2004 wildfire, but only the SI site was also burned in the 1998 wildfire.

Typically, the climate consists of short, warm summers, and long, cold winters. Fire frequencies in this area range from hundreds of years to 30 years between stand-replacing fires (Larsen and MacDonald, 1998; Stocks *et al.* 2002). The landscape consists of open wetlands, forested peatlands, and forested uplands, with black spruce (*Picea mariana* (Mill.) Britton, Sterns & Poggenb.), white spruce (*Picea glauca* (Moench) Voss), jack pine (*Pinus banksiana* Lamb.), and trembling aspen (*Populus tremuloides* Michx.) as the dominant tree species. The soils in this region are mostly classified as Mesisols, Gleysols, and Luvisols. The sample sites span a wide range of soil properties, with pH values ranging from 5.13–8.11 in the mineral horizons and 5.29–8.35 in the organic horizons, total C ranging from 0.4% (mineral horizon) to 44.6% (organic horizon), and a wide range of textures (Table 1). Detailed information on the study area and field sampling methods can be found in Whitman *et al.* (2019b).

Table 1. Soil properties across drainage class, short and long fire-free intervals, and organic and mineral soil horizons. Means reported (N = 3). SI=Short-interval, LI=Long-interval.

	Upland				Wetland			
	SI Organic	SI Mineral	LI Organic	LI Mineral	SI Organic	SI Mineral	LI Organic	LI Mineral
pH	6.83	6.78	6.52	6.25	6.92	7.34	7.02	6.94
EC (mScm^{-1})	0.32	0.33	0.25	0.09	0.81	0.23	0.66	0.23
% Clay		9		9		10		17
% Silt		32		24		26		54
% Sand		59		67		64		28
% Total N	0.45	0.07	0.58	0.14	1.69	0.27	1.53	0.55

% Total S	0.11	0.72	0.07	0.01	0.39	0.03	0.31	0.07
% Total C	10.52	1.75	15.77	2.29	32.45	4.63	35.70	11.11
% OM by LOI	27.40	5.10	31.53	4.72	68.62	3.95	80.68	5.40
CEC (cmol kg^{-1})	50.77	11.36	74.54	12.04	104.01	15.33	106.11	28.01
Ca (mg kg^{-1})	9,480	6,403	9,466	2,051	20,944	5,330	24,116	9,628
K (mg kg^{-1})	231	72	335	77	518	139	532	252
Mg (mg kg^{-1})	626	262	837	262	2,376	344	2,262	672
Na (mg kg^{-1})	95	15	91	13	230	14	232	46

Experimental design and site assessment

In 2016, 50 sites (25 pairs) were sampled; eight of these pairs were wetlands and 17 were

uplands. Paired sites shared the same drainage class, fire history, and their most recent fire,

which burned between 1995 and 2015, but had different previous fires, which allowed us to

characterize them as short fire-free interval fires (SI; fire-free interval was 4-17 years) or long

fire-free interval fires (LI; fire-free interval was 30-112 years). Fire history maps and fire-scarred

trees were used to date recent and previous fires (Whitman *et al*., 2019b). Time since last fire at

sampling varied across the dataset, ranging from 1 – 21 years.

Sites were selected > 100 m from roads. At each site, a 35 m transect oriented north-south was

used to collect vegetation data and soil samples. Vegetation survey methods are described in

detail in Whitman et al. (2019b). Briefly, along the transect, overstory structure was assessed by

sampling live and dead mature trees (greater than 1.3 m in height and greater than 3 cm diameter

at breast height (DBH)) stem density, basal area, and species composition. Understory structure

was assessed by collecting vegetation and small shrub abundance, tall shrub counts, the density,

species, and status (live or dead) of seedlings and saplings, surface fuel, and coarse woody

debris. At 0, 17.5, and 35 m along the transect, soil cores (5.5 cm diameter, 13.5 cm depth) were

sampled by gently extruding and separating the core into organic (O) horizons (where present, up

to 13.5 cm) and mineral (M) horizons (where present, up to 5 cm). Wetland mineral soils

underlying thin organic horizons were only present within the top 13.5 cm at three sites. The

three samples from the transect were pooled by horizon at each site and mixed gently by gloved

hand in a bag. From these site-level samples, sub-samples were collected for microbial

community analysis and stored in LifeGuard Soil Preservation solution (Qiagen, Germantown,

MD) in a 5 mL tube. Tubes were kept as cold as possible while in the field, then stored frozen.

The remaining soil samples were air-dried and analyzed for soil properties, as described in detail

in Whitman *et al.* (2019b) (Table 1).

DNA extraction, amplification, and sequencing

DNA extractions were performed for each sample, with a blank extraction every 24 samples,

using a DNEasy PowerLyzer PowerSoil DNA extraction kit (QIAGEN, Germantown, MD)

following manufacturer's instructions. LifeGuard Soil Preservation solution was removed from

samples by thawing the sample on ice and centrifuging for 2 minutes at 10,000 x g. Preservation

solution was gently pipetted off to remove as much as possible, and the sample was centrifuged a

second time for 30 seconds at 10,000 x g to remove any remaining preservation solution. The

sample was re-mixed with a spatula before weighing for extraction. Extracted DNA was

amplified in triplicate PCR reactions, targeting the 16S rRNA gene v4 region (henceforth,

"16S"). Reactions were performed in 96-well plates with each PCR mixture containing 12.5 µL

Q5 Hot-Start High-Fidelity 2X Master Mix (New England Biolabs, Ipswich, MA), 1.25 µL 10

µM 515F primer (AATGATACGGCGACCACCGAGATCTACAC-barcode-TATGGTAATT

GTGTGYCAGCMGCCGCGGTAA), 1.25 µL µM 806R primer (CAAGCAGAAGACGGCAT

ACGAGAT-barcode-AGTCAGCCAGCCGGACTACNVGGGTWTCTAAT) (Walters *et al.*

2015; Kozich *et al.* 2013), 1.25 µL 20 mg mL^{-1} BSA, 7.75 µL nuclease-free water, and 1 µL

DNA template. Positive control (bacterial isolate DNA) and negative control (nuclease-free

water) reactions were included on each plate. PCR mixtures were amplified on an Eppendorf

Mastercycler nexus gradient thermocycler (Eppendorf, Hamburg, Germany) at the following

conditions: 98°C for 2 minutes, (98°C for 10 seconds, 58°C for 15 seconds, 72°C for 10 seconds)

x 30 cycles, 72°C for 2 minutes, hold at 4°C. PCR amplification success was verified via gel

electrophoresis on a 1% agarose gel. The amplicon triplicates for samples and extraction blanks

were pooled and normalized using a SequalPrep Normalization Plate (96) Kit (ThermoFisher

Scientific, Waltham, MA), following manufacturer's instructions. Normalized samples were

pooled and library cleanup was performed using a Wizard SV Gel and PCR Clean-Up System

A9282 (Promega, Madison, WI). The purified library was submitted to the UW Madison

Biotechnology Center (UW-Madison, WI) for 2x250 paired end (PE) Illumina MiSeq

sequencing.

Sequence data processing and taxonomic assignments

The University of Wisconsin – Madison Biotechnology Center performed demultiplexing on

sequences. The total read count was 5,141,495 sequences; the minimum (non-blank) read count

was 18,297; the maximum read count was 52,378; and the mean read count (not including

blanks) was 30,826; sample blanks averaged 2,437 reads and did not produce any visible bands

in electrophoretic gels. Forward and reverse reads were imported into a Jupyter Notebook where

QIIME2 (QIIME2, v 2019.10) and dada2 (Callahan *et al.*, 2016) were used to filter, learn error

rates, denoise, and remove chimeras, to generate operational taxonomic units (OTUs) –

specifically, amplicon sequence variants. After quality control steps, a total of 3,593,901 reads

were retained. Taxonomy was assigned using the QIIME2 scikit-learn feature classifier trained

on the 515f-806r region of the 99% ID OTUs (Bokulich *et al.*, 2018) from the Silva138 database

(Quast *et al.*, 2013). All sequences are deposited in the NCBI SRA under accession numbers [to

be submitted upon paper acceptance].

Statistical analyses

Analyses and plotting were done with R (R Core Team, 2019) in Jupyter notebooks and RStudio,

using packages *phyloseq* (McMurdie and Holmes, 2013), *dplyr* (Wickham et al., 2019), and

ggplot2 (Wickham, 2016).

To test for the effect of reburn interval on bacterial community composition, we calculated the

Bray-Curtis dissimilarities for all sites from relative abundances, using the *vegan* package in R

(Oksanen et al., 2018, Bray and Curtis, 1957), and used permutational multivariate analysis of

variance (PERMANOVA) to determine whether SI *vs*. LI was a significant predictor of

community composition after controlling for paired sites. We plotted the dissimilarities using

NMDS and used the *envfit* function in *vegan* to map site characteristics onto the ordination. To

determine what factors to control for in subsequent data analyses, we used PERMANOVA to test

whether drainage class, soil horizon, and pH were also significant predictors. We then calculated

Bray-Curtis dissimilarities for each site-horizon pair. After controlling for soil horizon and

drainage class (except for when testing drainage class itself), we used ANOVA to test whether

the dissimilarities between paired sites were affected by 1) time since last fire, 2) vegetation

transition (*i.e.*, whether there was a change in leading species status, as inferred from different

leading species in the paired sites), 3) drainage class (wetland or upland), and 4) difference in

FFI between paired sites.

Because vegetation communities were affected by FFI in Whitman *et al.* (2019b), we also

examined the relationship between vegetation-related parameters and bacterial community

dissimilarities in LI *vs*. SI sites. These parameters included the effects of absolute difference in

density of small understory tree stems between paired sites (including all stems, broadleaf stems,

and conifer stems), difference in percent understory vegetation cover, and Bray-Curtis

dissimilarities of understory vegetation community. Additionally, in some figures, we indicate

sites where fires were likely to have been low-severity surface fires. This rough designation was

based on whether sites had live trees with large mean DBH (>5 cm), excluding sites where mean

height was <= 4 m, and TSLF >= 5 years, as trees could grow to this height/DBH in that time.

In order to determine whether bacterial richness was affected by short interval reburns, we

estimated the richness of bacterial communities by using the R package *breakaway* (Willis and

Bunge*,* 2015), using a weighted linear regression model (Rocchetti *et al*., 2011). To test for

differences in richness between paired SI and LI samples in the organic and mineral horizons, we

used paired *t*-tests. We then calculated the percent difference in richness estimates between

paired SI and LI sites and used an ANOVA to test for the effect of TSLF, vegetation transition,

drainage class, difference in FFI, and absolute difference in total number of stems, controlling

for soil horizon and drainage class (except for when testing drainage class itself). We also tested

whether including bacterial richness improved predictions of post-fire seedling recruitment by

using ANOVA, with live overstory stems, site moisture, TSLF, and interval as predictor

variables for the first model (as in Whitman *et al*. (2019b)), and live overstory stems, site

moisture, TSLF, interval, and bacterial richness in the second model.

In order to determine which specific taxa were associated with LI *vs*. SI sites, we estimated the

differential abundances of taxa in LI *vs*. SI sites using the R package *corncob* (Martin *et al.,*

2020) with the Wald testing procedure and a false discovery rate cutoff of 0.05, controlling for

sample pair and soil horizon, and analyzing wetlands and uplands separately. Using the

differential abundance estimates, we calculated the estimated $\log_2$-fold change in the relative

abundances of the significant taxa in the LI *vs*. SI sites. To identify responding OTUs that were

also identified in Whitman *et al.* (2019a), where we studied bacterial and fungal responses to fire

severity in the Canadian boreal forest at sites from the same region, we used BLAST (Camacho

et al., 2009). Every interval-responsive OTU from this study matched an OTU from the previous

study at 99.6-100% ID and all were classified according to their $\log_2$-fold change responses in

burned *vs*. unburned sites from the previous study: enriched in burned sites (positive), depleted in

burned sites (negative), or no significant change (neutral).

Results

Effects of short interval reburns on bacterial community composition

Short interval reburns, relative to long interval reburns, significantly affected bacterial

community composition (PERMANOVA, p = 0.002, Figure 2, Supplemental Table 1), after

controlling for pair ID. Drainage class (p = 0.001, Figure 2, Supplemental Table 2) and soil

horizon (p = 0.001, Figure 2, Supplemental Table 2) were also significant drivers of bacterial

community dissimilarities in the dataset, whereas pH was not (p = 0.28, Supplemental Table 2).

For the remaining analyses, mineral and organic soil horizons and wetlands and uplands were

analyzed separately or were controlled for by including these variables in regressions.

Figure 2. Non-metric multidimensional scaling plot of Bray-Curtis dissimilarities of the bacterial

community composition (k = 4, stress = 0.08, first two axes shown). Drainage class is identified

by fill (wetland = open symbols, upland = filled symbols), and soil horizon is identified by shape

(mineral (M) = circle, organic (O) = square). Reburn interval status is indicated by color (long =

green, short = orange). Light grey arrows connect site pairs, pointing from long to short fire-free

intervals. Symbols outlined in black indicate samples where fire was characterized as a low-

severity surface fire.

The second and fourth NMDS axes were most clearly associated with differences in LI and SI

sites. SI sites were associated with a shift away from conifer-dominated to broadleaf-dominated

and were also associated with higher soil pH (Figure 3).

Figure 3. Second and fourth axes of non-metric multidimensional scaling of Bray-Curtis dissimilarities for bacterial community composition (k = 4, stress = 0.08). Paired short and long interval sites are joined by thicker shaded arrows (green blunt end indicating LI site, orange pointed end indicating SI site). Light grey arrows indicate vectors for selected significant (p<0.05) site characteristics – pH, % conifer seedlings, and fire-free interval.

None of our factors of interest had significant effects on bacterial community dissimilarities between paired SI and LI sites. There was not a significant effect of TSLF on dissimilarities between bacterial communities from SI and LI sites (p = 0.19; Supplemental Figure 1 and Supplemental Table 3). Whether there was a vegetation transition between paired SI and LI sites was also not a significant factor driving bacterial community dissimilarities (p = 0.71; Supplemental Figures 1 and 2, Supplemental Table 3). Bacterial community dissimilarities between paired SI and LI sites were not different between wetlands and uplands (p = 0.13; Supplemental Figures 1 and 2, Supplemental Table 3). Lastly, bacterial community dissimilarities between paired SI and LI sites were not clearly affected by differences in FFI (p = 0.12; Supplemental Figure 2 and Supplemental Table 3).

Paired sites where the LI site had more total understory stems and a higher proportion of conifer

stems than the SI site had more dissimilar bacterial communities (Figure 4, Table 2).

Dissimilarities between bacterial communities in paired SI and LI sites were not affected by

difference in total understory vegetation cover (p = 0.91; Supplemental Figure 3, Supplemental

Table 4) or vegetation community dissimilarities (R^2 = -0.01, p-value = 0.96, Supplemental

Figure 4; Supplemental Table 4).

Table 2. ANOVA results for the effects of parameters of interest in individual models on
bacterial community dissimilarity after controlling for soil horizon and drainage class

Variable	Sum of Squares	F-value	p-value	R^2
Log difference in understory stems (all)	0.007	5.79	0.023 *	0.15
Log difference in understory stems (conifer)	0.056	4.47	0.043 *	0.12
Log difference in understory stems (broadleaf)	0.044	3.45	0.073	0.14

Figure 4. Bray-Curtis dissimilarities of bacterial community composition between paired short

interval (SI) and long interval (LI) reburn sites *vs*. the log difference (log(LI - SI) of a) all

understory stems per hectare, b) understory conifer stems per hectare, and c) understory

broadleaf stems per hectare. Drainage class is identified by color (wetland = light, upland =

dark), and soil horizon is identified by shape (mineral (M) = circle, organic (O) = square). Note

log scale on x-axis. Values at zero had no stems recorded at time of sampling, and therefore had

no differences in stem count. Pairs where the short interval fire was characterized as a low-

severity surface fire are indicated by a ˙ symbol, pairs where the long interval fire was

characterized as a low- severity surface fire are indicated by an * symbol, and pairs where both

fires were low- severity surface fire are indicated with a † symbol.

 Short-Interval and Long-Interval differences in richness

Richness was greater in SI sites for some pairs (up to 93% greater richness in SI sites) and lower

in SI sites for others (up to 53% lower richness in SI sites), but was not consistently significantly

different (Figure 5, paired t-test, p = 0.81, p = 0.39, p = 0.66, and p = 0.96, for upland mineral

soil, upland organic soil, wetland mineral soil, and wetland organic soil, respectively).

Figure 5. Boxplots of estimated bacterial community richness from paired (joined by grey lines)

short and long fire free interval samples in the a) upland mineral soil, b) upland organic soil, c)

wetland mineral soil, and d) wetland organic soil horizons (paired t-tests, p = 0.81, 0.39. 0.66,

and 0.96, respectively). Each point represents a sample. Points outlined in black indicate a

sample where fire was characterized as a low-severity surface fire.

None of our factors of interest had significant effects on bacterial community richness between

paired SI and LI sites. Percent change in estimated richness between paired SI and LI samples

was not associated with differences in FFI between paired sites (p = 0.08; Supplemental Table

5). Differences in estimated richness between paired SI and LI samples were also not associated

with TSLF (p = 0.80, Supplemental Table 5), vegetation transitions (p = 0.33, Supplemental

Table 5), difference in number of stems (p = 0.11, Supplemental Table 5), or drainage class (p =

0.93, Supplemental Table 5), after controlling for soil horizon and drainage class. Bacterial

richness did not improve predictions of seedling recruitment, as estimated by live understory

stem density (p = 0.32, Supplemental Table 6). After identifying differentially abundant bacteria

(results below), we also wanted to know whether including differentially abundant taxa of

interest would improve predictions of seedling recruitment. None of our top responding OTUs

(identified as *Blastococcus, Rhizobiaceae,* and *Callaberonia*) significantly improved predictions

of the recruitment of conifer and broadleaf seedlings in an ANOVA, after accounting for live

overstory stems, site moisture, TSLF, interval, and OTU abundance as predictor variables.

Differentially abundant bacteria between short interval and long intervals

We identified taxa that were differentially abundant in LI *vs*. SI sites in uplands (Figure 6a,

Supplemental Figure 6) and wetlands (Figure 6b, Supplemental Figure 5). Of the most

differentially abundant taxa, one taxon from the genus *Blastococcus* had 6.6 and 4.7 times

greater relative abundance in SI sites than LI sites, in both uplands and wetlands, respectively,

and was abundant across samples (uplands: $1.5\% \pm 1.6\%$; wetlands: $0.7\% \pm 1.2\%$). Of the taxa

that were most differentially abundant in LI sites over SI sites, for uplands, we identified a taxon

that had 3.9 times higher relative abundance in LI sites and was also relatively abundant overall

($0.75\% \pm 0.51\%$) from the family *Burkholderiaceae*. For wetlands, we identified a taxon that was

5.0 times more abundant in LI sites and was relatively abundant overall ($0.30\% \pm 0.26\%$) from

the family *Rhizobiaceae*.

Figure 6. Log$_2$-fold change in relative abundance in long *vs.* short fire-free intervals for (a)

upland and (b) wetland sites, after controlling for Site ID and soil horizon. Each point represents

a single OTU. Color and shape indicate the response the OTU had in Whitman *et al.* (2019a)

(green triangle = positive response to fire yellow circle = neutral response to fire, pink upside-

down triangle = negative response to fire,). The x-axis label indicates the finest-scale taxonomy

available. Size of points is scaled by the average relative abundance of taxa in short- and long-

interval sites. Only significantly differentially abundant taxa ($p_{FDR}<0.05$) and responses greater

than |1| are plotted. Solid line indicates no difference in relative abundance between long *vs*.

short interval sites, therefore points above the line indicate taxa that were significantly more

abundant in long-interval upland or wetland sites, and points below the line indicate taxa that

were significantly more abundant in short-interval upland or wetland sites.

Discussion

Shorter fire-free intervals alter bacterial communities, reflecting changes in vegetation and soil

properties

In addition to the clear differences in bacterial community composition between different

drainage classes and soil horizons (Figure 2), short interval reburns also shifted bacterial

community composition (Figure 3). The bacterial community changes between SI and LI were

associated with decreased conifer dominance, as well as higher soil pH (Figure 3). This echoes

the types of changes that have been seen in the aboveground communities in this region

(Whitman *et al.*, 2019b), indicating that soil bacterial community composition will also be

affected by predicted changes in this region – specifically, decreasing fire-free intervals. The

extent to which changes in the bacterial communities are being driven by changes in the

vegetation community, and vice-versa, is not possible to determine with this dataset and

experimental design. As discussed in more detail below, the presence of vegetation transitions

was not associated with greater differences in bacterial community composition between paired

SI and LI sites, nor did we find a strong association between aboveground vegetation and soil

bacterial communities. This lack of clear coupling may suggest that the primary drivers of

changes to plant and bacterial communities with shorter fire-free intervals differ. While soil pH

was not a significant predictor of community composition at the whole-dataset level after

controlling for drainage class and soil horizon, higher pH was associated with shifts in

community composition from LI to SI sites (Figures 2 and 3). This could be explained if short

interval reburns were either higher severity, increasing total combustion and ash production,

which would result in an increase in pH, particularly for combustion at higher temperatures

(Bodí et al., 2014), or just simply from the compound effects of the two most recent fires,

maintaining higher ash levels between the two. The shift toward increased soil pH in SI sites is

notable, since soil pH is regularly found to be a powerful predictor of bacterial community

composition in regional datasets (Rousk *et al*., 2010; Bahram *et al*., 2018). Given this

observation, it would be interesting to determine whether pH shifts are the primary driver of

shifts in bacterial community composition with shorter interval reburns. Since this was not our

primary question for this study, future studies could be designed to test this hypothesis.

We found that the effects of short interval reburns on paired site dissimilarities were not clearly

moderated by any of our predicted factors: TSLF, difference in FFI, vegetation transition, or the

influence of drainage class. We had predicted that increasing TSLF would decrease Bray-Curtis

dissimilarities between paired sites, as communities recover and converge on similar states post

fire. If this had been the case, it might have suggested that bacterial communities are resilient to

the effects of short interval reburns. However, our data do not indicate that increasing TSLF

allows communities to converge meaningfully, over the range of TSLF studied (1-21 years). This

is consistent with other studies of post-fire recovery in microbial communities, which have

indicated that it can take decades to over a century for soil microbial communities to return to

their pre-burn states (Dooley and Treseder, 2012; Ferrenberg *et al.,* 2013; Dove and Hart, 2017).

Thus, it may not be surprising that the effects of short interval reburns did not seem to decrease

community dissimilarity over the timescale of this study. That said, while we did not detect a

significant effect, we might speculate that such an effect could still emerge over longer

timescales. LI and SI pair dissimilarities at longer TSLF (12-21 years) ranged from 0.46-0.78

(more similar), while dissimilarities at shorter TSLF (1-2 years) ranged from 0.58 – 0.91 (less

similar). While the ranges overlap, they do suggest a trajectory towards more similar

communities after longer times since last fire. Future studies might be designed to target an even

broader distribution of TSLF across the SI-LI pairs to directly test this question, or to trace these

same sites over time. However, such a sample set may be difficult to collect, as it becomes

increasingly difficult to conclusively identify FFI as we move beyond the range of modern

satellite data records.

Since SI *vs*. LI was a significant predictor of bacterial community composition (Figures 2 and 3),

we expected that communities might become increasingly dissimilar as differences in FFI

increased. As differences in FFI increase, factors that could affect microbial community

composition, such as soil properties or vegetation composition, may also become increasingly

different, extending microbial community dissimilarities between the paired sites. However,

larger differences in FFI were not associated with more dissimilar communities. This suggests

that the differences in FFI do not scale consistently with differences in the direct or indirect

effects of fire on microbial community composition. This may also point to the range of possible

outcomes from a short interval fire. As a simplified example, there can be high-severity short

interval fires where a recent burn has left high fuel loads, supporting significant loss of biomass

and degradation of germination substrate, and leading to poor conifer generation. At the other

extreme, there can also be low-severity short interval fires where frequent recent fires lead to

surface fires with less tree mortality and SOM combustion, and safe germination sites. Thus, we

might expect there to be no consistent association with increasing differences in FFI and

bacterial community dissimilarity, as was observed here.

Vegetation can structure microbial communities through a wide range of mechanisms, such as

interactions in the rhizosphere or through differences in litter inputs (Berg and Smalla, 2009;

Merilä *et al*., 2010), so we had predicted that paired sites where differences between SI and LI

sites were so great that leading tree species were different ("Vegetation Transition") would also

be associated with greater differences in bacterial community composition. Counter to our

predictions, differences in the leading tree species between paired SI and LI sites were not

associated with more dissimilar bacterial communities. Furthermore, differences in total

vegetation cover or vegetation community dissimilarities were also not associated with bacterial

community dissimilarity. These findings are somewhat consistent with our previous work in the

region (Whitman *et al*., 2019a), where we found that understory vegetation and microbial

community dissimilarities were not related for upland soils, which dominate this dataset. They

are also consistent with the findings of Adkins et al. (2020) in a Sierra Nevada mixed conifer

forest, who suggested that post-wildfire vegetation effects on community-level soil microbial

properties were largely indirect and mediated through soil properties. However, when we

considered a more sensitive measure of change in the understory vegetation community -

difference in understory tree stem counts - this factor was significantly associated with

increasing bacterial community dissimilarities (Table 2 and Figure 4). This metric allows us to

discern sites with differences in tree regeneration, even if the leading species did not change.

Differences in tree regeneration could have important effects on local soil conditions, including

temperature and moisture, as sparsely vegetated sites or sites with more seedlings than mature

trees may be warmer and drier (Hoecker *et al.*, 2020). This finding raises the question, are

bacterial communities different because there was more regeneration, or was tree regeneration

more successful because of the bacterial community? Future experiments could be designed to

investigate this question directly.

Although we expected that microbial communities in wetlands would be buffered from potential

effects of short interval reburns more than uplands, we did not see an effect of drainage class on

community dissimilarities between paired LI and SI sites. Because sites were sampled more than

one year post fire, we were not able to generate accurate estimates of fire severity, which is

known to affect bacterial communities (Whitman *et al.*, 2019a; Day *et al.*, 2019). Our simplified

metric for inferring low-severity surface fires does not apply well here either, since trees are

generally less abundant in the wetland sites. However, if some wetlands had high burn severities

for short interval reburns, then we would not have expected our hypothesis to be supported. To

examine this possibility, we can consider two contrasting wetland pairs, which were both

sampled 8 years post fire. The first pair has the greatest dissimilarity between SI and LI

communities (Bray-Curtis dissimilarity 0.901 in the mineral soil and 0.827 in the organic soil).

For this pair, the organic horizon depth in SI was 88.45% lower than LI, and the dominant tree

type transitioned from white spruce to balsam poplar. These metrics may suggest a higher

severity fire in the SI. In contrast, the organic and mineral soil communities of the second pair

were more similar (Bray-Curtis dissimilarities of 0.699 and 0.575, in mineral and organic

horizons, respectively); in this pair, the organic depth in SI was only 19.45% lower than LI, and

the leading tree type did not change (trembling aspen-mixed forest). These metrics may suggest a

lower severity fire in SI compared to the first pair. The possibility that the first pair experienced a

relatively high severity fire in the SI (indicated by a greater difference in O horizon thickness and

change in leading tree type) may explain why the paired samples are more dissimilar, and would

be counter to our hypothesis that the wetlands would experience lower severity fires across the

board, leading to smaller effects of short interval reburns. Relative to severely burned SI uplands,
the biogeophysical differences between SI and LI wetlands may be lower (*e.g.*, thicker residual
organic horizons and greater post-fire conifer dominance; Whitman *et al.* 2019b), but repeated
wildfires nonetheless produce meaningful differences in both above and belowground
communities.

Bacterial richness is not consistently affected by shorter fire free intervals

We expected that bacterial richness would not be clearly affected by short interval reburns,
largely because soil bacterial communities are extremely rich to begin with and, despite advances
in sequencing technologies, methods for accurate richness estimation in high-throughput
amplicon sequencing datasets remain limited (Willis, 2019). This expectation was supported -
although most LI sites were richer than their SI pairs (Figure 5), this difference was not
significant and the percent difference in richness ranged widely - from 93% higher (SI richer
than LI) to 53% lower (LI richer than SI). Although fires often reach temperatures at the soil
surface that would be expected to kill all bacteria (Pingree and Kobziar, 2019), heating attenuates
rapidly with depth in the soil, meaning that lethal temperatures in the surface of the soil may be
accompanied by almost no changes just a few centimeters below the surface. Thus, it may not be
surprising that the effects of fire on bacterial richness have been inconsistent across studies, with
some studies reporting a decrease in bacterial richness with increasing fire severity (Sáenz de
Miera *et al.,* 2020), or reporting a lack of significant changes (Pressler *et al.* 2019; Whitman *et*
al., 2019a). Our findings indicate that, although SI *vs*. LI have different bacterial community
compositions (Figures 2 and 3), this difference is likely driven more by changes in the overall
community structure or by a few community members, rather than significant differences in
richness. While richness has often been linked to ecosystem multifunctionality (Delgado-
Baquerizo *et al.*, 2017) or resistance and resilience (Shade *et al.*, 2012), these relationships are
not always straightforward in soil communities, which generally have high functional
redundancy (Nunan et al., 2018). This functional redundancy is perhaps also reflected in our
finding that including bacterial richness in our models did not improve predictions of understory
stem regeneration. Thus, even for the paired sites where bacterial richness was lower in SI than
LI, we should not necessarily predict functional limitations.

Short- and long-interval communities have distinct responding bacteria

While the fire ecology of plants, and even many fungi, is well-characterized and has long been

studied (Cooper, 1961; Seaver, 1909), the bacterial equivalents of fire responder species like

Pinus banksiana or the fungus *Pyronema* are only just being established. By identifying specific

bacteria associated with SI or LI sites, we sought to both expand our understanding of

fundamental questions about fire ecology for bacteria, as well as to offer potential hypotheses

about the effects that changes to soil bacterial communities might have on ecosystem

functioning. For example, taxa that are more abundant in SI sites may be able to thrive under the

conditions characteristic of frequently burned areas. Thus, we might predict these microbes

would exhibit traits that allow them to consume pyrogenic organic matter more effectively,

thrive in areas that are more drought-prone, or be common in areas with less vegetation cover

and tree regeneration. Taxa that are more abundant in LI sites could be general fire responders

that are characteristic of normal fire regimes; these taxa may thrive in post-fire environments,

just not when their habitat is significantly altered by short interval reburns.

Even though our study sites spanned a wide range of post fire conditions, we were able to

identify several taxa that were consistently differentially abundant in SI and LI sites. Because of

this experimental design, we believe that these observed responses may be robust within this

region, rather than site-specific anomalies. Our most prominent responder was a *Blastococcus*

OTU, which was the most abundant responding OTU in both wetlands and in uplands (Figure 6).

It was over four times more abundant in SI soils than in LI soils, and consistently represented a

large proportion of the total community across sites. This was the same OTU (100% identical

over the sequenced region) as an OTU in Whitman *et al.* (2019a) that was enriched in burned

sites and was increasingly abundant in sites with increasing fire severity. In holm-oak forests of

Spain, members of *Blastococcus* also increased in abundance after wildfires in rhizosphere soils

versus neighboring unburned areas (Fernández-González *et al.*,2017; Cobo-Diaz *et al.*, 2015).

The sequence for this taxon was also 100% identical to a recently classified *Blastococcus deserti*

sp. isolated from a desert sample (Yang *et al.,* 2019). While clearly a fundamentally different

ecosystem, in their study, Yang *et. al.* noted that this isolate was able to survive at temperatures

up to 50 °C and utilize D-salacin as a C source, which is a compound found in the bark of

Populus and *Salix* species (Palo, 1984), when other related *Blastococcus* strains in their study

could not. Both of these traits offer mechanisms that could help this taxon thrive in the post-fire

environment: its potential ability to survive high temperatures during fires, and also metabolize

compounds characteristic of the organic horizon of an aspen-dominated forest – a common

successional species after short interval reburns in this region (Whitman *et al*., 2019b; Johnstone

and Chapin, 2006). Together, these studies highlight traits that allow us to propose this

Blastococcus OTU as a classical "pyrophilous" bacterium, which thrives in burned soils and

even becomes increasingly abundant with more frequent and more severe burns.

For wetland sites, the most prominent LI associated taxon was classified as belonging to the

family *Rhizobiaceae*. This OTU was detected in the Whitman *et al*. (2019a) study, but did not

show a negative or positive response to fire in that dataset. Members of family *Rhizobiaceae* are

often associated with nitrogen fixation (Spaink, Kondorosi, and Hooykaas, 1998), which could

be an important source of N in post-fire ecosystems (Smithwick et al., 2005). Some members of

the *Rhizobiaceae* family also perform other steps of the N cycle, including denitrification (Rich

et al., 2003). While these roles can be seen to be relevant post-fire, it is not immediately clear

why these taxa were more enriched in the LI wetland sites. One possibility is that if the SI

wetland sites were drier, then the LI sites may have had conditions more conducive to

denitrification (low oxygen due to high moisture, and high organic matter (Martínez-Espinosa *et*

al., 2021)) that selected for these taxa. Similarly, N fixation could also be more likely to be

supported under these conditions: it is a highly energy intensive process requiring both lots of C

for energy and at least locally anoxic conditions to protect the oxygen-sensitive nitrogenase

enzyme (Smercina *et al*., 2019). However, even though this *Rhizobiaceae* OTU is 100% identical

over the sequenced region to taxa that are known to be able to perform these functions, there may

still be important functional differences. We should consider these speculations on their role in

post-fire N cycling at LI sites as hypotheses for future testing.

For upland sites, the most prominent LI associated taxon, identified as being from the

Burkholderiaceae family, also matched an OTU in Whitman *et al.* (2019a) that was found to be

more abundant in burned sites, to the point of not being detected at any unburned sites. Upon

first consideration, this seems somewhat surprising – why would a positive burn responder be
more abundant at LI burned sites, whose soil properties would presumably be less influenced by
fire? In considering this question, it is important to remember that all of the sites in this study
were burned, so we should not necessarily expect that LI-associated taxa should also be
classified as negative burn responders. Furthermore, the dataset in Whitman *et al*. (2019a)
spanned a wide range of burned sites, including both long and short interval reburns across a
range of severity and vegetation communities. One possible explanation for this association
could be that members of the genus *Burkholderia*, from this same family, have been found to be
more abundant in lower pH soils (Stopnisek *et al*., 2014), and we observed here that short
interval reburns were associated with higher pH (Figure 3). Supporting this observation, the
corresponding OTU from Whitman *et al*. (2019a) clustered in a module associated with fire-
responsive taxa that were more abundant at lower pH sites in a co-occurrence network.
638
This LI-associated organism may also play an important role in post-fire plant establishment.
While the finest-scale matching taxonomy in the SILVA database placed this LI-associated OTU
within the *Burkholderiaceae* family, using NCBI BLAST, we found it is a 100% ID match for a
Caballeronia sordidicola that was isolated from a lichen in Svalbard Archipelago (AF512827.1;
Kim *et al*., 2017) and for a *Caballeronia sordidicola* spruce tree endophyte from a sub-boreal
forest in British Columbia, Canada (MG561776.1; Puri *et al*., 2018). Furthermore, the authors of
this second study identified that this isolate has the capacity to fix nitrogen (Puri *et al*., 2018),
readily colonized pine and spruce seedlings, increasing their biomass production 4-7-fold (Puri *et
al*., 2020a) and provided more than 50% of both lodgepole pine (*Pinus contorta*) and white
spruce seedlings' N requirements (Puri *et al*., 2020b). This is strongly consistent with our
observation that conifer seedlings were more abundant at LI sites. While our study design does
not allow us to separate the cause and effect between the increased presence of putative
endosymbiont diazotroph *Callaberonia sordidicola* and more abundant conifer seedlings at LI
sites, it clearly raises pressing questions. For example, linking our observations, we might ask
whether increases in pH due to short interval reburns shift the soil environment away from
conditions that are optimal for *Callaberonia sordidicola*, exacerbating other factors that
contribute to poor conifer seedling establishment, such as depleted seed banks or suboptimal
seedbed conditions associated with short interval reburns. Certainly, designing experiments to

probe the ecological implications of the depletion of this potentially highly influential spruce and

pine endosymbiont in short interval reburns would be an important next step for this system.

Conclusions

Short interval reburns result in altered soil bacterial communities. These altered short-interval

communities were associated with sites that had fewer understory stems, a lower percentage of

conifer seedlings, and higher pH. These findings suggest that, as has been seen in plant

communities, bacterial communities are also likely to change as fire-free intervals decrease.

Although there were changes in overall community structure, we did not see consistent effects of

short interval reburns on bacterial richness. We did, however, identify specific taxa that were

characteristic of SI *vs.* LI sites, including the abundant and strongly SI-associated

Blastococcus OTU. This common successional taxon in uplands as well as wetlands after short

interval reburns is a general fire-responder in this region and in other parts of the world.

Beginning to parse its specific fire ecology, future experiments could determine whether this is

due to an ability to survive higher temperatures and an ability to metabolize compounds

characteristic of the organic horizon of an aspen-dominated forest. Additionally, the depletion of

diazotrophic conifer endophyte *Callaberonia sordidicola* in short interval reburn sites raises

questions about whether its depletion is contributing to – or merely reflects – poor conifer

seedling recolonization post-fire at short-interval reburns.

There are numerous other future directions for this work. First, directly investigating some of the

hypotheses generated here is of interest, including the extent to which bacterial community shifts

due to shorter fire-free intervals are driven by changes in soil pH. Research is needed to elucidate

the relative influence of post-fire microbial communities in determining vegetation community

development and facilitating or impeding tree recruitment and, conversely, further research is

needed to identify the influence of post-fire vegetation on soil microbial community

development and the strength of these interacting effects. Additionally, while identifying specific

fire-responsive taxa is a critical first step in developing a fire ecology framework for bacteria, the

next steps will be to investigate the functional traits of these responding taxa in a mechanistic

way, in order to further understand their individual ecology and what effects changes in their
populations might have on ecosystem processes.

Acknowledgements

The field campaign in which these data were gathered was funded by the government of the
Northwest Territories, and the Natural Sciences and Engineering Research Council of Canada
(Funding Reference Number: CGSD3-471480-2015). Environment and Climate Change Canada,
Sam Haché, Parks Canada Agency, and Jean Morin provided in-kind support. Matthew Coyle,
Kathleen Groenwegen, Xianli Wang, Mary Stephens, Scott L. Stephens, Rodrigo Campos-Ruiz,
and Josh Gauthier provided indispensable help in the field. This research was performed using
the computing resources and assistance of the UW-Madison Center for High Throughput
Computing (CHTC) in the Department of Computer Sciences. The CHTC is supported by UW-
Madison, the Advanced Computing Initiative, the Wisconsin Alumni Research Foundation, the
Wisconsin Institutes for Discovery, and the National Science Foundation, and is an active
member of the Open Science Grid, which is supported by the National Science Foundation and
the US Department of Energy's Office of Science. T.W. and J. W. were partially supported by
the U.S. Department of Energy (DE-SC0020351). The authors have no conflicts of interest to
declare.

Supplemental Information

Supplemental information associated with this study is available. Code used for all analyses can
be found at https://github.com/WhitmanLab/WoodBuffalo2016. All sequences are deposited in
the NCBI SRA under accession numbers [to be submitted upon paper acceptance].

712
713

References

Adkins, J., Docherty, K. M., Gutknecht, J. L. M., & Miesel, J. R. (2020). How do soil microbial
communities respond to fire in the intermediate term? Investigating direct and indirect
effects associated with fire occurrence and burn severity. *Science of the Total Environment*,
745, 140957. https://doi.org/10.1016/j.scitotenv.2020.140957

Allison, S. D. & Martiny, J. B. H. (2009). Resistance, resilience, and redundancy in microbial
communities. *PNAS*, *105(S1)*, 11512-11519. https://doi.org/10.1073/pnas.0801925105

Bardgett, R. D. & Van Der Putten, W. H. (2014). Belowground biodiversity and ecosystem
functioning. *Nature*, *515*(7528), 505–511. https://doi.org/10.1038/nature13855

Berg, G. & Smalla, K. (2009). Plant species and soil type cooperatively shape the structure and
function of microbial communities in the rhizosphere. *FEMS Microbiology Ecology*, *68*, 1-
13. https://doi.org/10.1111/j.1574-6941.2009.00654.x

Bradshaw, C. J. A., & Warkentin, I. G. (2015). Global estimates of boreal forest carbon stocks
and flux. *Global and Planetary Change*, *128*, 24–30.
https://doi.org/10.1016/j.gloplacha.2015.02.004

Bray, J. R. & Curtis, J. T. (1957). An ordination of upland forest communities of southern
Wisconsin. *Ecological Monographs, 27*, 325-349. https://doi.org/10.2307/1942268

Brandt, J. P., Flannigan, M. D., Maynard, D. G., Thompson, I. D., & Volney, W. J. A. (2013).
An introduction to Canada's boreal zone: Ecosystem processes, health, sustainability, and
environmental issues1. *Environmental Reviews*, *21*(4), 207–226. https://doi.org/10.1139/er-
2013-0040

Brandt, J. P. (2009). The extent of the North American boreal zone. *Environmental Reviews*, *17*,
101–161. https://doi.org/10.1139/A09-004

Bogdanski, B. E. C. (2008). *Canada's boreal forest economy: economic and socio-economic*
issues and research opportunities. (Issue BC-X-414).

Bokulich, N.A., Kaehler, B.D., Rideout, J.R. *et al.* (2018). Optimizing taxonomic classification
of marker-gene amplicon sequences with QIIME 2's q2-feature-classifier plugin.
Microbiome, *6*, 90. https://doi.org/10.1186/s40168-018-0470-z

Callahan, B.J., McMurdie, P.J., Rosen, M.J., Han, A.W., Johnson, A.J.A., & Holmes, S.P.
(2016). DADA2: high-resolution sample inference from Illumina amplicon data. *Nature*
Methods 13, 581–583. https://doi.org/10.1038/nmeth.3869

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopouls, J., Bealer, K., & Madden, T.L.
(2009). BLAST+: architecture and applications. *BMC Bioinformatics*, *10*, 421.
https://doi.org/ 10.1186/1471-2105-10-421

Chandra, L. R., Gupta, S., Pande, V., & Singh, N. (2016). Impact of forest vegetation on soil
characteristics: a correlation between soil biological and physico-chemical properties. *3
Biotech*, *6*(2), 1–12. https://doi.org/10.1007/s13205-016-0510-y

Cobo-Díaz, J. F., Fernández-González, A. J., Villadas, P. J., Robles, A. B., Toro, N., &
Fernández-López, M. (2015). Metagenomic Assessment of the Potential Microbial Nitrogen
Pathways in the Rhizosphere of a Mediterranean Forest After a Wildfire. *Microbial
Ecology*, *69*(4), 895–904. https://doi.org/10.1007/s00248-015-0586-7

Cooper C.F. (1961). The Ecology of Fire. *Scientific American*, *204*, 150-160.

Day, N. J., Dunfield, K. E., Johnstone, J. F., et al. (2019). Wildfire severity reduces richness and
alters composition of soil fungal communities in boreal forests of western Canada. *Global
Change Biol*ogy, *25*, 2310– 2324. https://doi.org/10.1111/gcb.14641

Delgado-Baquerizo, M., Eldridge, D. J., Ochoa, V., Gozalo, B., Singh, B. K., & Maestre, F. T.
(2017). Soil microbial communities drive the resistance of ecosystem multifunctionality to
global change in drylands across the globe. *Ecology Letters*, *20*(10), 1295–1305.
https://doi.org/10.1111/ele.12826

Dooley, S. R. & Treseder, K. K. (2012). The effect of fire on microbial biomass: A meta-analysis
of field studies. *Biogeochemistry*, *109*(1–3), 49–61. https://doi.org/10.1007/s10533-011-
9633-8

Dove, N. C., Safford, H. D., Bohlman, G. N., Estes, B. L., & Hart, S. C. (2020). High-severity
wildfire leads to multi-decadal impacts on soil biogeochemistry in mixed-conifer forests.
Ecological Applications, *30*(4), 1–18. https://doi.org/10.1002/eap.2072

Fernández-González, A. J., Martínez-Hidalgo, P., Cobo-Díaz, J. F., Villadas, P. J., Martínez-
Molina, E., Toro, N., Tringe, S. G., & Fernández-López, M. (2017). The rhizosphere
microbiome of burned holm-oak: Potential role of the genus Arthrobacter in the recovery of
burned soils. *Scientific Reports*, *7*(1), 1–12. https://doi.org/10.1038/s41598-017-06112-3

Ferrenberg, S., O'Neill, S. P., Knelman, J. E., Todd, B., Duggan, S., Bradley, D., Robinson, T.,
Schmidt, S. K., Townsend, A. R., Williams, M. W., Cleveland, C. C., Melbourne, B. A.,
Jiang, L., & Nemergut, D. R. (2013). Changes in assembly processes in soil bacterial
communities following a wildfire disturbance. *ISME Journal*, *7*(6), 1102–1111.
https://doi.org/10.1038/ismej.2013.11

Gill, N. S., Sangermano, F., Buma, B., & Kulakowski, D. (2017). Populus tremuloides seedling
establishment: An underexplored vector for forest type conversion after multiple
disturbances. *Forest Ecology and Management*, *404*(May), 156–164.
https://doi.org/10.1016/j.foreco.2017.08.008

Hanes, C. C., Wang, X., Jain, P., Parisien, M. A., Little, J. M., & Flannigan, M. D. (2019). Fire-regime changes in canada over the last half century. *Canadian Journal of Forest Research*, *49*(3), 256–269. https://doi.org/10.1139/cjfr-2018-0293

Hart, S. C., DeLuca, T. H., Newman, G. S., MacKenzie, M. D., & Boyle, S. I. (2005). Post-fire vegetative dynamics as drivers of microbial community structure and function in forest soils. *Forest Ecology and Management*, *220*(1–3), 166–184. https://doi.org/10.1016/j.foreco.2005.08.012

Hoecker, T. J., Hansen, W. D., & Turner, M. G. (2020). Topographic position amplifies consequences of short-interval stand-replacing fires on postfire tree establishment in subalpine conifer forests. *Forest Ecology and Management*, *478*(August), 118523. https://doi.org/10.1016/j.foreco.2020.118523

Holden, S. R., Berhe, A. A., & Treseder, K. K. (2015). Decreases in soil moisture and organic matter quality suppress microbial decomposition following a boreal forest fire. *Soil Biology and Biochemistry*, *87*, 1–9. https://doi.org/10.1016/j.soilbio.2015.04.005

Holden, S. R. & Treseder, K. K. (2013). A meta-analysis of soil microbial biomass responses to forest disturbances. *Frontiers in Microbiology*, *4*(June), 1–17. https://doi.org/10.3389/fmicb.2013.00163

Hoy, E. E., Turetsky, M. R., & Kasischke, E. S. (2016). More frequent burning increases vulnerability of Alaskan boreal black spruce forests. *Environmental Research Letters*, *11*(9). https://doi.org/10.1088/1748-9326/11/9/095001

Jain, P., Wang, X., & Flannigan, M. D. (2017). Trend analysis of fire season length and extreme fire weather in North America between 1979 and 2015. *International Journal of Wildland Fire*, *26*(12), 1009–1020. https://doi.org/10.1071/WF17008

Johnstone, J. F. & Chapin, F. S. III (2006). Fire interval effects on successional trajectory in boreal forests of northwest Canada. *Ecosystems*, *9*, 268–277. https://doi.org/10.1007/s10021-005-0061-2

Johnstone, J. F., Allen, C. D., Franklin, J. F., Frelich, L. E., Harvey, B. J., Higuera, P. E., Mack, M. C., Meentemeyer, R. K., Metz, M. R., Perry, G. L. W., Schoennagel, T., & Turner, M. G. (2016). Changing disturbance regimes, ecological memory, and forest resilience. *Frontiers in Ecology and the Environment*, *14*(7), 369–378. https://doi.org/10.1002/fee.1311

Kasischke, E.S., O'Neill, K.P., French, N.H.F., & Bourgeau-Chavez, L.L., (2000). Controls on patterns of biomass burning in Alaskan boreal forests. In: Kasischke, E.S., Stocks, B.J. (Eds.), Fire, Climate Change, and Carbon Cycling in the Boreal Forest. Springer, New York, New York, USA, pp. 173–196.

Kelly, R., Chipman, M. L., Higuera, P. E., Stefanova, I., Brubaker, L. B., & Hu, F. S. (2013).
Recent burning of boreal forests exceeds fire regime limits of the past 10,000 years.
Proceedings of the National Academy of Sciences of the United States of America, *110*(32),
13055–13060. https://doi.org/10.1073/pnas.1305069110

Kim, J., Kwon, K. K., Kim, B. K., Hong, S. G. & Oh H.-M. (2017). Genome sequence of
Caballeronia sordidicola strain PAMC 26592 isolated from an arctic lichen species. *Korean*
Journal of Microbiology, 53: 64-66.

Knelman, J. E., Graham, E. B., Trahan, N. A., Schmidt, S. K., & Nemergut, D. R. (2015). Fire
severity shapes plant colonization effects on bacterial community structure, microbial
biomass, and soil enzyme activity in secondary succession of a burned forest. *Soil Biology*
and Biochemistry, *90*, 161–168. https://doi.org/10.1016/j.soilbio.2015.08.004

Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K., & Schloss, P. D. (2013).
Development of a dual-index sequencing strategy and curation pipeline for analyzing
amplicon sequence data on the MiSeq Illumina sequencing platform. *Applied and*
Environmental Microbiology, *79*, 5112–5120. https://doi.org/10.1128/AEM.01043-13

Larsen, C. P. S. & MacDonald, G. M. (1998). An 840-Year Record of Fire and Vegetation in a
Boreal White Spruce Forest. *Ecology*, *79*, 106–118. https://doi.org/10.1890/0012-
9658(1998)079[0106:AYROFA]2.0.CO;2

Larsen, C. P. S. (1997). Spatial and temporal variations in boreal forest fire frequency in northern
Alberta. *Journal of Biogeography*, *24*(5), 663–673. https://doi.org/10.1111/j.1365-
2699.1997.tb00076.x

Martin, B. D., Witten, D. & Willis, A. D. (2020). Modeling microbial abundances and dysbiosis
with beta-binomial regression. *Annals of Applied Statistics*, *14*(1), 94-115.
https://doi.org/10.1214/19-AOAS1283

Martínez-Espinosa, C., Sauvage, S., Al Bitar, A., Green, P.A., Vörösmarty, C.J. & Sánchez-
Pérez, J.M. (2021). Denitrification in wetlands: A review towards a quantification at global
scale. *Science of the Total Environment*, *754*, 142398.
https://doi.org/10.1016/j.scitotenv.2020.142398

McMurdie, P. J. & Holmes, S. (2013). Phyloseq: an R package for reproducible interactive
analysis and graphics of microbiome census data. *PLoS One*, *8*, e61217.
https://doi.org/10.1371/journal.pone.0061217

Merilä, P., Malmivaara-Lämsä, M., Spetz, P., Stark, S., Vierikko, K., Derome, J., & Fritze, H.
(2010). Soil organic matter quality as a link between microbial community structure and
vegetation composition along a successional gradient in a boreal forest. *Applied Soil*
Ecology, *46*, 259-267. https://doi.org/10.1016/j.apsoil.2010.08.003

Nunan, N., Leloup, J., Ruamps, L. X. O. S., Pouteau, V. X. R., & Chenu, C. (2017). Effects of habitat constraints on soil microbial community function. *Scientific Reports*, *7*, 4280. https://doi.org/10.1038/s41598-017-04485-z

Oksanen, J., Blanchet, F.G., Kindt, R., Legendre, P., Minchin, P.R., OHara, R.B., et al., (2018). Vegan: Community Ecology Package.

Palo, R.T. (1984). Distribution of birch (*Betula* SPP.), willow (*Salix* SPP.), and poplar (*Populus* SPP.) secondary metabolites and their potential role as chemical defense against herbivores. *Journal of Chemical Ecology*, *10*, 499–520. https://doi.org/10.1007/BF00988096

Parks, S. A., Parisien, M. A., Miller, C., Holsinger, L. M., & Baggett, L. S. (2018). Fine-scale spatial climate variation and drought mediate the likelihood of reburning. *Ecological Applications*, *28*(2), 573–586. https://doi.org/10.1002/eap.1671

Pellegrini, A. F. A. & Jackson, R. B. (2020). The long and short of it: A review of the timescales of how fire affects soils using the pulse-press framework. In *Advances in Ecological Research* (1st ed., Vol. 62). Elsevier Ltd. https://doi.org/10.1016/bs.aecr.2020.01.010

Pellegrini, A. F. A., Hobbie, S. E., Reich, P. B., Jumpponen, A., Brookshire, E. N. J., Caprio, A. C., Coetsee, C., & Jackson, R. B. (2020). Repeated fire shifts carbon and nitrogen cycling by changing plant inputs and soil decomposition across ecosystems. *Ecological Monographs*, *90*(4), 1–20. https://doi.org/10.1002/ecm.1409

Peterson, G. D. (2002). Contagious disturbance, ecological memory, and the emergence of landscape pattern. *Ecosystems*, *5*(4), 329–338. https://doi.org/10.1007/s10021-001-0077-1

Pingree, M. R. A. & Kobziar, L. N. (2019). The myth of the biological threshold: A review of biological responses to soil heating associated with wildland fire. *Forest Ecology and Management*, *432*, 1022–1029. https://doi.org/10.1016/j.foreco.2018.10.032

Pressler, Y., Moore, J. C., & Cotrufo, M. F. (2019). Belowground community responses to fire: meta-analysis reveals contrasting responses of soil microorganisms and mesofauna. *Oikos*, *128*(3), 309–327. https://doi.org/10.1111/oik.05738

Puri, A., Padda, K. P., & Chanway, C. P. (2018). Evidence of endophytic diazotrophic bacteria in lodgepole pine and hybrid white spruce trees growing in soils with different nutrient statuses in the West Chilcotin region of British Columbia, Canada. *Forest Ecology and Management*, *430*, 558-565. https://doi.org/10.1016/j.foreco.2018.08.049

Puri, A., Padda, K. P., & Chanway, C. P. (2020a). In vitro and in vivo analyses of plant-growth-promoting potential of bacteria naturally associated with spruce trees growing on nutrient-poor soils. *Applied Soil Ecology*, *149*(January), 103538. https://doi.org/10.1016/j.apsoil.2020.103538

Puri, A., Padda, K. P., & Chanway, C. P. (2020b). Can naturally-occurring endophytic nitrogen-fixing bacteria of hybrid white spruce sustain boreal forest tree growth on extremely nutrient-poor soils? *Soil Biology and Biochemistry*, *140*(July 2019), 107642. https://doi.org/10.1016/j.soilbio.2019.107642

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., & Glöckner, F.O., (2013). The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acids Research*, *41*(D1), D590–D596. https://doi.org/10.1093/nar/gks1219

R Core Team, 2019. R: A Language and Environment for Statistical Computing.

Ribeiro-Kumara, C., Köster, E., Aaltonen, H., & Köster, K. (2020). How do forest fires affect soil greenhouse gas emissions in upland boreal forests? A review. *Environmental Research*, *184*(February), 109328. https://doi.org/10.1016/j.envres.2020.109328

Rich, J. J., Heichen, R. S., Bottomley, P. J., Cromack, K., & Myrold, D. D. (2003). Community Composition and Functioning of Denitrifying Bacteria from Adjacent Meadow and Forest Soils. *Applied and Environmental Microbiology*, *69*(10), 5974 LP – 5982. https://doi.org/10.1128/AEM.69.10.5974-5982.2003

Rocchetti, I., Bunge, J. and Bohning, D. (2011). Population size estimation based upon ratios of recapture probabilities. *Annals of Applied Statistics*, *5*(2B), 1512-1533. http://dx.doi.org/10.1214/10-AOAS436

Rowe, J. S., & Scotter, G. W. (1973). Fire in the boreal forest. *Quaternary Research*, *3*(3), 444–464. https://doi.org/10.1016/0033-5894(73)90008-2

Sáenz de Miera, L. E., Pinto, R., Gutierrez-Gonzalez, J. J., Calvo, L., & Ansola, G. (2020). Wildfire effects on diversity and composition in soil bacterial communities. *Science of the Total Environment*, *726*. https://doi.org/10.1016/j.scitotenv.2020.138636

Saleem, M., Hu, J., & Jousset, A. (2019). More Than the Sum of Its Parts: Microbiome Biodiversity as a Driver of Plant Growth and Soil Health. *Annual Review of Ecology, Evolution, and Systematics*, *50*, 145–168. https://doi.org/10.1146/annurev-ecolsys-110617-062605

Seaver, F.J. (1909). Studies in Pyrophilous Fungi: I. The Occurrence and Cultivation of *Pyronema*. *Mycologia, 1*, 131–139.

Shade, A., Peter, H., Allison, S. D., Baho, D. L., Berga, M., Bürgmann, H., Huber, D. H., Langenheder, S., Lennon, J. T., Martiny, J. B. H., Matulich, K. L., Schmidt, T. M., & Handelsman, J. (2012). Fundamentals of microbial community resistance and resilience. *Frontiers in Microbiology*, *3*(Dec), 1–19. https://doi.org/10.3389/fmicb.2012.00417

Smercina, D. N., Evans, S. E., Friesen, M. L., & Tiemann, L. K. (2019). To fix or not to fix:
Controls on free-living nitrogen fixation in the rhizosphere. *Applied and Environmental*
Microbiology, *85*, e02546-18. https://doi.org/10.1128/AEM.02546-18

Stocks, B. J. et al. Boreal forest fire regimes and climate change. In: Remote sensing and climate
modelling: synergies and limitations (Eds Beniston, M. & Verstraete, M. M.) 233–246
(Kluwer Academic, 2001).

Stopnisek, N., Bodenhausen, N., Frey, B., Fierer, N., Eberl, L. and Weisskopf, L. (2014),
Preference of *Burkholderia* sp. for acid soils. *Environmental Microbiology*, *16*, 1503-1512.
https://doi.org/10.1111/1462-2920.12211

Van Der Heijden, M. G. A., Bardgett, R. D., & Van Straalen, N. M. (2008). The unseen majority:
Soil microbes as drivers of plant diversity and productivity in terrestrial ecosystems.
Ecology Letters, *11*(3), 296–310. https://doi.org/10.1111/j.1461-0248.2007.01139.x

Veraverbeke, S., Rogers, B. M., Goulden, M. L., Jandt, R. R., Miller, C. E., Wiggins, E. B., &
Randerson, J. T. (2017). Lightning as a major driver of recent large fire years in North
American boreal forests. *Nature Climate Change*, *7*(7), 529–534.
https://doi.org/10.1038/nclimate3329

Walker, X. J., Baltzer, J. L., Cumming, S. G., Day, N. J., Ebert, C., Goetz, S., Johnstone, J. F.,
Potter, S., Rogers, B. M., Schuur, E. A. G., Turetsky, M. R., & Mack, M. C. (2019).
Increasing wildfires threaten historic carbon sink of boreal forest soils. *Nature*, *572*(7770),
520–523. https://doi.org/10.1038/s41586-019-1474-y

Walters, W., Hyde, E. R., Berg-Lyons, D., Ackermann, G., Humphrey, G., Parada, A., et al.,
(2015). Improved bacterial 16S rRNA gene (V4 and V4-5) and fungal internal transcribed
spacer marker gene primers for microbial community surveys. *mSystems*, *1*, e00009–15.
https://doi.org/10.1128/mSystems.00009-15

Whitman, T., Whitman, E., Woolet, J., Flannigan, M. D., Thompson, D. K., & Parisien, M. A.
(2019a). Soil bacterial and fungal response to wildfires in the Canadian boreal forest across
a burn severity gradient. *Soil Biology and Biochemistry*, *138*(April), 107571.
https://doi.org/10.1016/j.soilbio.2019.107571

Whitman, E., Parisien, M. A., Thompson, D. K., & Flannigan, M. D. (2019b). Short-interval
wildfire and drought overwhelm boreal forest resilience. *Scientific Reports*, *9*(1), 1–12.
https://doi.org/10.1038/s41598-019-55036-7

Wickham, H. (2016). ggplot2: elegant graphics for data analysis. http://ggplot2.org.

Wickham, H., François, R., Henry, L., Müller, K. (2019). Dplyr: a grammar of data
manipulation. https://CRAN.R-project.org/package=dplyr.

Willis, A. & Bunge, J. (2015). Estimating diversity using frequency ratios. *Biometrics*, *71*(4),
1042-1049. https://doi.org/10.1111/biom.12332

Willis A. D. (2019). Rarefaction, Alpha Diversity, and Statistics. *Frontiers in Microbiology*, *10*,
1496. https://doi.org/10.3389/fmicb.2019.02407

Woolet, J. & Whitman, T. (2020). Pyrogenic organic matter effects on soil bacterial community
composition. *Soil Biology and Biochemistry*, *141*, 107678.
https://doi.org/10.1016/j.soilbio.2019.107678

Wotton, B. M. & Flannigan, M. D. (1993). Length of the fire season in a changing climate.
Forestry Chronicle, *69*(2), 187–192. https://doi.org/10.5558/tfc69187-2

Yang, Z. W., Asem, M. D., Li, X. *et al.* (2019). *Blastococcus deserti* sp. nov., isolated from a
desert sample. *Archives of Microbiology*, *201*, 193–198. https://doi.org/10.1007/s00203-
018-1604-1