Interactions during growth eclipse interactions during flowering for determining plant

fecundity in pollinator-sharing annual plants

Aubrie R. M. James and Monica A. Geber

Cornell University

Running head: Plant interactions during growth and flowering

Keywords: community ecology, species interactions, pollination ecology, pollinator-mediated

interactions, plant interactions, pollinator foraging

Abstract

Species interactions are foundational to ecological theory, but studies often reduce the complex

nature of species interactions. In plant ecology, the result is that interactions during vegetative

growth and flowering are considered separately, though both can affect fecundity. Here we use

a system of annual flowering plants in the genus *Clarkia* to ask how interactions during flowering

and growth contribute to plant interactions, and if pollinator behaviors explain apparent patterns

in plant interactions during flowering. We measure seed success and fecundity of *Clarkia* focal

plants in experimental interaction plots with the effect of pollinators experimentally removed or

retained. We also observe pollinator behaviors in the plots and experimental arrays. During

flowering, pollinators significantly changed the effect of *Clarkia* interactions on seed success in

31% of species interactions, and these changes corresponded to pollinator behaviors. Whole-

plant fecundity, however, did not depend on interactions between *Clarkia*; instead, non-*Clarkia*

forbs that grew earlier in the season limited fecundity, constituting a priority effect during

vegetative growth. Our study shows that interactions during vegetative growth can preclude the

effect of pollinator-mediated interactions on fecundity by limiting potential reproductive output,

and that simultaneously studying different modes of interaction allows for understanding the

contingency of ecological outcomes.

Introduction

Species interactions are foundational to the vast majority of ecological theory, including theories of the niche, population persistence, community network theory, density dependence, and coexistence theory. Despite outsize reliance on species interactions to explain natural patterns, however, the multifaceted nature of interactions can go unexamined in the literature (Bimler et al. 2018, Chesson 2018, Godoy et al. 2018, Bartomeus et al. 2021). In particular, studies suggesting a causal link between pollinator sharing and plant fecundity use proxies for pollination success but not whole-plant seed set, thus overlooking interactions over the course of growth in determining fecundity (Benadi and Pauw 2018, Opedal & Hegland 2019). On the other hand, studies suggesting a causal link between plant interactions over the course of growth typically overlook the importance of interactions during pollination and reproduction for fecundity (Chesson 2018, e.g., but see Lanuza et al. 2018). A natural step toward resolving this apparent discrepancy in approaches, and in bringing the multi-trophic complexity of plant interactions to bear in our theories of plant-pollinator systems, is to determine the components of natural history that might contribute to the net interaction between species, and the extent to which each component of the interaction actually determines plant performance (fecundity) (cf. Godoy et al. 2018).

Interactions between plants embedded in plant-pollinator networks can be broken down into two components: interactions over the course of vegetative growth that result from using resources like water, nitrogen, and sunlight, and interactions over the course of reproduction that result from pollinator sharing (Figure 1, Panel A). The net effect of growing together is thus a consequence of what transpires between plants over the course of both of these phases, despite the fact that they are rarely considered simultaneously. In the absence of pollinator sharing, plant interactions during vegetative growth -- be they positive, negative, or inconsequential -- should explain variation in performance (e.g. survival, fecundity) and thus

population persistence (Figure 1, Panel B). However, almost 90% of flowering plants are animal

pollinated (Ollerton et al. 2011), and pollinator sharing is a widespread phenomenon in plant

communities globally (Morales & Traveset 2008). As such, it is imperative to design studies that

accommodate interactions during growth and pollination to understand and explain plant

fecundity and understand population persistence (Bartomeus et al. 2021).

In flowering plant communities, pollinators have the potential to change the strength or direction

of plant interactions by making them either more or less competitive, or more or less facilitative.

The particular effect of pollinators on plant interactions depends on how pollinators behave

while foraging, as well as the composition of the pollinator community. First, pollinator sharing

can introduce or increase competition between plants. Two foraging behaviors in particular,

preference and constancy, determine successful conspecific (compatible) pollen transfer.

Preference is important because it represents a pollinator's choice to visit one plant species

over others when all are available (Figure 1, Panel C). Preference can introduce competition by

affecting the quantity of conspecific pollen that plants receive, where seed set declines with

declining pollinator preference (Pauw 2012, Song & Feldman 2014, Benadi 2015). Constancy is

the tendency of pollinators to visit one species of plant in a foraging bout (Figure 1, Panel D). If

pollinators are inconstant, then they can introduce incompatible, heterospecific pollen to co-

flowering plants, a form of reproductive interference between plants that decreases seed set.

Literature reviews have shown that heterospecific pollen transfer, a result of inconstant foraging

(for example), is not only widespread globally, but can have a large effect on individual plant

fecundity (Morales & Traveset 2008, Arceo-Gómez et al. 2016, Arceo-Gómez et al. 2019,

Moreira-Hernández & Muchhala 2019). Second, pollinator sharing may also have a facilitative

effect on a plant interaction by either alleviating a competitive interaction or amplifying a positive

interaction that has occurred during growth. Co-flowering plants have the potential to attract

more pollinators than when growing alone; for example, high plant density may result in high

competition over growth, but during flowering high floral densities in the same areas can

increase pollinator visitation rates and thus pollination service, a facilitative phenomenon known

as joint attraction (Figure 1, Panel E; Kunin and Iwasa 1996, Moeller 2004, Feldman et al. 2004,

Bizecki Robson 2013). The potential for such facilitation, however, should be contingent on low

interspecific pollen transfer, and thus high pollinator constancy. Third and finally, understanding

the composition of the pollinator community is critical in establishing the role of pollinators in

plant interactions, because different pollinator taxa are likely to exhibit differences in pollination

efficiency based on their behaviors (Waser et al. 1996; e.g. Eckhart et al. 2006, Barrios et al.

2016). Therefore, the effect of one plant species on another's successful pollination can greatly

depend on the specific pollinator species makeup of the community.

Though recent observational work corroborates that pollinators can affect flowering plant seed

set and flowering plant interactions (Lanuza et al. 2018, Bergamo et al. 2020), no experimental

study has quantified the contribution of pollinator sharing to the net effect of plant interactions

over growth and reproduction, nor demonstrated the link between interactions during

reproduction and pollinator foraging behavior. In this work, we do both in a system of four

sympatric and co-flowering winter annuals in the genus *Clarkia* (Onagraceae). These species

share bee pollinators specialized on the genus rather than any one species, and commonly co-

occur throughout their range of overlap, suggesting that they interact considerably over both

growth and reproduction (MacSwain, Raven, & Thorp 1973; Moeller 2005; Eisen and Geber

2018). Furthermore, these *Clarkia* flower late in the season after almost all other annual plants

in the area have senesced, such that the main interaction they experience during flowering is

brought on by sharing pollinators with each other (MacSwain, Raven, & Thorp 1973).

In this work, we investigate the separate and combined effects of plant-plant interactions over

growth and reproduction (via shared pollinators) among four species of *Clarkia*. To do so, we

use a combination of pairwise interaction plots with experimentally-sown *Clarkia* and

experimental pollen supplementation. In the pairwise interaction plots, *Clarkia* interact both

during growth and reproduction, with interactions during growth affecting plant size, and hence

reproductive potential, and interactions via shared pollinators during flowering affecting

successful pollination and seed set. The experimental pollen supplementation removes the

effect of pollinator interactions on final seed set, allowing the effect of interactions during growth

to be separated from the effect of interactions vias pollinators. We next investigate the link

between pollinator behavior and *Clarkia* interactions during flowering using pollinator

observations in our experimental interaction plots, and experimental floral arrays, to measure

pollinator preference, constancy, and visitation rate in response to floral density.

We predict that increasing background *Clarkia* density in plots will change the fecundity of focal

Clarkia plants sown into plots, independent of the effects of pollinators – i.e. *Clarkia* interactions

over the course of growth will affect total potential fecundity, absent the contribution of

pollinators. If sharing pollinators also contributes to the strength and direction of plant

interactions, then pollinators will render *Clarkia* interactions either more competitive or

facilitative -- i.e. pollinator-mediated *Clarkia* interactions over the course of flowering will also

affect the probability of seed set and total realized fecundity. Finally, the way pollinators change

Clarkia interactions should correspond to their foraging behavior. If interactions over

reproduction are competitive, then we predict pollinators will either be inconstant, exhibit strong

preferences for certain plant species, or both, because such behaviors result in heterospecific

pollen transfer or lower pollen receipt by less-preferred plants. If interactions over reproduction

are facilitative, then we predict pollinators will exhibit joint attraction and constant foraging.

Methods

System

Clarkia is a genus of outcrossing and generally self-compatible winter annual plants which

germinate in late winter and grow over the course of the spring. The four species of *Clarkia* in

this study (*C. cylindrica* ssp. *clavicarpa* (Jeps.) Lewis & Lewis*, C. speciosa* ssp. *polyantha* Lewis

& Lewis*, C. unguiculata* Lindl.*, C. xantiana* ssp. *xantiana* A. Gray, family Onagraceae) are

sympatric in the southernmost portion of the Sierra Nevada mountain range in the Kern River

Canyon (KRC; Kern County, California). *Clarkia* species co-occur in communities of varying

congeneric composition (from one to four *Clarkia* species), in low-elevation grassland,

chaparral, and oak woodlands. In the KRC, *Clarkia* begin to flower in early May, and flowering

lasts until mid- to late June. All four species overlap in flowering time for about two weeks in late

May, and share a group of bee pollinators specialized on the genus rather than any one species

(MacSwain, Raven, & Thorp, 1973, Singh 2013). Research has shown that at least one species,

C. xantiana, has higher seed set when it grows in multi-species *Clarkia* communities than at

sites where it occurs alone, and that this effect is explained by increased visitation from shared

pollinators at multi-species sites (Moeller 2004).

In this work, we use two different experiments to understand how pollinator-mediated

interactions contribute to the net interactions between *Clarkia*. First, we use an *in-situ* common

garden experiment to estimate the strength of *Clarkia* interactions over growth and flowering, as

well as estimate pollinator preference for *Clarkia* species. This is described in the 'Pairwise

Interaction Plots' sections below. Because pollinator constancy was impossible to estimate in

pairwise interaction plots due to their low species richness, we use experimental arrays of

flowering plants to estimate pollinator constancy ('Pollinator Arrays' sections below).

Experimental Design

Pairwise Interaction Plots

Interactions over growth and flowering

To determine the net effects of *Clarkia* growing and flowering together, we used an

experimental design closely modeled after work done to estimate competitive interactions in

annual plant communities (Godoy & Levine 2014; Godoy et al. 2014; Kraft et al. 2015;

Wainwright et al. 2019). In short, seeds of one *Clarkia* species were sown into a plot as a

competitive background, and then seeds of the same and different *Clarkia* species were sown at

low densities as monitored focal plants. This allowed us to estimate the effect of increasing

sown seed density of background *Clarkia* on focal plant fecundity.

Realized total plant fecundity is a function of both how many ovules a plant produces, which is

determined during growth and represents a plant's maximum potential fecundity, and the

success of converting those ovules to seeds, which is determined during flowering/pollination. In

our experiment, both of these phases are potentially determined by the density and identity of

background sown *Clarkia*. To isolate the contribution of interactions over flowering to the net

(combined growth and pollination) interactions of *Clarkia*, we added conspecific pollen to one

flower on each focal plant. This manipulation allowed us to compare seed production in fruits

with additional pollen and without additional pollen. The fruits that receive additional pollen as

flowers do not rely on pollinators for successful maximum seed set, so the difference in seed

production between these two pollination treatments is a direct quantification of the contribution

of pollinator-mediated interactions to realized focal plant fecundity. In addition, seed production

in the pollen-supplemented fruits can be interpreted as seed production in the absence of the

interactions mediated by pollinators, and therefore an estimate of how maximum potential

fecundity is a function of interactions over *Clarkia* growth.

In late June through early July 2016, we set up 160, 1X1 meter experimental plots in a

community where all four species naturally occur in the Kern River Canyon (35°31'45N,

118°39'36W; Area = 0.57ha). We arranged plots in blocks of four, where plots were separated

by ~1/4 meters, and blocks were separated by ~1/2 meter. We cleared all plots of all *Clarkia* to

prevent confounding natural seed rain into the plots. We left the non-*Clarkia* plants in place.

Clarkia seed was collected at various sites in the Kern River Canyon at the end of the 2016

growing season, after plants had senesced and fruits were just about to dehisce; this seed was

used in the sowing of the experiment. We extracted seeds using a hand deawner (Hoffman

Manufacturing, Inc, Fogelsville PA, USA) to break down the fruits, and a seed blower to clean

the seed.

We sowed the collected seed into experimental plots in October 2016. Each block was

haphazardly assigned a background *Clarkia* species, and each plot within the block was

haphazardly assigned one of five seeding densities (0, 0.04, 0.16, 0.64, and 2.56 g/m^2).

Background seeding densities were chosen based on previous estimates of naturally occurring

seed production at the field site (James, unpublished data). In total, our design yielded 20

background species X density treatments, with eight plot replicates per treatment.

Twenty focal seeds were sown into one of four designated locations (one for each species) into

each of the 160 plots. These locations were delimited using 2.5cm radius cabone craft rings

(CSS Industries, Inc., PA, USA) secured in the ground with two 3.8cm fence staples.

In the spring of 2017, we pruned the focal seedlings in the cabone rings to one focal *Clarkia*

plant. To prevent bias in selecting a focal seedling, we designated whatever seedling was

closest to the center of the ring as the focal plant and weeded the rest. In this same period, we

counted the number of background *Clarkia* that had germinated and the number of non-target

grasses and forbs growing within a 5cm radius from the focal seedling. We also weeded any

germinated, non-target *Clarkia* in the background of all plots.

During the two-week flowering phase in late May 2017, we haphazardly chose one flower on

each focal plant to be the 'no pollen addition' or control fruit and receive no additional pollen,

and another to be the 'pollen addition' fruit, receiving additional conspecific pollen. This was

done for all focal plants with $\geq$ 2 flowers. To control for any potential differences in the

environment to which paired flowers were exposed, we chose pairs of flowers that were open

and in a similar reproductive stage on the focal plant. To add conspecific pollen to the pollen

addition flowers, we harvested anthers from *Clarkia* before 8am from naturally-occurring *Clarkia*

plants growing nearby, before pollinators begin foraging for the day. This pollen was applied to

freshly opened and receptive stigmas on the selected flower on each focal plant immediately

after harvesting the pollen.

Focal fruits were collected at the end of the growing season, after the focal plant had senesced

but before its fruit capsules had dehisced. At the time of fruit collection, we counted the number

of mature fruits on all focal plants and collected the total above ground biomass of *Clarkia*

growing in all plot backgrounds. The number of seeds, aborted seeds, and ovules of each

treatment fruit were counted and recorded.

Pollinator visitation in the interaction plots

During the co-flowering period in 2017, we observed bee visitation behavior in the pairwise

interaction plots to quantify pollinator preferences and joint attraction. Due to time limitation in

the season and the fact that some plots did not have surviving focal plants, not all 1m^2 plots

were observed; however, at least one plot from each block was observed over the course of co-

flowering. Observations occurred in ten-minute periods. Once a bee entered a plot and visited a

flower, one person tracked and called the bee's visits to flowers in the plot while another person

recorded the information. The tracker identified the bee to either species or genus (when

species identification was impossible). After each observation period, we counted and recorded

the number of open flowers of each species in the observed plot.

Pollinator Arrays to assess constancy

We assembled cut-stem arrays of the four *Clarkia* species to observe pollinator behavior during

the 2015 and 2016 *Clarkia* growing seasons. To do so, we cleared a 5 x 5m patch of all *Clarkia*

in the middle of the experimental site. The array site was prepared in late April before *Clarkia*

plants had started flowering. We cleared the array site of all naturally-occurring *Clarkia* plants

and constructed a grid with nodes at every 0.5m increment for a total of 100 nodes to place cut

Clarkia stems. We buried a floral water pick at each node to fill with water and hold *Clarkia*

stems.

The data we present in this work are a subset from a larger experiment to observe pollinator

foraging behaviors. Here, each array treatment consists of one of four densities (in 2015: 10, 20,

40, or 100 plants; in 2016: 12, 24, 64, or 100 *Clarkia* stems) comprised of an equal number of

cut stems for all four species of *Clarkia.* This design allowed us to observe pollinator constancy

across a range of plant densities, holding diversity relatively constant. Because different bee

species forage at different times of day (Hart & Eckhart 2010), we ran most treatments in the

morning and afternoon on different days: one array run in the morning, and one array run in the

afternoon. For each array run, we gathered *Clarkia* stems before 8am to assemble in the grid.

We arranged *Clarkia* stems in the grid according to a randomly-generated map of a particular

treatment, where locations in the grid were either left 'blank' or received one *Clarkia* stem of a

prescribed *Clarkia* species depending on the treatment density.

During the array runs, we identified and monitored bees as they entered and foraged in arrays.

When a bee entered the array and landed on a flower, one person called out the grid location of

the flower/bee, identified the bee to species or genus, and tracked the bee's movements until it

left the array, while another person recorded. At the end of each array run, we gathered and

disposed of wilted *Clarkia* stems. Rarely, we re-used stems if they were visibly in 'good-shape'

(i.e., turgid petals and leaves), though never for more than two runs. Array treatments lasted 40

or 60 minutes; many treatments in 2016 were stopped at 40 minutes due to wilting. We did not

run the experiment on days when the predicted peak temperature was below 21°C or wind was

stronger than a light breeze.

Statistical Analysis

All calculations and statistical analyses were performed in R (R version 3.5.2, R Core Team,

2018).

Interactions during growth and reproduction

We used two different analyses to understand the effects of pollinators on *Clarkia* interactions.

First, we used generalized linear mixed effects models with a binomial error distribution to

determine what variables or combination of variables best explained the probability of

successful seed set in focal fruits. To do so, we first counted the number of fertilized seeds,

aborted seeds, and unfertilized ovules in the control and pollen-supplemented fruits harvested

from the focal plants. The sum of these three values represents the total potential fecundity of a

fruit. We then built a set of candidate models to explain variation in seed set success using four

fixed effects – the pollination treatment, the focal *Clarkia* species identity, the background

Clarkia species identity, and the biomass of background *Clarkia* in the plot. We did not include

non-*Clarkia* grasses or forbs in this analysis because only *Clarkia* were flowering/releasing

pollen during the experiment, and therefore would not affect the pollination interactions of

Clarkia. We also used one random effect, the identity of the plant, in all models. We defined

'success' as fertilized, unaborted seeds in the fruit, and 'failure' as everything else (ovules and

aborted seeds). We fit the data to all candidate models using the lme4 package (Bates et al.

2015). We then calculated the AICc of each candidate model, which we used to score model fit;

the model with the lowest AICc score is the best-fit model that we used for ecological inference.

The most complex model we used included an interaction of all four fixed effects, and

represents our prediction that successful seed set depends on the density-dependent effect of

the background *Clarkia* on the focal *Clarkia*, and varies according to pollination treatment.

After identifying the best-fit model, we used estimated marginal means to determine the

predicted density-dependent effect of background *Clarkia* on focal *Clarkia*, with and without

pollen addition (package emmeans, Lenth 2019). Here, the effect of pollinators on the

interaction is the change in the density-dependent effect of background *Clarkia* on focal *Clarkia*

seed set between the two pollination treatments (pollen supplementation and no pollen

supplementation). If pollinators change the strength or direction of the interaction, then the effect

of increasing background *Clarkia* biomass will be significantly different between the two

pollination treatments. We also tested if the estimated density-dependent effect of background

Clarkia on focal *Clarkia* was significantly different from zero for both pollination treatments. This

was to examine if and when pairwise *Clarkia* interactions were significantly different from neutral

(no effect of increasing density). Finally, we estimated pollen limitation to successful seed set of

each *Clarkia* species. To do so, we compared the probability of successful seed set according

to pollination treatment for each species, averaged across all background treatments. If species

exhibit pollen limitation to successful seed set, then there will be a significant difference

between pollination treatments.

The second analysis was to determine what variables best explained the total reproductive

output of focal plants in the interaction plots. Whereas the analysis described above is

concerned with the probability of seed set, this analysis is concerned with the total realized

fecundity of focal plants- that is, accounting for variation in fruit production, which itself may be a

function of the effects of background *Clarkia* density on focal *Clarkia* growth. We again used

generalized linear mixed effects models with a Poisson error distribution to model total

estimated reproductive output under two different scenarios: pollen addition (where pollinators

do not introduce changes in *Clarkia* interactions) and no pollen addition (where pollinators can

change *Clarkia* interactions). We included an observation-level random effect in these models to

account for overdispersion in the data. Total realized fecundity of the focal plant was estimated

for the two treatments (pollen addition and no pollen addition) by taking the product of the

number of fruits on the focal plant, the average potential fecundity of the two treatment fruits we

collected (i.e. the ovule+fertilized seed+aborted seed number), and the proportion of successful

seeds in the supplemented fruit and the un-supplemented fruit.

We built candidate models using six potential fixed effects: pollination treatment, background

Clarkia identity, focal *Clarkia* identity, background *Clarkia* biomass, and the number of forbs and

grasses in the 5cm radius ring around focal *Clarkia* seedlings. We used two independent

random effects: plant identity and (as mentioned) observation. The most complex candidate

model included an interaction between the pollination treatment, background *Clarkia* identity,

focal *Clarkia* identity, and background *Clarkia* biomass, as well as the additive effects of forb

and grasses. This candidate model represented our prediction that full plant seed production

depends on the density-dependent effect of the background *Clarkia* on the focal *Clarkia*, which

is modified by the activity of pollinators. Forb and grass counts were centered and standardized

to improve model performance. As before, we used AICc to score model fit, and used the model

with the lowest AICc score for ecological inference. With the best-fit model, we used the Wald

test to determine the significance of the predictors on full-plant fecundity.

Pollinator preference in the interaction plots

To quantify pollinator preference, we used plant-pollinator observations to build an interaction

network using the R package bipartite (Dormann et al. 2008). In the network, we only included

the first visit of each pollinator to a plot. This was because subsequent movements were

confounded by the experimental design: each plot was dominated by one background species,

and as such, subsequent movements within a plot were likely to be visits to the background

species, irrespective of pollinator preference (in the observation periods, visits to focal plants

against a background of a common species were rare). We first estimated d', or network

specialization, of the three most common pollinator taxa we observed visiting the plots

(Blüthgen et al. 2006). Higher values of d' indicate higher values of pollinator specialization

within the network, which we interpret as taxon-level pollinator preferences for different *Clarkia*.

We only included the three most common pollinator taxa because all other pollinators accounted

less than 2% of the sample (Table S1). Next, we used a chi-squared test to determine if net

visitation to *Clarkia* species deviated significantly from the null expectation of evenly-distributed

visits among plants. If a certain *Clarkia* species is preferred by the pollinator community overall,

then we would observe higher frequency of visitation to that species than under the null

expectation of random visitation.

Pollinator constancy in the experimental arrays

To understand the potential for pollinator (in)constancy to affect pairwise interactions between

species, we built a 4x4 matrix for each array run, where rows represent the *Clarkia* species first

visited by a pollinator, and columns represent the species visited next in a sequence of two

sequential moves. On-diagonal entries in the matrix thus represent pollinator movements to the

same species of *Clarkia*, and off-diagonal movements between different species. We divided

the sum of all movements in arrays by the sum of all inconstant movements to quantify the

proportion of movements that were inconstant, i.e. the proportion of movements that might lead

to incompatible pollen transfer in *Clarkia*. We also tabulated the number of inconstant

movements between each pair of *Clarkia* species and conducted a chi-squared test to

determine which pairs of species are at higher risk for incompatible pollen transfer. For this test,

we separated the pairs of species by the direction of the transition: for example, inconstant

movements from *C. cylindrica* to *C. speciosa* were considered separately from inconstant

movements from *C. speciosa* to *C. cylindrica.*

Joint attraction in the pairwise interaction plots

We used a linear model of visitation rate (visits per minute) and plotwise floral abundance to

determine the potential for joint attraction in the pairwise interaction plots. Floral abundance

from each array was log-transformed in the linear regression. If plant patches exhibit joint

attraction, then visitation rate should increase as a function of plotwise floral abundance.

Results

The effect of interactions over growth and flowering on seed production

We collected 371 fruits from 232 focal *Clarkia* plants in the pairwise interaction plots. The

uneven number of fruits collected was due to fruit loss caused by abortion and herbivory. We

found that the best-fit model of the probability of successful seed set included an interaction of

focal *Clarkia* species identity, background *Clarkia* species identity, background *Clarkia* biomass,

and pollination treatment (Table S2). In other words, the best fit model of the probability of seed

set depended on the species-specific, density-dependent effect of background *Clarkia* on focal

Clarkia, and this effect was modified by pollinators.

Pollinators significantly changed the density-dependent effect of background *Clarkia* on focal

Clarkia seed success in 31% of the interactions (five of 16 interactions; Figure 2, Table 1). In

three of these interactions, the density-dependent effect was more competitive when the effect

of pollinators was included (no pollen addition treatment) than when the effect of pollinators was

removed (pollen addition treatment). In other words, pollinators introduced a competitive effect

of the background *Clarkia* on focal *Clarkia* for the following three background/focal species

combinations: *C. unguiculata/C. xantiana; C. xantiana/C. unguiculata;* and *C. cylindrica/C.*

xantiana. Conversely, in the other two cases, the density-dependent effect of the background

was more positive with the effect of pollinators than without; that is, pollinators introduced a

facilitative effect of the background on focal *Clarkia* in *C. speciosa/C. cylindrica* and *C.*

speciosa/C. speciosa background/focal combinations.

Only two of the 32 density-dependent effects of background *Clarkia* density on the probability of

focal *Clarkia* seed set were significantly different from zero. One was the competitive effect of *C.*

unguiculata on *C. xantiana* in the pollen addition treatment ($z = -2.49$, $p=0.01$), and the other was

the competitive effect of *C. cylindrica* on *C. xantiana* in the no pollen addition treatment ($z= -$

4.89, $p<0.001$) (Figure 2).

Finally, the average pollen limitation to successful seed set varied by focal species. The effect of

the pollen addition treatment compared to the no additional pollen treatment was negligible in *C.*

speciosa, indicating that pollination service did not limit seed production for this species.

Additional pollen increased the probability of successful seed set in *C. xantiana, C. cylindrica,*

and *C. unguiculata*, in that order. This indicates that pollination service limited seed production

in these three species, but to varying degrees (Figure 3).

The best-fit model for full-plant fecundity included the additive fixed effects of (1) pollination

treatment and (2) the number of forbs surrounding the focal plant when it was a seedling (Table

S1). In other words, the full-plant fecundity of focal *Clarkia* plants was best predicted by the

effect that non-*Clarkia* forbs had on focal plants over the course of growth, and wholesale pollen

limitation in *Clarkia* during flowering (Figure 4). Full-plant fecundity strongly declined with

increasing forbs (-0.77 ± 0.15 SE log unit decrease; Wald p-value < 0.001), and was

significantly higher in the pollen addition treatment relative to the no pollen addition treatment

(1.17 ± 0.20 SE log unit increase; Wald p-value < 0.001).

Pollinator foraging behaviors

Preference and joint attraction in the interaction plots

We performed 209, 10-minute observation periods and observed 446 pollinators in the *Clarkia*

interaction plots in 2017. The three most abundant pollinator visitors, in order of abundance,

were *Diadasia angusticeps* (Apidae), *Hesperapis regularis* (Mellitidae), and *Lasioglossum sp.*

(Halictidae) bees, which made up more than 99% of the total observed pollinators (Table S1).

The preference of these three pollinators varied, where *D. anguisticeps* exhibited the most

specialized behavior, visiting *Clarkia speciosa* to the near exclusion of the other *Clarkia* (d' =

0.79), followed by *H. regularis* (d'=0.33), and *Lasioglossum sp.* (d'=17), both of which visited all

four *Clarkia*, but most commonly *C. xantiana* and *C. cylindrica* (Figure 5, Panel A).

Overall visitation to the different *Clarkia* was significantly different from the null expectation that

visitation to each species is proportional to their frequency in the network (χ^2(3)=64.4; p<0.001).

The residuals of the chi-squared analysis indicate strong overall pollinator preference for *C.*

speciosa and *C. xantiana* (Pearson residuals = 3.7 and 3.4, respectively), weak avoidance of *C.*

cylindrica (Pearson residual = -1.08), and strong avoidance of *C. unguiculata* (Pearson residual

= -6.1).

Pollinator constancy in the experimental arrays

We performed 17 array runs with all four *Clarkia* species, observing 664 pollinator movements

from 418 bees across 2015 and 2016. Of the 664 movements, only 100 (15%) were between

different plant species (and thus inconstant) movements. This indicates that *Clarkia* pollinators

are by and large constant foragers, even when equally exposed to all four *Clarkia* species.

Of the inconstant movements between species, some transitions between species pairs were

more common than others, indicating that certain species pairs were at higher risk for

interspecific pollen transfer than others. Interspecific movements were significantly different

from the null expectation of being evenly divided among species pairs (χ^2(11)=69.4; p<0.001).

The species pairs where switching occurred more frequently than the null expectation were (1)

C. xantiana and *C. unguiculata,* and (2) *C. xantiana* and *C. cylindrica*; and, for each pair,

switching was common in both directions (.e.g, X to U and U to X). Altogether, switches

involving these two pairs of *Clarkia* accounted for more than 60% of inconstant movements in

the dataset (Figure 5, Panel C).

Pollinator Joint Attraction

Finally, pollinator visitation rate (in visits per minute) increased with log floral abundance in the

interaction plots, indicating joint attraction (ß=0.27, Wald's p-value<0.001; Figure 5, Panel C).

Discussion

Theories in ecology necessarily reduce complexity to generate testable hypotheses of how the

natural world works. In community ecology, species interactions are the building blocks of

theories of community functioning and species persistence. Often, studies focus on single

aspects of species' life histories, such as interactions over growth or pollination, but not both.

This can lead to overlooking the multiplex nature of species interactions and overly simplistic

predictions about plant community functioning (Bartomeus et al. 2021). In this study we do our

best to reconcile that plant interactions during both vegetative growth and flowering can affect

interactions between co-flowering plant species in a system of four pollinator-sharing *Clarkia*

species. We experimentally addressed this in two parts: first, we grew *Clarkia* species together

and used hand pollination to determine the contribution of interactions over pollination to the

density-dependent effect of growing together for: (1) the probability of successful seed set and

(2) the total seed set of focal plants. Second, we investigated pollinator behaviors to determine

how the effect of species interactions we observed over pollination may be linked to patterns of

pollinator choice and movement.

The effect of pollinator-mediated interactions on potential fecundity

We find that the contribution of pollinator-mediated interactions during flowering to the realized

fecundity of plants is, above all, contingent on what transpires over growth. Whereas the

probability of successful seed set depended on pollinator-mediated *Clarkia* interactions

(discussed below), the realized fecundity of focal plants was best explained by only two

variables: the number of non-*Clarkia* forbs surrounding seedlings during growth, and pollination

treatment. We interpret this result to mean that the competitive effect of non-*Clarkia* forbs was

much more important in determining the realized fecundity of focal plants than any effect of

Clarkia density during growth or flowering. In particular, this result suggests that the competitive

effect exerted by non-*Clarkia* forbs on *Clarkia* seedlings during growth limited total potential

fecundity, and that this interaction overwhelmed the effects of interactions during flowering

(which can only limit the proportion of successful seeds out of total potential fecundity). Because

Clarkia are among the last to flower in the growing season in the region, this may be evidence

for a priority effect – e.g. non-*Clarkia* forbs draw down resources earlier in the season, thus

limiting the resources available later in the season when *Clarkia* plants grow and flower. This

result is consistent with other studies demonstrating that differences in phenology can tip the

balance of species interactions in annual plant systems (Godoy and Levine 2014): for example,

early-season plants can prevent the establishment or limit reproduction of late-season plants

due to resource draw down (specifically Nitrogen; Hooper & Dukes 2010, Mooney et al. 1986).

Despite not finding support for a significant effect of pollinator-mediated interactions on realized

plant fecundity, we still found evidence of strong pollen limitation to reproduction: realized full-

plant fecundity was higher in pollen-supplemented scenarios compared to no supplemental

pollen scenarios (Figure 4). However, this effect did not vary among species, suggesting that

differences in pollen limitation to successful seed set (at the fruit scale) were not great enough

among *Clarkia* species to materialize in species-specific differences in how pollen limitation

affected whole plant realized fecundity. Previous experimental work in other flowering plant

systems suggests that as pollinators become scarcer, pollinator-mediated competition increases

(Lázaro, Lundgren, and Totgren 2014; Ye et al., 2014). Taking the results of these studies and

the present study together, we suspect that pollinator abundance was high enough that

pairwise, pollinator-mediated interactions between plants did not bear on realized fecundity in

our experiment. Further, it is likely that the extent to which pollinator-mediated effects on

realized plant fecundity are a function of pollinator abundance, and as such may vary from year

to year.

Pollinator-mediated interactions correspond to foraging behavior and affect the

probability of successful seed set

The results at the scale of entire plants differ from results at the scale of fruits, where the

probability of seed set was governed by the density and identity of *Clarkia* surrounding a focal

plant. Significant shifts in species interactions between pollination treatments were apparent in

five of the 16 pairwise species interactions; put another way, pollinators changed the positive,

negative, or neutral effect of background *Clarkia* density on the probability of seed set. A few

patterns stand out from this analysis. First, the *Clarkia* pairs that exhibited density-dependent

interactions significantly different than zero (*C. cylindrica* background*/C. xantiana* focal*,* and *C.*

unguiculata background*/C. xantiana* focal; Figure 2) shared pollinators (Figure 5, Panel A), and

represented the two species pairs that pollinators switched between most often (Figure 5, Panel

B). Therefore, incompatible pollen transfer from pollinator sharing likely affects the probability of

successful seed set. Second, the only species that exhibited a significant intraspecific pollinator-

mediated interaction was *C. speciosa,* which does not share pollinators to a great extent with

the other three species (Figure 5, Panel A), and experienced lower rates of inconstant visitation

(Figure 5, Panel B). In this instance, increasing conspecific density trended toward competition

when the effect of pollinators was removed (i.e. there was a negative effect of background

density on seed set in the pollen addition treatment), but trended toward facilitation when the

effect of pollinators was retained (i.e. there was a positive effect of background density on seed

set in the no pollen addition treatment). This suggests that joint attraction of specialized

pollinators (carrying compatible pollen) has the potential to introduce positive density-

dependence in conspecific stands of flowering plants. Third and finally, the effect of *C. speciosa*

backgrounds on *C. cylindrica* seed set trended towards facilitative without the effect of

pollinators, but became less so with effect of pollinators; in this case, pollinators introduced a

competitive effect that dampened *C. speciosa* facilitation of *C. cylindrica*. Because these two

species do not share pollinators (Figure 4 Panel A), and there is rarely switching between the

two (Figure 4 Panel C), the competitive effect from pollinators is likely due to low pollinator

service to *C. cylindrica* when surrounded by *C. speciosa* (Table 1)*.*

Whereas five pairwise interactions were significantly different between pollination treatments,

only two of these effects were significantly different from zero. The greatest change between the

two pollination treatments was in plots where the background species was *C. unguiculata* and

the focal species was *C. xantiana*. In this interaction, the significantly competitive effect of *C.*

unguiculata density on *C. xantiana* ovule production during growth was neutralized when

including the effect of pollinators. To wit, pollinators acted as a facilitative force and 'rescued' *C.*

xantiana from the otherwise competitive effect of increasing *C. unguiculata* density*.* This result

may be explained by two pollinator behaviors we observed: first, joint attraction, where

increasing *Clarkia* density in the plot attracted more pollinators overall, and second, that

pollinators preferred *C. xantiana* much more than they did *C. unguiculata*, resulting in *C.*

xantiana receiving a disproportionate amount of pollinator visits. The other species interaction

significantly different from zero was that of *C. cylindrica* on *C. xantiana*, where the effect of *C.*

cylindrica density on focal *C. xantiana* seed set shifted from neutral without the effects of

pollinators, to significantly competitive when the effects of pollinators were retained – that is,

pollinators introduced a competitive effect of *C. cylindrica* on *C. xantiana*. We suspect this can

be explained simply by incompatible pollen transfer between plants from inconstant, highly-

shared pollinators. That being said, in these two species pairs, the fact that pollinators alleviated

competition in one case but exacerbated competition in another could alternatively be explained

by potential differences in how different species tolerate incompatible pollen. For example,

previous work has shown that flowering plants, including *Clarkia*, exhibit species-specific

sensitivity or tolerance to heterospecific pollens (Arceo-Gómez 2016, Arceo-Gómez et al. 2019);

it is possible that *C. xantiana* tolerates *C. unguiculata* pollen better than it does *C. cylindrica*

pollen*,* resulting in the differences in pollinator-mediated interactions*.* Future work could

investigate the extent to which pollinator foraging versus variation in heterospecific pollen

tolerance influence susceptibility to incompatible pollen transfer.

Finally, pollen limitation to successful seed set varied according to species. On average,

additional pollen did not affect the success of seeds in the fruits of *C. speciosa*, suggesting that

the pollination service to this species was high enough that additional pollen was (on average)

inconsequential for successful seed set. The behaviors of pollinators explain this result well:

high pollinator preference and low pollinator overlap (and therefore likely little heterospecific

pollen transfer) resulted in no effect of the experimental addition of pollen to *C. speciosa*

flowers' seed set. In contrast, additional pollen increased the average success of seeds in the

three other species: *C. xantiana* experienced a mild boost in seed success with additional

pollen, *C. cylindrica* a slightly higher boost, and *C. unguiculata* saw the greatest difference

between additional pollen and no additional pollen treatments. This is again explained by

pollinator behaviors, where the pollen limitation to successful seed set increased in order of

pollinator preference.

The importance of studying multiple life history stages to understand plant interactions

and fecundity

Studying different stages of the life history permits us a deeper understanding of how context-

dependency plays out in plant interactions and plant communities. Our approach of studying the

interactions of *Clarkia* species without removing the background of other grasses and forbs was

revealing of the ecology of the system: if we had taken a more traditional approach of studying

Clarkia pairwise interactions out of the context of the rest of the annual plant community (by

weeding, for example), we may have come to the conclusion that interactions over pollination

are strong drivers of *Clarkia* interaction dynamics because it would have been impossible to

detect the priority effect introduced by non-*Clarkia* forbs. To investigate how variable community

context could change the importance of interactions over growth versus flowering, this work

could be extended by using models to compare population persistence or coexistence

outcomes in neighborhoods that are dominated by non-*Clarkia* plants (where growth

interactions and priority effects may take precedence in determining fecundity) versus

neighborhoods comprising exclusively *Clarkia* plants (where pollinator-mediated interactions

and interference competition might be more important for fecundity).

Finally, though we have studied both growth and pollination aspects of life history in order to

understand how they work together to determine fecundity, the larger tasks of identifying

population and community-level ramifications of species interactions remains. Incorporating

germination rates and rates of seed survival in the seed bank, as well as a dynamic

understanding of both priority effects and pollinator population variation through time, could be

relevant for understanding the role of species interactions in population persistence. In the

future, studies should investigate the sensitivity of population growth rates to these variables, as

well as how coexistence dynamics could change according to variation in neighborhood

composition.

Acknowledgements

The authors would like to thank Cara Grula, Noel Graham, Liz Richards, Tim Burnette,

Anastasia Pulak, Louie Jacob, Aidan Healey, Vince Eckhart, David Moeller, and Christine

Solomon for their help in the field, Anurag Agrawal, Steve Ellner, and Tom Miller for comments

on earlier drafts of the manuscript, and Katherine Eisen, Gregor Seigmund, and the Geber lab

meeting group for both help in the field and comments on earlier drafts. This work was

conducted on occupied land historically belonging to the Tübatulabal tribe, who are currently

seeking federal recognition. This work was funded by NSF DEB 1754299 to MAG and the

Cornell Atkinson Center for Sustainable Biodiversity to ARMJ.

Literature Cited

Arceo-Gómez, G., Raguso, R.A., and Geber, M.A. 2016. Can Plants Evolve Tolerance

Mechanisms to Heterospecific Pollen Effects? An Experimental Test of the Adaptive Potential in

Clarkia Species. Oikos 125: 718–25.

Arceo-Gómez, G., Schroeder, A., Albor, C., Ashman, T.L., Knight, T.M., Bennett, J.M., Suarez,

B., and Parra-Tabla, V. 2019. Global Geographic Patterns of Heterospecific Pollen Receipt Help

Uncover Potential Ecological and Evolutionary Impacts across Plant Communities Worldwide.

Scientific Reports 9.

Barrios, B., Pena, S.R., Salas, A., and Koptur, S. 2016. Butterflies Visit More Frequently, but

Bees Are Better Pollinators: The Importance of Mouthpart Dimensions in Effective Pollen

Removal and Deposition. AoB PLANTS.

Bartomeus, I., Saavedra, S., Rohr, R.P., Godoy, O. 2021. Experimental Evidence of the

Importance of Multitrophic Sturcture for Species Persistence. Proceedings of the National

Academy of Sciences, 118: e2023872118.

Bates, D., Maechler, M., Bolker, B., Walker, S. 2015. Fitting Linear Mixed-Effects Models Using

lme4. Journal of Statistical Software, 671: 1-48.

Benadi G. 2015. Requirements For Plant Coexistence Through Pollination Niche Partitioning.

Proceedings of the Royal Society B 282:20150117.

Benadi, G., and Pauw, A. 2018. Frequency Dependence of Pollinator Visitation Rates Suggests

That Pollination Niches Can Allow Plant Species Coexistence. Journal of Ecology 106: 1892–

1901.

Bergamo, P.J., Streher, N.S., Traveset, A., Wolowski, M., and Sazima, M. 2020. Pollination

Outcomes Reveal Negative Density-Dependence Coupled with Interspecific Facilitation among

Plants. Ecology Letters.

Bimler, M. D., Stouffer, D., Lai, H. R., and Mayfield, M.M. 2018. Accurate Predictions of

Coexistence in Natural Systems Require the Inclusion of Facilitative Interactions and

Environmental Dependency. Journal of Ecology 106: 1839–52.

Bizecki Robson, D. 2013. An Assessment of the Potential for Pollination Facilitation of a Rare

Plant by Common Plants: Symphyotrichum sericeum (Asteraceae) as a Case Study. Botany 91:

34–42.

Blüthgen, N., Menzel, F., and Blüthgen, N. 2006. Measuring Specialization in Species

Interaction Networks. BMC Ecology 6.

Chesson, P. 2018. Updates on Mechanisms of Maintenance of Species Diversity. Journal of

Ecology 106: 1773–94.

Dormann, C.F., Fruend, J., Bluethgen, N. & Gruber B. 2009. Indices, graphs and null models:

analyzing bipartite ecological networks. The Open Ecology Journal, 2: 7-24.

Eckhart, V., Rushing, N.S., Hart G.M., and Hansen, J.D. 2006. Frequency-Dependent Pollinator

Foraging in Polymorphic Clarkia Xantiana Ssp. Xantiana Populations: Implications for Flower

Colour Evolution and Pollinator Interactions. Oikos 112: 412–21.

Eisen, K.E., and Geber, M.A. 2018. Ecological Sorting and Character Displacement Contribute

to the Structure of Communities of Clarkia Species. Journal of Evolutionary Biology 31: 1440–

58.

Feldman, T.S., Morris, W.F. and Wilson, W.G. 2004. When Can Two Plant Species Facilitate

Each Other 's Pollination? Oikos 105:197–207.

Godoy, O., and Levine, J.M. 2014. Phenology Effects on Invasion Success: Insights from

Coupling Field Experiments to Coexistence Theory. Ecology 95: 726–36.

Godoy, O., Bartomeus, I., Rohr, R.P., and Saavedra,S. 2018. Towards the Integration of Niche

and Network Theories. Trends in Ecology and Evolution 33:287–300.

Hart-Fredeluces, G. & Eckhart, V. 2010. Ecological Separation in Foraging Schedule And Food

Type Between Pollinators Of The California Wildflower, Clarkia xantiana ssp. xantiana. Journal

of Pollination Ecology 2.

Hooper, D.U., and Dukes, J.S. 2010. Functional Composition Controls Invasion Success in a

California Serpentine Grassland. Journal of Ecology 98: 764–77.

Kharouba, H. M., Ehrlén, J., Gelman, A., Bolmgren, K., Allen, J.M., Travers, S.E., and

Wolkovich, E.M. 2018. Global Shifts in the Phenological Synchrony of Species Interactions over

Recent Decades. Proceedings of the National Academy of Sciences of the United States of

America 115: 5211–16.

Kunin, W., and Iwasa, Y. 1996. Pollinator Foraging Strategies in Mixed Floral Arrays: Density

Effects and Floral Constancy. Theoretical Population Biology 49: 232–63.

Lanuza, J. B., Bartomeus, I. and Godoy, O. 2018. Opposing Effects of Floral Visitors and Soil

Conditions on the Determinants of Competitive Outcomes Maintain Species Diversity in

Heterogeneous Landscapes. Ecology Letters 21: 865–74.

Lázaro, A., Lundgren, R., and Totland, Ø. 2014. Experimental Reduction of Pollinator Visitation

Modifies Plant–Plant Interactions for Pollination. Oikos 123: 1037–48.

Lenth, R. 2020. emmeans: Estimated Marginal Means, aka Least-Squares Means. R package

version 1.4.4. https://CRAN.R-project.org/package=emmeans

MacSwain, J. W., Raven, P. H. and Thorp, R. W. 1973. Comparative Behaviour Of Bees And

Onagraceae. IV. Clarkia bees of the western United States. University of Calif. Publ. Entomol.

1:80.

Moeller, D.A. 2004. Facilitative Interactions among Plants via Shared Pollinators. Ecology 85:

3289–3301.

Moeller, D.A. 2005. Pollinator Community Structure And Sources Of Spatial Variation In Plant-

Pollinator Interactions In Clarkia xantiana ssp. xantiana. Oecologia 142: 28-37.

Mooney, H. A., Hobbs, R.J., Gorham, J., and Williams, K. 1986. Biomass Accumulation and

Resource Utilization in Co-Occurring Grassland Annuals. Oecologia 70: 555–58.

Moreira-Hernández, J.I., and Muchhala, N. 2019. Importance of Pollinator Mediated Interspecific

Pollen Transfer for Angiosperm Evolution. Annual Review of Ecology, Evolution, and

Systematics.

Morales, C.L., and Traveset, A. 2008. Interspecific Pollen Transfer: Magnitude, Prevalence and

Consequences for Plant Fitness, Critical Reviews in Plant Sciences, 27:221-238.

Ollerton, J., Winfree, R. and Tarrant, S. 2011. How Many Flowering Plants Are Pollinated by

Animals? Oikos 120: 321–26.

Opedal, Ø.H., and Hegland, S.J. 2019. Using Hierarchical Joint Models to Study Reproductive

Interactions in Plant Communities. Journal of Ecology 00 (October):1–11.

Pauw, A. 2012. Can pollination niches facilitate plant coexistence?. Trends in Ecology &

Evolution 28.

Singh, I. 2013. Master's Thesis. Cornell University.

Waser, N.M., Chittka, L., Price, M.V., Williams, N.M., and Ollerton, J. 1996. Generalization in

Pollination Systems, and Why It Matters. Ecology 77:1043–60.

Ye, Z.M., Dai, W.K., Jin, X.F., Gituru, R.W., Wang, Q.F., and Yang, C.F. 2014. Competition and

Facilitation among Plants for Pollination: Can Pollinator Abundance Shift the Plant-Plant

Interactions? Plant Ecology 215:3–13.

Zhiyuan, S. and Feldman, M.W. 2014. Adaptive Foraging Behaviour Of Individual Pollinators

And The Coexistence Of Co-Flowering Plants. Proceedings of the Royal Society B 281:

20132437.

Figure 1. (A) Flowering plants can interact with each other during growth, which determines

potential reproductive output, and flowering, which determines realized reproductive output (via

pollination). (B) Models and tests often focus on how interactions during growth affect realized

reproductive output without considering the effects of interactions during pollination. (C)

Pollinator preference for one species over another affects the realized reproductive output of

both species. (D) Pollinator constancy (solid line is a constant move; dashed line is an

inconstant move) can affect the extent to which plants experience reproductive interference

throughout heterospecific pollen. (E) Joint attraction of pollinators can be a positive effect of

flowering plant co-occurrence.

Figure 2. The effect of increasing density of background *Clarkia* on the probability of an ovule

becoming a successful seed in focal *Clarkia*. The effect (slope) varies according to species pairs

and interactions over pollination. Each row is a different background/focal species combination.

Points indicate the estimated effect of increasing background species' density on the probability

of seed set for the focal species, and lines are standard errors. Negative values indicate the

probability of seed set declines with increasing background density (a competitive effect of

background *Clarkia*), whereas positive values mean that the probability of seed set increases

with background density (there is a facilitative effect of background *Clarkia*). Letters stand for

the first letter of the species epithet: C- *Clarkia cylindrica*; S- *C. speciosa*; U-*C. unguiculata*; and

X- *C. xantiana*. Blue points are the treatment where the effect of pollinators is removed (pollen

addition). Orange points are the treatment where the effect of pollinators is retained (no

additional pollen). Bolded combinations are those that include an interaction between

background density and pollination treatment that is significantly different from zero; stars

indicate a significant effect of pollination treatment on the interaction (*, $p<0.05$; **, $p<0.01$; ***,

$p<0.001$).

Figure 3. Species-specific probability of seed success in fruits for both pollination treatments.

Points represent the estimated average across all background species treatments at the mean

background *Clarkia* density. Bars represent the 95% confidence interval for each estimate.

Figure 4. Predicted full-plant seed set as a function of pollination treatment and the number of

non-*Clarkia* forbs surrounding the focal as a seedling.

Figure 5. (A) The network of the plant and pollinator interactions observed during the pairwise

interaction experiment. (B) The proportion of inconstant moves for each transition type between

different *Clarkia*, colored by species pair, in the experimental array experiment. (C) Rate of

visitation (visits per minute) as a function of floral abundance in the experimental array.

Table 1. The effect of pollinators on the log-odds probability of seed set in the interactions

between Clarkia species (Figure 1), and the explanation of the change in terms of observed

pollinator behaviors (Figure 3). Only statistically significant pairwise interactions are shown in

this table.

Background	Focal	Change in effect of background on focal ± SE (no additional pollen treatment effect – pollen addition treatment effect)	Interpretation	Behavioral Explanation	p-value
Clarkia unguiculata (U)	*Clarkia xantiana* (X)	4.88 ± 0.93	Pollinators alleviate the competitive effect of U on X	Joint attraction to higher densities, but pollinator preference for *C. xantiana*	< 0.001 (***)
Clarkia cylindrica (C)	*Clarkia xantiana* (X)	-0.72 ± 0.21	Pollinators exacerbate the competitive effect of C on X	Incompatible pollen transfer	0.0011 (**)
Clarkia xantiana (X)	*Clarkia unguiculata* (U)	-0.44 ± 0.21	Pollinators exacerbate the competitive	Pollinator preference for *C. xantiana* and	0.039 (*)

		effect of X on U		incompatible pollen transfer	
Clarkia speciosa (S)	*Clarkia speciosa* (S)	1.40 ± 1.48	Pollinators alleviate the competitive effect of S on S	Pollinator attraction to higher densities	0.0091 (**)
Clarkia speciosa (S)	*Clarkia cylindrica* (C)	-1.41 ± 0.53	Pollinators reduce the facilitative effect of S on C	Low pollinator overlap and therefore low pollinator service to C	0.0073 (**)