Tension mediated mechanical activation and pocket delipidation lead to an
analogous MscL state

Bolin Wang,[1,2†] Benjamin J. Lane,[1,2†] Charalampos Kapsalis,[3] James R. Ault,[1,4] Frank
Sobott,[1,4] Hassane El Mkami,[5] Antonio N. Calabrese,[1,4] Antreas C. Kalli,[1,6] Christos
Pliotas[1,2,3]*

[1]Astbury Centre for Structural Molecular Biology, University of Leeds, UK

[2]School of Biomedical Sciences, Faculty of Biological Sciences, University of Leeds, UK

[3]Biomedical Sciences Research Complex, School of Biology, University of St Andrews, UK

[4]School of Molecular and Cellular Biology, Faculty of Biological Sciences, University of Leeds, UK

[5]School of Physics and Astronomy, University of St Andrews, UK

[6]Leeds Institute of Cardiovascular and Metabolic Medicine, School of Medicine, University of
Leeds, UK

[†]These authors contributed equally to this work

*Corresponding author: Christos Pliotas, c.pliotas@leeds.ac.uk

Author Contributions: C.P. conceived and designed the project. B.W., B.J.L., C.K., J.A., H.E.M.,
A.C.K. and C.P. performed research. B.W., B.J.L, C.K., J.A., F.S., H.E.M., A.N.C., A.C.K. and
C.P. processed and analysed data. C.P. wrote the manuscript assisted by B.W., B.J.L. with
contributions from H.E.M, A.N.C. and A.C.K.

Competing Interest Statement: The authors declare no conflicts of interest

Keywords: Lipids, MscL, MscS, mechanosensitive channels, EPR, HDX, MD

This PDF file includes:

Main Text

Figures 1 to 7

Tables 1 to 1

Abstract

The MscL channel gates in response to membrane tension changes to allow the exchange of molecules through its pore. Lipid removal from transmembrane pockets leads to a MscL response. However, it is unknown whether there is correlation between the tension mediated state and the state derived by pocket delipidation in the absence of tension. Transitions between MscL states may follow a similar pathway to cover the available conformational space but may not necessarily sample the same discrete intermediates. Here, we combined pulsed-EPR and HDX-MS measurements on MscL, coupled with molecular dynamics under membrane tension, to investigate the changes associated with the distinctively derived states. Whether it is tension or pocket delipidation, we find that MscL samples a similar expanded state, which is the final step of the delipidation pathway but only an intermediate stop of the tension mediated path. Our findings hint at synergistic modes of regulation in mechanosensitive channels.

Main Text

Introduction

Specific lipid binding events and changes in transbilayer pressure can modulate the structure of membrane proteins and regulate their function (1-11). Mechanosensitive (MS) ion channels are membrane proteins which sense and respond to tension changes in the cell membrane. Pressure-sensitive domains, or pockets, formed within transmembrane (TM) regions of bacterial MS channels have been proposed to play a role in their mechanical sensing and response (1, 2, 8, 12, 13). These pockets were first identified on the MS channel of small conductance MscS, leading to the lipids-move-first hypothesis, which predicts that lipids accessing the pockets physically inhibit (directly or allosterically) the movement of pore helices to gate the channel (2). When a modification (L89W) was introduced at the entrance of these pockets in MscL, steric lipid interactions with the backside of the pore-lining helix TM1 were significantly reduced and as a consequence MscL expanded, adopting an intermediate state, even in the absence of applied tension(8). MscL's mechanical response suggested that pocket targeting lipids could act as negative allosteric modulators for MS channels and that tension could be mimicked by molecules or gain-of-function modifications targeting the pocket region, which could disrupt the lipid pathway between the bulk membrane and the pockets to gate the channel (1, 8, 14, 15). Indeed, the function of MscL and MscS (5, 14) can be modulated by changes in molecule occupancy within these pockets (2, 8, 16-18) and even subtle structural differences within the same regions could lead to functional differences in different MscL proteins(19). This suggests a common regulatory mechanism, tailored to the unique structural landscape of MS channels, which may guide ligand specificity.
In mammals, the MS channel Piezo1 is activated by Yoda1, opening the possibility of endogenous agonists(20, 21), while TRAAK and TREK-1 are responsive to natural chemical ligands including arachidonic acid, polyunsaturated fatty acids, and lysophosphatidic acid(22-24). Lipids residing within similar inner-leaflet pockets have also been linked with mechanosensation in x-ray and cryo-EM structures of TRPV3(25), TRAAK(26), TREK-2(27), YnaI(28), MSL1(29) and MscS(2, 17, 30, 31), while their unique parallel to membrane plane orientation may be guided by an amphipathic helix that is important in the mechanosensitivity of MscL(32) and GPCRs(33). Pulsed electron-electron double resonance (PELDOR, also known as DEER)(34, 35) and x-ray studies on the LmrP multidrug transporter showed that lipids could provide a hydrophobic counterpart to partially fill binding pockets and participate in the coordination of ligand binding(36).
Attempts to model MscL opening have been previously implemented via steered MD simulations, with added tensions to pull the channel open (37-41), but the tension used in this modelling is an order of magnitude higher than that required to open MscL in giant unilamellar vesicles (GUVs) and spheroplasts(8, 42, 43). In other cases, MscL was subjected to extensive modifications or forces

were applied for relatively short periods of time(37, 44). MD simulations on TREK-2 have shown
that under increased bilayer tension the channel moves from a "down" to an "up" configuration
accompanied by cross sectional area changes, while its pockets are occupied by lipids only in the
absence of membrane stretch(45).
Hydrogen Deuterium eXchange Mass Spectrometry (HDX-MS) and 3 pulse Electron Spin Envelope
Echo Modulation (3pESEEM) spectroscopy are powerful tools in the investigation of membrane
protein structural dynamics. The former involves the exchange of protons on a protein with
deuterium to generate accessibility information, and has been used to provide key insights into the
role(s) of lipids in modulating membrane protein conformation(46-49). The latter measures weak
hyperfine coupling between unpaired electrons and nuclear spins to probe accessibility to solvent
(D_2O) and has been used to investigate the dynamics of membrane proteins(8, 19, 34, 35, 50-54).
Here we endeavored to investigate whether there is a structural analogy between the
physiologically relevant tension-activated state, and the one stabilized by modifications that result
in pocket lipid removal. To this end, we have combined untargeted (HDX-MS) and single-residue
(3pESEEM) methods to probe the architecture of MscL upon pocket delipidation and independently
generate a tension-activated MscL state by MD simulations. We find these two differently derived
states lead to an analogous state, suggesting a direct link between tension- and pocket lipid
removal- activation in MS channels. Further, our data suggest that upon tension-activation
additional structural changes can trigger substantial further opening of the channel pore, suggesting
that structural plasticity of these mechanosensitive ion channels enables them to respond
differently upon receipt of discrete stimuli.
102
103   **Results**
104
Channel gating by lipid pocket removal activation monitored by HDX-MS
The substitution of a tryptophan at position L89 in *Mycobacterium tuberculosis* MscL (TbMscL) has
been shown to restrict lipid access to channel pockets and destabilise the closed state, leading to
an expanded (sub-conducting) MscL state(8). In order to give a mechanistic insight into the
transitions occurring between the two states, we used HDX-MS to measure relative differences in
deuterium uptake between the WT (closed) and L89W (expanded by pocket delipidation) channel
proteins. We succeeded in obtaining 95% peptide coverage of the entire resolved MscL structure
where the gating occurs, i.e., residues 1-125, including the entire TM domain and the largest part
of the cytoplasmic helical bundle (83% overall coverage for the entire construct including the C-
terminal 6xHis-tag). Differences in uptake, detected at the peptide level, between the two states
allowed us to identify protein regions that became deprotected from exchange in the sub-
conducting L89W state (Fig 1). Three regions of MscL were identified and showed significant
changes between the two states. Peptides in the regions containing residues 37-53 (periplasmic
loop), 58-69 (top of TM2), and 97-111 (bottom of TM2, cytoplasmic loop and top of the cytoplasmic
helical bundle) had a significantly higher uptake of deuterium in L89W in comparison to WT TbMscL
(Fig 1, S1 and Table S1). Residues 37-53 run from the top of TM1 and end after the first β-sheet
in the loop connecting TM1 to TM2. Residues 58-69 correspond to the middle of the periplasmic
loop up until TM2, and residues 97-111 cover the bottom of TM2 to the top of the C-terminal helical
bundle (Fig 1 and S1). This gave us the first glimpses into MscL's pocket lipid removal activation
and, which protein domains are mostly impacted by the L89W (pocket entrance) modification
across the entire MscL's length.
126
Solvent accessibility mapping at single residue resolution by 3pESEEM
Next, we sought to investigate the feasibility of 3pESEEM to study the structure of MscL and identify
sites that are potentially dynamic during gating and solvent accessible. To this end, we generated
twenty-six single cysteine mutants, accounting for 20% of the total TbMscL length, spanning all
protein domains. Specifically, L2 and F5 on the S1 helix, N13, L17, V21, I23, V31 and F34 on TM1,
L42, and F48 on the periplasmic loop, residues N70, V71, L72, L73, S74, F79, F84, A85, Y87, F88,
L89, R98, K99, and K100 on TM2, and E102 and V112 on the cytoplasmic loop and helical bundle
(CHB) respectively, were mutated to Cys in a Cys-free WT TbMscL background (Fig 2). We

subsequently expressed, purified and spin labelled each one of these twenty-six mutants (MTSSL
modification is denoted as R1 hereafter) and performed 3pESEEM solvent accessibility
measurements (Fig S2). Solvent accessibility is based on the modulation depth of deuterium in the
time domain signal, which is proportional to its associated signal intensity in the frequency domain.
We used two independent analysis methods to determine solvent accessibility, both yielding very
similar results (Fig S3). Unlike PELDOR which requires high spin labelling efficiency to obtain high
sensitivity data acquisition(8, 13), 3pESEEM can be performed with no major losses in sensitivity
even if the sample of interest shows lower spin labelling efficiencies for individual sites. This allowed
us to obtain good quality spectra and quantify the solvent (D_2O) accessibility for all twenty-six sites
(Fig 2, S2 and S4). Residues in the S1 helix showed low accessibility, which is consistent with
previous reports suggesting that this helix is buried within the bilayer(32). TM1 pore-lining residues
presented low to intermediate accessibility, as would be expected for a closed channel pore(55).
I23, V31 and F34 on TM1 are intermediately solvent-exposed (or relatively buried) and positioned
on the same side of TM1 (Fig 2 and S4). This could be due either to the presence of native lipids,
detergent used for membrane protein extraction and/or hinderance by the presence of TM2 on the
TM1 interface. L42 and V48 are intermediately to highly exposed, as they form part of the
periplasmic loop connecting TM1 and TM2. N70, V71, L72, L73, and S74 present the largest
disparity across our whole data set. L72 is the most buried residue we measured (~ 3% compared
to the most solvent-exposed cytoplasmic-facing labelled residue K100R1) while N70 (~90%) and
V71 (~80%) are two of the most exposed (~90%), comparable to D_2O (solvent) accessibilities
measured for exposed cytoplasmic sites (Fig 2, S2 and S4). The remaining residues present
intermediate accessibilities to these two extremes, as expected for subsequent residues, which
form a helical turn with different space orientation. Residues N70 and V71 are solvent-exposed in
the closed MscL state, while residues L72 and L73 are buried. The latter pair of residues is known
to become exposed during opening, due to an anti-clockwise TM2 rotation(8, 19, 56). F79, F84 and
A85 are relatively buried and lie at the interface of TM helices between different MscL subunits.
Y87 has been identified as a lipid binding site(57) and along with F88 and L89 presented low to
intermediate accessibilities and forms part of the pocket region(8, 19, 58). Except for R98, which
showed intermediate accessibility, K99 and K100, presented high accessibilities, indicating these
residues are solvent-exposed in the closed state. R98, K99 and K100 form a positively charged
cluster for attracting negatively charged lipids(57) and our data suggest that the non-ionic DDM
binds either differently or at a different site. Substitution of these residues with the neutral Gln had
no effect on MscL's conformation, suggesting that specific lipid headgroup binding on this particular
site does not influence MscL gating(8). E102 (cytoplasmic loop) and V112, which is located in the
upper portion of CHB and is expected to move apart upon MscL opening (59), presented high
solvent accessibilities. Combined, this has enabled us to validate 3pESEEM as a suitable tool to
map the structure of MscL.
172
Identification of channel gating regions with single residue resolution
HDX-MS resolution is usually limited to the peptide level and, therefore, it is often unable to resolve
changes that occur at different residues within the same peptide. Here we have already shown that
the solvent accessibility of individual residues of MscL can be probed using 3pESEEM. This
generates higher resolution information pertaining to the regions of structural change identified by
HDX-MS, as well as to the channel pore vapour lock. In each of these regions, we introduced eight
distinct single cysteine substitutions in an L89W MscL background for comparison with identical
sites in the background of the wildtype protein. Specifically, N13 and V21 (pore), L42 in the
periplasmic loop, N70, V71, L72, L73 (top of TM2) and K100 (bottom of TM2 and cytoplasmic loop)
(Fig 3, S5 and S6). V21 forms the vapour lock (along with L17) that controls the channel gate and
N13 sits one turn of a helix below. No significant change in this region was highlighted via HDX-
MS, while due to the proximity of spin labels pointing towards the pore, distances are out of
PELDOR's distance measurement range (< 19 Å) (Fig 1 and 3B). However, 3pESEEM is highly
sensitive to the local spin label environment, does not need a reference state as it is the case for
HDX-MS, and is not subjected to distance restriction between sites like for PELDOR. Differences
in the vapour-lock region may not have been detected by HDX-MS as structural changes may have

resulted in complex changes to hydrogen bonding networks, whereby some residues were
deprotected from exchange and others became protected, resulting in only minor/no changes in
deuterium uptake at the peptide level (Fig 1 and S1). To resolve this, we labelled the V21 and N13
sites crucial for the channel's vapour lock, and six other sites within the regions identified by HDX-
MS, which undergo conformational changes when the L89W modification is present (Fig 3, S5 and
S6). Overall, the motivation behind this was to confirm the peptide-level HDX-MS data and
complement this with single residue resolution 3pESEEM data for crucial sites where changes may
have not been detected, due to limitations associated with peptide-level analysis in HDX-MS.
3pESEEM measurements at position L42, located in the periplasmic loop, showed no significant
difference in solvent accessibility between the WT and the L89W states. For N13 which points
towards the cytoplasm and is solvent-exposed in the closed state, we observed similar exposure
in the expanded L89W MscL state (Fig 3 and S7). In L89W MscL, the solvent accessibility increased
for V21, which is consistent with channel pore hydration as its side chain sits within the TM domain
in the closed state (Fig S7). An over ten-fold increase in solvent accessibility was observed for L72,
while a significant decrease was observed for K100, which sits at the C-terminal end of TM2 (Fig
3, S5 and S6). A decrease in solvent accessibility was observed for N70 and V71 in L89W MscL
with 2.4% and ~19% reduction, respectively. Despite being consecutive residues, 70-73 show
distinct changes in solvent accessibility in 3pESEEM, unlike HDX-MS, which reported an average
difference in deuterium uptake at the peptide level. A significant and smaller increase in
accessibility is seen for L72 and L73, which is opposite to the effect observed for N70 and V71,
while the bottom of TM2 (K100) becomes less accessible to solvent in the L89W background (Fig
3, S5 and S6). This result is consistent with a rotation of the top of TM2 which occurs during
mechanical activation and agrees with PELDOR distance measurements and x-ray structures(8,
19, 56). HDX-MS experiments showed deprotection on peptides inclusive of these residues and is
consistent with a dramatic increase in accessibility of residue L72 upon pocket lipid removal
activation (i.e. when the L89W mutation is present) in 3pESEEM measurements (Fig 3). V21 forms
the vapour-lock for TbMscL and shows an increase in solvent accessibility in an L89W background,
consistent with the pore gate hydration and the channel entering an expanded (and intermediate
for full opening) state. HDX-MS measurements show a deprotection of the TM2 bottom for L89W
versus WT suggesting an overall solvent exposure of this region (Fig 1 and 3). The MscL
cytoplasmic loop has been shown to participate in gating and provide access to streptomycin
entering the cell, while shortening of the loop influences MscL's function by irreversibly decreasing
its conductance(16, 60). Combined, HDX-MS and 3pESSEM allowed us to identify entire regions
and individual residues on MscL, which undergo substantial conformational changes upon pocket
delipidation (L89W background).
224
Membrane stretching promotes pore hydration in the WT channel
To mimic the naturally occurring activation of MscL by membrane tension, we implemented MD
simulations on the WT TbMscL in lipids, with and without tension applied to the lipid bilayer. We
performed all simulations with the same lipid composition in order to compare the pore properties
and overall channel architecture between the states. Our simulations were initiated from the
TbMscL X-ray structure (PDB 2OAR)(55) carrying no modifications. Previously, significantly higher
bilayer tension was required to open MscL in MD simulations over short periods of simulation
time(37-39), in comparison to the lower tension (~67.5 mN/m) and longer simulation times (300 ns)
used in this study, allowing us to capture intermediate states.
WT TbMscL was inserted in a POPC bilayer and simulations for 300 ns were performed in which
the x- and y- pressure of the box was set at -50 bar creating constant bilayer tension. During the
simulations, the bilayer could expand to adopt to the expansion occurring in the xy plane. As a
result, the bilayer thickness decreased approximately 1.2 nm compared to the POPC bilayer
without tension (Fig 4B and 5). The channel underwent large conformational changes reaching an
overall RMSD of ~14 Å compared to the initial closed structure to adapt to the changes in the bilayer
by tilting its TM helices towards the xy plane (Fig 4A, S8A and Video S1). In particular, TM1 and
TM2 gradually tilted by 30 and 15 degrees respectively towards the end of the simulations, while
the pressure-sensitive pocket surface and the area exposed to the bilayer decreased dramatically

(Fig 4, S8B and C). Increasing the timescales of the simulations was outside the scope of this
study, as our main interest was to trap and investigate in detail this intermediately open state and
the first stop in the MscL's tension mediated activation pathway. This has enabled us to compare
this model of a tension activated state with our experimentally derived state obtained by pocket
lipid removal, in the absence of membrane tension (8).
Analysis of pore hydration profiles using CHAP(61) showed opening and hydration of the WT
channel pore under tension in all of our simulations (Fig 5A). Whilst the higher (compared to
biologically relevant) membrane tension does not allow us to provide information about the
timescales of the channel transition from the closed to the hydrated state, it enables us to obtain
structural data for a key intermediate state, which has a hydrated partially open pore, essential in
MscL's gating process. For these pore hydration events we monitored the annular lipids, which
make direct contact or reside in close proximity to the channel and simultaneously interrogated the
pore radius profile using HOLE(62) (Fig 5B, C and 6C). For WT, we first observed that following
constant tension application for ~200 ns, the pore becomes hydrated, and the channel undergoes
major structural rearrangements (Fig 5A). Although a significant membrane thinning (~1.2 nm)
occurs, L89W's pore cannot open to allow water molecules to flow through and the pore radius is
still restricted to ~1 Å, despite similar global structure changes as the WT channel (Fig 5 B,C,S8
and Table S2). Following bilayer stretching the total pocket surface area decreases and limits
access and availability to bilayer lipids throughout our simulations (Fig 5B). Overall, the under-
tension WT MscL MD data agree well with the HDX-MS and 3pESEEM accessibility data obtained
for L89W (pocket delipidation by modification). In both cases, similar channel regions become
accessible to solvent suggesting a high similarity between the obtained expanded states (Fig 7A).
265
Safety-pin acting trapped lipids in the pockets do not allow MscL pore hydration
A direct consequence of the lipids-move-first model is that if lipids are trapped within the pockets,
then the channel pore would not be able to open under membrane tension application. To test this
hypothesis, we performed MD simulations with L89W TbMscL under bilayer tension and the same
conditions we used for the WT protein (POPC lipid bilayer, pressure value and simulation time).
We first tested whether L89W modification causes any distortion on TbMscL's secondary structure.
To this end, we calculated an overall RMSD of 1.8 Å between the L89W and WT TbMscL following
equilibration in lipids (and in the absence of applied tension), suggesting there is negligible impact
on MscL's structure imposed by the L89W mutation. We observed that under bilayer tension L89W
undergoes similar global structural changes as the WT TbMscL. In all our simulations modified
MscL's pore does not become hydrated, even under the tension conditions which previously
promoted WT pore hydration (Fig 5A, 7B, S8 and Table S2).
We observed that during the equilibration of our simulations, lipids intercalated into the pockets and
occupied them. Following tension application, lipids were pulled towards the membrane, but the
bulky tryptophan modification at the entrance disrupted the lipid exit path, despite tension
application sufficient to open the WT channel was applied (Fig 5). Interestingly, in the tension
activated state, F79 and L89 from adjacent subunits come very close to form two new smaller
pockets in the inner- and outer- leaflets of the now substantially thinner membrane (Fig 6A and B).
This spatial "refinement" may facilitate the next stage in MscL's activation, which follows this sub-
conducting state, and requires additional and highly localised pressure for full pore opening of 3 nS
and ~30 Å (2, 34, 35). This suggests that the expanded MscL state is an intermediate stop in the
tension mediated activation path and not the final destination, e.g. as in activation caused by pocket
lipid removal of MscL. Asymmetry in bilayer tension sensitivity has been previously observed for
MscL, MscS and other mammalian MS channels (63, 64) and we observed that inner-leaflet lipids
previously occupying MscL's pockets to keep the channel closed, could subsequently penetrate
either of the two newly created sub-pockets (Fig 5B, 6A and B). This change restricts the access
of new lipids from the bulk membrane to the pockets and immobilizes trapped lipids within the
pockets that entered initially during equilibration. In such a case these lipids should interact more
strongly with residue 89 in the modified (W89) than the WT (L89) channel. To test this hypothesis,
we calculated the pairwise energy forces between all lipids and residue 89, L and W, for the WT
and modified channels respectively (Table 1). We found that the lipid interaction (Lennard-Jones)

with residue 89 is twice as strong in the modified compared to WT channel (Table 1). This suggests
that lipids are "stuck" by the modification in the pockets, reducing their ability to exchange with the
bulk bilayer during tension application (Fig 6A and B). Despite the substantial global structural
rearrangements, we observed in L89W MscL under tension, when lipids remain tightly associated
with the pockets, its pore cannot hydrate (Fig 5 and 6C).
302
Lipid order and accessibility during tension application
To assess the lipid contact profile with the MscL TM domain we calculated the relative difference
of lipid contacts between the with and no tension TbMscL states (Fig 6 and S9). We found that the
relative lipid contacts within bilayer facing residues on the TM domain were reduced over the course
of the simulation. In contrast, the lipid contacts of hydrophobic residues that were not lipid exposed
in the closed state increased substantially (Fig 6 and S9). This is consistent with TM2 rotation
occurring during MscL opening(8) and with pockets decreasing their surface contact area and
becoming less accessible to lipids (Fig 4B). To understand lipid rearrangement occurring upon
tension application we calculated the lipid order before and after bilayer tension application and
found that all lipids (particularly the end of the lipid chains) orient substantially more horizontally in
the presence of tension (Fig S10A and C). However, the pocket lipids specifically orient marginally
more horizontal compared to the bulk bilayer lipids, in the presence of tension (Fig S10B). These
findings suggest that although tension application results in a more horizontal, in respect to the
membrane plane, lipid reorientation, additional horizontal tilting other than that caused by
membrane stretching is not required to favor lipid pocket preference or specificity (8, 10).
318
319 **Discussion**
320
According to the lipids move first model(2), MS channels respond and open their pore when under
membrane tension or when lipid occupancy within the pockets is reduced. Equally the channel's
pore will not open, if lipids are trapped within the pockets. It was not previously known whether the
functional subconducting intermediate states adopted by MscL upon either membrane tension or
pocket delipidation were analogous in MscL. Transitions between functional states of MscL may
follow a similar path to cover the available conformational space between these states but may not
necessarily sample the exact same discrete stable intermediates. Here we find that both activation
mechanisms (Mechanical or pocket lipid removal), result in MscL adopting a well-defined,
expanded state.
Our accumulative data suggest this could be the final destination of the pocket delipidation path,
but only an intermediate stop of the tension mediated activation path. Therefore, pocket lipid
removal constitutes a major component of the initial stages (to open or not to open) of tension-
mediated mechanical activation in MscL, consistent with pocket lipids acting as safety-pins of a
grenade (8). Following bilayer tension application to WT and L89W channels, we observe similar
global structural rearrangements (Fig 5 and 7). Despite this, only the WT channel pore increased
in diameter enough for pore hydration. The pore of the L89W channel was significantly smaller in
volume and diameter (Fig 2,3,5,6C and S8). Other previously reported loss-of-function or gain-of-
unction mutations at, or in proximity to the lipid binding pockets interfere with MscL protein-lipid
interactions and may stabilize MscL states by restricting lipid access to the pockets (14, 15).
Previously, when lipids were sterically blocked from entering the pockets, the probability of pore
opening increased and the activation threshold significantly decreased(8). Equivalently, in our
present MD simulations, the L89W mutation locks the exit door and traps the lipids in the pockets
by restricting the space available for lipids to move out. Consequently, this disallows channel pore
hydration under applied tension, which we indeed observe in our simulations of L89W TbMscL
(under membrane tension) (Fig 6A, B and Table 1). Lipids are loosely associated with the pockets
in the "hydrated" WT MscL, following tension activation, but these low pairwise force-contacts are
not sufficient to close the channel (Fig 5B, 6B and Table 1). We find that the pockets become
substantially smaller under tension and consequently less accessible to lipids (Fig 4B, 5B and 6).
A similar activation mechanism has been suggested for the small conductance MS channel MscS
where inner-leaflet lipids residing within the pockets are required to move before outer-leaflet lipids

can disengage to allow gating (25), and the degree of their occupancy within the pockets was linked
with distinct MscS states(17, 31). This agrees with our entropy-driven "lipids-move-first model" and
with pocket lipids being able to exchange with the bulk bilayer despite its distant positioning away
from the pockets (2, 17, 30, 31, 65-67).
In the modified L89W MscL, F79 and W89 form a "molecular" bridge resulting in a two-compartment
repartitioning of the smaller pockets (Fig 6A and B). The WT (under tension) expanded state is
substantially different from the closed state x-ray structure used (PDB 2OAR) (Fig 4, S8 and Video
S1). Importantly, TM2 movement (translation and expansion due to anticlockwise TM2 rotation)
under tension like that occurring upon pocket removal activation, consistent with the archaeal x-ray
structure of expanded MscL (56), and the expanded (and sub-conducting) state, revealed by
PELDOR (2) (Fig 7A).
Previously, cross-sectional expansions not associated with a conducting pore have been proposed
for the initial stages of MscL activation(41, 68), suggesting MscL could act as a membrane tension
dampener and undergo several transitions with the pore closed (69). Our data support the notion
that there are structural transitions that occur prior to channel opening and agree with these models.
We observed transitions which are part of the closed-closed expansion gating process, since the
overall RMSD between the L89W and WT structures following tension application was small, but
they were both significantly different from the closed state (PDB 2OAR) (Fig 4, 7B and S8). This is
substantiated by the fact that the channel is unable to reach the intermediate state under applied
tension after the lipids are trapped within the pockets *via* the L89W modification, by preventing pore
hydration and stabilizing a closed (non-hydrated pore) state (Fig 3, 5, 7).
Our data suggest that for MscL to reach the intermediate hydrated state, triggered either by tension
or molecules, the lipids will have to substantially loosen their contacts with the pockets (Fig 5 and
Table 1). Lipids could move under tension and adopt a more horizontal orientation in respect to the
membrane plane in the WT channel, in the absence of pocket modifications that disrupted lipid
exchange with the bulk bilayer (Fig S10). The addition of molecules which could compete for the
pockets may support or disrupt lipids acting as negative allosteric modulators within the pocket
sites (Fig 7C). Indeed, following similar modifications on MscL, which restrict lipid access to the
pockets, or when molecules that specifically bind these pockets (and compete with lipids) were
introduced, there was a reduction in MscL's activation threshold, prolonged openings, increased
open channel probability (8, 18, 70).
Disruption of lipid access into the TbMscL's pockets promoted a mechanical response in the
absence of applied tension, while subtle structural differences between these pockets in MscL
orthologues dramatically altered their function (8, 19). This pocket region in TbMscL is important
for lipid binding (2, 15) and binds specific molecules in EcMscL, which could modulate its
activity(16, 18).
Our model is consistent with the notion that other modifications or sites near or within the pocket
region could be stabilising MS state(s) by disrupting lipid access to the pockets even more
effectively than the L89W modification. Indeed, TbMscL L89 is located at the entrance of the
pockets and is structurally aligned to EcMscL M94 and in close proximity to I92, F93, I96 and K97,
modifications of which, influence MscL's activation by ultrasound pulses(71) and decrease its
tension threshold for use in a multicompartmental lipid vesicle framework(72). Molecules specific
to MscL are expected to act through disruption of lipid-protein interactions(32) within these pockets.
These did not fully open the channel, but caused a structural rearrangement of TM2's cytoplasmic
side accompanied by stretching of the cytoplasmic loop(16, 18). This is in agreement with the
structural changes observed by HDX and ESEEM for the modified L89W TbMscL channel (or the
structurally equivalent M94 EcMscL) and the tension activated WT channel (Fig 1,3,4,7 and S8).
V21R1 exposure increases in the intermediate (L89W) MscL state, and this increase is consistent
with a channel pore hydration and opening. N13R1's side chain is already solvent exposed in the
closed state, as it is pointing into the cytoplasm (Fig S7), and thus unaffected in the expanded
intermediate state (Fig 3).
Transition to the intermediate state increased solvent accessibility on the upper TM1 and TM2, the
cytoplasmic loops and the top portion of the cytoplasmic helical bundle (Fig 1 and 2), consistent

with stretching of the cytoplasmic loops, of which shortening influences MscL's gating properties,
pore conductance and oligomeric assembly(16, 42, 60, 73).
The top of TM1 (residues 36 - 41) partially unfolds within a single subunit of the MscL pentamer,
while the remaining four subunits bend without fully converting into a loop. This finding was
consistently observed across all three MD simulation repeats following the symmetric application
of tension along the membrane xy plane and suggests the presence of a subunit asymmetry during
MscL opening(41, 74, 75). In contrast to the WT channel, all five TM1 helices bend without a break
in helical symmetry or conversion into a loop in the L89W channel (Fig 4). For the top of the TM1
helix to partially unfold, lipids should become loosely associated with the pockets before
asymmetric opening could occur and eventually lead to full channel opening. Whether the lipids
are sequentially released (or loosen their contact) following single subunit disengagement remains
to be elucidated. Shifting of the equilibrium occurs in order to destabilize the closed state and
initiate symmetric (simultaneous) or asymmetric (sequential) pocket lipid removal from the five
MscL subunits. Our data are compatible with such an asymmetry in opening and this structural
effect may be the consequence of lipids moving out from MscL's five pockets sequentially(75).
Subsequently, following initial expansion and conversion into loops, these loops could then provide
membrane handles for MscL in order to transit to its full opening state (Video S1).
Despite a significant membrane thinning (~1.2 nm), L89W's pore does not allow water molecules
to flow through (Fig 4 and 5). Membrane thickness and hydrophobic mismatch energetically
contribute to MscL's gating, but our data suggest they are not the sole driving force and pocket
lipids have to be released first in order to destabilize MscL's closed state. However, it should be
noted that while tension application and channel gating occur within the 300 ns that our simulations
run, this may be faster compared to the timescale of MscL opening transitions *in situ*.
Lipids must adopt certain angles in respect to the membrane plane in order to enter (or exit) the
pockets and these angles should be consistent with lipids oriented horizontally, contrary to the bulk
bilayer lipids. Such lipids have been resolved in multiple x-ray and cryoEM structures of MscS(2,
17, 30), YnaI(28), MSL1(29), TRAAK(26), TREK-2(27) and TRPV3(25) channels. When we
calculated the lipid order parameters for our MD simulations, we found that under tension
application the lipids adopt a substantially more horizontal orientation (Fig S10). This is consistent
with significantly lower pulling forces required to remove single lipids from TbMscL's pockets when
they are applied at 45° in respect to the membrane plane(76).
In conclusion, here we have generated tension- and modification- activated MscL states and
interrogated their pore hydration properties, lipid-protein interactions and monitored their structural
dynamics by experimental (HDX-MS and 3pESEEM solvent accessibility measurements) and
computational (MD simulations) tools. Accumulatively, our data demonstrate that the intermediate
expanded MscL state derived by pocket delipidation (L89W) is structurally analogous to the tension
activated MscL state (mechanical)(Fig 7C). Our data further suggest that the expanded MscL state
is the final destination of MscL's pocket lipid removal activation path, but only an intermediate stop
for the tension mediated activation path, which leads to a fully open MscL state. It is quite intriguing
that whether it is by tension application or pocket lipid depletion, we find that MscL is sampling
similar defined states, as is also the case for MscS, whose states are also guided by pocket-lipid
availability (17).
The structural similarities of these two differently derived states suggest that lipids could act as
molecular triggers on specific sites and mimic the naturally occurring tension activation in MS
channels, with implications for MS channel evolution and multimodality.
449
450 **Materials and Methods**
451
452 Materials
453
N-Dodecyl-b-D-Maltopyranoside (DDM) anagrade was obtained from Anatrace or Glycon
(Germany). Isopropyl-β-D-thiogalactoside (IPTG) was obtained from Formedium, and (Tris(2-
carboxyethyl)phosphine (TCEP) was obtained from Thermo Scientific. The S-(2,2,5,5-tetramethyl-
2,5-dihydro-1H-pyrrol-3-yl)methyl methanesulfonothioate (MTSSL) spin label was obtained from

Toronto Research Chemicals. Lipids were purchased from Avanti Polar Lipids and Biobeads from
Biorad. All other chemicals unless otherwise stated were obtained from Sigma.
460
461
462 Methods
463
Mutagenesis and Protein Expression
Mutagenesis of TbMscL with a C-terminal 6xHis-tag in a pJ411:140126 vector allowed the
generation of single and double mutants. Primers were used for site-directed mutagenesis
reactions to introduce the point mutations. The wild type and mutant plasmids were transformed
into BL21(DE3) (ThermoFisher) E. coli cells. Cells were grown in 550 ml LB medium in a 2L flask
at 37 °C until they reached an OD600 ~0.8 and subsequently cooled down to 25 °C and induced
with 1 mM IPTG for 3.5 hrs. Cells were harvested by centrifugation at 4000 × g and stored at -
80 °C, until further use.
472
Protein Purification and Spin Labelling
The protocols followed in this study were similar to the ones previously described(8, 19, 35). In
brief, following protein expression, cell pellets were resuspended in phosphate-buffered saline and
subjected to lysis using a cell disrupter at 30 kpsi. To remove cell debris the suspension was
centrifuged at 4000 × g for 20 min and the resulting supernatant was centrifuged again at
100,000 × g for 1 h. The membrane pellet was resuspended and solubilized in buffer containing
50 mM sodium phosphate at pH 7.5, 300 mM NaCl, 10% v/v glycerol, 50 mM imidazole and 1.5%
w/v DDM (solubilization buffer) and incubated at 4 °C for 1 h. The sample was then centrifuged at
4000 × g for 20 min and the supernatant was passed through a +2Ni-NTA column containing
0.75 mL of +2Ni-NTA beads. The column was then washed with 10 mL buffer containing 50 mM
sodium phosphate of pH 7.5, 300 mM NaCl, 10% v/v glycerol and 0.05% w/v DDM (wash buffer)
and then with 5 mL wash buffer supplemented with 3 mM TCEP, for reduction of the cysteines.
Afterwards, MTSSL dissolved in wash buffer at a 10-fold excess of the expected protein
concentration was added to the column and left to react for 2 h at 4 °C. The protein was then eluted
from the column with 5 mL of wash buffer supplemented with 300 mM imidazole. Finally, the protein
was subjected to SEC using a Superdex 200 column (GE Healthcare) equilibrated with buffer
containing 50 mM sodium phosphate of pH 7.5, 300 mM NaCl and 0.05% w/v DDM. Collected
fractions of TbMscL were then concentrated to ~800 µM monomer concentration, which is suitable
for the EPR samples preparation
492
493 3pESEEM sample preparation
494 Purified TbMscL detergent or reconstituted samples were diluted by 50% with deuterated ethylene
495 glycol as cryoprotectant to a final monomer concentration of ~400 µM and a volume of 70 µL of the
496 mixture was then transferred into a 3 mm (OD) quartz tubes and flash frozen in liquid N2.
497
3pESEEM spectroscopy
X-band three pulse ESEEM measurements were performed on a Bruker ELEXSYS E580
spectrometer with a 4 mm dielectric resonator (MD4). Measurements were conducted at 80 K. The
3-pulse sequence used for the experiments is $\pi/2$-τ-$\pi/2$-T-$\pi/2$-stimulated echo with a pulse length
$t_{\pi/2}$ = 16 ns and inter-pulse delay τ that was adjusted to match either the blind spots of the proton
or deuterium. The delay T was incremented from 400ns in 12 ns steps. A four-step phase cycling
was used to eliminate the unwanted echoes. All the measurements were performed at the
maximum of the filed sweep spectrum of the nitroxide. For solvent accessibility determination, only
-3pESEEM data recorded with τ that corresponds to the proton blind spot were used. The obtained
time-domain traces were background-corrected, apodized with a hamming window and zero-filled
prior to Fourier transformation. The solvent accessibility can be determined by different analysis
methods from the deuterium 3pESEEM. In the present study, we used a model developed by
Jeschke and co-workers and it is based on deuterium modulation depth(51). The primary ESEEM

data was background corrected in a such way to obtain a deconvoluted and normalized nuclear
modulation function. We used two approaches to determine the solvent accessibility; the first one
by fitting the obtained nuclear modulation function for each mutant by a damped harmonic
oscillation function and the outcome from the fitting was used to estimate the solvent
accessibility(51) , the second approach by Fourier transformation of the nuclear modulation function
and in this case the water accessibility was determined directly form the intensities of the deuterium
peaks in the magnitude ESEEM spectra. The solvent accessibility parameters derived from both
methods are in a good agreement within the error bars as shown in Fig S3. Although the standard
deviations for the fitted parameters were less than 2% for all the mutants, we additionally accounted
for errors that might emanate from differences in relaxation behavior or in contribution from no-
water protons between the different mutants and use an error of 5%. All ESEEM raw data for this
study are within the following link: http://archive.researchdata.leeds.ac.uk/777/
523
HDX-MS sample preparation
For HDX WT or L89W mutant MscL were not spin-labelled. The cell disruption, membrane
preparation, solubilization, and application of the solubilized membranes to the Ni-NTA column
occurred as previously. The column was washed, as previously, with 10 mL buffer containing
50 mM sodium phosphate of pH 7.5, 300 mM NaCl, 10% v/v glycerol and 0.05% w/v DDM (wash
buffer). Following this step, the protein was eluted with 5 ml of wash buffer containing 300 mM
imidazole. The protein was not treated with TCEP or MTSL supplemented wash buffer as the
protein was not spin-labelled. The protein was then subjected to SEC using a Superdex 200 column
as previously described. The fractions were collected, and the protein samples were concentrated
to achieve 350 μl of protein at 16 μM.
534
HDX and LC-MS/MS
HDX-MS experiments were carried out using an automated HDX robot (LEAP Technologies, Fort
Lauderdale, FL, USA) coupled to an M-Class Acquity LC and HDX manager (Waters Ltd.,
Wilmslow, Manchester, UK). All samples were diluted to 16 μM in equilibration buffer (50 mM
potassium phosphate, 300 mM NaCl, pH 7.4, 0.05% DDM) prior to analysis. 5 μL sample was
added to 95 μl deuterated buffer (50 mM potassium phosphate, 300 mM NaCl pD 7.4, 0.05% DDM)
and incubated at 4 °C for 0.5, 1, 2, 10 or 60 min. Following the labelling reaction, samples were
quenched by adding 50 μl of the labelled solution to 100 μl quench buffer (50 mM potassium
phosphate, 300 mM NaCl, 0.1% DDM pH 2.2) giving a final quench pH ~2.5. 50 μl of quenched
sample was injected on to the HDX Manger and passed through immobilized pepsin and
aspergillopepsin columns connected in series (AffiPro, Czech Republic) at 115 μl min-1 (20 °C)
and loaded on to a VanGuard Pre-column Acquity UPLC BEH C18 (1.7 μm, 2.1 mm X 5 mm,
Waters Ltd., Wilmslow, Manchester, UK) for 3 mins in 0.3 % formic acid in water. The resulting
peptides were transferred to a C18 column (75 μm X 150 mm, Waters Ltd., Wilmslow, Manchester,
UK) and separated by gradient elution of 0-40 % MeCN (0.1 % v/v formic acid) in H2O (0.3 % v/v
formic acid) over 7 mins at 40 μl.min^{-1}. Trapping and gradient elution of peptides was performed at
0 °C to reduce back exchange. The HDX system was interfaced to a Synapt G2Si mass
spectrometer (Waters Ltd., Wilmslow, Manchester, UK). HDMSE and dynamic range extension
modes (Data Independent Analysis (DIA) coupled with IMS separation) were used to separate
peptides prior to CID fragmentation in the transfer cell (77). HDX data were analyzed using PLGS
(v3.0.2) and DynamX (v3.0.0) software supplied with the mass spectrometer. Restrictions for
identified peptides in DynamX were as follows: minimum intensity: 1000, minimum products per
MS/MS spectrum: 5, minimum products per amino acid: 0.3, maximum sequence length: 25,
maximum ppm error: 5, file threshold: 3/3. Following manual curation of the data, PAVED(78) and
Deuteros(79) were used to identify peptides with statistically significant increases/decreases in
deuterium uptake (applying 99% or 95% confidence intervals) and to prepare Wood's plots. The
raw HDX-MS data, have been deposited to the ProteomeXchange Consortium via the PRIDE
partner repository with the dataset identifier PXD021983. A summary of the HDX-MS data, as
recommended by reporting guidelines, is shown in Table S1. All data is available via
ProteomeXchange with identifier PXD021983. Project Name: Mechanical and molecular activation

lead to structurally analogous MscL states. Project accession: PXD021983 Username:
reviewer_pxd021983@ebi.ac.uk, Password: lnAkk2H3
567
568
Atomistic molecular dynamics simulations under no tension
The set up and parameters of MD simulations without tension were as previously described(8). In
brief, CHARMM-GUI was used to insert the TbMscL structure (2OAR) into a pre-equilibrated patch
of POPC bilayer containing approximately 387 lipids and occupying an area of 120 × 120 Å2. The
protein and membrane bilayer were solvated with TIP3P water and 150 mM NaCl. The simulations
were performed in an NPT ensemble at 303.15 K and 1bar pressure on all xyz axes using
GROMACS_2016.4 with CHARMM36 force field. The particle mesh Ewald (PME) method was
applied to calculate electrostatic forces, and the van der Waals interactions were smoothly switched
off at 10–12 Å by the force-switch manner. The time step was set to 2 fs in conjunction with the
LINCS algorithm. After the standard minimization and equilibration steps using the GROMACS
input scripts generated by CHARMM-GUI, 100 ns dynamic simulation was calculated.
580
Atomistic molecular dynamics simulations under bilayer tension
These MD simulations were also set up using CHARMM-GUI(80). The MscL structure (PDB 2OAR)
in the wild type and in a mutated form (L89W) was inserted in a symmetric bilayer containing 514
POPC lipids. Systems were neutralized with a 150 mM concentration of NaCl. The obtained system
was energy minimised for 5000 steps and subsequently equilibrated in 6 steps following CHARMM-
GUIs equilibration. Stretch-induced conformational changes in both the wild-type and mutant MscL
were investigated by unrestrained simulations of 300 ns (3 simulation repeats for the WT and 2 for
the L89W channel) where the bilayer plane (xy plane) pressure was changed semiisotropically to -
50 bars (which corresponds to a tension of ~67.5 mN/m). The pressure in the bilayer normal (z
direction) was kept at +1 bar and the temperature at 310 K. All the atomistic systems were
simulated using GROMACS 2016(81) with CHARMM36 force field(82) and a 2 fs time step. The
Nose-Hoover thermostat(83) and the Parrinello-Rahman barostat(84) were used for the stretch-
induced simulations. Long-range electrostatics were managed using the particle-mesh Ewald
method(85) and the LINCS algorithm was used to constrain bond lengths(86). All MD trajectories
and data for this study is within the following link: http://archive.researchdata.leeds.ac.uk/777/
596
597
598 **Data accessibility**
599
600 HDX mass spectrometry:
601 Data are available via ProteomeXchange with identifier PXD021983.
602 Project Name: Mechanical and molecular activation lead to structurally analogous MscL states
603 Project accession: PXD021983
604 Reviewer account details: Username: reviewer_pxd021983@ebi.ac.uk, Password: lnAkk2H3
605
606 3pESEEM and MD:
607 Data are available within the following link: http://archive.researchdata.leeds.ac.uk/777/
608
609 **Acknowledgments**
610

This project was supported by a Biotechnology and Biological Sciences Research Council
(BBSRC) grant (BB/S018069/1) to C.P., who also acknowledges support from the Wellcome Trust
(WT) (219999/Z/19/Z) and the Chinese Scholarship Council (CSC) in the form of studentships for
B.J.L. and B.W. respectively. ANC is a Sir Henry Dale Fellow jointly funded by the WT and the
Royal Society (220628/Z/20/Z). Funding from the BBSRC (BB/M012573/1) enabled the purchase
of mass spectrometry equipment.

617
618

References

1. Pliotas C & Naismith JH (2016) Spectator no more, the role of the membrane in regulating ion channel function. *Curr Opin Struct Biol* 45:59-66.
2. Pliotas C*, et al.* (2015) The role of lipids in mechanosensation. *Nat Struct Mol Biol* 22(12):991-998.
3. Teng J, Loukin S, Anishkin A, & Kung C (2015) The force-from-lipid (FFL) principle of mechanosensitivity, at large and in elements. *Pflugers Arch* 467(1):27-37.
4. Malcolm HR, Blount P, & Maurer JA (2015) The mechanosensitive channel of small conductance (MscS) functions as a Jack-in-the box. *Biochim Biophys Acta* 1848(1 Pt A):159-166.
5. Cox CD, Nakayama Y, Nomura T, & Martinac B (2015) The evolutionary 'tinkering' of MscS-like channels: generation of structural and functional diversity. *Pflugers Arch* 467(1):3-13.
6. Brohawn SG (2015) How ion channels sense mechanical force: insights from mechanosensitive K2P channels TRAAK, TREK1, and TREK2. *Ann N Y Acad Sci* 1352:20-32.
7. Naismith JH & Booth IR (2012) Bacterial mechanosensitive channels--MscS: evolution's solution to creating sensitivity in function. *Annu Rev Biophys* 41:157-177.
8. Kapsalis C*, et al.* (2019) Allosteric activation of an ion channel triggered by modification of mechanosensitive nano-pockets. *Nat Commun* 10(1):4619.
9. Patrick JW*, et al.* (2018) Allostery revealed within lipid binding events to membrane proteins. *Proc Natl Acad Sci U S A* 115(12):2976-2981.
10. Laganowsky A*, et al.* (2014) Membrane proteins bind lipids selectively to modulate their structure and function. *Nature* 510(7503):172-175.
11. Ridone P*, et al.* (2018) "Force-from-lipids" gating of mechanosensitive channels modulated by PUFAs. *J Mech Behav Biomed Mater* 79:158-167.
12. Ward R*, et al.* (2014) Probing the structure of the mechanosensitive channel of small conductance in lipid bilayers with pulsed electron-electron double resonance. *Biophys J* 106(4):834-842.
13. Pliotas C*, et al.* (2012) Conformational state of the MscS mechanosensitive channel in solution revealed by pulsed electron-electron double resonance (PELDOR) spectroscopy. *Proc Natl Acad Sci U S A* 109(40):E2675-2682.
14. Anishkin A, Chiang CS, & Sukharev S (2005) Gain-of-function mutations reveal expanded intermediate states and a sequential action of two gates in MscL. *J Gen Physiol* 125(2):155-170.
15. Iscla I, Wray R, Eaton C, & Blount P (2015) Scanning MscL Channels with Targeted Post-Translational Modifications for Functional Alterations. *PLoS One* 10(9):e0137994.
16. Wray R*, et al.* (2016) Dihydrostreptomycin Directly Binds to, Modulates, and Passes through the MscL Channel Pore. *PLoS biology* 14(6):e1002473.
17. Zhang Y*, et al.* (2021) Visualization of the mechanosensitive ion channel MscS under membrane tension. *Nature* 590:509-514.
18. Wray R, Iscla I, Kovacs Z, Wang J, & Blount P (2019) Novel compounds that specifically bind and modulate MscL: insights into channel gating mechanisms. *FASEB J* 33(3):3180-3189.
19. Kapsalis C, Ma Y, Bode BE, & Pliotas C (2020) In-Lipid Structure of Pressure-Sensitive Domains Hints Mechanosensitive Channel Functional Diversity. *Biophys J* 119(2):448-459.
20. Syeda R*, et al.* (2015) Chemical activation of the mechanotransduction channel Piezo1. *Elife* 4.
21. Botello-Smith WM*, et al.* (2019) A mechanism for the activation of the mechanosensitive Piezo1 channel by the small molecule Yoda1. *Nat Commun* 10(1):4503.

22. Bagriantsev SN, Peyronnet R, Clark KA, Honore E, & Minor DL, Jr. (2011) Multiple
modalities converge on a common gate to control K2P channel function. *EMBO J*
30(17):3594-3606.
23. Piechotta PL*, et al.* (2011) The pore structure and gating mechanism of K2P channels.
EMBO J 30(17):3607-3619.
24. Feliciangeli S, Chatelain FC, Bichet D, & Lesage F (2015) The family of K2P channels:
salient structural and functional properties. *J Physiol* 593(Pt 12):2587-2603.
25. Deng Z*, et al.* (2020) Gating of human TRPV3 in a lipid bilayer. *Nat Struct Mol Biol*.
26. Brohawn SG, Campbell EB, & MacKinnon R (2014) Physical mechanism for gating and
mechanosensitivity of the human TRAAK K+ channel. *Nature* 516(7529):126-130.
27. Dong YY*, et al.* (2015) K2P channel gating mechanisms revealed by structures of TREK-
2 and a complex with Prozac. *Science* 347(6227):1256-1259.
28. Flegler VJ*, et al.* (2020) The MscS-like channel YnaI has a gating mechanism based on
flexible pore helices. *Proc Natl Acad Sci U S A* 117(46):28754-28762.
29. Deng Z*, et al.* (2020) Structural mechanism for gating of a eukaryotic mechanosensitive
channel of small conductance. *Nat Commun* 11(1):3690.
30. Rasmussen T, Flegler VJ, Rasmussen A, & Bottcher B (2019) Structure of the
Mechanosensitive Channel MscS Embedded in the Membrane Bilayer. *J Mol Biol*.
31. Flegler VJ*, et al.* (2021) Mechanosensitive channel gating by delipidation. *Proc Natl Acad*
Sci U S A 118(33).
32. Bavi N*, et al.* (2016) The role of MscL amphipathic N terminus indicates a blueprint for
bilayer-mediated gating of mechanosensitive channels. *Nat Commun* 7:11984.
33. Erdogmus S*, et al.* (2019) Helix 8 is the essential structural motif of mechanosensitive
GPCRs. *Nat Commun* 10(1):5784.
34. Hartley AM, Ma Y, Lane BJ, Wang B, & Pliotas C (2021) Using pulsed EPR in the
structural analysis of integral membrane proteins. *Electron Paramag Reson* 27:74 -108.
35. Pliotas C (2017) Ion Channel Conformation and Oligomerization Assessment by Site-
Directed Spin Labeling and Pulsed-EPR. *Methods in enzymology* 594:203-242.
36. Debruycker V*, et al.* (2020) An embedded lipid in the multidrug transporter LmrP
suggests a mechanism for polyspecificity. *Nat Struct Mol Biol* 27(9):829-835.
37. Martinac AD, Bavi N, Bavi O, & Martinac B (2017) Pulling MscL open via N-terminal and
TM1 helices: A computational study towards engineering an MscL nanovalve. *PLoS One*
12(8):e0183822.
38. Gullingsrud J & Schulten K (2003) Gating of MscL studied by steered molecular
dynamics. *Biophys J* 85(4):2087-2099.
39. Bavi N*, et al.* (2017) Nanomechanical properties of MscL alpha helices: A steered
molecular dynamics study. *Channels* 11(3):209-223.
40. Jeon J & Voth GA (2008) Gating of the mechanosensitive channel protein MscL: the
interplay of membrane and protein. *Biophys J* 94(9):3497-3511.
41. Konijnenberg A*, et al.* (2014) Global structural changes of an ion channel during its gating
are followed by ion mobility mass spectrometry. *Proc Natl Acad Sci U S A*
111(48):17170-17175.
42. Herrera N, Maksaev G, Haswell ES, & Rees DC (2018) Elucidating a role for the
cytoplasmic domain in the Mycobacterium tuberculosis mechanosensitive channel of
large conductance. *Sci Rep* 8(1):14566.
43. Mukherjee N*, et al.* (2014) The activation mode of the mechanosensitive ion channel,
MscL, by lysophosphatidylcholine differs from tension-induced gating. *FASEB J*
28(10):4292-4302.
44. Katsuta H, Sawada Y, & Sokabe M (2019) Biophysical Mechanisms of Membrane-
Thickness-Dependent MscL Gating: An All-Atom Molecular Dynamics Study. *Langmuir*
35(23):7432-7442.
45. Aryal P*, et al.* (2017) Bilayer-Mediated Structural Transitions Control Mechanosensitivity
of the TREK-2 K2P Channel. *Structure* 25(5):708-718.e702.

724 46. Martens C, Shekhar M, Lau AM, Tajkhorshid E, & Politis A (2019) Integrating hydrogen-
725   deuterium exchange mass spectrometry with molecular dynamics simulations to probe
726   lipid-modulated conformational changes in membrane proteins. *Nat Protoc* 14(11):3183-
727   3204.
728 47. Ahdash Z*, et al.* (2019) HDX-MS reveals nucleotide-dependent, anti-correlated opening
729   and closure of SecA and SecY channels of the bacterial translocon. *Elife* 8.
730 48. Konermann L, Pan J, & Liu YH (2011) Hydrogen exchange mass spectrometry for
731   studying protein structure and dynamics. *Chem Soc Rev* 40(3):1224-1234.
732 49. Moller IR*, et al.* (2019) Conformational dynamics of the human serotonin transporter
733   during substrate and drug binding. *Nat Commun* 10(1):1687.
734 50. Cieslak JA, Focia PJ, & Gross A (2010) Electron spin-echo envelope modulation
735   (ESEEM) reveals water and phosphate interactions with the KcsA potassium channel.
736   *Biochemistry* 49(7):1486-1494.
737 51. Volkov A, Dockter C, Bund T, Paulsen H, & Jeschke G (2009) Pulsed EPR determination
738   of water accessibility to spin-labeled amino acid residues in LHCIIb. *Biophys J*
739   96(3):1124-1141.
740 52. Liu L*, et al.* (2016) Probing the Local Secondary Structure of Human Vimentin with
741   Electron Spin Echo Envelope Modulation (ESEEM) Spectroscopy. *J Phys Chem B*
742   120(48):12321-12326.
743 53. Zhang R*, et al.* (2015) Development of electron spin echo envelope modulation
744   spectroscopy to probe the secondary structure of recombinant membrane proteins in a
745   lipid bilayer. *Protein Sci* 24(11):1707-1713.
746 54. Matalon E*, et al.* (2013) Topology of the trans-membrane peptide WALP23 in model
747   membranes under negative mismatch conditions. *J Phys Chem B* 117(8):2280-2293.
748 55. Chang G, Spencer RH, Lee AT, Barclay MT, & Rees DC (1998) Structure of the MscL
749   homolog from Mycobacterium tuberculosis: a gated mechanosensitive ion channel.
750   *Science* 282(5397):2220-2226.
751 56. Li J*, et al.* (2015) Mechanical coupling of the multiple structural elements of the large-
752   conductance mechanosensitive channel during expansion. *Proc Natl Acad Sci U S A*
753   112(34):10726-10731.
754 57. Powl AM, East JM, & Lee AG (2005) Heterogeneity in the binding of lipid molecules to
755   the surface of a membrane protein: Hot spots for anionic lipids on the mechanosensitive
756   channel of large conductance MscL and effects on conformation. *Biochemistry*
757   44(15):5873-5883.
758 58. Michou M, Kapsalis C, Pliotas C, & Skretas G (2019) Optimization of Recombinant
759   Membrane Protein Production in the Engineered Escherichia coli Strains SuptoxD and
760   SuptoxR. *ACS Synth Biol* 8(7):1631-1641.
761 59. Bavi N*, et al.* (2017) Structural Dynamics of the MscL C-terminal Domain. *Sci Rep*
762   7(1):17229.
763 60. Yang LM*, et al.* (2012) Three routes to modulate the pore size of the MscL
764   channel/nanovalve. *ACS Nano* 6(2):1134-1141.
765 61. Klesse G, Rao S, Sansom MSP, & Tucker SJ (2019) CHAP: A Versatile Tool for the
766   Structural and Functional Annotation of Ion Channel Pores. *J Mol Biol* 431(17):3353-
767   3365.
768 62. Smart OS, Goodfellow JM, & Wallace BA (1993) The pore dimensions of gramicidin A.
769   *Biophys J* 65(6):2455-2460.
770 63. Perozo E, Kloda A, Cortes DM, & Martinac B (2002) Physical principles underlying the
771   transduction of bilayer deformation forces during mechanosensitive channel gating. *Nat
772   Struct Biol* 9(9):696-703.
773 64. Belyy V, Kamaraju K, Akitake B, Anishkin A, & Sukharev S (2010) Adaptive behavior of
774   bacterial mechanosensitive channels is coupled to membrane mechanics. *J Gen Physiol*
775   135(6):641-652.

65. Anishkin A, Akitake B, & Sukharev S (2008) Characterization of the resting MscS:
modeling and analysis of the closed bacterial mechanosensitive channel of small
conductance. *Biophys J* 94(4):1252-1266.
66. Reddy B, Bavi N, Lu A, Park Y, & Perozo E (2019) Molecular basis of force-from-lipids
gating in the mechanosensitive channel MscS. *Elife* 8.
67. Kefauver JM, Ward AB, & Patapoutian A (2020) Discoveries in structure and physiology
of mechanically activated ion channels. *Nature* 587(7835):567-576.
68. Betanzos M, Chiang CS, Guy HR, & Sukharev S (2002) A large iris-like expansion of a
mechanosensitive channel protein induced by membrane tension. *Nat Struct Biol*
9(9):704-710.
69. Boucher PA, Morris CE, & Joos B (2009) Mechanosensitive closed-closed transitions in
large membrane proteins: osmoprotection and tension damping. *Biophys J* 97(10):2761-
2770.
70. Iscla I, Wray R, Wei S, Posner B, & Blount P (2014) Streptomycin potency is dependent
on MscL channel expression. *Nat Commun* 5:4891.
71. Ye J*, et al.* (2018) Ultrasonic Control of Neural Activity through Activation of the
Mechanosensitive Channel MscL. *Nano Lett* 18(7):4148-4155.
72. Hindley JW*, et al.* (2019) Building a synthetic mechanosensitive signaling pathway in
compartmentalized artificial cells. *Proc Natl Acad Sci U S A* 116(34):16711-16716.
73. Reading E*, et al.* (2015) The Effect of Detergent, Temperature, and Lipid on the
Oligomeric State of MscL Constructs: Insights from Mass Spectrometry. *Chem Biol*
22(5):593-603.
74. Birkner JP, Poolman B, & Kocer A (2012) Hydrophobic gating of mechanosensitive
channel of large conductance evidenced by single-subunit resolution. *Proc Natl Acad Sci*
U S A 109(32):12944-12949.
75. Mika JT, Birkner JP, Poolman B, & Kocer A (2013) On the role of individual subunits in
MscL gating: "all for one, one for all?". *FASEB J* 27(3):882-892.
76. Vanegas JM & Arroyo M (2014) Force transduction and lipid binding in MscL: a
continuum-molecular approach. *PLoS One* 9(12):e113947.
77. Cryar A, Groves K, & Quaglia M (2017) Online Hydrogen-Deuterium Exchange Traveling
Wave Ion Mobility Mass Spectrometry (HDX-IM-MS): a Systematic Evaluation. *J Am Soc*
Mass Spectrom 28(6):1192-1202.
78. Cornwell O, Radford SE, Ashcroft AE, & Ault JR (2018) Comparing Hydrogen Deuterium
Exchange and Fast Photochemical Oxidation of Proteins: a Structural Characterisation of
Wild-Type and DeltaN6 beta2-Microglobulin. *J Am Soc Mass Spectrom* 29(12):2413-
2426.
79. Lau AMC, Ahdash Z, Martens C, & Politis A (2019) Deuteros: software for rapid analysis
and visualization of data from differential hydrogen deuterium exchange-mass
spectrometry. *Bioinformatics* 35(17):3171-3173.
80. Jo S, Kim T, Iyer VG, & Im W (2008) CHARMM-GUI: a web-based graphical user
interface for CHARMM. *J Comput Chem* 29(11):1859-1865.
81. Abraham MJ, T. Murtola, R. Schulz, S. Páll, J.C. Smith, B. Hess, E. Lindah, and E.
Lindahl (2015) GROMACS: High performance molecular simulations through multi-level
parallelism from laptops to supercomputers. *SoftwareX* 1-2:19-25.
82. Lee S*, et al.* (2014) CHARMM36 united atom chain model for lipids and surfactants. *J*
Phys Chem B 118(2):547-556.
83. Evans DJ, Holian B.L. (1985) The Nose-Hoover thermostat. *J Chem Phys* 83:4069–4074.
84. Parrinello M, Rahman A. (1981) Polymorphic transitions in single crystals: A new
molecular dynamics method. *J. Appl. Phys.* 52:7182–7190.
85. Darden T, York, D., Pedersen L. (1993) Particle mesh Ewald: An N·log(N) method for
Ewald sums in large systems. *J Chem Phys* 98:10089–10092.
86. Hess B, H. Bekker, H.J.C. Berendsen, J.G.E.M. Fraaije (1997) INCS: A linear constraint
solver for molecular simulations. *J. Comput Chem* 18:1463–1472.

829
830
831
832 **Figures and Tables**

A

B

Figure 1. Global changes on MscL induced by pocket lipid removal observed by HDX-MS. A.
Differences in the deuterium uptake of regions of TbMscL (PDB 2OAR) when comparing the WT
and the L89W (pocket lipid removal) modified protein. Regions highlighted in red are deprotected
following the L89W modification. Regions of the protein in grey show no significant difference
between the two conditions. L89W modification site is depicted as a cyan sphere. B. Wood's plots
showing the summed differences in deuterium uptake in MscL over all five HDX timepoints,
comparing WT with L89W MscL (Wood's plots were generated using Deuteros(79). Peptides
coloured in red, are deprotected from exchange in L89W MscL. No peptides were significantly
protected from exchange in L89W MscL compared with wild type MscL. Peptides with no significant
difference between conditions, determined using a 99% confidence interval (dotted line), are shown
in grey. Example deuterium uptake curves are shown in Fig S1.
857

A

B

Figure 2. Single-residue mapping of MscL by 3pESEEM. A. Solvent accessibility of spin-labelled residues are shown as red and pale-yellow spheres for the most and the least solvent accessible, respectively. Spin-labelled residues are coloured based on the quartile of their relative accessibility which has been normalised on scale of 0% and 100%, where the residue with the highest accessibility corresponds to 100%. B. Background-corrected time-domain 3pESEEM experimental spectra with fitting for representative spin-labelled mutants. Residue F5R1 is found on the S1 amphipathic helix, I23R1 and L42R1 on TM1, and K100R1 and E102R1 are at the interface between TM2 and the CHB.

Figure 3. Effect of pocket delipidation on MscL structure investigated by integrated 3pESEEM and HDX-MS. A Background-corrected time-domain 3pESEEM experimental spectra (orange) of the single mutants N13R1, V21R1, L42R1, N70R1, L72R1, L73R1, and K100R1 in DDM overlaid with their associated L89W double mutant spectra (purple). B Differences in solvent accessibility for TbMscL (PDB 2OAR) following L89W modification. Spin labelled mutation sites used for 3pESEEM accessibility measurements are represented by spheres, and peptides that demonstrate a change in accessibility in HDX-MS following the L89W modification are represented as highlighted helices. Red regions or spheres highlight areas that are deprotected, while blue spheres and regions show areas that are protected following the L89W modification. C Column bar charts represent the 2H accessibility derived from the fitting of 3pESEEM time-domain traces. Errors are calculated conservatively at 5% to compensate for the fitting error and differences in relaxation times of different spin labelled residues. There was no significant difference in solvent accessibility of N13R1, L42R1, and N70R1 compared to their 89W double mutant counterpart. Solvent accessibility increased for V21R1, L72R1, and decreased for L71R1 and K100R1 following the L89W modification.

A

30°

TM1

15°

TM2

B

Closed
WT - no tension

Open
WT - plus tension

Periplasm

Membrane

3.8 nm

Pockets

2.6 nm

Cytoplasm

Figure 4. **Tension-mediated expanded MscL state**. A. TM comparison between closed TbMscL
(tan, PDB 2OAR) and the open state WT MscL generated by MD simulations (cyan, under tension),
side, top and single subunit views. The latter shows a substantial tilting of TM1 and TM2 towards
the membrane plane. These conformational changes occurring under tension are more evident in
Video S1. B. The pockets surface dramatically decreases upon tension application, limiting lipid
access. Uniform membrane thinning of 1.2 nm (3.8 to 2.6 nm) also occurs during channel
expansion, due to tension application.
932
933

946
Figure 5. **MscL pore hydration investigation under applied bilayer tension**. WT under no
tension (left column), WT under bilayer tension (right column), L89W under bilayer tension (right
column). A. Solvent density profile of the MD simulations using the CHAP(61) software with the
vapour-lock position of MscL marked with blue arrows. B. Membrane becomes thinner when
tension is applied but only the WT TbMscL pore is hydrated (blue spheres) in contrast to L89W
channel. Lipid (olive sticks) availability is larger in the closed state, and lipids have easier access
within the pockets to provide force on the back of the vapour lock, keeping the MscL pore closed.
This is in contrast to WT MscL under tension (middle column) where the pockets become smaller,
while lipid availability decreases, and the pore becomes hydrated. However, when lipids are
trapped in the pockets of the L89W modified channel, the pore does not become hydrated despite
structural rearrangements occurring (right column). C. Surface visualization of the pore pathway
using the program HOLE(62). Red colour indicates a pore radius smaller than 1.15 Å (water
molecules cannot go through such an opening), blue represents a radius larger than 2.3 Å and
green between 1.15 to 2.3 Å.
961
962

Figure 6. The effect of membrane tension to MscL protein-lipid contact interactions. (A) and (B). F79 and L89W form a "molecular bridge" in the L89W under tension state and consequently trap the lipids within the pockets (right column). F79 and L89 come into greater proximity under bilayer tension in the WT channel (expanded MscL, middle column) compared to the closed state. However, they do not prevent the lipids from exchanging with the bulk bilayer during tension application. The L89W mutation locks lipids in the pockets, preventing the channel from transiting to a hydrated state. C. MscL pore radius profiles analysis and comparisons using HOLE(62). Although L89W MscL undergoes similar major rearrangements to WT MscL under tension its pore remains significantly smaller and almost identical to WT closed MscL under no tension. D. Relative changes in the number of lipid contacts following stimulated tension application in the membrane during MD. The blue regions show decrease in lipid contacts, while the red regions show an increase in lipid contacts. From left to right: Surface view, single subunit cartoon representation and internal MscL surface view. Internal channel regions become more exposed to lipids, while initially membrane exposed regions (closed state) become less exposed to lipids, suggesting a MscL TM helical rotation upon tension application and channel expansion/opening.

Figure 7. Comparison between tension mediated and pocket delipidation induced MscL
states and proposed model. TM1 and TM2 tilt and rotate to initiate MscL opening. Comparison
between expanded MaMscL (orchid, PDB ID: 4Y7J) and the open state WT TbMscL generated by
MD (cyan, under tension) (A) and the open WT TbMscL state generated by MD (cyan, under tesion)
and L89W TbMscL (green, under tension) (B). C. Proposed gating model for pocket lipid removal
and tension activated modes. Closed (vertical TM helices, large pockets, tightly associated annular
lipids and non-hydrated pore), Expanded (tilted TM helices, smaller pockets, loosely associated
annular lipids and hydrated pore). Extra membrane tension is required to fully open MscL.
1006
1007

1008
1009
1010
1011 **Table 1.** Pairwise force energy calculations between TbMscL 89 (L or W) and lipids in the WT (L)
1012 and pocket-entrance modified (W) channels under tension
1013
1014

89 - POPC	Coulomb (KJ/mol)	SD - Coulomb (KJ/mol)	Lennard Jones (KJ/mol)	SD - Lennard Jones (KJ/mol)	Total Energy (KJ/mol)	SD – Total Energy (KJ/mol)
WT	-14.50	1.76	-101.39	27.43	-115.89	29.19
L89W	-14.77	3.90	-192.48	3.39	-207.25	7.29

1020

1021

1022

1023