Mechanical and molecular activation lead to structurally analogous MscL states

Bolin Wang[1,2,3], Benjamin J. Lane[1,2,3], Charalampos Kapsalis[4], James R. Ault[2,5], Frank
Sobott[2,5], Hassane El Mkami[6], Antonio N. Calabrese[2,5], Antreas C. Kalli[2,7], Christos
Pliotas*[2,3,4]

[1]Equal first author contribution

[2]Astbury Centre for Structural Molecular Biology, University of Leeds, UK

[3]School of Biomedical Sciences, Faculty of Biological Sciences, University of Leeds, UK

[4]Biomedical Sciences Research Complex, School of Biology, University of St Andrews, UK

[5]School of Molecular and Cellular Biology, Faculty of Biological Sciences, University of Leeds, UK

[6]School of Physics and Astronomy, University of St Andrews, UK

[7]Leeds Institute of Cardiovascular and Metabolic Medicine, School of Medicine, University of
Leeds, UK

* Corresponding author: Christos Pliotas, c.pliotas@leeds.ac.uk

Author Contributions: C.P. conceived and designed the project. B.W., B.J.L, C.K., J.A., H.E.M.,
A.C.K. and C.P. performed research. B.W., B.J.L, C.K., J.A., F.S., H.E.M., A.N.C., A.C.K. and
C.P. processed and analysed data. C.P. wrote the manuscript assisted by B.W., B.J.L. with
contributions from H.E.M, A.N.C. and A.C.K.

Competing Interest Statement: The authors declare no conflicts of interest

Classification: Biological Sciences/Biophysics and Computational Biology

Keywords: MscL, mechanosensitive channels, EPR, HDX, ESEEM

This PDF file includes:

Main Text

Figures 1 to 7

Tables 1 to 1

Abstract

The mechanosensitive channel of large conductance MscL gates in response to tension changes
in the membrane to allow the exchange of molecules through its pore. Native ligands that bind and
modulate MscL are unknown and trapping an activated state has been challenging. Disruption of
lipid access to tension-sensitive transmembrane pockets by modification leads to a concerted
structural and functional MscL response. However, it is unknown whether there is structural
correlation between tension mediated and molecular activation in mechanosensitive channels.
Here, we combine HDX mass spectrometry and ESEEM solvent accessibility measurements on
MscL, coupled with molecular dynamics under bilayer tension, to investigate the structural changes
associated with the two distinctively derived states. Membrane thinning is not sufficient to hydrate

the MscL pore, when lipids are trapped in the pockets, under tensions capable to gate the native channel. Tension and molecule stabilised states present analogous MscL structures, suggesting a link between these two distinct activation mechanisms. These findings could hint at synergistic modes of regulation in mechanosensitive ion channels with implications for their multimodality.

Significance Statement

Bacterial mechanosensitive channels are gated by membrane tension and their function is modulated by lipids or molecules binding to tension sensitive domains. Eukaryotic mechanosensitive channels are multimodal and regulated both by tension and molecular triggers. However, whether molecular and tension activation present mechanistic similarities and whether such mechanisms have evolved through converged structural plasticity of a common origin is largely unclear. Here we use the mechanosensitive channel of large conductance MscL and compare its two distinctive gating dynamics by experimental (EPR and HDX-MS) and computational means (MD). We find that these two differently derived states are structurally analogous suggesting links between mechanosensitivity and ligand-gating in the regulation of mechanosensitive channels and with implications for their multimodality.

Main Text

Introduction

Mechanosensitive (MS) ion channels can sense and respond to tension changes in the cell membrane. Lipids are important for the folding, stability and structural integrity of membrane proteins (1), while structural responses mediated through lipid binding to exposed protein domains are allosterically linked to their regulation (2-4). Several models place the lipids as central to the allosteric regulation of membrane proteins (3, 5-10). Previously, hydrophobic pockets formed within transmembrane (TM) domains of bacterial MS channels have been proposed to play a role in the transduction of bilayer forces to the channel, directly forming pressure-sensitive domains, or pockets (2, 3, 5). These were first identified on the MS channel of small conductance MscS, leading to the lipid moves first hypothesis, which predicts that lipids access these pockets and apply pressure to the channel gate to keep it closed (5). When a modification (L89W) was introduced at the entrance of these pockets in MscL, lipid contacts with the backside of the pore-lining helices were reduced, generating a structural response in the absence of applied tension (2). Such a model suggests that molecular triggers could act as negative allosteric modulators and that tension could be mimicked by molecules in order to activate MS channels (2, 3).
Indeed, the function of MscL (11-13) and MscS (5, 14) can be modulated by molecule occupancy changes within these pockets. Subtle structural differences within the same regions lead to substantial functional differences in orthologous MscL proteins (15). This suggests a common regulatory mechanism, tailored to the unique structural landscape of MS channels, which may guide ligand specificity (2, 3).
In mammals, the MS channel Piezo1 is activated by Yoda1, opening up the possibility of endogenous agonists (16, 17), while TRAAK and TREK-1 are responsive to natural chemical ligands including arachidonic acid, polyunsaturated fatty acids, and lysophosphatidic acid (18-21). Lipids residing within similar inner-leaflet pockets have also been linked with mechanosensitivity in x-ray and cryo-EM structures of TRPV3 (22), TRAAK (23), TREK-2 (24) and MscS (25), while their unique parallel to membrane plane orientation may be guided by an amphipathic helix that is important for mechanosensitivity in MscL (26) and GPCRs (27). Pulsed electron-electron double resonance (PELDOR, also known as DEER) and x-ray studies on the LmrP multidrug transporter

showed that lipids could provide a hydrophobic counterpart to partially fill binding pockets and
participate in the coordination of ligand binding(28).
Attempts to model MscL opening have been previously implemented via steered MD simulations,
with pulling tensions (29-33) an order of magnitude higher than the ones required to open MscL in
giant unilamellar vesicles (GUVs) and spheroplasts (2, 34, 35). In other cases MscL was subjected
to substantial modifications or forces were applied for short periods of time (29, 36-38). MD
simulations on TREK-2 have shown that under increase bilayer tension the channel moves from a
"down" to an "up" configuration accompanied by cross sectional area changes, while its pockets
are occupied by lipids only in the absence of membrane stretch (39). Therefore, it seems that the
channel kind of "snaps open", unless it is pushed shut via lipids.
Hydrogen Deuterium eXchange Mass Spectrometry (HDX-MS) and 3 pulse Electron Spin Envelope
Echo Modulation (3pESEEM) spectroscopy are powerful tools in the investigation of membrane
protein structural dynamics. The former involves the exchange of protons on a protein with
deuterium generates accessibility information and has been used to provide key insights into the
role of lipids in determining the conformational state, but also secondary structure changes in
membrane transporters (40-44). The latter measures weak hyperfine coupling between unpaired
electrons and nuclear spins to probe accessibility to solvent (D_2O) and protons (residues, ions and
lipids) and has been used to investigate the dynamics of membrane proteins (2, 15, 45-51).
We hypothesized that there is a link between the physiologically relevant tension-activated state,
and the one stabilized by molecular triggers, and investigated whether there is structural analogy
between these two states. Here, we combine untargeted (HDX-MS) and single-residue (3pESEEM)
with molecular trigger activation (by modifying the pocket entrance with a bulky amino acid
substitution, L89W) on MscL and independently generate a tension-activated state by MD
simulations to interrogate whether there is a link between the two differently derived states. We find
these states are structurally analogous, suggesting an intimate relationship between tension- and
ligand- activation in MS channels, potentially linked to their functional multimodality.
110
111 **Results**
112
Channel gating by molecular activation monitored by HDX-MS
The substitution of a bulky tryptophan residue at position L89 in the Mycobacterium tuberculosis
MscL (TbMscL) has been shown to restrict lipid access to channel pockets and destabilised the
closed state, leading to an expanded (sub-conducting) MscL state(2). In order to give a mechanistic
insight into the transitions occurring between the two states, we used HDX-MS to measure relative
differences in deuterium uptake between the WT (closed) and L89W (expanded by molecular
activation) channels in DDM. We succeeded in obtaining 95% peptide coverage of the entire
resolved MscL structure where the gating occurs, i.e. residues 1-125, including the entire TM
domain and the largest part of the cytoplasmic helical bundle (83% overall coverage for the entire
construct including the C-terminal 6xHis-tag). Differences in uptake, detected at the peptide level,
between the two states allowed us to identify protein regions that became deprotected from
exchange in the sub-conducting L89W state (Fig 1). Three regions of MscL were identified and
showed significant changes between the two states. Peptides in the regions containing residues
37-53 (periplasmic loop) and, 58-69 (top of TM2), and 97-111 (bottom of TM2, cytoplasmic loop
and top of the cytoplasmic helical bundle) had a significantly higher uptake of deuterium in L89W
in comparison to WT TbMscL (Fig 1, S1 and Table S1). Residues 37-53 run from the top of TM1
and end after the first β-sheet in the loop connecting TM1 to TM2. Residues 58-69 correspond to
the middle of the periplasmic loop up until TM2, and residues 97-111 cover the bottom of TM2 to
the top of the C-terminal helical bundle (Fig 1 and S1). This gave us the first glimpses into which
MscL domains are mostly impacted by the L89W (pocket entrance) modification across MscL's
length.
134
Solvent accessibility mapping at single residue resolution by 3pESEEM
We first aimed to investigate the feasibility of 3pESEEM on MscL and identify potential dynamic
sites. To this end, we generated twenty-six single cysteine mutants, accounting for over 20% of the

total TbMscL resolved length, spanning all protein domains. In particular, L2 and F5 on the S1 helix,
N13, L17, V21, I23, V31 and F34 on TM1, L42, and F48 on the periplasmic loop, residues N70,
V71, L72, L73, A74, F79, F84, A85, Y87, F88, L89, R98, K99, and K100 on TM2, and E102 and
V112 on the cytoplasmic loop and helical bundle (CHB) respectively, were mutated in the Cys-free
WT TbMscL as the background (Fig 2). We subsequently expressed, purified and spin labelled
each one of these twenty-six mutants (MTSSL modification is denoted as R1 thereafter) and
performed 3pESEEM solvent accessibility measurements, based on the modulation depth of
deuterium in the time domain signal, which is proportional to its associated signal intensity in the
frequency domain. Unlike PELDOR which requires high spin labelling efficiency to obtain high
sensitivity data acquisition (2), 3pESEEM can be performed with no major losses in sensitivity even
if the samples of interest show lower spin labelling efficiencies for individual sites. This has allowed
us to obtain good quality spectra and quantify the solvent (D_2O) accessibility for all twenty-six sites.
Residues in the S1 helix showed low accessibility, which is consistent with previous reports
suggesting that this helix is buried within the bilayer (53). TM1 pore-lining residues presented low
to intermediate accessibility (Fig 2 and S2), as would be expected for a closed channel pore (2, 54,
55). I23, V31 and F34 on TM1 are intermediately solvent-exposed (or relatively buried) and
positioned on the same side of TM1 (Fig 2 and S2). This could be due either to the presence of
native lipids, DDM and/or hinderance by the presence of TM2 on the TM1 interface. L42 and V48
are intermediately to highly exposed, as they form part of the periplasmic loop connecting TM1 and
TM2. N70, V71, L72, L73, and A74 present the largest disparity across our whole data set. L72 is
the most buried residue we measured (~ 3% compared to the most solvent-exposed cytoplasmic-
facing labelled residue K100R1) while N70 (~90%) and V71 (~80%) are two of the most exposed
(~90%), comparable to D_2O (solvent) accessibilities measured for exposed cytoplasmic sites (Fig
2 and S2). The rest present intermediate accessibilities to these two extremes, as expected for
subsequent residues, which form a helical turn with different space orientation. Residues N70 and
V71 are solvent-exposed in the closed MscL state, while residues L72 and L73 are buried. The
latter pair of residues is known to become exposed during opening, due to an anti-clockwise TM2
rotation (2, 56). F79, F84 and A85 are relatively buried and lie at the interface of TM helices
between different MscL subunits. Y87 has been identified as a lipid binding site (57) and along with
F88 and L89 presented low to intermediate accessibilities and forms part of the pocket region (2,
15). Except for R98, which showed intermediate accessibility, K99 and K100, presented high
accessibilities, indicating these residues are solvent-exposed in the closed state. These three
residues form a positively charged cluster for attracting negatively charged lipids (57) and our data
suggest that the non-ionic DDM binds either differently or at a different site. Substitution of these
residues with the neutral Gln had no effect on MscL's conformation, suggesting that specific lipid
headgroup binding on this particular site does not influence MscL gating (2). E102 (cytoplasmic
loop) and V112 (cytoplasmic helical bundle) presented high solvent accessibility as it would be
expected for cytoplasmic sites.
176
Identification of channel gating regions with single residue resolution
HDX-MS resolution is usually limited to the peptide level and, therefore, is unable to resolve
changes that occur at different residues within the same peptide. Solvent accessibility can be
probed at the single residue level using 3pESEEM allowing us to examine the relative solvent
accessibility of individual residues. This generates higher resolution information pertaining to the
regions of structural change identified by HDX-MS, as well as to the channel pore vapour lock. In
each of these regions, we introduced eight distinct single cysteine substitutions in an L89W MscL
background. Specifically, N13 and V21 (pore), L42 in the periplasmic loop, N70, V71, L72, L73 (top
of TM2) and K100 (bottom of TM2 and cytoplasmic loop) (Fig 3). V21 forms the vapour lock (along
with L17) that controls the channel gate and N13 sits one turn of a helix below. No significant
change in this region was highlighted via HDX-MS, while due to the close proximity of spin labels
pointing towards the pore (Fig S3), distances are out of PELDOR's distance measurement range
(>20 A). However, 3pESEEM is highly sensitive to the local spin label environment of specific sites,
does not need a reference channel state as it is the case for HDX-MS, and is not subjected to
distance restrictions as PELDOR. Differences in the vapour-lock region may not have been

detected by HDX-MS as structural changes may have resulted in complex changes to hydrogen
bonding networks, whereby some residues were deprotected from exchange and others became
protected, resulting in only minor/no changes in deuterium uptake at the peptide level (Fig 1). To
resolve this, we labelled the crucial V21 site and N13, and six other sites within the regions identified
by HDX-MS, which undergo conformational changes when the L89W modification is present (Fig
3). N13R1's side chain points towards the cytoplasm and is expected to be solvent-exposed in the
closed state (Fig S2 and S3). Overall, the motivation behind this was to confirm the peptide-level
HDX-MS data and complement this with single residue resolution data for cases where crucial sites
have been undetected potentially due to limitations imposed by peptide-level HDX-MS analysis.
3pESEEM measurements at positions N13 and L42 showed no significant difference in solvent
accessibility between the WT and the L89W background. In L89W, the accessibility increased for
V21, which is consistent with a pore hydration, and an over ten-fold increase observed for L72. A
significant decrease was observed for K100 though, which sits at the C-terminal end of TM2 (Fig
3). A decrease in solvent accessibility was observed for N70 and V71 in presence of the
modification (molecular activation) with 2.4% and ~19% reduction, respectively. Despite being
consecutive residues, 70-73 show distinct changes in solvent accessibility in 3pESEEM, unlike
HDX-MS, which averages across the top of TM2 region and relies on single peptide coverage of
these residues. A significant and smaller increase in accessibility is seen for L72 and L73, and
opposite to the effect observed for N70 and V71, while the bottom of TM2 (K100) becomes less
accessible to solvent in the L89W background (Fig 3). This result is consistent with a rotation of the
top of TM2 which occurs during mechanical activation and agrees with PELDOR distance
measurements and x-ray structures (2, 15, 56). HDX-MS experiments showed deprotection on
peptides inclusive of these residues and is consistent with a dramatic increase in accessibility of
residue L72 for L89W in 3pESEEM measurements (Fig 3). V21R1 forms the vapour-lock for
TbMscL and shows an increase in solvent accessibility in an L89W background, consistent with
the pore gate hydration and the channel entering a sub-conducting state. HDX-MS measurements
show a deprotection of the TM2 bottom for L89W versus WT suggesting an overall solvent
exposure of this region (Fig 1 and 3). The MscL cytoplasmic loop has been shown to participate
in gating and provide access to streptomycin for entering the cell, while shortening of the loop
influences MscL's function by irreversibly decreasing its conductance (13, 58).
222
Membrane tension stretching promotes pore hydration in the WT channel
In order to mimic the naturally occurring activation of MscL and estimate how far the proposed
transitions determined in DDM micelles are comparable to simulated transitions in a lipid
membrane, we implemented MD simulations on the WT TbMscL, with tension and no tension
applied to the lipid bilayer. We performed the no tension simulations within the same lipid
composition in order to compare the pore properties and overall channel architecture between
these states. Our simulations were initiated from the TbMscL X-ray structure (PDB 2OAR)(55)
carrying no modifications. Previously, significantly higher bilayer tension was required to open MscL
in MD simulations over short periods of simulation time(29-31), in comparison to the lower tension
(~67.5 mN/m) and longer simulation times (300 ns) used in this study, allowing us to capture
intermediate states.
WT TbMscL was inserted in a POPC bilayer and simulations for 300 ns were performed in which
the x- and y- pressure of the box was set at -50 bars constant bilayer tension. During the
simulations, the bilayer could expand in order to adopt the membrane expansion occurring in the
xy plane. As a result the bilayer thickness decreased approximately 1.2 nm compared to the POPC
bilayer without tension (Fig 4). The channel underwent large conformational changes reaching an
overall RMSD of ~14 Å compared to the initial closed structure (Fig S4A) to adapt to the changes
in the bilayer by tilting its TM helices towards the xy plane (Fig 4A and Video S1). In particular, TM1
and TM2 gradually tilted by 30 and 15 degrees respectively towards the end of the simulations,
while the pressure-sensitive pocket surface and the area exposed to the bilayer decreased
dramatically (Fig 4, S4B and C).
Analysis of pore hydration profiles using CHAP (59) showed opening and hydration of the WT
channel pore under tension in all of our simulations (Fig 5A). Whilst the higher (compared to

biologically relevant) membrane tension does not allow us to provide information about the
timescales of the channel transition from the closed to the hydrated state, it enables us to obtain
structural data for an important intermediate state, which has a hydrated open pore, essential in
MscL's gating process. For these pore hydration events we monitored the annular lipids, which
make direct contact or reside in close proximity to the channel and simultaneously interrogated the
pore radius profile using HOLE(60) (Fig 5B,C and 6C). We first observed that following constant
tension application for ~200 ns, the pore becomes hydrated and the channel undergoes major
structural rearrangements (Fig 5A). Although a significant membrane thinning (~1.2 nm) occurs
during tension application, L89W's pore cannot open to allow water molecules to flow through and
the pore radius is still restricted to ~1 Å, even though the changes in global secondary structure
are similar to WT channel under tension (Fig 5 B,C,S4 and Table S2). Following bilayer stretching
the total pocket surface area decreases and limits access and availability to bilayer lipids
throughout our simulations (Fig 5B). Overall, the under tension MD data agree well with the HDX-
MS and 3pESEEM accessibility data obtained for L89W (molecular activation). In both cases,
similar channel regions become accessible to solvent suggesting a similar opening mechanism for
the obtained expanded and sub-conducting state.
262
Safety-pin acting trapped lipids in the pockets do to allow for pore hydration
A direct consequence of the lipid moves first model would be that if lipids are trapped within and
cannot become loosely associated with the pockets, under bilayer tension, the channel will not be
able to open. To test this hypothesis we performed MD simulations with L89W TbMscL under
tension and identical to WT conditions (POPC lipid bilayer, tension value and simulation time).
L89W under tension undergoes similar global structural changes to the WT. However, its pore does
not become hydrated, even under the tension conditions which previously promoted WT pore
hydration (Fig 5A, 7B, S4 and Table S2).
During the equilibration of our simulations lipids intercalated into the pockets and occupied them.
Following tension application, lipids were pulled towards the membrane, but the bulky tryptophan
modification at the entrance disrupted the lipid exit path, despite tension application sufficient to
open the WT channel was applied (Fig 5).
Interestingly, in the tension activated state, F79 and L89 from adjacent subunits come very close
to form two new smaller pockets in the inner- and outer- leaflets of the now substantially thinner
membrane (Fig 5C and D). This spatial "refinement" may facilitate the next stage in MscL's
activation, which follows this sub-conducting state, that requires highly localised pressure for full
opening of 3 nS and ~30 Å (2, 34, 35). Asymmetry in bilayer tension sensitivity has been observed
for other mammalian mechanosensitive channels (61). Further, we observed that inner-leaflet
lipids which could previously occupy MscL's pockets and keep the channel closed, now could
penetrate either of the two newly created sub-pockets (Fig 5B and 6 A,B).
This change restricts the access of new lipids from the bulk membrane to the pockets, and
immobilizes trapped lipids within the pockets that entered initially during equilibration. In such a
case these lipids should interact more strongly with residue 89 in the modified (W89) than the WT
(L89) channel. To test this hypothesis we calculated the pairwise energy forces between all lipids
and residue 89, L and W, for the WT and modified channels respectively (Table 1). We found that
the lipid interaction (Lennard-Jones) with residue 89 are twice as strong in the modified compared
to WT channel (Table 1). This finding suggests that lipids are "stuck" by the modification in the
pockets, reducing their ability to exchange with the bulk bilayer during tension application (Fig 3C
and D). Despite the substantial global channel rearrangements L89W undergoes, lipids remain
tightly associated with the pockets and the pore cannot hydrate (Fig 5).
293
Lipid order and contact accessibility changes during tension application
To assess the lipid contact profile with the MscL TM domain we calculated the relative difference
of lipid contacts between the under- and no tension TbMscL states (Fig 6 and S5). We found that
the relative lipid contacts within bilayer facing residues on the TM domain were reduced over the
course of the simulation. In contrast, the lipid contacts of hydrophobic residues that were not lipid
exposed in the closed state increased substantially (Fig 6 and S5). This is consistent with TM2

rotation occurring during MscL opening (2) and with pockets decreasing their surface contact area
and becoming less accessible to lipids during opening (Fig 4B). To understand lipid rearrangement
occurring upon tension application we calculated the lipid order before and after tension application.
We found that all lipids (particularly the lipid chain ends) orient substantially more horizontally and
that the pocket lipids specifically orient marginally more horizontal compared to the bulk bilayer
lipids in the presence of tension (Fig S6).
306
307 **Discussion**
308
According to the lipid moves first model (5), MS channels open when lipids reduce contacts with
the pockets. Equally, the model predicts that if lipids are trapped within the pockets, then the
channel will not be able to open. Following application of bilayer tension to WT and L89W channels,
we observe similar global structural rearrangements. Despite this, only the WT channel pore
increased in diameter enough for pore hydration. The pore of the L89W channel was significantly
smaller in volume and diameter. (Fig 2,3,5,6C and S4). Previously, when lipids were disrupted from
entering the pockets, due to the L89W mutation located at the pocket entrance, the pore opening
probability increased and the activation threshold significantly decreased (2). In the present set up,
the L89W mutation locks the exit door and traps the lipids in the pockets by applying twice as strong
steric forces which constrict the space for lipids to move out, compared to L89 in the WT channel
(Fig 6A, B and Table 1). Lipids are also present in the pockets in the "hydrated" WT, following
tension activation, but these are relatively loose and are not sufficient to close the channel (Fig 5B,
6B and Table 1). We find that the pockets become substantially smaller under tension and
consequently less accessible to lipids (Fig 4B, 5B and 6). In the modified channel, F79 and W89
form a "molecular" bridge resulting in a two-compartment repartitioning of the smaller pockets (Fig
6A and B). The WT (under tension) expanded state is substantially different from the closed state
x-ray structure used (PDB 2OAR) (Fig 4, S4 and Video S1). Importantly, TM2 movement
(translation and expansion due to anticlockwise TM2 rotation) under tension is similar to that
occurring upon molecular activation. When lipids are disrupted from entering the pockets, the state
generated is similar to the archaeal x-ray expanded MscL model (56), which is also consistent with
the expanded (and sub-conducting) L89W state, revealed by PELDOR (2) (Fig 7A). The L89W
(molecular) and WT (tension) activated states are similar (Fig 7A), suggesting that MscL's gating
(from closed to expanded) is independent of the trigger used.
Previously, cross-sectional expansions, not associated to a conducting pore have been proposed
for the initial stages of MscL activation (33, 62), suggesting MscL could act as a membrane tension
dampener and undergo through closed-closed pore conformations (63). Our accumulative
experimental (structural and biophysical) and computational data are in agreement with these
models. Our data support the notion that there are structural transitions that occur prior to channel
opening. Support is also provided for intimate lipid interactions with TM pockets that act as pressure
safety pins to keep the channel closed until a certain tension threshold is reached (2). This was
substantiated further when the channel was unable to gate after the lipids were trapped within these
pockets via the L89W modification (Fig 5). Despite undergoing large conformational changes
consistent with those seen for WT under tension, the trapped lipids prevent pore hydration and
stabilize a closed (non-hydrated pore) but expanded state (Fig 3, 5, 7).
Our data suggest that for MscL to reach a hydrated state, promoted either by tension or molecular
triggers, the lipids will have to loosen their contacts with the pockets (Fig 5 and Table 1). Lipids
could move under tension and adopt a more horizontal orientation in respect to the membrane
plane in the WT channel, in the absence of pocket modifications that disrupted lipid exchange with
the bulk bilayer (Fig S6). The addition of molecules which could compete for the pockets may
support or disrupt lipids acting as negative allosteric modulators within the pocket sites (Fig 7C).
Following similar modifications at MscL's pocket entrance, or when molecules that specifically bind
these pockets were introduced, they caused a reduction in MscL's activation threshold, prolonged
openings and increased open channel probability (2, 12, 64).
Disruption of lipid access into the TbMscL's pockets promoted a mechanical response in the
absence of applied tension (2), while subtle structural differences between these pockets in MscL

orthologues dramatically altered their function (15). This pocket region in TbMscL, which is
important for lipid binding (2, 15), also binds, specific to EcMscL molecules, which could modulate
its activity (12, 13). TbMscL L89 is located at the entrance of the pockets and is structurally aligned
to EcMscL M94 and in close proximity to I92, F93, I96 and K97, modifications of which, influence
MscL's activation by ultrasound pulses (65), decrease its tension threshold for use in a
multicompartmental lipid vesicle framework (66), and modulate its function (12) . Molecules specific
to MscL are expected to act through disruption of lipid-protein interactions (26) within these pockets.
These did not fully open the channel, but caused a structural rearrangement of TM2's cytoplasmic
side accompanied by stretching of the cytoplasmic loop (12, 13). This is in agreement with the
structural changes observed by HDX and ESEEM for the modified L89W TbMscL channel (or the
structurally equivalent M94 EcMscL) and the tension activated WT channel (Fig 1,3,4,7 and S4).
V21 exposure increase in the expanded (by molecular means) MscL state is consistent with a
channel pore hydration and opening. N13R1's side chain sits one and two turn(s) of a helix below
V17 and V21 on TM1 respectively. N13 is already exposed in the closed state, due to its side chain
being solvent accessible (Fig S2 and S3), thus unaffected in the expanded L89W modified state.
Transition to a sub-conducting state (L89W in DDM) increased solvent accessibility on the upper
TM1 and TM2 and the cytoplasmic loops on top of the helical bundle (Fig 1 and 2). Widening of
MscL's cytoplasmic interface causes a stretching of the loops, of which shortening or modification
has been shown to influence MscL's gating properties, pore conductance and oligomeric assembly
(13, 34, 58, 67, 68).
The top of TM1 (residues 36 - 41) partially unfolds within a single subunit of the MscL pentamer,
while the remaining four subunits bend without fully converting into a loop. This finding was
consistently observed across all three MD simulation repeats following the symmetric application
of tension along the membrane xy plane and suggests the presence of a subunit asymmetry during
MscL opening (33, 69). In contrast to the WT channel, all five TM1 helices bend without a break in
helical symmetry or conversion into a loop in the L89W channel (Fig 4). For the top of TM1 helix to
partially unfold, lipids should become loosely associated with the pockets before asymmetric
opening could occur and eventually lead to channel opening. A similar activation mechanism has
been suggested for the small conductance MS channel MscS where inner-leaflet lipids residing
within TM pockets are required to move before outer-leaflet lipids can disengage to allow gating
(25). Whether the lipids are sequentially released (or loosen their contact) following single subunit
activation remains to be elucidated. Our data are compatible with such an asymmetry in opening
and this structural effect may be the consequence of lipids moving out from MscL's five pockets
sequentially. Subsequently, following initial expansion and conversion into loops, these loops could
then provide membrane handles for MscL in order to transit to its full opening state (Video S1).
Despite a significant membrane thinning (~1.2 nm), L89W's pore cannot allow water molecules to
flow through (Fig 4). Membrane thickness and hydrophobic mismatch energetically contribute to
MscL's gating, but our data suggest they may not be the sole driving force and pocket lipids have
to be released first in order to destabilize MscL's closed state. However, we should note that while
tension application and channel gating occurs within the 300 ns that our simulations run, this may
be faster compared to the timescale of MscL opening transitions.
Molecular activation due to binding of molecules acts in domains which are important for lipid-
protein interactions and are highly sensitive to membrane tension. Lipids must adopt certain angles
in respect to the membrane plane in order to enter (or exit) the pockets and these angles should
be consistent with lipids oriented horizontally, contrary to the bulk bilayer lipids. Such lipids have
been resolved in multiple x-ray and cryoEM structures of MscS (5, 25), TRAAK (23), TREK-2 (24)
and TRPV3 (22) channels. When we calculated the lipid order parameters for our MD simulations,
we found that under tension application the lipids adopt a substantially more horizontal orientation
(Fig S6). This is consistent with significantly lower pulling forces required to remove single lipids
from TbMscL's pockets when they are applied at 45° in respect to the membrane plane (70). Such
lipid rearrangement may be part of the closed-closed expansion gating process, since the overall
RMSD between the L89W and WT structures following tension application was small, while both
differ significant from the closed state (PDB 2OAR) (Fig 7B and S4). These findings are consistent
with the pocket lipids acting as negative allosteric modulators . Shifting of the equilibrium occurs in

order to destabilize the closed state and initiate symmetric (simultaneous) or asymmetric
(sequential) pocket lipid removal from the five MscL subunits (2, 69).
Accumulatively, our structural, biophysical and computational data demonstrate that MscL state
derived by modification of pressure sensitive TM regions is analogous to the tension activated MscL
state (mechanical), summarized in our proposed model (Fig 7C). We generated tension- and
modification- activated MscL states and interrogated their pore hydration properties, lipid-protein
interactions and monitored their structural dynamics by experimental (HDX-MS and 3pESEEM
solvent accessibility measurements) and computational (MD simulations) tools. The structural
similarities of these two derived states suggests that molecular triggers could act on specific sites
of MscL to mimic naturally occurring tension activation, with implications for the multimodality and
ligand activation of MS channels.
419
420 **Materials and Methods**
421
422 For *Materials see* the *SI Appendix*
423
424 Mutagenesis and Protein Expression
425 For detailed protocols *see* the *SI Appendix*
426
427 Protein Purification and Spin Labelling
428 The protocols followed in this study were similar to the ones previously described (2, 15, 54). For
429 details and also HDX-MS sample preparation *see* the *Appendix*.
430
3pESEEM spectroscopy
X-band three pulse ESEEM measurements were performed on a Bruker ELEXSYS E580
spectrometer with a 4 mm dielectric resonator (MD4). Measurements were conducted at 80 K. The
3-pulse sequence used for the experiments is $\pi/2$-τ-$\pi/2$-T-$\pi/2$-stimulated echo with a pulse length
$t_{\pi/2}$ = 16 ns and inter-pulse delay τ that was adjusted to match either the blind spots of the proton
or deuterium. The delay T was incremented from 400ns in 12 ns steps. A four-step phase cycling
was used to eliminate the unwanted echoes. All the measurements were performed at the
maximum of the filed sweep spectrum of the nitroxide. For solvent accessibility determination, only
-3pESEEM data recorded with τ that corresponds to the proton blind spot were used. The obtained
time-domain traces were background-corrected, apodized with a hamming window and zero-filled
prior to Fourier transformation. The solvent accessibility can be determined either from the
modulation depth in time domain or the corresponding amplitude of the deuterium peak in
frequency domain. In the present work, the modulation depth for each mutant was determined by
fitting the corresponding time-domain trace by a damped harmonic oscillation function and the
outcome from the fitting was used to estimate the solvent accessibility (47).
446
HDX and LC-MS/MS
HDX-MS experiments were carried out using an automated HDX robot (LEAP Technologies, Fort
Lauderdale, FL, USA) coupled to an M-Class Acquity LC and HDX manager (Waters Ltd.,
Wilmslow, Manchester, UK). All samples were diluted to 16 µM in equilibration buffer (50 mM
potassium phosphate, 300 mM NaCl, pH 7.4, 0.05% DDM) prior to analysis. 5 µL sample was
added to 95 µl deuterated buffer (50 mM potassium phosphate, 300 mM NaCl pD 7.4, 0.05% DDM)
and incubated at 4 °C for 0.5, 1, 2, 10 or 60 min. Following the labelling reaction, samples were
quenched by adding 50 µl of the labelled solution to 100 µl quench buffer (50 mM potassium
phosphate, 300 mM NaCl, 0.1% DDM pH 2.2) giving a final quench pH ~2.5. 50 µl of quenched
sample was injected on to the HDX Manger and passed through immobilized pepsin and
aspergillopepsin columns connected in series (AffiPro, Czech Republic) at 115 µl min-1 (20 °C)
and loaded on to a VanGuard Pre-column Acquity UPLC BEH C18 (1.7 µm, 2.1 mm X 5 mm,
Waters Ltd., Wilmslow, Manchester, UK) for 3 mins in 0.3 % formic acid in water. The resulting
peptides were transferred to a C18 column (75 µm X 150 mm, Waters Ltd., Wilmslow, Manchester,

UK) and separated by gradient elution of 0-40 % MeCN (0.1 % v/v formic acid) in H2O (0.3 % v/v
formic acid) over 7 mins at 40 µl min-1. Trapping and gradient elution of peptides was performed
at 0 oC to reduce back exchange. The HDX system was interfaced to a Synapt G2Si mass
spectrometer (Waters Ltd., Wilmslow, Manchester, UK). HDMSE and dynamic range extension
modes (Data Independent Analysis (DIA) coupled with IMS separation) were used to separate
peptides prior to CID fragmentation in the transfer cell (71). HDX data were analyzed using PLGS
(v3.0.2) and DynamX (v3.0.0) software supplied with the mass spectrometer. Restrictions for
identified peptides in DynamX were as follows: minimum intensity: 1000, minimum products per
MS/MS spectrum: 5, minimum products per amino acid: 0.3, maximum sequence length: 25,
maximum ppm error: 5, file threshold: 3/3. Following manual curation of the data, PAVED(72) and
Deuteros (52) were used to identify peptides with statistically significant increases/decreases in
deuterium uptake (applying 99% or 95% confidence intervals) and to prepare Wood's plots. The
raw HDX-MS data, have been deposited to the ProteomeXchange Consortium via the PRIDE
partner repository with the dataset identifier PXD021983. A summary of the HDX-MS data, as
recommended by reporting guidelines, is shown in Table S1.

Atomistic Molecular Dynamics Simulations
MD simulations under no tension: The set up and parameters of MD simulations without tension
were as previously described (2). For more details *see* the *SI Appendix*.
MD simulations under tension: These MD simulations were also set up using CHARMM-GUI (73).
The MscL structure (PDB 2OAR) in the wild type and in a mutated form (L89W) was inserted in a
symmetric bilayer containing 514 POPC lipids. Systems were neutralized with a 150 mM
concentration of NaCl. The obtained system was energy minimised for 5000 steps and
subsequently equilibrated in 6 steps following CHARMM-GUIs equilibration. Stretch-induced
conformational changes in the MscL model were investigated by unrestrained simulations of 300
ns (3 simulation repeats for each system) were the bilayer plane (xy plane) pressure was changed
semiisotropically to -50 bars (which corresponds to a tension of ~67.5 mN/m). The pressure in the
bilayer normal (z direction) was kept at +1 bar. All the atomistic systems were simulated using
GROMACS 2016 (74) with CHARMM36 force field (75) and a 2 fs time step. The Nose-Hoover
thermostat (76) and the Parrinello-Rahman barostat(77) were used for the stretch-induced
simulations. All simulations were performed at 310 K. Long-range electrostatics were managed
using the particle-mesh Ewald method(78) and the LINCS algorithm was used to constrain bond
lengths(79).

Data accessibility

HDX mass spectrometry:
Data are available via ProteomeXchange with identifier PXD021983.
Project Name: Mechanical and molecular activation lead to structurally analogous MscL states
Project accession: PXD021983
Account details: Username: reviewer_pxd021983@ebi.ac.uk, Password: lnAkk2H3

3pESEEM and MD:
Data are available within the following link: http://archive.researchdata.leeds.ac.uk/777/

Acknowledgments

This project was supported by a Biotechnology and Biological Sciences Research Council
(BBSRC) grant (BB/S018069/1) to C.P., who also acknowledges support from the Wellcome Trust
(WT) (219999/Z/19/Z) and the Chinese Scholarship Council (CSC) in the form of studentships for
B.J.L. and B.W. respectively. ANC is a Sir Henry Dale Fellow jointly funded by the WT and the
Royal Society (220628/Z/20/Z). Funding from the BBSRC (BB/M012573/1) enabled the purchase
of mass spectrometry equipment.

References

515
516
517
1. Laganowsky A, et al. (2014) Membrane proteins bind lipids selectively to modulate their
structure and function. Nature 510(7503):172-175.
2. Kapsalis C, et al. (2019) Allosteric activation of an ion channel triggered by modification of
mechanosensitive nano-pockets. Nat Commun 10(1):4619.
3. Pliotas C & Naismith JH (2016) Spectator no more, the role of the membrane in regulating ion
channel function. Curr Opin Struct Biol 45:59-66.
4. Patrick JW, et al. (2018) Allostery revealed within lipid binding events to membrane proteins.
Proc Natl Acad Sci U S A 115(12):2976-2981.
5. Pliotas C, et al. (2015) The role of lipids in mechanosensation. Nat Struct Mol Biol 22(12):991-
998.
6. Teng J, Loukin S, Anishkin A, & Kung C (2015) The force-from-lipid (FFL) principle of
mechanosensitivity, at large and in elements. Pflugers Arch 467(1):27-37.
7. Malcolm HR, Blount P, & Maurer JA (2015) The mechanosensitive channel of small
conductance (MscS) functions as a Jack-in-the box. Biochim Biophys Acta 1848(1 Pt A):159-166.
8. Cox CD, Nakayama Y, Nomura T, & Martinac B (2015) The evolutionary 'tinkering' of MscS-like
channels: generation of structural and functional diversity. Pflugers Arch 467(1):3-13.
9. Brohawn SG (2015) How ion channels sense mechanical force: insights from mechanosensitive
K2P channels TRAAK, TREK1, and TREK2. Annals of the New York Academy of Sciences
1352:20-32.
10.Naismith JH & Booth IR (2012) Bacterial mechanosensitive channels--MscS: evolution's
solution to creating sensitivity in function. Annual review of biophysics 41:157-177.
11.Wray R, Wang J, Iscla I, & Blount P (2020) Novel MscL agonists that allow multiple antibiotics
cytoplasmic access activate the channel through a common binding site. PLoS One
15(1):e0228153.
12.Wray R, Iscla I, Kovacs Z, Wang J, & Blount P (2019) Novel compounds that specifically bind
and modulate MscL: insights into channel gating mechanisms. FASEB journal : official publication
of the Federation of American Societies for Experimental Biology 33(3):3180-3189.
13.Wray R, et al. (2016) Dihydrostreptomycin Directly Binds to, Modulates, and Passes through
the MscL Channel Pore. PLoS biology 14(6):e1002473.
14.Rasmussen T, et al. (2019) Interaction of the Mechanosensitive Channel, MscS, with the
Membrane Bilayer through Lipid Intercalation into Grooves and Pockets. J Mol Biol 431(17):3339-
3352.
15.Kapsalis C, Ma Y, Bode BE, & Pliotas C (2020) In-Lipid Structure of Pressure-Sensitive Domains
Hints Mechanosensitive Channel Functional Diversity. Biophys J 119(2):448-459.
16.Syeda R, et al. (2015) Chemical activation of the mechanotransduction channel Piezo1. Elife 4.
17.Botello-Smith WM, et al. (2019) A mechanism for the activation of the mechanosensitive Piezo1
channel by the small molecule Yoda1. Nat Commun 10(1):4503.
18.Bagriantsev SN, Peyronnet R, Clark KA, Honore E, & Minor DL, Jr. (2011) Multiple modalities
converge on a common gate to control K2P channel function. EMBO J 30(17):3594-3606.
19.Li B, Rietmeijer RA, & Brohawn SG (2020) Structural basis for pH gating of the two-pore domain
K(+) channel TASK2. Nature 586(7829):457-462.
20.Piechotta PL, et al. (2011) The pore structure and gating mechanism of K2P channels. EMBO
J 30(17):3607-3619.
21.Feliciangeli S, Chatelain FC, Bichet D, & Lesage F (2015) The family of K2P channels: salient
structural and functional properties. J Physiol 593(Pt 12):2587-2603.
22.Deng Z, et al. (2020) Gating of human TRPV3 in a lipid bilayer. Nat Struct Mol Biol.
23.Brohawn SG, Campbell EB, & MacKinnon R (2014) Physical mechanism for gating and
mechanosensitivity of the human TRAAK K+ channel. Nature 516(7529):126-130.
24.Dong YY, et al. (2015) K2P channel gating mechanisms revealed by structures of TREK-2 and
a complex with Prozac. Science 347(6227):1256-1259.

25.Rasmussen T, Flegler VJ, Rasmussen A, & Bottcher B (2019) Structure of the
Mechanosensitive Channel MscS Embedded in the Membrane Bilayer. J Mol Biol.
26.Bavi N, et al. (2016) The role of MscL amphipathic N terminus indicates a blueprint for bilayer-
mediated gating of mechanosensitive channels. Nature Communications 7.
27.Erdogmus S, et al. (2019) Helix 8 is the essential structural motif of mechanosensitive GPCRs.
Nat Commun 10(1):5784.
28.Debruycker V, et al. (2020) An embedded lipid in the multidrug transporter LmrP suggests a
mechanism for polyspecificity. Nat Struct Mol Biol 27(9):829-835.
29.Martinac AD, Bavi N, Bavi O, & Martinac B (2017) Pulling MscL open via N-terminal and TM1
helices: A computational study towards engineering an MscL nanovalve. PLoS One
12(8):e0183822.
30.Gullingsrud J & Schulten K (2003) Gating of MscL studied by steered molecular dynamics.
Biophys J 85(4):2087-2099.
31.Bavi N, et al. (2017) Nanomechanical properties of MscL alpha helices: A steered molecular
dynamics study. Channels (Austin) 11(3):209-223.
32.Jeon J & Voth GA (2008) Gating of the mechanosensitive channel protein MscL: the interplay
of membrane and protein. Biophys J 94(9):3497-3511.
33.Konijnenberg A, et al. (2014) Global structural changes of an ion channel during its gating are
followed by ion mobility mass spectrometry. Proc Natl Acad Sci U S A 111(48):17170-17175.
34.Herrera N, Maksaev G, Haswell ES, & Rees DC (2018) Elucidating a role for the cytoplasmic
domain in the Mycobacterium tuberculosis mechanosensitive channel of large conductance.
Scientific reports 8(1):14566.
35.Mukherjee N, et al. (2014) The activation mode of the mechanosensitive ion channel, MscL, by
lysophosphatidylcholine differs from tension-induced gating. FASEB journal : official publication of
the Federation of American Societies for Experimental Biology 28(10):4292-4302.
36.Katsuta H, Sawada Y, & Sokabe M (2019) Biophysical Mechanisms of Membrane-Thickness-
Dependent MscL Gating: An All-Atom Molecular Dynamics Study. Langmuir 35(23):7432-7442.
37.Sawada Y & Sokabe M (2015) Molecular dynamics study on protein-water interplay in the
mechanogating of the bacterial mechanosensitive channel MscL. Eur Biophys J 44(7):531-543.
38.Sawada Y, Murase M, & Sokabe M (2012) The gating mechanism of the bacterial
mechanosensitive channel MscL revealed by molecular dynamics simulations: from tension
sensing to channel opening. Channels (Austin) 6(4):317-331.
39.Aryal P, et al. (2017) Bilayer-Mediated Structural Transitions Control Mechanosensitivity of the
TREK-2 K2P Channel. Structure 25(5):708-718.e702.
40.Martens C, Shekhar M, Lau AM, Tajkhorshid E, & Politis A (2019) Integrating hydrogen-
deuterium exchange mass spectrometry with molecular dynamics simulations to probe lipid-
modulated conformational changes in membrane proteins. Nat Protoc 14(11):3183-3204.
41.Ahdash Z, et al. (2019) HDX-MS reveals nucleotide-dependent, anti-correlated opening and
closure of SecA and SecY channels of the bacterial translocon. Elife 8.
42.Konermann L, Pan J, & Liu YH (2011) Hydrogen exchange mass spectrometry for studying
protein structure and dynamics. Chem Soc Rev 40(3):1224-1234.
43.Moller IR, et al. (2019) Conformational dynamics of the human serotonin transporter during
substrate and drug binding. Nat Commun 10(1):1687.
44.Martens C, et al. (2018) Direct protein-lipid interactions shape the conformational landscape of
secondary transporters. Nat Commun 9(1):4151.
45.Cieslak JA, Focia PJ, & Gross A (2010) Electron spin-echo envelope modulation (ESEEM)
reveals water and phosphate interactions with the KcsA potassium channel. Biochemistry
49(7):1486-1494.
46.Pliotas C (2017) Ion Channel Conformation and Oligomerization Assessment by Site-Directed
Spin Labeling and Pulsed-EPR. Methods in enzymology 594:203-242.
47.Volkov A, Dockter C, Bund T, Paulsen H, & Jeschke G (2009) Pulsed EPR determination of
water accessibility to spin-labeled amino acid residues in LHCIIb. Biophys J 96(3):1124-1141.
48.Liu L, et al. (2016) Probing the Local Secondary Structure of Human Vimentin with Electron
Spin Echo Envelope Modulation (ESEEM) Spectroscopy. J Phys Chem B 120(48):12321-12326.

49. Zhang R, et al. (2015) Development of electron spin echo envelope modulation spectroscopy
to probe the secondary structure of recombinant membrane proteins in a lipid bilayer. Protein Sci
24(11):1707-1713.
50. Matalon E, et al. (2013) Topology of the trans-membrane peptide WALP23 in model membranes
under negative mismatch conditions. J Phys Chem B 117(8):2280-2293.
51. Hartley AM, Ma Y, Lane BJ, Wang B, & Pliotas C (2021) Using pulsed EPR in the structural
analysis of integral membrane proteins. Electron Paramag Reson 27:74 -108.
52. Lau AMC, Ahdash Z, Martens C, & Politis A (2019) Deuteros: software for rapid analysis and
visualization of data from differential hydrogen deuterium exchange-mass spectrometry.
Bioinformatics 35(17):3171-3173.
53. Bavi N, et al. (2016) The role of MscL amphipathic N terminus indicates a blueprint for bilayer-
mediated gating of mechanosensitive channels. Nat Commun 7:11984.
54. Michou M, Kapsalis C, Pliotas C, & Skretas G (2019) Optimization of Recombinant Membrane
Protein Production in the Engineered Escherichia coli Strains SuptoxD and SuptoxR. ACS Synth
Biol 8(7):1631-1641.
55. Chang G, Spencer RH, Lee AT, Barclay MT, & Rees DC (1998) Structure of the MscL homolog
from Mycobacterium tuberculosis: a gated mechanosensitive ion channel. Science
282(5397):2220-2226.
56. Li J, et al. (2015) Mechanical coupling of the multiple structural elements of the large-
conductance mechanosensitive channel during expansion. Proc Natl Acad Sci U S A
112(34):10726-10731.
57. Powl AM, East JM, & Lee AG (2005) Heterogeneity in the binding of lipid molecules to the
surface of a membrane protein: Hot spots for anionic lipids on the mechanosensitive channel of
large conductance MscL and effects on conformation. Biochemistry 44(15):5873-5883.
58. Yang LM, et al. (2012) Three routes to modulate the pore size of the MscL channel/nanovalve.
ACS Nano 6(2):1134-1141.
59. Klesse G, Rao S, Sansom MSP, & Tucker SJ (2019) CHAP: A Versatile Tool for the Structural
and Functional Annotation of Ion Channel Pores. J Mol Biol 431(17):3353-3365.
60. Smart OS, Goodfellow JM, & Wallace BA (1993) The pore dimensions of gramicidin A. Biophys
J 65(6):2455-2460.
61. Clausen MV, Jarerattanachat V, Carpenter EP, Sansom MSP, & Tucker SJ (2017) Asymmetric
mechanosensitivity in a eukaryotic ion channel. Proc Natl Acad Sci U S A.
62. Betanzos M, Chiang CS, Guy HR, & Sukharev S (2002) A large iris-like expansion of a
mechanosensitive channel protein induced by membrane tension. Nature structural biology
9(9):704-710.
63. Boucher PA, Morris CE, & Joos B (2009) Mechanosensitive closed-closed transitions in large
membrane proteins: osmoprotection and tension damping. Biophys J 97(10):2761-2770.
64. Iscla I, Wray R, Wei S, Posner B, & Blount P (2014) Streptomycin potency is dependent on
MscL channel expression. Nat Commun 5:4891.
65. Ye J, et al. (2018) Ultrasonic Control of Neural Activity through Activation of the
Mechanosensitive Channel MscL. Nano Lett 18(7):4148-4155.
66. Hindley JW, et al. (2019) Building a synthetic mechanosensitive signaling pathway in
compartmentalized artificial cells. Proc Natl Acad Sci U S A 116(34):16711-16716.
67. Reading E, et al. (2015) The Effect of Detergent, Temperature, and Lipid on the Oligomeric
State of MscL Constructs: Insights from Mass Spectrometry. Chem Biol 22(5):593-603.
68. Liu Z, Gandhi CS, & Rees DC (2009) Structure of a tetrameric MscL in an expanded
intermediate state. Nature 461(7260):120-124.
69. Birkner JP, Poolman B, & Kocer A (2012) Hydrophobic gating of mechanosensitive channel of
large conductance evidenced by single-subunit resolution. Proc Natl Acad Sci U S A
109(32):12944-12949.
70. Vanegas JM & Arroyo M (2014) Force transduction and lipid binding in MscL: a continuum-
molecular approach. PLoS One 9(12):e113947.

71.Cryar A, Groves K, & Quaglia M (2017) Online Hydrogen-Deuterium Exchange Traveling Wave
Ion Mobility Mass Spectrometry (HDX-IM-MS): a Systematic Evaluation. J Am Soc Mass Spectrom
28(6):1192-1202.
72.Cornwell O, Radford SE, Ashcroft AE, & Ault JR (2018) Comparing Hydrogen Deuterium
Exchange and Fast Photochemical Oxidation of Proteins: a Structural Characterisation of Wild-
Type and DeltaN6 beta2-Microglobulin. J Am Soc Mass Spectrom 29(12):2413-2426.
73.Jo S, Kim T, Iyer VG, & Im W (2008) CHARMM-GUI: a web-based graphical user interface for
CHARMM. Journal of computational chemistry 29(11):1859-1865.
74.Abraham MJ, T. Murtola, R. Schulz, S. Páll, J.C. Smith, B. Hess, E. Lindah, and E. Lindahl
(2015) GROMACS: High performance molecular simulations through multi-level parallelism from
laptops to supercomputers. SoftwareX 1-2:19-25.
75.Lee S, et al. (2014) CHARMM36 united atom chain model for lipids and surfactants. J Phys
Chem B 118(2):547-556.
76.Evans DJ, Holian B.L. (1985) The Nose-Hoover thermostat. J. Chem. Phys. 83:4069–4074.
77.Parrinello M, Rahman A. (1981) Polymorphic transitions in single crystals: A new molecular
dynamics method. J. Appl. Phys. 52:7182–7190.
78.Darden T, York, D., Pedersen L. (1993) Particle mesh Ewald: An N·log(N) method for Ewald
sums in large systems. J. Chem. Phys. 98:10089–10092.
79.Hess B, H. Bekker, H.J.C. Berendsen, J.G.E.M. Fraaije (1997) INCS: A linear constraint solver
for molecular simulations. J. Comput. Chem. 18:1463–1472.
694
695

696
697 **Figures and Tables**

706
707
708
Figure 1. Observed changes in solvent accessibility and secondary structure by HDX-MS following
L89W pocket-entrance modification. A. Differences in the deuterium uptake of regions of TbMscL
(PDB 2OAR) when comparing the WT and the L89W modified protein. Regions highlighted in red
are deprotected following the L89W modification. Regions of the protein in grey show no significant
difference between the two conditions. L89W modification site is depicted as a cyan sphere. B.
Wood's plots showing the summed differences in deuterium uptake in MscL over all five HDX
timepoints, comparing WT with L89W MscL (Wood's plots were generated using Deuteros(52).
Peptides coloured in red, are deprotected from exchange in L89W MscL. No peptides were
significantly protected from exchange in L89W MscL compared with wild type MscL. Peptides with
no significant difference between conditions, determined using a 99% confidence interval (dotted
line), are shown in grey. Example deuterium uptake curves are shown in Fig S1.
720

Figure 2. Single residue solvent accessibility mapping of MscL by 3pESEEM. Spin-labelled residues are shown as red and pale yellow spheres for the most and the least solvent accessible residues respectively. Spin-labelled residues are coloured based on the quartile of their relative accessibility which has been normalised on scale of 0% and 100%, where the residues with the highest accessibility corresponds to 100%.

730
731
732

Figure 3. Molecular activation of MscL investigated by integrated 3pESEEM and HDX-MS solvent
accessibility measurements. A Time-domain 3pESEEM fitted experimental spectra (orange) of the
single mutants N13R1, V21R1, L42R1, N70R1, L72R1, L73R1, and K100R1 in DDM overlaid with
the L89W double mutant fitted spectra (purple) (for fitting see Methods). B Differences in solvent
accessibility for TbMscL (PDB 2OAR) following L89W modification. Spin labelled mutation sites
used for 3pESEEM accessibility measurements are represented by spheres, and peptides that
demonstrate a change in accessibility in HDX-MS following the L89W modification are represented
as highlighted helices. Red regions or spheres highlight areas that are deprotected, while blue
spheres and regions show areas that are protected following the L89W modification C Column bar
charts represent the 2H accessibility derived from the fitting of 3pESEEM time-domain traces.
Errors are calculated conservatively at 5% to compensate for the fitting error and differences in
relaxation times of different spin labelled residues. There was no significant difference in solvent
accessibility of N13R1, L42R1, and N70R1 compared to their 89W double mutant counterpart.
Solvent accessibility increased for V21R1, L72R1, and decreased for L71R1 and K100R1 following
the L89W modification.
749
750

A

B

Closed
WT - no tension

Open
WT - plus tension

Periplasm

Membrane

Cytoplasm

3.8 nm

2.6 nm

Pockets

TM1

TM2

30°

15°

774
Figure 5. Pore hydration under applied bilayer tension. WT under no tension (left column), WT
under bilayer tension (right column), L89W under bilayer tension (right column). A. Solvent density
profile of the MD simulations using the CHAP(59) software with the vapour-lock position of MscL
marked with blue arrows. B. Membrane becomes thinner when tension is applied but only the WT
TbMscL pore is hydrated (blue spheres) in contrast to L89W channel. Lipid (olive sticks) availability
is larger in the closed state, and lipids have easier access within the pockets to provide force on
the back of the vapour lock, keeping the MscL pore closed. This is in contrast to WT MscL under
tension (middle column) where the pockets become smaller, while lipid availability decreases and
the pore becomes hydrated. However, when lipids are trapped in the pockets of the L89W modified
channel, the pore does not become hydrated despite structural rearrangements occurring (right
column). C. Surface visualization of the pore pathway using the program HOLE(60). Red colour
indicates a pore radius smaller than 1.15 Å (water molecules cannot go through such an opening),
blue represents a radius larger than 2.3 Å and green between 1.15 to 2.3 Å.
788
789

A

Closed
WT - no tension

Open
WT - plus tension

Closed
L89W - plus tension

C

B

POPC

D

Periplasm

Membrane

Cytoplasm

Peri Loop

TM1

TM2

S1

CHB

Increase

No Change

Decrease

797
798
799
800
Figure 6. The effect of membrane tension to MscL lipid contact interactions. (A) and (B). F79 and
L89W form a "molecular bridge" in the L89W under tension state and consequently trap the lipids
within the pockets (right column). F79 and L89 come into greater proximity under bilayer tension in
the WT channel (expanded MscL, middle column) compared to the closed state. However, they do
not prevent the lipids from exchanging with the bulk bilayer during tension application. The L89W
mutation locks lipids in the pockets, preventing the channel from transiting to a hydrated state. C.
MscL pore radius profiles analysis and comparisons using HOLE(60). Although L89W MscL
undergoes similar major rearrangements to WT MscL under tension its pore remains significantly
smaller and almost identical to WT closed MscL under no tension. D. Relative changes in the
number of lipid contacts following stimulated tension in the membrane during MD simulations. The
blue regions show decrease in lipid contacts, while the red regions show an increase in lipid
contacts. From left to right: Surface view, single subunit cartoon representation and internal MscL
surface view. Internal channel regions become more exposed to lipids, while initially membrane
exposed regions (closed state) become less exposed to lipids, suggesting a MscL TM helical
rotation upon tension application and channel expansion/opening.
816
817

Figure 7. Structural comparison between tension and molecular MscL activation and proposed
model. TM1 and TM2 tilt and rotate to initiate MscL opening. Comparison between expanded
MaMscL (orchid, PDB ID: 4Y7J) and the open state WT TbMscL generated by MD (cyan, under
tension) (A) and the open WT TbMscL state generated by MD (cyan, under tesion) and L89W
TbMscL (green, under tension) (B). C. Proposed gating model for molecular and tension activated
modes. Closed (vertical TM helices, large pockets, tightly associated annular lipids and non-
hydrated pore), Expanded (tilted TM helices, smaller pockets, loosely associated annular lipids and
hydrated pore). Extra membrane tension is required to fully open MscL.
834
835

836
837
838
839 **Table 1.** Pairwise force energy calculations between TbMscL 89 (L or W) and lipids in the WT (L)
840 and pocket-entrance modified (W) channels and under tension

89 - POPC	Coulomb (KJ/mol)	SD - Coulomb (KJ/mol)	Lennard Jones (KJ/mol)	SD - Lennard Jones (KJ/mol)	Total Energy (KJ/mol)	SD – Total Energy (KJ/mol)
WT	-14.50	1.76	-101.39	27.43	-115.89	29.19
L89W	-14.77	3.90	-192.48	3.39	-207.25	7.29