Journal Name: Ecology

Manuscript type:

Manuscript title: Disentangling the roles of inter- and intraspecific variation on leaf trait

distributions across the eastern United States

Sergio Marconi[1,2], Benjamin G. Weinstein[2], Jeremy W. Lichstein[3], Stephanie A. Bohlman[1],

Aditya Singh[4], Ethan P. White[2]

[1]School of Forest, Fisheries, & Geomatics, University of Florida, Gainesville, Florida, USA

[2]Dept. of Wildlife Ecology & Conservation, University of Florida, Gainesville, Florida, USA

[3]Dept. of Biology, University of Florida, Gainesville, Florida, USA

[4]Dept Agricultural & Biological Engineering, University of Florida, Gainesville, FL 32611, USA

Keywords: leaf traits, intra-species variability, joint distribution, cross-scale, NEON, FIA.

Corresponding Author: Sergio Marconi **Email**: s.marconi@ufl.edu

Open Research statement: All data and data products are archived and accessible on

Zenodo (https://doi.org/10.5281/zenodo.4651366). Code to replicate the analyses and retrieve

data is stored on GitHub (https://github.com/MarconiS/Disentangling-the-role-of-phylogeny-

and-climate-on-joint-leaf-trait-distributions-across-Eastern-Uni/tree/0.1) and Zenodo

(https://zenodo.org/badge/latestdoi/353383665).

Abstract

Functional traits are influenced by phylogenetic constraints and environmental conditions, but previous large-scale studies modeled traits either as species weighted averages or directly from the environment, precluding analyses of the relative contributions of inter- and intraspecific variation across regions. We developed a joint model integrating phylogenetic and environmental information to understand and predict the distribution of eight leaf traits across the eastern USA. This model explained 68% of trait variation, outperforming both species-only and environment-only models, with variance attributable to species alone (23%), the environment alone (13%), and their combined effects (25%). The importance of the two drivers varied by trait. Predictions for the eastern USA produced accurate estimates of intraspecific variation and deviated from both species-only and environment-only models. Predictions revealed that intraspecific variation holds information across scales, affects relationships in the leaf economic spectrum and is key for interpreting trait distributions and ecosystem processes within and across ecoregions.

Introduction

Global change is expected to cause extensive changes in ecosystems, driving unprecedented redistribution of species and altering their traits (Diaz and Cabido, 2001). Understanding how these changes in the environment will drive adaptations within and across species is essential for management and conservation. Functional traits are among the most important properties of biological organisms because they are involved in key ecosystem processes from local community assembly (McGill et al., 2006, Sterck et al., 2011) to global biogeochemical cycles (Bonan et al., 2002; Verheijen et al., 2013), and these processes are interconnected across scales (Reichstein et al., 2014, Peaucelle et al., 2019). Biotic interactions, micro-climate, and soil conditions can affect species co-occurrence and influence local trait distributions (Bruelheide et

al., 2018, Simpson et al., 2016), and variation in climate within species ranges can affect realized

niches and drive trait responses (Chave, 2013). Relationships between traits, like the the leaf

economic spectrum (LES) in plants (Wright et al., 2004), reveal information about biological

constraints in resource allocation than impact ecophysiology and are a key component of

ecosystem models (Fisher et al, 2015). Given their central role across levels of organization,

understanding how traits vary within and among species across scales and environments is

essential for conserving ecosystem function (Violle et al., 2014).

Traits vary geographically through a combination of shifts in species abundances and

intraspecific trait variation (Leps et al., 2011, Laughlin et al., 2012, Münzbergová et al., 2017).

Thus, understanding how traits respond to the environment across wide geographic areas

requires approaches that integrate both inter- and intraspecific variation. This is challenging

because individual level trait data are geographically and taxonomically limited, making it hard

for traditional methods to identify the relative importance of inter- and intraspecific variation at

large scales (Henn et al., 2018). In addition, due to limited data availability, studying traits at

broad geographic scales may require predicting trait values for species that have not been

sampled throughout their geographic range, or even species that have not been sampled at all.

Given these challenges, most current approaches to studying broad-scale trait variation focus on

predicting community weighted mean (CWM) trait values, which circumvents data limitations

by either focusing directly on trait–environment relationships (based on direct relationships

between environment and CWMs without explicitly considering species distributions or

abundances) or by estimating CWMs directly from species mean trait values (without accounting

for intraspecific variation, and considering environmental variables only for the purpose of

predicting species distributions) (Miller & Ives, 2019).

Models based on direct trait-environment relationships offer the advantage of predicting

trait distributions without requiring data on species abundances. A corollary of this approach is

that it ignores well-established species and phylogenetic signals in trait variation driven by

biological, physical, and historical constraints (Wright et al., 2004, Ordoñez et al., 2009,

Anderegg et al., 2018), and implicitly assumes that the environment captures relevant changes in

species distribution and abundance. However, since species-environment relationships are

typically noisy (Clark 2016), failing to explicitly account for species distribution and abundance

inevitably introduces prediction errors. Furthermore, models that do not explicitly incorporate

species effects cannot be used to study inter- and intraspecific contributions to trait variation,

which has important implications for functional ecology, ecophysiology, and ecosystem

modeling (Anderegg et al., 2018, Osnas et al., 2018, Tautenhahn et al., 2020).

An alternative to pure environmental models is to predict traits directly from species-

level means, which may provide better estimates of large-scale trait distributions (Clark 2016,

Wieczynski et al., 2019, Swenson & Weiser, 2010). Species-based approaches assume that

environmental drivers affect trait distributions only indirectly by shaping community assembly

and therefore that species distributions themselves are the best predictor of traits and associated

ecosystem function. Species-based models can be used to make predictions for traits over large

areas by leveraging national-scale forest inventories or other vegetation networks and can be

combined with species distribution models to forecast shifts in trait distributions (Swenson &

Weiser, 2010, Clark 2016). However, species-level methods ignore intraspecific variation, which

can be larger than interspecific variation for broadly distributed species (Niinemets, 2015,

Messier et al., 2017).

Both pure environment and pure species approaches have contributed to our

understanding of trait variation and allow for predicting community level trait distributions

without requiring extensive field surveys of traits. However, as noted above, neither approach is

designed to study intraspecific trait variation: pure environment approaches cannot separate

inter- and intraspecific variation, and pure species approaches completely ignore intraspecific

variation. Also, these approaches often fail to account for phylogenetic signals among closely

related species (but see Swenson et al., 2017), which is potentially important for generating more

robust trait predictions for species that are sparsely sampled, or not sampled at all (Blomberg et

al., 2003, Swenson 2013, Swenson et al., 2014). These limitations prevent assessment of the

relative importance of inter- and intraspecific variation on trait distributions across a continuum

of geographical scales, reducing our ability to generalize and understand mechanisms driving

trait distributions and inter-trait relationships.

To address these limitations, we developed a model that combines species identity

(including phylogenetic relationships from the Tree of Life; Hinchliff et al., 2015) and

environmental drivers (climate and topography) with leaf trait data from the National Ecological

Observatory Network (NEON) (NEON, 2020). Our approach makes it possible to estimate the

relative contributions of inter- and intraspecific variation to trait distributions at any location or

scale where species abundance data are available. This allows us to address whether changes in

the environment have a direct effect on trait distributions and inter-trait relationships, within and

across species. We jointly modeled eight leaf traits: nitrogen (N%), carbon (C%), chlorophyllA

(ChlA%), chlorophyllB (ChlB%), carotenoids (Crt%), leaf mass per area (LMA, g m^{-2}), lignin

(%) and cellulose (%). We compared our combined species-environment model to models based

on only environmental drivers or only on species and phylogenetic information. We integrated

the combined model with Forest Inventory and Analysis (FIA) data (Bechtold & Patterson. 2005)

and Daymet data (Thornton et al., 2018) to make leaf trait predictions for ~1.2 million trees

across the eastern USA and validated the predictions against independent field datasets.

We compared predictions from our combined model to predictions from environment-

only and species-only approaches to assess the influence of inter- vs. intraspecific variation on

trait distributions across scales, species, and functional types. Specifically, we present four

applications, in which we: (1) quantify the relative contributions of environmental factors vs.

species and phylogenetic information to leaf trait variation at the continental scale; (2) assess

drivers of ecoregional differences in trait distributions; (3) analyze large-scale intraspecific trait

variation driven by environmental drivers; and (4) explore how intraspecific trait variation

impacts the relationship between N% and LMA (one of the LES relationships) and the

implications for representing leaf trait variation in ecosystem models.

Materials and Methods

Data

We used data from NEON (National Ecological Observatory Network, 2020), the

Botanical Information and Ecology Network (BIEN, Maitner et al., 2020) and TRY (Kattge et

al., 2020) to link information on leaf traits, species identity, and approximate locations for

individual trees. We used Foliar Physical and Chemical Properties (DP1.10026.001) and

Vegetation Structure data (DP1.10098.001) from NEON to build joint trait distribution models

with environmental drivers (climate and topography) alone, phylogenetic drivers (species

identity and phylogeny) alone, and both (combined model). Linking the two different NEON

datasets produced individual tree data with stem geolocation and measures of eight leaf traits

(LMA, chlorophyll A and B, carotenoids, lignin, cellulose, C, N) for 477 trees in 20 sites across

the USA (Fig. S.1). Since foliar trait concentrations vary significantly with phenology and

canopy position (Niinemets et al., 2015), foliar samples were collected at the "peak of

greenness" and from the sunlit portion of the canopy. We tested the generalizability of our

approach outside of NEON by evaluating predictions from independent (out-of-sample) eastern

USA data available from the BIEN and TRY datasets (Appendix S1). These two datasets provide

measures for C, N and LMA for a total of 353 individual trees. We used data from the Open Tree

of Life (Redelings 2017) to measure phylogenetic distances among species, which we used to

inform the covariance structure of our combined and species-only models.

Data for environmental drivers included average monthly climate data from 1995 to 2015

(Appendix S1) extracted from Daymet (Thornton et al., 2018) and topographic variables

(elevation, slope, and aspect) reported in the NEON and FIA datasets. For three common eastern

USA tree species (*Acer rubrum*, *Fagus grandifolia*, and *Abies balsamea*), we used all publicly

available leaf N% data from the TRY database to quantify intraspecific variation in leaf N%

across each species' geographic range in the USA. We selected these three species because: (1)

Abies balsamea is the needleleaf species with the most leaf N% data in TRY in the USA; (2)

Fagus grandifolia is the broadleaf species with the most leaf N% data in TRY in the USA; and

(3) *Acer rubrum* occurs throughout much of the eastern USA in a wide variety of habitats (e.g.,

from xeric to mesic) and has abundant leaf N% data in TRY. We combined our trait modeling

approach with data from the USA Forest Inventory and Analysis (https://www.fia.fs.usda.gov/)

(FIA) to estimate traits for all trees surveyed in the FIA across the eastern USA from 2016 to

2019. We used Lv.3 ecoregions and Lv.2 ecoprovinces as defined by the Environmental

Protection Agency (Omernik et al., 2014) to define analysis regions at different scales.

Overview of Models

We modeled the joint multivariate distribution of the eight leaf traits (the response

variables) using three different approaches: (1) environment-only model using climate and

topography as fixed effects; (2) species-only model using species as random effects, with

covariances among the random effects structured by the phylogeny for all woody species

recorded in the FIA database for the eastern USA; and (3) 'combined model' including both

environmental and species effects (throughout this paper, 'species effects' in our models refers to

phylogenetically structured species random effects). Here, we briefly summarize the modeling

framework; full details are in Appendix S2. All approaches used the same joint-multilevel

Bayesian framework and were evaluated using 5-fold cross-validation. The joint structure of the

model allows for modeling all traits simultaneously (Wilkinson et al., 2021), considering the

correlation structure across traits (i.e., inter-trait relationships, such as those typical of the LES).

Environmental effects were fitted using generalized additive models (GAMs) to account for non-

linear relationships. We used thin plate regression splines to estimate the smooth terms using the

brms R package (Bürkner, 2017). Phylogenetic relationships across species were modeled by

including species as a random effect and accounting for their phylogenetic relationships by

estimating their correlation structure from cross-species cophenetic distances (Paradis et al.,

2019). The distance matrix was used to estimate the correlation structure across taxa, allowing

parameter estimates for the lead traits of rare or unsampled taxa to borrow strength from widely

sampled species (de Villemeruil & Nakagawa, 2014). We used multivariate normal families and

weakly informative priors in all cases (Appendix S2).

To reduce collinearity and the number of climate predictors, we performed a PCA for

each climate variable (net radiation, precipitation, vapor pressure, maximum and minimum

temperature) using monthly averages from 1985 to 2015. We used the first component of each

PCA to represent each climate variable in the environment-only and combined models. To

quantify uncertainty in model accuracy, we used the 95% prediction interval of the Bayesian R^2

(Gelman et al., 2018). See additional methods details in Appendix S1-S3.

Partitioning trait variation

We used two factors variance partitioning (Ribas et al., 2006) to estimate the relative

contribution of inter- vs. intraspecific trait variation at the continental scale by comparing each

single-driver model (environment-only and species-only) to the combined model following

Muñoz & Real (2006). Previous studies partitioned inter- vs. intraspecific trait variation using

measured trait values (Leps et al., 2011 de Bello et al., 2011). To adapt this framework to the

modeling context, we quantified the proportion of variance explained (Bayesian R^2) for each trait

by each of our three models (environment, species, and combined models), and we then

quantified the proportion of variance attributable to species effects alone ($\sigma^2_{species} = R^2_{combined} -$

R^2_{env}), environmental effects alone ($\sigma^2_{env} = R^2_{combined} - R^2_{species}$), and joint species-environment

effects ($\sigma^2_{joint} = R^2_{combined} - \sigma^2_{species} - \sigma^2_{env}$). Note that the pure environment effect (σ^2_{env}) is a

component of intraspecific variation, because σ^2_{env} is, by definition, independent of species-level

(interspecific) effects ($\sigma^2_{species}$). However, not all intraspecific variation can be attributed to

measured environmental variables; thus, intraspecific variation contributes not only to σ^2_{env} , but

also to unexplained variation. In summary, trait variation can be conceptualized as including

pure species effects ($\sigma^2_{species}$), pure environmental effects (σ^2_{env}, which is equivalent to

intraspecific variation explained by the measured environmental variables), joint species-

environment effects (σ^2_{joint}), and unexplained variation (which includes measurement error as

well as intraspecific variation associated with errors or omissions in the environmental data).

Analysis methods for quantifying inter- and intraspecific variation at scale

To study inter- and intraspecific patterns in trait variation, including inter-trait

relationships, we used the combined model to estimate leaf traits for all individual trees in FIA

plots in the eastern USA (all USA states bordering or to the east of the Mississippi River). For

each individual tree (~1.2 million trees in ~30,000 inventory plots), we estimated its posterior

median trait values using each of our three models. For the environment-only and combined

models, we used the climate principal components (see above) at the publicly reported location

of each FIA plot, and we used the topographic variables (elevation, slope, and aspect) reported

by FIA for each plot. Predictions from our species-only model produced broad-scale geographic

patterns like those from previous species-only models (Swenson & Weiser, 2010; Clark, 2016)

(Fig. S.2). We compared predictions from our species-only model (which ignores intraspecific

variation) to predictions from the combined model (which includes both inter- and intraspecific

variation) to explore the consequences of intraspecific variation for broad-scale distributions.

To quantify relationships between N% and LMA, we used ordinary least squares (OLS)

regression slopes based on log-transformed predictions of N% and LMA from our combined

model for individual trees in FIA plots. To quantify differences in N% vs. LMA relationships

within and among species and plant functional types, we compared the OLS slopes from

regressions based on individuals vs. species-level means, for all eastern USA species and

separately for broadleaf and needleleaf plant functional types. We used OLS regression instead

of standard major axis (SMA) regression because (1) this facilitated comparisons with the OLS

slopes reported by Osnas et al., (2018), including inter- and intraspecific trait relationships for

different species groups; (2) SMA slopes are unstable when OLS slopes are shallow, as observed

for a large fraction species; and (3) we expect only small errors in the x-variable (LMA) in these

regressions (see additional details on the N% vs. LMA regressions in Supplement 6).

Results and Discussion

Model evaluation: explained variation at continental scale

We fit the species-only, environment-only, and combined (species and environment)

models using leaf trait data from NEON and evaluated their explanatory power using the

Bayesian R^2 of the predicted values for 88 out-of-sample test trees. For all 8 leaf traits, the

combined model explained the largest amount of variance in the out-of-sample test data (average

R^2 across the 8 traits = 0.64), substantially outperforming both the environment-only (average R^2

= 0.35) and species-only (average R^2 = 0.52) models (Fig. 1, Fig. S.3). The combined model had

the highest performance for predicting LMA (R^2 = 0.81) and the lowest performance for

predicting ChlBA (R^2 = 0.51). Uncertainty in predictions was accurately estimated across all

traits and for all models (Fig. S.4), with the combined model's mean 95% coverage values

ranging from 94.3% to 98.8%. Residuals from the combined model – which represent variation

that is unexplained by environmental factors, species identity, or phylogeny – revealed patterns

like those previously described across species, suggesting the presence of inherent trait trade-offs

and correlations that occur both across species, and also within species in a given environment.

Combined-model residuals revealed a strong negative correlation between N% and LMA, as in

the LES (Wright et al., 2004), and the presence of two main trait clusters (Fig. S.5): traits mainly

involved in photosynthesis (N%, Carotenoids, ChlA and ChlB) and traits related to leaf structure

(LMA, C%, Cellulose%, Lignin%).

Model evaluation: geographic transferability

To determine the appropriateness of using the model to make predictions outside of the

scope of the NEON dataset, we tested the performance of the combined model on novel locations

and novel species using independent data from the Botanical Information and Ecology Network

(BIEN) and TRY. These datasets include trait data on LMA, N% and C% for 62 eastern USA

tree species, including 27 species unavailable in training data (Fig. S.1, supplement S1). The

combined model showed good transferability to other data sources (mean $R^2 = 0.54$, 95%

coverage = 91%, Fig. S.6). Accounting for phylogenetic relationships, in addition to

environmental predictors, yielded substantial model improvements when predicting traits for

species not sampled at NEON sites (Fig. 2). This improvement occurs because accounting for

phylogenetic structure when modeling species random effects (see Methods) allows parameter

estimates for unsampled species to borrow strength from closely related species for which

observations are available (Evans et al., 2016). The above transferability analyses suggest that

the combined model is suitable for broad-scale applications.

Model evaluation: intraspecific variation across wide geographic ranges

The combined model produced realistic ranges of intraspecific variation when compared

with available independent data for widely distributed species (Fig. 3). Specifically, intraspecific

ranges of N% predicted from the combined model were similar to those observed in field

samples for three widespread and abundantly sampled species in the NEON, FIA and TRY

datasets (*Abies balsamea, Acer rubrum*, and *Fagus grandifolia*). For these species (Fig. 3b-d),

combined-model predictions of N% spanned an average range of 1.12 N%, which is as wide as

the average difference between evergreen needleleaf and deciduous broadleaf species in the

eastern USA (1.13 N% based on NEON field data). This large magnitude of intraspecific

variation supports the idea that adaptation to the environment accounts for a large proportion of

community-level variation and plays an important role in community-mean trait shifts across

environmental gradients (Albert et al., 2010, Violle et al., 2014, Siefert et al., 2015).

271 Application 1: Role of species and environment in predicting traits at the continental scale

The importance of different environmental drivers varied among traits, supporting an

important role of climate in driving leaf economics in local communities (Ordoñez et al., 2009).

Precipitation and temperature were mainly important for traits involved in photosynthesis (N%,

ChlA, ChlB and Carotenoids), generally having a positive effect on their concentration (except

for ChlB, Fig. S.7). Net radiation showed a negative effect on N%, while vapor pressure

generally had a negative effect on pigments but a positive effect on traits associated with leaf

toughness and durability (cellulose, lignin and C%). Elevation was the most important

topographic predictor and the strongest environmental driver of LMA, consistent with previous

studies (Reich & Oleksyn, 2004, Poorter et al., 2009, Kitajima et al., 2016).

On average, interspecific variation (phylogenetically structured species-level effects,

controlling for environmental variation; see $\sigma^2_{species}$ in Methods) accounted for 25% of the total

explained variation across the 8 traits, intraspecific variation (pure environmental effects,

controlling for species effects; see σ^2_{env} in Methods) accounted for 13%, and joint species-

environment effects accounted for 23% (Fig. 1). However, variance partitioning differed for

photosynthetic traits (pigments and N%) vs. structural traits (cellulose%, C%, lignin%, and

LMA). For photosynthetic traits, the largest variance component was the joint species-

environment component (Fig. 1); i.e., for these traits, much of the explained variation was

associated with coordinated shifts in the environment and species composition. In contrast, for

structural traits, pure species effects (i.e., interspecific variation that is independent of

environmental variation) accounted for most of the trait variation (Fig. 1). It is important to note,

however, that data limitations likely lead us to underestimate the importance of pure

environmental (intraspecific variation) effects. Firstly, our analysis is restricted to the upper (sun-

lit) canopy. Although this is a common practice (Pérez-Harguindeguy et al., 2013), it ignores

trait variation across the canopy light gradient, a major source of intraspecific variation (Lusk et

al., 2008; Osnas et al., 2018). Secondly, 62% of species in our analysis were only sampled within

a single NEON site. Thirdly, as in any observational study, we have only limited environmental

data, which inevitably leads to underestimate environmentally structured intraspecific variation.

Despite the likely underestimation of intraspecific variation in our analysis, pure

environmental effects (a component of intraspecific variation; see Methods) accounted for a

substantial fraction of explained variation for all traits except for N% and LMA (Fig. 1). Thus,

although species-only models accounted for the majority of explained variation (i.e., the sum of

the joint effects and species effects in Fig. 1) – consistent with previous studies highlighting the

important role of species distributions in predicting traits at broad geographic scales (Siefert, et

al., 2015; Clark 2016, Yang & Swenson, 2018) – predictions for most traits can be substantially

improved by a combined species-environment modeling approach. This conclusion – i.e., that

accounting for intraspecific variation can enhance trait predictions − would likely be

strengthened by improved data on traits and environmental variables.

Application 2: Drivers of ecoregional differences in traits distribution

Predictions from the combined model show broad-scale patterns associated with shifts in

forest communities (Fig. S.8) and large-scale climatic and topographic patterns across latitudinal

and altitudinal gradients (Fig. 4). In some cases, trait distributions shifted abruptly between

neighboring ecoregions due to a combination of shifts in local environmental conditions and in

community assembly (independent of the environment). For example, changes in species

composition may explain trait differences between the Mississippi Alluvial Plains and

neighboring ecoregions (Fig. S.9). The Mississippi Alluvial Plains ecoregion is characterized by

agricultural land use, and forests are often limited to riparian ecosystems favoring bottomland

broadleaf species (e.g., *Celtis laevigata*, *Fraxinus pennsylvanica*, *Salix nigra*) in contrast to

needleleaf species (e.g., *Juniperus virginiana*, *Pinus taeda*, and *Pinus echinata*) more common in

the neighboring ecoregions. Being that broadleaf species are generally characterized by higher

N% and lower LMA, this change in community assembly translates into predicted regional trait

patterns. Other ecoregion boundaries show little change in species composition, in which case

geographic variation in traits may be attributed to environmental effects on intraspecific

variation. For example, in the Appalachian Mountains, opposing patterns are exhibited at high

altitudes in the Blue Ridge (lower N% and pigment; higher LMA and C%) compared to valleys

piedmont in the neighboring ecoregions (higher N% & pigments, lower LMA & C%) (Fig. S.10).

Predictions from the combined model differed from the species-only and environment-

only models, with the direction and magnitude of divergences varying across ecoregions (Fig. 5,

Fig. S.11). Species-only and environment-only model predictions differed significantly from the

combined model ($p < 0.0001$ in paired t-tests) in 80% cases (different combinations of traits,

ecoregions, and models). Similarly, predictions from the species-only and environment-only

models differed from each other ($p < 0.0001$ in paired t-tests) for 93% of ecoregion-trait

combinations, which demonstrates the distinct effects of species and environmental drivers on

trait distributions and shows the importance of a combined approach for prediction & inference.

Patterns of divergence from the combined model indicate how species and environmental

effects vary geographically across the eastern USA (Fig. 4). In regions with no significant

divergence (e.g., the Ridge and Valley region), species-only and environment-only models yield

similar results to each other and to the combined model. This convergence suggests a limited role

for intraspecific variation combined with strong environmental filtering of species assembly.

However, for most ecoregions, species-only and environment-only models diverge from the

combined model in opposite directions (i.e., positive divergence for one and negative divergence

for the other) (Fig. 5d, 5e, 5f, S.11). Significant divergence for the species-only model indicates

that continental-scale species averages fail to correctly represent trait values at finer scales, due

to local environmental conditions and/or competition shifting their values away from the mean.

Areas where the combined model predicts higher N% than the species-only model (blue

shading in Fig. 5a) suggest the presence of environmental effects that increase N% above species

means. Conversely, orange-shaded areas in Fig. 5a indicate environmental effects that lower N%

below species means. These patterns may be explained by regulatory mechanisms adjusting

allocation to proteins, pigments, or structural compounds to balance photosynthetic capacity,

toughness, and chemical defense across environmental gradients (Weih & Karlsson, 2001,

Tjoelker et al., 2001, Albert et al., 2010). In contrast, divergence between the combined and

environment-only models (Fig. 5b) could be explained by (i) environmental effects that have a

phylogenetic signal not captured by the environmental variables included in our analysis; and/or

(ii) stochastic factors (e.g., disturbance history and dispersal limitation) that have resulted in

species distribution patterns that are decoupled from current environmental conditions (Burns &

Strauss, 2012, McIntyre et al., 1999). Testing mechanistic hypotheses for the above divergence

patterns is beyond the scope of our study. Nevertheless, identifying these biogeographic patterns

is a step towards better understanding trait distributions and is only possible using modeling

approaches that combine phylogenetic and environmental information.

Application 3: Large scale intraspecific trait variation driven by environmental drivers

In addition to mapping species and environmental trait signals, another essential reason

for a combined approach is that it allows for predictions of intraspecific variation for individual

species (Fig. 3). Models based on community-weighted means, by definition, do not account for

intraspecific variation. In contrast, environment-only models implicitly capture at least some

component of intraspecific variation (see Methods) but cannot separate this intraspecific

variation from trait variation due to shifts in species composition (Hulshof & Swenson, 2010).

Our combined modeling approach makes this separation possible for hundreds of species-trait

combinations. Quantifying intraspecific variation in a comprehensive manner (i.e., for all species

across a large geographic region) is a necessary step towards testing hypotheses about the role of

individual trait variation in species coexistence and interspecific competition (Hart et al., 2016).

Here, we explore the potential for using a combined species-environment approach to studying

intraspecific variation by evaluating how its distribution vary among eastern USA species.

Predicted patterns of intraspecific variation varied widely across species. For N%, the

ratio of predicted intraspecific to total observed interspecific variation ranged from less than 10%

for species with limited geographic ranges (e.g., *Populus heterophylla*, *Sabal palmetto*, and

Gleditsia aquatica) to over 60% for broadleaf species with broad geographic ranges (e.g., *Cercis*

canadensis, Betula lenta, and *Carpinus caroliniana*). Intraspecific responses of N% to

temperature also showed a variety of patterns among species (Fig. 3a, Fig. S.12), including (1)

bell-shaped patterns (n = 85) like those observed in compilations of field data (Reich & Oleksyn,

2004, Laughlin et al., 2012); (2) negative relationships (n = 7); and (3) cases with no significant

relationship (n = 108). Note that our modeling approach does not explicitly include species-by-

environment interactions. Thus, differences among species in their magnitudes of intraspecific

variation and in their trait-environment relationships emerge from how species distributions in

environmental space combine with non-linear trait-environment relationships.

Application 4: Inter- and intraspecific trait relationships: implications for the leaf

economics spectrum and terrestrial ecosystem models

In terrestrial ecosystem models, accurately representing leaf functional diversity (e.g.,

biogeochemical and/or land-surface models) requires capturing variation within and among

species (Bonan et al., 2002, Osnas et al., 2018). The LES provides a simple, one-dimensional

framework for understanding leaf functional variation from local to global scales (Wright et al.,

2004) and for representing plant functional diversity in terrestrial ecosystem models

(Sakschewski et al., 2015), including the land components of Earth System Models (Bonan et al.,

2002, Fisher et al., 2015). On the other hand, divergent patterns in trait variation observed within

species, among species, and among functional or taxonomic groups (Osnas et al., 2018) and at

different taxonomic scales (Anderegg et al., 2018) suggest that a one-dimensional framework

may be too simplistic. There is growing interest in enhanced representations of trait diversity in

terrestrial ecosystem models (Fisher et al., 2015, Sakschewski et al., 2015, Dantas de Paula et al.,

2021), but these model developments are currently data-limited due to limited information on

trait distributions at broad geographic scales (see details below).

When expressed per-unit leaf mass, leaf traits related to photosynthesis, metabolism, and

nutrient concentrations typically have strong negative covariance with LMA across species

(Wright et al., 2004, Lloyd et al., 2013, Osnas et al., 2013) but are often roughly invariant within

species across canopy light gradients (Ellsworth and Reich 1993, Evans and Poorter 2001,

Niinemets et al., 2015). Constant mass-normalized traits across canopy light gradients −

equivalent to a roughly linear increase in the area-normalized value of these same traits as LMA

increases from shade to sun (Osnas et al., 2018) − is accounted for in some terrestrial ecosystem

models (e.g., Sellers et al., 1992, Bonan et al., 2012). Intraspecific variation in leaf traits across

broad environmental gradients can be substantial as well, with important consequences for

ecosystem models (Reich et al., 2014, Dong et al., 2017). Our broad-scale systematic mapping of

N% and LMA, for individual trees (summarized at the FIA plot level in Fig. 4, Fig. S.13) enables

further understanding of intraspecific trait relationships across geographic space.

For the eastern USA, our results show that slopes of N% vs LMA within and among

species fall between the shallow slopes found within species across light gradients (typically

between 0 and −0.3; Supplement 6), and the steeper slopes recorded among global species (−0.6;

Supplement 6). These results were consistent for both deciduous broadleaf (DB) and evergreen

needleleaf (EN) plant functional types (PFTs; Table S.2). Specifically, in our analysis,

intraspecific slopes had median values around −0.37 (Fig. 6a-b), slopes across all individuals

within PFTs were -0.36 for DB and -0.48 for EN (Fig. 6c), and slopes across species means were

-0.4 for DB and -0.45 for EN (Fig. 6d). These intermediate values (i.e., steeper than those

observed across intraspecific light gradients, but shallower than observed across global species)

may be explained by partially compensating factors. LMA variation associated with intraspecific

LL variation across space may largely involve variation in low-nutrient, structural components of

LMA (as occurs across species; Kitajima et al., 2021), which would lead to steep N% vs. LMA

slopes (Osnas et al., 2018). LMA variation associated with certain climatic responses (e.g.,

sensitivity of leaf water potential to tissue water content; Niinemets 2015) may also largely

involve low-nutrient structural mass, which would again lead to steep N% vs. LMA slopes. In

contrast, water loss can be reduced at dry sites (which tend to have high LMA) if N% and thus

photosynthetic efficiency are also high (Wright et al., 2001), which would lead to positive N%

vs. LMA slopes across moisture gradients. This combination of mechanisms, and perhaps similar

mechanisms operating along geographic gradients (e.g., temperature extremes & season length),

may result in intraspecific leaf trait relationships across space that are intermediate to those

driven by photosynthetic responses to light vs. those driven by interspecific variation in LL.

433   Our results suggest that to accurately represent leaf functional variation, terrestrial

ecosystem models may need to consider three different types of variation: intraspecific variation

across light gradients, intraspecific variation across geographic environmental gradients, and

interspecific relationships (which may differ among PFTs). Given the impracticality of

parameterizing species-level global ecosystem models, we expect these models to continue to

explore approaches to representing trait variation within broadly defined PFTs without explicitly

modeling individual species (e.g., Fisher et al., 2015, Sakschewski et al., 2015). Accurately

representing the different intra- and interspecific patterns described above does not necessarily

require an explicit species-level approach. A greater understanding of the anatomical and

physiological mechanisms underlying trait variation (e.g., Niinemets et al., 2015) could lead to a

generalized, species-independent model that captures divergent patterns within & among PFTs.

444   Our results have implications for interpreting interspecific leaf trait relationships. PFT-

level slopes across individuals (Fig. 6c) differed from slopes across species means (Fig. 6d).

These differences in slope are greatly reduced when species are rarified to equal abundance (Fig.

S.14), suggesting that differences in slope are largely due to differences in species abundance:

more common species exert greater influence in the individual-level analysis (Fig. 6c), whereas

all species contribute equally to the species-level analysis (Fig. 6d). Therefore, accounting for

within-species relationships may be relevant for some specific applications. For example, when

parametrizing or testing ecosystem models, trait relationships estimated from individual-level

datasets (that preserve species differences in abundance and intraspecific variation) should more

accurately capture ecosystem-level responses compared to analyses based on species means.

Conclusions

Both phylogenetic and environmental effects are fundamental to understanding drivers

and distribution of plant traits. Combining both in a single model is challenging due to data

limitations but is possible by leveraging large scale datasets. This approach allows for improved

traits predictions compared to models that rely on either species means or environmental

conditions in isolation and allows for robust predictions for species and regions not included in

the training data. Across the eastern USA, interspecific trait variation, geographic variation in

species' abundances, and intraspecific variation all have important effects on trait distributions

and LES relationships. The influence of these drivers varies by species, trait, ecoregion, & scale.

Our approach overcomes data limitations by integrating multiple sources of biological

and environmental information to create a single integrated model. As new traits, phylogenetic,

and vegetation inventory data become available globally, combined modeling approaches, such

as the one presented here, can be applied to other geographic regions to study intraspecific

variation and trait-species-environment relationships. Such analyses are already possible using

national forest inventories in some regions (e.g., Canada, New Zealand & European countries;

Schelhaas et al, 2006, Gills et al., 2005, Paul et al., 2021) in combination with ever-growing

plant trait data (e.g., TRY; https://www.try-db.org/). Expanding this work beyond the USA may

contribute to further understanding the mechanisms driving trait distributions across scales and

the link between traits, species distributions, community assembly, and ecosystem function.

Acknowledgements: This work was supported by the Gordon and Betty Moore Foundation's

Data-Driven Discovery Initiative grant GBMF4563 to E. P. White; National Science Foundation

(NSF) grant 1926542 to E. P. White, S. A. Bohlman, A. Zare, D. Z. Wang, and A. Singh; NSF

grant DEB-1442280 to J. W. Lichstein and S. A. Bohlman; USDA/NIFA McIntire-Stennis

program grant FLA-FOR-005470 to S. A. Bohlman; University of Florida Biodiversity Institute

(UFBI) and Informatics Institute (UFII). No additional funding was received for this study.

Statement of authorship: SM, JL and EW designed the experiment and the methods; SM

performed the analysis; EW, AS, SB and BW supervised the work, helped with experimental

design, and supervised analysis and technical aspects. All authors contributed to the manuscript.

References

Albert, Cécile Hélène, Wilfried Thuiller, Nigel Gilles Yoccoz, Alex Soudant, Florian

Boucher, Patrick Saccone, and Sandra Lavorel. "Intraspecific functional variability: extent,

structure and sources of variation." Journal of Ecology 98, no. 3 (2010): 604-613.

Anderegg, L. D., Berner, L. T., Badgley, G., Sethi, M. L., Law, B. E., &

HilleRisLambers, J. (2018). Within-species patterns challenge our understanding of the leaf

economics spectrum. Ecology letters, 21(5), 734-744.

Bechtold, W. A., and P. L. Patterson. 2005. The enhanced Forest Inventory and Analysis

program - National sampling design and estimation procedures. General Technical Report SRS-

80. USDA Forest Service, Southern Research Station, Asheville, NC.

Bonan, G.B., Levis, S., Kergoat, L. & Oleson, K.W. (2002). Landscapes as patches of

plant functional types: an integrating concept for climate and ecosystem models. Global

Biogeochem. Cy., 16, 1021.

Bruelheide, H., Dengler, J., Purschke, O., Lenoir, J., Jiménez-Alfaro, B., Hennekens,

S.M., Botta-Dukát, Z., Chytrý, M., Field, R., Jansen, F. and Kattge, J., 2018. Global trait–

environment relationships of plant communities. Nature ecology & evolution, 2(12), p.1906.

Bürkner P.C. (2017). brms: An R Package for Bayesian Multilevel Models Using Stan.

Journal of Statistical Software, 80(1), 1-28. doi:10.18637/jss.v080.i01

Burns, J. H., & Strauss, S. Y. (2012). Effects of competition on phylogenetic signal and

phenotypic plasticity in plant functional traits. Ecology, 93(sp8), S126-S137.

Chave, J., 2013. The problem of pattern and scale in ecology: what have we learned in 20

years?. Ecology letters, 16, pp.4-16.

Clark, J. S. (2016). Why species tell more about traits than traits about species: predictive

analysis. Ecology, 97(8), 1979-1993.

De Bello, F., et al., 2011. Quantifying the relevance of intraspecific trait variability for

functional diversity. - Methods in Ecology and Evolution 2: 163-174.

De Villemeruil P. & Nakagawa, S. (2014) General quantitative genetic methods for

comparative biology. Modern phylogenetic comparative methods and their application in

evolutionary biology: concepts and practice (ed. Garamszegi L.) Springer, NY. pp. 287-303.

Dantas de Paula, M., M. Forrest, L. Langan, J. Bendix, J. Homeier, A. Velescu, W.

Wilcke, T. Hickler. 2021. Nutrient cycling drives plant community trait assembly and ecosystem

functioning in a tropical mountain biodiversity hotspot. New Phytologist 232:551–566.

Diaz, S., and M. Cabido. 2001. Vive la difference: plant functional diversity matters to

ecosystem processes. Trends in Ecology & Evolution 16:646–655.

Dong, N., Prentice, I. C., Evans, B. J., Caddy-Retalic, S., Lowe, A. J., & Wright, I. J.

(2017). Leaf nitrogen from first principles: field evidence for adaptive variation with climate.

Biogeosciences, 14(2), 481-495.

Evans, E., Merow, C., Record, S., McMahon, S.M. Enquist, B.J., 2016. Towards process-

based range modeling of many species. Trends in Ecology & Evolution, 31(11), pp.860-871.

Fisher, R. A., Muszala, S., Verteinstein, M., Lawrence, P., Xu, C., McDowell, N. G., ... &

Bonan, G. (2015). Taking off the training wheels: the properties of a dynamic vegetation model

without climate envelopes, CLM4. 5 (ED). Geoscientific Model Development, 8(11), 3593-3619.

Gelman, Ben Goodrich, Jonah Gabry & Aki Vehtari. (2018). R-squared for Bayesian

regression models, The American Statistician.

Hart, S. P., et al., 2016. How variation between individuals affects species coexistence. -

Ecology Letters 19: 825-838.

Henn, J.J., Buzzard, V., Enquist, B.J., Halbritter, A.H., Klanderud, K., Maitner, B.S.,

Michaletz, S.T., Pötsch, C., Seltzer, L., Telford, R.J. and Yang, Y., 2018. Intraspecific trait

variation and phenotypic plasticity mediate alpine plant species response to climate change.

Frontiers in Plant Science, 9, p.1548.

Hinchliff, Cody E., et al., "Synthesis of phylogeny and taxonomy into a comprehensive

tree of life." Proceed. of the Nat. Academy of Sciences 112.41 (2015): 12764-12769.

https://doi.org/10.1073/pnas.1423041112

Kattge, J., Bönisch, G., Díaz, S., Lavorel, S., Prentice, I.C., Leadley, P., Tautenhahn, S.,

Werner, G.D., Aakala, T., Abedi, M. and Acosta, A.T., 2020. TRY plant trait database–enhanced

coverage and open access. Global change biology.

Kitajima K, Wright SJ, Westbrook JW (2016) Leaf cellulose density as the key

determinant of inter- and intra-specific variation in leaf fracture toughness in a species-rich

tropical forest. Interface Focus 6:20150100.

Laughlin, D. C., et al., 2012. A predictive model of community assembly that

incorporates intraspecific trait variation. - Ecology Letters 15: 1291-1299.

Lepš, J., et al., 2011. Community trait response to environment: disentangling species

turnover vs intraspecific trait variability effects. - Ecography 34: 856-863.

Lusk, C.H., Reich, P.B., Montgomery, R.A., Ackerly, D.D. & Cavender-Bares, J. (2008).

Why are evergreen leaves so contrary about shade? Trends Ecol. Evol., 23, 299–303.

Maitner B. (2020). BIEN: Tools for Accessing the Botanical Information and Ecology

Network Database. R package version 1.2.4. https://CRAN.R-project.org/package=BIEN

McGill, B.J., Enquist, B.J., Weiher, E. and Westoby, M., 2006. Rebuilding community

ecology from functional traits. Trends in ecology & evolution, 21(4), pp.178-185.

McIntyre, S., Díaz, S., Lavorel, S., & Cramer, W. (1999). Plant functional types and

disturbance dynamics–Introduction. Journal of Vegetation Science, 10(5), 603-608.

Miller, J. E., Damschen, E. I., & Ives, A. R. (2019). Functional traits and community

composition: A comparison among community-weighted means, weighted correlations, and

multilevel models. Methods in Ecology and Evolution, 10(3), 415-425.

Münzbergová, Z., Hadincová, V., Skálová, H. and Vandvik, V., 2017. Genetic

differentiation and plasticity interact along temperature and precipitation gradients to determine

plant performance under climate change. Journal of Ecology, 105(5), pp.1358-1373.

Muñoz A.-R. & Real R. (2006) Assessing the potential range expansion of the exotic

monk parakeet in Spain. Diversity and Distributions 12: 656-665.

NEON (National Ecological Observatory Network), 2020. Plant foliar traits

(DP1.10026.001). Provisional data http://data.neonscience.org on 2 Jan 2020. Boulder, CO, USA

Niinemets, Ü., Keenan, T.F. and Hallik, L., 2015. A worldwide analysis of within-canopy

variations in leaf structural, chemical and physiological traits across plant functional types. New

Phytologist, 205(3), pp.973-993.

Omernik, J. M., & Griffith, G. E. (2014). Ecoregions of the conterminous United States:

evolution of a hierarchical spatial framework. Environmental management, 54(6), 1249-1266.

Ordoñez, J. C., et al., "A global study of relationships between leaf traits, climate and soil

measures of nutrient fertility." Global Ecology and Biogeography 18.2 (2009): 137-149.

Osnas, J.L., Katabuchi, M., Kitajima, K., Wright, S.J., Reich, P.B., Van Bael, S.A., Kraft,

N.J., Samaniego, M.J., Pacala, S.W. and Lichstein, J.W., 2018. Divergent drivers of leaf trait

variation within species, among species, and among functional groups. Proceed. of the Nat.

Academy of Sciences, 115(21), pp.5480-5485.

Paradis, E., Blomberg, S., Bolker, B., Brown, J., Claude, J., Cuong, H.S. and Desper, R.,

2019. Package 'ape'. Analyses of phylogenetics and evolution, version, 2(4).

Peaucelle, M., Bacour, C., Ciais, P., Vuichard, N., Kuppel, S., Peñuelas, J., Belelli

Marchesini, L. et al., 2019. Covariations between plant functional traits emerge from

constraining parameterization of a terrestrial biosphere model. Global Ecol. and Biogeography.

Pérez-Harguindeguy, N., et al., 2013. New handbook for standardised measurement of

plant functional traits worldwide. - Australian Journal of Botany 61: 167-234.

Poorter, H., Niinemets, Ü., Poorter, L., Wright, I.J. and Villar, R., 2009. Causes and

consequences of variation in leaf mass per area (LMA): a meta-analysis. New phytologist,

182(3), pp.565-588.

Redelings BD, Holder MT. "A supertree pipeline for summarizing phylogenetic and

taxonomic information for millions of species." PeerJ. 2017;5:e3058.

Reich, Peter B. "The world-wide 'fast–slow' plant economics spectrum: a traits

manifesto." Journal of Ecology 102.2 (2014): 275-301.

Reich, P.B., Oleksyn., J., 2004. Global patterns of plant leaf N and P in relation to

temperature and latitude. Proceed. of the Nat. Academy of Sciences, 101 (30) 11001-11006;

Reichstein, M., Bahn, M., Mahecha, M.D., Kattge, J. and Baldocchi, D.D., 2014. Linking

plant and ecosystem functional biogeography. Proceed. of the Nat. Academy of Sciences,

111(38), pp.13697-13702.

Ribas A., Barbosa A.M., Casanova J.C., Real R., Feliu C. & Vargas J.M. (2006)

Geographical patterns of the species richness of helminth parasites of moles (Talpa spp.) in

Spain: separating the effect of sampling effort from those of other conditioning factors. Vie et

Milieu 56: 1-8.

Schelhaas, M.J., Varis, S., Schuck, A. and Nabuurs, G.J., 2006, EFISCEN Inventory

Database, European Forest Institute, Joensuu, Finland, http://www.efi.int/

Siefert, A., et al., 2015. A global meta-analysis of the relative extent of intraspecific trait

variation in plant communities. - Ecology Letters 18: 1406-1419.

Simpson, A.H., Richardson, S.J. and Laughlin, D.C., 2016. Soil–climate interactions

explain variation in foliar, stem, root and reproductive traits across temperate forests. Global

Ecology and Biogeography, 25(8), pp.964-978.

Sterck, F., Markesteijn, L., Schieving, F. and Poorter, L., 2011. Functional traits

determine trade-offs and niches in a tropical forest community. Proceed. of the Nat. Academy of

Sciences, 108(51), pp.20627-20632.

Swenson, N.G., Weiser, M.D., Mao, L., Araújo, M.B., Diniz-Filho, J.A.F., Kollmann, J.,

Nogués-Bravo, D., et al., 2017. Phylogeny and the prediction of tree functional diversity across

novel continental settings. Global Ecology and Biogeography, 26(5), pp.553-562.

Swenson, N.G., 2014. Phylogenetic imputation of plant functional trait databases.

Ecography, 37(2), pp.105-110.

Swenson, N.G., 2013. The assembly of tropical tree communities–the advances and

shortcomings of phylogenetic and functional trait analyses. Ecography, 36(3), pp.264-276.

Swenson, N. G., & Weiser, M. D. (2010). Plant geography upon the basis of functional

traits: an example from eastern North American trees. Ecology, 91(8), 2234-2241.

Tautenhahn, S., Migliavacca, M. & Kattge, J. (2020). News on intra-specific trait

variation, species sorting, and optimality theory for functional biogeography and beyond. New

Phytologist, 228, 6–10.

Tjoelker, M. G., Oleksyn, J., & Reich, P. B. (2001). Modelling respiration of vegetation:

evidence for a general temperature dependent Q10. Global change biology, 7(2), 223-230.

Thornton, M.M., P.E. Thornton, Y. Wei, B.W. Mayer, R.B. Cook, and R.S. Vose. 2018.

Daymet: Annual Climate Summaries on a 1-km Grid for North America, Version 3. ORNL

DAAC, Oak Ridge, Tennessee, USA. https://doi.org/10.3334/ORNLDAAC/1343

USDA Forest Service, 2001. Forest Inventory and Analysis National Core Field Guide,

Volume I: Field Data Collection Procedures For Phase 2 Plots, Version 1.5. US Department of

Agriculture, Forest Service, Washington, DC.

Verheijen, L.M., Brovkin, V., Aerts, R., Boenisch, G., Cornelissen, J.H.C., et al., (2013).

Impacts of trait variation through observed trait-climate relationships on performance of an Earth

system model: a conceptual analysis. Biogeosciences, 10, 5497–5515.

Violle, C., Reich, P.B., Pacala, S.W., Enquist, B.J. and Kattge, J., 2014. The emergence

and promise of functional biogeography. Proceed. of the Nat. Academy of Sciences, 111(38),

pp.13690-13696.

Weih, M., & Karlsson, P. S. (2001). Growth response of Mountain birch to air and soil

temperature: is increasing leaf-nitrogen content an acclimation to lower air temperature?. New

Phytologist, 150(1), 147-155.

Weng, Ensheng, Caroline E. Farrior, Ray Dybzinski, and Stephen W. Pacala. "Predicting

vegetation type through physiological and environmental interactions with leaf traits: evergreen

and deciduous forests in an earth system modeling framework." Global change biology 23, no. 6

(2017): 2482-2498.

Wieczynski, D.J., Boyle, B., Buzzard, V., Duran, S.M., Henderson, A.N., Hulshof, C.M.,

Kerkhoff, A.J., McCarthy, et al., 2019. Climate shapes and shifts functional biodiversity in

forests worldwide. Proceed. of the Nat. Academy of Sciences, 116(2), pp.587-592.

Wilkinson, D. P., Golding, N., Guillera-Arroita, G., Tingley, R., & McCarthy, M. A.

(2021). Defining and evaluating predictions of joint species distribution models. Methods in

Ecology and Evolution, 12(3), 394-404.

Wright, I.J., Reich, P.B., Westoby, M., Ackerly, D.D., Baruch, Z., Bongers, F.,

Cavender-Bares, J., Chapin, T., Cornelissen, J.H., Diemer, M. and Flexas, J., 2004. The

worldwide leaf economics spectrum. Nature, 428(6985), p.821.

Yang, J., Cao, M. and Swenson, N.G., 2018. Why functional traits do not predict tree

demographic rates. Trends in ecology & evolution, 33(5), pp.326-336

Fig. 1. Variance partitioning between pure species effects, pure environmental effects

(intraspecific variation), and joint species-environment effects based on the R^2 for independent

validation data. Pure environmental variation is independent of species variation, and vice versa.

Joint variation is jointly explained by species and environmental effects and cannot be uniquely

ascribed to either. LMA is leaf dry mass per area, and all other traits are % mass.

Fig. 2. Observed vs. predicted values from the combined model for (a) LMA (g m-2), (b) N%,

and (c) C%, either assuming phylogenetically-structured covariance among species random

effects (azure), or assuming independent species random effects (orange). Ellipses are 95%

confidence ellipses. (d) RMSE (normalized by standard deviation) showing the difference

between accounting for (azure) or ignoring (orange) phylogenetic relationships when predicting

traits for species not included in the training data.

Fig. 3. Intraspecific variation of predicted leaf N% for tree species in the eastern USA. (a)

Relationships (estimated using GAMs) of N% (median predicted values) vs. maximum monthly

temperature for the 91 species (each species has a separate line) showing significant effects of

temperature on N% (the darkness of the curve increases with the strength of the relationship

between N% and temperature). (b-d) Geographic patterns and range of intraspecific N% for two

widespread deciduous broadleaf (*Acer rubrum* and *Fagus grandifolia*) and one widespread

needleleaf evergreen (*Abies balsamea*) tree species. Mean (points) and 99% intraspecific ranges

(lines) in b-d are from: TRY data (within the eastern USA), NEON data (within the eastern

USA), the combined model, and random error from the species-only model. Red crosses

represent TRY sample locations.

Fig. 4. Distributions across the eastern USA for leaf structural (LMA, lignin, cellulose and C%),

macro-nutrient (N%), and pigment (carotenoids, chlorophyll A and B) traits. Values represent

trait averages at each FIA plot (n = 30,331 plots), calculated across all individual trees sampled

within the plot.

Fig. 5. Divergence in N% between the combined model and the species-only (a) and

environment-only (b) models. Dots represent FIA plots where predicted N% was higher in the

combined (blue), environment (orange), or species (green) model. White areas represent plots

with little to no divergence (p > 0.0001), or areas with no FIA plots recorded. (c-f) Examples of

divergence patterns in four ecoregions (see the outlined ecoregions in the maps, labeled c-f

according to their corresponding histogram): combined model minus environment model

(orange) and combined model minus species model (green). Vertical black lines represent zero

divergence. Ecoregion examples include: all three models capturing similar information with

ΔN% near 0 (c); species and environment models similar to each other but diverging from the

combined model (d); and and divergence among all three models (e-f).

Fig. 6. Relationships between leaf nitrogen percent mass (N%) and leaf mass per area (LMA) across the eastern USA, based on combined-model predictions of N% and LMA for individual trees in eastern USA Forest Inventory and Analysis (FIA) plots. (A) Fitted intraspecific ordinary least squares (OLS) regression lines for eastern USA deciduous broadleaf (DB: blue) and evergreen needleleaf (EN: orange) species. Each line is an individual species. (B) Distribution of intraspecific OLS slopes for DB and EN species; bars are medians, boxes span the interquartile range, whiskers span 95% of slopes, and points show outliers. (C) OLS slopes (± s.e.) across all individuals within each PFT. (D) OLS slopes (± s.e.) across species means within each PFT. Individual trait values were log_{10}-transformed for analyses in (A-C), and species means were log_{10}-transformed for analyses in (D). Dashed and dotted black lines in panels C and D are from Osnas et al., (2018): the dashed line at -0.18 represents the intraspecific slope across the canopy light gradient, and the dotted black line at -0.62 represents the slope across global species (see details in Supplement 6).