Deficiency of the paternally inherited gene *Magel2* alters the development of separation-

induced vocalization in mice

Authors:

Gabriela M. Bosque Ortiz [1,2], Gustavo M. Santana[1,2], Marcelo O. Dietrich [1,2,3,4]*

[1] Laboratory of Physiology of Behavior, Department of Comparative Medicine, Yale School of

Medicine, New Haven, CT, USA.

[2] Interdepartmental Neuroscience Program, Biological and Biomedical Sciences Program,

Graduate School in Arts and Sciences, Yale University, New Haven, CT.

[3] Program in Integrative Cell Signaling and Neurobiology of Metabolism, Department of

Comparative Medicine, Yale School of Medicine, New Haven, CT, USA.

[4] Department of Neuroscience, Yale School of Medicine, New Haven, CT, USA.

* Correspondence to: marcelo.dietrich@yale.edu

Abstract

Offspring behavior results from the combined expression of maternal and paternal genes.

Genomic imprinting, however, silences some genes in a parent-of-origin specific manner, a

process that, among all animals, occurs only in mammals. How genomic imprinting affects the

behavior of mammalian offspring remains poorly understood. Here we studied the effects of the

loss of the paternally inherited gene *Magel2* on the emission of separation-induced ultrasonic

vocalization (USV) by mouse pups. Using quantitative analysis of more than one hundred

thousand USVs, we characterized the rate of vocalizations as well as their spectral features

from postnatal days 6 to 12 (P6-P12), a critical phase during mouse development when pups

fully depend on the mother for survival. Our analyses show that *Magel2* deficient offspring emit

separation-induced vocalizations at lower rates and with altered spectral features. Using

methods for a holistic analysis of the entire vocal repertoire of pups, we found that *Magel2*

deficient mice at postnatal day 8 (P8) emit USVs that resemble the vocal repertoire of wildtype

mice at older ages (P10-12). These results suggest that the deficiency of this paternally

inherited gene impairs the expression of separation-induced vocalization in pups, a behavior

that supports the pups' growth and development.

Keywords

Genomic imprinting; offspring-parent conflict; Prader Willi Syndrome; Autism spectrum

disorders; behavior development.

Introduction

For the normal development of mammals, offspring need copies from both maternal and paternal genomes. Some genes, however, are expressed in a parent-of-origin specific manner. In other words, some genes are always expressed when inherited from the mother and some genes are always expressed when inherited from the father [1–3]. The process that regulates the expression of genes in a parent-of-origin specific manner is called genomic imprinting [2,3].

Genomic imprinting depends on epigenetic modifications of the genome. These modifications do not alter the sequence of the DNA but the chemical structure of the DNA, thereby leading to altered gene expression [4]. An imprinted gene can be silenced in the maternal genome and, therefore, only the paternal allele will be expressed in the offspring. Consider, for example, a series of imprinted genes in human chromosome 15, which illustrates the importance of paternally inherited genes for normal mammalian development [5–9]. The deletion of the paternally inherited genes in chromosome 15 leads to neurodevelopmental disorders, such as Prader-Willi syndrome (PWS). PWS presents with hypotonia and poor feeding early in life, followed by hyperphagia, alteration in social behavior, and cognitive deficits. It should be noted, however, that because PWS involves several genes, it masks the relative contribution of single imprinted genes on the phenotype of the offspring.

Among the PWS-related genes, MAGEL2 is a candidate gene for some of the clinical features of PWS. Humans with loss-of-function mutations in MAGEL2 present clinical aspects of PWS and autism spectrum disorders [10–12], suggesting that this single paternally inherited gene supports at least some of the developmental alterations found in PWS. In agreement with the clinical features of MAGEL2 deficiency in humans, *Magel2* deficient mice show impairments in growth and adult social behaviors which are, however, of small-effect size [10,13,14].

In this study, we investigated whether the loss of paternally inherited *Magel2* in mice alters

behaviors that are ecologically relevant for the development of the offspring. We chose to study

the vocal behavior of mouse pups when separated from their dams [15–23], as separation-induced

vocalizations signal the needs of the pups to the dams [19,24–29]. In contrast to human babies,

mouse pups vocalize in the ultrasonic frequency range (45 - 120 kHz) [20,21,23,30], which humans

cannot hear. In order to survey the vocal behavior of mice, we recorded the emission of

ultrasonic vocalizations (USVs) when pups were separated from the home nest. We performed

these studies at postnatal days 6, 8, 10, and 12, since it is during this phase of mouse

development that the peak expression of separation-induced vocalizations typically occurs

[15,31,32]. We then used VocalMat [33], a software developed by our group, to perform quantitative

analysis of mouse vocal behavior. Our analysis shows that the deficiency of *Magel2* in mice

impairs the expression of separation-induced vocalizations, a behavior that supports pups'

growth and development.

Results

Early waning of vocal behavior in *Magel2*$^{m+/p-}$ deficient pups

To investigate the effects of paternally inherited *Magel2* on the vocal behavior of infant mice, we

crossed heterozygote males for *Magel2* deficiency with wildtype females. From this cross, we

generated *Magel2* deficient offspring (*Magel2*$^{m+/p-}$) that carry the null allele from the father (p−)

and the imprinted allele from the mother (m+). This cross also generates wildtype littermates

(*Magel2*$^{m+/p+}$), used as experimental controls. As previously reported[10], *Magel2*$^{m+/p-}$ pups display

lower body weight compared to controls (genotype: $F_{1, 182}$ = 19.91, $P < 10^{-4}$; age: $F_{3, 182}$ = 70.09,

$P < 10^{-9}$; genotype x age: $F_{3, 182}$ = 0.02, P = 0.996; two-way ANOVA).

We recorded the emission of USVs during 20 minutes of separation from the home nest at

different postnatal ages (P6, P8, P10 and P12; **Figure 1A**). First, we analyzed the total number

of USVs emitted during the period of separation using two-way ANOVA. We found a significant

effect of genotype, age, and interaction between genotype and age (genotype: $F_{1, 181}$ = 20.61, P

$< 10^{-4}$; age: $F_{3, 181}$ = 11.80, $P < 10^{-6}$; genotype x age: $F_{3, 181}$ = 3.90, P = 0.009; **Figure 1B**). Post-

hoc analysis (Sidak's multiple comparisons test; **Figure 1B**) shows that the total number of

USVs is similar among groups at P6 (control: 991 ± 139 USVs, n = 16; *Magel2*$^{m+/p-}$: 787 ± 130

USVs, n = 20; P = 0.75), P10 (control: 503 ± 45 USVs, n = 23; *Magel2*$^{m+/p-}$: 466 ± 70 USVs, n =

20; P = 0.98), and P12 (control: 594 ± 71 USVs, n = 30; *Magel2*$^{m+/p-}$: 345 ± 46 USVs, n = 28; P

= 0.06). Compared to controls, however, *Magel2*$^{m+/p-}$ pups show a ≈53% reduction in the

emission of USVs at P8 (control: 1045 ± 100 USVs, n = 24; *Magel2*$^{m+/p-}$: 495 ± 65 USVs, n = 28;

$P < 10^{-4}$). It is worth noting that P8 *Magel2*$^{m+/p-}$ pups emit a similar number of USVs compared to

P10 and P12 control pups, suggesting an early decline in vocal behavior. We also performed

these analyses in females and males separately and found similar effects of genotype and age

(**Figure 1C-D** and **Supplementary Table 1**). These results show age-specific reductions in the

emission of USVs in *Magel2*[m+/p−] mice, suggesting a non-sex specific effect for paternally

inherited *Magel2* on the vocal behavior of the offspring.

The emission of USVs occurs when the breathing musculature contracts, expelling air from the

lungs and propelling it through the larynx [29,34]. Since previous reports found that *Magel2*[m+/p−]

mice display hypotonia [35], we considered the hypothesis that the low rate of USV emission in

Magel2 deficient pups is due to a lower capacity to expel air from the lungs. To rule out this

hypothesis, we measured the intensity (or volume, in decibels) of the USVs. Since the intensity

of the USVs relates to the pressure by which the air is expelled through the larynx [29,34], a lower

intensity is expected in cases of hypotonia. This analysis shows that the intensity of the emitted

USVs between *Magel2*[m+/p−] mice and controls is similar in all ages tested (genotype: $F_{1,\,175}$ =

0.82, P = 0.63; age: $F_{3,\,175}$ = 2.64, P = 0.05; genotype x age: $F_{3,\,175}$ = 0.68, P = 0.56; two-way

ANOVA; **Figure 1E**). Thus, hypotonia does not seem to be a factor of primary significance for

the lower rate of USV emission in *Magel2*[m+/p−] mice [34].

***Magel2*[m+/p−] mice emit vocalizations with distinct spectral features**

In addition to the rate of separation-induced vocalizations, the spectral features of the USVs

also correlate with altered maternal care [36]. To test the extent to which *Magel2* deficiency affects

the spectral features of USVs across ages, we used two-way ANOVA to analyze the frequency

characteristics (pitch) and duration (**Figure 2A**) of USVs [33]. We found significant effects of

genotype and age for maximal frequency (genotype: $F_{1,\,182}$ = 20.38, $P < 10^{-4}$; age: $F_{3,\,182}$ = 2.79,

P = 0.04; genotype x age: $F_{3,\,182}$ = 6.45, P = 0.0004; **Figure 2B**) and bandwidth (genotype: $F_{1,\,182}$

= 9.79 P = 0.002; age: $F_{3,\,182}$ = 6.94, P = 0.0002; genotype x age: $F_{3,\,182}$ = 5.88, P = 0.0007;

Figure 2E). When comparing *Magel2*[m+/p−] to control mice, post-hoc analysis (Sidak's multiple

comparisons test) shows that at P8—but not at P6, P10, or P12—there is a 3% reduction in

maximal frequency (control: 88.4 ± 1.5 kHz, n = 24; *Magel2*[m+/p−]: 85.6 ± 1.1 kHz, n = 28; $P < 10^{-}$

[4]; **Figure 2B**) and a 40% reduction in bandwidth (control: 17.3 ± 1.2 kHz, n = 24; *Magel2*[m+/p−]:

10.4 ± 0.7 kHz, n = 28; $P < 10^{-5}$; **Figure 2E**). Moreover, we found a significant effect of genotype

for mean frequency (genotype: $F_{1, 181} = 4.86$, $P = 0.02$; age: $F_{3, 181} = 0.24$, $P = 0.87$; genotype x

age: $F_{3, 181} = 0.41$, $P = 0.74$; **Figure 2D**), but not for minimal frequency (genotype: $F_{1, 182} = 0.54$,

$P = 0.46$; age: $F_{3, 182} = 1.97$, $P = 0.12$; genotype x age: $F_{3, 182} = 0.47$, $P = 0.83$ **Figure 2C**) or

duration (genotype: $F_{1, 175} = 3.54$, $P = 0.06$; age: $F_{3, 175} = 5.78$, $P = 0.0009$; genotype x age: $F_{3,}$

$_{175} = 1.98$, $P = 0.11$; **Figure 2F**). In addition to the main frequency component, USVs can

contain harmonics (**Figure 2G**). We calculated the percentage of USVs with harmonics and

found a significantly lower number in *Magel2*[m+/p−] mice compared to controls at P8 (control: 9.1 ±

1.6 % , n = 24; *Magel2*[m+/p−]: 3.4 ± 0.8 %, n = 28; $U = 146$, $P_{2-tailed} = 0.0003$, Mann-Whitney test;

Figure 2H) but not at P6 (control: 2.0 ± 0.2 % , n = 16; *Magel2*[m+/p−]: 2.1 ± 0.2 %, n = 20; $U =$

138, $P_{2-tailed} = 0.49$, Mann-Whitney test; **Figure 2H**), P10 (control: 5.7 ± 0.8 % , n = 23;

Magel2[m+/p−]: 6.1 ± 1.2 %, n = 20; $U = 240$, $P_{2-tailed} = 0.98$, Mann-Whitney test; **Figure 2H**), or P12

(control: 3.6 ± 0.6 % , n = 30; *Magel2*[m+/p−]: 4.9 ± 0.1 %, n = 28; $U = 408.5$, $P_{2-tailed} = 0.86$, Mann-

Whitney test; **Figure 2H**). In sum, these results suggest that the loss of paternally inherited

Magel2 in mice causes discrete changes in the features of separation-induced vocalizations that

are most evident at postnatal day eight.

Discrete changes in the use of syllable types by *Magel2*[m+/p−] mice

Mouse pups emit USVs of distinct classes—i.e., syllable types. Thus, the emission of different

syllable types could explain the discrete changes in the spectro-temporal features of USVs in

Magel2[m+/p−] (**Figure 3A**) [33,37]. We used machine learning to automatically categorize each USV

into one of eleven syllable types based on the morphology of the main component of the

vocalization. The output of the method was the probability for each USV to be of a certain

syllable type. The highest probability (P_1) defined the syllable type for a given USV (**Figure 3B**).

Using this approach, we did not find any significant differences in the distribution of syllable

types emitted by control and *Magel2*$^{m+/p-}$ mice at P6, P10, and P12 (**Figure 3C, 3E, 3F**). At P8,

however, *Magel2*$^{m+/p-}$ pups emit 54% less USVs of the type *chevron* (control: 8.7 ± 1.6%; n =

24; *Magel2*$^{m+/p-}$: 4.0 ± 0.9 %; n = 28; *P* = 0.008), 71% less USVs of the type *step-down* (control:

0.7 ± 0.1 %; n = 24; *Magel2*$^{m+/p-}$: 0.2 ± 0.05 %; n = 28; *P* < 10^{-4}), 44% less USVs of the type

step-up (control: 14.5 ± 1.4 %; n = 24; *Magel2*$^{m+/p-}$: 8.1 ± 1.3 %; n = 28; *P* = 0.001), and 87%

less USVs of the type *two-steps* (control: 3.6 ± 1.1 %; n = 24; *Magel2*$^{m+/p-}$: 0.5 ± 0.2 %; n = 28; *P*

= 0.005) compared to controls (**Figure 3D**). Conversely, *Magel2*$^{m+/p-}$ pups emit 36% more USVs

of the type *flat* (control: 9.7 ± 0.9 %; n = 24; *Magel2*$^{m+/p-}$: 15.2 ± 1.2 %; n = 28; *P* = 0.007) and

33% more USVs of the type *short* compared to controls (control: 23.8 ± 1.8 %; n = 24;

Magel2$^{m+/p-}$: 35.5 ± 2.0 %; n = 28; *P* < 10^{-4}; **Figure 3D)**. (**Supplementary Table 2** provides a

detailed analysis of the spectro-temporal features of each syllable type across all ages tested in

controls and *Magel2*$^{m+/p-}$ pups.) In summary, we found that *Magel2*$^{m+/p-}$ mice at P8 use simpler

vocalizations that fall under the 'flat' and 'short' classifications. These findings are in line with

our previous results **(Figure 1-2)** demonstrating that the largest differences in vocal behavior

occur in eight-day-old *Magel2*$^{m+/p-}$ pups.

Altered vocal repertoire of *Magel2*$^{m+/p-}$ mice

As stated above, the vocal analysis pipeline outputs the probability for each USV to be classified

as each of the eleven syllable types (P_1, P_2, P_3, … $P_{11;}$ **Figure 3A-B**). This distribution of

probabilities allows the qualitative and quantitative comparison of the vocal classification among

groups [33]. By considering the distribution of probabilities to classify each USV, it is possible to

estimate how similar the vocal repertoire of one group of mice is to another group. To compare

the vocal repertoire of mice across all ages studied, we used diffusion maps—a dimensionality

reduction technique that decreases the number of dimensions of the probability distribution from

eleven classes to three dimensions in a Euclidean space (**Figure 4A**) [33]. Using pairwise

comparisons (**Figure 4B**), we estimated the similarity between the vocal repertoire of mice of

different ages and genotypes. Using this method, we found that control pups at P6 and P8

(Cohen's coefficient: κ = 0.99) and control pups at P10 and P12 (Cohen's coefficient: κ = 0.95)

display vocal repertoires that are similar to each other (**Figure 4**). These two age groups (P6-P8

and P10-P12), however, present lower pairwise similarities when compared to each other with κ

ranging from 0.67 to 0.77 (**Figure 4**). These results suggest that the vocal repertoire of control

pups undergoes significant changes between P8 and P10.

Next, we analyzed the same transitions in the vocal repertoire of $Magel2^{m+/p-}$ pups. The

comparison between the vocal repertoire of $Magel2^{m+/p-}$ pups at P6 and P8 show lower pairwise

similarity (κ = 0.80) compared to control pups (**Figure 4**). In contrast to littermate controls,

$Magel2^{m+/p-}$ pups at P8 show a higher pairwise similarity with P10 (κ = 0.84) and P12 (κ = 0.99)

pups. These findings suggest that $Magel2^{m+/p-}$ pups switch their vocal repertoire at a younger

age (P8) to a vocal repertoire that control pups emit at an older age (P10-P12).

Finally, we directly compared the vocal repertoire of $Magel2^{m+/p-}$ and control pups. $Magel2^{m+/p-}$

pups, at P6, show high pairwise similarity when compared to controls at P6 (κ = 1.00) and P8 (κ

= 1.00), but not at P10 (κ = 0.80) and P12 (κ = 0.72). At P8, $Magel2^{m+/p-}$ pups show relatively

low pairwise similarities with control pups at P6 (κ = 0.76) and P8 (κ = 0.78) but show high

similarities with control pups at P10 (κ = 1.00) and P12 (κ = 0.95). At P10, $Magel2^{m+/p-}$ pups

show relatively lower pairwise similarities with control pups at P6 (κ = 0.67) and P8 (κ = 0.70)

than at P10 (κ = 0.82) and P12 (κ = 0.81). This pattern is more evident in P12 $Magel2^{m+/p-}$ pups,

which show lower pairwise similarities with control pups at P6 (κ = 0.74) and P8 (κ = 0.75) than

at P10 (κ = 0.99) and P12 (κ = 0.97). Altogether, these analyses suggest that the vocal

repertoire of $Magel2^{m+/p-}$ pups transforms at a faster pace than control pups—with $Magel2^{m+/p-}$

pups at P8 resembling control pups at P10-P12. Thus, the critical phase of development

between P8 and P10 seems to mark an important period for the effect of the maternally

imprinted gene, *Magel2*, on the vocal behavior of the offspring.

Discussion

In this study, we recorded and analyzed vocalizations from *Magel2* deficient pups and their wildtype littermates during a 20-minute period at postnatal days 6, 8, 10 and 12. Using custom-built software to automatically analyze vocalizations [33], we counted the number of vocalizations and measured the spectro-temporal features of each vocalization, including its intensity, duration, bandwidth, mean frequency, maximum frequency, minimum frequency, and use of harmonic components. We also assigned a syllable type for each vocalization based on its morphological features in the time-frequency plane. We further used quantitative methods to analyze the vocal repertoire of mice across groups and ages. These methods shed light on discrete changes in the development of separation-induced vocalizations in *Magel2* deficient mice.

Using our software allowed us to expand the period of analysis to 20 minutes. By comparison, previous studies quantified vocal behavior during a much shorter period—typically between 1 to 5 minutes. The longer period of analysis enabled us to characterize more than a hundred thousand vocalizations, which provided an in-depth view of the vocal repertoire of pups. With regards to the vocal behavior of wildtype mice, our results demonstrated that the emission of separation-induced USVs gradually decreases from P6-P8 to P10-P12. This result agrees with previous studies, which show an inverse-U shape profile for separation-induced vocalizations in mice during the first two weeks of life [15,29,31,32,38]. Based on the large number of vocalizations we analyzed, we also found that wildtype mice use simpler vocalizations at older ages (P10-P12) compared to younger ages (P6-P8). These findings in wildtype mice provide the basis for comparisons with *Magel2* deficient pups.

Intriguingly, *Magel2* deficient pups show different dynamics for separation-induced

vocalizations. At P6, these pups vocalize comparably to wildtype littermates, but at P8, their

vocal number and features resemble wildtype littermates that are older (P10-P12). An

explanation for these results is that *Magel2* deficient pups are less responsive to certain social

cues. In socially isolated pups, therefore, the deprivation of these cues would not induce the

behavior to the same degree as in wildtype pups. An alternative explanation for these results is

that for *Magel2* deficient pups' vocal behavior does not have the same fitness value compared

to wildtype pups. In the latter case, the behavior begins to change at younger ages due to the

lack of reinforcement. While it is difficult to test these interpretations experimentally, the fact that

Magel2 deficient pups have lower body weight during early development suggests a decrease in

the fitness of these animals and would support the idea that the development of vocal behavior

in these mice is impaired. Whether the change in vocal behavior and the decrease in body

weight are causally related warrants further investigation.

The discrepancies in expression of vocal behavior between wildtype and *Magel2* deficient pups

support a more general role for imprinted genes on offspring behavior. Consider, for example,

previous studies of other imprinted genes. Deletion of the paternally inherited imprinted gene

Peg3 lowers vocal rate in mouse pups [39]. Conversely, deletion of the maternally inherited gene

Gabrb3 increases vocal rate [40]. Thus, imprinted genes inherited from the father seem to

increase vocal rate while imprinted genes inherited from the mother seem to decrease vocal

rate. In evolutionary terms, these findings support the theory that genomic imprinting evolved to

balance the cost of the phenotype for the offspring and for the mother, as well as to balance the

best interests of mothers and fathers in altering offspring's phenotype. In the case of paternally

inherited genes, therefore, the expression of these genes favors the use of maternal resources,

which is in the best interests of the father, by increasing vocal behavior to increase maternal

care. Conversely, the loss of these paternally inherited genes favors the conservation of

maternal resources, which is in the best interests of the mother, by decreasing vocal behavior

and, consequently, the demand for maternal care [1,41]. Again, the lower body weight of *Magel2*

pups supports this argument [10]. Future studies should test this theoretical view more directly by

systematically investigating the effect of imprinted genes on the behavior of offspring and on the

behavior of mothers towards their offspring.

Acknowledgements

We thank Yuexuan Li, Delva Leão, Dr. Onur Iyilikci, and Dr. Mahdieh Godazgar for critical feedback on the project and the manuscript. Special thanks to: Delva Leão for her contribution to our analysis; Jeremy Bober for helping with animal maintenance and care; as well as Antonio Fonseca who built the VocalMat software used for our analysis of ultrasonic vocal recordings. We thank David Gillich, Susan Andranovich, Valeria Krizsan, and Vickie Clark for administrative support. M.O.D. was supported by a NARSAD Young Investigator Grant ID 22709 from the Brain & Behavior Research Foundation, by the National Institute of Diabetes and Digestive and Kidney Diseases of the National Institutes of Health (R01DK107916, P30 DK045735), by the Yale Center for Clinical Investigation Scholar Award, by the Whitehall Foundation, by the Charles H. Hood Foundation, Inc. (Boston, MA), by a pilot grant from the Modern Diet and Physiology Research Center (The John B. Pierce Laboratory), by a grant of the Foundation for Prader-Willi Research, and by the Reginald and Michiko Spector Award in Neuroscience. G.M.B.O. was supported by the HHMI Gilliam Fellowship. G.M.S. was supported by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES). We thank David Bruin for copyediting the manuscript. The authors declare no conflict of interest.

Author contributions

M.O.D and G.M.B.O. conceived the hypothesis, designed the study and wrote the manuscript. G.M.B.O. performed experiments. G.M.B.O. and G.M.S. analyzed and plotted the data. All authors read and edited the manuscript.

Methods

Experimental models and subject details

All preweaning mice used in the experiments were 6 to 12 days old from both sexes (see table

below for total number of recording used under each age specifying sex and genotype). Litters

were provided from 4 separate breeding pairs. Dams used were 2 to 6 months old. To generate

experimental pups, we used the following cross: $Magel2^{m+/p-}$ (Jax #009062) dams bred with

C57BL/6J (Jax #000664) males. Offspring from this cross were either $Magel2^{m+/p-}$ or wildtype

($Magel2^{m+/p+}$). All mice were kept in temperature- and humidity-controlled rooms, in a 12/12 hr.

light/dark cycle, with lights on from 7:00 AM–7:00 PM. Studies took place during the light cycle.

Food (Teklad 2018S, Envigo) and water were provided ad libitum. All procedures were

approved by IACUC (Yale University).

AGE	*Control* (F)	*Control* (M)	$Magel2^{m+/p-}$ (F)	$Magel2^{m+/p-}$ (M)
P06	9	7	10	10
P08	15	9	15	13
P10	10	13	11	10
P12	15	15	16	12

Behavior test

Pups from the same litter were placed individually in a soundproof chamber containing fresh

bedding material [36]. An UltraSoundGate Condenser Microphone CM 16 (Avisoft Bioacoustics,

Berlin, Germany) was placed 10 cm above the recording chamber and connected to the

UltraSoundGate 416 USGH device to record ultrasonic vocalizations. The recording sessions

lasted 20 minutes. Four to eight chambers were recorded simultaneously. After testing, mice

were placed back in their home cage with the dam. Pups were tested at postnatal days 6, 8, 10,

and 12. Because pups that were naïve for the test—only tested at one specific age—show

similar results as pups tested at multiple ages, we pooled all mice together for our analysis.

Vocalization analysis

Vocalization analysis

Ultrasonic vocalizations were automatically extracted from audio recordings using a custom-built

tool [33]. In brief, audio recordings were converted from the time-domain to the frequency-domain

using a 1024-point Fast Fourier Transform (FFT) through a 512-width hamming window with

50% overlap. Spectrograms were computed from the FFT and processed as images. Each pixel

in the spectrogram corresponded to the intensity of each time-frequency component. Next, we

applied a series of image-processing techniques (e.g., contrast enhancement, binarization,

median filter, and morphological operations) to obtain segmentation of candidate vocalizations.

A single spectrogram was generated for each candidate vocalization detected. Candidate

vocalizations were classified as noise or real vocalization using a local median noise filter. The

remaining vocalization candidates are further labeled under one of eleven call type

classifications [37] using a Convolutional Neural Network (CNN), or as noise. The CNN was

trained using a curated vocalization dataset, containing over 20,000 noise samples and 40,000

vocalization samples. Finally, the tool produces one spectrogram centralized on each

vocalization for visual inspection, and a table (*xlsx* format) containing spectro-temporal features

for each USV, such as time, duration, bandwidth, frequency, and intensity (minimum, mean, and

maximum) values.

Quantification and statistical analysis

Prism 8.0 or above was used to analyze data and plot figures. Shapiro-Wilk normality test was

used to assess normal distribution of the data. Then, data were analyzed using two-way

ANOVA or mixed-effects analysis. Sidak's multiple comparisons test was used to find post hoc

differences among groups and to calculate the 95% confidence intervals to report effect size. To

analyze differences in the use of harmonics, we used the non-parametric Mann-Whitney Test

with Bonferroni correction to find statistically different effects. Statistical data are provided in text

and in the figures. In the text, values are provided as mean ± SEM. $P < 0.05$ was considered

statistically significant and, when necessary and as described above, was corrected using

Bonferroni's method.

REFERENCES

1. Perez JD, Rubinstein ND, Dulac C. New Perspectives on Genomic Imprinting, an Essential and Multifaceted Mode of Epigenetic Control in the Developing and Adult Brain. *Annu Rev Neurosci*. 2016. doi:10.1146/annurev-neuro-061010-113708

2. Barlow DP, Bartolomei MS. Genomic imprinting in mammals. *Cold Spring Harb Perspect Biol*. 2014. doi:10.1101/cshperspect.a018382

3. Ferguson-Smith AC. Genomic imprinting: The emergence of an epigenetic paradigm. *Nat Rev Genet*. 2011. doi:10.1038/nrg3032

4. Morison IM, Ramsay JP, Spencer HG. A census of mammalian imprinting. *Trends Genet*. 2005. doi:10.1016/j.tig.2005.06.008

5. Cassidy SB, Driscoll DJ. Prader-Willi syndrome. *Eur J Hum Genet*. 2009. doi:10.1038/ejhg.2008.165

6. Bittel DC, Butler MG. Prader-Willi syndrome: Clinical genetics, cytogenetics and molecular biology. *Expert Rev Mol Med*. 2005. doi:10.1017/S1462399405009531

7. Miller JL, Lynn CH, Driscoll DC, et al. Nutritional phases in Prader-Willi syndrome. *Am J Med Genet Part A*. 2011;155(5):1040-1049. doi:10.1002/ajmg.a.33951

8. Koenig K, Klin A, Schultz R. Deficits in Social Attribution Ability in Prader?Willi Syndrome. *J Autism Dev Disord*. 2004;34(5):573-582. doi:10.1007/s10803-004-2551-z

9. Nicholls RD, Knoll JHM, Butler MG, Karam S, Lalande M. Genetic imprinting suggested by maternal heterodisomy in non-deletion Prader-Willi syndrome. *Nature*. 1989. doi:10.1038/342281a0

10. Bischof JM, Stewart CL, Wevrick R. Inactivation of the mouse Magel2 gene results in growth abnormalities similar to Prader-Willi syndrome. *Hum Mol Genet*. 2007. doi:10.1093/hmg/ddm225

11. Schaaf CP, Gonzalez-Garay ML, Xia F, et al. Truncating mutations of MAGEL2 cause Prader-Willi phenotypes and autism. *Nat Genet*. 2013;45(11):1405-1408.

doi:10.1038/ng.2776

12.   Kozlov S V., Bogenpohl JW, Howell MP, et al. The imprinted gene Magel2 regulates

normal circadian output. *Nat Genet*. 2007. doi:10.1038/ng2114

13.   Michelsson K, Michelsson O. Phonation in the newborn, infant cry. In: *International*

Journal of Pediatric Otorhinolaryngology. ; 1999. doi:10.1016/S0165-5876(99)00180-9

14.   Lee S. Expression and imprinting of MAGEL2 suggest a role in Prader-Willi syndrome

and the homologous murine imprinting phenotype. *Hum Mol Genet*. 2000.

doi:10.1093/hmg/9.12.1813

15.   Branchi I, Santucci D, Alleva E. Ultrasonic vocalisation emitted by infant rodents: A tool

for assessment of neurobehavioural development. In: *Behavioural Brain Research*. ;

2001. doi:10.1016/S0166-4328(01)00277-7

16.   Branchi I, Santucci D, Alleva E. Analysis of ultrasonic vocalizations emitted by infant

rodents. *Curr Protoc Toxicol*. 2006. doi:10.1002/0471140856.tx1312s30

17.   Ehret G. Infant rodent ultrasounds - A gate to the understanding of sound communication.

In: *Behavior Genetics*. ; 2005. doi:10.1007/s10519-004-0853-8

18.   Hahn ME, Lavooy MJ. A review of the methods of studies on infant ultrasound production

and maternal retrieval in small rodents. In: *Behavior Genetics*. ; 2005.

doi:10.1007/s10519-004-0854-7

19.   Noirot E. Ultrasounds and maternal behavior in small rodents. *Dev Psychobiol*. 1972.

doi:10.1002/dev.420050410

20.   Nyby J, Wysocki CJ, Whitney G, Dizinno G. Pheromonal regulation of male mouse

ultrasonic courtship (Mus musculus). *Anim Behav*. 1977. doi:10.1016/0003-

3472(77)90009-4

21.   Nyby J, Dizinno G, Whitney G. Sexual dimorphism in ultrasonic vocalizations of mice

(Mus musculus): Gonadal hormone regulation. *J Comp Physiol Psychol*. 1977.

doi:10.1037/h0077411

22.   Sales GD, Smith JC. Comparative studies of the ultrasonic calls of infant murid rodents.

Dev Psychobiol. 1978. doi:10.1002/dev.420110609

23.   Zippelius H-M, Schleidt WM. Ultraschall-Laute bei jungen M usen. *Naturwissenschaften*.

1956;43(21):502-502. doi:10.1007/BF00632534

24.   Carlier M, Roubertoux P, Cohen-Salmon C. Differences in patterns of pup care in Mus

musculus domesticus I-Comparisons between eleven inbred strains. *Behav Neural Biol*.

1982. doi:10.1016/S0163-1047(82)91213-4

25.   Cohen-Salmon C. Differences in patterns of pup care in Mus musculus domesticus. VIII

Effects of previous experience and parity in XLII inbred mice. *Physiol Behav*. 1987.

doi:10.1016/0031-9384(87)90204-6

26.   Noirot E. Changes in responsiveness to young in the adult mouse: The effect of external

stimuli. *J Comp Physiol Psychol*. 1964. doi:10.1037/h0042864

27.   Smith JC. Responses of adult mice to models of infant calls. *J Comp Physiol Psychol*.

1976. doi:10.1037/h0077287

28.   Hofer MA, Brunelli SA, Shair HN. Potentiation of isolation-induced vocalization by brief

exposure of rat pups to maternal cues. *Dev Psychobiol*. 1994;27(8):503-517.

doi:10.1002/dev.420270804

29.   Hernandez-Miranda LR, Ruffault P-L, Bouvier JC, et al. Genetic identification of a

hindbrain nucleus essential for innate vocalization. *Proc Natl Acad Sci U S A*.

2017;114(30):8095-8100. doi:10.1073/pnas.1702893114

30.   Nyby J, Dizinno GA, Whitney G. Social status and ultrasonic vocalizations of male mice.

Behav Biol. 1976. doi:10.1016/S0091-6773(76)92198-2

31.   Castellucci GA, Calbick D, McCormick D. The temporal organization of mouse ultrasonic

vocalizations. *PLoS One*. 2019. doi:10.1371/journal.pone.0199929

32.   Elwood RW, Keeling F. Temporal organization of ultrasonic vocalizations in infant mice.

Dev Psychobiol. 1982. doi:10.1002/dev.420150306

33.   Fonseca AHO, Santana GM, Bampi S, Dietrich MO. Analysis of ultrasonic vocalizations

418   from mice using computer vision and machine learning. *bioRxiv*. 2020.

419   doi:10.1101/2020.05.20.105023

420   34.   Riede T. Subglottal pressure, tracheal airflow, and intrinsic laryngeal muscle activity

421   during rat ultrasound vocalization. *J Neurophysiol*. 2011. doi:10.1152/jn.00478.2011

422   35.   Kamaludin AA, Smolarchuk C, Bischof JM, et al. Muscle dysfunction caused by loss of

423   Magel2 in a mouse model of Prader-Willi and Schaaf-Yang syndromes. *Hum Mol Genet*.

424   2016. doi:10.1093/hmg/ddw225

425   36.   Zimmer MR, Fonseca AHO, Iyilikci O, Pra RD, Dietrich MO. Functional Ontogeny of

426   Hypothalamic Agrp Neurons in Neonatal Mouse Behaviors. *Cell*. 2019.

427   doi:10.1016/j.cell.2019.04.026

428   37.   Grimsley JMS, Monaghan JJM, Wenstrup JJ. Development of social vocalizations in

429   mice. *PLoS One*. 2011. doi:10.1371/journal.pone.0017460

430   38.   Winslqw JT, Insel TR. *Endogenous Opioids: Do They Modulate the Rat Pup's Response

431   to Social Isolation?* Vol 105.; 1991.

432   39.   McNamara GI, Creeth HDJ, Harrison DJ, et al. Loss of offspring Peg3 reduces neonatal

433   ultrasonic vocalizations and increases maternal anxiety in wild-type mothers. *Hum Mol

434   Genet*. 2018. doi:10.1093/hmg/ddx412

435   40.   Jiang YH, Pan Y, Zhu L, et al. Altered ultrasonic vocalization and impaired learning and

436   memory in Angelman syndrome mouse model with a large maternal deletion from Ube3a

437   to Gabrb3. *PLoS One*. 2010. doi:10.1371/journal.pone.0012278

438   41.   Wilkins JF, Haig D. What good is genomic imprinting: The function of parent-specific

439   gene expression. *Nat Rev Genet*. 2003. doi:10.1038/nrg1062

Figure 1. *Magel2* deficiency affects the emission of ultrasonic vocalizations in mice. (**A**)
Schematic of the protocol used to record separation-induced USVs in mice (from P6 to P12); pups
are separated from the home nest in a new chamber equipped with an ultrasonic microphone and
recorded for 20 minutes. (**B**) Total number of USVs emitted by control (blue) and *Magel2* deficient
(purple) littermates at P6, P8, P10, and P12; right panel denotes the 95% confidence intervals as
a measure of effect size. (**C**) Similar to (B), but only considering female pups. (**D**) Similar to (B),
but only considering male pups. (**E**) Average intensity of the USVs measured in decibels; right
panel denotes the 95% confidence intervals as a measure of effect size. Bars represent mean
value with error bars representing SEM and round symbols representing individual values. When
plotting the effect sizes, squared symbols and black lines represent 95% confidence intervals
calculated as the different between *Magel2* deficient and control pups. *P* values are provided in
the figures as calculated using Sidak's multiple comparison test. The sample sizes for control and
Magel2 deficient pups are: P6, n = 16 and 20; P8, n = 24 and 28; P10, n = 23 and 20; and P12,
n = 30 and 28, respectively.

Figure 2. *Magel2* **deficient pups emit ultrasonic vocalizations of distinct spectro-temporal features.** (**A**) Illustration of a spectrogram with the spectro-temporal features measured for each USV. (**B**) Maximum frequency of the USVs emitted by control and *Magel2* deficient littermates at P6, P8, P10, and P12; right panel denotes the 95% confidence intervals as a measure of effect size. (**C**) Similar to (**B**) but plotting the minimum frequency of the USVs. (**D**) Similar to (**B**) but plotting the mean frequency of the USVs. (**E**) Similar to (**B**) but plotting the bandwidth of the USVs. (**F**) Similar to (**B**) but plotting the duration of the USVs. (**G**) Illustration of the spectrogram of a single USV with a harmonic component. (**H**) Ratio of harmonic across all USVs emitted by control and *Magel2* deficient littermates. Bars represent mean value with error bars representing SEM and round symbols representing individual values. When plotting the effect sizes, squared symbols and black lines represent 95% confidence intervals calculated as the different between *Magel2* deficient and control pups. In B-F, *P* values are provided in the figures as calculated using Sidak's multiple comparison test post hoc analysis from two-way ANOVA test. In H, *P* values are provided as calculated using Mann-Whitney test. The sample sizes for control and *Magel2* deficient pups are: P6, n = 16 and 20; P8, n = 24 and 28; P10, n = 23 and 20; and P12, n = 30 and 28, respectively.

Figure 3. *Magel2* deficient pups emit ultrasonic vocalizations with discrete changes in the distribution of syllable types. (**A**) Illustration of the convolutional neural network used to classify each USV into one of eleven syllable types based on their morphology in spectrograms. (**B**) Spectrograms representing each of the eleven syllable types. (**C**) Distribution of syllable types in P6 pups—control in blue and *Magel2* deficient in purple. Data are showed as fraction of the total number of USVs; right panel denotes the 95% confidence intervals as a measure of effect size. (**D**) Similar to (C), but for P8 pups. (**E**) Similar to (C), but for P10 pups. (**F**) Similar to (C), but for P12 pups. Bars represent mean value with error bars representing SEM and round symbols representing individual values. When plotting the effect sizes, squared symbols and black lines represent 95% confidence intervals calculated as the different between *Magel2* deficient and control pups. *P* values are provided in the figures as calculated using Sidak's multiple comparison test as a post hoc analysis after two-way ANOVA. The sample sizes for control and *Magel2* deficient pups are: P6, n = 16 and 20; P8, n = 24 and 28; P10, n = 23 and 20; and P12, n = 30 and 28, respectively.

489
Figure 4. Analysis of the vocal repertoire of pups across ages. (**A**) Illustration of the output
of the convolutional neural network, with a distribution of eleven probabilities for vocal
classification (one probability for each of the eleven syllable types, with the highest probability
defining the syllable type). Using diffusion maps, a dimensionality reduction technique, these
eleven dimensions are reduced to three dimensions in the Euclidian space. (**B**) Illustration of a
pairwise comparison of the vocal repertoire of pups using diffusion maps and 3D alignment of
the manifolds (see methods for more details). (**D**) Comparison of the pairwise distance matrix
between control pups at different ages using Cohen's Kappa coefficient. (**D-E**) Examples of two
pairwise comparisons with high and low alignment. (**F-H**) Similar to (**D-E**), but for *Magel2*
deficient pups. (**I-K**) Similar to (C-H), but comparing control and *Magel2* deficient pups across

ages. The sample sizes for control and *Magel2* deficient pups are: P6, n = 16 and 20; P8, n = 24
and 28; P10, n = 23 and 20; and P12, n = 30 and 28, respectively.