Deficiency of the paternally inherited gene *Magel2* alters the development of separation-

induced vocalization and maternal behavior in mice

Authors:

Gabriela M. Bosque Ortiz [1, 2], Gustavo M. Santana[1, 2], Marcelo O. Dietrich [1, 2, 3, 4 *]

[1] Laboratory of Physiology of Behavior, Department of Comparative Medicine, Yale School of

Medicine, New Haven, CT, USA.

[2] Interdepartmental Neuroscience Program, Biological and Biomedical Sciences Program,

Graduate School in Arts and Sciences, Yale University, New Haven, CT.

[3] Program in Integrative Cell Signaling and Neurobiology of Metabolism, Department of

Comparative Medicine, Yale School of Medicine, New Haven, CT, USA.

[4] Department of Neuroscience, Yale School of Medicine, New Haven, CT, USA.

* Correspondence to: marcelo.dietrich@yale.edu

Abstract

The behavior of offspring results from the combined expression of maternal and paternal genes. Genomic imprinting silences some genes in a parent-of-origin specific manner, a process that, among all animals, occurs only in mammals. How genomic imprinting affects the behavior of mammalian offspring, however, remains poorly understood. Here, we studied how the loss of the paternally inherited gene *Magel2* in mouse pups affects the emission of separation-induced ultrasonic vocalizations (USV). Using quantitative analysis of more than one hundred thousand USVs, we characterized the rate of vocalizations as well as their spectral features from postnatal days 6 to 12 (P6-P12), a critical phase of mouse development during which pups fully depend on the mother for survival. Our analyses show that *Magel2* deficient offspring emit separation-induced vocalizations at lower rates and with altered spectral features. We also show that dams display altered behavior towards their own *Magel2* deficient offspring. In a test to compare the retrieval of two pups, dams retrieve wildtype control pups first and faster than *Magel2* deficient offspring. These results suggest that the loss of *Magel2* impairs the expression of separation-induced vocalization in pups as well as maternal behavior, both of which support the pups' growth and development.

Keywords

Genomic imprinting; offspring-parent conflict; Prader Willi Syndrome; Autism spectrum disorders; behavior development.

Introduction

For the normal development, mammalian offspring need copies from both maternal and paternal genomes. Some genes, however, are expressed in a parent-of-origin specific manner. In other words, some genes are always expressed when inherited from the mother and some genes are always expressed when inherited from the father[1-3]. The process that regulates the expression of genes in a parent-of-origin specific manner is called genomic imprinting[4].

Genomic imprinting depends on epigenetic modifications of the genome. These modifications do not alter the sequence of the DNA but the chemical structure of the DNA, thereby leading to altered gene expression[1,4]. For example, an imprinted gene can be silenced in the maternal genome, and only the paternal allele will be expressed in the offspring (or vice-versa). Thus far, genomic imprinting has only been found in flowering plants and mammals[2,5-7].

The adaptive value of genomic imprinting remains a matter of intense theorization. A prevailing theory on the effects of genomic imprinting, known as the kinship theory[5-7], posits that maternal genes balance the energy investment of the mother between offspring survival and her own while paternal genes favor offspring survival alone. For example, the expression of paternal genes in the offspring would favor growth, while the expression of maternal genes would stunt growth[2,8-11]. The influence of imprinted genes is not restricted to growth, however. Imprinted genes are also primarily involved in brain development and in social behaviors[2,8-10].

Consider, for example, a series of imprinted genes in human chromosome 15 that are paternally inherited[5–9]. The deletion of these paternally inherited genes in chromosome 15 leads to neurodevelopmental disorders, such as Prader-Willi syndrome (PWS). PWS presents with hypotonia and poor feeding early in life, followed by hyperphagia, alteration in social behavior, and cognitive deficits. It should be noted that because PWS involves several genes in human

chromosome 15, it masks the relative contribution of single imprinted genes on the phenotype of

the offspring. Among the PWS-related genes, MAGEL2 is a candidate gene for some of the

clinical features of PWS. Humans with loss-of-function mutations in MAGEL2 present clinical

aspects of PWS and autism spectrum disorders[10–12], suggesting that this single paternally

inherited gene supports at least some of the developmental alterations found in PWS. In

agreement with the clinical features of MAGEL2 deficiency in humans, *Magel2* deficient mice

show impairments in growth and adult social behaviors[10,13,14]. Despite these previous studies, a

more systematic investigation of the effects of imprinted genes on behavior is necessary to

understand the adaptive value of these genetic modifications in mammals.

We reasoned that investigating the effects of imprinted genes on offspring behavior during the

early postnatal period—when the exchange of resources between the mother and the offspring is

most important—can help shed light on the adaptive value of genomic imprinting. Towards this

end, we studied how the loss of paternally inherited *Magel2* affects the vocal behavior of mouse

pups when separated from their dams[15–23], as separation-induced vocalizations signal the needs

of the pups to the dams[19,24–29]. In contrast to human babies, mouse pups vocalize in the ultrasonic

frequency range (45 - 120 kHz)[20,21,23,30], which humans cannot hear. In order to survey the vocal

behavior of mice, we recorded the emission of ultrasonic vocalizations (USVs) when pups were

separated from the home nest. We performed these studies at postnatal days 6, 8, 10, and 12,

since it is during this phase of mouse development that the peak expression of separation-induced

vocalizations typically occurs[15,31,32]. We then used VocalMat[33] to perform quantitative analysis of

mouse vocal behavior. Moreover, we employed a maternal retrieval assay to test the effects of

the loss of paternally inherited *Magel2* on maternal behavior towards the offspring.[12-17] Our

analysis shows that in mouse pups the deficiency of *Magel2* impairs the expression of separation-

induced vocalizations. This deficiency also reduces maternal retrieval behavior towards *Magel2*

deficient pups compared to non-deficient siblings.

Results

Early waning of vocal behavior in *Magel2*[m+/p−] deficient pups

To investigate the effects of paternally inherited *Magel2* on the vocal behavior of infant mice, we

crossed heterozygote males for *Magel2* deficiency with wildtype females. From this cross, we

generated *Magel2* deficient offspring (*Magel2*[m+/p−]) that carry the null allele from the father (p−)

and the imprinted allele from the mother (m+). This cross also generates wildtype littermates

(*Magel2*[m+/p+]), used as experimental controls. As previously reported[15], *Magel2*[m+/p−] pups display

lower body weight compared to controls (genotype: $F_{1, 181}$ = 19.71, P < 10^{-4}; age: $F_{3, 181}$ = 69.64,

P < 10^{-10}; genotype x age: $F_{3, 181}$ = 0.02, P = 0.99; two-way ANOVA; **Figure S1**). Post-hoc

analysis shows a significant difference in body weight between genotypes at P12 (control: 6.08

± 0.14 g; *Magel2*[m+/p−]: 5.51 ± 0.20 g; P = 0.04; Holm-Šídák test; **Figure S1**).

We recorded the emission of USVs during 20 minutes of separation from the home nest at

different postnatal ages (P6, P8, P10, and P12; **Figure 1A**). First, we analyzed the total number

of USVs emitted during the period of separation using two-way ANOVA. We found a significant

effect of genotype, age, and interaction between genotype and age (genotype: $F_{1, 181}$ = 20.61, P

< 10^{-4}; age: $F_{3, 181}$ = 11.80, P < 10^{-6}; genotype x age: $F_{3, 181}$ = 3.90, P = 0.01; **Figure 1B**). Post-

hoc analysis (Holm-Šídák test) shows that the total number of USVs is similar among groups at

P6 (control: 991 ± 139 USVs; *Magel2*[m+/p−]: 787 ± 130 USVs; P = 0.75), P10 (control: 503 ± 45

USVs; *Magel2*[m+/p−]: 466 ± 70 USVs; P = 0.98), and P12 (control: 594 ± 71 USVs; *Magel2*[m+/p−]:

345 ± 46 USVs; P = 0.06) (**Figure 1B**). Compared to controls, however, *Magel2*[m+/p−] pups show

a ≈53% reduction in the emission of USVs at P8 (control: 1045 ± 100 USVs; *Magel2*[m+/p−]: 495 ±

65 USVs; P < 10^{-4}). We also analyzed the data in 5-minute intervals and found similar effects of

Magel2 deficiency on the rate of vocalizations (**Figure S2**). Moreover, separating our analysis in

females and males show similar effects of genotype and age (**Figure 1C-D** and **Table S1**).

(Because we did not find differences in the rate of vocalization, we have pooled males and

females in all further analysis). In sum, our results thus far show age-specific reductions in the

emission of USVs in *Magel2*[m+/p−] mice, suggesting a non-sex specific effect for paternally

inherited *Magel2* on the vocal behavior of the offspring.

The emission of USVs occurs when the breathing musculature contracts, expelling air from the

lungs and propelling it through the larynx[12,18]. Since previous reports found that *Magel2*[m+/p−]

mice display hypotonia[19], we considered the hypothesis that the low rate of USV emission in

Magel2 deficient pups is due to a lower capacity to expel air from the lungs. To rule out this

hypothesis, we measured the intensity (or volume, in decibels) of the USVs. Since the intensity

of the USVs relates to the pressure by which the air is expelled through the larynx[18], a lower

intensity is expected in cases of hypotonia. This analysis shows that the intensity of the emitted

USVs between *Magel2*[m+/p−] mice and controls is similar in all ages tested (genotype: $F_{1, 181}$ =

2.67, P = 0.10; age: $F_{3, 181}$ = 2.33, P = 0.08; genotype x age: $F_{3, 181}$ = 0.78, P = 0.51; two-way

ANOVA; **Figure 1E**). Thus, hypotonia does not seem to be a factor of primary significance for

the lower rate of USV emission in *Magel2*[m+/p−] mice[16].

***Magel2*[m+/p−] mice emit vocalizations with distinct spectral features**

In addition to the rate of separation-induced vocalizations, the spectral features of the USVs

also correlate with altered maternal care[20]. To test the extent to which *Magel2* deficiency affects

the spectral features of USVs across ages, we used two-way ANOVA to analyze the frequency

characteristics (pitch) and duration (**Figure 2A**) of USVs[21]. We found significant effects of

genotype and age for maximal frequency (genotype: $F_{1, 181}$ = 20.51, $P < 10^{-4}$; age: $F_{3, 181}$ = 3.23,

P = 0.02; genotype x age: $F_{3, 181}$ = 6.18, $P < 10^{-3}$; **Figure 2B**), bandwidth (genotype: $F_{1, 181}$ =

9.91, $P < 10^{-2}$; age: $F_{3, 181}$ = 6.97, $P < 10^{-3}$; genotype x age: $F_{3, 181}$ = 5.92, $P < 10^{-3}$; **Figure 2E**),

and duration (genotype: $F_{1, 181}$ = 4.08, P = 0.04; age: $F_{3, 181}$ = 6.35, $P < 10^{-3}$; genotype x age: $F_{3,}$

$_{181}$ = 2.05, P = 0.11; **Figure 2F**). In addition, we found a significant effect of genotype for mean

frequency (genotype: $F_{1, 181}$ = 4.86, P = 0.02; age: $F_{3, 181}$ = 0.24, P = 0.87; genotype x age: $F_{3, 181}$

= 0.41, P = 0.74; **Figure 2D**), but not for minimal frequency (genotype: $F_{1, 181}$ = 0.47, P = 0.47;

age: $F_{3, 181}$ = 1.96, P = 0.12; genotype x age: $F_{3, 181}$ = 0.48, P = 0.69; **Figure 2C**). We then used

post-hoc analysis (Holm-Šídák test) and found that at P8—but not at P6, P10, or P12—

Magel2$^{m+/p-}$ mice show significant differences compared to control mice in maximal frequency,

bandwidth, and duration (see panels plotting the 95% confidence intervals in **Figure 2**).

In addition to the main frequency component, USVs can contain harmonics (**Figure 2G**). We

calculated the percentage of USVs with harmonics and found a significantly lower number in

Magel2$^{m+/p-}$ mice compared to controls at P8 (control: 9.0 ± 1.7 %; *Magel2*$^{m+/p-}$: 3.5 ± 0.8 %; U =

145, $P_{\text{2-tailed}}$ = 0.002, Mann-Whitney test; **Figure 2H**) but not at P6 (control: 4.7 ± 0.9 %;

Magel2$^{m+/p-}$: 3.8 ± 0.6 %; U = 142.5, $P_{\text{2-tailed}}$ = 0.93, Mann-Whitney test; **Figure 2H**), P10

(control: 5.8 ± 0.9 %; *Magel2*$^{m+/p-}$: 6.1 ± 1.2 %; U = 240, $P_{\text{2-tailed}}$ = 0.98, Mann-Whitney test;

Figure 2H), or P12 (control: 3.6 ± 0.7 %; *Magel2*$^{m+/p-}$: 5.0 ± 1.5 %; U = 408.5, $P_{\text{2-tailed}}$ = 0.98,

Mann-Whitney test; **Figure 2H**). In sum, these results suggest that the loss of paternally

inherited *Magel2* in mice causes discrete changes in the features of separation-induced

vocalizations that are most evident at postnatal day eight.

Discrete changes in the use of syllable types by *Magel2*$^{m+/p-}$ mice

Mouse pups emit USVs of distinct classes—i.e., syllable types. Thus, the emission of different

syllable types could explain the discrete changes in the spectro-temporal features of USVs in

Magel2$^{m+/p-}$ [21,22]. We used a validated software to automatically categorize each USV into one of

eleven syllable types based on the morphology of the main component of the vocalization

(**Figure 3A-B**) [21]. The output of the method was the probability for each USV to be of a certain

syllable type. The highest probability (P_1) defined the syllable type for a given USV (**Figure 3A**).

The percent use of each syllable type per recording was compared between genotypes at each

age using two-way ANOVA. Using this approach, we did not find any significant differences in

the distribution of syllable types emitted by control and $Magel2^{m+/p-}$ mice at P6 (genotype: $F_{1, 374}$

= 2.5 x 10^{-16}, P > 0.99; class: $F_{10, 374}$ = 93.25, P < 10^{-4}; genotype x class: $F_{10, 374}$ = 1.00, P = 0.44)

and P12 (genotype: $F_{1, 616}$ = 7.7 x 10^{-16}, P > 0.99; class: $F_{10, 616}$ = 278, P < 10^{-4}; genotype x class:

$F_{10, 616}$ = 0.81, P = 0.62; **Figure 3C, 3F**). At P8, however, $Magel2^{m+/p-}$ pups emit vocalizations of

different syllable types (genotype: $F_{1, 539}$ = 2.2 x 10^{-15}, P > 0.99; class: $F_{10, 539}$ = 167, P < 10^{-6};

genotype x class: $F_{10, 539}$ = 9.51, P < 10^{-6}; **Figure 2D**), which include: 55% less USVs of the type

chevron (control: 9.0 ± 1.6 %; $Magel2^{m+/p-}$: 4.0 ± 0.01 %; P = 0.02, Holm-Šídák test); 44% less

USVs of the type *step-up* (control: 14.3 ± 1.4 %; $Magel2^{m+/p-}$: 8.1 ± 1.3 %; P < 10^{-3}, Holm-Šídák

test); 36% more USVs of the type *flat* (control: 9.6 ± 0.9 %; $Magel2^{m+/p-}$: 15.2 ± 1.2 %; P =

0.007, Holm-Šídák test); and 33% more USVs of the type *short* compared to controls (control:

23.6 ± 1.8 %; $Magel2^{m+/p-}$: 35.5 ± 2.0 %; P < 10^{-4}, Holm-Šídák test; **Figure 3D**). At P10, we also

identified discrete differences in the emission of USVs of different syllable types between groups

(genotype: $F_{1, 462}$ = 8.1 x 10^{-17}, P > 0.99; class: $F_{10, 462}$ = 149, P < 10^{-4}; genotype x class: $F_{10, 462}$ =

2.11, P = 0.02; **Figure 2E**): $Magel2^{m+/p-}$ pups emit 19% less *down frequency modulation*

(control: 31.0 ± 2.1 %; $Magel2^{m+/p-}$: 25.0 ± 1.5 %; P = 0.02; Holm-Šídák test) and emit 53%

more *up frequency modulation* (control: 4.8 ± 0.8 %; $Magel2^{m+/p-}$: 10.3 ± 2.0 %; P = 0.04; Holm-

Šídák test; **Figure 3E**). (**Table S1** provides a detailed analysis of the spectro-temporal features

of each syllable type across all ages tested in controls and $Magel2^{m+/p-}$ pups).

In summary, we found that $Magel2^{m+/p-}$ mice at P8 use simpler vocalizations that fall under the

'flat' and 'short' classifications instead of multicomponent USVs. These findings are in line with

our previous results **(Figures 1-2)** demonstrating that the largest differences in vocal behavior

occur in eight-day-old $Magel2^{m+/p-}$ pups.

Altered vocal repertoire of *Magel2*^m+/p− mice

As stated above, the vocal analysis pipeline outputs the probability for each USV to be classified

as each of the eleven syllable types (P_1, P_2, P_3, … P_{11}; **Figure 3A-B**). This distribution of

probabilities allows the qualitative and quantitative comparison of the vocal classification among

groups[21]. By considering the distribution of probabilities to classify each USV, it is possible to

estimate how similar the vocal repertoire of one group of mice is to another group. To compare

the vocal repertoire of mice across all ages studied, we used diffusion maps—a dimensionality

reduction technique that decreases the number of dimensions of the probability distribution from

eleven classes to three dimensions in a Euclidean space (**Figure 4A**)[21]. Using pairwise

comparisons (**Figure 4B**), we estimated the similarity between the vocal repertoire of mice of

different ages and genotypes. Using this method, we found that control pups at P6 and P8

(Cohen's coefficient: κ = 0.99) and control pups at P10 and P12 (Cohen's coefficient: κ = 0.95)

display vocal repertoires that are similar to each other (**Figure 4C-D**). These two age groups

(P6-P8 and P10-P12), however, present lower pairwise similarities when compared to each

other with κ ranging from 0.67 to 0.77 (**Figure 4E**). These results suggest that the vocal

repertoire of control pups undergoes significant changes between P8 and P10.

Next, we analyzed the same transitions in the vocal repertoire of *Magel2*^m+/p− pups. The

comparison between the vocal repertoire of *Magel2*^m+/p− pups at P6 and P8 show lower pairwise

similarity (κ = 0.80) compared to control pups (**Figure 4F**). In contrast to littermate controls,

Magel2^m+/p− pups at P8 show a higher pairwise similarity with P10 (κ = 0.84) and P12 (κ = 0.99)

pups (**Figure 4F-H)**. Finally, we directly compared the vocal repertoire of *Magel2*^m+/p− and

control pups. *Magel2*^m+/p− pups, at P6, show high pairwise similarity when compared to controls

at P6 (κ = 1.00) and P8 (κ = 1.00), but not at P10 (κ = 0.80) and P12 (κ = 0.72). At P8,

Magel2^m+/p− pups show relatively low pairwise similarities with control pups at P6 (κ = 0.76) and

P8 (κ = 0.78) but show high similarities with control pups at P10 (κ = 1.00) and P12 (κ = 0.95).

At P10, *Magel2*[m+/p−] pups show relatively lower pairwise similarities with control pups at P6 (κ =

0.67) and P8 (κ = 0.70) than at P10 (κ = 0.82) and P12 (κ = 0.81). This pattern is more evident

in P12 Magel2[m+/p−] pups, which show lower pairwise similarities with control pups at P6 (κ =

0.74) and P8 (κ = 0.75) than at P10 (κ = 0.99) and P12 (κ = 0.97).

Altogether, these analyses suggest a different dynamic for the ontogeny of the vocal repertoire

of *Magel2*[m+/p−] compared to control pups—with *Magel2*[m+/p−] pups at a younger age (i.e., P8)

resembling control pups at an older age (i.e., P10-P12). Thus, the period of development

between P8 and P10 seems to mark an important period for the effect of the maternally

imprinted gene, *Magel2*, on the vocal behavior of the offspring.

Dams prioritize their control offspring compared to *Magel2*[m+/p−] in a retrieval test

The emission of vocalization by infants draws caregiver's attention and care[12,13,15,17,22,23] while

genetically mute mouse pups are neglected by their dams[12]. Together, these observations led

us to speculate that the deficiency of *Magel2* in pups, which alters vocal behavior, could lead to

altered maternal behavior. In order to test this idea, we used a behavior assay to quantify

maternal behavior towards their own control and *Magel2*[m+/p−] offspring. In this assay, dams are

first placed in the middle of a three-chamber apparatus that contains their home nest in the

middle chamber (**Figure 5A**). After a period of acclimation, in the next stage, one pup of each

genotype is placed at the opposite ends of the apparatus. The time (latency) to retrieve each

pup back to the nest is then recorded (**Figure 5A**).

We initially evaluated whether dams show a preference to retrieve pups from one of the

genotypes. This analysis shows that control pups were more likely to be retrieved first (11/15

trials; 73.3%) compared to *Magel2*[m+/p−] pups (4/15 trials; 26.7%; **Figure 5B and Table S2**).

Accordingly, on average, the latency to retrieve control pups was less than half of the latency to

retrieve *Magel2*$^{m+/p-}$ pups (control: 51.9 ± 12.9 s; *Magel2*$^{m+/p-}$: 104.6 ± 19.1 s; $W = 82$, $P_{2\text{-tail}} =$

0.02, Wilcoxon test; **Figure 5C**). Because eleven out of fifteen *Magel2*$^{m+/p-}$ pups were retrieved

second, we also ran experiments placing wildtype control pups in both ends of the apparatus to

compare the latency to retrieve the second pup in each of the experiments. Using this

comparison, dams took twice as long to retrieve *Magel2*$^{m+/p-}$ pups compared to control pups

(control: 64.57 ± 12.3 s; *Magel2*$^{m+/p-}$: 126.3 ± 22.23 s; $W = 22$, $P_{2\text{-tail}} = 0.08$, Mann-Whitney test;

Figure 5D). In summary, the deficiency of Magel2 in offspring alters not only offspring behavior

but also the behavior of the mother towards their own offspring.

Discussion

In this study, we recorded and analyzed vocalizations from *Magel2* deficient pups and their wildtype littermates at postnatal days 6, 8, 10 and 12. Using custom-built software to automatically quantify vocalizations[21], we counted the number of vocalizations and measured the spectro-temporal features of each vocalization, including its intensity, duration, bandwidth, mean frequency, maximum frequency, minimum frequency, and use of harmonic components. We also assigned a syllable type for each vocalization based on its morphological features in the time-frequency plane. We used further quantitative methods to analyze the vocal repertoire of mice across groups and ages. These methods shed light on discrete changes in the development of separation-induced vocalizations in *Magel2* deficient mice.

With regard to the vocal behavior of wildtype mice, our results demonstrate that the emission of separation-induced USVs gradually decreases from P6-P8 to P10-P12. This result agrees with previous studies, which show an inverse-U shape profile for separation-induced vocalizations in mice during the first two weeks of life [15,29,31,32,38]. Based on the large number of vocalizations we analyzed, we also found that wildtype mice use simpler vocalizations at older ages (P10-P12) compared to younger ages (P6-P8). These findings in wildtype mice provide the basis for comparisons with *Magel2* deficient pups.

Magel2 deficient pups show different dynamics for separation-induced vocalizations. At P6, these pups vocalize comparably to wildtype littermates, but at P8, their vocal number and features resemble wildtype littermates that are older (P10-P12). An explanation for these results is that *Magel2* deficient pups are less responsive to certain social cues. In socially isolated pups, therefore, the deprivation of these cues would not induce the behavior to the same degree as in wildtype pups. An alternative explanation for these results is that for *Magel2* deficient

pups, vocal behavior does not have the same fitness value compared to wildtype pups. In the

latter case, the behavior begins to change at younger ages due to the lack of reinforcement

(maternal care). While it is difficult to prove these interpretations experimentally, the delayed

latency for dams to retrieve *Magel2* deficient pups compared to wildtype littermates supports the

idea that the deficiency of this imprinted gene impairs the fitness of the offspring. Moreover, the

fact that *Magel2* deficient pups have lower body weight during early development further

suggests a decrease in the fitness of these animals. Whether the change in vocal behavior and

the decrease in body weight and maternal behavior are causally related warrants further

investigation.

Interestingly, the deficiency of *Magel2* is not the only example of an imprinted gene affecting the

emission of USVs by mouse pups. Consider, for example, previous studies of other imprinted

genes. Deletion of the paternally inherited imprinted gene *Peg3* lowers vocal rate in mouse pups

[39]. Conversely, deletion of the maternally inherited genes *Gabrb3* and *Ube3a* increases vocal

rate[9]. Moreover, duplication of the paternal imprint loci on chromosome 15 increases the

separation-induced vocalization of mouse pups[24]. Based on these findings, it is tempting to

speculate that imprinted genes inherited from the father increase vocal rate while imprinted genes

inherited from the mother decrease vocal rate. To test this generalization more formally, future

studies will need to test the effect of *all* imprinted genes on the vocal behavior of the offspring.

P8 *Magel2*$^{m+/p-}$ vocalize at a low rate, mainly emitting short and flat vocals similar to those of

older pups. Moreover, when we estimate the similarity between the vocal repertoire of mutant

and wildtype pups across ages tested, we find that P8 *Magel2*$^{m+/p-}$ USVs are most similar to

those produced by wildtype P10 and P12 pups. Interestingly, reduction in USV rate is

associated to ASD, which is highly comorbid with MAGEL2$^{m+/p-}$ patients[25]. Further studies are

necessary to describe whether deviation in USV emission for *Magel2* deficient mice represents

some kind of curve shift in the behavior. For example, it could be that the critical phase in which

pups use separation-induced vocalization to gain maternal attention is shortened in offspring

deficient for *Magel2.* Thus, our findings could suggest that alteration in the emission of USVs is

part of a social behavior deficit in *Magel2*$^{m+/p-}$ pups that continues into adulthood[26].

In broader terms, we posit that our findings support the theory that genomic imprinting evolved

to balance the cost of the phenotype for the offspring and for the mother, as well as to balance

the best interests of mothers and fathers in altering offspring's phenotype[3,29]. In the case of

paternally inherited genes, the expression of these genes increases vocal behavior, thus,

increasing maternal care and favoring the use of maternal resources, which is in the best

interests of the father. Conversely, the loss of these paternally inherited genes decreases vocal

behavior and, consequently, the demand for maternal care, thus, conserving maternal

resources, which is in the best interests of the mother[2,5-7,27]. This theoretical framework is

supported by our findings in a test that measures the latency to retrieve isolated pups back to

the home nest, which demonstrate impaired maternal care towards Magel2 deficient pups

compared to wildtype littermates[12,13,15,17,28]. As alluded above, future studies should test this

theoretical view more directly by systematically investigating the effect of imprinted genes on the

behavior of offspring and on the behavior of mothers towards their offspring. These efforts will

help elucidate behavior phenotypes that occur in neurodevelopmental disorders as well as

expand our understanding of the evolutionary adaptation of genomic imprinting.

Acknowledgements

We thank the Dietrich Lab members for critical feedback on the project and the manuscript. Special thanks to: Delva Leão for her contribution to the data analysis; Jeremy Bober for helping with animal maintenance and care; as well as Antonio Fonseca who developed VocalMat software used for the analysis of vocal recordings. We thank David Gillich, Susan Andranovich, Valeria Krizsan, and Vickie Clark for administrative support. M.O.D. was supported by the National Institute of Mental Health of the National Institutes of Health under Award Number (R01MH125008) and by a grant of the Foundation for Prader-Willi Research. G.M.B.O. was supported by the HHMI Gilliam Fellowship. G.M.S. was supported by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES). We thank David Bruin for copyediting the manuscript. The authors declare no conflict of interest.

Author contributions

M.O.D and G.M.B.O. conceived the hypothesis, designed the study and wrote the manuscript. G.M.B.O. performed experiments. G.M.B.O. and G.M.S. analyzed and plotted the data. All authors read and edited the manuscript.

Material and Methods

Experimental models and subject details

All preweaning mice used in the experiments were 6 to 12 days old from both sexes (see table below for total number of recording used under each age specifying sex and genotype). Litters were provided from 9 separate breeding pairs. Separate litters were used for isolation and maternal retrieval tests. Details of litters tested found in **Table S1 and S2**. Dams used were 2 to 6 months old. To generate experimental pups, we used the following cross: $Magel2^{m+/p-}$ (Jax #009062) dams bred with C57BL/6J (Jax #000664) males. Offspring from this cross were either $Magel2^{m+/p-}$ or wildtype ($Magel2^{m+/p+}$). All mice were kept in temperature- and humidity-controlled rooms, in a 12/12 hr. light/dark cycle, with lights on from 7:00 AM–7:00 PM. Studies took place during the light cycle. Food (Teklad 2018S, Envigo) and water were provided ad libitum. All procedures were approved by IACUC (Yale University).

Behavior test

Pups from the same litter were placed individually in a soundproof chamber containing fresh bedding material [19]. An UltraSoundGate Condenser Microphone CM 16 (Avisoft Bioacoustics, Berlin, Germany) was placed 10 cm above the recording chamber and connected to the UltraSoundGate 416 USGH device to record ultrasonic vocalizations. The recording sessions lasted 20 minutes. Four to eight chambers were recorded simultaneously. After testing, mice were placed back in their home cage with the dam. Pups were tested at postnatal days 6, 8, 10, and 12. Because pups that were naïve for the test—only tested at one specific age—show similar results as pups tested at multiple ages, we pooled all mice together for our analysis. Moreover, literature in the field have reported that repeated isolation testing does not affect infant vocal behavior[29,30]. Details of individual behavior measurements are found in **Table S1**.

Vocalization analysis

Ultrasonic vocalizations were automatically extracted from audio recordings using a custom-built

tool [14]. In brief, audio recordings were converted from the time-domain to the frequency-domain

using a 1024-point Fast Fourier Transform (FFT) through a 512-width hamming window with

50% overlap. Spectrograms were computed from the FFT and processed as images. Each pixel

in the spectrogram corresponded to the intensity of each time-frequency component. Next, we

applied a series of image-processing techniques (e.g., contrast enhancement, binarization,

median filter, and morphological operations) to obtain segmentation of candidate vocalizations.

A single spectrogram was generated for each candidate vocalization detected. Candidate

vocalizations were classified as noise or real vocalization using a local median noise filter. The

remaining vocalization candidates are further labeled under one of eleven call type

classifications [20] using a Convolutional Neural Network (CNN), or as noise. The CNN was

trained using a curated vocalization dataset, containing over 20,000 noise samples and 40,000

vocalization samples. Finally, the tool produces one spectrogram centralized on each

vocalization for visual inspection, and a table (*xlsx* format) containing spectro-temporal features

for each USV, such as time, duration, bandwidth, frequency, and intensity (minimum, mean, and

maximum) values.

Maternal Retrieval Test

The maternal preference test was performed in a three-chamber apparatus (65 x 42 x 23 cm)

and comprised of three stages: Stage 1 – acclimation: the dam was allowed to explore the

apparatus without the presence of pups for five minutes. In the middle of the apparatus, home

cage nesting under an infrared igloo. Stage 2 – exploration: two P8 mice were placed on each

side of the apparatus and the dam was allowed to explore the pups for five minutes. Stage 3 -

preference: dam was recorded while she retrieved pups back to her home cage nesting. Groups

were randomly alternated between both sides to avoid preference for one side of the chamber.

Latency to retrieve each pup was timed from videos taken of the retrieval tests. Summary of

results found in **Table S2**.

Quantification and statistical analysis

Prism 8.0 or above was used to analyze data and plot figures. Shapiro-Wilk normality test was

used to assess normal distribution of the data. To analyze differences in the use of harmonics,

we used the non-parametric Mann-Whitney Test with Bonferroni correction to find statistically

different effects. Comparison of diffusion maps were analyzed through pairwise comparison

using Cohen's kappa coefficient. Maternal retrieval latency for retrieval of each pup from the

start of test used Wilcoxon test. Analysis of second retrieval latency from first retrieval was

calculated using Welch's t test. The rest of the data was analyzed using two-way ANOVA or

mixed-effects analysis. Sidak's multiple comparisons test was used to find post hoc differences

among groups and to calculate the 95% confidence intervals to report effect size. In the text,

values are provided as mean ± SEM. $P < 0.05$ was considered statistically significant and, when

necessary and as described above, was corrected using Bonferroni's method. Statistical data

are provided in text and in the figures.

Figure 1. *Magel2* deficiency affects the emission of ultrasonic vocalizations in mice. (**A**)

Schematic of the protocol used to record separation-induced USVs in mice (from P6 to P12); pups

are separated from the home nest in a new chamber equipped with an ultrasonic microphone and

recorded for 20 minutes. (**B**) Total number of USVs emitted by control (blue) and *Magel2* deficient

(purple) littermates at P6, P8, P10, and P12; right panel denotes the 95% confidence intervals as

a measure of effect size. (**C**) Similar to (B), but only considering female pups. (**D**) Similar to (B),

but only considering male pups. (**E**) Average intensity of the USVs measured in decibels; right

panel denotes the 95% confidence intervals as a measure of effect size. Bars represent mean

value with error bars representing SEM and round symbols representing individual values. When

plotting the effect sizes, squared symbols and black lines represent 95% confidence intervals

calculated as the different between *Magel2* deficient and control pups. *P* values are provided in

the figures as calculated using Sidak's multiple comparison test. The sample sizes for control and

Magel2 deficient pups are: P6, n = 16 and 20; P8, n = 23 and 28; P10, n = 23 and 20; and P12,

n = 30 and 28, respectively.

Figure 2. *Magel2* deficient pups emit ultrasonic vocalizations of distinct spectro-temporal

features. (A) Illustration of a spectrogram with the spectro-temporal features measured for each

USV. **(B)** Maximum frequency of the USVs emitted by control and *Magel2* deficient littermates

at P6, P8, P10, and P12; right panel denotes the 95% confidence intervals as a measure of

effect size. **(C)** Similar to (B) but plotting the minimum frequency of the USVs. **(D)** Similar to (B)

but plotting the mean frequency of the USVs. **(E)** Similar to (B) but plotting the bandwidth of the

USVs. **(F)** Similar to (B) but plotting the duration of the USVs. **(G)** Illustration of the spectrogram

of a single USV with a harmonic component. **(H)** Ratio of harmonic across all USVs emitted by

control and *Magel2* deficient littermates. Bars represent mean value with error bars representing

SEM and round symbols representing individual values. When plotting the effect sizes, squared

symbols and black lines represent 95% confidence intervals calculated as the different between

Magel2 deficient and control pups. In B-F, *P* values are provided in the figures as calculated

using Sidak's multiple comparison test post hoc analysis from two-way ANOVA test. In H, *P*

values are provided as calculated using Mann-Whitney test. The sample sizes for control and

Magel2 deficient pups are: P6, n = 16 and 20; P8, n = 23 and 28; P10, n = 23 and 20; and P12,

n = 30 and 28, respectively.

Figure 3. *Magel2* deficient pups emit ultrasonic vocalizations with discrete changes in the distribution of syllable types. (**A**) Illustration of the convolutional neural network used to classify each USV into one of eleven syllable types based on their morphology in spectrograms. (**B**) Spectrograms representing each of the eleven syllable types. (**C**) Distribution of syllable types in P6 pups—control in blue and *Magel2* deficient in purple. Data are showed as fraction of the total number of USVs; right panel denotes the 95% confidence intervals as a measure of effect size. (**D**) Similar to (C), but for P8 pups. (**E**) Similar to (C), but for P10 pups. (**F**) Similar to (C), but for P12 pups. Bars represent mean value with error bars representing SEM and round symbols representing individual values. When plotting the effect sizes, squared symbols and black lines represent 95% confidence intervals calculated as the different between *Magel2* deficient and control pups. *P* values are provided in the figures as calculated using Sidak's multiple comparison test as a post hoc analysis after two-way ANOVA. The sample sizes for control and *Magel2* deficient pups are: P6, n = 16 and 20; P8, n = 23 and 28; P10, n = 23 and 20; and P12, n = 30 and 28, respectively.

Figure 4. Analysis of the vocal repertoire of pups across ages. (**A**) Illustration of the output

of the convolutional neural network, with a distribution of eleven probabilities for vocal

classification (one probability for each of the eleven syllable types, with the highest probability

defining the syllable type). Using diffusion maps, a dimensionality reduction technique, these

eleven dimensions are reduced to three dimensions in the Euclidian space. (**B**) Illustration of a

pairwise comparison of the vocal repertoire of pups using diffusion maps and 3D alignment of

the manifolds (see methods for more details). (**D**) Comparison of the pairwise distance matrix

between control pups at different ages using Cohen's Kappa coefficient. (**D-E**) Examples of two

pairwise comparisons with high and low alignment. (**F-H**) Similar to (**D-E**), but for *Magel2*

deficient pups. (**I-K**) Similar to (C-H), but comparing control and *Magel2* deficient pups across

ages. The sample sizes for control and *Magel2* deficient pups are: P6, n = 16 and 20; P8, n = 23

and 28; P10, n = 23 and 20; and P12, n = 30 and 28, respectively.

Figure 5. Maternal retrieval behavior is biased towards wildtype versus *Magel2* deficient

pups. (**A**) Diagram of maternal retrieval protocol with P8 pups on opposite ends of the

apparatus and mother at nest in the middle compartment. (**B**) Pie chart showing percentages of

first retrieval of control and *Magel2* deficient pups from all trials. (**C**) Welch's t-test comparing

the retrieval latency of control and *Magel2* deficient pups. (**D**) Welch's t-test comparing the

latency for the retrieval of the second pup comparing *Magel2* deficient pups and wildtype

controls that were picked second in the wildtype versus wildtype trial. Significant *P* values are

shown in the graphs. The number *Magel2* deficient pups and controls trials are: P8, n = 15 and

n = 7.

References

1.   Barlow DP, Bartolomei MS. Genomic imprinting in mammals. *Cold Spring Harb Perspect*
Biol. 2014;6(2).

2.   Perez JD, Rubinstein ND, Dulac C. New Perspectives on Genomic Imprinting, an
Essential and Multifaceted Mode of Epigenetic Control in the Developing and Adult Brain. *Annu*
Rev Neurosci. 2016;39:347-384.

3.   Ferguson-Smith AC. Genomic imprinting: the emergence of an epigenetic paradigm. *Nat*
Rev Genet. 2011;12(8):565-575.

4.   Morison IM, Ramsay JP, Spencer HG. A census of mammalian imprinting. *Trends*
Genet. 2005;21(8):457-465.

5.   Haig D. Coadaptation and conflict, misconception and muddle, in the evolution of
genomic imprinting. *Heredity (Edinb).* 2014;113(2):96-103.

6.   Patten MM, Ross L, Curley JP, Queller DC, Bonduriansky R, Wolf JB. The evolution of
genomic imprinting: theories, predictions and empirical tests. *Heredity (Edinb).* 2014;113(2):119-
128.

7.   Spencer HG, Clark AG. Non-conflict theories for the evolution of genomic imprinting.
Heredity (Edinb). 2014;113(2):112-118.

8.   Li L, Keverne EB, Aparicio SA, Ishino F, Barton SC, Surani MA. Regulation of maternal
behavior and offspring growth by paternally expressed Peg3. *Science.* 1999;284(5412):330-
333.

9.   Jiang YH, Pan Y, Zhu L, et al. Altered ultrasonic vocalization and impaired learning and
memory in Angelman syndrome mouse model with a large maternal deletion from Ube3a to
Gabrb3. *PLoS One.* 2010;5(8):e12278.

10.   Hernandez A, Martinez ME, Fiering S, Galton VA, St Germain D. Type 3 deiodinase is
critical for the maturation and function of the thyroid axis. *J Clin Invest.* 2006;116(2):476-484.

11.   Martinez ME, Charalambous M, Saferali A, et al. Genomic imprinting variations in the
mouse type 3 deiodinase gene between tissues and brain regions. *Mol Endocrinol.*
2014;28(11):1875-1886.

12.   Hernandez-Miranda LR, Ruffault PL, Bouvier JC, et al. Genetic identification of a
hindbrain nucleus essential for innate vocalization. *Proc Natl Acad Sci U S A.*
2017;114(30):8095-8100.

13.   Zimmerberg B, Rosenthal AJ, Stark AC. Neonatal social isolation alters both maternal
and pup behaviors in rats. *Dev Psychobiol.* 2003;42(1):52-63.

14.   Uematsu A, Kikusui T, Kihara T, et al. Maternal approaches to pup ultrasonic
vocalizations produced by a nanocrystalline silicon thermo-acoustic emitter. *Brain Res.*
2007;1163:91-99.

15.   Hahn ME, Lavooy MJ. A review of the methods of studies on infant ultrasound
production and maternal retrieval in small rodents. *Behav Genet.* 2005;35(1):31-52.

16.   Asaba A, Kato M, Koshida N, Kikusui T. Determining Ultrasonic Vocalization
Preferences in Mice using a Two-choice Playback Test. *J Vis Exp.* 2015(103).

17.   Noirot E. Ultrasounds and maternal behavior in small rodents. *Dev Psychobiol.*
1972;5(4):371-387.

18.   Riede T. Subglottal pressure, tracheal airflow, and intrinsic laryngeal muscle activity
during rat ultrasound vocalization. *J Neurophysiol.* 2011;106(5):2580-2592.

19.   Kamaludin AA, Smolarchuk C, Bischof JM, et al. Muscle dysfunction caused by loss of
Magel2 in a mouse model of Prader-Willi and Schaaf-Yang syndromes. *Hum Mol Genet.*
2016;25(17):3798-3809.

20.   Zimmer MR, Fonseca AHO, Iyilikci O, Pra RD, Dietrich MO. Functional Ontogeny of
Hypothalamic Agrp Neurons in Neonatal Mouse Behaviors. *Cell.* 2019;178(1):44-59 e47.

21.   Fonseca AH, Santana GM, Bosque Ortiz GM, Bampi S, Dietrich MO. Analysis of
ultrasonic vocalizations from mice using computer vision and machine learning. *Elife.* 2021;10.

22.   Grimsley JM, Monaghan JJ, Wenstrup JJ. Development of social vocalizations in mice.
PloS one. 2011;6(3):e17460.

23.   Branchi I, Santucci D, Alleva E. Ultrasonic vocalisation emitted by infant rodents: a tool
for assessment of neurobehavioural development. *Behav Brain Res.* 2001;125(1-2):49-56.

24.   Nakatani J, Tamada K, Hatanaka F, et al. Abnormal behavior in a chromosome-
engineered mouse model for human 15q11-13 duplication seen in autism. *Cell.*
2009;137(7):1235-1246.

25.   Takahashi T, Okabe S, Broin PO, et al. Structure and function of neonatal social
communication in a genetic mouse model of autism. *Mol Psychiatry.* 2016;21(9):1208-1214.

26.   Fountain MD, Tao H, Chen CA, Yin J, Schaaf CP. Magel2 knockout mice manifest
altered social phenotypes and a deficit in preference for social novelty. *Genes Brain Behav.*
2017;16(6):592-600.

27.   Wilkins JF, Haig D. What good is genomic imprinting: the function of parent-specific
gene expression. *Nat Rev Genet.* 2003;4(5):359-368.

28.   Farrell WJ, Alberts JR. Maternal responsiveness to infant Norway rat (Rattus
norvegicus) ultrasonic vocalizations during the maternal behavior cycle and after steroid and
experiential induction regimens. *J Comp Psychol.* 2002;116(3):286-296.

29.   Barnes TD, Rieger MA, Dougherty JD, Holy TE. Group and Individual Variability in
Mouse Pup Isolation Calls Recorded on the Same Day Show Stability. *Front Behav Neurosci.*
2017;11:243.

30.   Verjat A, Rodel HG, Feron C. Isolation calls in house mouse pups: Individual
consistency across time and situations. *Dev Psychobiol.* 2019;61(8):1135-1145.

Figure S1. Body weight of pups across tested ages. Individual body weights of control (blue)

and *Magel2* deficient (purple) littermates at P6, P8, P10, and P12. Line with error bars represent

mean and SEM. Round symbols representing individual values. *P* values are provided in the main

text. The sample sizes for control and *Magel2* deficient pups are: P6, n = 16 and 20; P8, n = 23

and 28; P10, n = 23 and 20; and P12, n = 30 and 28, respectively.

Figure S2. Vocal rate in 5-minute bins across ages. Average USV emission of control (blue)

and *Magel2* deficient (purple) littermates at (A) P6, (B) P8, (C) P10, and (D) P12. Connecting lines

trace USV means for each genotype across 5-minute bins from the 20-minute isolation recording.

Line with error bars represent mean and SEM of weights. Round symbols represent mean values.

2way ANOVA results for each age are found on the tables on the right of each graph. The sample

sizes for control and *Magel2* deficient pups are: P6, n = 16 and 20; P8, n = 23 and 28; P10, n =

23 and 20; and P12, n = 30 and 28, respectively.

Figure S3. Rate of USVs in naïve versus experienced mouse pups. Average USV emission

of pups tested repeatedly (purple) and naïve pups (blue). Bars represent USV means for each

genotype and condition. Line with error bars represent mean and SEM. Round symbols

represent rate for individual pups. Multiple Mann-Whitney test was used to compare rates. *P*

values are the following: P8 WT = 0.36; P8 MUT = 0.46; P10 WT = 0.46; P10 MUT = 0.58.