Communication

The Exon Junction Complex Core Represses Caner-specific Mature mRNA Re-splicing: A Potential Key Role in Terminating Splicing

Yuta Otani[1,2], Toshiki Kameyama[1,3] and Akila Mayeda[1,*]

[1] Division of Gene Expression Mechanism, Institute for Comprehensive Medical Science, Fujita Health University, Toyoake, Aichi 470-1192, Japan
[2] Laboratories of Discovery Research, Nippon Shinyaku Co., Ltd., Kyoto, Kyoto 601-8550, Japan
[3] Present address: Department Physiology, Fujita Health University, School of Medicine, Toyoake, Aichi 470-1192, Japan

* Correspondence: mayeda@fujita-hu.ac.jp (A.M.)

Abstract: Using the TSG101 pre-mRNA, we previously discovered cancer-specific re-splicing of mature mRNA that generates aberrant transcripts/proteins. The fact that mRNA is aberrantly re-spliced in various cancer cells implies there must be an important mechanism to prevent deleterious re-splicing on the spliced mRNA in normal cells. We thus postulated that the mRNA re-splicing is controlled by specific repressors and we searched for repressor candidates by siRNA-based screening for mRNA re-splicing activity. We found that knock-down of EIF4A3, which is a core component of the exon junction complex (EJC), significantly promoted mRNA re-splicing. Remarkably, we could recapitulate cancer-specific mRNA re-splicing in normal cells by knock-down of any of the core EJC proteins, EIF4A3, MAGOH or RBM8A (Y14), implicating the EJC core as the repressor of mRNA re-splicing often observed in cancer cells. We propose that the EJC core is a critical mRNA quality control factor to prevent over-splicing of mature mRNA.

Keywords: Pre-mRNA splicing; mRNA re-splicing; EJC; TSG101; EIF4A3; MAGOH; RBM8A (Y14)

1. Introduction

Although the basic mechanism of how pre-mRNA splicing is initiated and proceeded has been characterized in detail, it is less well known about how splicing is the terminated. *Cis*-acting splicing signals, such as 5′/3′ splice sites and branch site followed by poly-pyrimidine tract, are not highly conserved, and thus they are necessary but not sufficient to be utilized [reviewed in 1]. Because of this redundancy, there remain many splice site-like sequences in fully spliced mRNA. However these spurious splice sites are usually never used and the mature mRNA is safely exported to the cytoplasm for translation. These processes are definitely critical to maintain the integrity of coding mRNA. Whereas in the vertebrate multi-step splicing pathways, such as recursive splicing [2,3], intrasplicing [4], and nested splicing [5,6], it is evident that the initially spliced intermediates can serve to be spliced again, indicating that the splicing process is not terminated when certain conditions are met. One component in the termination process could be the DEAH-box family RNA helicases involved in the dissociation of spliceosome, such as DHX8 (also known as Prp22) and DHX15 (Prp43) [reviewed in 7]. However, the exact mechanism to control splicing termination is poorly understood. Here we address this thorny problem by taking advantage of an aberrant multi-step splicing, where a norally spliced mRNA can also be a substrate for further splicing often in cancer cells, termed cancer-specific mRNA re-splicing.

Aberrant splicing that occurs over very long distances, skipping many exons with flanking huge introns generating annotated truncated mRNAs, are often observed in cancer cells [reviewd in 8]; e.g., TSG101 and FHIT pre-mRNAs were often reported as typical cases. Using these two substrates, we discovered that the mRNA re-splicing event occurs on mature spliced mRNA, in which the normal splicing removes all authentic splice sites and bringing the weak re-splice sites into close proximity [9]. The re-spliced TSG101 mRNA, TSG101Δ154-1054 mRNA (Figure 1A), produces an aberrant protein product that upregulates the full-size TSG101 by interfering with the ubiquitin-dependent degradation of TSG101 protein [10]. Importantly, the overexpression of TSG101 protein enhances cell proliferation and promotes malignant tumor formation in nude mice [10,11]. TSG101 is essential functions for cell proliferation and survival, while its overexpression is closely associated with aggressive metastasis and progression in various cancers [Reviewed in 12]. It is thus critical to control this kind of aberrant mRNA re-splicing pathway in normal cells to prevent production of potentially deleterious protein products.

Here we postulated that the control of the aberrant mRNA re-splicing is promoted by a specific repressor in normal cells. To test this hypothesis, we performed siRNA screening of repressor candidates based on cancer-specific mRNA re-splicing activity of the TSG101 pre-mRNA. This approach identified EIF4A3, whose knock-down significantly stimulated re-splicing of the TSG101 mRNA. EIF4A3 is one of three essential core components of the exon junction complex (EJC) along with MAGOH and RBM8A (Y14). The EJC is deposited upstream of the spliced exon-exon junctions as a consequence of pre-mRNA splicing and the EJC together with the recruited peripheral factors play important roles in multiple post-splicing events, such as mRNA export, nonsense-mediated mRNA decay (NMD), and translation [reviewed in 13,14]. Here we demonstrate that EJC core per se, but not the EJC peripheral factors, is necessary to repress mRNA re-splicing in normal cells.

2. Results

2.1. Re-Splicing of mRNA Is Enhanced by EIF4A3 Knock-down

Previously, it was demonstrated that the induction and overexpression of the tumor suppressor gene *TP53* (*p53*) prevents cancer-specific aberrant splicing of TSG101 pre-mRNA [10,15], i.e., generation of the re-spliced TSG101Δ154-1054 mRNA [9]. Therefore, this mRNA re-splicing event is under the control of *TP53*, presumably indirectly through other regulators; i.e., upregulation of an mRNA re-splicing repressor and/or downregulation of an mRNA re-splicing activator. To identify such repressor or activator candidates, we activated TP53, using highly selective MDM2 inhibitor RG7388 [16] to repress aberrant re-splicing, and searched for up- or down-regulated genes on a microarray (Supplementary Tables S1). We confirmed that the activation of *TP53* by RG7388 indeed prevents re-splicing activity of TSG101 pre-mRNA (Supplementary Figure S1).

We selected 48 candidate proteins including hnRNP proteins, RNA helicases, a splicing-related kinase, and other RNA-binding proteins (Table S2) from the up- and down-regulated genes upon activation of TP53. Then we performed siRNA-mediated knock-down of each protein (Table S2 for the siRNA sequences) in the human breast cancer cell line MCF-7, in which mRNA re-splicing was previously demonstrated [9]. We found that the knocking down EIF4A3 markedly increased the aberrant re-spliced product of TSG101 mRNA (TSG101Δ154-1054) over twenty times more than in the control (Figure 1B). Our results indicate that EIF4A3 is a potent repressor of mRNA re-splicing. Since EIF4A3 expression was not up-regulated, but rather down-regulated, under the RG7388-mediated activation of TP53 (Table S1), EIF4A3 was evidently not a TP53-dependent repressor of mRNA re-splicing (see Discussion).

2.2. The EJC Core Factors Were Identified as Repressors of mRNA Re-Splicing

Remarkably, EIF4A3 protein, which appers to repress mRNA re-splicing, is one of the core components of the EJC that is deposited onto mature mRNA [reviewed in 13,14]. Next, we test whether EIF4A3 functions solely or EIF4A3 functions as a core component in the EJC. We thus knocked-down the other EJC core components, MAGOH, RBM8A and CASC3 (MLN51) using siRNAs in MCF-7 cells.

Like for EIF4A3, we found that knock-down of both MAGOH and RBM8A markedly activated the aberrant re-splicing of TSG101 mRNA, whereas CASC3 knock-down had no effect (Figure 1C). Since EIF4A3, MAGOH and RBM8A are essential, but CASC3 is dispensable, for the assembly of a stable EJC core [17], it is reasonable to assume that an intact EJC core assembly is necessary to repress mRNA re-splicing. We confirmed the actual binding of the EJC core on TSG101 mRNA by RNA-immunoprecipitation assay using anti-RBM8A antibody (Figure S2). Together, we conclude that the formation of the EJC core *per se* is a substantial repressor of mRNA re-splicing.

2.3. The EJC Core Represses mRNA Re-Splicing in Normal Cells

If the EJC core alone is necessary to repress mRNA re-splicing in normal cells, the knock-down of each EJC core factor in normal cells should activate mRNA re-splicing as observed in cancer cells.

We tested this assumption using MCF-10A cells, which is a non-cancerous human mammary epithelial cell line. We could not detect re-spliced mRNA product in MCF-10A cells (Figure 2A), just as we had not in normal mammary epithelial cells [9]. Remarkably, mRNA re-spliced product was generated under efficient siRNA-mediated knock-down of each EJC core factor, EIF4A3, MAGOH or RBM8A (Figure 2B). We conclude that knockdown of a single EJC core factor is sufficient to activate mRNA re-splicing in normal cells.

2.4. Repression of mRNA Re-Splicing Is Not Due To EJC-Mediated NMD

The EJC contains peripheral factors responsible for nonsense-mediated mRNA decay (NMD) such as the UPF proteins (UPF1, 2, 3) [reviewed in 13,14]. As the re-spliced product, TSG101Δ154-1054 mRNA, contains a premature termination codon (PTC) downstream of re-spliced junction, it

could be a target of NMD [9]. Therefore, knock-down of EJC core factors could also dissociate UPF proteins so that it potentially represses NMD of re-spliced mRNA leading to its eventual increase.

Mature mRNA re-splicing, generating TSG101Δ154-1054 mRNA, is a cancer-specific event and it is barely detectable in normal cells [9; and references therein]. First, it is critical to determine whether the re-spliced mRNA is indeed not generated or it is the result of the degradation of the generated mRNA by NMD in normal cells. We thus performed siRNA-mediated knock-down of essential NMD factor UPF1 in normal MCF-10A cells. This knock-down effectively depleted UPF1 but it did not generate the re-spliced TSG101Δ154-1054 product (Figure 3A), confirming that the re-spliced TSG101Δ154-1054 product barely exists in normal cells, but that this is not due to NMD.

We therefore quantified mRNA re-splicing activity under the knock-down of UPF1 and EIF4A3 using HeLa cells (in which siRNA-mediated interference is more efficient than that in MCF-10A cells). The mRNA re-spliced product was markedly increased by the depletion of EIF4A3, but not at all by the depletion of UPF1 (Figure 3B, middle graph). In this assay, the level of GAS5 mRNA, a well-known endogenous NMD target [18,19], was significantly increased in both EIF4A3 and UPF1 depleted cells (Figure 3B, right graph), supporting that NMD is inhibited almost equally by knocking down of either EIF4A3 or UPF1. Together, we conclude that the observed increase of mRNA re-spliced product by the knock-down of EJC core factor is attributed to the intrinsic function of EJC as a repressor of mRNA re-splicing, but is not due to the repression of EJC-mediated NMD.

2.5. mRNA Re-splicing Does Not Depend on EJC-mediated mRNA Export or Splicing Regulation

As well as several NMD factors, the EJC also interacts peripherally with several mRNA export factors and splicing regulators [reviewed in 13,14]. It is conceivable that knock-down of EJC core factor could dissociate mRNA export factor such as NXF1 (TAP), which might lead to the nuclear retention of mature mRNA. Then the nuclear-retained mRNA could be a target of post-transcriptional re-splicing. It is also possible that the dissociation of other EJC-associated peripheral factors such as ACIN1 (ACINUS), PNN (Pinin), RNPS1, and SAP18, known as potential splicing regulators, could affect the efficiency of mRNA re-splicing [reviewed in 13,14].

We thus tested these EJC peripheral factors by siRNA-mediated knock-down in MCF-7 cells (Figure 4A). The mRNA re-splicing was not significantly activated (Figure 4B) as we observed in the depletion of EJC core factors (Figure 1B, C). The knock-down of other EJC-associated mRNA export factors, UAP56 and URH49, also had no significant effects on mRNA re-splicing activity (unpublished results). Our results clearly implicate that mRNA re-splicing event is not due to the extra splicing on the nuclear retained mRNA caused by inhibition of mRNA export. We conclude that the formed EJC core itself, but not the associated EJC-peripheral factors, is involved in the prevention of mRNA re-splicing.

2.6. Reduced RBM8A Protein Caused mRNA Re-splicing in MCF7 Cells

TSG101 mRNA is frequently re-spliced in various cancer tissues for unknown reasons. Since we have discovered the EJC core factor as a repressor of mRNA re-splicing, it is reasonable to predict reduced expression of an EJC core factor in cancer cells. We indeed observed a modest decrease of RBM8A protein in cancer MCF-7 cells compared to that in non-cancerous MCF-10A cells (Figure 5A, B). Remarkably, overexpression of RBM8A in stable cell line significantly restored the repression of mRNA re-splicing (Figure 5C, D), indicating that the observed repression in MCF-7 cells, at least, is indeed due to downregulation of RBM8A.

3. Discussion

Here we demonstrate that the minimum three core EJC factors, without any EJC peripheral factors, is responsible for repressing deleterious re-splicing on mature mRNA in normal cells, implicating the novel role of EJC core on mature mRNA in terminating constitutive splicing (Figure 6). Two intriguing questions remain; namely, why mRNA re-splicing often occurs in cancer cells but not in normal cells, and what is the exact mechanism underlying the EJC-triggered repression of mRNA re-splicing.

Using MCF-7 cells, we showed that the downregulation of EJC core factor, RBM8A, is one of the

causes of mRNA re-splicing.   We need extensive investigations to examine whether the repression of mRNA re-splicing is due to a scarcity of viable EJC cores using different kinds of cancer cells. Since EIF4A3 was not a TP53-dependent re-splicing repressor, there could be another unidentified regulator of mRNA re-splicing, which is under the control of the tumor suppressor gene *TP53*.   All these repressors, together with conceivable cancer-specific activators, of mRNA re-splicing should be comprehensively investigated to elucidate the cause of cancer-specific mRNA re-splicing frequently observed in various cancer cells/tissues (Figure S3).

Regarding the involved mechanism in the EJC-triggered repression of mRNA re-splicing, we propose the following two hypotheses; (i) The non-canonical binding of the ECJ core to the corresponding alternative splice sites on the spliced mRNA causes steric hindrance to prevent mRNA re-splicing.   (ii) The EJC promotes spliced mRNA into highly compacted mRNA particle (mRNP) structures that interfere with re-assembly of early spliceosome to initiate mRNA re-splicing (Figure 6).

Considering the hypothesis (i), recent studies uncovered EJC-triggered RNPS1-dependent and RNPS1-independent mechanisms that repress the use of spurious and recursive splice sites [20,21]. According to this RNPS1-independent mechanism, which is our case in the repression of mRNA re-splicing (Figure 4), the EJC core must directly bind upstream of the 3′ re-splice sites to prevent re-splicing.   However, reported data of cross-linking and immunoprecipitation coupled to high-throughput sequencing (CLIP-Seq) in TSG101 mRNA indicated that the significant binding sites of the EJC core factor, EIF4A3, were located more than 18 nucleotides downstream from the 3′ re-splice sites [22].   Therefore the possibility of masking the re-spliced 3' splice sites by the EJC core is unlikely.

The hypothesis (ii) is more promising.   After the release of the spliced mRNA from the spliceosome by the helicase DHX8, bare mature mRNA has a risk of nuclease digestion. Biochemical analysis of the components of mature mRNPs revealed that EJCs multimerize together with other RNA-binding proteins including SR proteins to form mega-Dalton sized higher-order complexes that protect spliced mRNA from endonuclease digestion [23,24].   This fact prompted us to assume that the EJC core depletion could prevent the formation of such highly compacted structures on spliced mRNA so that it allows re-assembly of the dissociated spliceosome to initiate mRNA re-splicing (Figure 6).   Further work is under way to address this attractive hypothesis.

It is conceivable that the EJC core is a common key factor controlling different types of the multi-step splicing process [20,21].   Using TSG101 pre-mRNA as a model substrate of cancer-specific aberrant mRNA re-splicing, we demonstrate that the deposition of the EJC safeguards against the deleterious over-splicing event.   We propose that EJC functions as a key signal to terminate pre-mRNA splicing and thus plays a further role to maintain quality of the protein-coding transcriptome (Figure S3).

4. Materials and Methods

4.1. Cell Culture and siRNA-mediated Knock-down

MCF-7 cells, obtained from the Cell Resource Center (http://www2.idac.tohoku.ac.jp/dep/ccr/index_en.html), were cultured in Dulbecco's modified Eagle's medium (Fujifilm Wako Chemicals, Kyoto, Japan) supplemented with 10% fetal bovine serum (Sigma-Aldrich, St. Louis, MO, USA).   MCF-10A cells, obtained from the American Type Culture Collection (ATCC), were cultured in a special medium (MEGM with MEGM SingleQuots; Lonza, Walkersville, MD, USA).

MCF-7 and MCF-10A cells were transiently transfected with the indicated siRNAs (Table S3) using Lipofectamine RNAi MAX (Thermo Fisher Scientific, Waltham, MA, USA), according to the manufacturer's instructions.   At 48 h post-transfection, total RNA was isolated from siRNA-treated cells using an RNeasy Mini Kit (QIAGEN, Venlo, Nederland) according to the manufacturer's instructions.

4.2. Establishment of Stable MCF-7 Cell Lines and Induction of RBM8A

All the synthetic oligonucleotides used as primers were purchased (Integrated DNA Technologies, Tokyo, Japan).   The coding-fragment of RBM8A protein was amplified by PCR on cDNA prepared

from MCF-7 cells using KOD-Plus-Neo polymerase (Toyobo, Osaka, Japan) with the specific primer set (Table S3).   The amplified DNA was subcloned into a TA cloning vector pGEM-T (Promega, Madison, WI) using TaKaRa Ex Taq polymerase (Takara Bio, Kusatsu, Japan).   The obtained pGEM-T-RBM8A plasmid was cleaved with Sfi I and subcloned into the Sfi I site of pSBtet-GP vector (Addgene, Watertown, MA, USA) [25].

In order to establish stable cell lines, the pSBtet-GP constructs (pSB-DsRed and pSB-RBM8A) were co-transfected with the *Sleeping Beauty* transposon vector pCMV(CAT)T7-SB100X (Addgene) [26] using Lipofectamine 3000 (Thermo Fisher Scientific), according to the manufacturer's instructions.   At 48 h post-transfection, cells were selected with1 µg/ml puromycin (Sigma-Aldrich) for ~10 days and terminated when most of cells showed GFP-derived green fluorescence.   To induce RBM8A and control DsRed proteins, 50 ng/ml of doxycycline (DOX) were added to each cell lines and cultured for 48 h.

4.3. RT–PCR and RT–qPCR Analyses

The cDNA was synthesized from total RNA using SuperScript III reverse transcriptase (Thermo Fisher Scientific) with Oligo(dT)18 primer (Thermo Fisher Scientific), and analyzed by PCR using TaKaRa Ex Taq polymerase (Takara Bio) with target-specific primers (Integrated DNA Technologies, Coralville, IA, USA; Table S3).   The PCR products were visualized by 2% agarose gel electrophoresis.   The PCR products were further quantified by Applied Biosystems 7900HT Fast Real Time PCR System (Thermo Fisher Scientific) using THUNDERBIRD SYBR qPCR Mix (Toyobo).

4.4. Immunoblotting Analysis

Cells were lysed with RIPA buffer supplemented with Protease Inhibitor Cocktail (Nacalai Tesque, Kyoto, Japan) according to the manufacturer's instructions, and separated by 15% SDS-polyacrylamide gel electrophoresis and electro-blotted onto a Immun-Blot PVDF Membrane using a Trans-Blot SD Semi-Dry Transfer Cell (Bio-Rad, Hercules, CA, USA).   The membranes were blocked with 2% skimmed milk and probed with antibodies, essentially as described [27].   The primary antibodies against GAPDH (1:2000 dilution; Medical & Biological Laboratories, Nagoya, Japan), EIF4A3 (1:1000; Abcam, Cambridge, UK), RBM8A (1:1000; GeneTex, Irvine, CA, USA), and MAGOH (1:1000; MBL) were used.   The secondary antibodies were the anti-rabbit IgG-HRP (1:2500; Sigma-Aldrich) and anti-mouse IgG-HRP (1:2500; Sigma-Aldrich).   Immuno-reactive proteins were visualized using the Immobilon Western Chemiluminescent HRP Substrate (Merck Millipore, Burlington, MA, USA).   The detected bands on the scanned blots were quantitated with ImageJ freeware.

4.5 RNA Immunoprecipitation (RIP) Assay

RIP assay with MCF-7 cells was performed using the RIP-Assay Kit (Medical & Biological Laboratories) with anti-RBM8A antibody (Medical & Biological Laboratories) according to the manufacturer's instructions.   The immunoprecipitated RNAs were amplified by RT–PCR using the primers targeting TSG101 mRNA (Table S3) and visualized by 2% agarose gel electrophoresis.

4.6. Statistical Analysis

Analyzed expression levels from two groups were compared using Welch's t-test.   One-way analysis of variance (ANOVA) followed by post-hoc tests were used to analyze results among multiple groups with a normal distribution and equal variance.   Statistical significance was set at $P < 0.05$.   Data were presented as mean ± standard deviations.

Author Contributions: Organizer of project, A.M.; planning of research, Y.O. and T.K.; experiments, Y.O.; data analyses, Y.O., T.K. and A.M.; writing—original draft preparation, Y.O.; writing—review and editing, Y.O., T.K. and A.M.; project supervision and administration, A.M.   All authors have read, edited, and agreed to the published version of the manuscript.

Funding: Y.O. was supported by Nippon Shinyaku Co., Ltd.   T.K. was partly supported by Grants-in-Aid for Scientific Research (C) [Grant number: JP17K07182] from the Japan Society for the

Promotion of Science (JSPS) and a research grant from the Aichi Cancer Research Foundation. A.M. was partly supported by Grants-in-Aid for Scientific Research (B) [Grant number: JP16H04705], Grants-in-Aid for Challenging Exploratory Research [Grant number: JP16K14659] from JSPS, the Austrian–Japanese Joint Projects funding from the Austrian Science Fund (FWF) and JSPS, the MEXT-Supported Program for the Strategic Research Foundation at Private Universities, and a research grant from the Aichi Cancer Research Foundation.

Acknowledgments: We are grateful to Drs. H. Naito, K. Takagaki, H. Fujiwara, and A. Matsuura in Nippon Shinyaku Co., Ltd. for their support and encouragements for Y.O. We thank all members of the Mayeda and Ishi labotarories for their constructive discussions, and Drs. H. Naito, K. Takagaki and J. Venables for critical reading of the manuscript.

Conflicts of interest: Y.O. was delegated to the Mayeda Laboratory as a visiting student from Nippon Shinyaku Co., Ltd., and he received financial support from the company. The other authors have no financial conflicts of interest.

References

1. Will, C.L.; Lührmann, R. Spliceosome structure and function. *Cold Spring Harb. Perspect Biol.* **2011**, *3*, a003707.

2. Kelly, S.; Georgomanolis, T.; Zirkel, A.; Diermeier, S.; O'Reilly, D.; Murphy, S.; Langst, G.; Cook, P.R.; Papantonis, A. Splicing of many human genes involves sites embedded within introns. *Nucleic Acids Res.* **2015**, *43*, 4721-4732.

3. Sibley, C.R.; Emmett, W.; Blazquez, L.; Faro, A.; Haberman, N.; Briese, M.; Trabzuni, D.; Ryten, M.; Weale, M.E.; Hardy, J.; Modic, M.*, et al.* Recursive splicing in long vertebrate genes. *Nature* **2015**, *521*, 371-375.

4. Parra, M.K.; Tan, J.S.; Mohandas, N.; Conboy, J.G. Intrasplicing coordinates alternative first exons with alternative splicing in the protein 4.1R gene. *EMBO J.* **2008**, *27*, 122-131.

5. Suzuki, H.; Kameyama, T.; Ohe, K.; Tsukahara, T.; Mayeda, A. Nested introns in an intron: evidence of multi-step splicing in a large intron of the human dystrophin pre-mRNA. *FEBS Lett.* **2013**, *587*, 555-561.

6. Gazzoli, I.; Pulyakhina, I.; Verwey, N.E.; Ariyurek, Y.; Laros, J.F.; t Hoen, P.A.; Aartsma-Rus, A. Non-sequential and multi-step splicing of the dystrophin transcript. *RNA Biol.* **2016**, *13*, 290-305.

7. De Bortoli, F.; Espinosa, S.; Zhao, R. DEAH-box RNA helicases in pre-mRNA splicing. *Trends Biochem. Sci.* **2020**.

8. Venables, J.P. Aberrant and alternative splicing in cancer. *Cancer Res.* **2004**, *64*, 7647-7654.

9. Kameyama, T.; Suzuki, H.; Mayeda, A. Re-splicing of mature mRNA in cancer cells promotes activation of distant weak alternative splice sites. *Nucleic Acids Res.* **2012**, *40*, 7896-7906.

10. Chua, H.H.; Huang, C.S.; Weng, P.L.; Yeh, T.H. TSGΔ154-1054 splice variant increases TSG101 oncogenicity by inhibiting its E3-ligase-mediated proteasomal degradation. *Oncotarget* **2016**, *7*, 8240-8252.

11. Chua, H.H.; Kameyama, T.; Mayeda, A.; Yeh, T.H. Cancer-Specifically Re-Spliced TSG101 mRNA Promotes Invasion and Metastasis of Nasopharyngeal Carcinoma. *Int. J. Mol. Sci.* **2019**, *20*.

12. Ferraiuolo, R.M.; Manthey, K.C.; Stanton, M.J.; Triplett, A.A.; Wagner, K.U. The Multifaceted Roles of the Tumor Susceptibility Gene 101 (TSG101) in Normal Development and Disease. *Cancers (Basel)* **2020**, *12*.

13. Le Hir, H.; Sauliere, J.; Wang, Z. The exon junction complex as a node of post-transcriptional networks. *Nat. Rev. Mol. Cell Biol.* **2016**, *17*, 41-54.

14. Schlautmann, L.P.; Gehring, N.H. A Day in the Life of the Exon Junction Complex. *Biomolecules* **2020**, *10*.

15. Moyret-Lalle, C.; Duriez, C.; Van Kerckhove, J.; Gilbert, C.; Wang, Q.; Puisieux, A. p53 induction prevents accumulation of aberrant transcripts in cancer cells. *Cancer Res.* **2001**, *61*, 486-488.

16. Ding, Q.; Zhang, Z.; Liu, J.J.; Jiang, N.; Zhang, J.; Ross, T.M.; Chu, X.J.; Bartkovitz, D.; Podlaski, F.; Janson, C.; Tovar, C*., et al.* Discovery of RG7388, a potent and selective p53-MDM2 inhibitor in clinical development. *J. Med. Chem.* **2013**, *56*, 5979-5983.

17. Gehring, N.H.; Lamprinaki, S.; Hentze, M.W.; Kulozik, A.E. The hierarchy of exon-junction complex assembly by the spliceosome explains key features of mammalian nonsense-mediated mRNA decay. *PLoS Biol.* **2009**, *7*, e1000120.

18. Smith, C.M.; Steitz, J.A. Classification of *gas*5 as a multi-small-nucleolar-RNA (snoRNA) host gene and a member of the 5'-terminal oligopyrimidine gene family reveals common features of snoRNA host genes. *Mol. Cell. Biol.* **1998**, *18*, 6897-6909.

19. Tani, H.; Torimura, M.; Akimitsu, N. The RNA degradation pathway regulates the function of GAS5 a non-coding RNA in mammalian cells. *PLoS One* **2013**, *8*, e55684.

20. Boehm, V.; Britto-Borges, T.; Steckelberg, A.L.; Singh, K.K.; Gerbracht, J.V.; Gueney, E.; Blazquez, L.; Altmuller, J.; Dieterich, C.; Gehring, N.H. Exon Junction Complexes Suppress Spurious Splice Sites to Safeguard Transcriptome Integrity. *Mol. Cell* **2018**, *72*, 482-495 e487.

21. Blazquez, L.; Emmett, W.; Faraway, R.; Pineda, J.M.B.; Bajew, S.; Gohr, A.; Haberman, N.; Sibley, C.R.; Bradley, R.K.; Irimia, M.; Ule, J. Exon Junction Complex Shapes the Transcriptome by Repressing Recursive Splicing. *Mol. Cell* **2018**, *72*, 496-509 e499.

22. Saulière, J.; Murigneux, V.; Wang, Z.; Marquenet, E.; Barbosa, I.; Le Tonquèze, O.; Audic, Y.; Paillard, L.; Roest Crollius, H.; Le Hir, H. CLIP-seq of eIF4AIII reveals transcriptome-wide mapping of the human exon junction complex. *Nat. Struct. Mol. Biol.* **2012**, *19*, 1124-1131.

23. Singh, G.; Kucukural, A.; Cenik, C.; Leszyk, J.D.; Shaffer, S.A.; Weng, Z.; Moore, M.J. The cellular EJC interactome reveals higher-order mRNP structure and an EJC-SR protein nexus. *Cell* **2012**, *151*, 750-764.

24. Metkar, M.; Ozadam, H.; Lajoie, B.R.; Imakaev, M.; Mirny, L.A.; Dekker, J.; Moore, M.J. Higher-Order Organization Principles of Pre-translational mRNPs. *Mol. Cell* **2018**, *72*, 715-726 e713.

25. Kowarz, E.; Loscher, D.; Marschalek, R. Optimized Sleeping Beauty transposons rapidly generate stable transgenic cell lines. *Biotechnol. J.* **2015**, *10*, 647-653.

26. Mates, L.; Chuah, M.K.; Belay, E.; Jerchow, B.; Manoj, N.; Acosta-Sanchez, A.; Grzela, D.P.; Schmitt, A.; Becker, K.; Matrai, J.; Ma, L*., et al.* Molecular evolution of a novel hyperactive Sleeping Beauty transposase enables robust stable gene transfer in vertebrates. *Nat. Genet.* **2009**, *41*, 753-761.

27. Harlow, E.; Lane, D. Immunoblotting. In *Antibodies: A Laboratory Manual*, Harlow, E.; Lane, D., Eds.; Cold Spring Harbor Laboratory: 1988; pp 471–510.

Figure 1. Identification of EJC-core factors as repressor candidates of cancer-specific mRNA re-splicing. (**a**) The schematic structure of normally spliced TSG101 mRNA and cancer-specifically re-spliced TSG101Δ154–1054 mRNA. The primer sets for the detection were indicated with arrows. (**b**) To measure the generated normal TSG101 and aberrant TSGΔ154-1054 mRNAs, the extracted total RNAs were analyzed by RT–qPCR using the indicated specific primer sets in (A). To knock-down the expression of the endogenous *EIF4A3* gene, MCF-7 cells were transfected with EIF4A3 siRNAs together with control siRNA. The cell extracts were then analyzed by RT–qPCR and the relative amounts of EIF4A3 mRNA are plotted. The histograms represent the means ± standard deviations of three replicates (*P < 0.05). (**c**) To knock-down the expression of endogenous EJC core factor genes, MCF-7 cells were transfected with siRNA targeting EIF4A3, MAGOH, RBM8A, CASC3 or control siRNA (cont.). To observe the knocked-down mRNAs and generated normal TSG101 and aberrant TSG101 mRNAs, the extracted total RNAs were analyzed by RT–PCR.

Figure 2. Depletion of EJC-core factors can induce cancer-specific mRNA re-splicing in non-cancerous MCF-10A cells. (**A**) Total RNAs extracted from MCF-10A or MCF-7 cells were analyzed by RT-PCR using specific primer sets (see Figure 1b). (**B**) To knock-down the expression of endogenous EJC core factor genes, MCF-10A cells were transfected with the indicated siRNAs. To observe the knocked-down efficiency and TSG101 re-splicing activity, the extracted total RNAs were analyzed by RT–PCR.

Figure 3. Activation of mRNA re-splicing by the knock-down of EJC core factors is not due to the repression of EJC-mediated NMD. (**a**) To knock-down the expression of endogenous NMD factor UPF1, MCF-10A cells were transfected with the indicated siRNAs. To observe the knocked-down mRNA and TSG101 re-splicing activity, the extracted total RNAs were analyzed by RT–PCR. MCF-7 cells (control) shows the re-spliced mRNA that lacks in other lanes using MCF-10A cells. (**b**) HeLa cells were transfected with EIF4A3 or UPF1 siRNAs (or control siRNA). To evaluate the re-splicing activity with TSG101 mRNA and the NMD efficiency with GAS5 mRNA, the extracted total RNAs were analyzed by RT–qPCR using specific primer sets. The histograms represent the means ± standard deviations of three replicates (*$P < 0.05$, n.s.=not statistically significant $P > 0.05$).

Figure 4. Depletion of an mRNA export factor or EJC-peripheral factors has no significant effects on mRNA re-splicing in MCF-7 cells. (**a, b**) To knock-down the expression of endogenous mRNA export factor (NXF1) and EJC-peripheral factors (ACIN1, PNN, RNPS1, SAP18), MCF-7 cells were transfected with the indicated siRNAs. To observe the knocked-down efficiency and TSG101 re-splicing activity, the extracted total RNAs were analyzed by RT–PCR.

Figure 5. RBM8A protein expression is lower in cancer cells as compared with that in non-cancerous cells, and RBM8A overexpression represses mRNA re-splicing in cancer cells. (**a**) Whole cell extracts from MCF-10A and MCF-7 cells were analyzed by immunoblotting with the indicated specific antibody. GAPDH was used as an endogenous control. (**b**) RBM8A protein expression was normalized to that of GAPDH and quantified by densitometry. The histograms represent the means ± standard deviations of three replicates (*$P < 0.05$). (**c**) The stable MCF-7 cell lines expressing control and RBM8A proteins (pSB-DsRed and pSB-RBM8A) were cultured with or without doxycycline (DOX) for 48 h. The total RNAs were analyzed by RT–PCR using the indicated primer sets. (**d**) To evaluate the re-splicing activity with TSG101 mRNA, the total RNAs were analyzed by qRT-PCR using specific primer sets. The histograms represent the means ± standard deviations of three replicates (*$P < 0.05$, n.s. $P > 0.05$).

Figure 6. Model of the EJC-triggered pathway leading to the termination of splicing. The spliced mRNA was shown to form packed and compacted structure of mRNP by bindings between EJC cores and among EJC and SR proteins (EJC-SR nexus) together with other RNA-binding proteins (RBPs) [23,24]. We assume that splicing-dependent binding of EJC core to mature mRNA is essential to initiate the formation of this highly compacted mRNP structure.

Communication

The Exon Junction Complex Core Represses Caner-specific Mature mRNA Re-splicing: A Potential Key Role in Terminating Splicing

Yuta Otani[1,2], Toshiki Kameyama[1,3] and Akila Mayeda[1,*]

[1] Division of Gene Expression Mechanism, Institute for Comprehensive Medical Science, Fujita Health University, Toyoake, Aichi 470-1192, Japan
[2] Laboratories of Discovery Research, Nippon Shinyaku Co., Ltd., Kyoto, Kyoto 601-8550, Japan
[3] Present address: Department Physiology, Fujita Health University, School of Medicine, Toyoake, Aichi 470-1192, Japan

* Correspondence: mayeda@fujita-hu.ac.jp (A.M.)

Contents

Table S3 List of the synthetic oligonucleotides used in the experiments (see also Table S2).

siRNAs (sense sequences) for cellular knockdown (T: deoxyribonucleotide)		
EIF4A3 siRNA #1	ID: s18876 (Thermo Fisher Scientific)	Knockdown (KD) of EIF4A3 (shared in Table S2)
EIF4A3 siRNA #2	ID: s18877 (Thermo Fisher Scientific)	
RBM8A (Y14) siRNA #1	ID: s532199 (Thermo Fisher Scientific)	KD of RBM8A
RBM8A (Y14) siRNA #2	ID: s532200 (Thermo Fisher Scientific)	
MAGOH siRNA #1	ID: s226590 (Thermo Fisher Scientific)	KD of MAGOH
MAGOH siRNA #2	ID: s8466 (Thermo Fisher Scientific)	
CASC3 (MNL51) siRNA #1	ID: s22392 (Thermo Fisher Scientific)	KD of CASC3
CASC3 (MNL51) siRNA #2	ID: s22393 (Thermo Fisher Scientific)	
UPF1 siRNA #1	ID: s11927 (Thermo Fisher Scientific)	KD of UPF1
UPF1 siRNA #2	ID: s11928 (Thermo Fisher Scientific)	
NXF1 siRNA #1	ID: s20533 (Thermo Fisher Scientific)	KD of UPF1
NXF1 siRNA #2	ID: s20532 (Thermo Fisher Scientific)	
ACINUS siRNA	5′-gaggccuucuggauugacaTT -3′ (Integrated DNA Technologies)	KD of ACINUS
PININ siRNA	5′-ggagcaagggcauuuacuaTT-3′ (Integrated DNA Technologies)	KD of PININ
RNPS1 siRNA	5′-caaaggcuaugcguacguaTT-3′ (Integrated DNA Technologies)	KD of RNPS1
SAP18 siRNA	5′-gaacugacaagcuuaguaaTT-3′ (Integrated DNA Technologies)	KD of SAP18
Primer DNAs for plasmid construction (F: forward primer, R: reverse primer)		
RBM8A coding-F	5′-ACAAGTGGCCTCTGAGGCCCCACCATGGCGGACGTGCTAGATCT-3′	Construction for final pSB-RBM8A
RBM8A coding-R	5′-TGACGGGCCTGACAGGCCTCAGCGACGTCTCCGGTCTG-3′	
Primer DNAs for mRNA products assay (F: forward primer, R: reverse primer)		
GAPDH-F	5′-GCACCGTCAAGGCTGAGAAC-3′	PCR for detection of GAPDH cDNA
GAPDH-R	5′-TGGTGAAGACGCCAGTCCA-3′	
EIF4A3-F	5′-GATCTTGGCTCCCACAAGAG-3′	PCR for detection of EIF4A3 cDNA
EIF4A3-R	5′-TAGTCACCGAGAGCAAGCAG-3′	
TSG101-F	5′-GAAGTTATCATTCCCACAGCTC-3′	PCR for detection of spliced TSG101
TSG101-R	5′-GACGTACATGCTTCAGGAAGAC-3′	
TSG101Δ154-1054-F	5′-TACAAATACAGAGACCTAACTCTCCC-3′	PCR for detection of re-spliced TSG101
TSG101Δ154-1054-R	5′-GACGTACATGCTTCAGGAAGAC-3′	
RBM8A-F	5′-ATTCATCTCAACCTCGACAGG-3′	PCR for detection of RBM8A cDNA
RBM8A-R	5′-CAAATCCTGGCCATTGAGTC-3′	
MAGOH-F	5′-CAAAGAGGATGATGCATTGTG-3′	PCR for detection of MAGOH cDNA
MAGOH-R	5′-TGTGTTCATCTCCAATGACGA-3′	
CASC3-F	5′-ACCACCGCCTCATCTGTATC-3′	PCR for detection of CASC3 cDNA
CASC3-R	5′-TGGGCGGGGTTATAGTAGGT-3′	
UPF1-F	5′-GCAACGGACGTGGAAATACT-3′	PCR for detection of UPF1 cDNA
UPF1-R	5′-CTTGTGCAGGGTCACCTCTT-3′	
GAS5-F	5′-CTTGCCTGGACCAGCTTAAT-3′	PCR for detection of GAS5 cDNA
GAS5-R	5′-CAAGCCGACTCTCCATAC-3′	
NXF1-F	5′-GATTCTTTGCGAGCCTTCACC-3′	PCR for detection of NXF1 cDNA
NXF1-R	5′-AGGAGGAGATGACAGACGACAACC-3′	
ACINUS-F	5′-GACGATCATCTAGGGTCAGACAG-3′	PCR for detection of ACINUS cDNA
ACINUS-R	5′-GGAAGGTGTTTCTTGATCCTCTT-3′	

Left, middle, and right columns: names, sequences, and purpose of the oligonucleotides, respectively.

Figure S1. Activation of TP53 by the MDM2 inhibitor RG7388 induces the repression of TSG101 mRNA re-splicing activity. To evaluate the re-splicing activity with TSG101 mRNA, the extracted total RNAs from MCF-7 cells were analyzed by RT–qPCR using specific primer sets (see Figure 1a). The RG7388 concentrations and culture time after RG7388 addition were indicated. The histograms represent the means ± standard deviations of three replicates (*$P < 0.05$).

Figure S2. The EJC core factor RBM8A binds TSG101 mRNAs in MCF-7 cells. (**a**) RBM8A-bound mRNAs were separated from cell lysates of MCF-7 cells by RNA immunoprecipitation (RIP). Normal rabbit IgG was used as a negative control of the immunoprecipitation. Two independent RIP assays were performed and a representative immunoblot is shown here. (**b**) The RIP samples were analyzed by RT–PCR using a primer set targeting spliced TSG101 mRNA. The PCR products were visualized by agarose gel electrophoresis.

Figure S3. Model of the regulation system of cancer-specific mRNA re-splicing. Unidentified factors are indicated with question marks (?). Our discovery of EJC-core as a repressor of mRNA re-splicing implies a mechanism for terminating pre-mRNA splicing in normal cells, which is essential to maintain quality of mRNA.