Molecular diagnosis of new isolate of tomato yellow leaf curl virus in Iraq

Ali Alabde, Osamah Alisawi*, Fadhal Al Fadhal

Department of Plant Protection, Faculty of Agriculture, Kufa University, Najaf, Iraq

*Corresponding Author

Osamah Alisawi
Department of Plant Protection
University of Kufa
Najaf-Iraq
Tel: 009647816353396

E-mail: osama.alisawi@uokufa.edu.iq

Keywords: Tomato yellow leaf curl virus; next generation sequencing; bioinformatics; tomato
16   genome.

Abstract

Tomato yield and quality in Iraq have been threatened by a variable range of infections caused

by tomato yellow leaf curl virus. In previous studies, the TYLCV isolates were partially

characterized using molecular tests for small fragments not the entire length of the virus. Sample

of TYLCV-infected tomato has applied in this study to diagnose complete sequence of TYLCV

isolate. Three sets of primers that belong to three well-identified strains in Iraq were used in a

PCR technique and interestingly the results were negative. A new Iraqi isolate has been

characterized as a first novel Iraqi isolate detected ever using next generation (NGS) and

bioinformatics techniques. The NGS platform has produced about 78,232,062 paired reads of the

TYLCV-infected tomato var. Oula F1. The complete raw reads of the infected variety have been

analyzed using RepeatExplorer pipeline and Map to reference. The full sequence of TYLCV was

reconstructed and extracted to consist of 2770 nt and then deposited in Genbank under accession

number MT583814. Copy numbers and genome proportion of this sequence have been calculated that were 3523 and 0,086% respectively. Phylogenetic analyses of full nucleotide sequences confirmed close relationship to Iranian isolate (TYLCV-Kahnooj) than other published viruses. Additionally, a and β DNA satellites have not discovered in the TYLCV-infected sample.

Introduction

The family Geminiviridae is recently divided into seven genera: *Becurtovirus*, *Begomovirus*, *Curtovirus*, *Eragrovirus*, *Mastrevirus*, *Topocuvirus* and *Turncurtovirus.* Host range, insect vector, genome organization and genome-wide pairwise sequence identities have been applied to classify this family (Varsani *et al*., 2017). Tomato yellow leaf curl virus (TYLCV) belongs to genus *Begomovirus* and has a monopartite single-stranded DNA (ssDNA) encapsidated in geminate particles. TYLCV is transmitted by the whitefly, *Bemisia tabaci* (Genn.) that effectively spread virus particles through persistent manner (Panno *et al*., 2016). The disease caused by this virus is the most important disease on tomato in Iraq and worldwide (Mansoor *et al*., 2003; Salati *et al.,* 2002; Al-Kuwaiti *et al*., 2013). Infected plants become stunted with chlorotic leaves, curled-up margins, and abscised flowers. Since the first infection was recorded in Israel 1939-1940, multiple regions have been identified to have isolates from this virus over the world and classified to different viral species (Antignus and Cohen 1994; Delatte *et al*., 2004; Pico *et al*., 1996; Polston and Anderson 1997). Tomato yellow leaf curl Sardinia virus TYLCSV infected tomatoes in Israel and Sardinia, and was the first sequenced isolate in 1991(Kheyr-Pour *et al*., 1991; Navot *et al*., 1991), and since that date many isolates have been sequenced from various locations (Moriones and Navas-Castillo 2000). In Iraq, the disease was detected based on symptoms assay and serological tests. Recently, most popular molecular works on this virus have been achieved by PCR based on amplified short fragments of TYLCV genome such as coat protein domain (Al-Ani *et al*., 2010; Reddy *et al*., 2005; Maruthi *et al*., 2007). Further, many tomato genotypes have been screened against TYLCV isolates to characterize similarities and differences through these genotypes (Al Tamimi, 2020). TYLCV has been isolated not only from tomato but also from various plants such as eggplant, melon, cucumber, sesame, cowpea in addition to weeds as a primary source of

infection like *Malva parviflora* , *Chenopodium murale* and *Chenopodium quinoa* (Al- Jubouri *et*

al ., 2014). Al-Kuwaiti *et al* (2013) have reported complete sequence of TYLCV that showed

99% identity to Spain strain (Acc. AJ489258) confirming its existence in Iraq. In this study, we

utilized next generation sequencing and bioinformatics techniques for the first time in Iraq to

find out new isolate with complete sequence of TYLCV. Furthermore, phylogeny analyses have

been performed to understand how these isolates related to each other.

Materials and methods

Plant material

From open fields of Najaf governorate, the main region of tomato production in Iraq, fresh and

young leaves of TYLCV-infected tomato variety Oula F1 (Seminis Seeds Company, USA) were

harvested. This variety is considered the most popular and TYLCV-sensitive commercial variety

of tomato in the middle of Iraq.

DNA isolation, PCR and NGS sequencing

To extract the DNA, the young leaves (about 2 gm) of the infected tomato were applied using

cetyl-trimethylammonium bromide (CTAB) method with minor modifications (Doyle and Doyle

1990). Three sets of primers that belong to strains of TYLCV-IS, TYLCV-ES and TYLCV-Mld

were applied in a PCR technique to check their infection within examined sample (Table 1)

(Anfoka *et al*., 2005).

Table 1. Three primers of main strains of TYLCV in Iraq that used to amplify and characterize

virus sequence in the examined sample.

Primers	Sequence (5'- 3')	Size (bp)	Strains
TYv2337 TYc138	ACGTAGGTCTTGACATCTGTTGAGCTC AAGTGGGTCCCACATATTGCAAGAC	634	TYLCV-IS Israel
TYAlmv2516 TYAlmcl15	TTTTATTTGTTGGTGTTTGTAGTTGAAG ATATTGATGGTTTTTCAAAACTTAGAAG	433	TYLCSV-ES Spain Almeria

| TYv2664 | ATTGACCAAGATTTTTACACTTATCCC | 316 | TYLCV – mld |
| TYc138 | AAGTGGGTCCCACATATTGCAAGAC | | Normal mild |

PCR conditions were an initial denaturation of 10 min at 95°C, 49 cycles of 30 s at 95°C, 20 s at

55-63°C and 30 s at 72°C followed by a final extension of 30 s at 72°C. To obtain the whole

genome sequence, the extracted genomic DNA sample was sent to sequencing at DNA-link

Company, Republic of Korea. The genome was sequenced by using the Novaseq6000 2x150bp

reads technique and application WGS (PCR Free550) based on the manufacturer's procedure.

RepeatExplorer program

To collect virus sequences from the whole data (NGS) of infected tomato, RepeatExplorer

pipeline was applied to recognize these kinds of sequences in forms of clusters within repetitive

DNA blocks (Novák *et al*. 2013). The targeted clusters of RepeatExplorer were extracted, and

then Repbase dataset (Jurka *et al*. 2005), and Basic Local Alignment Search Tool (Altschul *et al*.

1990) were used to identify contigs of each cluster. Furthermore, the extracted sequences were

mapped to well-identified plant virus sequences from DPVweb (Adams and Antoniw 2005).

Map to reference

Using Geneious program V. 11 (Kearse *et al*. 2012), the raw reads were assembled against

TYLCV Iraq (Acc. JQ354991.1) as a reference sequence that diagnosed by Al-Kuwaiti *et al*.

(2013) and then consensus sequence was collected. Mapping to reference was repeated against

consensus sequence (TYLCV) to confirm and obtain the actual number of assembled reads.

Also, the raw reads were applied to assemble against a and β DNA satellites. The outcomes

showed in a report that has a number of assembled reads and total used reads in addition to the

frequent overlapped reads. This data used to calculate copy numbers (Number of assembled

reads x read length/reference sequence length) and genome proportions (Number of assembled

reads / numbers of total NGS reads x 100) (Mustafa *et al*. 2018). Finally, the two consensus

sequences from RepeatExplorer and Map to reference were aligned using pairwise alignment in

Geneious program to collect the right length of TYLCV isolate.

Phylogenetic analysis

The maximum likelihood (ML) method was applied to find phylogeny model using MEGA7

(Tamura *et al*. 2013). The Geneious program V. 11 (Kearse *et al*. 2012) was used for alignment

and optimized manually. The entire nucleotide sequences of 13 main TYLCV isolates including

our isolate of interest, Kahnooj and Iraq isolates were applied to phylogeny study. ClustalW

alignment was used for extracting sequences about 2770 nt for each. The phylogenetic tree

reconstructed using General Time Reversible (GTR). Bayesian phylogeny inference was used for

analysis with Bayesian inference of phylogeny (MrBayes 3.2.6) (Huelsenbeck and Ronquist

2001).

Results

The extracted DNA concentration and quality were 418 ng\ml and 1.92 which was in a high

quality level for sequencing approach (Fig. 1. A). PCR has surprisingly failed to get amplified

sequences for the three TYLCV isolates indicating that our sample has probably a new virus

sequence. High quality raw reads have been obtained from NGS Illumina platform with about

78,232,062 paired reads; 150 bp read length and 36.9% GC content (Fig. 1. B).

RepeatExplorer has been clustered the whole uploaded raw reads that showed tandemly repeated

sequences, TYLCV sequences involved in cluster number 76 as a putative satellite with a

genome proportion 0,084% and the consensus sequence was 2763 nt (Fig. 1. C). On the other

side, the report of Map to reference revealed that 65,303 reads have been assembled to count

genome proportion and copy numbers of the isolate that were 0,086% and 3523 respectively

whereas the consensus sequence was 2770 nt. In order to obtain complete sequence, the aligned

consensus sequences from both RepeatExplorer and Map to reference showed 100% similarity

and the longest sequence was extracted. Lastly, we determined 2770 nt of TYLCV and named as

Najaf isolate and then deposited in Genbank under accession number MT583814.1.

Phylogenetic analyses showed what we have proven earlier by pairwise alignment between Najaf

isolate and each of Kahnooj and Iraq isolates. Kahnooj was most likely similar to Najaf isolate

than Iraq. TYLCV nucleotide and amino acid sequence alignments showed that Najaf isolate is

closely related to Kahnooj isolate the Iranian TYLCV in 93.8% pairwise identity while it was

90.3% with Iraq isolate (Fig. 2). However, polymorphic sequences have shown higher rate in

intergenic region (IR) (75%), V2 (91.9%), C1 (Rep) (93.1%), and C4 (88.3%) than other coding

regions V1 (97.3%), C3 (95.9%) and C2 (96.5%).

Our research has been extended trying to find DNA-a and β satellites using Map to reference tool

but we did not get any positive hits.

Discussion

The deep sequencing of whole genome sequencing technique has revolutionized genomic

research collecting a massive data of the host genome with complete sequences of cellular

components. Additionally, this method is considered as unbiased approach to detect plant viruses

with no previous knowledge about the suspected elements. The NGS technique does not use

specific primers, antibodies, and virus-specific reagents to diagnose pathogenic viruses (Adams

et al., 2009). In this method, full image of plant viruses and complex infection could be

investigated in the host genome unlikely to biased methods that focus on almost one expected

virus (Jones *et al*., 2017). In Iraq, most of molecular tests concentrated on partial sequences of

TYLCV such as coat protein domain to detect the virus and apply these amplified parts in

multiple alignments and phylogeny analyses. The only full-length sequenced of this virus from

Iraq was Iraq isolate (JQ354991.1) that achieved by Al-Kuwaiti *et al* (2013) and named as

TYLCV Iraq but it shared   99% similarity to Spain strain (Acc. AJ489258), Reunion Islands

(Acc. AM409201) and Mauritius (Acc. HM448447). Additionally, Sadeq Al-Kuwaiti (2013) has

applied rolling circle amplification (RCA) and non-degenerate primers to catch the full length of

virus.   Therefore, our new isolate is considered the first full sequenced Iraqi isolate of TYLCV

with totally new data yet. We found Kahnooj isolate from Iran is closer than other isolates even

from Iraq with 93.8 % similarity. It is highly expected that Najaf and Kahnooj isolates are

sharing the same ancestor and most likely the virus was imported into Iraq from Iran by

international trading (Mali *et al*., 2003; Gibbs *et al*., 2008; Al-Kuwaiti 2013).

Mapping reads to the reference has revealed the estimated copy numbers of this virus was 3523

within infected cell. Copy number value has been always considered as a good sign for variation

and variability of endogenous components within host genome especially transposable elements
including viruses (Mustafa *et al*. 2018; Alisawi 2019). Interestingly, this is one of NGS and
bioinformatics benefits to calculate copy numbers of the components of interest and this
approach has been applied in many studies to predict viral copies (Yang *et al*., 2017; Alisawi
2019). So far, the number of copies indicates virus particles indeed infected the host genome. In
fact, this little amount of copies strongly explains the early stage of symptoms emerging in the
infected sample. However, further works need to be achieved regarding isolate distribution and
diversity over tomato varieties and different regions in Iraq.

We could not reveal the existence of DNA satellites in our infected sample probably due to
moderate level of infection in the examined sample. A and β DNA satellites are often associated
with begomoviruses in their replication, movement within infected plant and horizontal
transmission between plants, as well as involved in induction of virus symptoms and
suppression of transcriptional gene silencing. This result is consistent with most published
outcomes that reported the importance of DNA satellites for disease severity and inducing
symptoms. However, we do expect to find these satellites in severely TYLCV-infected plants as
no more researches have been covered DNA satellites in Iraq (Laufs *et al*., 1995; Saunders *et al*.,
2004; Cui *et al*., 2005; Yang *et al*., 2011; Amin *et al*., 2011; Zhou, 2013).

Conclusion

This work concluded that whole genome sequencing technique alongside bioinformatics is
highly recommended to detect complete sequence of TYLCV isolate within infected tomato
genome raw reads. Consequently, a clear insight of virus pathway and identity has been revealed
by phylogenetic analysis. More investigations need to be done for detecting full sequences of
highly pathogenic plant viruses in Iraq.

Conflict of Interest Statement
The authors have no conflicts of interest to declare.

Funding Sources

There are no funding sources to report.

Statement of Ethics

The authors have no ethical conflicts to disclose.

Author Contributions

A.A. carried out the experiment. O.A. analyzed the NGS data and wrote the manuscript with
support from F.A. F.A. helped supervise the study.

References

Adams, I. P., Glover, R. H., Monger, W. A., Mumford, R., Jackeviciene, E., Navalinskiene, M.,
... & Boonham, N., 2009. Next-generation sequencing and metagenomic analysis: a universal
diagnostic tool in plant virology. *Molecular plant pathology*, *10*(4), 537-545.

Adams M.J. & Antoniw J.F, 2005. DPVweb: An open access internet resource on plant viruses
and virus diseases. *Outlooks on Pest Management* 16, 268-270.

Al-Ani, R. A., Diwan, S. N. H., & Adhab, M. A, 2010. Efficiency of *Thuja orientalis* and
Artimisia campestris extracts to control of *Potato leaf roll virus* (PLRV) in potato
plants. *Agriculture and Biology Journal of North America*, *1*(4), 579-583.

Alisawi, O. N, 2019. Virus integration and tandem repeats in the genomes of *Petunia* (Doctoral
dissertation, University of Leicester).

Al-Jubouri, A. A., Al-Fadhal, F. A., & Samaka, H. M, 2014. Detection of *Tomato yellow leaf*
curl virus (TYLCV) and its strains on tomato plants and host rang determine by Polymerase
Chain Reaction. *Kufa Journal for Agricultural Sciences*, *6*(4), 99-117.

Al-Kuwaiti, N.; B. Otto; C. Collins; S. Seal and Maruthi, M, 2013. Molecular characterization
and first complete genome sequence of *Tomato yellow leaf curl virus* (TYLCV) infecting
tomato in Iraq. *New Disease Reports*, 27: 17.

Altschul S.F., Gish W., Miller W., Myers E.W. & Lipman D.J, 1990. Basic local alignment
search tool. *Journal of Molecular Biology* 215, 403-410.

Al Tamimi, S. Q, 2020. Molecular identification of some isolates of *Tomato yellow leaf curl*

virus and test response some tomato genotypes and their control(Doctoral dissertation,

University of Kufa)(In Arabic).

Amin, I., Hussain, K., Akbergenov, R., Yadav, J. S., Qazi, J., Mansoor, S., ... & Briddon, R. W,

2011. Suppressors of RNA silencing encoded by the components of the cotton leaf curl

begomovirus-betasatellite complex. *Molecular plant-microbe interactions*, *24*(8), 973-983.

Anfoka, G.H., Abhary, M. and Nakhla, M.K., 2005. Molecular identification of species of the

Tomato yellow leaf curl virus complex in Jordan. *Journal of Plant Pathology*, pp.65-70.

Antignus, Y., & Cohen, S, 1994. Complete nucleotide sequence of an infectious clone of a mild

isolate of *Tomato yellow leaf curl virus* (TYLCV). *Phytopathology*, *84*(7), 707-712.

Cui, X., Li, G., Wang, D., Hu, D., & Zhou, X., 2005. A begomovirus DNAβ-encoded protein

binds DNA, functions as a suppressor of RNA silencing, and targets the cell nucleus. *Journal of*

Virology, *79*(16), 10764-10775.

Delatte H, Naze F, Granier M, Reynaud B, Peterschmitt M, Lett M., 2004. Tomato-infecting

begomoviruses in the South Western islands of the Indian Ocean. *Presented at the 4th*

International Geminivirus Symposium, W3-4.

Doyle J. & Doyle J., 1990. Isolation of DNA from small amounts of plant tissues. *BRL Focus*

12, V15.

Gibbs, A. J., Ohshima, K., Phillips, M. J., & Gibbs, M. J., 2008. The prehistory of potyviruses:

their initial radiation was during the dawn of agriculture. *PloS one*, *3*(6), e2523.

Huelsenbeck J.P. & Ronquist F., 2001. MRBAYES: Bayesian inference of phylogenetic trees.

Bioinformatics 17, 754-755.

Jones, S., Baizan-Edge, A., MacFarlane, S., & Torrance, L., 2017. Viral diagnostics in plants

using next generation sequencing: computational analysis in practice. *Frontiers in Plant*

Science, *8*, 1770.

Jurka J., Kapitonov V.V., Pavlicek A., Klonowski P., Kohany O. & Walichiewicz J., 2005.

Repbase Update, a database of eukaryotic repetitive elements. *Cytogenetic and Genome*

Research 110, 462-467.

Kheyr-Pour A, Bendahmane M, Matzeit V, Accotto GP, Crespi S, Gronenborn B., 1991. *Tomato*

yellow leaf curl virus from Sardinia is a whitefly-transmitted monopartite geminivirus. *Nucleic*

Acids Res 19:6763–6769.

Kearse M., Moir R., Wilson A., Stones-Havas S., Cheung M., Sturrock S., Buxton S., Cooper A.,
Markowitz S. & Duran C., 2012. Geneious Basic: an integrated and extendable desktop software
platform for the organization and analysis of sequence data. *Bioinformatics* 28, 1647-1649.

Laufs J., Traut W., Heyraud F., Matzeit V., Rogers S. G., Schell J., Gronenborn B., 1995. In
vitro cleavage and joining at the viral origin of replication by the replication initiator protein of
Tomato yellow leaf curl virus. *Proc Natl Acad Sci U S A* 92:3879–3883.

Mali, V. R., Subr, Z., & Kudela, O., 2003. Seed transmission of como and potyviruses in faba
bean and vetch cultivars introduced into Slovakia. *Acta phytopathologica et entomologica*
hungarica, *38*(1-2), 87-97.

Mansoor, S., Briddon, R. W., Zafar, Y., & Stanley, J., 2003. Geminivirus disease complexes: an
emerging threat. *Trends in plant science*, *8*(3), 128-134.

Maruthi M, Rekha A, Mirza S, Alam S, Colvin J., 2007. PCR-based detection and partial
genome sequencing indicate high genetic diversity in Bangladeshi begomoviruses and their
whitefly vector, *Bemisia tabaci*. *Virus Genes* 34, 373–385.

Moriones E, Navas-Castillo J., 2000. *Tomato yellow leaf curl virus*, an emerging virus complex
causing epidemics worldwide. *Virus Res* 71:123–134.

Mustafa S.I., Schwarzacher T. & Heslop-Harrison J., 2018. Complete mitogenomes from
Kurdistani sheep: abundant centromeric nuclear copies representing diverse ancestors.
Mitochondrial DNA Part A, 1-14.

Navot N, Pichersky E, Zeidan M, Zamir D, Czosnek H., 1991. *Tomato yellow leaf curl virus*: a
whitefly-transmitted geminivirus with a single genomic component. *Virology* 185:151–161.

Novák P., Neumann P., Pech J., Steinhaisl J. & Macas J., 2013. RepeatExplorer: a Galaxy based
web server for genome-wide characterization of eukaryotic repetitive elements from next-
generation sequence reads. *Bioinformatics* 29, 792-793.

Panno, S., Iacono, G., Davino, M., Marchione, S., Zappardo, V., Bella, P., ... & Davino, S., 2016.
First report of *Tomato leaf curl New Delhi virus* affecting zucchini squash in an important
horticultural area of southern Italy. *New Dis. Rep*, *33*(6), 2044-0588.

Pico B, Diez MJ, Nuez F., 1996. Viral diseases causing the greatest economic losses to the
tomato crop. II. The *Tomato yellow leaf curl virus*. *Sci Hort* 67:151–196.

Polston JE, Anderson PK., 1997. The emergence of white fly transmitted geminiviruses in
tomato in the Western Hemisphere. *Plant Dis* 81:1358–1369.

Reddy, R. C., Colvin, J., Muniyappa, V., & Seal, S., 2005. Diversity and distribution of
begomoviruses infecting tomato in India. *Archives of virology*, *150*(5), 845-867.

Sadeq Al-Kuwaiti, N. A., 2013. Molecular characterization of viruses infecting potato and
vegetables in Iraq (Doctoral dissertation, University of Greenwich).

Salati, R., Nahkla, M. K., Rojas, M. R., Guzman, P., Jaquez, J., Maxwell, D. P., & Gilbertson, R.
L., 2002. *Tomato yellow leaf curl virus* in the Dominican Republic: characterization of an
infectious clone, virus monitoring in whiteflies, and identification of reservoir
hosts. *Phytopathology*, *92*(5), 487-496.

Saunders, K., Norman, A., Gucciardo, S., & Stanley, J., 2004. The DNA β satellite component
associated with ageratum yellow vein disease encodes an essential pathogenicity protein
(βC1). *Virology*, *324*(1), 37-47.

Tamura K., Stecher G., Peterson D., Filipski A. & Kumar S., 2013. MEGA6: molecular
evolutionary genetics analysis version 6.0. *Molecular Biology and Evolution* 30, 2725- 2729.

Varsani, A., Roumagnac, P., Fuchs, M., Navas-Castillo, J., Moriones, E., Idris, A., ... & Martin,
D. P., 2017. *Capulavirus* and *Grablovirus*: two new genera in the family
Geminiviridae. *Archives of Virology*, *162*(6), 1819-1831.

Yang, Z., Mammel, M., Papafragkou, E., Hida, K., Elkins, C. A., & Kulka, M., 2017.
Application of next generation sequencing toward sensitive detection of enteric viruses isolated
from celery samples as an example of produce. *International journal of food microbiology*, *261*,
73-81.

Yang, X., Xie, Y., Raja, P., Li, S., Wolf, J. N., Shen, Q., *et al*., 2011. Suppression of
methylation-mediated transcriptional gene silencing by βC1-SAHH protein interaction during
geminivirus-betasatellite infection. *PLoS Pathog*. 7:e1002329. doi:
10.1371/journal.ppat.1002329.

Zhou, X., 2013. Advances in understanding begomovirus satellites. *Annual review of*
phytopathology, *51*.

Figures

Fig. 1. A: Gel electrophoresis of genomic DNA of TYLCV-infected tomato var. Oula F1 (T), in comparison with a control genomic DNA (C). B: Raw reads mapped to complete sequence of TYLCV-Najaf isolate (2770 nt), the assembled reads were 65,303 that used to calculate genome proportion and copy numbers. C: RepeatExplorer report showed cluster number 76 for TYLCV sequence (2763 nt) that picked up as a putative satellite due to tandemly arranged feature within raw reads.

Fig. 2. Phylogenetic tree of 13 full lengths TYLCV isolates showing close relationship between Najaf and Kahnooj isolates. The tree was generated using Geneious tree builder with bootstrape analyses and 50% support, the out group member was *Tomato leaf curl Sudan virus*.

309   Words count: 3566