Biome-scale woody encroachment threatens conservation potential and sustainability of U.S. rangelands

Scott L. Morford[a,b,*], Brady W. Allred[a,b], Dirac Twidwell[c], Matthew O. Jones[a,b], Jeremy D. Maestas[d], David E. Naugle[b]

[a] Numerical Terradynamic Simulation Group, University of Montana, Missoula, MT 59812, USA

[b] University of Montana, W. A. Franke College of Forestry and Conservation, Missoula, MT 59812, USA

[c] Department of Agronomy and Horticulture, University of Nebraska–Lincoln, Lincoln, NE 68588, USA

[d] US Department of Agriculture, Natural Resources Conservation Service, Portland, OR 97232, USA

*Corresponding Author: Scott L. Morford, Numerical Terradynamic Simulation Group. ISB 415, University of Montana, Missoula MT, 59812. (406) 461-4910.

Email: scott.morford@ntsg.umt.edu

Author Contributions: B.A, D.T., and D.N. conceived the project; B.A. and D.N. secured funding. S.M. and B.A. designed the analysis with additional input from D.T. and M.J. S.M. developed the modeling framework and performed the analysis. All authors assisted in evaluating and interpreting analysis results. S.M. wrote the paper with contributions from B.A., D.N., and D.T, and M.J. All authors reviewed and provided input on the manuscript draft

Competing Interest Statement: None.

Classification: Biological Sciences: Sustainability science

Keywords: herbaceous production, yield gap, global change, agricultural land use, afforestation

Preprint server: bioarxiv.org;

This PDF file includes:

> Main Text
> Figures 1 to 4

Abstract

Rangelands of the United States provide ecosystem services that sustain biodiversity
and rural economies. Native tree encroachment is a recognized and long-standing
conservation challenge to these landscapes, but its impact is often overlooked due to the
slow pace of tree invasions and the positive public perception of trees. Here we show
that tree encroachment is a dominant change agent in U.S. rangelands; tree cover has
increased by more than 77,000 km2 over 30 years, and more than 25% of U.S.
rangelands are now experiencing sustained tree cover expansion. Further, we use
machine learning methods to estimate the potential herbaceous production (forage) lost
to tree encroachment. Since 1990 roughly 300 Tg of herbaceous biomass has been lost,
totaling some $5 billion in foregone revenue to agricultural producers. These results
suggest that tree encroachment is similar in scale and magnitude to row-crop
conversion, another primary cause of rangeland loss in the U.S. Prioritizing conservation
efforts to prevent tree encroachment in rangelands can bolster ecosystem and economic
sustainability of these landscapes, particularly among privately-owned lands threatened
by land-use conversion.

Significance Statement

Rangeland biomes are being rapidly lost and degraded due to expansion of row-crop
agriculture, the built environment, and a proliferation of exotic plants. Tree expansion in
rangelands exacerbates these losses but has been difficult to track at large scales due
to the slow, incremental pace of tree spread. Here we use improved satellite technology
to map the scale of tree cover expansion and its impact on forage production across
U.S. rangelands. We reveal that the pace of tree expansion is similar in magnitude to
grassland cultivation (tillage) and establish that forage losses from tree encroachment
may threaten the economic sustainability and conservation value of working rangelands
that are critical habitat to some of the most imperiled species in North America.

Main Text

Introduction

Native trees are invading global rangeland biomes (1, 2). Fire suppression, livestock overgrazing, nutrient pollution, and increasing CO_2 emissions (3) contribute to extensive tree encroachment to rangelands (here defined as grasslands, shrublands, and open woodlands). These invasions exacerbate already substantial losses of rangelands to cropland conversion (4) and threaten the resilience and conservation potential of grazing land biomes worldwide (5–7).

Tree encroachment negatively impacts a myriad of rangeland ecosystem services, including water storage and supply (8), habitat quality (9) biodiversity (10, 11), and land surface albedo (12, 13). Most importantly, tree cover is a key regulator of herbaceous production (the combined production of grasses and forbs; i.e., forage) upon which both wildlife and livestock depend (10). Substantial effort has been invested in disentangling the complex relationship between herbaceous production and tree cover (10, 14, 15). Yet, critical knowledge gaps still exist for how herbaceous production responds to tree cover expansion at landscape to biome scales.

Identifying where tree encroachment impacts landscapes and ecosystem services has traditionally relied on methods such as historical image analysis, paired site investigations, and evaluations of pollen records (16–18). These methods have documented substantial tree canopy expansion in Africa, the Americas, Asia, and Australia (1). Unfortunately, critical intelligence on patterns, pace, and magnitude of tree invasions to rangelands remains absent at large scales. While satellite remote sensing tools are applied globally to track forest cover changes, these analyses typically lack the sensitivity to track tree cover gains in rangelands (19) or omit analysis of tree cover in rangelands altogether (20).

Two-thirds of western U.S. rangelands are under private ownership, and conservation of these lands requires addressing linkages between ecological and economic sustainability. In particular, declining ranch incomes hastens the conversion of intact grasslands to row-crop agriculture, energy production, and subdivision (21, 22). For example, row-crop conversion has accelerated in recent years due to commodity market instability and declining economic returns from livestock production (23). Tree

encroachment is thought to exacerbate declines in ranching profitability as invading

trees out-compete and displace grasses and forbs (15). Revealing the economic cost of

tree encroachment on forage production could help motivate livestock producers to

implement proactive, conservation-focused tree management to prevent woody

encroachment. However, scale-appropriate data and tools to evaluate the interaction of

tree expansion and herbaceous production are not yet available.

Recent technological advances in remote sensing collectively suggest large-scale tree

cover expansion in U.S. rangelands (24–26), consistent with decades of empirical

observations (2). Related innovations in tracking annual herbaceous production (27, 28)

now allow for an integrated spatiotemporal analysis across the western United States.

To assess how tree cover expansion impacts herbaceous production, we jointly

analyzed the two across 17 western states for 1990 - 2019. This analysis *1)* provides a

spatial and temporal evaluation of tree encroachment in rangelands and *2)* quantifies the

difference between attainable and actual herbaceous production as a function of tree

cover expansion, a difference termed yield gap (29). Finally, yield gap estimates derived

in this analysis are combined with grazing rental rates from the United States

Department of Agriculture (USDA) to estimate foregone revenue to livestock producers

from tree encroachment.

Results and Discussion

Tree cover gains between 1990 and 2019 in western U.S rangelands were substantial.

Rangelands added 77,323 ± 1,222 km^2 of trees (**Figure 1**), increasing absolute tree

cover from 154,502 km^2 to 231,825 km^2 (**Supplemental Table 1**). Tree cover grew on

average 2,577 km^2 per year, with annual increases observed in all years on record.

The pace of tree encroachment is similar in magnitude to row-crop conversion, another

primary threat to conservation and sustainability of U.S. rangelands. Cropland

conversion rates from 2008 to 2016 in the western U.S. ranged from 1,649 to 4,385 km^2

annually (median = 2,777 km^2) (23) while the median annual loss of rangelands to tree

encroachment was 1,899 km^2 over this same period. Together, these data suggest

rangelands of the western U.S. are being lost at a rate of 0.89 hectares (2.2 acres) per

minute, losses that are 68% higher than estimates based solely on row-crop conversion.

The observed increase in absolute tree cover does not fully convey the impact and

extent that invading trees have on grazing lands. We observe a net sustained tree cover

expansion across roughly 25% of all western U.S.rangelands, illustrating that tree

encroachment is broadly distributed and not limited to a particular geography, climate, or

ownership. Outsized impacts to wildlife habitat (9, 30), hydrology (8), and land-surface

albedo (13) provide compelling evidence for how small increases in tree density impact

the function of U.S rangelands.

Our analysis also revealed sizable losses in potential herbaceous production resulting

from tree cover expansion (**Figure 2**). Production losses totaled 302 ± 30 Tg (dry

biomass) from 1990 through 2019. In 2019, the yield gap totaled 20.0 Tg, representing

5% - 6% of the potential forage production for all western U.S. rangelands (**Figure 2b**).

For context, the 2019 yield gap of 20 Tg is enough forage to sustain 1.5 million bison or

1.9 million cattle.

Spatially, yield gaps are highest in the southern Great Plains, where production potential

is greatest due to favorable climate conditions and where woody encroachment has

expanded most rapidly. However, forage losses in the northern Great Plains and Great

Basin are no less substantial; proportionally, losses in these regions can be similar to or

exceed losses from more productive grazing lands of the southern Great Plains

(**Supplemental Figure 1**). Lost production identified among the Dakotas, Montana,

Nebraska, and Wyoming is particularly concerning because these states are central to

ongoing efforts to protect the northern Great Plains biome's connectivity and biodiversity

and are focal areas for preventing cropland conversion (21, 31). Moreover, these lands

are home to the last remaining large-scale migrations of land mammals in the

contiguous United States (32) and critical habitat for grassland birds, some of the most

imperiled wildlife on the continent (11, 33). California was the only state in our dataset

that did not develop a yield gap over the 30-year record (**Supplemental Table 5**).

To explore how lost herbaceous production may impact ranch economics, we used our

yield gap data in combination with pasture rental rates to estimate lost revenues owing

to tree cover expansion. Pasture rental rates compiled by the USDA (34) represent the

private rate that ranchers pay to lease grazing land, so they are a meaningful proxy for

the going-market value of forage. We found that on an inflation-corrected basis, potential

revenue losses across the 17 states totaled $4.83 ± 0.53 billion over the 30 years

analyzed; losses exceeded $300 million in 2019 and are increasing at a rate of roughly

$11.3 million per year (**Figure 2c**). These coarse estimates suggest that revenue losses

from tree encroachment to rural agricultural economies are at a minimum on the order of

$100s of millions of dollars per year. Moreover, this calculation is likely conservative

because it uses a tree cover basis from 1990 and excludes forage losses due to shrub

encroachment.

Importantly, primary production is not fixed or constant in an era of global environmental

change (35). We observed sizable increases in herbaceous production among

rangelands where tree cover remained constant (averaging 195.7 kg ha^{-1} yr^{-1} more

production in 2019 than 1990), consistent with other studies showing increasing

continental-scale productivity (35–37). Consequently, many regions of the western U.S.

with increasing tree cover saw only moderate losses in total herbaceous production

(averaging 37.7 kg ha^{-1} yr^{-1} less production in 2019 than in 1990; **Supplemental Figure**

2). Investigating local-scale yield gaps in combination with tree encroachment data

provides further clarity regarding the interaction between tree cover and herbaceous

production. Generally, we observed a lag between the inception of tree encroachment

and losses of herbaceous production, consistent with data and theory from invasion

ecology (38). However, in the absence of disturbance or tree management, tree cover

will eventually cross a site-specific threshold where yield gaps rapidly develop (**Figure**

3). For example, at the Loess Canyon site (**Figure 3 - Panel C** and **Figure 4**) we

observed that a herbaceous production yield gap took roughly 20 years to develop (1990

- 2010) as tree cover grew from 2 to 20%. Over the following five years (2010 - 2014) the

yield gap expanded dramatically, resulting in a 30% loss of potential production. These

yield gap data are consistent with field-based investigations in the Loess Canyons (39),

experimental woody-encroachment data from the Konza LTER (**Supplemental Text**),

and literature-based reviews of tree-grass interactions in dryland systems (10).

These site-specific data illustrate both challenges and opportunities to address tree

encroachment in rangelands. The time-lag between when trees colonize and when yield

gaps emerge may promote a perverse incentive among some land managers to tolerate

tree encroachment until an invasion reaches more advanced stages. In these cases,

managing tree encroachment becomes much more expensive and complex due to

removing additional tree biomass and established seed stores (40). Conversely, the time
lag provides an opportunity to implement cost-effective management to prevent further
tree encroachment, productivity decline, and economic harm to agricultural producers,
particularly if pre-emptive management fails to control the reproductive source of tree
encroachment (40).

USDA conservation programs direct nearly $90 million annually in public and private
funds towards tree and brush removal to promote rangeland health (41). However, these
interventions have generally been opportunistic rather than targeted, resulting in little
mitigation of tree invasions at biologically relevant scales (40). Directing these funds
using state-of-the-art targeting, valuation, and monitoring tools (25, 42) will help prioritize
projects that can simultaneously promote sustainable agricultural economics while
maintaining and growing core, intact rangelands at scale. To this end, we include a
simple web-enabled map application that visualizes and investigates yield losses
attributable to tree cover expansion (https://rangelands.app/yield-gap/).

Our modeling approach is not without limitations. The hindcast modeling method used
here does not consider future climate scenarios, so may not accurately inform future
trajectories of production responses to tree encroachment. However, herbaceous
productivity is forecasted to increase across much of western U.S. rangelands through
2100 (37, 43), lending some confidence that our findings are generalizable. Second, our
monetary estimates for lost revenues should be considered preliminary as they do not
consider economic leakage or policy and social barriers that may limit the adoption of
management alternatives at scale. Locally, our estimates for forage value agree well
with independent agricultural economic data (**Supplemental text**). Still, more
sophisticated analyses will be required to assess how managing tree encroachment
benefits ranching economics.

Loss of rangeland production and corresponding impacts to ecosystem services need to
be considered as part of global scientific advocacy efforts calling for the afforestation of
rangelands (44). Based on the current state of climate-vegetation science, increasing
tree cover in rangelands is not a climate mitigation solution (45). Tree encroachment can
increase carbon storage in rangelands (16, 46), but these carbon storage gains are
typically offset by reducing land-surface albedo at mid- to high- latitudes (12, 47). For

example, across the northern Great Plains, tree cover would need to reach 95+% to

contribute to climate cooling (12); such a scenario would result in the wholescale

collapse of the northern Great Plains and its biodiversity therein (4). The most effective

climate mitigation strategy for rangelands remains preventing grassland conversion to

row-crop agriculture (5).

Identifying where tree encroachment creates yield gaps and how forage loss impacts

ranching revenues provides a mechanism to focus private-land conservation efforts on

economically sustainable outcomes that prevent land-use conversion. Herbaceous

production is the cornerstone of food and fiber economics in working rangelands and

evaluating it similar to other agricultural products provides a means to stack market

incentives with other conservation subsidies. Preventing yield gaps in rangelands is a

novel and landscape-scale conservation strategy to benefit nature and society.

Materials and Methods
198
The analysis area included the 17 western-most states in the contiguous United States,

consistent with other data products generated from the Rangelands Analysis Platform

(24). Our calculations only consider production lost to trees and do not include or

consider additional production losses attributable to woody shrubs. The total area

included in this analysis was 4.73 million km^2, comprising 6.9 billion pixels. Model inputs

and outputs consist of annualized data from 1990 to 2019 unless otherwise stated.

Modeling evaluation, data sources, and computational libraries used in this analysis are

provided in the supplemental.

1. Mapping Rangelands

The spatial extent of rangelands in the western United States was determined from

Landfire Biophysical Settings (BPS), version 1.4.0. Landfire BPS represents the

dominant vegetation system on the landscape prior to Euro-American settlement.

Grasslands and shrublands were identified using the BPS categorization grouping

attribute, while open woodlands were selected based on the BPS name component

where 'woodland' was the primary attribute. Map units described as barren were

excluded. Our rangeland classification layer from Landfire BPS was further modified to

exclude crop agricultural and developed lands from the U.S. Geological Survey National

Land Cover Database (2016) the National Agricultural Statistical Service Cropland Data

Layer (NASS CDL). All production and monetary calculations provided in this manuscript

omit riparian and primary forests map units.

2. Tree Cover Modeling

Annual tree cover was taken from the *Rangeland Analysis Platform V2 (RAP)* cover

products (24) for the years 1990 through 2019. These data were processed using the

LandTrendr (LT) segmentation algorithm in Google Earth Engine (GEE) to denoise the

time-series data (48). The LandTrendr parameters used in this analysis are summarized

in **Table S2**.

To remove spurious tree-cover trends attributable to prediction uncertainty or model

error, we applied two independent rules in our tree cover trend analysis. First, for the

analysis to tally a change in pixel tree cover, the absolute difference between the

maximum tree cover and minimum tree cover in the pixel time series had to be greater

than 2.8%, corresponding to the mean absolute error (MAE) of the tree cover product

(24). Second, we evaluated each pixel segment from the LT-GEE segmentation product

to determine if the change in tree cover was an artifact of prediction uncertainty/variance

(See Bayesian approximation discussion in ref. 24). To accomplish this, we performed a

Welch T-test using the tree cover and prediction variance estimates from the first and

last observation in the pixel segment ($n = 4$, two-tailed, p-value $= 0.01$). For tree cover

change to be tallied in a pixel, it had to have one or more segments where the t-value <

p-value. While imperfect, this approach was an effective screening tool to filter change

estimates for pixels with high prediction uncertainty without implementing high

computational cost Monte Carlo simulations.

3. Herbaceous Production & Yield Gap Modeling

We used XGBoost, a supervised machine learning library, to predict herbaceous

biomass production as a function of tree cover and other biophysical variables. Biomass

in our model was derived from total annual herbaceous production reported by the

Rangeland Analysis Platform (27); we term this biomass variable *Observed Production*.

Conceptually and numerically, our analysis used the following general framework

(***equation 1***).

$$\textit{Observed Production(YR)} \ = \ f \, (YR, \ MU, \ LP, \ TC_{(yr),}, \ SP) \qquad \textbf{eqn 1.}$$

In our framework, *YR* represents year, MU = soil map unit, LP = landscape position,

$TC_{(yr)}$ = tree cover for year YR, SP = spatial position/coordinates. Factor variables were

encoded as integers during training and inference. Model variables, source data, and a

rationale for their inclusion are presented in **Supplemental Table 3**.

Model training and inference were performed using the XGBoost (49) implementation

from the open-source H2O.ai library v. To improve computational performance, we

enabled LightGBM emulation and implemented the dart booster drop-out method to limit

overfitting. We used a cartesian grid search for hyper-parameter selection, optimized for

the lowest MAE on 10 million randomly selected records. The optimization (loss) function

used for model training was mean squared error (MSE) with L2 regularization. Our

ensemble used 60 trees with a maximum tree depth of 15. A complete list of model

parameters is included in **Supplemental Table 4**.

Model training: We split our analysis area into 53 units (models) using the spatial extent

of Level-3 EcoRegions developed by the US EPA. The size of the training data for each

unit was limited to 71,582,780 records due to computational limitations; the remaining

training data were used as hold-out (test) data. The models were trained with 3.8 billion

records, randomly sampled from our dataset; the remaining 5.5 billion records were used

for holdout evaluation. Evaluation of the training metrics showed that holdout MAE

values closely matched training MAE values, suggesting the model did not overfit the

data (**Supplemental Figure 3)**.

Model inference: For each pixel-year combination, the XGBoost model generates two

predictions. Prediction one estimates herbaceous production for a given year using the

corresponding tree cover for that year*;* we term this *Estimated Production (**equation 2**)*,

270   *Estimated Production$(YR, TC_{(yr)})$ = E[m| YR, MU, LP, $TC_{(yr)}$, SP]*   **eqn 2.**

271   where ***m*** is the model and *$TC_{(yr)}$* is the tree cover for year *YR*. Prediction two estimates

272   the herbaceous production for the year *YR* using the tree cover from 1990. We term

273   prediction two *Achievable Production (**equation 3**)*, it represents the herbaceous

274   production for year *YR* using 1990 tree-cover.

275   *Achievable Production$(YR, TC_{(1990)})$ = E[m| YR, MU, LP, $TC_{(1990)}$, SP]*   **eqn 3.**

276   The yield gap attributable to tree cover expansion since 1990 is calculated using the

277   difference of these two predictions for each pixel-year combination (**equation 4**). Note

278   that the yield gap can be negative in cases where tree cover declines over time.

279   *Yield Gap(YR) = Achievable Production(YR,TC$_{(1990)}$) –*

280   *Estimated Production(YR,TC$_{(yr)}$)*   **eqn 4.**

281   Prediction error for each pixel was calculated as the absolute difference between

282   *Observed Production and Estimated Production **(equation 5).** Model Error* was

283   represented as the MAE **(equation 6)**, aggregated by year and map unit to provide

284   some localization; *n* in **equation 6** is the number of observations for each MU.

285   *Prediction Error(YR) =*

286   *|Observed Production(YR) – Estimated Production(YR,TC$_{(yr)}$)|*   **eqn 5.**

Model Error(MU | YR) =[∑ *Prediction Error*(MU | YR)] / n   **eqn 6.**

For the analysis, the yield gap is only tallied when *Yield Gap* > *Model Error* for 3 or more

consecutive years. This prevents spikes in yield gap attributable to data noise or

modeling error from being included in lost potential production estimates. *Potential*

Production in our analysis and map products is presented as the *Observed Production*

plus *Yield Gap.* Additional information regarding sensitivity and uncertainty in our model

predicts are presented in the Supplemental.

4. Monetary calculations

We calculated the value of forage lost to tree cover expansion using our yield gap

estimates with grazing rental rates from USDA National Agricultural Statistics Service

(NASS) (34). We used the annual state-level data reported as dollars per animal unit

month (AUM) because these records were the most complete for our analysis (1990 -

2019). Military and protected non-grazing lands (e.g. National Parks) were excluded

from the monetization calculations, but other federal grazing lands (i.e., BLM and USFS

allotments) were included. Missing data were imputed using a linear-interpolation of the

time series data. The value of forage production on a mass basis was calculated as

follows:

$$\frac{\$}{kg\ production} = \frac{\$}{AUM} \times \frac{AUM}{kg\ forage} \times \frac{kg\ forage}{kg\ production}$$

eqn. 7

In this calculation, dollars per AUM was taken from NASS tables; AUM per kg forage

was assumed to equal to 1 / 344.73, equivalent to 760 lbs per AUM; kg forage per kg

production represents the utilization factor (*UF*), and was assumed to equal 0.40 across

all locations. The selection of the UF was data-driven: it was derived from an analysis of

available NASS rental rates in combination with herbaceous production data (***equation***

8). The general form of the equation was:

$$UF = \frac{kg\ forage}{AUM} \times \frac{\$}{acre} \times \frac{AUM}{\$} \times \frac{acre}{kg\ production}$$

eqn. 8

where *$/acre* and *$/AUM* represent rental rates extracted from NASS data for years

2008 - 2019. Finally, dollar estimates were inflation corrected to 2019 dollars using the

consumer price index from the Bureau of Labor Statistics (50). A summary of annual

state-level yield gap and forage value data is provided in **Supplemental Table 5**.

The $90 million figure presented as tree removal conservation expenditures in the text

reflects 2019 EQIP spending on *Practice 314* plus 25% private matching funds.

5. Production trends and local scale production responses to tree encroachment.

Production trends presented in **Figure 3** and **Supplemental Figure 2** represent

Observed Production values extracted directly from the Rangeland Analysis Platform

and were independent of yield gap calculations. In both cases, the data presented is

based on a long-term pixel-wise trend analysis using a linear fit in Google Earth Engine.

Production trends from **Supplemental Figure 2** have been masked to rangelands with

increasing, stable, and decreasing tree cover using the methods described in sections 1

and 2 of the methods.

Acknowledgments

This work was made possible by the USDA-Natural Resources Conservation Service's

(NRCS) Conservation Effects Assessment Project - Wildlife Component, the NRCS's

Working Lands for Wildlife, and the Bureau of Land Management. Model data will be

made available via our web application and archived at Zenodo. Caleb P. Roberts

contributed to this draft. Code to replicate the analysis will be archived on Zenodo. Any

use of trade, firm, or product names is for descriptive purposes only and does not imply

endorsement by the U.S. Government.

References

1. L. L. Nackley, A. G. West, A. L. Skowno, W. J. Bond, The Nebulous Ecology of Native Invasions. *Trends Ecol. Evol.* **32**, 814–824 (2017).
2. S. R. Archer, *et al.*, "Brush Management as a Rangeland Conservation Strategy: A Critical Evaluation" in *Conservation Benefits of Rangeland Practices: Assessment, Recommendations, and Knowledge Gaps.*, D. D. Briske, Ed. (USDA Natural Resources Conservation Service, 2011), p. 66.
3. G. P. Asner, A. J. Elmore, L. P. Olander, R. E. Martin, A. T. Harris, Grazing systems, ecosystem responses, and global change. 45 (2004).
4. J. M. Hoekstra, T. M. Boucher, T. H. Ricketts, C. Roberts, Confronting a biome crisis: global disparities of habitat loss and protection: Confronting a biome crisis. *Ecol. Lett.* **8**, 23–29 (2004).
5. J. E. Fargione, *et al.*, Natural climate solutions for the United States. *Sci. Adv.* **4**, eaat1869 (2018).
6. C. Kremen, A. M. Merenlender, Landscapes that work for biodiversity and people. *Science* **362**, eaau6020 (2018).
7. J. W. Veldman, *et al.*, Where Tree Planting and Forest Expansion are Bad for Biodiversity and Ecosystem Services. *BioScience* **65**, 1011–1018 (2015).
8. T. E. Huxman, *et al.*, Ecohydrological implications of woody plant encroachment. *Ecology* **86**, 308–319 (2005).
9. S. Baruch-Mordo, *et al.*, Saving sage-grouse from the trees: A proactive solution to reducing a key threat to a candidate species. *Biol. Conserv.* **167**, 233–241 (2013).
10. S. R. Archer, K. I. Predick, An ecosystem services perspective on brush management: research priorities for competing land-use objectives. *J. Ecol.*, 14 (2014).
11. K. V. Rosenberg, *et al.*, Decline of the North American avifauna. *Science* **366**, 120–124 (2019).
12. P. M. Mykleby, P. K. Snyder, T. E. Twine, Quantifying the trade-off between carbon sequestration and albedo in midlatitude and high-latitude North American forests. *Geophys. Res. Lett.* **44**, 2493–2501 (2017).
13. J. Ge, C. Zou, Impacts of woody plant encroachment on regional climate in the southern Great Plains of the United States. *J. Geophys. Res. Atmospheres* **118**, 9093–9104 (2013).
14. R. J. Scholes, Convex Relationships in Ecosystems Containing Mixtures of Trees and Grass. *Environ. Resour. Econ.* **26**, 559–574 (2003).
15. J. D. Anadon, O. E. Sala, B. L. Turner, E. M. Bennett, Effect of woody-plant encroachment on livestock production in North and South America. *Proc. Natl. Acad. Sci.* **111**, 12948–12953 (2014).
16. G. P. Asner, S. Archer, R. F. Hughes, R. J. Ansley, C. A. Wessman, Net changes in regional woody vegetation cover and carbon storage in Texas Drylands, 1937–1999. *Glob. Change Biol.* **9**, 316–335 (2003).
17. K. C. Lubetkin, A. L. Westerling, L. M. Kueppers, Climate and landscape drive the pace and pattern of conifer encroachment into subalpine meadows. *Ecol. Appl.* **27**, 1876–1887

(2017).
18. S. Archer, C. Scifres, C. R. Bassham, R. Maggio, Autogenic Succession in a Subtropical
Savanna: Conversion of Grassland to Thorn Woodland. *Ecol. Monogr.* **58**, 111–127 (1988).
19. M. C. Hansen, *et al.*, High-Resolution Global Maps of 21st-Century Forest Cover Change.
Science **342**, 850–853 (2013).
20. M. Rigge, *et al.*, Quantifying Western U.S. Rangelands as Fractional Components with
Multi-Resolution Remote Sensing and In Situ Data. *Remote Sens.* **12**, 412 (2020).
21. T. J. Lark, Protecting our prairies: Research and policy actions for conserving America's
grasslands. *Land Use Policy* **97**, 104727 (2020).
22. B. W. Allred, *et al.*, Ecosystem services lost to oil and gas in North America. *Science* **348**,
401–402 (2015).
23. T. J. Lark, S. A. Spawn, M. Bougie, H. K. Gibbs, Cropland expansion in the United States
produces marginal yields at high costs to wildlife. *Nat. Commun.* **11** (2020).
24. B. W. Allred, *et al.*, Improving Landsat predictions of rangeland fractional cover with
multitask learning and uncertainty. *Methods Ecol. Evol.* **0** (2021).
25. M. O. Jones, *et al.*, Beyond Inventories: Emergence of a New Era in Rangeland Monitoring.
Rangel. Ecol. Manag. **73**, 577–583 (2020).
26. S. K. Filippelli, *et al.*, Monitoring pinyon-juniper cover and aboveground biomass across the
Great Basin. *Environ. Res. Lett.* **15**, 025004 (2020).
27. M. O. Jones, *et al.*, Annual and 16-day rangeland production estimates for the western
United States. *bioRxiv*, 2020.11.06.343038 (2020).
28. N. P. Robinson, *et al.*, Rangeland Productivity Partitioned to Sub-Pixel Plant Functional
Types. *Remote Sens.* **11**, 1427 (2019).
29. M. K. van Ittersum, *et al.*, Yield gap analysis with local to global relevance—A review. *Field
Crops Res.*, 14 (2013).
30. B. T. Hamilton, B. L. Roeder, M. A. Horner, Effects of Sagebrush Restoration and Conifer
Encroachment on Small Mammal Diversity in Sagebrush Ecosystem. 10 (2019).
31. J. H. Haggerty, M. Auger, K. Epstein, Ranching Sustainability in the Northern Great Plains:
An Appraisal of Local Perspectives. *Rangelands* **40**, 83–91 (2018).
32. K. Joly, *et al.*, Longest terrestrial migrations and movements around the world. *Sci. Rep.* **9**,
15333 (2019).
33. L. A. Brennan, Kublesky, William P., North American grassland Birds: an unfolding
conservation crisis? *J. Wildl. Manag.* **69**, 1–13 (2005).
34. United States Department of Agriculture, *National Agriculture Statistics Service Quick Stats
Database* (2020).
35. J. A. Hicke, *et al.*, Trends in North American net primary productivity derived from satellite
observations, 1982-1998. *Glob. Biogeochem. Cycles* **16**, 2-1-2–14 (2002).
36. M. C. Reeves, B. B. Hanberry, H. Wilmer, N. E. Kaplan, W. K. Lauenroth, An Assessment of
Production Trends on the Great Plains from 1984 to 2017. *Rangel. Ecol. Manag.*,
S1550742420300117 (2020).
37. R. B. Boone, R. T. Conant, J. Sircely, P. K. Thornton, M. Herrero, Climate change impacts on
selected global rangeland ecosystem services. *Glob. Change Biol.* **24**, 1382–1393 (2018).
38. J. A. Crooks, Lag times and exotic species: The ecology and management of biological
invasions in slow-motion1. *Écoscience* **12**, 316–329 (2005).
39. C. Bielski, R. Scholtz, V. M. Donovan, C. R. Allen, D. Twidwell, Overcoming an "irreversible"
threshold: a 15-year fire experiment. *J. Environ. Manage.* **(Accepted)**.

40. C. P. Roberts, D. R. Uden, C. R. Allen, D. Twidwell, Doublethink and scale mismatch polarize
policies for an invasive tree. *PLOS ONE* **13**, e0189733 (2018).
41. United States Department of Agriculture, *EQIP Data Dashboards*.
42. D. R. Uden, *et al.*, Spatial Imaging and Screening for Regime Shifts. *Front. Ecol. Evol.* **7**, 407
(2019).
43. M. C. Reeves, A. L. Moreno, K. E. Bagne, S. W. Running, Estimating climate change effects
on net primary production of rangelands in the United States. *Clim. Change* **126**, 429–442
(2014).
44. J.-F. Bastin, *et al.*, The global tree restoration potential. *Science* **365**, 76–79 (2019).
45. P. Friedlingstein, M. Allen, J. G. Canadell, G. P. Peters, S. I. Seneviratne, Comment on "The
global tree restoration potential." *Science* **366**, eaay8060 (2019).
46. R. K. Connell, J. B. Nippert, J. M. Blair, Three Decades of Divergent Land Use and Plant
Community Change Alters Soil C and N Content in Tallgrass Prairie. *J. Geophys. Res.*
Biogeosciences **125** (2020).
47. G. Bala, *et al.*, Combined climate and carbon-cycle effects of large-scale deforestation.
Proc. Natl. Acad. Sci. **104**, 6550–6555 (2007).
48. R. Kennedy, *et al.*, Implementation of the LandTrendr Algorithm on Google Earth Engine.
Remote Sens. **10**, 691 (2018).
49. T. Chen, C. Guestrin, XGBoost: A Scalable Tree Boosting System in *Proceedings of the 22nd*
ACM SIGKDD International Conference on Knowledge Discovery and Data Mining, (ACM,
2016), pp. 785–794.
50. U.S. Bureau of Labor Statistics, *Consumer Price Index (CPI) Database* (2020).
https://www.bls.gov/cpi/, [Accessed Feb 16, 2021].

Figures and Tables

Figure 1. Tree cover expansion in western U.S. rangelands (1990 - 2019). a) Net tree canopy
expansion in rangelands shown in green; converted agricultural lands and built environment
shown in white. b) Cumulative net tree cover expansion; error bands represent the cumulative
95% prediction interval. c) Annual net tree cover change; error bands represent the 95%
prediction interval. The red time-series represents the rate of agricultural conversion for 2008 -
2016 from ref. 23.

Figure 2. Herbaceous production yield gap attributable to tree cover expansion: 1990 - 2019. a)
Map of 2019 yield gap; converted agricultural lands and built environment shown in white. b)
Total annual yield gap; includes dry herbaceous biomass (grass and forb); c) Annual lost revenue
to ranching operations from tree cover expansion in the western United States.

Figure 3. Localized relationships between tree cover change and herbaceous production from years 1990 - 2019 for five diverse locations across U.S rangelands. Impacts on herbaceous production from tree encroachment vary by location due to local biophysical conditions and climate. These data are calculated at the county scale and are truncated at ±20% to emphasize dominant threshold behavior.

Figure 4. Local-scale evaluation of tree encroachment and yield gap development in the Loess
Canyons, Nebraska (W100.44° N40.95°). Herbaceous production decreased by 34% percent
between 1990 and 2019. **(Top)** Time series aerial imagery shows the dramatic expansion of
Eastern Red Cedar (*Juniperus virginiana*). (**Bottom**) Time-series and spatial trends in achievable
and observed forage production show the forage yield gap's evolution as a function of tree cover
expansion.