Litter commensal bacteria can limit the horizontal gene transfer of antimicrobial

resistance to *Salmonella* in chickens

Running Title: Commensal bacteria reduce HGT

Adelumola Oladeinde[1], Zaid Abdo[2], Benjamin Zwirzitz[3¶], Reed Woyda[2¶], Steven M.

Lakin[2¶], Maximilian O. Press[4¶*], Nelson A. Cox[1], Jesse C. Thomas IV[6], Torey Looft[7],

Michael J. Rothrock[1], Gregory Zock[8], Jodie Plumblee Lawrence[1], Denice Cudnik[1],

Casey Ritz[8] and Samuel E. Aggrey[8], Ivan Liachko[4], Jonas R. Grove[4], Crystal Wiersma[2]

Authors' Affiliation

[1]U.S. National Poultry Research Center, USDA-ARS, Athens, GA, USA. [2]Department of

Microbiology, Immunology and Pathology, Colorado State University, Fort Collins,

Colorado, USA. [3]Institute of Food Science, University of Natural Resources and Life

Sciences, Vienna, Austria. [4]Phase Genomics Inc, Seattle, WA, 98109, USA. [5]Office of

National Programs, USDA-ARS, Beltsville, Maryland, USA. [6]Division of STD Prevention,

National Center for HIV/AIDS, Viral Hepatitis, STD and TB Prevention, Center for

Disease and Control, Atlanta, Georgia, USA. [7]National Animal Disease Center, USDA-

ARS, Ames, IA, USA,[8]Poultry Science Dept, University of Georgia, Athens, GA, USA

Corresponding authors ade.oladeinde@usda.gov; zaid.abdo@colostate.edu

[¶]These authors contributed equally to this work.

*Present address: Maximilian O. Press, Inscripta, Inc., Boulder, Colorado, USA.

Abstract

Fostering a 'balanced' gut microbiome through the administration of beneficial microbes that can competitively exclude pathogens has gained a lot of attention and use in human and animal medicine. However, little is known about how microbes affect the horizontal gene transfer of antimicrobial resistance (AMR). To shed more light on this question, we challenged neonatal broiler chicks raised on reused broiler chicken litter – a complex environment made up of decomposing pine shavings, feces, uric acid, feathers, and feed, with *Salmonella* Heidelberg (*S.* Heidelberg), a model pathogen. Neonatal chicks challenged with *S.* Heidelberg and raised on reused litter were more resistant to *S.* Heidelberg cecal colonization than chicks grown on fresh litter. Furthermore, chicks grown on reused litter were at a lower risk of colonization with *S.* Heidelberg strains that encoded AMR on IncI1 plasmids. We used 16S rRNA gene sequencing and shotgun metagenomics to show that the major difference between chicks grown on fresh litter and reused litter was the microbiome harbored in the litter and ceca. The microbiome of reused litter samples was more uniform and enriched in functional pathways related to the biosynthesis of organic and antimicrobial molecules than fresh litter samples. We found that *E. coli* was the main reservoir of plasmids encoding AMR and that the IncI1 plasmid was maintained at a significantly lower copy per cell in reused litter compared to fresh litter. These findings support the notion that commensal bacteria play an integral role in the horizontal transfer of plasmids encoding AMR to pathogens like *Salmonella*.

Importance/Significance

Antimicrobial resistance spread is a worldwide health challenge, stemming in large part, from the ability of microorganisms to share their genetic material through horizontal gene transfer. To address this issue, many countries and international organization have adopted a One health approach to curtail the proliferation of antimicrobial resistant bacteria. This includes the removal and reduction of antibiotics used in food animal production and the development of alternatives to antibiotics. However, there is still a significant knowledge gap in our understanding of how resistance spreads in the absence of antibiotic selection and the role commensal bacteria play in reducing antibiotic resistance transfer. In this study, we show that commensal bacteria play a key role in reducing the horizontal gene transfer of antibiotic resistance to *Salmonella* and provide the identity of the bacterial species that potentially perform this function in broiler chickens and also postulate the mechanism involved.

Introduction

Horizontal gene transfer (HGT) is recognized as the main mechanism by which bacteria

acquire antimicrobial resistance (AMR) and exposure to antibiotics has been shown to

drive antimicrobial resistance gene (ARG) transfer [1]. Consequently, the rise in AMR in

bacteria from hospital settings has been linked to the overuse of antibiotics in humans

and their use in food animal production [2, 3]. These public health concerns have led to

a reduction in antibiotics used for raising food animals [4, 5], including a ban on

antibiotics' use in Europe [6]. We previously showed that neonatal broiler chicks

challenged with a nalidixic acid resistant *S*. Heidelberg strain and raised antibiotic-free

on fresh litter composed of pine shavings were colonized at a high rate with *S*.

Heidelberg strains that harbored Incl1 plasmids encoding AMR [7]. We selected *S*.

Heidelberg as the model pathogen because of its promiscuity to plasmids carrying AMR

[8, 9].

There is limited research on if and how commensal bacteria reduce AMR in food-

borne pathogens [10-16], such as *Salmonella enterica*, and the role it plays in AMR

reduction and transfer. Therefore, the goal of this study was to determine the role

commensal bacteria play in limiting *Salmonella enterica* from acquiring AMR and to

provide information on the bacterial species and mechanism involved. To do this, we

compared the dynamics of AMR transfer in neonatal broiler chicks raised on fresh litter

to chicks raised on reused litter. We chose reused litter because it is a complex

environment made up of decomposing plant-based bedding (e.g., wood shavings,

sawdust and rice or peanut hulls) mixed with chicken feces, uric acid, feathers, feed,

insects, and other broiler-sourced materials. Therefore, reused litter carries a unique

and complex population of bacteria, fungi, and viruses [17-19], interacting with various

forms of eukaryotes, making it a suitable environment to study competitive exclusion.

Broiler chickens are commonly raised on litter but how litter is managed differs between

countries, producers, and farmers. For instance, the practice of reusing litter over

multiple flocks of broiler chickens is a widespread practice in the US and Brazil while

Canada [20] and Europe [21] recommend fresh litter bedding for every flock. One

argument against litter reuse is that it harbors pathogenic bacteria that can be

transferred to the next flock [22]. Contrastingly, proponents of litter reuse argue that it

confers competitive exclusion against pathogens when effectively managed and it is

cost-effective [23-26]. Beyond broiler chicken production, competitive exclusion by

commensal bacteria has received enormous attention in the 20th century, resulting in

the bloom of commercially marketed probiotics [27] and the underlying theory behind

the application of fecal microbiota transplantation in human medicine [28].

Results

S. **Heidelberg abundance and prevalence in ceca and litter.**

We confirmed that neonatal chicks were *Salmonella*-free by testing the chick pads used

for transportation from the hatchery for *Salmonella*. None of the chick pads were

positive for *Salmonella.* The reused litter for this study was confirmed to be *Salmonella*-

free in earlier studies [29, 30]. To determine if the microbiome offers protection against

Salmonella colonization, we challenged neonatal broiler chicks with a nalidixic resistant

(nal^R) strain of *S.* Heidelberg and raised the chicks on either fresh pine shavings (here

after referred to as fresh litter) (n = 75) or reused litter (litter previously used to raise

three flocks of broiler chickens) (n =75). Chicks were challenged either through oral

gavage (n = 25) , cloacal inoculation (n =25), or by the seeder method (i.e., a few chicks

[n = 5] were challenged orally and comingled with unchallenged chicks [n = 20]) [7].

Unchallenged chicks on fresh (n =25) and reused litter (n =25) were used as controls.

Broiler chicks on fresh and reused litter were housed separately for 14 days (four

separate pens for each treatment in a house) and chicks were not administered any

medication or antibiotics for the duration of the study.

Afterwards, we determined the concentration of nal^R S. Heidelberg in the ceca and

litter of chicks 14 days after they were challenged. Average nal^R S. Heidelberg

concentration in the ceca (n=10 for oral and cloacal, n=15 for seeder) was not

significantly different between chicks raised on fresh litter compared to reused litter (W-

statistic, P. value; oral = 41, 0.15; cloacal= 41, 0.96; seeder = 5, 1.00) (Fig.1a).

However, the ceca of chicks raised on fresh litter were more likely to be positive for nal^R

S. Heidelberg (X_2= 16.07, df = 2, P = 0.0003), than chicks on reused litter. The

percentage of cecal samples positive for nal^R S. Heidelberg was 100%, 100% and 50%,

respectively for oral, cloacal and seeder chicks raised on fresh litter, compared to 80%,

100% and 33.3 %, for chicks raised on reused litter. For the seeder treatment on fresh

litter, one seeder was lost due to premature mortality and three of the four seeders were

positive for nal^R S. Heidelberg, while the ceca of four of the ten uninoculated contact

chicks were positive. In contrast, the ceca of the five orally challenged seeders raised

on reused litter were positive for nal^R S. Heidelberg, but all contact chicks were negative

(n = 10).

For the litter, oral and seeder treatments carried higher levels of nal^R S. Heidelberg

in fresh litter compared to reused litter, while cloacally inoculated chicks had higher

levels in reused litter compared to fresh litter (Fig.1b). All litter samples from challenged

chicks were positive for nal^R S. Heidelberg by direct culture or after enriching in buffered

peptone water (BPW), except for one reused litter sample from the seeder treatment

that was negative. For unchallenged chicks used as controls, one litter sample from

fresh and reused litter tested positive after enrichment in BPW. These data suggests

that litter age/type did not have a significant effect on the abundance of nal^R S.

Heidelberg in the ceca, but chicks raised on reused litter (66%) had a lower *Salmonella*

positivity rate compared to chicks on fresh litter (79%).

Broiler litter age/type affected the horizontal transfer of AMR.

Next, we questioned if the litter plays a role in the HGT of AMR. First, we performed

antibiotic susceptibility testing (AST) on S. Heidelberg isolates recovered from the ceca

and litter of broiler chicks raised on fresh litter (n=158) and reused litter (n=141). As

expected, all S. Heidelberg isolates were resistant to nalidixic acid (Fig. 2). Thirty-one

percent of isolates from the ceca (37/118) and 37.5 % from the litter (15/40) of chicks on

fresh litter acquired resistance to gentamicin, tetracycline, and streptomycin [7] (Fig.

2a). Contrastingly, 3% of S. Heidelberg isolates from the ceca of chicks on reused litter

acquired resistance to either cefoxitin (1/114) or tetracycline (2/114), while ~18% of

isolates from reused litter acquired resistance to either ciprofloxacin (1/27), tetracycline

(2/27) or streptomycin (2/27) (Fig. 2a). The percentage of S. Heidelberg isolates that

acquired AMR in fresh litter differed by the route of inoculation [7]. For instance, 40% of

isolates from orally challenged chicks acquired AMR compared to 24% for cloacal [7].

We determined if *S*. Heidelberg isolates recovered from chicks raised on fresh or

reused litter differed in their core genome or if they acquired new mobile elements like

plasmids. To answer this question, we performed whole genome sequence (WGS)

analysis on *S*. Heidelberg isolates (n=29) recovered from the ceca and litter of chicks

that were cloacally challenged. We chose cloacal samples because only this route of

challenge resulted in 100% colonization of the ceca of chicks raised on fresh and

reused litter. There was no difference in the core genome multilocus sequence types

(cgMLST) of evolved *S*. Heidelberg isolates from fresh and reused litter, however, 28 of

the 29 evolved isolates had a different cgMLST from the ancestral nal^R *S*. Heidelberg

(Fig. 2b). We used the genomes sequenced to construct a maximum likelihood (ML)

tree based on the pangenome and mutations of *S*. Heidelberg strains recovered. The

core genome (genes present in $\geq$ 95% of the strains) and accessory genome (genes

present in $<$ 95% of the strains) was composed of 4,373 and 356 genes, respectively.

After removing 10 isolates with identical mutations, a total of 118 informative sites

(SNPs and indels) was used for ML SNP tree construction.

The three constructed ML trees did not group isolates by litter age/type (fresh

versus reused) (Fig. 2b, Fig. S1a and b), but *S*. Heidelberg isolates that acquired IncI1

plasmids (plasmid MLST 26; hereafter referred to as IncI1-pST26) (n=5) formed a

separate clade on the accessory genome tree (Fig. 2b). The IncI1-pST26 plasmid

harbored the aminoglycoside (*aadA1*), tetracycline (*tetA*) and mercury (*mer* operon)

resistance genes [7] (Fig. S2). Isolates from the ceca (n=3) and litter (n=1) of chicks on

fresh litter that harbored IncI1-pST26 displayed the antibiotic resistance phenotype

predicted by WGS (Fig. 2b). Contrastingly, one isolate from reused litter carried IncI1-

pST26 but was found to be susceptible to all antibiotics tested except for the ancestral

nalidixic resistance. These results indicated that chicks grown on reused litter are less

likely to carry *S*. Heidelberg isolates harboring AMR compared to chicks on fresh litter.

The microbiome of neonatal chicks differed for fresh and reused litter.

We used 16S rRNA gene sequencing to examine the differences in the microbiome

between neonatal chicks raised on fresh and reused litter. The beta diversity of the

cecal (n = 59) and litter (n = 22) microbiome of chicks was significantly different

between fresh and reused litter (ceca, P = 0.0002; litter, P = 0.0002) (Fig. 3a).

Furthermore, the route of inoculation used for *S*. Heidelberg affected the beta diversity

of the ceca and litter for fresh and reused litter (P < 0.05). For example, chicks

challenged orally (n = 10) or through the cloaca (n =10) and raised on fresh litter

harbored a significantly different beta diversity than uninoculated chicks (n = 10) in the

ceca (P < 0.002) (Fig. 3a). Likewise, the beta diversity of the ceca of chicks challenged

orally (n = 10) differed from cloacal (n = 9) and uninoculated controls (n = 10) for reused

litter (P < 0.05). In the litter, pair wise comparisons between routes of inoculation were

not significant (P > 0.05) for beta diversity. Nevertheless, fresh litter samples (n = 3)

from orally challenged chicks clustered next to litter samples (n = 3) from seeder

treatments, while litter samples (n = 2) from cloacally inoculated chicks were different

from oral and seeder treatments (Fig. 3a). For reused litter samples, litter from orally (n

= 3) and cloacally (n = 3) challenged chicks clustered together, while litter (n = 2) from

seeder treatments was similar to litter samples (n = 3) from control chicks. Lastly, the

bacterial community structure (assessed using beta dispersion) of reused litter was less

variable than fresh litter (P = 0.001), however there was no difference in variability for

the ceca ($P = 0.15$). Moreover, the route of challenge affected the bacterial community

beta dispersion in the ceca and litter of chicks on fresh and reused litter ($P = 0.001$).

The alpha diversity (i.e., the number of amplicon sequence variants (ASVs)) of the

microbiome present in the ceca and litter of chicks on reused litter was higher for the

Observed species measure of diversity (ceca, $P < 0.001$; litter, $P = 0.001$) but not for

Shannon (ceca, $P = 0.25$; litter, $P = 0.05$) and Simpson (ceca, $P = 0.97$; litter, $P = 0.28$)

indices. The route of *S.* Heidelberg challenge affected the alpha diversity of the cecal

and litter microbiome ($P < 0.05$) (Fig. 3b). The alpha diversity of the ceca of challenged

and control chicks raised on reused litter was higher than chicks grown on fresh litter for

Observed species measure of diversity ($P < 0.01$). Cloacal challenged chicks raised on

reused litter had higher alpha diversity in their ceca compared to chicks on fresh litter for

Shannon and Simpson indices ($P < 0.05$). Uninoculated control chicks on fresh litter had

higher alpha diversity in their ceca than control chicks on reused litter for Simpson

index, (Fig. 3b). The litter of cloacal, seeder, and control chicks on reused litter had

higher alpha diversity than the litter of chicks on fresh litter for Shannon measure of

diversity ($P < 0.05$) (Fig. 3b). In contrast, the litter of orally challenged chicks on fresh

litter had higher alpha diversity than the litter of gavaged chicks on reused litter for

Shannon index ($P = 0.0001$). There was no significant difference in alpha diversity

between fresh and reused litter for Observed and Simpson indices ($P > 0.05$).

On the taxonomic level, orally challenged chicks on reused litter carried higher

abundance of Actinobacteria in the ceca compared to chicks on fresh litter ($P = 0.003$),

while uninoculated control chicks on fresh litter had higher relative abundance of

Actinobacteria in the ceca compared to control chicks on reused litter ($P = 0.004$) (Fig.

3c). Proteobacteria was higher in the ceca of oral, cloacal and control chicks on reused

litter compared to chicks on fresh litter ($P < 0.01$). Firmicutes had higher relative

abundance in the ceca of chicks on fresh litter compared to reused litter for all

treatments ($P < 0.01$) while Bacteroidetes had higher relative abundance in the ceca of

cloacal and control chicks on reused litter compared to chicks on fresh litter ($P < 0.05$)

(Fig. 3c). For the litter, Actinobacteria was higher in reused litter compared to fresh litter

($P < 0.05$) while Proteobacteria and Bacteroidetes were higher in fresh litter compared

to reused litter ($P < 0.05$). There was no significant difference in phyla abundance

between routes of inoculation ($P > 0.05$) for the litter. These results shows that the

microbiome of chicks raised on fresh litter differed from chicks grown on reused litter,

and reused litter cecal and litter samples of challenged chicks were more uniform than

fresh litter samples.

S. Heidelberg challenge modulated the gut microbiome.

We determined that the core microbiome (i.e., ASVs > 0.1% relative abundance and

present in at least 80 % of samples) of fresh litter differs from reused litter (Table S1)

and found that *S*. Heidelberg challenge perturbed the cecal and litter microbiome of

broiler chicks (Fig. 4). For example, Enterobacteriaceae (*Klebsiella pneumonaie* and

Proteus mirabilis) were part of the core microbiome in fresh litter, while they were not

part of the core microbiome of reused litter (Table S1). Contrastingly, Actinobacteria (23

ASVs) was part of the core microbiome of reused litter, while only one ASV classified as

Actinobacteria was determined to be a core member of fresh litter (Table S1).

Unsurprisingly, the bacterial ASVs that increased or decreased in abundance after *S*.

Heidelberg challenge differed between chicks on fresh and reused litter (Fig. 4a and b).

253 To determine the ASVs that were modulated by *S.* Heidelberg challenge (i.e., ASVs

254 that increased or decreased after challenge), we compared the relative abundance of

255 ASVs in the ceca and litter of challenged chicks to uninoculated control chicks (Fig. 4).

256 In the ceca of chicks on fresh litter, the ASVs significantly modulated were assigned to

257 five bacterial families (Fig. 4a). Three ASVs matching family Clostridiales vadinBB6

258 group, nine ASVs classified as Ruminococcaceae and seven ASVs classified as

259 Lachnospiraceae increased in abundance in challenged chicks compared to control

260 chicks. Contrastingly, ASVs classified as Lactobacillaceae, Enterococcaceae,

261 Ruminococcaceae (n=8) and Lachnospiraceae (n=10) decreased in abundance in

262 challenged chicks compared to control chicks (Fig. 4a).

263  For the ceca of chicks on reused litter, the ASVs modulated were assigned to

264 twelve bacterial families. Amplicon sequence variants matching Bifidobacteriaceae,

265 order Clostridiales, Moraxellaceae, Rikenellaceae, Eggerthellaceae, Enterococcaceae,

266 Family XIII, Lactobacillaceae (n=2), Ruminococcaceae (n=2) and Lachnospiraceae

267 (n=2) increased in abundance in challenged chicks compared to control chicks. In

268 contrast, ASVs classified as Enterobacteriaceae, Enterococcaceae, Clostridiales

269 vadinBB6 group (n=2), Ruminococcaceae (n=3) and Lachnospiraceae (n=7) decreased

270 in abundance in challenged chicks compared to control chicks (Fig. 4a).

271  For the litter, the ASVs that were significantly modulated in fresh litter were assigned

272 to twenty-one bacterial families (Fig. 4b). The ASVs that increased in abundance in

273 fresh litter of challenged chicks belonged to Caulobacteraceae, Dysgonomonadaceae,

274 Burkholderiaceae, Flavobacteriaceae , Cellvibrionaceae, Enterobacteriaceae,

275 Enterococcaceae (n=2), Lactobacillaceae (n=2), Sphingobacteriaceae (n=2),

Pseudomonadaceae (n=3), Ruminococcaceae (n=3), Moraxellaceae (n=4), Clostridiales

vadinBB6 group (n=4), and Lachnospiraceae (n=4). Conversely, ASVs assigned to

Staphylococcaceae, Brevibacteriaceae, order Micrococcales, Corynebacteriaceae,

Xanthomonadaceae, Weeksellaceae, Bacillaceae (n=2), Ruminococcaceae (n=6) and

Lachnospiraceae (n=6) decreased in abundance in the litter of challenged chicks

compared to control chicks (Fig 4b).

Amplicon sequence variants modulated in reused litter after challenge were grouped

into twenty-four bacterial families. The ASVs that increased in abundance in reused

litter of challenged chicks were assigned to Lactobacillaceae, Saprospiraceae,

Bifidobacteriaceae, Rikenellaceae, Pseudomonadaceae, Sphingobacteriaceae,

Leuconostocaceae, Glycomycetaceae, Streptococcaceae, Enterobacteriaceae,

Wohlfahrtiimonadaceae, Bacteroides (n=2), Ruminococcaceae (n=3), Planococcaceae

(n=3), Lachnospiraceae (n=3) and Clostridiaceae 1 (n=5). The ASVs that decreased in

the reused litter of challenged chicks were classified as Burkholderiaceae,

Microbacteriaceae, Micrococcaceae, Moraxellaceae (n=2), Wohlfahrtiimonadaceae

(n=2), Aeromonadaceae (n=3), Clostridiales vadinBB6 group (n=3), Ruminococcaceae

(n=4), Lachnospiraceae (n=4) and Bacillaceae (n=6),

Not all bacterial families were modulated by the cloacal route of *S*. Heidelberg

challenge. For instance, Enterococcaceae was only modulated by oral challenge in the

ceca of chicks grown on fresh and reused litter (Fig. 4a). Similarly, Pseudomonadaceae

was only modulated in the litter of chicks that were orally challenged (Fig. 4b). Together,

these results show that the litter microbiome modulated by *S*. Heidelberg challenge

differed between chicks on fresh litter and reused litter, and the route of challenge

affected the ASVs that increased or decreased in abundance.

Enterobacteriaceae and Clostridiales were the major bacterial hosts for AMR.

We used Hi-C metagenomics to explore the bacterial reservoir of AMR in the cecal

microbiome of 2 cecal samples each from cloacally challenged chicks on fresh and

reused litter. We previously used this approach to identify *E. coli* as the main bacterial

reservoir of transferable plasmids in fresh litter [7]. Here, we extended our analysis to

cecal samples from reused litter. The average assembly size and number of contigs for

the four samples was 1,238,395,273±167,820 bp and 1,797,423±281,253, respectively.

The number of metagenome assembled genomes (MAGs) found in cecal samples from

fresh and reused litter was 456±422. We searched the MAGs for sequences matching

plasmid incompatibility (*inc*) groups and replicons available on the PlasmidFinder

database. The *inc* groups found matched common Enterobacteriaceae plasmids

including IncF, IncI, IncX, IncH, IncY, IncB/O/K/Z, p0111 and Col-like plasmids (Table

1). Three plasmid replicons were found for Gram-positive bacteria including *rep2*,

rep18b and *repA* (Table 1). The low representation of plasmids is expected, since the

PlasmidFinder database is much more comprehensive for Enterobacteriaceae than

other bacterial species, and within the well-studied Proteobacteria most plasmids

cannot be typed [31].

Next, we used proximity-ligation to find the bacterial host of ARGs and to determine

if they are encoded on plasmids or the chromosome. One hundred ARGs were found in

the MAGs and the majority (~ 50%) were harbored by family Enterobacteriaceae (Fig.

5). *Escherichia coli* was the only Enterobacteriaceae found to be the putative host for

these ARGs in fresh and reused litter. Members of the order Clostridiales were the

putative hosts for 27% of ARGs, while unclassified bacterial species (n = 6) and

members of the phyla Firmicutes (n = 9), Bacteriodetes (n = 3) and Actinobacteria (n =

1) were the hosts for the remaining ARGs. Fifty-four percent of the ARGs were found to

be encoded on the chromosome while 46% were found on plasmids. One cecal sample

from fresh litter (Fig. 5) harbored *E. coli* MAGs that encoded silver and copper

resistance genes (*silABCFRS* and *pcoABCDRS*) on the chromosome, while virulence

genes (*iroBCDEN*) were found on plasmids. A cecal sample from a chick raised on

reused litter harbored *E. coli* MAGs that encoded ampicillin (*bla*$_{TEM1B}$), tetracycline

(*tetA*), and mercury resistance (*merCPRT*) genes on plasmids (Fig. 5). This result

suggests that the microbiome of chicks on fresh and reused litter harbored AMR, and

Enterobacteriaceae and Clostridiales were the putative bacterial hosts of the ARGs.

E. coli isolates harbored AMR on plasmids and genomic islands.

In our earlier work, we found that *E. coli* MAGs were the primary hosts of IncI1 by

retrospectively screening the cecal contents of chicks on fresh litter for *E. coli* isolates

[7]. Using this approach, we confirmed that the IncI1-pST26 plasmid acquired by *S.*

Heidelberg was identical to the IncI1-pST26 plasmid present in *E. coli* strains (Fig. S2).

Here, we performed AST and WGS on additional *E. coli* isolates recovered from the

ceca of chicks used for Hi-C metagenomics (Table S2) and a pangenome analysis was

done on nineteen of them. The isolates were randomly selected from either

CHROMagar™ plates supplemented with or without gentamicin and tetracycline.

The inferred phylogeny of the *E. coli* strains using their core (n = 3385) and

accessory genes (n = 6024) partially grouped the isolates by the litter used for raising

the chicken host (Fig. 6). The ML trees grouped the isolates into three main clades

(Clade I - III). One *E. coli* isolate from the ceca of chicks on reused litter made up Clade

I on the core and accessory gene tree, while the number of isolates in Clade II and III

strains differed between core (Fig. 6a) and accessory gene tree (Fig. 6b). Clade II

isolates were clonal and recovered from chicks on fresh litter. Clade III was divided into

two subclades represented by isolates from fresh (subclade IIIa) and reused litter

(subclade IIIb) and subclade IIIb isolates were more clonal compared to subclade IIIa.

E. coli Isolates classified as phylogroup A, D and F were more likely to be found in

chicks on fresh litter, while phylogroup B1 was found in the ceca of chicks from reused

litter (Fig. 6, Table S2). Likewise, cgMLST classification showed that ST69, ST2705 and

ST6858 were associated with *E. coli* strains originating from chicks on fresh litter, while

ST1403 was found in reused litter. These results suggest that the *E. coli* strains from

this study have evolved to persist in either fresh or reused litter.

Some *E. coli* strains from fresh litter harbored ARGs on the chromosome and

plasmids. *E. coli* strains classified as pandemic lineage of extraintestinal pathogenic

ST69 from fresh litter harbored a ~128 kb genomic island encoding virulence genes

(pyelonephritis-associated pili (*pap)* operon) and ARGs for aminoglycoside (*strAB* and

aph(3')-Ia) tetracycline (*tetB*), sulfonamide (*sul2*) and metal resistance (silver (*sil*) and

copper (*pco)* operons) (Table S2; Fig. S3). The presence of a genomic island encoding

AMR suggests that the ARGs are mobile. We found DNA regions encoding identical

metal resistance genes (10 - 40 kb) as seen in the genomic island in p0111/IncH1B

plasmids harbored by *E. coli* strains from this study and in *E. coli* genomes found in the

NCBI database (Fig. S3). In addition, ST69 *E. coli* strains harbored a multireplicon IncF

plasmid encoding bla_{TEM-1B}, *aac(3)-IId* and virulence genes (colicin M and catecholate

siderophore uptake system (*iroBCDEN*)), and an untypeable IncI1 plasmid. (See Data

Set S1 in the supplemental material; Data Set S1 to S5 are available in the Dryad

repository online at https://doi.org/10.5061/dryad.c866t1g6c).

As previously reported [7], *E. coli* ST6858 from fresh litter carried an identical *S.*

Heidelberg IncI1-pST26 plasmid, and harbored IncI2, IncF, and cryptic Col-type

plasmids (Data Set S2). Antibiotic susceptible ST2705 *E. coli* isolates from fresh litter

harbored no plasmid encoding antibiotic resistance genes but carried a phage-like

plasmid classified as p0111 and a multireplicon IncF plasmid encoding metal resistance

and virulence genes (Data Set S3). The common ARG carried by ST1403 *E. coli* strains

from reused litter was bla_{TEM-1B}. The gene was encoded on an IncX1 plasmid (Data Set

S4), and it conferred ampicillin resistance. Additionally, ST1403 *E. coli* strains harbored

IncFIB, IncI2 and ColRNAi plasmids. One *E. coli* isolate (ST93) from reused litter

harbored an IncH1B/p0111 plasmid encoding ARGs for aminoglycoside (*aadA1, aac(3)-*

VIa), tetracycline (*tetB*), sulfonamide (*sul1*) and mercury resistance (*mer* operon) and an

IncF plasmid encoding colicin M and *iroBCDEN* genes (Data Set S5).

Bacterial hosts from reused litter maintained IncI1 plasmid at low copies.

The low rate of AMR acquisition by *S.* Heidelberg isolates from reused litter made us

hypothesize that IncI1-pST26 plasmids are present at lower copies in reused litter. To

test this hypothesis, we used qPCR assays targeting the region upstream of the *repA* of

IncI1 plasmid including the *inc*RNAI to determine their copy number in the ceca of

chicks on fresh and reused litter. We normalized the abundance of IncI1 against

glyceraldehyde-3-phosphate dehydrogenase A (*gapA*) housekeeping gene encoded on

the chromosome of several enteric bacteria [32] including *Salmonella* and *E. coli.* We

targeted Enterobacteriaceae with *gapA* because *E. coli* was determined to be the hosts

for AMR by metagenomics and WGS in this study. The relative abundance of

Enterobacteriaceae bacterial population carrying IncI1 was higher in the ceca of orally

challenged chicks on fresh litter compared to chicks on reused litter (W = 85, *P* =

0.00041) and was not significantly different for cloacally inoculated (W = 23, *P* = 0.24)

and uninoculated control chicks (W = 63, *P* = 0.16) (Fig. 7).

To further explore the effect of low copy IncI1 plasmids on AMR transfer, we

selected the *S*. Heidelberg isolate from reused litter harboring IncI1-pST26 that was

unexpectedly susceptible to antibiotics predicted by ARGs encoded on the plasmid. The

isolate (SH-16-0-5B) was confirmed twice by broth microdilution to be susceptible to

gentamicin, streptomycin, and tetracycline and once on Mueller Hinton agar

supplemented with varying concentrations (0, 2, 4, 8 and 16 mg/l) of gentamicin (see

supplementary methods). To ensure that the plasmid contig identified as IncI1-pST26 in

this strain was not an artifact of short read sequencing, we performed qPCR on plasmid

DNA and targeted four coding DNA sequence on IncI1-pST26 including its *inc*RNAi,

class-1 integron (*intI1*), aminoglycoside (*aadA1*) and tetracycline resistance (*tetA*)

genes (Table S3). This confirmed that the strain carried IncI1-pST26 (qPCR cycle

threshold ranged from 20.2 – 27.9 for the four targets). Afterwards, we used WGS

(depth of coverage of IncI1-pST26 contig/depth of coverage of the largest chromosome

contig) to compare the copy number of IncI1-pST26 in SH-16-0-5B to the copy number

of IncI1-pST26 in four *S*. Heidelberg isolates (hereafter referred to as SH-IncI1-FL)

recovered from the ceca of cloacally inoculated chicks on fresh litter. This revealed that

SH-16-0-5B maintained Incl1-pST26 at ~0.2 copies/cell (i.e., 1 of 5 cells harbored

Incl1) compared to 2.8±0.2 copies/cell in SH-Incl1-FL from fresh litter.

To investigate if the copy number difference determined using WGS exist in relevant

environmental conditions, we first acclimatized the isolates in pre-reduced cecal extract

(pH 6.5) for 2 h before exposing them to pre-reduced cecal extract at pH 2.5 under

microaerophilic conditions. We reasoned that the low pH of the upper gastrointestinal

tract (GIT) of the chicken host will interfere with the copy number of plasmids kept in a

bacterial cell, which would have a direct effect on the rate of HGT. The copy number of

Incl1-pST26 was significantly lower (0.00005 - 0.21 copies /per cell i.e., 1 in 20,000 to 1

in 5 cells carried Incl1) in SH-16-0-5B compared to SH-Incl1-FL (4.94 - 6.65 copies /per

cell) for pH 6.5 (W = 9, P = 0.1) and pH 2.5 (W = 81, P = 4.114e-05) (Table 2). These

results show that $E.\ coli$ and $S.$ Heidelberg populations in reused litter harbored lower

copies of Incl1-pST26 than the populations in fresh litter.

Discussion

Litter is commonly used as a bedding material for raising poultry and it is ingested

by chickens during pecking activities. Therefore, the microbiome of the litter is one of

the first inoculum that colonizes the GIT of broiler chicks. Litter can be managed as

single use i.e., complete removal of litter from a broiler house after each flock or it can

be reused over multiple flocks. The benefits of litter reuse include lower cost for growing

chickens and the sustainable management of litter waste [33]. It has been shown that

chickens raised on reused litter harbor a different microbiome than chicks grown on

fresh litter [17, 34] and that reused litter stimulates higher humoral and cell-mediated

immune responses than fresh litter in chickens [35]. Muniz et al. [23] and Roll et al. [26]

showed that the number of litter samples positive for *Salmonella* significantly decreases

as the number of litter reuses increased compared with the first use of the litter.

In this study, we corroborate these findings by showing that neonatal chicks

challenged with *S*. Heidelberg and raised on reused litter were more resistant to S.

Heidelberg colonization i.e., lower positivity rate compared to chicks on fresh litter. In

addition, we found that chicks raised on reused litter were less likely to be colonized

with *S*. Heidelberg isolates harboring plasmids encoding AMR compared to chicks on

fresh litter. We determined that the major difference between chicks grown on fresh and

reused litter was the bacterial community harbored in the litter and ceca.

Pathogen induced gut microbiome modulation can create an imbalance in the

abundance of bacterial species i.e., some species outgrow known resident bacteria,

leading to a dysbiosis of the gut microbiome and successful *Salmonella* colonization

and AMR transfer [36]. However, gut conditions that allow the maintenance of a diverse

and beneficial microbiome are likely to succeed in establishing homeostasis and limit

the transfer of AMR [37, 38]. In this study, we found that the microbiome of reused litter

was associated with a lower HGT of AMR to *S*. Heidelberg populations in neonatal

chicks. The microbiome data led us to draw one main conclusion that the litter used for

growing chicks and the route of *Salmonella* challenge affected which ASVs increased or

decreased in abundance in the ceca and litter.

This was the case for the ASV assigned to *Bifidobacterium* that increased in the

ceca and litter of orally challenged chicks raised on reused litter but was not modulated

in cloacally inoculated chicks or in chicks on fresh litter. Consequently, *Bifidobacterium*

was not part of the core microbiome of the ceca or litter in this study (Table S1) and

most likely evolved with neonatal chicks. *Bifidobacterium* are strict anaerobic

Actinobacteria and are among the first microbes to colonize the GIT of vertebrates [39].

Additionally, *Bifidobacterium* has been shown to affect the abundance of AMR and

virulence genes in the GIT of infants [40-42]. Casaburi et al [42] reported that infants fed

a probiotic strain of *B. longum* harbored lower levels of ARG compared to non-fed

controls. Alignment of shotgun metagenomic reads generated from the ceca of broiler

chicks (n = 4) to Kraken database [43] revealed that 21% and 11% of the reads

mapping to the genus *Bifidobacterium*, were aligned to *B. longum* and *B. catenulatum,*

respectively (Fig. S4). Their increase in abundance for only orally challenged chicks on

reused litter suggests that the route *S.* Heidelberg used for gaining entry into the GIT

and the litter microbiome affected how *Bifidobacterium* was perturbed. Therefore, it is

conceivable that reused litter microbiome created conditions that allowed

Bifidobacterium to flourish and proliferate, while the microbiome in fresh litter led to a

reduction or elimination of *Bifidobacterium* species.

We used metagenomics and WGS to show that *E. coli* was the main reservoir of

AMR in the cecal microbiome of fresh and reused litter. Importantly, we found the

identical IncI1-pST26 plasmid, acquired by *S.* Heidelberg, to be present in *E. coli*,

suggesting that the plasmid was transferred in vivo within chicks raised on fresh litter.

The possible sources of the *E. coli* strains in this study include the chicken host and the

broiler house environment. Although, we did not establish if the meconium harbored *E.*

coli, it is plausible that post-hatch chicks were colonized with *E.coli* from the hatchery

[44]. The evolution of these ancestral *E. coli* strains would be shaped by selective

pressures such as exposure to antibiotics and metals, and competition with resident

microbiota in the environment. For chicks placed on fresh litter, the bacterial community

of the first fecal droppings will compete with mostly pine shaving/plant associated

microbiome. In contrast, chicks placed on reused litter will compete with microbes and

metabolites deposited from earlier flocks including resident *E. coli* and *Salmonella*

strains that are well-adapted to the broiler house environment. Also, the physio-

chemical properties of the litter will affect the litter structure, the native microbiome and

the survival of the invading pathogen. The difference in litter physio-chemical

parameters in this study was inferred using pH and moisture. The pH of reused litter

was higher (7.12 ± 0.26) than fresh litter (6.54 ± 0.17) (P = 0.028), while fresh litter had

higher moisture (20.83 ± 3.54 %) than reused litter (14.65± 3.51 %) (P = 0.11). Litter

moisture and pH are known factors that affect pathogen survival and microbial diversity

[45, 46].

We did not administer antibiotics to chicks in this study and have shown that metals

did not significantly affect the metabolism of *S*. Heidelberg [7]. Hence, the litter

microbiome and physio-chemical properties are the major selection pressure that could

explain the evolutionary trajectory of *E. coli* in this study. This hypothesis was further

supported after a functional protein pathway analysis was performed on the cecal

metagenome of chicks raised on fresh and reused litter. Pathways related to the

biosynthesis of organic and antimicrobial molecules were enriched to a greater degree

in the ceca of chicks on reused litter than chicks on fresh litter (Fig. S5). Of note was the

enrichment of biosynthetic pathways for antimicrobials and secondary metabolites in

reused litter (e.g., macrolides, tetracyclines, carbapenems, vancomycin and

polyketides), suggesting that reused litter harbored a higher abundance of microbial

species with antimicrobial properties than fresh litter.

Contrastingly, pathways associated with the biosynthesis of secondary metabolites

of plants including flavonoids, indole-alkaloids and stilbenoid, diarylheptanoid and

gingerol were enriched in samples from fresh litter (Fig. S5). Though we measured only

two cross-sectional timepoints for reused and fresh litter, the relative abundance of

plant-metabolic related pathways in fresh litter compared to the enrichment of

antimicrobial molecules in reused litter suggests that a selective pressure gradient

exists over time for the colonizing microbiome in reused litter. Thus, the microbiome of

reused litter has the antimicrobial capability to competitively exclude invading pathogens

including strains carrying AMR encoding plasmids.

Plasmid carriage is expected to impose a fitness cost on the host, but plasmid

bearing microbes are pervasive in nature [47]. We showed previously that carriage of

IncI1-pST26 by *S*. Heidelberg isolates from fresh litter presented a variable fitness cost

to the host [7]. We found here that Enterobacteriaceae populations in the ceca of

chicks on reused litter harbored lower copies of the IncI1 plasmid per cell compared to

chicks on fresh litter. In addition, we did not isolate any *E. coli* isolate carrying IncI1 from

the ceca of chicks raised on reused litter. The limited number of Enterobacteriaceae

IncI1 donor population would affect the rate of transfer to *S*. Heidelberg recipients as

direct contact is a requirement for conjugation [48]. Nevertheless, we found one *S*.

Heidelberg isolate from the ceca of one chick on reused litter that harbored IncI1-

pST26, however, only a fraction of its population harbored IncI1 plasmid. This isolate

did not exhibit the resistance phenotype predicted by the ARGs encoded on the IncI1-

pST26 plasmid. These results suggest that IncI1 plasmid carriage in reused litter posed

a fitness cost on the bacterial host, but this cost was ameliorated by maintaining IncI1 at

low copies in our study.

It is important to mention that the litter used for this study is not representative of

all broiler litter. Litter composition may differ between farms and there are multiple

physio-chemical factors that can affect the litter microbiome, e.g., moisture and pH,

antibiotic usage, and feed additives. Furthermore, the age of the litter i.e., numbers of

flocks that have been grown on the litter, and the length of the downtime between flocks

are examples of management practices that may influence microbiome succession.

Nonetheless, our study showed that the litter microbiome significantly affects the

bacterial diversity of broiler chicks and the HGT of AMR.

Materials and Methods

Details of methods used for preparing the *S*. Heidelberg inocula, challenging neonatal

chicks, bacterial and DNA analyses, whole genome, and metagenome sequencing and

bioinformatics have been described before for neonatal chicks raised on fresh litter [7] .

We briefly redescribe these methods and present others below.

Determining if post-hatch chicks were *Salmonella*-free.

Chick pad that conveyed chicks from the hatchery were pre-enriched in 500 ml of

Buffered Peptone Water (BPW) for 18-24h at 37° C. Two different enrichments broths

were used to isolate *Salmonella* from the BPW pre-enrichment broths: Tetrathionate

(TT; Becton-Dickinson, Sparks, MD) broth and Rappaport-Vassiliadis (RV; Becton

Dickinson) media. After overnight incubation at 42°C in both enrichment broths, 10 µl

aliquots from each enrichment broth was spread on two different differential media:

Brilliant Green Sulfa (BGS; Becton Dickinson) agar and xylose lysine tergitol-4 (XLT-4;

Becton Dickinson) agar and incubated for 18-24 h at 37° C. Isolated colonies

characteristic of *Salmonella* were stabbed and streaked onto triple sugar iron agar (TSI;

Becton-Dickinson) and lysine iron agar fermentation (LIA; Becton-Dickinson) and

incubated for 18-24 h at 37° C for biochemical confirmation.

***S*. Heidelberg inoculum preparation.**

The *S*. Heidelberg strain was made resistant to 200 ppm of nalidixic acid (nal^R *S*.

Heidelberg) for selective enumeration and was grown overnight in poultry litter extract,

centrifuged, and resuspended in 1X phosphate buffered saline (PBS). The resuspended

cells were used as inocula. The complete genome of the ancestor to nal^R *S*. Heidelberg

is available under GenBank accession number : CP066851.

Challenging broiler chicks with nal^R *S*.*S*. Heidelberg.

One-day-old Cobb 500 broiler chicks were either uninoculated (n =25), gavaged (n =25)

or (n =25) cloacally inoculated with ~10^6 colony forming units (CFU) of nal^R *S*.

Heidelberg. We also included a seeder-bird colonization method, whereby five chicks

were gavaged and mingled with twenty uninoculated chicks. Afterwards, chicks were

placed in floor pens at a stocking density of 0.65 m^2/chick on fresh pine shavings (fresh

litter, n =100) or reused litter (n =100). The reused litter was previously used to raise

three flocks of broiler chicken's antibiotic-free under simulated commercial poultry

production conditions and was top dressed with 0.5 cm of fresh pine shavings before

the placement of each flock. Broiler chicks on fresh and reused litter were housed

separately and were raised for 14 days on antibiotic-free starter diet and water. At 14

days, forty-five chicks from each fresh and reused litter (ten chicks from gavaved,

cloacal and uninoculated groups and fifteen chickens from the seeder method (5 seeder

and 10 contact chicks) were euthanized to determine the extent of nal^R *S*. Heidelberg

colonization in ceca. Additionally, litter samples were collected as grab samples from

each pen after chicks were euthanized. The experiments were performed in April 2018.

The study was approved by the University of Georgia's Office of Animal Care and Use

under Animal Use Protocol: A2017 04-028-A2.

Determination of nal^R *S*. Heidelberg concentration in the ceca and litter.

Ceca were removed from the eviscera of broiler chicks and stomached for 60 seconds

after the addition of 3X volume to the weight (v/w) of BPW. Litter was collected as grab

samples from seven locations (4 corners of the pen and 3 locations under the waterer)

in each pen after chicks were removed. The litter samples were pooled and 30g was

subsampled in duplicates from each pen as previously described [29]. Serial dilutions of

the cecal and litter slurry were plated onto BGS containing 200-ppm nalidixic acid and

plates were incubated for 24 h at 37°C.

After incubation, colonies were counted and calculated per gram of ceca or litter dry

weight. When no colonies appeared, pre-enriched cecal and litter slurry was streaked

onto BGS agar supplemented with nalidixic acid and incubated overnight. These plates

were then examined for the presence/absence of *Salmonella* colonies. Two to six single

colonies were randomly selected from each plate and archived in 30% LB glycerol at -

80 ^{o}C. In addition, cecal slurry was saved at a 4:1 ratio in Luria Bertani (LB) broth (BD

Difco, MD, USA) containing 30% glycerol at –80°C, while litter samples were stored in

vacuum sealed whirl pack bags at -20°C. Litter pH and moisture was determined as

described previously [29].

Antibiotic resistance phenotype determination.

Antibiotic susceptibility testing was done on *S.* Heidelberg and *E. coli* isolates by following the National Antimicrobial Resistance Monitoring System (NARMS) protocol for Gram-Negative bacteria. Results were interpreted according to clinical and laboratory standards institute guidelines and breakpoints established by NARMS. In addition, we used agar dilution to determine the minimum inhibitory concentration of gentamicin for one *S.* Heidelberg isolate (SH-16-0.5B) (see supplementary methods).

Genomic, environmental and plasmid DNA extraction.

Unless otherwise noted, DNA was extracted and purified from bacterial cultures using FastDNA™ spin Kit for Soil (MP Biomedicals, LLC, CA, USA), while 250 mg of cecal slurry (previously saved in LB broth containing glycerol) and litter were extracted with Qiagen DNeasy Power-Soil DNA kit (Hilden, Germany). Extracted DNA were used for qPCR, 16S rRNA gene sequencing, shotgun metagenomics and whole genome sequencing. Plasmid DNA was extracted from two *S.* Heidelberg isolates that harbored IncI1. Isolates were cultured overnight on sheep blood agar and plasmid DNA was extracted using Qiagen Plasmid Midi kit (Qiagen Inc, Germantown, MD) as per manufacturer's instructions. Plasmid DNA was used to construct a calibration curve for IncI1 plasmid copy number determination.

Real-time quantitative PCR.

Real-time qPCR amplification was performed as described previously (45) using a CFX96 Touch Real-Time PCR Detection System (Bio-Rad Inc., Hercules, CA). Reaction mixtures contained 1X SsoAdvanced Universal SYBR Green Supermix (Bio-Rad Inc., Hercules, CA), 600 nM (each) primers) and 2 µl of DNA. Primers used in this study are

shown in Table S3. Unless otherwise stated, primers were designed with beacon

designer (Premier Biosoft, Palo Alto, CA) and synthesized by Integrated DNA

Technologies (Coralville, IA). Calibration curves used for converting qPCR cycle

threshold values to gene copies per gram of ceca was determined using the genomic

DNA from an *E. coli* strain that harbored IncI1-pST26 plasmid. To convert qPCR cycle

threshold values to IncI1 copies per cell of *S.* Heidelberg, plasmid DNA of a relevant *S.*

Heidelberg strain harboring IncI1-pST26 was used for calibration curve construction

(Table S3). Two primer sets specific to the *gapA* gene and the region upstream of the

repA of IncI1 including the *inc*RNAI were used to determine the gene copies of IncI1 in

the ceca. The *gapA* is a housekeeping gene encoded on the chromosome of enteric

bacteria including *E. coli*, *Klebsiella*, *Citrobacter* and *Salmonella* [32]. The gene copies

of IncI1 per cell was determined as the copy ratio of *inc*RNAi to *gapA* [49]

Whole genome sequencing and analysis.

Whole genome sequencing libraries were prepared using either Nextera™ XT or

Nextera™ DNA Flex library preparation kits (Illumina, Inc., CA, USA) following the

manufacturers protocol. Libraries were sequenced on the Illumina MiSeq platform with

150 or 250-bp paired end reads. Additionally, five *E. coli* isolates were selected for long

read sequencing using Sequel II System (PacBio Biosciences Inc.) or MinION device

(Oxford Nanopore Technology) (File S1). Preparation and sequencing of long read

libraries were done by sequencing core centers of University of Georgia and Colorado

State University. The method used for read quality control and demultiplexing has been

reported [7].

Genome assembly, resistome characterization, and quality assessment of

assemblies was done using Reads2Resistome pipeline v.1.1.1 [50]. For tools used for

genome annotation, bacteria and plasmid typing, see supplementary methods. MAFFT

v. 1.4.0 [51] implemented in Geneious Prime® v 2020.0.1, was used to align and

compare sequences. Roary [52] was used for pan genome analysis and phylogenetic

trees were constructed using the maximum likelihood method implemented in RAxML-

NG v 1.0.0 [53]. IslandViewer [54] was used to predict genomic islands.

Identification of single nucleotide polymorphism (SNP) and Indels present in *S.*

***Heidelberg* isolates**.

Alignment of raw FASTQ reads to *S.* Heidelberg (Genbank accession number:

CP016573) was done using BWA (v0.7.17) [55] and SNPs/indels were called using

Genome Analysis Toolkit [56]. Variant call format (VCF) files of SNPs/Indels and the

script   used   are   available   in   Dryad   Digital

Repository: https://doi.org/10.5061/dryad.c866t1g6c.

16S rRNA gene community analysis.

The V4 hypervariable region of the 16S rRNA gene was sequenced using the paired-

end (250 × 2) method on the Illumina MiSeq platform. The cecal (n = 59; 30 and 29

chicks from fresh litter and reused litter, respectively) and litter (n = 22, 11 samples each

from fresh and reused litter) samples were part of a larger sequencing run and

processed together with other samples. Thus, detailed sequence processing

parameters are described in [7]. To avoid bias introduced by spurious amplicon

sequence variants (ASVs) or from samples with low sequencing depth, any ASVs with

less than 5 reads and samples with less than 5000 reads were removed from the

dataset before further analysis.

Statistical analysis of microbial communities was performed in the R environment

using the packages "phyloseq", "Ampvis2", and "vegan". Alpha diversity indices were

calculated with a dataset rarefied to the minimum sample size (8740 sequences) and

compared using the Wilcoxon Signed Rank test. Principal coordinates analysis (PCoA)

based on Bray-Curtis distances was performed with no initial data transformation and

after removing ASVs present with relative abundance less than 0.1% in any sample.

The core microbiome was determined as ASVs 0.1% relative abundance and present

in at least 80 % of samples.

Cecal metagenome sequencing.

Two hundred and fifty milligrams of cecal slurry from chicks cloacally challenged with *S.*

Heidelberg and raised on fresh (n = 2) or reused litter (n = 2) were used for generating

shotgun and Hi-C DNA library. For shotgun library, cecal DNA was extracted with

Qiagen DNeasy Power-Soil DNA kit (Hilden, Germany) and Nextera™ XT library

preparation kit was used to make the library. Hi-C library was made using Phase

Genomics (Seattle, WA, USA) ProxiMeta Hi-C Microbiome Kit following the

manufacturer's instructions. Libraries were sequenced by Novogene corporation

(Sacramento, CA, USA) on the Illumina HiSeq platform using 150bp paired end reads.

Two libraries were sequenced per HiSeq flowcell lane resulting in a total of ~111 million

shotgun reads, and ~213 million Hi-C reads per sample. Quality controlled shotgun

reads were classified using Kraken2 (v2.0.8 beta) [43] to create a count profile of the

metagenome. We used cumulative sum scaling implemented in metagenomeSeq to

normalize read counts [57].

Metagenome assembled genomes and associated plasmids and ARGs

The details of the methods used for proximity-guided metagenome assembly and

deconvolution have been described before [7]. Briefly, shotgun metagenomics

sequence reads were filtered and trimmed for quality and normalized before assembly

with metaSPAdes [58] using default options. Hi-C sequence reads were then aligned to

the assembly following the ProxiMeta Hi-C kit manufacturer's recommendations

(https://phasegenomics.github.io/2019/09/19/hic-alignment-and-qc.html).   Metagenome

deconvolution was performed with ProxiMeta [59], resulting in the creation of putative

genome and genome fragment clusters. Hi-C clusters were assessed for quality using

CheckM [60] and assigned taxonomic classifications with Mash v2.1 [61] resulting in the

metagenome assembled genomes (MAGs). We used PlasmidFinder (v2.1.1) [62] to

identify plasmids present in MAGs.

To view AMR gene connections to hosts, AMR genes in the metagenomic

assembly were annotated using NCBI's AMRFinderPlus software (version: 3.10.5) with

the --plus option specified, and all other options set to defaults. The AMR assembly

annotations were then used in combination with the MAG taxonomic annotations and

the mobile element to host association matrices generated by ProxiMeta to annotate

which AMR genes are present in each MAG, and specify whether the gene originated

from a genomic, viral or plasmid contig. This was done by making a NxM matrix C,

where $N = |AMRa|$ and $M = |MAG|$, where $MAG = \{$set of individual metagenomic bins$\}$

and $$AMRa = \{x \subset AMR : x \in \text{Assembly}\},$$ where

$AMR = \{\text{set of all AMR genes in database}\}$. The matrix ($C$) was then filled as:

For i in $|AMRa|$:

For j in $|MAG|$:

711
$$C_{i,j} = \begin{cases} 0 & \text{if } AMR_i \notin MAG_j \\ S & \text{if } AMR_i \in MAG_j \end{cases}, \text{ where } S = \begin{cases} 1 & \text{if } AMR_i \in AMRg \\ 2 & \text{if } AMR_i \in AMRv \\ 3 & \text{if } AMR_i \in AMRp \end{cases}$$

and where, $AMRg = \{x \subset AMRa : x \in \text{Genomic Contigs}\}$, $AMRv = \{x \subset AMRa : x \in$

Viral Contigs$\}$, $AMRp = \{x \subset AMRa : x \in \text{Plasmid Contigs}\}$. The matrix, C, was then

filtered to remove rows and columns with only 0 values and plotted using a color

encoding system indicating whether the AMR gene was of genomic (cyan) or plasmid

(gold) origin.

Functional Pathway Analysis.

Both shotgun sequences and ProxiMeta-deconvoluted genomes were also utilized to

assess enrichment of functional pathways. Shotgun metagenomic sequences were

aligned to the Hi-C genomic contigs using BWA-MEM algorithm as implemented in the

BWA (v0.7.17) [55]. The Hi-C genomic contigs were then annotated using Prokka

(v1.13) [63]. For each annotated region on the Hi-C genomic contigs, sequences

overlapping the annotated region in the sequence alignments were counted using a

custom Python script to produce a count matrix of genes for all samples. Gene counts

were normalized by gene length and using total sum scaling normalization to control for

differences in sequencing depth [64]. Enzyme commission identifiers were then mapped

to KEGG functional pathways according to the KEGG ontology. For each pathway, an

adjacency graph was reconstructed from the KEGG XML files, and connected

components were determined using depth-first search [65]. Z-scores were calculated for

pathways and connected components for each sample and statistical testing was then

performed by comparing scores to the normal distribution. Details of this analysis is

described in the supplementary methods.

***E. coli* isolation from ceca.**

The method used for screening frozen cecal contents has been reported [7]. Briefly, two

cecal slurry from challenged chicks raised on fresh litter and reused litter were spread

plated onto CHROMagar™ plates supplemented with or without gentamicin (8 ppm) and

tetracycline (8 ppm) or ampicillin (16 ppm) and cefoxitin (16 ppm) (Thermo Fisher,

Waltham, MA). After 24 h incubation, blue green and blue-cream colonies were counted

as presumptive *E. coli* and 4 - 5 colonies from each spread plate were picked and

restruck for colony purification. No colonies grew on CHROMagar™ plates

supplemented with ampicillin and cefoxitin. Colonies were preserved in LB broth

containing glycerol and frozen before they were used for including solid agar

competition experiment.

Determination of plasmid copy number of IncI1.

To compare the copy number of IncI1 harbored by *S.* Heidelberg strain (SH-16-0-5B)

from reused litter to the copy number of IncI1 in three SH-IncI1-FL isolates from fresh

litter, we exposed each isolate to acidified filter sterilized cecal extract. Cecal extract

(CE) was prepared from the cecal content of 2-weeks old broiler chickens (see

supplementary methods). Cecal extracts were pre-reduced by covering tubes with gas

permeable paper strip and incubating overnight under microaerophilic conditions (5%

O_2, 10% CO_2, and 85% N2) at 42^oC. Three single colonies of each strain were selected

from overnight cultures grown on sheep blood agar and transferred to a microcentrifuge

tube containing 900 µl of pre-reduced CE (pH = 6.52), i.e., one tube per for each SH-

IncI1-FL strain and three tubes for SH-16-0-5B.

After transfer, tubes were vortexed, covered with a gas permeable paper strip and

incubated at 41^{o}C under microaerophilic conditions for 2 h. After incubation, tubes were

vortexed and 100 µl of the suspension was transferred to a microcentrifuge tube

containing 900 µl of CE (pH of CE was adjusted to 2.52 using 1M HCl (Spectrum

Chemical Mfg. Corp., CA, USA) and 1M NaOH (Fisher Chemical, NJ, USA) Afterwards,

tubes were vortexed, and incubation continued for another 24 h. To determine the copy

number of IncI1, one replicate tube per strain was removed at timepoints 2 h for pH 6.5

and 0.5, 4 and 24 h for pH 2.5 and used for DNA extraction. DNA was extracted from

500 - 800 µl of CE samples using FastDNA™ SPIN Kit for Soil. Calibration curves for

gapA and *inc*RNAI were generated using genomic and plasmid DNA extracted from

ancestral *S.* Heidelberg (GenBank accession number : CP066851) and one SH-IncI1-

FL isolate, respectively. Plasmid copy number of IncI1 was determined as the copy ratio

of *inc*RNAi to *gapA* [49]

Statistical analyses.

Continuous variables were log-transformed before any statistical tests were performed.

Moreover, continuous variables did not meet the assumption of a normal distribution;

therefore, non-parametric testing for direct comparisons was performed using Wilcoxon

rank sum and signed rank tests, and the Kruskal-Wallis rank sum test was used for one-

way analysis of variance tests. Additionally, a generalized linear model with a binomial

distributed outcome and a log link function was performed as described earlier [66] to

determine if there are significant differences in *Salmonella* prevalence

(presence/absence) and the litter used for raising broiler chicks. The significance of the

model was established using a likelihood ratio test (R function ANOVA with argument

test set to "Chisq"). Multiple comparisons of means, i.e., regression coefficients for route

of inoculation and litter age/type were done using the multcomp package in R. The mcp

function was used to specify linear hypotheses and the glht function was used to make

Tukey contrasts. Statistical analyses were performed using R (v4.0.3).

Ethics statement.

All animal experiments were approved by the University of Georgia Office of Animal

Care and Use under Animal Use Protocol: A2017 04-028-A2.

Data availability.

All raw short and long FASTQ reads for *S.* Heidelberg are publicly available under NCBI

accession number: PRJNA683658, while *E. coli* FASTQ reads are available under NCBI

accession number: PRJNA684578. Shotgun and Hi-C reads from fresh litter and reused

litter cecal samples are publicly available under NCBI accession number:

PRJNA688069 and 16S rRNA gene sequences under NCBI accession number:

PRJNA669215. The complete genome assemblies for *E. coli* strains Ec-FL1-1X and Ec-

FL1-2X were previously published and are available under GenBank accession

numbers CP066836 and JAFCXR000000000. Complete genome of *E. coli* strain Ec-

RL2-1X has been made available under GenBank accession numbers CP066839.

Variant call format (VCF) files of identified SNPs/indels and the Linux/Unix shell script

used have been deposited in Dryad Digital Repository:

https://doi.org/10.5061/dryad.c866t1g6c.

Acknowledgements.

We are grateful to Marlo Sommers, Carolina Hall, Jeromey Jackson, and Latoya Wiggins for their logistical and technical help. This work was supported by USDA Agricultural Research Service (Project Number: 6040-32000-010-00-D), non-assistance cooperative agreement (58-6040-6-030) between USDA Agricultural Research Service and University of Georgia, Research Foundation, and research service agreement (58-6040-8-035) between USDA Agricultural Research Service and Colorado State University.

This study was supported in part by resources and technical expertise from the Georgia Advanced Computing Resource Center, a partnership between the University of Georgia's Office of the Vice President for Research and Office of the Vice President for Information Technology. Any opinions expressed in this paper are those of the authors and do not necessarily reflect the official positions and policies of the USDA or the National Science Foundation, and any mention of products or trade names does not constitute recommendation for use. IL, MOP, and JRG are employees of Phase Genomics, Inc., and have been supported in part by NIAID grants R44AI162570 and R44AI150008. The remaining authors declare no competing commercial interests in relation to the submitted work. USDA is an equal opportunity provider and employer.

Author's contributions.

A.O., and Z.A. designed the study. A.O., G.Z., D.E.C. and C.R. performed live-broiler chicken studies. M.J.R contributed to the design of the solid agar mating experiment. A.O., J.L., G.Z., and D.C. performed bacteriological analyses, antibiotic susceptibility testing, DNA extraction and qPCR. G.Z. and A.O. performed Illumina whole genome

sequencing. A.O. performed cecal shotgun and Hi-C library preparation. T.L. made 16S

rRNA gene libraries and sequencing. B.Z. performed 16S rRNA bacterial community

analysis and interpretation. A.O., Z.A., M.O.P., I.L., J.R.G., C.W., J.C.T., S.M.L., and

R.W. performed bioinformatic analyses and data curation. A.O., J.L. and D.C.

performed solid agar mating experiment. A.O., and B.Z. performed statistical analyses.

A.O., Z.A., N.A.C., B.Z., R.W., M.O.P., J.R.G., S.M.L., G.Z., and J.L. drafted the

manuscript, which was reviewed and edited by all authors. A.O., Z.A., and S.A.A.

supervised the study.

850 Table 1. Plasmid incompatibility groups and replicons found in cecal metagenome assembled genomes
851 using PlasmidFinder[a].
852

	Hi-C-FL1	Hi-C-FL2	Hi-C-RL1	Hi-C-RL2
Gram-negative	Col(MG828)	Col(MG828)	Col(MG828)	Col(MG828)
	Col156	ColpVC	Col(pHAD28)	Col156
	ColpVC	ColRNAI	Col156	IncFIB
	ColRNAI	IncB/O/K/Z	ColRNAI	IncFII(pCoo)
	IncFIB	IncFIB	IncFIB	IncHI1B(pNDM-CIT)
	IncFIC(FII)	IncFII(pSFO)	IncFII(pSFO)	IncI1-I(Alpha)
	IncFII(pRSB107)	IncI2	IncI1-I(Alpha)	

	IncHI1B(pNDM-CIT) IncI1-I(Alpha) IncI2(Delta) IncX1 IncX2 p0111	IncX1 IncX2	IncI2 IncX1 IncX2 IncY	
Gram-positive	rep2 repA(pB82)	None	rep18b	repA(pB82)

853 [a] Sample IDs; FL = Fresh litter, RL = Reused litter.
854

867 Table 2. IncI1 copy number in *S*. Heidelberg isolates recovered from fresh and reused litter.

S. Heidelberg Strain ID	Time (h)	pH of cecal extract	IncI1 copy number/cell[a](S.D)	Coefficient of variation (%)
SH-16-0-5B-RL	2	6.52	0.001±0.001	155.8
SH-IncI1-FL	2	6.52	5.594±0.582	10.4
SH-16-0-5B-RL	0.5	2.52	0.001±0.0003	47.9
	4	2.52	0.141±0.064	46.0
	24	2.52	0.084±0.038	46.4

SH-Incl1-FL	0.5	2.52	6.304±0.165	2.6
	4	2.52	6.301±0.343	5.4
	24	2.52	5.892±0.378	6.4

868 [a]Three individual *S*. Heidelberg isolates from fresh litter (FL) were used to determine the copy number of
869 Incl1 for SH-Incl1-FL, while Incl1 copy number in *S*. Heidelberg strain SH-16-0-5B from reused litter (RL)
870 was determined using three replicates.

876
877

Figure Captions

Fig 1. *S*. Heidelberg concentration in the ceca and litter of broiler chicks. Box plot
of nal[R] *S*. Heidelberg concentration in the (**a**) ceca (n=10 for oral and cloacal, n=15 for
seeder) (**b**) and litter (n = 2 subsamples from pooled litter) of chicks raised on fresh or
reused litter ($P > 0.05$; Wilcoxon signed-rank test). Cecal and litter samples were

collected 14 days after challenging chicks with nal[R] *S.* Heidelberg through oral gavage,
cloacal inoculation or through the seeder method.
895

Fig 2. Broiler chicks raised on reused litter carried lower levels of antimicrobial resistant *S.* Heidelberg. (**a**) Distribution of antibiotic resistance acquired by nal[R] *S.* Heidelberg isolates from the ceca and litter of chicks on fresh (n=158) and reused litter (n=141). (**b**) Maximum likelihood tree constructed using accessory genes present in *S.* Heidelberg isolates (*n* = 30) recovered from the ceca and litter of cloacally inoculated chicks raised on fresh and reused litter. The GTR model of nucleotide substitution and the GAMMA model of rate heterogeneity were used for sequence evolution prediction. Numbers shown next to the branches represent the percentage of replicate trees where associated isolates cluster together based on ~100 bootstrap replicates. All *S.* Heidelberg strains were assembled using Illumina short reads except SH4-3A and SH-ancestral that were assembled by combining Illumina short reads with PacBio or MinION long reads. The tree was rooted with the ancestral *S.* Heidelberg strain (SH-ancestral, Genbank accession no: CP066851). Green colored subclade shows strains that harbored IncI1-pST26. All isolates are expected to be resistant to Nalidixic acid. (Nal, nalidixic acid; Gen, gentamicin; Str, streptomycin; Tet, tetracycline; Fox, Cefoxitin; Cip, Ciprofloxacin)

Fig 3. Reused litter samples harbored a uniform and diverse microbiome than fresh litter samples. (**a**) Principal coordinates analysis of Bray-Curtis distances based on 16S rRNA gene libraries obtained from ceca and litter samples. Each point is values from individual libraries with colors denoting the route used for *S.* Heidelberg challenge for respective samples. (b) Average alpha diversity indices of rarefied ceca and litter samples grouped by the litter age/type and the route of *S.* Heidelberg challenge. Boxes indicate the interquartile range (75th to 25th) of the data. Whiskers extend to the most extreme value within 1.5 * interquartile range and dots represent outliers beyond that range. Black bar represents the median. (**c**) Phylum-level classification of 16S rRNA gene sequence reads in each ceca and litter sample grouped by the route of *S.* Heidelberg challenge. 16S rRNA gene sequencing was performed on the ceca (59 chicks from fresh litter (n = 30) and reused litter (n = 29)) and litter (n = 22 (11 samples each from fresh and reused litter)).

Fig 4. *S.* Heidelberg challenge modulated the microbiome of the ceca and litter of broiler chicks. Plots of amplicon sequence variants (ASVs) that were significantly differentially abundant ($p_{adj} < .05$) in the (**a**) ceca and (**b**) litter of challenged chicks on fresh and reused litter compared to uninoculated controls. Significant ASVs are plotted individually and colored according to their family-level classification. The route of *S.* Heidelberg challenge is shown on the edge of each plot as colored squares.

Fig 5. Enterobacteriaceae and Clostridiales were the bacterial hosts of AMR in broiler chicks. AMR gene hosts found by Hi-C contacts. AMR genes (horizontal axis of heatmap) associated with metagenome-assembled genomes (MAGs) present in cecal samples from chicks on fresh litter (Hi-C-FL1 and Hi-C-FL2) and reused litter (Hi-C-RL1 and Hi-C-RL2). MAGs and AMR gene sources were derived from Hi-C based

deconvolution of the metagenome assembly and placed into a bacterial phylogeny
using Mash and CheckM by the ProxiMeta platform. The MAG assigned to family
Enterobacteriaceae in each sample matches most closely to an *Escherichia*
coli genome. (Legend: Gold - AMR genes that are associated with plasmid sequences
on the same contigs, Cyan – AMR genes that are likely to be genomically integrated,
Black – no contact above statistical thresholds)

Fig 6. *Escherichia coli* was the main reservoir of plasmids and AMR in broiler chicks.
Pan genome analysis was performed on *E. coli* strains (n =19) recovered from the ceca
of broiler chicks challenged with *S.* Heidelberg and raised on fresh (FL) (n = 2) or
reused litter (RL) (n=2). Maximum likelihood tree constructed using the core (**b**) and
accessory (**c**) genes of the *E. coli* strains. Illumina short reads were combined with
either PacBio or MinION long reads to assemble the genomes of *E. coli* strains Ec-FL1-
1X, Ec-FL1-2X, Ec-FL1-3, Ec-RL2-1 and Ec-RL2-1X. GTR model of nucleotide
substitution and GAMMA model of rate heterogeneity was used for sequence evolution
prediction. Numbers shown next to the branches represent the percentage of replicate
trees where associated isolates cluster together based on ~100 bootstrap replicates.
Clade numbers were assigned arbitrarily to ease discussion on the differences in
phylogeny between isolates. Tree was rooted with Ec-RL2-1X. (Note: cgMLST – core
genome Multi Locus Sequence Type). Black rectangular box shows the *E. coli* strain
carrying identical IncI1-pST26 as *S.* Heidelberg.

Fig 7. Relative abundance of IncI1 in Enterobacteriaceae bacterial population of
the ceca. Box plot of the ratio of IncI1 gene copies/gram of ceca to gene copies of *gapA*
/gram of ceca for chicks raised on fresh (oral = 9, cloacal = 8, control = 9) and reused
(oral = 10, cloacal = 9, control = 10) litter. (***$P < 0.001$; Wilcoxon signed-rank test).

965
966
967
968

974   **Reference:**

975   1 Von Wintersdorff, C. J., Penders, J., Van Niekerk, J. M., Mills, N. D., Majumder, S.,

976   Van Alphen, L. B., Savelkoul, P. H., Wolffs, P. F. 2016 Dissemination of antimicrobial

resistance in microbial ecosystems through horizontal gene transfer. *Frontiers in microbiology*. **7**, 173.

Ventola, C. The antibiotic resistance crisis: part 1: causes and threats. PT 40 (4): 277–283. 2015.

Angulo, F. J., Baker, N. L., Olsen, S. J., Anderson, A., Barrett, T. J. Year Antimicrobial use in agriculture: controlling the transfer of antimicrobial resistance to humans. Seminars in pediatric infectious diseases; 2004: Elsevier; 2004. p. 78-85.

Tang, K. L., Caffrey, N. P., Nóbrega, D. B., Cork, S. C., Ronksley, P. E., Barkema, H. W., Polachek, A. J., Ganshorn, H., Sharma, N., Kellner, J. D. 2017 Restricting the use of antibiotics in food-producing animals and its associations with antibiotic resistance in food-producing animals and human beings: a systematic review and meta-analysis. *The Lancet Planetary Health*. **1**, e316-e327.

Singer, R., Porter, L. Estimates of On-Farm Antimicrobial Usage in Broiler Chicken and Turkey Production in the United States, 2013-2017. . 2019 [cited; Available from: Retrieved from https://www.uspoultry.org/poultry-antimicrobial-use-report/docs/USPOULTRY_Antimicrobial-Report.pdf

Kirchhelle, C. 2018 Swann Song: Antibiotic Regulation in British Livestock Production (1953-2006). *B Hist Med*. **92**, 317-351. (10.1353/bhm.2018.0029)

Oladeinde, A., Abdo, Z., Press, M. O., Cook, K., Cox, N. A., Zwirzitz, B., Woyda, R., Lakin, S. M., Thomas, J. C. t., Looft, T.*, et al.* 2021 Horizontal Gene Transfer Is the Main Driver of Antimicrobial Resistance in Broiler Chicks Infected with Salmonella enterica Serovar Heidelberg. *mSystems*. **6**, e0072921. (10.1128/mSystems.00729-21)

8 Gieraltowski, L., Higa, J., Peralta, V., Green, A., Schwensohn, C., Rosen, H., Libby,

T., Kissler, B., Marsden-Haug, N., Booth, H.*, et al.* 2016 National Outbreak of Multidrug

Resistant Salmonella Heidelberg Infections Linked to a Single Poultry Company. *PLoS*

One. **11**, e0162369. (10.1371/journal.pone.0162369)

9 Edirmanasinghe, R., Finley, R., Parmley, E. J., Avery, B. P., Carson, C., Bekal, S.,

Golding, G., Mulvey, M. R. 2017 A Whole-Genome Sequencing Approach To Study

Cefoxitin-Resistant Salmonella enterica Serovar Heidelberg Isolates from Various

Sources. *Antimicrob Agents Chemother.* **61**, (10.1128/AAC.01919-16)

10 Ceccarelli, D., van Essen-Zandbergen, A., Smid, B., Veldman, K. T., Boender, G. J.,

Fischer, E. A. J., Mevius, D. J., van der Goot, J. A. 2017 Competitive Exclusion

Reduces Transmission and Excretion of Extended-Spectrum-beta-Lactamase-

Producing Escherichia coli in Broilers. *Appl Environ Microbiol.* **83**,

(10.1128/AEM.03439-16)

11 Chantziaras, I., Smet, A., Filippitzi, M. E., Damiaans, B., Haesebrouck, F., Boyen, F.,

Dewulf, J. 2018 The effect of a commercial competitive exclusion product on the

selection of enrofloxacin resistance in commensal E. coli in broilers. *Avian Pathology.*

47, 443-454.

12 Raffatellu, M. 2018 Learning from bacterial competition in the host to develop

antimicrobials. *Nature medicine.* **24**, 1097-1103.

13 Achard, C. S., Dupouy, V., Siviglia, S., Arpaillange, N., Cauquil, L., Bousquet-Mélou,

A., Zemb, O. 2019 Variability of the ability of complex microbial communities to exclude

microbes carrying antibiotic resistance genes in rabbits. *Frontiers in microbiology.* **10**,

1503.

14 Baumgartner, M., Bayer, F., Pfrunder-Cardozo, K. R., Buckling, A., Hall, A. R. 2020

Resident microbial communities inhibit growth and antibiotic-resistance evolution of

Escherichia coli in human gut microbiome samples. *PLoS Biol.* **18**, e3000465.

(10.1371/journal.pbio.3000465)

15 Dame-Korevaar, A., Fischer, E. A., van der Goot, J., Velkers, F., Ceccarelli, D.,

Mevius, D., Stegeman, A. 2020 Early life supply of competitive exclusion products

reduces colonization of extended spectrum beta-lactamase-producing Escherichia coli

in broilers. *Poultry science.* **99**, 4052-4064.

16 Methner, U., Rösler, U. 2020 Efficacy of a competitive exclusion culture against

extended-spectrum β-lactamase-producing Escherichia coli strains in broilers using a

seeder bird model. *BMC veterinary research.* **16**, 1-7.

17 Wang, L., Lilburn, M., Yu, Z. 2016 Intestinal Microbiota of Broiler Chickens As

Affected by Litter Management Regimens. *Front Microbiol.* **7**, 593.

(10.3389/fmicb.2016.00593)

18 Figueroa, A., Derksen, T., Biswas, S., Nazmi, A., Rejmanek, D., Crossley, B.,

Pandey, P., Gallardo, R. 2020 Persistence of low and highly pathogenic avian influenza

virus in reused poultry litter, effects of litter amendment use, and composting

temperatures. *Journal of Applied Poultry Research.* 100096.

19 Wadud, S., Michaelsen, A., Gallagher, E., Parcsi, G., Zemb, O., Stuetz, R.,

Manefield, M. 2012 Bacterial and fungal community composition over time in chicken

litter with high or low moisture content. *British poultry science.* **53**, 561-569.

20 Council, N. F. A. C. 2016 Code of Practice: For the care and handling of hatching

eggs, breeders, chickens, and turkeys.

21 ADAS, Ltd. 2016 Comparison of the Regulatory Framework and Key Practices in the

Poultry Meat Supply Chain in the EU and USA : a study for the association of poultry

processors

and poultry trade in the EU Countries.

22 Dornelas, K. C., Mascarenhas, N. M. H., Rodrigues, H. C. S., do Nascimento, R. T.,

Furtado, D. A., do Nascimento, J. W. B. 2020 Chicken bed: a review on reuse,

treatment and influence on ambience. *Poultry Science*.

23 Muniz, E., Mesa, D., Cuaspa, R., Souza, A. M., Santin, E. 2014 Presence of

Salmonella spp. in reused broiler litter. *Revista Colombiana de Ciencias Pecuarias*. **27**,

12-17.

24 Yamak, U. S., Sarica, M., Boz, M. A., Ucar, A. 2016 Effect of reusing litter on broiler

performance, foot-pad dermatitis and litter quality in chickens with different growth rates.

Kafkas Univ. Vet. Fak. J. **22**, 85-91.

25 Casey, R., Fairchild, B., Lacy, M. Litter quality and broiler performance. In: C. e. S.

Bulletin, ed. 1267 ed 2005.

26 Roll, V. F., Dai Pra, M. A., Roll, A. P. 2011 Research on Salmonella in broiler litter

reused for up to 14 consecutive flocks. *Poult Sci.* **90**, 2257-2262. (10.3382/ps.2011-

01583)

27 Zucko, J., Starcevic, A., Diminic, J., Oros, D., Mortazavian, A. M., Putnik, P. 2020

Probiotic–friend or foe? *Current Opinion in Food Science*. **32**, 45-49.

28 Gupta, S., Allen-Vercoe, E., Petrof, E. O. 2016 Fecal microbiota transplantation: in

perspective. *Therapeutic advances in gastroenterology*. **9**, 229-239.

29 Oladeinde, A., Cook, K., Orlek, A., Zock, G., Herrington, K., Cox, N., Plumblee

Lawrence, J., Hall, C. 2018 Hotspot mutations and ColE1 plasmids contribute to the

fitness of Salmonella Heidelberg in poultry litter. *PLoS One.* **13**, e0202286.

(10.1371/journal.pone.0202286)

30 Bucher, M. G., Zwirzitz, B., Oladeinde, A., Cook, K., Plymel, C., Zock, G., Lakin, S.,

Aggrey, S. E., Ritz, C., Looft, T.*, et al.* 2020 Reused poultry litter microbiome with

competitive exclusion potential against Salmonella Heidelberg. *J Environ Qual.* **49**, 869-

881. (10.1002/jeq2.20081)

31 Pfeifer, E., de Sousa, J. A. M., Touchon, M., Rocha, E. P. C. 2021 Bacteria have

numerous distinctive groups of phage-plasmids with conserved phage and variable

plasmid gene repertoires. *Nucleic Acids Research.* **49**, 2655-2673.

(10.1093/nar/gkab064)

32 Wertz, J. E., Goldstone, C., Gordon, D. M., Riley, M. A. 2003 A molecular phylogeny

of enteric bacteria and implications for a bacterial species concept. *J Evol Biol.* **16**,

1236-1248. (10.1046/j.1420-9101.2003.00612.x)

33 Wiedemann, S. 2015 Litter reuse: an evidence-based guide to reusing litter. Vol.

project PRJ-006440, publication 14/095. *Rural Industries Reserach and Development*

Corporation (RIRDC): Canberra, ACT). Available at https://www*. agrifutures. com.*

au/product/litter-reuse-an-evidence-based-guide-to-reusing-litter/[Verified   19   March

2019].

34 Cressman, M. D., Yu, Z., Nelson, M. C., Moeller, S. J., Lilburn, M. S., Zerby, H. N.

2010 Interrelations between the microbiotas in the litter and in the intestines of

commercial broiler chickens. *Appl Environ Microbiol.* **76**, 6572-6582.

(10.1128/AEM.00180-10)

35 Lee, K., Lillehoj, H., Jang, S., Lee, S., Bautista, D., Siragusa, G. 2013 Effect of

Bacillus subtilis-based direct-fed microbials on immune status in broiler chickens raised

on fresh or used litter. *Asian-Australasian journal of animal sciences.* **26**, 1592.

36 Stecher, B., Denzler, R., Maier, L., Bernet, F., Sanders, M. J., Pickard, D. J., Barthel,

M., Westendorf, A. M., Krogfelt, K. A., Walker, A. W. 2012 Gut inflammation can boost

horizontal gene transfer between pathogenic and commensal Enterobacteriaceae.

Proceedings of the National Academy of Sciences. **109**, 1269-1274.

37 Gagliardi, A., Totino, V., Cacciotti, F., Iebba, V., Neroni, B., Bonfiglio, G.,

Trancassini, M., Passariello, C., Pantanella, F., Schippa, S. 2018 Rebuilding the Gut

Microbiota Ecosystem. *Int J Environ Res Public Health.* **15**, (10.3390/ijerph15081679)

38 Montassier, E., Valdes-Mas, R., Batard, E., Zmora, N., Dori-Bachash, M., Suez, J.,

Elinav, E. 2021 Probiotics impact the antibiotic resistance gene reservoir along the

human GI tract in a person-specific and antibiotic-dependent manner. *Nat Microbiol.* **6**,

1043-1054. (10.1038/s41564-021-00920-0)

39 O'callaghan, A., Van Sinderen, D. 2016 Bifidobacteria and their role as members of

the human gut microbiota. *Frontiers in microbiology.* **7**, 925.

40 Casaburi, G., Frese, S. A. 2018 Colonization of breastfed infants by Bifidobacterium

longum subsp. infantis EVC001 reduces virulence gene abundance. *Human*

Microbiome Journal. **9**, 7-10.

41 Taft, D. H., Liu, J., Maldonado-Gomez, M. X., Akre, S., Huda, M. N., Ahmad, S. M.,

Stephensen, C. B., Mills, D. A. 2018 Bifidobacterial Dominance of the Gut in Early Life

and Acquisition of Antimicrobial Resistance. *mSphere.* **3**, (10.1128/mSphere.00441-18)

42 Casaburi, G., Duar, R. M., Vance, D. P., Mitchell, R., Contreras, L., Frese, S. A.,

Smilowitz, J. T., Underwood, M. A. 2019 Early-life gut microbiome modulation reduces

the abundance of antibiotic-resistant bacteria. *Antimicrobial Resistance & Infection*

Control. **8**, 1-18.

43 Wood, D. E., Lu, J., Langmead, B. 2019 Improved metagenomic analysis with

Kraken 2. *Genome biology.* **20**, 257.

44 Poulsen, L. L., Thofner, I., Bisgaard, M., Christensen, J. P., Olsen, R. H.,

Christensen, H. 2017 Longitudinal study of transmission of Escherichia coli from broiler

breeders to broilers. *Vet Microbiol.* **207**, 13-18. (10.1016/j.vetmic.2017.05.029)

45 de Toledo, T. D., Roll, A. A. P., Rutz, F., Dallmann, H. M., Pra, M. A. D., Leite, F. P.

L., Roll, V. F. B. 2020 Y An assessment of the impacts of litter treatments on the litter

quality and broiler performance: A systematic review and meta-analysis. *Plos One.* **15**,

(ARTN e0232853

10.1371/journal.pone.0232853)

46 Joerger, R. D., Ganguly, A., de Los Santos, M., Li, H. 2020 Effect of sodium bisulfate

amendments on bacterial populations in broiler litter. *Poult Sci.* **99**, 5560-5571.

(10.1016/j.psj.2020.08.013)

47 Brockhurst, M. A., Harrison, E. 2021 Ecological and evolutionary solutions to the

plasmid paradox. *Trends Microbiol.* (10.1016/j.tim.2021.11.001)

48 Virolle, C., Goldlust, K., Djermoun, S., Bigot, S., Lesterlin, C. 2020 Plasmid Transfer

by Conjugation in Gram-Negative Bacteria: From the Cellular to the Community Level.

Genes (Basel). **11**, (10.3390/genes11111239)

49 Lee, C., Kim, J., Shin, S. G., Hwang, S. 2006 Absolute and relative QPCR

quantification of plasmid copy number in Escherichia coli. *J Biotechnol.* **123**, 273-280.

(10.1016/j.jbiotec.2005.11.014)

50 Woyda, R. R., Oladeinde, A., Abdo, Z. 2020 Reads2Resistome: An adaptable and

high-throughput whole-genome sequencing pipeline for bacterial resistome

characterization. *bioRxiv.* 2020.2005.2018.102715. (10.1101/2020.05.18.102715)

51 Katoh, K., Standley, D. M. 2013 MAFFT multiple sequence alignment software

version 7: improvements in performance and usability. *Mol Biol Evol.* **30**, 772-780.

(10.1093/molbev/mst010)

52 Page, A. J., Cummins, C. A., Hunt, M., Wong, V. K., Reuter, S., Holden, M. T.,

Fookes, M., Falush, D., Keane, J. A., Parkhill, J. 2015 Roary: rapid large-scale

prokaryote pan genome analysis. *Bioinformatics.* **31**, 3691-3693.

(10.1093/bioinformatics/btv421)

53 Kozlov, A., Darriba, D., Flouri, T., Morel, B., Stamatakis, A. 2018 RAxML-NG: A fast,

scalable, and user-friendly tool for maximum likelihood phylogenetic inference. *bioRxiv.*

(10.1101/447110)

54 Bertelli, C., Laird, M. R., Williams, K. P., Simon Fraser University Research

Computing, G., Lau, B. Y., Hoad, G., Winsor, G. L., Brinkman, F. S. L. 2017

IslandViewer 4: expanded prediction of genomic islands for larger-scale datasets.

Nucleic Acids Res. **45**, W30-W35. (10.1093/nar/gkx343)

55 Li, H., Durbin, R. 2009 Fast and accurate short read alignment with Burrows-

Wheeler transform. *Bioinformatics.* **25**, 1754-1760. (10.1093/bioinformatics/btp324)

56 McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A.,

Garimella, K., Altshuler, D., Gabriel, S., Daly, M.*, et al.* 2010 The Genome Analysis

Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data.

Genome Res. **20**, 1297-1303. (10.1101/gr.107524.110)

57 Paulson, J. N., Stine, O. C., Bravo, H. C., Pop, M. 2013 Differential abundance

analysis for microbial marker-gene surveys. *Nat Methods.* **10**, 1200-1202.

(10.1038/nmeth.2658)

58 Nurk, S., Meleshko, D., Korobeynikov, A., Pevzner, P. A. 2017 metaSPAdes: a new

versatile metagenomic assembler. *Genome Res.* **27**, 824-834.

(10.1101/gr.213959.116)

59 Press, M. O., Wiser, A. H., Kronenberg, Z. N., Langford, K. W., Shakya, M., Lo, C.-

C., Mueller, K. A., Sullivan, S. T., Chain, P. S., Liachko, I. 2017 Hi-C deconvolution of a

human gut microbiome yields high-quality draft genomes and reveals plasmid-genome

interactions. *biorxiv.* 198713.

60 Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P., Tyson, G. W. 2015

CheckM: assessing the quality of microbial genomes recovered from isolates, single

cells, and metagenomes. *Genome Res.* **25**, 1043-1055. (10.1101/gr.186072.114)

61 Ondov, B. D., Treangen, T. J., Melsted, P., Mallonee, A. B., Bergman, N. H., Koren,

S., Phillippy, A. M. 2016 Mash: fast genome and metagenome distance estimation using

MinHash. *Genome Biol.* **17**, 132. (10.1186/s13059-016-0997-x)

62 Carattoli, A., Zankari, E., Garcia-Fernandez, A., Larsen, M. V., Lund, O., Villa, L.,

Aarestrup, F. M., Hasman, H. 2014 In Silico Detection and Typing of Plasmids using

PlasmidFinder and Plasmid Multilocus Sequence Typing. *Antimicrob Agents Ch.* **58**,

3895-3903. (10.1128/Aac.02412-14)

63 Seemann, T. 2014 Prokka: rapid prokaryotic genome annotation. *Bioinformatics.* **30**,

2068-2069. (10.1093/bioinformatics/btu153)

64 Paulson, J. N., Stine, O. C., Bravo, H. C., Pop, M. 2013 Differential abundance

analysis for microbial marker-gene surveys. *Nature methods.* **10**, 1200.

65 Csardi, G., Nepusz, T. 2006 The igraph software package for complex network

research. *InterJournal, complex systems.* **1695**, 1-9.

66 Rothrock, M. J., Guard, J. Y., Oladeinde, A. 2021 Salmonella Diversity Along the

Farm-to-Fork Continuum of Pastured Poultry Flocks in the Southeastern United States.

Frontiers in Animal Science. **2**, (10.3389/fanim.2021.761930)

1190

1191

1192

1193

1194

1195

1196

1197

1198

1199

1200

1201

1202

1203

1204
1205
1206

A

B

Strain	Litter age/type	Isolation Source	AMR phenotype	Acquired ARG	Plasmids	cgMLST
SH4-2A	Fresh	Ceca	Nal	None	IncX1	186980
SH-FL2-0-5A	Fresh	Litter	Nal	None	IncX1	186980
SH20-4A	Reused	Ceca	Nal	None	IncX1	186980
SH-FL1-3A	Fresh	Litter	Nal	None	IncX1	186980
SH-FL1-2A	Fresh	Litter	Nal	None	IncX1	186980
SH6-2B	Fresh	Ceca	Nal	None	IncX1	186980
SH6-1C	Fresh	Ceca	Nal	None	IncX1	186980
SH6-2A	Fresh	Ceca	Nal	None	IncX1	186980
SH16-1B	Reused	Ceca	Nal	None	IncX1	186980
SH16-1A	Reused	Ceca	Nal	None	IncX1	186980
SH16-0-5C	Reused	Ceca	Nal	None	IncX1	186980
SH20-0-5C	Reused	Ceca	Nal	None	IncX1	186980
SH20-4C	Reused	Ceca	Nal	None	IncX1	186980
SH-RL1-0-5A	Reused	Litter	Nal, Str	None	IncX1	186980
SH-RL2-4A	Reused	Litter	Nal	None	IncX1	186980
SH-RL2-0-5A	Reused	Litter	Nal	None	IncX1	186980
SH6-2C	Fresh	Ceca	Nal	None	IncX1	186980
SH16-0-5A	Reused	Ceca	Nal	None	IncX1	186980
SH20-0-5A	Reused	Ceca	Nal	None	IncX1	186980
SH6-1B	Fresh	Ceca	Nal	None	IncX1	186980
SH20-0-5B	Reused	Ceca	Nal	None	IncX1,Col8282, ColRNAI	186980
SH4-0-5A	Fresh	Ceca	Nal	None	IncX1,Col8282, ColRNAI	186980
SH4-1A	Fresh	Ceca	Nal	None	IncX1, Col8282, ColRNAI	186980
SH4-0-5B	Fresh	Ceca	Nal	None	IncX1,Col8282, ColRNAI	171575
SH-FL1-0-5A	Fresh	Litter	Nal, Gen, Str, Tet	aac(3)-Via, aadA1, tetA	IncX1, IncI1	186980
SH16--0-5B	Reused	Ceca	Nal	aac(3)-Via, aadA1, tetA	IncX1, IncI1, Col8282	186980
SH4-3A	Fresh	Ceca	Nal, Gen, Str, Tet	aac(3)-Via, aadA1, tetA	IncX1, IncI1, Col8282	186980
SH4-1B	Fresh	Ceca	Nal, Gen, Str, Tet	aac(3)-Via, aadA1, tetA	IncX1,IncI1	186980
SH6-1A	Fresh	Ceca	Nal, Gen, Str, Tet	aac(3)-Via, aadA1, tetA	IncX1, IncI1, Col8282	186980
SH-ancestral	NA	NA	Nal	None	IncX1	171516

0.08

A

B

C

A

B