Heterogeneous non-canonical nucleosomes predominate in yeast cells *in situ*

Zhi Yang Tan[1,3], Shujun Cai (蔡舒君)[1,3], Alex J. Noble[2,3], Jon K. Chen[1,3], Jian Shi[1], and

Lu Gan (甘露)[1,4,*]

[1] Department of Biological Sciences and Center for BioImaging Sciences, National

University of Singapore, Singapore 117543

[2] National Resource for Automated Molecular Microscopy, Simons Electron Microscopy

Center, New York Structural Biology Center, New York, United States

[3] These authors contributed equally

[4] Lead contact

* Correspondence: lu@anaphase.org, @MitosisLab

SUMMARY

Nuclear processes depend on the organization of chromatin, whose basic units are

cylinder-shaped complexes called nucleosomes. A subset of mammalian nucleosomes

in situ (inside cells) resembles the canonical structure determined *in vitro* 25 years ago.

Nucleosome structure *in situ* is otherwise poorly understood. Using cryo-ET and 3-D

classification analysis of yeast cells, here we find that canonical nucleosomes account

for less than 10% of total nucleosomes expected *in situ*. In a strain in which H2A-GFP is

the sole source of histone H2A, class averages that resemble canonical nucleosomes

both with and without an extra density are found *ex vivo*, but not *in situ*. These data

suggest that the yeast intranuclear environment favors multiple non-canonical

nucleosome conformations. Using the structural observations here and the results of

previous genomics and biochemical studies, we propose a model in which the average

yeast nucleosome's DNA is partially detached *in situ*.

Keywords

chromatin, nucleosome, yeast, GFP, cryo-ET, cryo-sectioning, cryo-FIB milling

INTRODUCTION

Eukaryotic chromosomes are polymers of DNA-protein complexes called nucleosomes. An octamer of proteins, consisting of a heterotetramer of histones H3 and H4 and two heterodimers of histones H2A and H2B, resides at the nucleosome's center (Luger et al., 1997). Canonical nucleosomes resemble 10 nm wide, 6 nm thick cylinders and have 145 – 147 base pairs of DNA bent in 1.65 left-handed superhelical gyres around the histone octamer (Zhou et al., 2019; Zlatanova et al., 2009). In contrast, non-canonical nucleosomes have either partially detached DNA, partially detached histones, fewer than 8 histones, or a combination of these features (Zlatanova et al., 2009). Both X-ray crystallography and single-particle cryo-EM have shown that reconstituted nucleosomes, either alone or within a complex, are largely canonical *in vitro* (Zhou et al., 2019). Nucleosome structures *in situ* inside cells remain mysterious.

Molecular-resolution (2 – 4 nm) studies of unique objects like cells may be obtained by cryo-electron tomography (cryo-ET), a form of cryo-electron microscopy (cryo-EM) that generates 3-D reconstructions called cryotomograms. These studies reveal life-like snapshots of macromolecular complexes because the samples are prepared and then imaged in an unfixed, unstained, frozen-hydrated state. Most eukaryotic cells are too thick for cryo-ET, so thinner frozen-hydrated samples are made by cutting by cryomicrotomy or thinning by cryo-focused ion beam (cryo-FIB) milling (Ng and Gan, 2020; Strunk et al., 2012). These two approaches respectively produce cryosections and plank-like samples called cryolamellae. Subvolumes called subtomograms contain independent copies of the cells' macromolecular complexes. These subtomograms can

be further studied by averaging, which increases the signal-to-noise ratio, and

classification, which facilitates the analysis of heterogeneity. Large macromolecular

complexes such as ribosomes and proteasomes have been identified *in situ* by 3-D

classification followed by comparison of the class averages to known structures – an

approach called purification *in silico* (Beck and Baumeister, 2016).

Using the purification *in silico* approach, we previously showed that canonical

nucleosomes exist in cryotomograms of yeast cell lysates *ex vivo* and in a HeLa cell

cryolamella (Cai et al., 2018a; Cai et al., 2018b; Cai et al., 2018c). Herein, we use the

term *ex vivo* to describe nucleosomes from lysates instead of the term *in vitro*, which is

more commonly used to describe either reconstituted or purified mononucleosomes.

However, our 3-D structural analysis did not generate canonical nucleosome structures

from cryosectioned fission yeast cells (Cai et al., 2018b). The discrepancy between

nucleosome class averages *ex vivo* and *in situ* could have either technical or biological

origins. As a biological explanation for the absence of canonical nucleosome class

averages *in situ*, we hypothesized that yeast nucleosomes are either conformationally

or constitutionally heterogeneous.

In this work, we test this heterogenous-nucleosome hypothesis by using cryo-ET to

image both wild-type cells and strains that have nucleosomes bearing GFP as a density

tag. We use the budding yeast *S. cerevisiae*, herein called yeast, because it has only

two copies of each histone gene and because it is more amenable to gene editing. Our

work compares the chromatin in both lysates and thin cellular cryo-EM samples

prepared using primarily cryo-FIB milling. To obtain more information about

nucleosomes *in situ*, we create one strain in which H2A-GFP is the sole source of H2A,

meaning that the nucleosomes are expected to project one or two extra densities from

their surface. Canonical nucleosomes are abundant in nuclear lysates and, in the H2A-

GFP-expressing strain lysates, have extra densities consistent with GFP. In contrast,

canonical nucleosomes account for less than 10% of the expected number of total

nucleosomes in wild-type cell cryolamellae. Furthermore, neither canonical

nucleosomes nor nucleosome-like particles with extra protruding densities were

detected in the H2A-GFP-expressing strain. These findings suggest that the yeast

intracellular environment disfavors the canonical nucleosome conformation.

RESULTS

Canonical nucleosomes are abundant in wild-type lysates

The crystal structure of the reconstituted yeast nucleosome (White et al., 2001) shows a

canonical structure that is largely indistinguishable from the first published one (Luger et

al., 1997). To describe the various views of the nucleosome, herein we use the compact

nomenclature introduced by (Zhou et al., 2019): the disc view is along the superhelical

axis, and the gyre view is along the pseudo-dyad axis (Figure 1A). The side view, which

was not defined by Zhou et al, is orthogonal to both the disc and gyre views.

The original RELION subtomogram-analysis workflow (Bharat and Scheres, 2016)

involves 2-D classification, followed by 3-D classification (Figure S1A), whereas in the

alternative approach, 2-D classification is bypassed and subtomograms are subjected

directly to 3-D classification (Figure S1B); for brevity, we term this alternative method

"direct 3-D classification". Direct 3-D classification is limited by computer hardware

(Kimanius et al., 2016), but can detect more canonical nucleosomes (Cai et al., 2018a);

see the Methods for more details. In this study, the original workflow is used on a subset

of samples to show example 2-D class averages for comparison with other studies.

However, the conclusions in this paper are drawn from direct 3-D classification, which

detects more canonical nucleosomes.

Our previous subtomogram analysis of nuclear lysates (Cai et al., 2018c) revealed that

nucleosomes from the wild-type strain YEF473A (Bi and Pringle, 1996) adopt the

canonical structure *ex vivo*. We repeated this experiment on the strain BY4741

(Brachmann et al., 1998), which serves as the wild-type and parent strain for the

histone-GFP tagging mutants described later. In this experiment, yeast nuclei are

isolated, lysed, then deposited on an EM grid. Cryotomograms of BY4741 nuclear

lysates reveal the crowded nucleosome-like particles and other cellular debris (Figure

S2). Next, we template matched for nucleosome-like particles using a nucleosome-

sized featureless cylinder as a reference. Direct 3-D classification (Figure S3, A and B)

followed by 3-D refinement produced an 18 Å-resolution subtomogram average of a

BY4741 canonical nucleosome class (Figure S3C). The subtomogram average of wild-

type yeast nucleosomes in nuclear lysates has longer linker DNA (Figure 1B) than the

crystal structure, which was reconstituted using a 146-bp DNA fragment.

Canonical nucleosomes are rare in wild-type cells *in situ*

Cryo-FIB milling is a compression-free method to thin frozen-hydrated cells (Hayles et

al., 2007; Mahamid et al., 2015; Marko et al., 2006; Medeiros et al., 2018; Rigort et al.,

2010; Villa et al., 2013). This technique uses a beam of gallium ions to thin a cell under

cryogenic conditions, producing a frozen-hydrated plank-like cellular sample called a

cryolamella. Canonical nucleosomes are detectable in a HeLa cell cryolamella (Cai et

al., 2018a), meaning that cryo-FIB milling does not grossly perturb canonical

nucleosomes *in situ*. We prepared cryolamellae of wild-type BY4741 cells.

Cryotomograms of yeast cryolamellae showed that nuclei were packed with

nucleosome-like particles (Figure S4). Two-dimensional class averages of template-

matched nucleosome-like particles reveal densities that have the approximate size and

shape of nucleosomes (Figure S5A). We then subjected the particles belonging to the

most nucleosome-like 2-D class averages to 3-D classification, following the original

RELION classification workflow. However, none of these 3-D class averages resemble

canonical nucleosomes (Figure S5B). While many of the nucleosome-like class

averages have dimensions similar to the canonical nucleosome, none of them have

densities that resemble the distinctive 1.65 left-handed gyres of DNA. To rule out the

possibility that canonical nucleosomes were missed during 2-D classification, we

performed direct 3-D classification using 100 classes. None of the resultant 100 class

averages resemble a canonical nucleosome (Figure S6).

To determine if our template-matching and classification workflow missed the canonical

nucleosomes, we re-did both template matching and 3-D classification of BY4741 wild-

type yeast cryolamellae with intentionally biased references. Instead of a featureless

cylinder, we used the yeast nucleosome crystal structure (White et al., 2001) as the

reference (Figure S7A). If canonical nucleosomes are abundant in yeast, they should be

detected as a class average that resembles a low-resolution nucleosome crystal

structure. No canonical nucleosome class averages were seen in this control

experiment (Figure S7B and C).

Our previous study of a HeLa cell (Cai et al., 2018a), which detected canonical

nucleosomes, used a Volta phase plate (VPP). VPP data has more low-resolution

contrast than defocus phase-contrast data. To test if canonical nucleosomes in yeast

cryolamellae are detectable in VPP data, we recorded VPP tilt series and reconstructed

tomograms of BY4741 cell cryolamellae (Figure S8). Subtomogram analysis of the VPP

tomograms by 2-D classification (Figure S9A), followed by 3-D classification did not

reveal a nucleosome class average in BY4741 (Figure S9B). When we performed direct

3-D classification using 100 classes, we detected one class average that resembles a

canonical nucleosome (Figure S10A, Movie S1). A second round of classification

revealed two types of class averages, one of which resembles a slightly elongated

nucleosome (Figure S10B, Movie S2). Refinement of these two types of nucleosomes

produced 24 Å-resolution averages that differ in the amount of linker DNA visible

(Figures 1C and S10C). The class average that has more ordered linker DNA vaguely

resembles the chromatosome, a form of the nucleosome that has linker DNA crossing

at the entry-exit site and in contact with a linker histone (Bednar et al., 2017; Zhou et al.,

2015). However, the DNA does not appear to cross at the DNA entry-exit site, and, at

the present resolution, it is not possible to determine if the linker histone is present.

Unlike plunge-frozen complexes, which interact with the air-water interface and have

biased orientations (Noble et al., 2018), complexes *in situ* are not subject to such

biases. Visualization of the angular distribution of the canonical nucleosomes shows

that the disc views are undersampled and likely missed by the classification analysis

(Figure S10D), which we also observed in our analysis of HeLa nucleosomes *in situ*

(Cai et al., 2018a). This missing hemisphere of views results in roughly half of the

canonical nucleosomes going undetected. In summary, the use of VPP and relatively

thin (≤ 160 nm) cryolamellae made it possible to detect two canonical nucleosome

classes that differ in the amount of ordered linker DNA, similar to what we observed in a

HeLa cell (Cai et al., 2018a).

Following our previous work (Cai et al., 2018a), we performed a negative control by

analyzing a tomogram of a region in the cytoplasm (Figure S11), which does not have

any nucleosomes. We performed template matching using the same reference as for

our analysis of nuclei and then direct 3-D classification into 100 classes. None of the

resultant class averages resemble a canonical nucleosome (Figure S12), confirming

that our analysis was not biased by the cylindrical reference.

Our classification detected only 769 canonical nucleosomes. If we account for the

undersampling of disc views (Figure S10D), we estimate there are ~1,500 canonical

nucleosomes detected in the 5 tomograms. In comparison, we estimate that in the

single HeLa cryolamella cryotomogram (Cai et al., 2018a), there were more than 2,000

nucleosomes in a nucleus volume ~1/6th of the total analyzed here in BY4741. The

percentage of HeLa nucleosomes that are non-canonical *in situ* is unknown and will

require further study. To visualize the distribution of canonical nucleosomes, we

remapped the two class averages back into their positions in the original tomogram that

contains the largest number of canonical nucleosomes (Figure 2). The remapped model

shows that the canonical nucleosomes are scattered throughout the sampled nuclear

volume. There are no large clusters of canonical nucleosomes like what we saw near

the nuclear envelope of a HeLa cell. Using experimentally determined values for

nucleosome number and chromatin volume (Oberbeckmann et al., 2019; Uchida et al.,

2011), the tomograms we analyzed are expected to hold 25,000 nucleosomes.

Therefore, the nucleosomes (both canonical and non-canonical) should pack with an

order-of-magnitude higher density than visualized in our remapped model (Figure 2B).

In summary, our calculations suggest that the vast majority (> 90%) of the nucleosomes

in BY4741 yeast are non-canonical.

Histone GFP-tagging strategy

Our previous cryo-ET analysis of nucleosomes in a HeLa cell revealed that

subtomogram 3-D classification is sensitive to features much smaller than the

nucleosomes, as evidenced by the separation of canonical-nucleosome class averages

that differ by ~10 bp of linker DNA near the dyad (Cai et al., 2018a). Furthermore,

studies of flagella (Oda and Kikkawa, 2013) and pilus machines (Chang et al., 2016)

showed that subtomogram averages of complexes *in situ* can reveal either the presence

or absence of protein densities as small as fluorescent proteins. These observations led

us to attempt to use the GFP tag to facilitate nucleosome identification *in situ*. Our

strategy is to compare subtomogram averages of nucleosome-like particles in strains

that express only wild-type histones versus those that express only GFP-tagged

histones. Note that this tagging strategy did *not* work as intended because we could not

detect tagged nucleosomes 3-D classes *in situ*. However, this negative result provided

an important clue about the nature of nucleosomes inside yeast cells (see below).

Histones can accept a genetically encoded GFP tag at either the N- or C-terminus. An

N-terminal GFP tag is not expected to be visible in subtomogram averages because it

would be separated from the histone's globular domain by the long, flexible N-terminal

"tail". Therefore, we chose to tag the histone C-terminus, which does not have a flexible

tail (Figure S13, A and B). To further confine the tag, we eliminated the peptide linker

that is included in popular GFP-tagging modules (Figure S13C). *S. cerevisiae* contains

two copies of each histone (Figure S13D), which are arranged as gene pairs. To

maximize our chances of detecting nucleosome class averages that have an extra

density, we sought to create strains in which a histone-GFP fusion is the sole source of

one class of histones (Figure S13, E and F).

In budding yeast, the H2A and H2B genes are arranged as gene pairs *HTA1-HTB1* and

HTA2-HTB2 (Hereford et al., 1979). We first deleted the entire *HTA2-HTB2* gene pair to

generate LGY0012. This deletion prevents *HTA2-HTB2*, which is flanked by two

retrotransposon elements, from being amplified in circular DNA molecules (Libuda and

Winston, 2006). We then inserted the GFP gene at the 3' end of *HTA1*, without a linker,

to generate LGY0016, making H2A-GFP the sole source of H2A. We confirmed

LGY0016's genotype by PCR analysis (Figure S14, A and B) and Sanger sequencing.

Immunoblots using anti-H2A or anti-GFP antibodies detected a single ~40 kilodalton

band, which is the expected size of the H2A-GFP fusion protein (Figure S14C).

Accordingly, the LGY0016 nuclei showed bright fluorescence (Figure S14D). We also

constructed a strain, LGY0015, that expresses an H2B-GFP fusion without a linker

peptide (Figure S14, E – H). LGY0015 also has an untagged copy of H2B. We were

unable to create strains that have either H2B-GFP, H3-GFP, or H4-GFP as the sole

H2B, H3, and H4 sources, respectively. Consistent with this low tolerance for a GFP-

tagged histone as the sole source of a histone type, the LGY0016 doubling time is

~50% longer than for wild-type BY4741 (130 minutes versus 85 minutes) in rich media.

GFP is detectable on nucleosome subtomogram averages *ex vivo*

To determine if GFP tags are detectable on the nucleosome's surface, we imaged the

nuclear lysates of LGY0016 and LGY0015 cells by cryo-ET (Figures S15 and S16). The

2-D class averages of LGY0016 and LGY0015 nuclear lysates resemble those seen in

single-particle cryo-EM studies of reconstituted nucleosomes (Chua et al., 2016),

though with lower-resolution features (Figure 3A, S17A). Note that 2-D classification

uses a circular mask, meaning that it will not bias the shape of the resultant class

averages to resemble, for example, the double-lined motifs seen in nucleosome side

and gyre views. To increase the number of detected nucleosomes, we used direct 3-D

classification into 40 classes. We obtained canonical nucleosome 3-D class averages

this way in the lysates of both strains (Figure S18 and S19; Movies S3 and S4). These

class averages have the unmistakable structural motifs of canonical nucleosomes, such

as a 10-nm diameter, 6-nm thick cylindrical shape, and the left-handed path of the DNA

densities. There is more DNA than the crystal structure's 1.65 gyres because lysate

samples have linker DNA. All these properties are consistent with the subtomogram

analysis of nucleosomes from wild-type strains BY4741 (Figure 1B) and YEF473A (Cai

et al., 2018c).

Subsequent classification rounds revealed nucleosome class averages that had an

extra density projecting from one or both faces (Figures S18B and S19B). One example

of each class (zero, one, or two extra densities) was refined to ~25 Å resolution (Figure

3B, S17B). The position of the extra density is consistent with the location of the H2A C-

terminus being closer to the DNA entry-exit point (Figure 3C) and the H2B C-terminus

being far from the DNA entry-exit point (Figure S17C). In principle, LGY0015 cells can

assemble nucleosomes that have two copies of H2B-GFP. The absence of LGY0015

nucleosome classes with two densities suggest that they are either unstable or too rare

to detect by 3-D classification. We focused our *in situ* cryo-ET analysis on LGY0016

cells because the GFP tags are easier to recognize on nucleosomes from nuclear

lysates of this strain.

Canonical nucleosome classes are not detected in LGY0016 cells *in situ*

In an attempt to detect more yeast nucleosomes *in situ*, we did three types of imaging

experiments on LGY0016 cells. We performed cryo-ET of cryolamellae with and without

the VPP (Figures S20 and S21) and we also imaged cryosections with the VPP (Figure

S22). The cryolamellae benefit from the absence of compression artifacts while the

cryosections benefit from being thinner on average. We performed template matching

and then subjected the hits directly to 3-D classification, which was needed to detect

canonical nucleosomes in BY4741 above. The 3-D class averages from all three

samples resembled neither canonical nucleosomes nor cylindrical bodies with one or

more protruding densities that is expected from the lysate samples (Figures 4, S23,

S24, Movie S5). Some class averages have two linear motifs that resemble the double

DNA gyres opposite the DNA entry-exit point, but these DNA-like densities do not go

1.65 times around the center of mass as expected of canonical nucleosomes (Luger et

al., 1997). Small variations in the classification parameters such as mask size, class

number did not reveal any canonical nucleosome classes in LGY0016 cell

cryotomograms. We therefore conclude that canonical nucleosomes are rare inside

budding yeast nuclei and that non-canonical nucleosomes are the vast majority.

Ribosome control for sample, data, and analysis pathologies

We may have missed canonical nucleosomes if there were pathologies with either our

lamellae, data, or workflow, which would result in grossly misclassified and misaligned

particles. To test this hypothesis, we performed template matching, 3-D classification

and alignment on cytoplasmic ribosomes. Because ribosomes are so large (> 3

megadaltons) and have been studied extensively *in situ*, any of the pathologies listed

above would result in either the absence of ribosome class averages or extremely low

resolution. Using our control tomogram of the cytoplasm, which is densely packed with

ribosomes, we obtained a subtomogram average at between ~33 Å resolution based on

the Fourier shell correlation (FSC) = 0.5 cutoff criterion (or 28 Å using the FSC = 0.143

criterion), from 1,150 particles (Figure S25). As a comparison, we simulated densities

with the yeast ribosome crystal structure between 15 Å and 30 Å resolution and found

that our average has density features consistent with this resolution range. The

resolutions of our nucleosome and ribosome averages (~24 Å and 33 Å) are

comparable to recent *in situ* subtomogram averages using similar numbers of particles

(~500 to 1,500) (Laughlin et al., 2022; van den Hoek et al., 2022). Therefore, our

lamellae, data, and workflow do not show evidence of pathologies.

GFP tagging of H3 and H4 also does not reveal nucleosomes

In cryolamellae of LGY0016 cells, the absence of nucleosome-like class averages that

have an extra density bump suggested that previously unappreciated properties of the

H2A-H2B heterodimer make this histone pair a poor candidate for the GFP fusion tag

strategy (see Discussion). We therefore revisited GFP tagging of H3 and H4, which are

found in all known and speculated forms of nucleosomes and non-canonical

nucleosomes (Zlatanova et al., 2009). Because all attempts to make H3- or H4-GFP

"sole source" strains failed, we tested strains that had one wild-type copy and one GFP-

tagged copy of one of these histones. We tagged members of the *HHF1-HHT1* gene

pair, which encode histone H4 (*HHF1*) and H3 (*HHT1*). *HHF1-HHT1* is flanked by the

same transposon elements as *HTA2-HTB2*, so gene amplification here would not result

in the wild-type copy outnumbering the tagged copy. The genotypes and phenotypes

were validated by confirmation PCRs, western blots, and fluorescence confocal

microscopy (Figures S26 and S27). We created H3-GFP strains without a linker

(LGY0007) and with linker sequences RIPGLIN (LGY0002) and GGSGGS (LGY0070);

this latter linker was introduced previously (Verzijlbergen et al., 2010). For H4, we could

only obtain a strain with a H4-GFP-expressing strain by using the flexible GGSGGS

linker (LGY0071; Figure S27). Flexible linkers are not expected to facilitate tag-based

identification because the tag can occupy a much larger volume and get blurred out as

a result but was the only one tolerated in our H4 tagging attempts.

We next prepared nuclear lysates of each of these strains, performed cryo-ET, and

subjected them to direct 3-D classification analysis (Figures S28 – S31). Like the other

strains, the lysates of each of these four strains had large numbers of canonical

nucleosomes (Figures S28A, S29A, S30A, S31A). Of the canonical nucleosomes, we

were only able to detect class averages that have the extra density in the LGY0007

lysates (Figure S28); the nuclear lysates of the other strains do not have nucleosome

class averages with a GFP density. Finally, to test if nucleosomes, canonical or non-

canonical, with the GFP density bump, could be detected *in situ*, we performed cryo-ET

of LGY0007 cryolamellae with the VPP (Figure S32). We then performed template

matching and subjected the hits to direct 3-D classification analysis. Compared to the

BY4741 (wild type) cell cryolamella VPP class averages (Figure S10A), the 3-D class

averages in LGY0007 cell cryolamellae resemble neither canonical nucleosomes nor

cylindrical bodies with a protruding density (Figure S33). Therefore, H3- and H4-GFP

fusions cannot be used to detect non-canonical nucleosomes and much more work

needs to be done to identify non-canonical nucleosomes.

DISCUSSION

Cryo-EM of cells has made the structural analysis of chromatin organization *in situ*

feasible. The earliest *in situ* studies were done with projection cryo-EM images and 2-D

Fourier analysis of cryosectioned cells, which revealed that long-range order is absent

in chromatin *in situ* (Eltsov et al., 2008; McDowall et al., 1986). Cryo-ET studies later

showed that short-range order is present in isolated chicken erythrocyte nuclei (Scheffer

et al., 2011), but absent in picoplankton and budding yeast (Chen et al., 2016; Gan et

al., 2013). These early cryo-ET studies revealed little about nucleosome structure *in situ*

because, as Eltsov *et al* observed, the nucleosomes in those samples appeared like

smooth ellipsoids (Eltsov et al., 2018). Recent cryo-ET studies using state-of-the-art

electron-counting cameras and energy filters have revealed that in cryotomographic

slices of cell nuclei, there are densities that resemble nucleosome side and gyre views

in which gyre-like features are resolved (Cai et al., 2018a; Cai et al., 2018b; Eltsov et

al., 2018). This higher-fidelity data made it possible to use 3-D classification to detect

canonical nucleosomes in nuclear envelope-associated chromatin in a HeLa cell (Cai et

al., 2018a).

Our use of 3-D classification detected canonical nucleosomes in wild-type yeast nuclear

lysates and canonical nucleosomes both with and without GFP tags in nuclear lysates

of some strains that bear histone-GFP fusions. We also detected a canonical

nucleosome class average in wild-type cell cryolamellae imaged with a VPP. The *in situ*

detections suggest that there are only ~1,500 canonical nucleosomes (taking account of

the undersampling of nucleosomes in the disc view) out of the 25,000 nucleosomes

expected of the total sampled nuclear volume. Note that the percentage of canonical

nucleosomes in lysates cannot be accurately estimated because we cannot determine

how many chromosomes' worth of nucleosomes are in each field of view. The estimates

from the lamellae are more reliable because the expected numbers of nucleosomes

(canonical or not) can be estimated from genomics, biochemical, and X-ray tomography

data (see Methods). When we analyzed LGY0016 cryolamellae, in which H2A-GFP is

the sole source of H2A, we did not detect canonical nucleosome classes or any

cylindrical nucleosome-sized structures that have extra density bumps. Likewise,

analysis of cryolamellae of LGY0007 did not reveal either a canonical nucleosome-like

class average or any cylindrical nucleosome-sized density with an extra density bump,

even though H3-GFP should be incorporated into the H3-H4 tetramer, the central

component of the nucleosome.

The absence of either a canonical nucleosome-like class average that has an extra

density bump, or a cylinder with an extra density bump in the LGY0016 strain suggests

that the H2A-H2B heterodimer is mobile *in situ*. By "mobility", we are not implying that

H2A-H2B is dissociated. The H2A/H2B dimers are in contact with the H3/H4

heterotetramer because crosslinking and single-particle fluorescence imaging

experiments showed that approximately 80% of H2B – and presumably H2A to which it

stably dimerizes – is bound to chromatin *in situ* (Mohan et al., 2018; Ranjan et al.,

2020). H2A-H2B mobility was hinted at by the observations from X-ray crystallography

that yeast nucleosomes lack the hydrogen bonds that stabilize the two H2A-H2B

heterodimers in metazoan nucleosomes (White et al., 2001) and that interfaces

between H2A-H2B and H3 are moderately exposed to small-molecule probes (Marr et

al., 2021). Furthermore, recent simulations show that the H2A/H2B heterodimer may

adopt numerous orientations and small displacements, with either fully or partially

unwrapped DNA, while remaining in contact with H3-H4 (Ishida and Kono, 2022). We

also cannot rule out the possibility that expression of H2A-GFP nucleosome structure is

less ordered *in situ*.

Like LGY0016 (H2A-GFP sole source), none of the LGY0007 (H3 plus H3-GFP)

nucleosome-like class averages had an extra density consistent with the GFP density

tag, unlike the lysates. Furthermore, there were no canonical nucleosome class

averages – either without or with the GFP density – seen among the class averages of

LGY0007 lamella nucleosome-like particles. These observations suggest that the

expression of H3-GFP makes nucleosomes less ordered *in situ*. Altogether, our

experiments show that in budding yeast, canonical nucleosomes are rare *in situ* and

that the expression of GFP-tagged histones leads to further perturbations of

nucleosome structure *in situ*.

Our data is consistent with a model in which yeast canonical nucleosomes are abundant

ex vivo (Figure 5A). They adopt multiple non-canonical conformations *in situ* (Figure 5,

B and C) and rarely adopt the canonical one (Figure 5D). Note that the blurring in these

panels implies the heterogeneity in the positions of the DNA and proteins, not their

actual motion. In addition to H2A-H2B and DNA conformational heterogeneity, diverse

nucleosome-associated proteins that bind to multiple positions (Figure 5C) would further

increase the heterogeneity; in this figure, the blurring implies constitutional

heterogeneity, not motion. A high abundance of non-canonical nucleosomes means that

more of the genome would be accessible, consistent with yeast having high levels of

transcription. Nucleosome heterogeneity is also consistent with the absence of

chromatin long-range order *in situ* because crystalline oligonucleosome arrays can only

form if sequential nucleosomes adopt nearly identical conformations (Ekundayo et al.,

2017; Robinson et al., 2006; Song et al., 2014). Further investigation is needed to

identify the biochemical and biophysical factors responsible for the abundance of non-

canonical nucleosomes in yeast and to determine their diverse structures (Zlatanova et

al., 2009).

In combination with our previous work on a HeLa cell (Cai et al., 2018a), we show that

globular complexes as small as canonical nucleosomes (~200 kilodaltons) can be

detected by 3-D classification of subtomograms from cryolamellae that were imaged

with a VPP. Given the low throughput of cryo-FIB milling, the low yield of cryolamellae

thinner than 160 nm, and the technical difficulties of VPP imaging, it is not yet feasible

to systematically determine the conditions needed to detect sub-200 kDa complexes by

purification *in silico*. More depositions of eukaryote cryolamellae cryo-ET datasets in the

public database EMPIAR (Iudin et al., 2016) may enable a thorough search of data-

collection and processing-parameter space.

Our *in situ* cryo-ET-based model adds to the body of work on non-canonical

nucleosomes. ChIP-seq analysis has detected sub-nucleosomes *in situ*, though the

abundance was unknown (Rhee et al., 2014). MNase-seq has also detected

nucleosomes in a partially unwrapped and partially disassembled state *in situ*

(Ramachandran et al., 2017). A recent Hi-C variant "Hi-CO" presented evidence that <

147 bp of yeast nucleosomal DNA is protected from MNase attack (Ohno et al., 2019),

which is consistent with the partial detachment of core DNA. Molecular dynamics

advances in all-atom and coarse-grained simulations are now showing that

nucleosomes are far more dynamic than previously appreciated (Armeev et al., 2021;

Brandani et al., 2021; Farr et al., 2021; Huertas et al., 2021; Ishida and Kono, 2021).

The DNA-unwrapping associated with nucleosome breathing was shown to disfavor

ordered helical oligonucleosome structures (Farr et al., 2021). Nucleosome breathing is

also evident in high-throughput atomic-force microscopy experiments, which revealed

that only ~30% of nucleosomes are fully wrapped, and that approximately half of the

nucleosomes have an opening angle (measured between the entry/exit DNA arms) 60°

larger than the fully wrapped one (Konrad et al., 2021). Nucleosomes that have non-

canonical nucleosome properties, such as lower stability or exposure of internal

surfaces, have been reported in fission yeast (Koyama et al., 2017; Sanulli et al., 2019),

which may explain why we did not observe canonical nucleosomes in cryosections in

those cells either (Cai et al., 2018b). Some nucleosomes in human and fly cell

cryosections appear "gaping", i.e., with the inter-DNA-gyre distance slightly larger than

~2.7 nm (Eltsov et al., 2018). Partial DNA detachment has been seen in complexes

between nucleosomes and remodelers and transcription factors (Eustermann et al.,

2018; Farnung et al., 2017; Liu et al., 2017; Sundaramoorthy et al., 2018;

Sundaramoorthy et al., 2017; Willhoft et al., 2018), methyltransferases (Bilokapic and

Halic, 2019; Jang et al., 2019), or transcription-related complexes (Dodonova et al.,

2020; Kujirai et al., 2018). Larger amounts (up to ~25 bp) of DNA detachment have

been seen very rarely in cryo-EM structures (Bilokapic et al., 2018; Zhou et al., 2021). A

recent study of cryosectioned fly embryos has also presented evidence of nucleosome-

like structures such as hemisomes and three-gyre structures (Fatmaoui et al., 2022).

The prevalence and functional consequences of non-canonical nucleosomes *in situ*

remain to be studied in other organisms.

Alternative hypotheses

We consider the alternative hypothesis that canonical nucleosomes are the dominant

form *in situ* and that we have missed them as a result of our image processing. Many of

the 3-D class averages appear to have multiple gyre-like densities, which may arise

from nucleosome stacking. However, classes with multiple gyre-like densities are also

found in the class averages from the cytoplasm, which does not have nucleosomes

(Figure S12). These are "junk" classes that result from the averaging of different particle

species into the same class. Importantly, stacked canonical nucleosomes are so

conspicuous that they could not have been missed in tomographic slices. For instance,

two stacked canonical nucleosomes have twice the mass of a single canonical

nucleosome and would have dimensions 12 nm by 10 nm (Figure S34). Multiple

stacked canonical nucleosomes would appear like 10-nm-thick filaments.

Another alternative hypothesis is that the lack of disc views, corresponding to

nucleosomes whose "face" is parallel to the lamella surface, resulted in canonical

nucleosomes going undetected. This hypothesis would be supported only if the

orientations of canonical nucleosomes were biased. However, biased orientation in

cryo-EM experiments is usually a result of interactions with the air-water interface

(Noble et al., 2018), which does not exist for nucleosomes inside the nucleus of a cell.

Another hypothesis is that the missing views resulted in elongated class averages

because of the missing wedge or missing cone in Fourier space. This artifact would only

affect the distribution of canonical versus non-canonical nucleosomes if canonical

nucleosomes have a preferred orientation, which we have just argued against due to

the nature of our cellular samples. The class averages from non-disc-view nucleosomes

do not have a missing-wedge/cone artifact because they include nucleosomes whose

superhelical axes are in the X-Y plane; when averaged together, the Fourier transforms

of these nucleosomes "fill in" Fourier space because they sample numerous rotations

about their super-helical axes. Missing-wedge-free reconstructions are exemplified by

plunge-frozen actin filaments, which lie parallel to the EM grid. As illustrated in a recent

study (Merino et al., 2018), reconstructions of filamentous actin are free of missing-

wedge distortions even though the axial views are completely absent.

Some of the particles in the tomographic slices resemble donuts. Because nucleosome

disc views also resemble donuts, it is possible that we completely missed these

particles in either our template matching or classification analyses. As stated above,

even if we missed all the disc views, our conclusions would not change because the

orientations of canonical nucleosomes inside of cellular samples are not biased.

Furthermore, the donut-like particles cannot be canonical nucleosomes because they

are too wide (> 12 nm). A subset of them were detected by template matching and

classification and separated into their own class, which resembles a cylinder with a

rounded cap (Movie S5, row 5 column 1).

Another hypothesis for the low numbers of detected canonical nucleosomes is that the

nucleoplasm is too crowded. However, crowding is an unlikely explanation because we

were able to detect canonical nucleosome class averages in our most-crowded nuclear

lysates, which are so crowded that most nucleosomes are butted against others

(Figures S15 and S16).

Limitations of the study

Because naked linker DNA cannot yet be seen *in situ* in yeast, the connectivity between

the nucleosomes (both canonical and non-canonical) could not be followed. The

subtomogram analysis was unable to resolve the structures of the non-canonical

nucleosomes. Because of the low resolution, we could not assess the variability in the

structure and composition of the canonical nucleosomes. All of these limitations could

potentially be addressed by increasing the data quality and the numbers of classes and

subtomograms per class, which require advances at all stages of structural cell biology.

These advances include improvements in cryo-FIB milling throughput and

reproducibility (Buckley et al., 2020; Tacke et al., 2021; Zachs et al., 2020), cryo-EM

cameras, laser phase plates (Turnbaugh et al., 2021), template-matching /

segmentation software (Bepler et al., 2020; Moebel et al., 2021), and subtilt refinement

software such as emClarity (Himes and Zhang, 2018), EMAN2 (Chen et al., 2019),

Warp/M (Tegunov et al., 2021) and RELION 4 (Zivanov et al., 2022). Because structural

cell biology is still a nascent field, the optimum combination of sample prep, imaging,

and data analysis will require thorough exploration of the parameter space at each step.

The nucleosome Volta subtomogram averages presented here were limited to ~24 Å,

even though the data was recorded close to focus as suggested by a higher-resolution

study of purified ribosomes (Khoshouei et al., 2017). Limiting factors include lower

particle numbers, alignment inaccuracy, VPP charging, and the lack of contrast transfer

function (CTF) modeling. In single-particle cryo-EM analysis, high-resolution analysis (3

Å or better) requires accurate modeling and then compensation for the CTF. A key first

step of CTF modeling is the visualization of Thon rings in Fourier power spectra. We

plotted Fourier power spectra in both 2-D with CTFFIND (Rohou and Grigorieff, 2015)

and as a 1-D rotational average with IMOD (Mastronarde, 1997), but could not see

Thon rings (Figure S35), meaning that CTF compensation is not feasible. A study of

VPP single-particle cryo-EM analysis recommends intentionally underfocussing to 500

nm to increase the number of Thon rings, making CTF modeling easier (Danev et al.,

2017). However, the samples used in that study were much thinner (plunge-frozen

proteasomes) and the dose used per image was much higher (~40 electrons / $Å^2$).

Because the subtomogram analysis of smaller complexes inside cryolamellae faces

multiple challenges, more investigation is needed to determine the optimum parameters

for increased resolution.

Another challenge is the positive identification of the non-canonical nucleosome

species. The approaches attempted here (the use of a GFP tag) failed. An alternative

approach would be to remap the various averages and show that the linker DNA of

sequential non-canonical nucleosomes point to each other, as we have previously done

for HeLa chromatin *in situ* (Cai et al., 2018a). Unfortunately, this approach is not

suitable for yeast because only the canonical class averages have linker-DNA densities

– none of the other class averages have densities that resemble linker DNA. Because

non-canonical yeast nucleosomes *in situ* are unknown structures, we would have to

identify individual instances using a "GFP of cryo-EM", in the form of a compact fusion

protein-like structure that can be directly visualized in tomographic slices without

classification and subtomogram averaging. Several attempts have been made in recent

decades (Diestra et al., 2009; Mercogliano and DeRosier, 2007; Nishino et al., 2007;

Wang et al., 2011). More work is needed to test the suitability of such tags as "GFPs of

cryo-EM" *in situ*.

SUPPLEMENTAL INFORMATION

Supplemental information can be found online at doi-xyz

ACKNOWLEDGMENTS

We thank Kerry Bloom, Vu Nguyen, and Carl Wu for advice on fluorescent-protein tagging of histones; Bill Rice and Ed Eng for help with cryo-EM data collection; John Heumann for discussions about PEET and for implementing a parallelized duplicate-removal routine; Rado Danev, Kliment Verba, and Shenping Wu for advice on the VPP. The Quadro P6000 used in this work was kindly donated by the NVIDIA Corporation. ZYT, SC, JKC, and LG were supported by a Singapore Ministry of Education Tier 1 grant R-154-000-A49-114 and Tier 2 grant MOE2019-T2-1-140. AJN was supported by a grant from the NIH National Institute of General Medical Sciences (NIGMS) (F32GM128303). Some of this work was performed at the Simons Electron Microscopy Center and National Resource for Automated Molecular Microscopy located at the New York Structural Biology Center, supported by grants from the Simons Foundation (SF349247), NYSTAR, and the NIH National Institute of General Medical Sciences (GM103310) with additional support from NIH S10 RR029300.

AUTHOR CONTRIBUTIONS

SC, ZYT, AJN, JKC, LG – project design, experiments, writing; JS, LG – training.

DECLARATION OF INTERESTS

The authors declare no conflicts of interest.

STAR⊡Methods

Detailed methods are provided in the online version of this paper and include the

following:

● KEY RESOURCES TABLE

● RESOURCE AVAILABILITY

○ Lead contact

○ Materials availability

○ Data and code availability

● EXPERIMENTAL MODEL AND SUBJECT DETAILS

● METHOD DETAILS

○ Cell culture and strain construction

○ DNA gels and immunoblots

○ Fluorescence microscopy

○ Preparation of nuclear lysates

○ Preparation of cryosections

○ Preparation of cryolamellae

○ Cryo-ET data collection and reconstruction

○ Nucleosome template matching, classification, and subtomogram

averaging

○ Biased-reference classification control

○ Estimation of nucleosomes sampled per cryolamella

○ Simulations of nucleosome tomographic slices

●   QUANTIFICATION AND STATISTICAL ANALYSIS

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and

will be fulfilled by the lead contact, Lu Gan (lu@anaphase.org).

Materials availability

All *S. cerevisiae* strains generated in this study are available upon request.

Data and code availability

●   A subtomogram average of a BY4741 canonical nucleosome *ex vivo*, a double-

GFP tagged LGY0016 nucleosome *ex vivo*, and the two BY4741 canonical

nucleosome classes *in situ* have been deposited at EMDB as entry EMD-31086.

All raw cryo-ET data, reconstructed tomograms and BY4741 cryolamellae VPP *in*

situ class averages have been deposited in EMPIAR under entry EMPIAR-

10678.

●   All auxiliary scripts have been deposited at GitHub and are publicly available as

of the date of publication. URLs of resources are listed in the key resources

table.

●   Any additional information required to reanalyze the data reported in this

paper is available from the lead contact upon request.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Yeast Strain and Growth Conditions

All yeast strains listed in the key resources table were streaked on yeast extract

peptone dextrose (YPD) agar plates (2% w/v peptone, 1% w/v yeast extract, 2% w/v

glucose for liquid media; additional 2% w/v agar for agar plates) at 30°C and cultured in

YPD in conical flasks shaking at 200 – 250 RPM at 30°C. All yeast strains are of mating

type a. Modifications or lack thereof to all histone genes were authenticated by PCR

and Sanger sequencing (Bio Basic Asia Pacific Pte Ltd, Singapore) in all yeast strains.

Bacterial Growth Conditions

The plasmids shown in the key resources table are provided in DH5-Alpha *Escherichia*

coli. The bacteria were streaked on LB agar plates with ampicillin (40 g/L LB Broth with

agar (Miller), 100 µg/mL ampicillin) at 37°C and cultured in LB liquid medium with

ampicillin (25 g/L LB Broth (Miller), 100 µg/mL ampicillin) in vent cap tubes shaking at

200 – 250 RPM at 37°C.

METHOD DETAILS

Plasmid extraction and linearization

The pFA6a plasmids (Lee et al., 2013; Longtine et al., 1998) used are shown in the key

resources table. The plasmid pFA6a-GFP(S65T)-His3MX6 was a gift from John Pringle

(Addgene 41598; Addgene, Watertown, MA) and pFA6a-link-yoTagRFP-T-Kan was a

gift from Wendell Lim & Kurt Thorn (Addgene 44906); both were given in the form of

bacterial stabs. Note that pFA6a-link-yoTagRFP-T-Kan is the source of the KanR

marker used to delete HTB2-HTA2. Five mL of bacteria were cultured overnight, then

plasmids were extracted with the QIAprep Spin Miniprep Kit (QIAGEN, Hilden,

Germany) following the manufacturer's instructions.

Extracted plasmids were linearized by double digestion with a reaction containing 1 µg

of plasmid DNA, 5 µL of 10× rCutSmart™ Buffer (New England BioLabs, Ipswich, MA),

10 units each of *SalI* and *EcoRV* restriction enzymes (New England BioLabs) topped up

to 50 µL with nuclease-free water. The reaction mixture was heated at 37°C for 15

minutes for the digestion reaction, then 80°C for 20 minutes to inactivate the enzymes.

Strain construction

The strain details are shown in Table S1. Primers were from IDT (Integrated DNA

Technologies, Inc., Singapore) and listed in Table S2. Q5 PCR Master Mix (New

England BioLabs) was used for PCR reactions. BY4741 (Brachmann et al., 1998)

served as the wild-type strain. Tagging and deletion cassettes were amplified from

linearized pFA6a plasmids (Lee et al., 2013; Longtine et al., 1998) with a PCR reaction

containing 1 ng of template DNA and 0.5 µM of each primer, using a PCR program of

98°C for 30 seconds, 30 cycles of 98°C for 5 seconds, 60°C for 10 seconds and 72°C

for 1.5 minutes, then 72°C for 5 minutes.

Cells were transformed using the lithium acetate / PEG4000 method reported in

(Nishimura and Kanemaki, 2014). Overnight cell culture in YPD was diluted to 0.1 OD_{600}

in 25 mL of YPD and grown to 0.3 OD_{600}. Ten mL of cells were collected, centrifuged at

1,600 $\times$ g at 25°C for 3 minutes, and the supernatant was removed. The cells were then

washed twice with 10 mL sterile water with centrifugation at 1,600 $\times$ g at 25°C for 3

minutes. The cells were resuspended in 1 mL sterile water, transferred to a new 1.5 mL

collection tube, centrifuged at 17,900 $\times$ g at 25°C for 1 minute, washed in 1 mL of

TE/LiAc (10 mM tris, 1 mM ethylenediaminetetraacetic acid (EDTA), 100 mM lithium

acetate), with centrifugation at 17,900 $\times$ g for 1 minute, and resuspended in 50 µL of the

same buffer. Fifty µL of cell suspension was transferred to a new 1.5 mL collection tube

containing 5 µL of 10 mg/mL salmon sperm DNA (Sigma-Aldrich, Burlington, MA) plus 5

µL of PCR-amplified cassette DNA. 360 µL of TE/LiAc/PEG (10 mM tris, 1 mM EDTA,

100 mM lithium acetate, 40% w/v Polyethylene glycol 4000) was added and incubated

with shaking at 25°C for 30 minutes. Forty µL of dimethyl sulfoxide was added, the

suspension was incubated at 42°C for 15 minutes in a water bath, then cooled on ice for

2 minutes. The suspension was centrifuged at 13,000 x g for 1 minute, the liquid was

removed, and the cells were resuspended in 300 µL of 1$\times$ TE buffer, pH 8.0 (10 mM tris,

1 mM EDTA). Two hundred µL of this cell suspension was plated on a selection plate

and incubated for several days at 30°C. Histidine selection plates were created with 6.7

g/L yeast nitrogen base without amino acids (Sigma-Aldrich), 1.92 g/L yeast synthetic

drop-out medium supplements without histidine (Sigma-Aldrich), 2% w/v glucose and

2% w/v agar. G418 selection plates were created by adding G418 to the molten agar

(YPD for G418 single selection, histidine auxotrophy medium for double selection) to a

concentration of 200 mg/L before it was poured.

Transformants were validated by PCR and Sanger sequencing. Genomic DNA was

extracted with the DNeasy Blood & Tissue Kit (QIAGEN) following the manufacturer's

instructions. Confirmation PCR was performed with 1 µg of template genomic DNA and

0.5 µM of each primer, using a PCR program of 94°C for 2 minutes, 30 cycles of 94°C

for 1 minute, 62°C for 1 minute and 72°C for 3.5 minutes, then 72°C for 5 minutes. PCR

products were purified with the QIAquick PCR Purification Kit (QIAGEN) following the

manufacturer's instructions, with water used for the final elution before they were sent

for Sanger sequencing.

DNA gels and immunoblots

PCR products were electrophoresed in 2% agarose in Tris-acetate-EDTA and

visualized with FloroSafe DNA Stain (Axil Scientific Pte Ltd, Singapore). The

electrophoresis was performed at 100 Volts for 60 – 80 minutes before visualization with

a G:Box (Syngene).

To generate protein samples for immunoblot analysis, ~20 OD_{600} units of yeast cells

were pelleted and stored at –80°C for at least 1 hour. Cells were then resuspended in

200 µL of ice-cold 20% trichloroacetic acid (TCA) and vortexed with glass beads at 4°C

for 1 minute 4 times. For each sample, 500 µL of ice-cold 5% TCA was added, mixed

with the pellet, then transferred to a new 1.5 mL collection tube. Another 500 µL of ice-

cold 5% TCA was mixed with each pellet and transferred to the same 1.5 mL collection

tube as before, so that each collection tube had 1 mL total volume. The tubes were then

left on ice for 10 minutes and centrifuged at 15,000 $\times g$ at 4°C for 20 minutes. The TCA

was aspirated from the tube, then the pellet was resuspended in 212 µL of Laemmli

sample buffer with 2-mercaptoethanol added (Bio-Rad, Hercules, CA). To neutralize the

residual TCA, 26 µL of 1 M Tris, pH 8 was added. This mixture was heated at 95°C for 5

minutes, centrifuged at 25°C at 15,000 $\times$ g for 10 minutes, then 5 µL of the supernatant

was subjected to SDS-PAGE.

The primary and secondary antibodies used for immunoblots are shown in Table S3.

10% Mini-PROTEAN® TGX™ Precast Protein Gels (Bio-Rad) were used, with either

Precision Plus Protein™ WesternC™ Standards (Bio-Rad) or Invitrogen™ MagicMark™

XP Western Protein Standard (Thermo Fisher Scientific, TFS, Waltham, MA) used as

the ladder. The proteins were electrophoresed at 100 Volts for 1 hour at 25°C. The gels

were transferred onto Immun-Blot® PVDF Membranes (Bio-Rad) at 4°C in transfer

buffer (3.02 g/L tris, 14.4 g/L glycine, 20% methanol). The transfer was performed at

100 Volts for 30 minutes. The membranes were then blocked with 2% BSA in 1× Tris-

Buffered Saline, 0.1% Tween® 20 Detergent (TBST) for 1 hour. This was followed by 50

µg/mL avidin in TBST for 30 minutes, then a wash with TBST for 20 minutes if Precision

Plus Protein™ WesternC™ Standards ladder was used. All antibody dilution factors are

reported in Table S3. The membranes were probed with primary antibodies at the

stated dilutions in 2% BSA in TBST for 1 hour at 25°C. Membranes were washed with

TBST for 20 minutes three times at 25°C. The membranes were then probed with

secondary antibodies in 2% BSA in TBST for 1 hour, with additional Precision Protein™

StrepTactin-HRP Conjugate (Bio-Rad) at 1:10,000 dilution if Precision Plus Protein™

WesternC™ Standards ladder was used. Finally, the membranes were washed with

TBST for 10 minutes 3 times, then treated with a 50:50 mixture of Clarity Western

Peroxide Reagent and Clarity Western Luminol/Enhancer Reagent (Bio-Rad) for 5

minutes before visualization by chemiluminescence on an ImageQuant LAS 4000

(Cytiva, Marlborough, MA).

Fluorescence microscopy

Cells were grown to log phase ($OD_{600} = 0.1 - 1.0$), of which 2 OD_{600} units of cells were

collected, pelleted at 5,000 $\times$ g for 1 minute, then resuspended in 1 mL of YPD. Four µL

of cell culture was then applied to a glass slide and pressed against a coverslip. The

cells were imaged live at 23°C with an Olympus FV3000 Confocal Laser Scanning

Microscope (Olympus, Tokyo, Japan) equipped with a 1.35 NA 60× oil-immersion

objective lens. GFP fluorescence was acquired using the 488 nm laser-line, with a DIC

image recorded in parallel with the fluorescence image. Images were captured as Z-

stacks thick enough to sample the GFP signals through all the nuclei in each stage

position. Additional details regarding data collection are shown in Table S4.

Preparation of nuclear lysates

Yeast nuclei were prepared with reagents from the Yeast nuclei isolation kit (Abcam

206997, Cambridge, United Kingdom), unless noted otherwise. Yeast cells (30 mL,

OD_{600} ~1) were pelleted at 3,000 $\times$ g for 5 minutes at 25°C. The pellet was washed

twice with 1 mL water (3,000 $\times$ g, 1 minute). The pellet was then resuspended in 1 mL

Buffer A (pre-warmed to 30°C) containing 10 mM dithiothreitol. The suspension was

incubated in a 30°C water bath for 10 minutes. Cells were then pelleted at 1,500 $\times$ g for

5 minutes and then resuspended in 1 mL Buffer B (prewarmed to 30°C) containing lysis

enzyme cocktail (1:100 dilution). The suspension was incubated in a 30°C shaker for 15

minutes for cell-wall digestion and then pelleted at 1,500 $\times g$ for 5 minutes at 4°C. The

pellet was resuspended in 1 mL pre-chilled Buffer C with protease inhibitor cocktail

(1:1,000 dilution). The cells were lysed by 15 up-and-down strokes with a pre-chilled

glass Dounce homogenizer on ice. The lysate was incubated with shaking for 30

minutes at 25°C. The cell debris was pelleted at 500 $\times g$ for 5 minutes at 4°C. The

supernatant was then transferred to a new tube. The nuclei were pelleted at 20,000 $\times g$

for 10 minutes at 4°C. The nuclear pellet was resuspended in 10 – 20 µL pre-chilled

lysis buffer (50 mM EDTA and 1:1,000 protease inhibitor cocktail dilution) and incubated

for 15 minutes on ice.

Nuclei lysates (3 µL) were added to a glow-discharged CF-4/2-2C-T grid (Protochips,

Morrisville, NC). The grid was plunge-frozen using a Vitrobot Mark IV (blot time: 1

second, blot force: 1, humidity: 100%, temperature: 4°C).

Preparation of cryosections

Self-pressurized freezing was done based on a modified version of a published protocol

(Yakovlev and Downing, 2011). Yeast cells (30 mL, OD_{600} = 0.2 – 0.6) were pelleted

and resuspended in a dextran stock (40 kilodalton, 60% w/v, in YPD) to a final

concentration of 30%. Cells were then loaded into a copper tube (0.45 / 0.3 mm outer /

inner diameters). Both ends of the tube were sealed with flat-jaw pliers. The tube was

held horizontally and dropped into the liquid-ethane cryogen. The tube's ends were

removed under liquid nitrogen with a tube-cut tool (Engineering Office M. Wohlwend,

Sennwald, Switzerland).

Gold colloid solution (10-nm diameter, 5 µL at 5.7×10^{12} particles/mL in 0.1 mg/mL BSA)

was applied to a continuous-carbon grid (10-nm thick carbon) and then air-dried

overnight. Cryosections were controlled by a custom joystick-based micromanipulator

(MN-151S, Narishige Co., Ltd., Tokyo, Japan) (Ladinsky et al., 2006; Ng et al., 2020).

Seventy nm-thick frozen-hydrated sections were cut at −150°C in a Leica UC7/FC7

cryo-ultramicrotome (Leica Microsystems, Vienna, Austria). The EM grid was positioned

underneath the ribbon using a Leica micromanipulator (Studer et al., 2014). The

cryosection ribbon (~3 mm long) was then attached to the grid by operating the Crion

(Leica Microsystems) in "charge" mode for ~30 seconds (Pierson et al., 2010).

Preparation of cryolamellae

Cells were plunge-frozen and then cryo-FIB milled using the method of Medeiros and

Bock (Medeiros et al., 2018) as follows. Immediately before plunge-freezing, mid-log

phase (OD_{600} ~0.6) yeast cells were pelleted at 4,000 $\times$ g for 5 minutes. They were then

resuspended in YPD media containing 3% (v/v) dimethyl sulfoxide as cryo-protectant to

a final OD_{600} of approximately 2.5. Four µL of the cells were subsequently deposited

onto Quantifoil R2/4 200 mesh copper grids (Quantifoil Micro Tools GmbH, Jena,

Germany), which were then manually blotted from the back with Whatman® Grade 1

filter paper for approximately 3 – 5 seconds. The grids were then plunged into a 63/37

propane/ethane mixture (Tivol et al., 2008) using a Vitrobot Mark IV (humidity: 100%,

temperature: 4°C). Cryo-FIB milling was performed on a Helios NanoLab 600

DualBeam (Thermo Fisher Scientific, TFS, Waltham, MA) equipped with a Quorum

PolarPrep 2000 transfer system (Quorum Technologies, Laughton, United Kingdom).

Plunge-frozen yeast samples were coated with a layer of organometallic platinum using

the in-chamber gas injection system and the cold-deposition method (Hayles et al.,

2007). Cryolamellae were then generated as follows: bulk material was first removed

using the FIB at 30 kV 2.8 nA, followed by successive thinning of the cryolamellae at

lower currents of 0.28 nA and 48 pA.

Cryo-ET data collection and reconstruction

All cryo-ET data were collected on Titan Krioses (TFS). Tilt series were collected with

either TFS Tomo4, Leginon (Suloway et al., 2009), SerialEM (Mastronarde, 2003), or

PACE-tomo (Eisenstein et al., 2022). Images were recorded either on a Falcon II (TFS)

in integration mode or as movie frames on a K2 or K3 summit camera (Gatan,

Pleasanton, CA) in super-resolution mode. The pixel sizes for the K2 and K3 data were

chosen so that when binned to the same level, they closely match the ~7 Å pixel size in

our previous *in situ* study of HeLa chromatin (Cai et al., 2018a). Smaller pixel sizes

were chosen for Falcon II data because this camera has lower detective quantum

efficiency than the K-series cameras (Ruskin et al., 2013). Movies were aligned with

either MotionCor2 (Zheng et al., 2017) or IMOD alignframes (Mastronarde, 1997). Prior

to starting data collection of cryolamellae, the stage was pre-tilted to either −10° or −15°

to account for the milling angle. Additional data-collection details are shown in Table S5.

Volta Phase Plate (VPP) imaging was done using the protocol of Fukuda and

colleagues {Fukuda, 2015}. The VPP position and condenser stigmator were adjusted

such that the Ronchigram (the projected pattern visible in the camera) was free of

elongated features, indicating no astigmatism, and is flat rather than grainy, indicating

that the VPP is on-plane. Notably, the VPP needs to be heated, or else the phase shift

will rapidly exceed $\pi/2$ radians {Danev, 2014}. Two different VPP assemblies were

used, which we will refer to as the NYSBC-2019 and NUS-2020 ones. These VPPs

required different heater power settings to maintain a stable phase shift. The NYSBC-

2019 and NUS-2020 VPP heaters were operated at ~100 mW and ~370 mW,

respectively.

Cryotomograms were reconstructed using IMOD's *eTomo* workflow (Mastronarde,

1997). Tilt series of lysates and cryosections were aligned using the gold beads as

fiducials while those of cryolamellae were aligned with patches as fiducials. Only the tilt

series that exhibited the minimal amount of drift and sample warping were analyzed. To

further improve the tilt series alignment around the chromatin, fiducials (beads or

patches) were chosen on the chromatin regions. For cryolamella patch tracking, tilt

series were binned to a pixel size of 6.8 Å, the "Break contours into pieces w/ overlap"

was set to 10, and the low-frequency rolloff sigma and cutoff radius were set to 0.03 and

0.1 $pixel^{-1}$, respectively. A boundary model was created so that only the chromatin was

enclosed. Fiducial patches were manually deleted if they overlapped with debris like ice

crystals or if they mistracked. We did not detect a correlation between the alignment

residual (Table S6) and our ability to detect canonical nucleosomes.

CTF estimation and phase flipping were done on the defocus phase contrast datasets

using IMOD's *ctfplotter* and *ctfphaseflip* programs. Prior to reconstruction, the

cryosection and cryolamellae tilt series were binned to a final pixel size of 6.8 Å using

the *eTomo* antialiasing option. Two tomogram versions were reconstructed for each tilt

series. For visualization purposes, the tilt series were low-pass filtered to attenuate

spatial frequencies beyond 25 Å to 30 Å resolution, prior to tomogram reconstruction.

For classification analysis, the tilt series were low-pass filtered with a Gaussian rolloff

starting at 15 Å resolution for lysates and 20 Å for cryosections and cryolamellae. In the

classification jobs, the resolution of the data was further limited to 25 Å or 20 Å (see

next section for details). More details of the datasets analyzed in this paper are shown

in Table S6.

Nucleosome template matching, classification, and subtomogram averaging

A featureless round-edged 10 nm-diameter, 6 nm-thick cylindrical template was created

using the Bsoft program *beditimg* (Heymann and Belnap, 2007). A cubic search grid

with a 12-nm spacing was created with the PEET program *gridInit* (Heumann, 2016).

Regions that had high-contrast artifacts from surface contaminants and ice crystals

were excluded. Template matching was done using PEET (Heumann, 2016; Heumann

et al., 2011; Nicastro et al., 2006), with a duplicate-removal cutoff distance of 6 nm. To

accelerate the runs, no orientation search was done around the cylindrical axis and the

resolution was attenuated starting at 70 Å on account of the smooth appearance of the

template. Candidate hit lists of different cross-correlation cutoffs were generated using

the PEET program *createAlignedModel*, then visualized together with the tomograms in

3dmod. The cross-correlation cutoff that eliminated spurious densities (primarily empty

nucleoplasm) was chosen. The final numbers of subtomograms analyzed for each

sample are in Table S7 and in the figures and figure legends.

Classification and subtomogram analysis were done with RELION (Kimanius et al.,

2016; Scheres, 2012), following the workflows in Figure S1. In the published workflow

(Figure S1A) (Bharat and Scheres, 2016), each subtomogram is first averaged by

projecting the entire volume (16 nm along the Z axis), which introduces contributions

from densities above and below the candidate nucleosome. To minimize the influence

of other densities, pseudo-projections were created by averaging ~12 nm along the Z

axis, using the *ot_relion_project.py* script. 2-D classification using mask diameters

ranging from 120 Å to 140 Å produced clear nucleosome-like classes, though the

smaller masks included fewer adjacent densities. Note that in RELION, only circular

masks are available for 2-D classification, meaning that it is not possible for the pseudo-

projected densities to appear cylinder-like due to truncation by the mask. Densities that

belonged to the most nucleosome-like classes were exported for 3-D classification, split

into 30 classes. The resolution cutoffs were 25 Å for 2-D and 20 Å for 3-D classification.

To eliminate the influence of adjacent densities during 3-D classification, a smooth

cylindrical mask with a cosine-shaped edge was applied. The mask was created using

beditimg and *relion_mask_create*. Because the GFP densities protrude from the

nucleosome surface, we used a 9 nm-tall cylindrical mask for the analysis of

nucleosomes with GFP tags, such as LGY0016 nucleosomes. The percentage of

subtomograms belonging to each class was extracted with the script

count_particles.awk from Guillaume Gaullier (Gaullier, 2021).

As observed in our previous study (Cai et al., 2018a), some nucleosomes were lost in

the 2-D classification process. We therefore used the alternative workflow in which the

template matching hits were directly classified in 3-D (Figure S1B). To accommodate

the increased diversity of complexes (some of which would have been removed had 2-D

classification been done), we used 40 classes for BY4741, LGY0015, and LGY0016

lysates and 100 classes for cryolamellae and lysates of strains in which H3 or H4 were

GFP tagged, without prior 2-D classification. These jobs crashed frequently because the

RELION memory usage scales up with the number of classes (Kimanius et al., 2016).

We were able to eliminate the crash problem by using higher-memory GPUs and by

decreasing the number of translational search steps from 5 to 3 or 4. The canonical

nucleosome classes were subjected to "Gold-standard" 3-D refinement (Henderson et

al., 2012). No map sharpening was applied. Subtomogram class-average volumes were

visualized with UCSF Chimera (Pettersen et al., 2004).

Biased-reference classification control

A 14 Å-resolution density map was simulated from the yeast nucleosome crystal

structure PDB 1ID3 (White et al., 2001) using the Bsoft program *bgex*. To account for

the artifacts associated with defocus phase contrast and CTF correction, a 6 μm

underfocus was applied and then "corrected" for using the Bsoft program *bctf*. The map

was also subjected to the 20 Å-resolution low-pass filter that was used on the tilt series

using the IMOD program *mtffilter*. Template matching was done using this reference,

including data to higher resolution (28 Å) than that for the less-biased search above and

including a search around all Euler angles. The hits were subjected to 2-D classification

to remove obvious non-nucleosomal densities. Next, the hits were 3-D classified using

the simulated map as an initial alignment reference. To maximize the model bias, the

template was only low-pass filtered to 20 Å-resolution instead of the recommended 60 Å

(Bharat and Scheres, 2016).

Estimation of nucleosomes sampled per cryolamella

First, the average concentration of nucleosomes in the chromatin was estimated. The

absolute number of nucleosomes per cell determined from genomics is 60,000

(Oberbeckmann et al., 2019). Soft X-ray tomography measurements revealed that the

average G1 nucleus volume is 2 μm^3, of which 20% is nucleolus (Uchida et al., 2011).

Accordingly, chromatin (the nuclear volume not taken by the nucleolus), which contains

the vast majority of nucleosomes, occupies ~1.6 μm^3. These two experimental values

give an average nucleosome concentration of 37,500 per μm^3. Next, the nuclear volume

sampled by subtomogram analysis was determined by first drawing one closed contour

around the chromatin using 3dmod; this closed contour encloses only the portion of the

tomogram that was analyzed by template matching. The volume of this closed contour,

which is one-voxel thick, was extracted using the command:

imodinfo -F model.mod

This command outputs the quantity "Cylinder volume", in cubic pixels (voxels). The total

tomographic volume sampled (Table S8) was obtained by multiplying Cylinder volume,

the voxel volume (0.31 nm^3 for 0.68 nm pixel size), and the number of tomographic

slices that contain chromatin. The BY4741 VPP tomograms summed to 0.67 μm^3, which

yields ~25,000 nucleosomes. For the HeLa cell in (Cai et al., 2018a) (EMPIAR-10179),

the nucleus volume analyzed was 0.12 μm^3.

Simulations of nucleosome tomographic slices

Atomic models of the nucleosome (PDB 1KX5) (Davey et al., 2002) were manually

positioned in UCSF Chimera and edited to remove the N-terminal tails. A 3-D density

map was calculated with the Bsoft program *bgex*. There is no software that simulates

Volta contrast, so we approximated the Volta-induced phase shift by setting the

amplitude contrast to 100% and the defocus to zero in the Bsoft program *bctf*, followed

by "correction", also done with *bctf*. A tilt series was calculated from the simulated map

using the IMOD program *xyzproj*. The tilt series was then aligned by cross-correlation

and then backprojected using IMOD's *eTomo* workflow. Parameters such as pixel size,

tilt range, and tilt angle were kept as close to the experimental ones as possible.

Tomographic slices were made in IMOD slicer at the same thickness as for the real

cryotomograms.

Ribosome subtomogram analysis control

Ribosomes were analyzed using the same software packages as for nucleosomes, as

detailed in the workflow in Figure S25. Candidate ribosomes were template matched in

a single VPP tomogram (from Figure S11), using a 25-nm-diameter sphere as a

reference. The candidate ribosomes were filtered by cross correlation so that obvious

false positives (vacuum) were excluded, leaving 3,816 hits. These particles were

subjected to direct 3-D classification with k = 10, resulting in six non-empty classes.

Particles belonging to the two ribosome classes (1,150 total) were pooled and "gold-

standard" refined, yielding a density map at 28 Å resolution (FSC = 0.143) / 33 Å

resolution (FSC = 0.5). Density maps were simulated at 15 Å, 20 Å, and 30 Å resolution

using the Bsoft program *bgex* and the yeast ribosome crystal structure (Ben-Shem et

al., 2011). To approximate the use of the Volta phase plate, the amplitude contrast was

set to 90% and the defocus to –1 µm.

REFERENCES

Armeev, G.A., A.S. Kniazeva, G.A. Komarova, M.P. Kirpichnikov, and A.K. Shaytan. 2021. Histone dynamics mediate DNA unwrapping and sliding in nucleosomes. *Nat Commun*. 12:2387.

Beck, M., and W. Baumeister. 2016. Cryo-Electron Tomography: Can it Reveal the Molecular Sociology of Cells in Atomic Detail? *Trends Cell Biol*. 26:825-837.

Bednar, J., I. Garcia-Saez, R. Boopathi, A.R. Cutter, G. Papai, A. Reymer, S.H. Syed, I.N. Lone, O. Tonchev, C. Crucifix, H. Menoni, C. Papin, D.A. Skoufias, H. Kurumizaka, R. Lavery, A. Hamiche, J.J. Hayes, P. Schultz, D. Angelov, C. Petosa, and S. Dimitrov. 2017. Structure and Dynamics of a 197 bp Nucleosome in Complex with Linker Histone H1. *Mol Cell*. 66:384-397 e388.

Ben-Shem, A., N. Garreau de Loubresse, S. Melnikov, L. Jenner, G. Yusupova, and M. Yusupov. 2011. The structure of the eukaryotic ribosome at 3.0 A resolution. *Science*. 334:1524-1529.

Bepler, T., K. Kelley, A.J. Noble, and B. Berger. 2020. Topaz-Denoise: general deep denoising models for cryoEM and cryoET. *Nat Commun*. 11:5208.

Bharat, T.A., and S.H. Scheres. 2016. Resolving macromolecular structures from electron cryo-tomography data using subtomogram averaging in RELION. *Nat Protoc*. 11:2054-2065.

Bi, E., and J.R. Pringle. 1996. ZDS1 and ZDS2, genes whose products may regulate Cdc42p in Saccharomyces cerevisiae. *Mol Cell Biol*. 16:5264-5275.

Bilokapic, S., and M. Halic. 2019. Nucleosome and ubiquitin position Set2 to methylate H3K36. *Nat Commun*. 10:3795.

Bilokapic, S., M. Strauss, and M. Halic. 2018. Histone octamer rearranges to adapt to DNA unwrapping. *Nat Struct Mol Biol*. 25:101-108.

Brachmann, C.B., A. Davies, G.J. Cost, E. Caputo, J. Li, P. Hieter, and J.D. Boeke. 1998. Designer deletion strains derived from Saccharomyces cerevisiae S288C: a useful set of strains and plasmids for PCR-mediated gene disruption and other applications. *Yeast*. 14:115-132.

Brandani, G.B., C. Tan, and S. Takada. 2021. The kinetic landscape of nucleosome assembly: A coarse-grained molecular dynamics study. *PLoS Comput Biol*. 17:e1009253.

Buckley, G., G. Gervinskas, C. Taveneau, H. Venugopal, J.C. Whisstock, and A. de Marco. 2020. Automated cryo-lamella preparation for high-throughput in-situ structural biology. *J Struct Biol*. 210:107488.

Cai, S., D. Böck, M. Pilhofer, and L. Gan. 2018a. The in situ structures of mono-, di-, and trinucleosomes in human heterochromatin. *Mol Biol Cell*. 29:2450-2457.

Cai, S., C. Chen, Z.Y. Tan, Y. Huang, J. Shi, and L. Gan. 2018b. Cryo-ET reveals the
macromolecular reorganization of S. pombe mitotic chromosomes in vivo. *Proc Natl*
Acad Sci U S A. 115:10977-10982.

Cai, S., Y. Song, C. Chen, J. Shi, and L. Gan. 2018c. Natural chromatin is heterogeneous and
self-associates in vitro. *Mol Biol Cell*. 29:1652-1663.

Chang, Y.W., L.A. Rettberg, A. Treuner-Lange, J. Iwasa, L. Sogaard-Andersen, and G.J.
Jensen. 2016. Architecture of the type IVa pilus machine. *Science*. 351:aad2001.

Chen, C., H.H. Lim, J. Shi, S. Tamura, K. Maeshima, U. Surana, and L. Gan. 2016. Budding
yeast chromatin is dispersed in a crowded nucleoplasm in vivo. *Mol Biol Cell*. 27:3357-
3368.

Chen, M., J.M. Bell, X. Shi, S.Y. Sun, Z. Wang, and S.J. Ludtke. 2019. A complete data
processing workflow for cryo-ET and subtomogram averaging. *Nat Methods*. 16:1161-
1168.

Chua, E.Y., V.K. Vogirala, O. Inian, A.S. Wong, L. Nordenskiold, J.M. Plitzko, R. Danev, and S.
Sandin. 2016. 3.9 A structure of the nucleosome core particle determined by phase-plate
cryo-EM. *Nucleic Acids Res*. 44:8013-8019.

Danev, R., D. Tegunov, and W. Baumeister. 2017. Using the Volta phase plate with defocus for
cryo-EM single particle analysis. *Elife*. 6.

Davey, C.A., D.F. Sargent, K. Luger, A.W. Maeder, and T.J. Richmond. 2002. Solvent mediated
interactions in the structure of the nucleosome core particle at 1.9 a resolution. *J Mol*
Biol. 319:1097-1113.

Diestra, E., J. Fontana, P. Guichard, S. Marco, and C. Risco. 2009. Visualization of proteins in
intact cells with a clonable tag for electron microscopy. *J Struct Biol*. 165:157-168.

Dodonova, S.O., F. Zhu, C. Dienemann, J. Taipale, and P. Cramer. 2020. Nucleosome-bound
SOX2 and SOX11 structures elucidate pioneer factor function. *Nature*. 580:669-672.

Eisenstein, F., H. Yanagisawa, H. Kashihara, M. Kikkawa, S. Tsukita, and R. Danev. 2022.
Parallel cryo electron tomography on in situ lamellae. *bioRxiv*.

Ekundayo, B., T.J. Richmond, and T. Schalch. 2017. Capturing Structural Heterogeneity in
Chromatin Fibers. *J Mol Biol*. 429:3031-3042.

Eltsov, M., D. Grewe, N. Lemercier, A. Frangakis, F. Livolant, and A. Leforestier. 2018.
Nucleosome conformational variability in solution and in interphase nuclei evidenced by
cryo-electron microscopy of vitreous sections. *Nucleic Acids Res*. 46:9189-9200.

Eltsov, M., K.M. MacLellan, K. Maeshima, A.S. Frangakis, and J. Dubochet. 2008. Analysis of
cryo-electron microscopy images does not support the existence of 30-nm chromatin
fibers in mitotic chromosomes in situ. *Proceedings of the National Academy of Sciences*.
105:19732-19737.

Eustermann, S., K. Schall, D. Kostrewa, K. Lakomek, M. Strauss, M. Moldt, and K.P. Hopfner.
2018. Structural basis for ATP-dependent chromatin remodelling by the INO80 complex.
Nature. 556:386-390.

Farnung, L., S.M. Vos, C. Wigge, and P. Cramer. 2017. Nucleosome-Chd1 structure and
implications for chromatin remodelling. *Nature*. 550:539-542.

Farr, S.E., E.J. Woods, J.A. Joseph, A. Garaizar, and R. Collepardo-Guevara. 2021.
Nucleosome plasticity is a critical element of chromatin liquid-liquid phase separation
and multivalent nucleosome interactions. *Nat Commun*. 12:2883.

Fatmaoui, F., P. Carrivain, D. Grewe, B. Jakob, J.-M. Victor, A. Leforestier, and M. Eltsov. 2022.
Cryo-electron tomography and deep learning denoising reveal native chromatin
landscapes of interphase nuclei. *bioRxiv*.

Gan, L., M.S. Ladinsky, and G.J. Jensen. 2013. Chromatin in a marine picoeukaryote is a
disordered assemblage of nucleosomes. *Chromosoma*. 122:377-386.

Gaullier, G. 2021. cryoEM-scripts. Vol. 2021.

Hayles, M.F., D.J. Stokes, D. Phifer, and K.C. Findlay. 2007. A technique for improved focused
ion beam milling of cryo-prepared life science specimens. *J Microsc*. 226:263-269.

Henderson, R., A. Sali, M.L. Baker, B. Carragher, B. Devkota, K.H. Downing, E.H. Egelman, Z.
Feng, J. Frank, N. Grigorieff, W. Jiang, S.J. Ludtke, O. Medalia, P.A. Penczek, P.B.
Rosenthal, M.G. Rossmann, M.F. Schmid, G.F. Schroder, A.C. Steven, D.L. Stokes, J.D.
Westbrook, W. Wriggers, H. Yang, J. Young, H.M. Berman, W. Chiu, G.J. Kleywegt, and
C.L. Lawson. 2012. Outcome of the first electron microscopy validation task force
meeting. *Structure*. 20:205-214.

Hereford, L., K. Fahrner, J. Woolford, Jr., M. Rosbash, and D.B. Kaback. 1979. Isolation of
yeast histone genes H2A and H2B. *Cell*. 18:1261-1271.

Heumann, J.M. 2016. PEET. Vol. 2021. University of Colorado Boulder.

Heumann, J.M., A. Hoenger, and D.N. Mastronarde. 2011. Clustering and variance maps for
cryo-electron tomography using wedge-masked differences. *J Struct Biol*. 175:288-299.

Heymann, J.B., and D.M. Belnap. 2007. Bsoft: image processing and molecular modeling for
electron microscopy. *J Struct Biol*. 157:3-18.

Himes, B.A., and P. Zhang. 2018. emClarity: software for high-resolution cryo-electron
tomography and subtomogram averaging. *Nat Methods*. 15:955-961.

Huertas, J., H.R. Scholer, and V. Cojocaru. 2021. Histone tails cooperate to control the
breathing of genomic nucleosomes. *PLoS Comput Biol*. 17:e1009013.

Ishida, H., and H. Kono. 2021. Torsional stress can regulate the unwrapping of two outer half
superhelical turns of nucleosomal DNA. *Proc Natl Acad Sci U S A*. 118.

Ishida, H., and H. Kono. 2022. Free Energy Landscape of H2A-H2B Displacement From
Nucleosome. *J Mol Biol*. 434:167707.

Iudin, A., P.K. Korir, J. Salavert-Torres, G.J. Kleywegt, and A. Patwardhan. 2016. EMPIAR: a
public archive for raw electron microscopy image data. *Nat Methods*. 13:387-388.

Jang, S., C. Kang, H.S. Yang, T. Jung, H. Hebert, K.Y. Chung, S.J. Kim, S. Hohng, and J.J.
Song. 2019. Structural basis of recognition and destabilization of the histone H2B
ubiquitinated nucleosome by the DOT1L histone H3 Lys79 methyltransferase. *Genes*
Dev. 33:620-625.

Khoshouei, M., S. Pfeffer, W. Baumeister, F. Forster, and R. Danev. 2017. Subtomogram
analysis using the Volta phase plate. *J Struct Biol*. 197:94-101.

Kimanius, D., B.O. Forsberg, S.H. Scheres, and E. Lindahl. 2016. Accelerated cryo-EM
structure determination with parallelisation using GPUs in RELION-2. *Elife*. 5.

Konrad, S.F., W. Vanderlinden, W. Frederickx, T. Brouns, B.H. Menze, S. De Feyter, and J.
Lipfert. 2021. High-throughput AFM analysis reveals unwrapping pathways of H3 and
CENP-A nucleosomes. *Nanoscale*. 13:5435-5447.

Koyama, M., W. Nagakura, H. Tanaka, T. Kujirai, Y. Chikashige, T. Haraguchi, Y. Hiraoka, and
H. Kurumizaka. 2017. In vitro reconstitution and biochemical analyses of the
Schizosaccharomyces pombe nucleosome. *Biochem Biophys Res Commun*. 482:896-
901.

Kujirai, T., H. Ehara, Y. Fujino, M. Shirouzu, S.I. Sekine, and H. Kurumizaka. 2018. Structural
basis of the nucleosome transition during RNA polymerase II passage. *Science*.
362:595-598.

Ladinsky, M.S., J.M. Pierson, and J.R. McIntosh. 2006. Vitreous cryo-sectioning of cells
facilitated by a micromanipulator. *J Microsc*. 224:129-134.

Laughlin, T.G., A. Deep, A.M. Prichard, C. Seitz, Y. Gu, E. Enustun, S. Suslov, K. Khanna, E.A.
Birkholz, E. Armbruster, J.A. McCammon, R.E. Amaro, J. Pogliano, K.D. Corbett, and E.
Villa. 2022. Architecture and self-assembly of the jumbo bacteriophage nuclear shell.
Nature.

Lee, S., W.A. Lim, and K.S. Thorn. 2013. Improved blue, green, and red fluorescent protein
tagging vectors for S. cerevisiae. *PLoS One*. 8:e67902.

Libuda, D.E., and F. Winston. 2006. Amplification of histone genes by circular chromosome
formation in Saccharomyces cerevisiae. *Nature*. 443:1003-1007.

Liu, X., M. Li, X. Xia, X. Li, and Z. Chen. 2017. Mechanism of chromatin remodelling revealed by
the Snf2-nucleosome structure. *Nature*. 544:440-445.

Longtine, M.S., A. McKenzie, 3rd, D.J. Demarini, N.G. Shah, A. Wach, A. Brachat, P.
Philippsen, and J.R. Pringle. 1998. Additional modules for versatile and economical
PCR-based gene deletion and modification in Saccharomyces cerevisiae. *Yeast*.
14:953-961.

Luger, K., A.W. Mader, R.K. Richmond, D.F. Sargent, and T.J. Richmond. 1997. Crystal
structure of the nucleosome core particle at 2.8 A resolution. *Nature*. 389:251-260.

Mahamid, J., R. Schampers, H. Persoon, A.A. Hyman, W. Baumeister, and J.M. Plitzko. 2015. A
focused ion beam milling and lift-out approach for site-specific preparation of frozen-
hydrated lamellas from multicellular organisms. *J Struct Biol*. 192:262-269.

Marko, M., C. Hsieh, W. Moberlychan, C.A. Mannella, and J. Frank. 2006. Focused ion beam
milling of vitreous water: prospects for an alternative to cryo-ultramicrotomy of frozen-
hydrated biological samples. *J Microsc*. 222:42-47.

Marr, L.T., J. Ocampo, D.J. Clark, and J.J. Hayes. 2021. Global histone protein surface
accessibility in yeast indicates a uniformly loosely packed genome with canonical
nucleosomes. *Epigenetics Chromatin*. 14:5.

Mastronarde, D.N. 1997. Dual-axis tomography: an approach with alignment methods that
preserve resolution. *J Struct Biol*. 120:343-352.

Mastronarde, D.N. 2003. SerialEM: A program for automated tilt series acquisition on Tecnai
microscopes using prediction of specimen position. *Microscopy and Microanalysis*.
9:1182CD.

McDowall, A.W., J.M. Smith, and J. Dubochet. 1986. Cryo-electron microscopy of vitrified
chromosomes in situ. *EMBO J*. 5:1395-1402.

Medeiros, J.M., D. Bock, G.L. Weiss, R. Kooger, R.A. Wepf, and M. Pilhofer. 2018. Robust
workflow and instrumentation for cryo-focused ion beam milling of samples for electron
cryotomography. *Ultramicroscopy*. 190:1-11.

Mercogliano, C.P., and D.J. DeRosier. 2007. Concatenated metallothionein as a clonable gold
label for electron microscopy. *J Struct Biol*. 160:70-82.

Merino, F., S. Pospich, J. Funk, T. Wagner, F. Kullmer, H.D. Arndt, P. Bieling, and S. Raunser.
2018. Structural transitions of F-actin upon ATP hydrolysis at near-atomic resolution
revealed by cryo-EM. *Nat Struct Mol Biol*. 25:528-537.

Moebel, E., A. Martinez-Sanchez, L. Lamm, R.D. Righetto, W. Wietrzynski, S. Albert, D.
Lariviere, E. Fourmentin, S. Pfeffer, J. Ortiz, W. Baumeister, T. Peng, B.D. Engel, and C.
Kervrann. 2021. Deep learning improves macromolecule identification in 3D cellular
cryo-electron tomograms. *Nat Methods*. 18:1386-1394.

Mohan, C., L.M. Kim, N. Hollar, T. Li, E. Paulissen, C.T. Leung, and E. Luk. 2018. VivosX, a
disulfide crosslinking method to capture site-specific, protein-protein interactions in yeast
and human cells. *Elife*. 7.

Ng, C.T., and L. Gan. 2020. Investigating eukaryotic cells with cryo-ET. *Mol Biol Cell*. 31:87-
100.

Ng, C.T., M. Ladinsky, and L. Gan. 2020. Serial Cryomicrotomy of Saccharomyces cerevisiae
for Serial Electron Cryotomography. *Bio-Protocol*. 10.

Nicastro, D., C. Schwartz, J. Pierson, R. Gaudette, M.E. Porter, and J.R. McIntosh. 2006. The molecular architecture of axonemes revealed by cryoelectron tomography. *Science*. 313:944-948.

Nishimura, K., and M.T. Kanemaki. 2014. Rapid Depletion of Budding Yeast Proteins via the Fusion of an Auxin-Inducible Degron (AID). *Curr Protoc Cell Biol*. 64:20 29 21-16.

Nishino, Y., T. Yasunaga, and A. Miyazawa. 2007. A genetically encoded metallothionein tag enabling efficient protein detection by electron microscopy. *J Electron Microsc (Tokyo)*. 56:93-101.

Noble, A.J., V.P. Dandey, H. Wei, J. Brasch, J. Chase, P. Acharya, Y.Z. Tan, Z. Zhang, L.Y. Kim, G. Scapin, M. Rapp, E.T. Eng, W.J. Rice, A. Cheng, C.J. Negro, L. Shapiro, P.D. Kwong, D. Jeruzalmi, A. des Georges, C.S. Potter, and B. Carragher. 2018. Routine single particle CryoEM sample and grid characterization by tomography. *Elife*. 7.

Oberbeckmann, E., M. Wolff, N. Krietenstein, M. Heron, J.L. Ellins, A. Schmid, S. Krebs, H. Blum, U. Gerland, and P. Korber. 2019. Absolute nucleosome occupancy map for the Saccharomyces cerevisiae genome. *Genome Res*. 29:1996-2009.

Oda, T., and M. Kikkawa. 2013. Novel structural labeling method using cryo-electron tomography and biotin-streptavidin system. *J Struct Biol*. 183:305-311.

Ohno, M., T. Ando, D.G. Priest, V. Kumar, Y. Yoshida, and Y. Taniguchi. 2019. Sub-nucleosomal Genome Structure Reveals Distinct Nucleosome Folding Motifs. *Cell*. 176:520-534 e525.

Pettersen, E.F., T.D. Goddard, C.C. Huang, G.S. Couch, D.M. Greenblatt, E.C. Meng, and T.E. Ferrin. 2004. UCSF Chimera--a visualization system for exploratory research and analysis. *J Comput Chem*. 25:1605-1612.

Pierson, J., J.J. Fernandez, E. Bos, S. Amini, H. Gnaegi, M. Vos, B. Bel, F. Adolfsen, J.L. Carrascosa, and P.J. Peters. 2010. Improving the technique of vitreous cryo-sectioning for cryo-electron tomography: electrostatic charging for section attachment and implementation of an anti-contamination glove box. *J Struct Biol*. 169:219-225.

Ramachandran, S., K. Ahmad, and S. Henikoff. 2017. Transcription and Remodeling Produce Asymmetrically Unwrapped Nucleosomal Intermediates. *Mol Cell*. 68:1038-1053 e1034.

Ranjan, A., V.Q. Nguyen, S. Liu, J. Wisniewski, J.M. Kim, X. Tang, G. Mizuguchi, E. Elalaoui, T.J. Nickels, V. Jou, B.P. English, Q. Zheng, E. Luk, L.D. Lavis, T. Lionnet, and C. Wu. 2020. Live-cell single particle imaging reveals the role of RNA polymerase II in histone H2A.Z eviction. *Elife*. 9.

Rhee, H.S., A.R. Bataille, L. Zhang, and B.F. Pugh. 2014. Subnucleosomal structures and nucleosome asymmetry across a genome. *Cell*. 159:1377-1388.

Rigort, A., F.J. Bauerlein, A. Leis, M. Gruska, C. Hoffmann, T. Laugks, U. Bohm, M. Eibauer, H. Gnaegi, W. Baumeister, and J.M. Plitzko. 2010. Micromachining tools and correlative approaches for cellular cryo-electron tomography. *J Struct Biol*. 172:169-179.

Robinson, P.J., L. Fairall, V.A. Huynh, and D. Rhodes. 2006. EM measurements define the
dimensions of the "30-nm" chromatin fiber: evidence for a compact, interdigitated
structure. *Proc Natl Acad Sci U S A*. 103:6506-6511.

Rohou, A., and N. Grigorieff. 2015. CTFFIND4: Fast and accurate defocus estimation from
electron micrographs. *J Struct Biol*. 192:216-221.

Ruskin, R.S., Z. Yu, and N. Grigorieff. 2013. Quantitative characterization of electron detectors
for transmission electron microscopy. *J Struct Biol*. 184:385-393.

Sanulli, S., M.J. Trnka, V. Dharmarajan, R.W. Tibble, B.D. Pascal, A.L. Burlingame, P.R. Griffin,
J.D. Gross, and G.J. Narlikar. 2019. HP1 reshapes nucleosome core to promote phase
separation of heterochromatin. *Nature*. 575:390-394.

Scheffer, M.P., M. Eltsov, and A.S. Frangakis. 2011. Evidence for short-range helical order in
the 30-nm chromatin fibers of erythrocyte nuclei. *Proceedings of the National Academy*
of Sciences. 108:16992-16997.

Scheres, S.H. 2012. RELION: implementation of a Bayesian approach to cryo-EM structure
determination. *J Struct Biol*. 180:519-530.

Song, F., P. Chen, D. Sun, M. Wang, L. Dong, D. Liang, R.M. Xu, P. Zhu, and G. Li. 2014. Cryo-
EM study of the chromatin fiber reveals a double helix twisted by tetranucleosomal units.
Science. 344:376-380.

Strunk, K.M., K. Wang, D. Ke, J.L. Gray, and P. Zhang. 2012. Thinning of large mammalian
cells for cryo-TEM characterization by cryo-FIB milling. *J Microsc*. 247:220-227.

Studer, D., A. Klein, I. Iacovache, H. Gnaegi, and B. Zuber. 2014. A new tool based on two
micromanipulators facilitates the handling of ultrathin cryosection ribbons. *J Struct Biol*.
185:125-128.

Suloway, C., J. Shi, A. Cheng, J. Pulokas, B. Carragher, C.S. Potter, S.Q. Zheng, D.A. Agard,
and G.J. Jensen. 2009. Fully automated, sequential tilt-series acquisition with Leginon. *J*
Struct Biol. 167:11-18.

Sundaramoorthy, R., A.L. Hughes, H. El-Mkami, D.G. Norman, H. Ferreira, and T. Owen-
Hughes. 2018. Structure of the chromatin remodelling enzyme Chd1 bound to a
ubiquitinylated nucleosome. *Elife*. 7.

Sundaramoorthy, R., A.L. Hughes, V. Singh, N. Wiechens, D.P. Ryan, H. El-Mkami, M.
Petoukhov, D.I. Svergun, B. Treutlein, S. Quack, M. Fischer, J. Michaelis, B. Bottcher,
D.G. Norman, and T. Owen-Hughes. 2017. Structural reorganization of the chromatin
remodeling enzyme Chd1 upon engagement with nucleosomes. *Elife*. 6.

Tacke, S., P. Erdmann, Z. Wang, S. Klumpe, M. Grange, J. Plitzko, and S. Raunser. 2021. A
streamlined workflow for automated cryo focused ion beam milling. *J Struct Biol*.
213:107743.

Tegunov, D., L. Xue, C. Dienemann, P. Cramer, and J. Mahamid. 2021. Multi-particle cryo-EM
refinement with M visualizes ribosome-antibiotic complex at 3.5 A in cells. *Nat Methods*.
18:186-193.

Tivol, W.F., A. Briegel, and G.J. Jensen. 2008. An improved cryogen for plunge freezing.
Microsc Microanal. 14:375-379.

Turnbaugh, C., J.J. Axelrod, S.L. Campbell, J.Y. Dioquino, P.N. Petrov, J. Remis, O. Schwartz,
Z. Yu, Y. Cheng, R.M. Glaeser, and H. Mueller. 2021. High-power near-concentric
Fabry-Perot cavity for phase contrast electron microscopy. *Rev Sci Instrum*. 92:053005.

Uchida, M., Y. Sun, G. McDermott, C. Knoechel, M.A. Le Gros, D. Parkinson, D.G. Drubin, and
C.A. Larabell. 2011. Quantitative analysis of yeast internal architecture using soft X-ray
tomography. *Yeast*. 28:227-236.

van den Hoek, H., N. Klena, M.A. Jordan, G. Alvarez Viar, R.D. Righetto, M. Schaffer, P.S.
Erdmann, W. Wan, S. Geimer, J.M. Plitzko, W. Baumeister, G. Pigino, V. Hamel, P.
Guichard, and B.D. Engel. 2022. In situ architecture of the ciliary base reveals the
stepwise assembly of intraflagellar transport trains. *Science*. 377:543-548.

Verzijlbergen, K.F., V. Menendez-Benito, T. van Welsem, S.J. van Deventer, D.L. Lindstrom, H.
Ovaa, J. Neefjes, D.E. Gottschling, and F. van Leeuwen. 2010. Recombination-induced
tag exchange to track old and new proteins. *Proc Natl Acad Sci U S A*. 107:64-68.

Villa, E., M. Schaffer, J.M. Plitzko, and W. Baumeister. 2013. Opening windows into the cell:
focused-ion-beam milling for cryo-electron tomography. *Curr Opin Struct Biol*. 23:771-
777.

Wang, Q., C.P. Mercogliano, and J. Lowe. 2011. A ferritin-based label for cellular electron
cryotomography. *Structure*. 19:147-154.

White, C.L., R.K. Suto, and K. Luger. 2001. Structure of the yeast nucleosome core particle
reveals fundamental changes in internucleosome interactions. *EMBO J*. 20:5207-5218.

Willhoft, O., M. Ghoneim, C.L. Lin, E.Y.D. Chua, M. Wilkinson, Y. Chaban, R. Ayala, E.A.
McCormack, L. Ocloo, D.S. Rueda, and D.B. Wigley. 2018. Structure and dynamics of
the yeast SWR1-nucleosome complex. *Science*. 362.

Yakovlev, S., and K.H. Downing. 2011. Freezing in sealed capillaries for preparation of frozen
hydratedsections. *J Microsc*. 244:235-247.

Zachs, T., A. Schertel, J. Medeiros, G.L. Weiss, J. Hugener, J. Matos, and M. Pilhofer. 2020.
Fully automated, sequential focused ion beam milling for cryo-electron tomography.
Elife. 9.

Zheng, S.Q., E. Palovcak, J.P. Armache, K.A. Verba, Y. Cheng, and D.A. Agard. 2017.
MotionCor2: anisotropic correction of beam-induced motion for improved cryo-electron
microscopy. *Nat Methods*. 14:331-332.

Zhou, B.R., J. Jiang, H. Feng, R. Ghirlando, T.S. Xiao, and Y. Bai. 2015. Structural Mechanisms
of Nucleosome Recognition by Linker Histones. *Mol Cell*. 59:628-638.

Zhou, K., G. Gaullier, and K. Luger. 2019. Nucleosome structure and dynamics are coming of
age. *Nat Struct Mol Biol.* 26:3-13.

Zhou, M., L. Dai, C. Li, L. Shi, Y. Huang, Z. Guo, F. Wu, P. Zhu, and Z. Zhou. 2021. Structural
basis of nucleosome dynamics modulation by histone variants H2A.B and H2A.Z.2.2.
EMBO J. 40:e105907.

Zivanov, J., J. Oton, Z. Ke, A. von Kugelgen, E. Pyle, K. Qu, D. Morado, D. Castano-Diez, G.
Zanetti, T.A.M. Bharat, J.A.G. Briggs, and S.H.W. Scheres. 2022. A Bayesian approach
to single-particle electron cryo-tomography in RELION-4.0. *Elife.* 11.

Zlatanova, J., T.C. Bishop, J.M. Victor, V. Jackson, and K. van Holde. 2009. The nucleosome
family: dynamic and growing. *Structure.* 17:160-171.
1303

1304

1305
Figure 1. Canonical nucleosomes *in vitro* and *in situ*
(A) Space-filling model of the reconstituted yeast nucleosome crystal structure (PDB
1ID3) (White et al., 2001), showing from left to right, the disc, side, and gyre views. The
pseudo-dyad axis is indicated by the arrow. The DNA is rendered as light blue and the
histones in the core are shaded blue (H3), green (H4), red (H2B), and yellow (H2A). (B)
Subtomogram average of nucleosomes from wild-type (BY4741) yeast nuclear lysates.
The linker DNA is indicated by the short arrows and the DNA gyre motifs are indicated
by the arrowheads. (C) Subtomogram averages of nucleosomes in wild-type
cryolamellae, oriented similarly to the nucleosomes in the other two panels. The upper
(blue) class has more ordered linker DNA than the lower (magenta) class. Note that the
subtomogram average in panel B looks different from those in panel C (and in Figures 3
and S17) because it is at higher resolution (18 Å vs 24 Å).

1318
Figure 2. Canonical nucleosomes are rare in wild-type cells
(A) VPP tomographic slice (12 nm) of a BY4741 cell cryolamella. Large subcellular
structures are labeled: nuclear pore complex (NPC), nuclear megacomplex (M), nuclear
microtubule (MT), nuclear envelope (NE), and ribosome (R). The inset is a 4-fold
enlargement of the boxed area, and a nucleosome-like particle (n) is indicated. (B)
Remapped model of the two canonical nucleosome class averages in the tomogram
from panel A. The approximate location of the nuclear envelope is indicated by the blue
dashed line. The insets B1 and B2 show 4-fold enlargements of the corresponding
boxed areas. Note that the remapped model projects the full 150 nm thickness of this
cryolamella.

1329
Figure 3. Visualization of GFP-tagged nucleosomes *in vitro*
(A) Example 2-D class averages of nucleosome-like particles that were template-
matched with a featureless cylinder reference. Out of 88,896 template-matched
particles, 66,328 were retained after 2-D classification. (B) Class averages (3-D) of
nucleosomes from nuclear lysates. Solid arrowheads indicate the extra (GFP) densities.
Open arrowheads indicate the positions that lack this density. These class averages
were obtained after classification directly from subtomogram averaging, without an
intervening 2-D classification step. (C) The approximate positions of the H2A C-termini
are rendered in yellow and indicated by arrows in the crystal structure of the yeast
nucleosome (White et al., 2001). Note that in the crystal structure, fewer of the H2A C-
terminal amino acids were modeled than for H2B, meaning that the H2A C-terminus is
not perfectly ordered. To facilitate comparison, this structure is oriented like the class
averages in panel B. The nucleosome densities in panel B are longer along the pseudo-
dyad axis (horizontal) because they have linker DNA, which is absent in the
nucleosome crystal structure.

Figure 4. Canonical nucleosome class averages are not detected in LGY0016 (H2A-GFP) *in situ*

Class averages (3-D) of nucleosome-like particles in VPP cryotomograms of LGY0016 cryolamellae. The starred classes have two linear motifs. Movie S5 shows the progress of this classification job.

1351
Figure 5. **Models of yeast nucleosome heterogeneity**
Schematics of DNA (light blue) and histones (shaded pie slices) in the nucleosome disc
view. The cartoons only illustrate the 147 bp of "core" DNA. (A) Canonical nucleosome,
in which all the histones and 147 bp of DNA are part of an ordered complex. (B)
Nucleosome with alternative histone H2A-H2B (yellow, red) conformations and partially
dissociated DNA. (C) Nucleosome bound to non-histone proteins (gray). The blurred
gray box represents different proteins that can bind, thereby contributing to
constitutional heterogeneity. The blurred appearance represents a large range of
positions and orientations that protein and DNA components adopt inside cells, which
would result in the absence of a class average resembling a canonical nucleosome. (D)
Canonical nucleosomes are a minority conformation *in situ*.

1363
Figure S1. Subtomogram classification workflow
Classification starts with subtomograms that contain nucleosome-like particles, template matched in PEET (Cai et al., 2018). (A) In the workflow recommended by Bharat (Bharat et al., 2015), the subtomograms are first subjected to sequential rounds of 2-D classification to remove particles that belong to 'bad' classes. Once the bad classes are removed, the remaining set is subjected to 3-D classification. (B) In the alternative "direct 3-D" classification workflow, the subtomograms are subjected directly to 3-D classification. The 2-D classification steps (greyed out) are bypassed.

Figure S2. Overview of BY4741 (Wild type) nuclear lysate
Tomographic slice (12 nm) of BY4741 nuclear lysates. Some non-chromatin features are indicated: granule (G), coated vesicle (V), naked DNA (DNA), membrane fragments (membrane), carbon support film (carbon), virus-like particle (VLP), and gold fiducial (Au). The abundant granular densities in this field of view are nucleosome-like particles. One example is indicated in the 4-fold enlargement in the inset (n). The linear features at the lower left and upper right are back-projection artifacts from the image borders. These regions are excluded from analysis.

1381
Figure S3. Direct 3-D classification of BY4741 (Wild type) nuclear lysates
(A) Class averages (3-D) of two types of BY4741 nucleosomes. The canonical
nucleosome class averages are shaded blue and the percent of particles belonging to
the canonical classes are indicated under the density map. (B) Second round of
classification, using the classes indicated in panel A. (C) The resolution of the refined
canonical nucleosome is ~18 Å by the FSC = 0.5 criterion. The refined density map is
reproduced from Figure 1B.

1389

Figure S4. Overview of a BY4741 (Wild type) cryolamella
Tomographic slice (12 nm) of a BY4741 cryolamella. Some non-chromatin features are indicated: nuclear envelope (NE), nuclear microtubule (MT), nucleosome-like particle (n), megacomplex (M), ribosome (R), and a putative vacuole (Vac). The high-contrast linear features at the lower left are back-projection artifacts from image borders.

1395
Figure S5. Classification of BY4741 (Wild type) nucleosome-like particles *in situ*
(A) Class averages (2-D) of template-matched nucleosome-like particles in BY4741
(wild-type) cells. The class averages whose member particles were selected for
additional rounds of classification are indicated by a green triangle in the box's lower
right corner. The starting dataset consisted of 62,428 template-matched candidate
BY4741 nucleosomes from 3 tomograms, from which 16,608 particles were selected by
2-D classification. (B) Class averages (3-D) of nucleosome-like particles in BY4741
cells.

1404
1405 **Figure S6. Direct 3-D classification of BY4741 (Wild type) cryolamellae densities**
1406 Class averages (3-D) of nucleosome-like particles in cryotomograms of BY4741
1407 cryolamellae. In this experiment, 2-D classification was bypassed.

1408
Figure S7. Classification using the nucleosome crystal structure reference
(A) A featureless cylinder (left) and a simulated EM map from a crystal structure of the
nucleosome (right). (B) Class averages (2-D) of BY4741 (wild type) nucleosome-like
particles that were template matched using the nucleosome crystal structure. Classes
that were pooled for 3-D classification are indicated by a green triangle in the lower right
corner of its box. Out of 72,190 template-matched particles, 6,439 were selected by 2-D
classification. (C) Class averages (3-D) of the most nucleosome-like particles selected
by 2-D classification. The classification was performed using the same dataset as for
Figures S5 and S6.

BY4741 (Wild type), VPP

1420

1421 **Figure S8. Overview of a BY4741 (Wild type) cryolamella, VPP data**
1422 Tomographic slice (12 nm) of a BY4741 cryolamella imaged with a VPP. The nuclear
1423 envelope (NE), nuclear pore complex (NPC), ribosome (R), and megacomplex (M) are
1424 indicated. The inset shows a threefold enlargement of the boxed area. A nucleosome-
1425 like complex (n) is indicated.

1426
Figure S9. Classification of nucleosome-like particles *in situ* in BY4741 (Wild type) VPP data

(A) Class averages (2-D) of template-matched nucleosome-like particles in VPP cryotomograms of BY4741 (wild-type) cell cryolamellae. The class averages whose member particles were selected for additional rounds of classification are indicated by a green triangle in the box's lower right corner. The starting dataset consisted of 129,473 template-matched candidate BY4741 nucleosomes from 3 tomograms. 12,551 particles were selected by 2-D classification. (B) Class averages (3-D) of nucleosome-like particles in BY4741 cells image with a VPP.

1436
Figure S10. Direct 3-D classification of BY4741 (Wild type) nuclei in VPP
tomograms of cryolamellae
(A) Class averages (3-D) of BY4741 nucleosomes. The canonical nucleosome class
average is shaded blue and the percent of particles belonging is indicated under the
density map. Note that this percentage is relative to the 129,473 template-matched
nucleosome-like particles, not the ~25,000 expected number of nucleosomes in the 5
tomograms. (B) Second round of classification from the canonical nucleosome class
identified in panel A. (C) The resolution of both class averages is ~24 Å by the FSC =
0.5 criterion. The refined density maps are reproduced from Figure 1C and the number
of particles per class is labeled in black. (D) Angular-distribution plot of the canonical
nucleosome class from panel A. Movies S1 and S2 show the progress of this
classification jobs in panels A and B, respectively.

1449
Figure S11. Overview of a BY4741 (Wild type) cryolamella imaged in the cytoplasm, VPP data
Tomographic slice (12 nm) of a BY4741 cryolamella imaged in the cytoplasm. The vacuoles (V), a ribosome (R), and cell wall (CW) are indicated.

Figure S12. Classification of BY4741 (Wild type) cryolamellae VPP densities from the cytoplasm

Class averages (3-D) of "nucleosome-like" particles that were template matched from the cytoplasm. Both the template-matching and 3-D classification parameters were identical to the ones used to analyze the nucleus. 2-D classification was bypassed. One class has no contributing particles and was removed.

Figure S13. Strategy to tag H2A with GFP

(A) Crystal structure of the yeast nucleosome, PDB 1ID3 (White et al., 2001), in the disc view (left) and side view (right). DNA, dark purple; histone octamer, light gray. One set of histone C-termini are labeled and rendered as space-filling models. Two key points of reference, pseudo-dyad and opposite pseudo-dyad, are indicated by the arrowhead and arrow tail, respectively. The other set of C-termini is related by a 180° rotation around the pseudo-dyad axis. (B) Two views of the GFP crystal structure, PDB 1GFL (Yang et al., 1996), rendered at the same scale as panel A. The N-terminus is rendered as a black space-filling model. (C) Details of the sequence at the H2A-GFP fusion, with (7aa) and without (0aa) the 7 amino-acid linker. The H2A C-terminal sequence is shaded yellow, GFP N-terminal sequence is shaded green, and the flexible linker is shaded gray. LGY0016 has a 0aa linker. (D) Histone genotype of the parent strain BY4741. (E) Homologous recombination strategy to C-terminally tag the sole copy of the H2A gene with GFP. Upper – replacement of the HTA2-HTB2 locus with KanR. Lower – Tagging of HTA1 with GFP. The junction between the HTA1 (H2A) and GFP is boxed. (F) Histone genotype of LGY0016.

1478
Figure S14. Experimental verification of H2A-GFP and H2B-GFP tagging
(A) Maps of the two H2A-H2B loci in the parent (wild-type) strain BY4741 and the H2A-
GFP strain LGY0016. Primers used for PCR verification are indicated with the half
arrow symbols. (B) Agarose gel of PCR amplicons expected (or not) from LGY0016

genomic DNA, in which the HTB2-HTA2 locus is deleted and the HTA1 locus is tagged
with GFP. (C) Immunoblot analysis of strains LGY0016 (expresses H2A-GFP only) and
LGY0012 (expresses H2A only, control). The α-GFP antibody correctly detected the
large H2A-GFP fusion protein in LGY0016 (lane 4) but not in LGY0012 (lane 3, negative
control). The α-H2A antibody detected H2A-GFP in LGY0016 and H2A in LGY0012. (D)
DIC and GFP fluorescence confocal microscopy for LGY0016 cells. In the left panel, the
GFP signals are overlaid in green. In the right panel, GFP signals are rendered with
inverted contrast. (E) Map of the HTA1-HTB1 locus before (upper) and after (lower) the
GFP insertion, with validation primers indicated. The color scheme is the same as in
panel A. Note that LGY0015 still has the HTA2-HTB2 gene pair and therefore
expresses untagged H2B. (F) Validation PCR for LG0015. (G) Validation immunoblots
for strains BY4741 (wild type) and LGY0015 (expresses H2B and H2B-GFP). The α-
GFP detected a ~40 kDa protein (H2B-GFP) in LGY0015 but not BY4741. The α-H2B
antibody detects the 40 kDa (H2B-GFP) in LGY0015 but not BY4741. Note that the α-
H2B antibody has poorer specificity and does not generate a strong signal for the
untagged H2B. (H) In the left panel, the GFP signals are overlaid in green. In the right
panel, GFP signals are rendered with inverted contrast.

Figure S15. Overview of LGY0016 (H2A-GFP) nuclear lysate
Tomographic slice (12 nm) of LGY0016 nuclear lysates. Some non-chromatin features
are indicated: carbon support film (carbon), membrane fragments (membrane), gold
fiducial (Au), and virus-like particle (VLP). The abundant granular densities in this field
of view are nucleosomes. One small subarea (boxed) is enlarged 3-fold in the inset. A
nucleosome-like particle (n) is indicated.

1507
Figure S16. Overview of LGY0015 (H2B, H2B-GFP) nuclear lysate
Tomographic slice (12 nm) of LGY0015 nuclear lysates. Some non-chromatin features
are indicated: carbon support film (carbon), membrane fragments (membrane), gold
fiducial (Au), and virus-like particle (VLP). The abundant granular densities in this field
of view are nucleosomes. One small subarea (boxed) is enlarged 3-fold in the inset. A
nucleosome-like particle (n) is indicated.

1514
Figure S17. Classification of LGY0015 (H2B, H2B-GFP) nuclear lysates
(A) Class averages (2-D) of nucleosome-like particles. Out of 96,979 template-matched
particles from 6 tomograms, 56,872 particles were selected by 2-D classification. (B)
Class averages (3-D) of two types of LGY0015 nucleosomes. The solid arrowhead
indicates the position of the GFP tag whereas the open arrowheads indicate the
positions that lack this density. These are the same class averages shown in the overall
workflow in Figure S19. (C) Two views of the nucleosome crystal structure, indicating
the location of the H2B C-terminus (salmon).

Figure S18. Direct 3-D classification of LGY0016 (H2A-GFP) lysates
(A) Class averages (3-D) of nucleosome-like particles from nuclear lysates of LGY0016 cells. The canonical nucleosome-like class averages are shaded blue while the non-canonical nucleosome averages are shaded gray. Movie S3 shows the convergence of the classification and shows more views of these class averages. (B) The second round of 3-D classification, using the canonical nucleosomes from panel A. See Movie S4 for more details of this classification job. (C) Refined densities of three types of LGY0016 nucleosomes isolated from nuclear lysates; reproduced from Figure 3B. The class numbers correspond to those in panel B (blue text) while the number of particles per class are labeled in black. The Fourier shell correlation (FSC) plot of the three refined class averages is labeled with the same numbering scheme. The resolution is ~26 Å by the FSC = 0.5 criterion.

1536
Figure S19. Direct 3-D classification of LGY0015 (H2B, H2B-GFP) lysates
(A) Class averages of nucleosome-like particles from nuclear lysates of LGY0015 cells.
The canonical nucleosome class averages are shaded blue while the non-canonical
nucleosome averages are shaded gray. (B) The second round of 3-D classification,
using the canonical nucleosomes from panel A. Note that some class averages, such as
the gray one, have a density that is not connected to the nucleosome. This density is
from a nearby particle that protruded into the mask. (C) The two types of LGY0015
nucleosomes are reproduced from Figure S17B. The numbers of particles per class are
indicated below the density maps in black. The FSC plot of the LGY0015 nuclear
lysates nucleosome class averages in panel D uses the same numbering scheme. The
resolution is ~26 Å by the FSC = 0.5 criterion.

1548
1549 **Figure S20. Overview of a LGY0016 (H2A-GFP) cryolamella**
1550 Tomographic slice (12 nm) of a LGY0016 cryolamella. Some non-chromatin features
1551 are indicated: nuclear envelope (NE), nuclear microtubule (MT), megacomplex (M), and
1552 ribosome (R). The inset is a threefold enlargement of the boxed area. A nucleosome-
1553 like particle (n) is indicated.

LGY0016 (H2A-GFP), VPP

1554

1555 **Figure S21. Overview of a LGY0016 (H2A-GFP) cryolamella, VPP data**
1556 Tomographic slice (12 nm) of a LGY0016 cryolamella. The nuclear envelope (NE),
1557 nuclear microtubule (MT), and a megacomplex (M) are indicated. The inset is a
1558 threefold enlargement of the boxed area. A nucleosome-like particle (n) is indicated.

1559
1560 **Figure S22. Overview of a LGY0016 (H2A-GFP) cryosection, VPP data**
1561 Tomographic slice (12 nm) of a LGY0016 cryosection. Some non-chromatin features
1562 are indicated: nuclear envelope (NE), megacomplex (M), and ribosome (R). The inset is
1563 a threefold enlargement of the boxed area. A nucleosome-like particle (n) is indicated.
1564 The smeared features at the lower left are back-projection artifacts from image borders.

1565
1566 **Figure S23. Direct 3-D classification of LGY0016 (H2A-GFP) cryolamellae**
1567 **densities**
1568 Class averages (3-D) of nucleosome-like particles in cryotomograms of LGY0016
1569 cryolamellae. The starred classes have two linear motifs.

Figure S24. Direct 3-D classification of LGY0016 (H2A-GFP) cryosection densities in VPP data

Class averages (3-D) of nucleosome-like particles in VPP cryotomograms of LGY0016 cryolamellae. The starred classes have two linear motifs.

1575
Figure S25. Control VPP subtomogram analysis of ribosomes *in situ*
(A) Candidate ribosomes were template matched from a single VPP tomogram of the
cytoplasm from Figure S11, using a 25-nm-diameter sphere as a reference. Direct 3-D
classification into ten classes led to six non-empty classes, of which two were of
ribosomes. Note that template matching is insufficient to uniquely identify a
macromolecular complex, particularly when the reference is a featureless body like a
sphere or cylinder. A simple reference was used to reduce the model bias. Template
matching done with featureless references generates substantial numbers of false
positives, which are confidently removed by classification. Complexes are only
considered as identified after the 3-D classification step produces recognizable class
averages. Therefore, the term "candidate ribosomes" is used to describe the 3,816
template matching hits. (B) The 1,150 ribosome subtomograms were pooled and 3-D
refined, producing an average at 28 Å resolution by the "Gold-standard" FSC (0.143
cutoff) criterion and 33 Å resolution with an FSC = 0.5 cutoff. (C) Comparison of the *in*
situ ribosome subtomogram average (STA) with density maps simulated from the yeast
ribosome crystal structure (Ben-Shem et al., 2011) at three different resolutions.

1592
Figure S26. Verification of (H3, H3-GFP) strains
(A) Map of the HHT1-HHF1 locus in the parent (Wild type) strain BY4741 and the H3-
GFP strains LGY0002 (H3, H3-RIPGLIN-GFP), LGY0007 (H3, H3-0aa-GFP) and
LGY0070 (H3, H3-GGSGGS-GFP). Primers used for PCR verification are indicated with
the half arrow symbols. (B) Agarose gel of PCR amplicons expected from LGY0002,
LGY0007 and LGY0070 genomic DNA, in which the HHT1 locus is tagged with GFP.
The only difference between the strains is the linker separating HHT1 and the GFP,
which is too small to observe a difference in PCR amplicon sizes. (C) Immunoblot
analysis of strains LGY0002, LGY0007 and LGY0070. The α-GFP antibody correctly

detected the large H3-GFP fusion protein in LGY0002 (lane 3), LGY0007 (lane 4) and
LGY0070 (lane 5), but not in BY4741 (lane 2, negative control). The α-H3 antibody
detected H3 in all lanes, but failed to detect H3-GFP in LGY0002, LGY0007 and
LGY0070. (D) DIC and GFP fluorescence confocal microscopy for LGY0002, LGY0007
and LGY0070 cells. In the left panel, the GFP signals are overlaid in green. In the right
panel, GFP signals are rendered with inverted contrast.

Figure S27. Verification of (H4, H4-GFP) strain

(A) Map of the HHT1-HHF1 locus in the parent (wild-type) strain BY4741 and the H4-GFP strain LGY0071. Primers used for PCR verification are indicated with the half arrow symbols. (B) Agarose gel of PCR amplicons expected from LGY0071 genomic DNA, in which the HHF1 locus is tagged with GFP. (C) Immunoblot analysis of strain LGY0071. The α-GFP antibody correctly detected the large H3-GFP fusion protein in LGY0071 (lane 3), but not in BY4741 (lane 2, negative control). The α-H4 antibody detected H4-GFP in LGY0071 and H4 in both lanes. (D) DIC and GFP fluorescence confocal microscopy for LGY0002, LGY0007 and LGY0070 cells. In the left panel, the GFP signals are overlaid in green. In the right panel, GFP signals are rendered with inverted contrast.

1621
Figure S28. Direct 3-D classification of LGY0007 (H3, H3-0aa-GFP) nuclear lysates
(A) Class averages (3-D) of nucleosome-like particles from nuclear lysates of LGY0007
cells, which express a wild-type copy of H3 and an H3-GFP that has no linker
sequence. The canonical nucleosome-like class averages with an additional GFP
density are shaded blue while the non-canonical nucleosome averages and canonical
nucleosome-like class averages without any additional density are shaded gray. (B) The
second round of 3-D classification, using the canonical nucleosomes with an additional
GFP density from panel A. The remove duplicates function of RELION was used to
remove particles (defined as particles within a radius of 180 Å from another particle)
from the indicated class average for the 3-D refinement. Twenty-seven duplicate
particles were removed before refining. (C) Refined density of LGY0007 nucleosomes
isolated from nuclear lysates. The number of particles in the chosen class is labeled in
black. The resolution is ~23 Å by the FSC = 0.5 criterion.

1635
1636 **Figure S29. Direct 3-D classification of LGY0002 (H3, H3-RIPGLIN-GFP) nuclear**
1637 **lysates**
1638 Class averages (3-D) of nucleosome-like particles from nuclear lysates of LGY0002
1639 cells, which express a wild-type copy of H3 and an H3-GFP that has a RIPGLIN linker
1640 sequence. In this experiment, 2-D classification was bypassed.

Figure S30. Direct 3-D classification of LGY0070 (H3, H3-GGSGGS-GFP) nuclear lysates

Class averages (3-D) of nucleosome-like particles from nuclear lysates of LGY0070 cells, which express a wild-type copy of H3 and an H3-GFP that has a GGSGGS linker sequence. In this experiment, 2-D classification was bypassed.

1647
Figure S31. Direct 3-D classification of LGY0071 (H4, H4-GGSGGS-GFP) nuclear lysates

Class averages (3-D) of nucleosome-like particles from nuclear lysates of LGY0071 cells, which express a wild-type copy of H4 and an H4-GFP that has a GGSGGS linker sequence. In this experiment, 2-D classification was bypassed.

LGY0007 (H3, H3-GFP), VPP

M→

n→

10 nm

NE

10 nm

1654

 Figure S32. Overview of a LGY0007 (H3, H3-GFP) cryolamella
 Tomographic slice (12 nm) of a LGY0007 cryolamella. Some non-chromatin features
 are indicated: nuclear envelope (NE), megacomplex (M), and ribosome (R). The inset is
 a threefold enlargement of the boxed area. A nucleosome-like particle (n) is indicated.

1660
1661 **Figure S33. Direct 3-D classification of LGY0007 (H3, H3-GFP) cryolamellae**
1662 **densities**
1663 Class averages (3-D) of nucleosome-like particles in cryotomograms of LGY0007
1664 cryolamellae. In this experiment, 2-D classification was bypassed.

Figure S34. Simulated tomographic slices of single and stacked nucleosomes
Atomic models (upper, from PDB 1KX5) were used to simulate tomographic slices (12 nm, lower). (A and B) Mononucleosomes in the gyre and side views, respectively. (C and D) Two examples of stacked nucleosomes. In panel C, both nucleosomes have the same orientation. In panel D, the upper nucleosome is rotated 90° along the Y axis relative to the lower nucleosome. The stacked nucleosomes were separated by 55 Å center-to-center along the Y axis to emulate a worst-case (for image processing) scenario. Note that these simulated tomographic slices are not intended to accurately model the image-formation process. They serve to show how different side/gyre views of stacked nucleosomes appear.

1676
Figure S35. Fourier power spectra analysis of tilt series images
An image corresponding to a ~12° pre-tilt (to make the lamella perpendicular to the
electron-optical axis to maximize the signal-to-noise ratio) was extracted from tilt series
20211117_004 and then Fourier transformed using (A) CTFFIND and (B) IMOD
ctfplotter. Thon rings could not be detected in the CTFFIND 2-D diagnostic image or the
IMOD 1-D radial average of the Fourier power spectrum.
1683

Movie S1. Direct 3-D classification of BY4741 (Wild type) nucleosome-like particles in VPP tomograms of cell cryolamellae, round 1

The progress of 30 rounds of 3-D classification is shown. There are 100 classes, initialized with a smooth nucleosome-sized cylindrical reference. The final iteration (30) is also shown in Figure. S10, with only the most nucleosome-like classes shaded.

Movie S2. Direct 3-D classification of BY4741 (Wild type) nucleosome-like particles in VPP tomograms of cell cryolamellae, round 2

The most nucleosome-like classes from round 1 were selected and subjected to a second round of classification, with 4 classes, again initialized with a smooth nucleosome-sized cylindrical reference.

Movie S3. Direct 3-D classification of LGY0016 (H2A-GFP) lysate nucleosomes, round 1

The progress of 30 rounds of 3-D classification is shown. There are 40 classes, initialized with a smooth nucleosome-sized cylindrical reference. The final iteration (30) is also shown in Figure. S18, with only the most nucleosome-like classes shaded.

Movie S4. Direct 3-D classification of LGY0016 (H2A-GFP) lysate nucleosomes, round 2

The most nucleosome-like classes from round 1 were selected and subjected to a second round of classification, with 10 classes, again initialized with a smooth nucleosome-sized cylindrical reference. Two "junk" classes were removed.

Movie S5. Direct 3-D classification of LGY0016 (H2A-GFP) nucleosome-like particles in VPP tomograms of cell cryolamellae, round 1

3-D classification analysis of LGY0016 (H2A-GFP) nucleosome-like particles from cryolamellae imaged with a VPP. The initial reference is a featureless cylinder. There are 100 classes.

 Table S1. Genotypes of strains used in this paper

Strain	Parent	Genotype	Source
BY4741	--	*MATa his3D1 leu2Δ0 met15Δ0 ura3Δ0*	EUROSCARF
LGY0012	BY4741	*(hta2-htb2)Δ0::KANMX*	This paper
LGY0015	BY4741	*HTB1-GFP(S65T)(0aa linker)-HIS3MX*	This paper
LGY0016	LGY0012	*(hta2-htb2)Δ0::KANMX HTA1-GFP(S65T)(0aa linker)-HIS3MX*	This paper
LGY0002	BY4741	*HHT1-GFP(S65T)(RIPGLIN linker)-HIS3MX*	This paper
LGY0007	BY4741	*HHT1-GFP(S65T)(0aa linker)-HIS3MX*	This paper
LGY0070	BY4741	*HHT1-GFP(S65T)(GGSGGS linker)-HIS3MX*	This paper
LGY0071	BY4741	*HHF1-GFP(S65T)(GGSGGS linker)-HIS3MX*	This paper

1717

 Table S2. PCR primers, 5' → 3'

Editing	
HTA1-GFP Tag F	AAAGAAGTCTGCCAAGGCTACCAAGGCTTCTCAAGAATTAAGTAAAGGAGAAGAACTTTT
HTA1-GFP Tag R	TTTAGTTCCTTCCGCCTTCTTTAAAATACCAGAACCGATCGAATTCGAGCTCGTTTAAAC
HTB1-GFP Tag F	TACTAGAGCTGTTACCAAGTACTCTTCCTCTACTCAAGCAAGTAAAGGAGAAGAACTTTT
HTB1-GFP Tag R	TAAATAATAATATTAATTATAACCAAAGGAAGTGATTTCAGAATTCGAGCTCGTTTAAAC
HAB2 Del F	AAGAATGTTTGATTTGCTTTGTTTCTTTTCAACTCAGTTCCAGATCCGCTAGGGATAACA
HAB2 Del R	AAAAGAAAACATGACTAAATCACAATACCTAGTGAGTGACTCGATGAATTCGAGCTCG
HHT1-RIPGLIN-GFP Tag F	GGATATCAAGTTGGCTAGAAGATTAAGAGGTGAAAGATCACGGATCCCCGGGTTAATTAA
HHT1-GFP Tag F	GGATATCAAGTTGGCTAGAAGATTAAGAGGTGAAAGATCAAGTAAAGGAGAAGAACTTTT
HHT1-GFP Tag R	TTTTGTTCGTTTTTTACTAAAACTGATGACAATCAACAAAGAATTCGAGCTCGTTTAAAC
HHT1-GGSGGS-GFP Tag F	TCCAAAAGAAGGATATCAAGTTGGCTAGAAGATTAAGAGGTGAAAGATCAGGTGGATCTGGTGGATCTAGTAAAGGAGAAGAACTTTT
HHT1-GFP Tag R (Long)	TTTATTGTGTTTTTGTTCGTTTTTTACTAAAACTGATGACAATCAACAAAGAATTCGAGCTCGTTTAAAC
HHF1-GGSGGS-GFP Tag F	TTGTTTATGCTTTGAAGAGACAAGGTAGAACCTTATACGGTTTCGGTGGTGGTGGATCTGGTGGATCTAGTAAAGGAGAAGAACTTTT
HHF1-GFP Tag R	CGAATCCCAAATATTTGCTTGTTGTTACCGTTTTCTTAGAATTAGCTAAAGAATTCGAGCTCGTTTAAAC
Confirmation	
FA1	CGGTGGTAAAGGTGGTAAAG
RA1	TCGTTTCTGATAAACCAGGT
RG	CCGTTTCATATGATCTGGGT
FH	GACCATTTGCTGTAATCGAC
RK	CCTTATTTTTGACGAGGGGA
RB2	ATTAACCGGGATTCACTGAC
RA2	CAGTTCTTGAGAAGCTTTGG
RA2.2	CTGGACGAAGACGAAGTAAT
FB1	ATGTCTGCTAAAGCCGAAAA
RB1	AGTCAGCGACATCTGTCTTT
FT1	AAGCAAACAGCAAGAAAGTC
RT1	CTTCTGACAGCAAGGGTATT
FF1	ATGTCCGGTAGAGGTAAAGG
RF1	ACACACGAAAATCCTGTGAT
Sequencing	
FA1	CGGTGGTAAAGGTGGTAAAG
RA1	TCGTTTCTGATAAACCAGGT
RG	CCGTTTCATATGATCTGGGT
Histone-GFP Seq R1.2	CTGACCTACAGGAAAGAGTT
Histone-GFP Seq F2.1	CTGTCCTTTTACCAGACAAC

Histone-GFP Seq F2.2	CTCTTATTGACCACACCTCT
Histone-GFP Seq R2.2	CTTCTTTGCAGGAAGAATCG
Histone-GFP Seq F3.1	TCACATCACATCCGAACATA
FH	GACCATTTGCTGTAATCGAC
RA2.2	CTGGACGAAGACGAAGTAAT
HAB2Del Seq R1	ATTACCCTGTTATCCCTAGC
HAB2Del Seq R2	CAGGAGTACGGATAAAATGC
HAB2Del Seq R3	TATCGAATCGACAGCAGTAT
RK	CCTTATTTTTGACGAGGGGA
RB2	ATTAACCGGGATTCACTGAC
FB1	ATGTCTGCTAAAGCCGAAAA
RB1	AGTCAGCGACATCTGTCTTT
FT1	AAGCAAACAGCAAGAAAGTC
RT1	CTTCTGACAGCAAGGGTATT
FF1	ATGTCCGGTAGAGGTAAAGG
RF1	ACACACGAAAATCCTGTGAT

1719

1720 **Table S3. Antibodies for immunoblots**

Antigen	1⁰ antibody	2⁰ antibody	Dilution	
			1⁰	**2⁰**
H2A	Active Motif 39235	CST 7074S	1:1000	1:5000
H2B	Abcam ab1790	CST 7074S	1:1000	1:5000
H3	Abcam ab1791	CST 7074S	1:1000	1:5000
H4	Abcam ab10158	CST 7074S	1:1000	1:5000
GFP	Santa Cruz sc9996	CST 7076S	1:1000	1:5000

1721 CST = Cell Signaling Technology

1722

Table S4. Confocal microscopy details

General

Instrument	Olympus FV3000
Pinhole	1 Airy unit
X, Y pixel	0.087 [µm]
Z pixel	0.410 [µm]

Acquisition

Objective lens	UPLSAPO 60XO
Objective lens magnification	60×
Objective lens NA	1.35
Scan device	Galvano
Scan direction	One way
Dwell time	2.0 [µs/pixel]
Sequential mode	Line
Integration type	None
Integration count	0
Zoom	×2.37

GFP channel settings

Emission wavelength	510 [nm]
PMT voltage	530 – 615 [V]
C.A.	210 [µm]
Bits/pixel	12 [bits]
Laser wavelength	488 [nm]
Laser transmissivity	0.03 or 0.1 [%]
AOTF/AOM transmissivity	0.3 [%]
Laser ND filter	10 [%]
Detection wavelength	500 – 600 [nm]

DIC channel settings

PMT voltage	340 or 380 [V]
C.A.	210 [µm]
Bits/pixel	12 [bits]
Laser wavelength	488 [nm]
Laser transmissivity	0.01 or 0.03 [%]
AOTF/AOM transmissivity	0.3 [%]
Laser ND filter	10 [%]

1723

 Table S5. Cryo-ET details

Sample preparation	
EM grids	Lysates: C-flat CF-4/2-2C or Quantifoil R2/4 200
	Cryosections: continuous carbon
	Cryolamellae: Quantifoil R2/4 200
Plunge freezer	Vitrobot Mark IV
Cryomicrotome	UC7/FC7
Attachment device	Crion
Micromanipulators	Leica micromanipulator; Narishige MN-151-S
Cryomicrotome feed	70 or 100 nm
Cryo-FIB-SEM	Helios NanoLab 600 DualBeam
Cryo-transfer device	Quorum PP2000T
Milling currents	Rough: 2.8 nA; Intermediate: 0.28 nA; Polishing: 48 pA

Cryo-ET data collection	
Microscope	Titan Krios
Energy	300 keV
Camera: recording mode	Falcon II: integration
	K2, K3: super-resolution, movie frames
Energy filter width	K2, K3: 20 eV
Tomography software	TFS Tomo4, Leginon, SerialEM 3.8.6, PACE-tomo
Unbinned pixel size	Falcon II: 5.8Å, 4.6Å; K2: 2.8Å; K3: 3.4Å
Contrast mechanism	Defocus phase contrast (lysates and cryolamellae)
	Volta phase contrast (cryosections and cryolamellae)
Defocus (nominal)	Defocus phase contrast: –4 to –10 µm
	Volta phase contrast: 0 to –0.25 µm
Cumulative dose	$100 - 120 \, e^- / Å^2$
Dose fractionation	Falcon II & K2: 1 / cosine
	K3: (1/cosine)^(1/y), where y = 2 or 4
Tilt range	Lysates: ±60°; bidirectional, negative angles first (Falcon II)
	Lysates: ±60°; dose-symmetric (Hagen et al., 2017) (K3)
	Cryosections: ±60°; bidirectional, negative angles first
	Cryolamellae: –70° to +50°, dose-symmetric (Hagen et al., 2017)
Tilt increment	1°, 2°, or 3°

Cryo-ET data analysis	
Tomogram processing	IMOD 4.11
Template matching	PEET 1.15
Reference creation	Bsoft 1.8.8
Mask creation	Bsoft 1.8.8, RELION 3.0.8
Subtomogram analysis	RELION 3.0.8
Tomogram visualization	UCSF Chimera 1.13.1, IMOD 4.11
Auxiliary scripts	https://github.com/anaphaze/ot-tools
	https://github.com/Guillawme/cryoEM-scripts
Calculations	Google sheets
Figure/movie editing	Adobe Photoshop, Illustrator, and Premiere Pro CC

1725

 Table S6. Cryotomogram details

Tomogram	Strain	Sample	Fig	Dose (e/Å²)	Pixel size (Å)	Δf * (µm)	Δtilt (°)	Cam	VPP	thick (nm)	resid
20210601_012	BY4741	lysate	-	120	3.4	5-7	2	K3	–	100	0.48
20210601_014	BY4741	lysate	S2	120	3.4	5-7	2	K3	–	20-70	0.66
20210601_016	BY4741	lysate	-	120	3.4	5-7	2	K3	–	10-60	0.66
20210601_017	BY4741	lysate	-	120	3.4	5-7	2	K3	–	20-60	0.34
20210601_018	BY4741	lysate	-	120	3.4	5-7	2	K3	–	10-50	0.58
20210601_019	BY4741	lysate	-	120	3.4	5-7	2	K3	–	10-70	0.50
20210601_020	BY4741	lysate	-	120	3.4	5-7	2	K3	–	10-40	0.50
20210601_023	BY4741	lysate	-	120	3.4	5-7	2	K3	–	10-40	0.46
20210601_024	BY4741	lysate	-	120	3.4	5-7	2	K3	–	10-30	0.43
20201224_01	BY4741	lamella	-	100	3.4	7	1	K3	–	130-210	0.14
20201224_02	BY4741	lamella	-	100	3.4	8.5	1	K3	–	160	0.42
20201224_03	BY4741	lamella	S4	100	3.4	9	1	K3	–	190	0.18
20211117_004	BY4741	lamella	2	122	3.4	0.1	2	K3	+	150	0.27
20211117_009	BY4741	lamella	-	122	3.4	0.1	2	K3	+	140	0.33
20211117_037	BY4741	lamella	-	122	3.4	0.1	2	K3	+	120	0.28
20211117_049	BY4741	lamella	S11	122	3.4	0.1	2	K3	+	150	0.29
20211117_070	BY4741	lamella	S8	122	3.4	0.1	2	K3	+	150	0.23
20211117_075	BY4741	lamella	-	122	3.4	0.1	2	K3	+	110	0.23
20180730_01	LGY0016	lysate	S15	100	4.6	10	2	FII	–	40-100	0.17
20180730_02	LGY0016	lysate	-	100	4.6	9.2	2	FII	–	40-110	0.38
20180730_08	LGY0016	lysate	-	100	4.6	9.5	2	FII	–	40-100	0.27
20180730_09	LGY0016	lysate	-	100	4.6	9.5	2	FII	–	50-90	0.19
20180823_02	LGY0015	lysate	-	100	5.8	9.5	2	FII	–	20-80	0.29
20180823_03	LGY0015	lysate	S16	100	5.8	9.0	2	FII	–	20-85	0.37
20180823_04	LGY0015	lysate	-	100	5.8	10	2	FII	–	30-50	0.53
20180823_05	LGY0015	lysate	-	100	5.8	9.5	2	FII	–	20-50	0.38
20180823_06	LGY0015	lysate	-	100	5.8	9.5	2	FII	–	20-60	0.24
20180823_07	LGY0015	lysate	-	100	5.8	9.5	2	FII	–	40-80	0.32
20201207_10	LGY0016	lamella	-	120	3.4	7.5	1	K3	–	130	0.13
20201207_12	LGY0016	lamella	S20	120	3.4	8.5	1	K3	–	80	0.28
20201207_13	LGY0016	lamella	-	110	3.4	4.5	1	K3	–	100	0.27
20201207_17	LGY0016	lamella	-	120	3.4	5	1	K3	–	100	0.49
20211102_031	LGY0016	lamella	-	122	3.4	0	2	K3	+	80-180	0.32
20211102_032	LGY0016	lamella	-	122	3.4	0	2	K3	+	100	0.19
20211102_035	LGY0016	lamella	S21	122	3.4	0	2	K3	+	160	0.22
20181109_15	LGY0016	section	-	100	2.8	0.25	3	K2	+	75-90	0.81
20181109_42	LGY0016	section	-	100	2.8	0.25	3	K2	+	75-90	0.44
20181109_45	LGY0016	section	-	100	2.8	0.25	3	K2	+	75-90	0.75
20181109_47	LGY0016	section	-	100	2.8	0.25	3	K2	+	75-90	0.69
20190118_13	LGY0016	section	-	100	2.8	0.25	2	K2	+	75-90	0.83
20190118_16	LGY0016	section	-	100	2.8	0.25	2	K2	+	75-90	0.70
20190120_44	LGY0016	section	S22	100	2.8	0.25	2	K2	+	85-100	0.81
20220805_batch 04_ts_001	LGY0002	lysate	-	120	3.4	9-11	2	K3	–	90-115	0.14
20220805_batch 04_ts_002	LGY0002	lysate	-	120	3.4	10-12	2	K3	–	10-85	0.21
20220805_batch	LGY0002	lysate	-	120	3.4	9-13	2	K3	–	75	0.24

04_ts_006											
20220805_batch 06_ts_003	LGY0002	lysate	-	120	3.4	8-11	2	K3	–	10-60	0.26
20220805_batch 07_ts_001	LGY0002	lysate	-	120	3.4	10-11	2	K3	–	95-135	0.36
20220805_batch 22_ts_001	LGY0007	lysate	-	120	3.4	10	2	K3	–	10-100	0.17
20220805_batch 22_ts_002	LGY0007	lysate	-	120	3.4	9-10	2	K3	–	10-95	0.15
20220805_batch 23_ts_001	LGY0007	lysate	-	120	3.4	10	2	K3	–	10-105	0.20
20220805_batch 24_ts_001	LGY0007	lysate	-	120	3.4	9-11	2	K3	–	10-70	0.17
20220805_batch 24_ts_003	LGY0007	lysate	-	120	3.4	9-12	2	K3	–	60-100	0.19
20220805_batch 26_ts_004	LGY0007	lysate	-	120	3.4	10	2	K3	–	10-100	0.22
20220805_batch 29_ts_001	LGY0007	lysate	-	120	3.4	10-11	2	K3	–	10-105	0.17
20220805_batch 30_ts_001	LGY0007	lysate	-	120	3.4	10-11	2	K3	–	70	0.14
20220805_batch 31_ts_003	LGY0007	lysate	-	120	3.4	9-12	2	K3	–	40	0.16
20220805_batch 31_ts_005	LGY0007	lysate	-	120	3.4	10-11	2	K3	–	35-75	0.20
20220805_batch 17_ts_001	LGY0070	lysate	-	120	3.4	9-14	2	K3	–	60-90	0.25
20220805_batch 17_ts_003	LGY0070	lysate	-	120	3.4	9-12	2	K3	–	50-150	0.33
20220805_batch 17_ts_004	LGY0070	lysate	-	120	3.4	9-12	2	K3	–	10-60	0.68
20220805_batch 18_ts_002	LGY0070	lysate	-	120	3.4	9-11	2	K3	–	10-80	0.31
20220805_batch 18_ts_007	LGY0070	lysate	-	120	3.4	9-11	2	K3	–	10-90	0.20
20220805_batch 18_ts_008	LGY0070	lysate	-	120	3.4	9-10	2	K3	–	10-90	0.13
20220805_batch 19_ts_001	LGY0070	lysate	-	120	3.4	9-11	2	K3	–	10-155	0.75
20220805_batch 20_ts_001	LGY0070	lysate	-	120	3.4	9-10	2	K3	–	10-95	0.27
20220805_batch 08_ts_004	LGY0071	lysate	-	120	3.4	8-11	2	K3	–	10-60	0.12
20220805_batch 09_ts_003	LGY0071	lysate	-	120	3.4	10-12	2	K3	–	10-40	0.17
20220805_batch 09_ts_004	LGY0071	lysate	-	120	3.4	9-11	2	K3	–	10-60	0.38
20220805_batch 11_ts_002	LGY0071	lysate	-	120	3.4	10-11	2	K3	–	30-50	0.40
20220805_batch 12_ts_001	LGY0071	lysate	-	120	3.4	10-11	2	K3	–	25-70	0.21
20220805_batch 13_ts_001	LGY0071	lysate	-	120	3.4	9-12	2	K3	–	60-80	0.18

20220805_batch 13_ts_004	LGY0071	lysate	-	120	3.4	9-10	2	K3	–	10-80	0.19
20220805_batch 13_ts_006	LGY0071	lysate	-	120	3.4	9-12	2	K3	–	55-85	0.25
20220805_batch 15_ts_001	LGY0071	lysate	-	120	3.4	9-10	2	K3	–	10-65	0.21
20220805_batch 15_ts_003	LGY0071	lysate	-	120	3.4	7-10	2	K3	–	55-105	0.27
20221121_057	LGY0007	lamella	-	120	3.4	0	2	K3	+	150	0.68
20221121_060	LGY0007	lamella	-	120	3.4	0	2	K3	+	145	0.44
20221121_065	LGY0007	lamella	S32	120	3.4	0	2	K3	+	115	0.25
20221125_050	LGY0007	lamella	-	120	3.4	0	2	K3	+	175	0.61

All data reported in this table were used for subtomogram analysis and were deposited as EMPIAR-10678. The K2 and K3 raw data were collected in super-resolution mode, with ½ the pixel size reported in the table. Pixel size therefore refers to the camera's "bin ×1" pixel. * Refined defocus (Δf) values are reported for defocus phase-contrast data while nominal defoci are reported for Volta phase-contrast (VPP) data. Δtilt = tilt increment. Fig = figures that show this dataset; those without a figure number were used for subtomogram averaging. Camera (Cam): FII = Falcon II, K2 = K2-GIF, K3 = K3-GIF. thick = thickness, measured from the reconstructed tomogram. For cryosections, the thickness is variable due to the presence of crevasses. resid = alignment residual, in nanometers.

Table S7. Total nucleosome-like particles analyzed

Sample	Contrast	Tomograms	Particles
Lysates (BY4741)	Defocus	9	49,413
Lysates (LGY0015)	Defocus	6	96,979
Lysates (LGY0016)	Defocus	4	88,896
Lysates (LGY0002)	Defocus	5	75,960
Lysates (LGY0007)	Defocus	10	93,104
Lysates (LGY0070)	Defocus	8	89,795
Lysates (LGY0071)	Defocus	10	108,754
Cryolamella (BY4741)	Defocus	3	62,428
Cryolamella (BY4741)	Volta	5	129,473
Cryolamella (LGY0016)	Defocus	4	76,169
Cryolamella (LGY0016)	Volta	3	92,989
Cryosections (LGY0016)	Volta	4	33,194
Cryolamella (LGY0007)	Volta	4	166,582

Table S8. Nucleus volume sampled for subtomogram analysis

Tomogram	Volume (nm^3)
20211117_004	203,763,976
20211117_009	126,593,404
20211117_037	37,130,529
20211117_070	193,990,991
20211117_075	106,392,372
Total	667,871,272

SUPPLEMENTAL REFERENCES

Ben-Shem, A., N. Garreau de Loubresse, S. Melnikov, L. Jenner, G. Yusupova, and M.
    Yusupov. 2011. The structure of the eukaryotic ribosome at 3.0 A resolution. *Science*.
    334:1524-1529.

Bharat, T.A., C.J. Russo, J. Lowe, L.A. Passmore, and S.H. Scheres. 2015. Advances in Single-
    Particle Electron Cryomicroscopy Structure Determination applied to Sub-tomogram
    Averaging. *Structure*. 23:1743-1753.

Cai, S., Y. Song, C. Chen, J. Shi, and L. Gan. 2018. Natural chromatin is heterogeneous and
    self-associates in vitro. *Mol Biol Cell*. 29:1652-1663.

Hagen, W.J.H., W. Wan, and J.A.G. Briggs. 2017. Implementation of a cryo-electron
    tomography tilt-scheme optimized for high resolution subtomogram averaging. *J Struct
    Biol*. 197:191-198.

White, C.L., R.K. Suto, and K. Luger. 2001. Structure of the yeast nucleosome core particle
    reveals fundamental changes in internucleosome interactions. *EMBO J*. 20:5207-5218.

Yang, F., L.G. Moss, and G.N. Phillips, Jr. 1996. The molecular structure of green fluorescent
    protein. *Nat Biotechnol*. 14:1246-1251.