# Heterogeneous but not random: Cargo degradation in phagosomes is kinetically coupled to their intracellular mobility

Yanqi Yu[1]†, Zihan Zhang[1]†, Glenn F. W. Walpole[2,3], and Yan Yu[1]*

[1]Department of Chemistry, Indiana University, Bloomington, IN 47405-7102

[2]Program in Cell Biology, The Hospital for Sick Children, Toronto, Ontario, Canada

[3]Department of Biochemistry, University of Toronto, Ontario, Canada

† Y.-q. Y. and Z. Z. contributed equally to this work

*Corresponding Author: Yan Yu, Department of Chemistry, Indiana University, Bloomington, IN 47405-7102, phone 812-855-0593 Email, yy33@iu.edu

**Abstract**

Immune cells degrade internalized pathogens in phagosomes through sequential biochemical changes. The degradation must be fast enough for effective infection control. The presumption is that each phagosome degrades cargos autonomously with a distinct but stochastic kinetic rate. Here we report that the degradation kinetics of individual phagosomes is not stochastic but coupled to their intracellular motility. By engineering RotSensors that are optically anisotropic, magnetic responsive, and fluorogenic in response to degradation activities in phagosomes, we monitored cargo degradation kinetics in single phagosomes simultaneously with their translational and rotational dynamics. We show that phagosomes that move faster centripetally are more likely to encounter and fuse with lysosomes, thereby acidifying faster and degrading cargos more efficiently. The degradation rates increase nearly linearly with the translational and rotational velocities of

phagosomes. Our results indicate that the centripetal motion of phagosomes functions as a clock

for controlling the progression of cargo degradation.

**Introduction**

Innate immune cells, such as macrophages, ingest pathogens and dead cells via a phagocytosis

process. This is an essential process in infection control and tissue homeostasis. After recognizing

ligands on pathogens, immune cells internalize them into phagosomes and transport them from the

cell periphery to the perinuclear region[1, 2, 3, 4, 5]. During this intracellular transport, phagosomes

degrade the entrapped pathogens through sequential biochemical transformations, including the

acidification of the lumen[6], the activation of hydrolytic enzymes[7], and the generation of reactive

oxygen species (ROS)[8]. Phagosomes must degrade pathogens efficiently to prevent them from

escaping. Many pathogens, such as the bacterium *Legionella pneumophila*, were found to evade the

degradative environment in phagosomes by hijacking the intracellular trafficking process[9, 10, 11].

These examples highlight that the degradative function of phagosomes is tightly associated with

their dynamics in cells. Understanding the relationship between the degradative function and

dynamics of phagosomes is critical for developing therapeutics against infections and

inflammation. It is, however, challenging due to the autonomous nature of the two processes. Each

phagosome degrades its cargo at its own distinct kinetic rate, which was assumed to be stochastic.

The velocity at which phagosomes are transported along microtubules is also heterogeneous and

believed to be stochastic[2, 12]. Therefore, those dynamic and biochemical changes must be quantified

at the single phagosome level to reveal their relationships, but there is currently no method to do

so. The dynamics of phagosomes in cells has been extensively studied using single-particle

imaging[13, 14]. The degradative activities, including lumen acidification and protein digestion, in

phagosomes have been studied separately in population-based measurements[15]. In those ensemble-

average studies, the individuality of phagosomes was overlooked. Regarding the relationship

between the degradative function and dynamics of phagosomes, studies have relied on approaches that inhibit phagosome dynamics by disrupting the cytoskeleton or molecular motor functions. The disruption of actin was shown to attenuate phagosome-endosome fusion[16, 17, 18]. Microtubule depolymerization was found to disrupt: i) phagosome-lysosome fusion[19], ii) content delivery from late endosomes to phagosomes[20], and iii) acquisition of lysosome markers[21]. These observations demonstrated that dynamics of phagosomes in cells is important for their degradative function, but none of those studies identified the quantitative relationship between phagosome dynamics and kinetics of the biochemical activities during cargo degradation.

In this study, we develop a single-phagosome imaging toolset for monitoring both degradative function and dynamics of individual phagosomes. The RotSensor particles are optically anisotropic and fluorogenic in response to acidification in phagosomes and phagosome-lysosome fusion, two key degradative activities. Using the RotSensors as cargos in phagosomes, we imaged simultaneously the biochemical changes during phagosome degradation and their translational and rotational dynamics in living cells. By combining this toolset with magnetic tweezers to control the dynamics of phagosomes, we obtained direct evidence that the microtubule-based centripetal transport of phagosomes determines the kinetic rates of their degradation process. More mobile phagosomes fuse with lysosomes and acidify in their lumen more rapidly, following a nearly linear relationship. Our results reveal the orchestration of the dynamical and biochemical processes that occur during pathogen degradation in phagosomes, providing a possible mechanism by which some pathogens may evade immune clearance by disrupting the intracellular dynamics of residing phagosomes.

**Results**

**pH-responsive rotational particle sensors (pH-RotSensors) for measuring phagosome**

**acidification and dynamics**

The acidification of phagosome lumen is an early indicator of its degradative function and required

for many subsequent events, including the activation of degradative enzymes[22, 23]. Therefore, we

first designed pH-responsive RotSensors to simultaneously measure the acidification of single

phagosomes and their intracellular transport dynamics, including both translational and rotational

movements (Fig. 1a). Each RotSensor consists of a pH-responsive particle (1 μm diameter)

covalently conjugated with a smaller green particle (100 nm diameter) with non-overlapping

fluorescence emission (Fig. 1b). Once engulfed into phagosomes, the "snowman"-like RotSensors

report both the translational and the rotational motion of the residing phagosome. The azimuthal

(φ) and polar (θ) angles of each RotSensor can be measured, respectively, from the orientation and

projection distance between the pair of the two particles, as previously described (Fig. 1b)[24].

To monitor phagosome acidification, the 1 μm particle in each RotSensor was biotinylated and

subsequently conjugated with two types of streptavidin, one type labeled with the pH-indicator

pHrodo Red and the other with the reference dye CF640R. The dye pHrodo Red fluoresces intensely

in an acidic environment, making it an ideal indicator of phagosome acidification. CF640R was

chosen as a reference dye because it is pH insensitive and photostable (Supplementary Fig. 1).

Streptavidin is critical in the RotSensor design because it acts as a cushion layer to separate the

pHrodo Red dye from the particle surface. Without the streptavidin linker, pHrodo Red that was

directly conjugated onto the particle surface exhibited little pH sensitivity. The pH-sensitive

RotSensors were then coated with immunoglobulin G (IgG) via non-specific adsorption to initiate

Fc receptor (FcR)-dependent phagocytosis in RAW264.7 macrophage cells.

To quantify the pH response of the RotSensors, we measured the ratio of fluorescence emission

intensities of pHrodo Red and of the reference dye CF640R; $I_{pHrodo}/I_{ref}$, as we varied pH inside

phagosomes in live cells and in aqueous buffers (Supplementary Fig. 2). For both cases, $I_{pHrodo}/I_{ref}$

of the RotSensors increased linearly with decreasing pH, consistent with previous reports[25]. The pH

responses of individual RotSensors, including the initial $I_{pHrodo}/I_{ref}$ ratio and slope, varied slightly

from one to another, because of the different amounts of dyes on particles and the slightly different

phagosome lumen environments[26, 27, 28, 29]. To eliminate the effect of such variation in pH

measurements, we calibrated our pH measurements for each phagosome. We measured

$I_{pHrodo}/I_{ref}$ of each phagosome versus time and then converted the intensity ratio to pH based on

$I_{pHrodo}/I_{ref}$ of the same phagosome at pH 4.5 and 7.3 (details in Methods section).

**pH-RotSensors reveal correlation between phagosome acidification rate and transport**

**dynamics**

Before tracking the dynamics of phagosomes, we first asked whether the rotation of RotSensors

faithfully reports the rotation of their residing phagosomes. Transmission electron microscopy

(TEM) images show that ≈ 86% of internalized RotSensors remained tightly wrapped by the

phagosome membrane at the end of our experiments (50 min after particles were added to cells)

(Supplementary Fig. 3). Based on the tight membrane fitting and the presence of ligand-receptor

binding between the RotSensors and phagosome membranes, the RotSensors are unlikely to

freely rotate inside phagosomes. Thus, their rotation is expected to report the rotation of their

residing phagosomes.

We then simultaneously imaged the pH response and the rotational and translational dynamics of

single phagosomes encapsulating RotSensors (Fig. 1c). Phagosomes acidified in three steps: 1) an

initial standby period during which the phagosome pH remained at neutral pH of ≈ 7.3 (Fig. 1d),

2) a rapid acidification period when the pH quickly dropped over a period of a few minutes, and 3) an eventually plateau at pH 4.5-5.0. This acidic pH is a prerequisite for activating degradative enzymes[8, 30]. While individual phagosomes reached slightly different final pH values, their acidification kinetics mostly (31 of 42 phagosomes) followed a sigmoidal-Boltzmann function:

$$pH = pH_{final} + \frac{pH_{initial} - pH_{final}}{1 + exp\left(\frac{t - t_0}{dt}\right)} \qquad (1)$$

in which $t_0$ is the half-response point, and $t_{initial}$ and $t_{final}$ denote the beginning and the end of rapid acidification, respectively (Fig. 1d). Slope at $t_0$ is the acidification rate by definition. We found that phagosomes acidified at an average rate of 0.33 ± 0.28 pH unit/min (N = 42). The acidification of phagosomes requires intact microtubules, as the acidification was abolished upon microtubule disruption by nocodazole (Supplementary Fig. 4).

For transport dynamics, we observed that a majority (~70%) of phagosomes exhibited a slow-to-fast movement transition both translationally and rotationally (Fig. 1e and f). The remaining 30% of phagosomes exhibited only slow movements. This initially slow movements are not dependent on microtubules, as the average velocities were not affected by microtubule depolymerization (Supplementary Fig. 5). After a few minutes of slow movements, phagosomes moved rapidly centripetally with velocities as high as 0.1 µm/s (Fig. 1e). This range of phagosome velocities is consistent with literature reports for 1 µm bead-containing phagosomes in macrophage cells[13]. Meanwhile, phagosomes also rotated frequently in between segments of translational motion with rotational velocities up to 0.15 rad/s (Fig. 1f). These dynamic behaviors, particularly phagosome rotation, are characteristic of cargo transport driven by microtubule-based molecular motors, as we previously reported[31]. This was further confirmed in control experiments, in which the rapid translational and rotational dynamics disappeared in cells treated with nocodazole (Supplementary Fig. 4).

Notably, the beginning of the rapid acidification period appeared to coincide with the onset of fast

microtubule-based transport (Fig. 1e and f). To explore the possible correlation between the two

processes, we calculated the acidification rate of single phagosomes based on the sigmoidal fitting

of their individual pH-time plots (Fig. 1d). We then plotted the acidification rate of single

phagosomes against their individual translational and rotational velocities during the rapid

acidification phase (from $t_{initial}$ to $t_{final}$) (Fig. 1g and h). The acidification rate varied broadly

among phagosomes, even those from the same cells. This reflected the individuality of each

phagosome in degradation. Despite this heterogeneity, the acidification rates of individual

phagosomes follow a linear relationship with both their translational and rotational velocities with

a Pearson's coefficient of 0.78 and 0.81, respectively. Those observations apply not only to

phagosomes from different cells but also to those from the same cells. This is shown in Fig. 1g and

h, in which data of phagosomes from the same cells are the same color. This is also a general

phenomenon independent of the activation status of macrophage cells (Supplementary Fig. 6).

Further, the acidification rates of single phagosomes also follow a linear relationship with the

percentage, maximum amplitude, and frequency of active rotation of phagosomes during the rapid

acidification phase (from $t_{initial}$ to $t_{final}$) with a Pearson's coefficient of 0.76, 0.60 and 0.61,

respectively (Fig. 1i and j, and Supplementary Fig. 7). For this analysis, active rotation of

phagosomes was distinguished from passive random rotation by using a Wavelet analysis method

reported previously (Supplementary Fig. 8; Methods)[32]. The percentage of active rotation was

defined as the total time during which the phagosome underwent active rotation divided by the time

duration of its rapid acidification (from $t_{initial}$ to $t_{final}$). The results altogether indicate that more

mobile phagosomes acidify faster. As an additional evidence supporting this conclusion, we also

observed that the ~30% of phagosomes that never underwent rapid microtubule-based transport

exhibited minimal acidification (Supplementary Fig. 19a-c). Surprisingly however, the final pH

attained by individual phagosomes showed no correlation with either the translational or the

rotational velocities of those phagosomes (Supplementary Fig. 10). Because the final acidic pH

determines whether a phagosome can successfully degrade its cargo, this result suggests that the

degradative capacity of a phagosome is likely determined prior to its transport on microtubules, but

that the rate of degradation is correlated with the transport dynamics.

**FRET-RotSensors for measuring phagosome-lysosome fusion**

During cargo degradation, phagosome acidification relies on its continuous fusion with lysosomes,

which deliver proton-pumping vacuolar H+-ATPase (V-ATPase) to it[28, 33, 34, 35, 36]. Since we found

that more mobile phagosomes acidify more rapidly, we asked whether phagosome-lysosome fusion

is likewise correlated with phagosome mobility. To do this, we measured the kinetics of

phagosome-lysosome fusion by modifying the RotSensor design into a Förster Resonance Energy

Transfer (FRET)-based assay. This fluorescence assay can be used to determine when two

fluorophores, a donor inside lysosomes and an acceptor inside phagosomes, are in close proximity

to one another. FRET-sensitive RotSensors were prepared in the same way as pH-sensitive

RotSensors, but, in this case, the sensor particles were labeled with streptavidin that was conjugated

with the donor fluorophore Alexa568 (Fig. 2a). For FRET, lysosomes were loaded with fluid-phase

biotinylated bovine serum albumin (BSA) that was conjugated with the acceptor fluorophore

Alexa647 using a pulse–chase procedure[37]. We confirmed that the BSA-Alexa647-biotin was

contained inside lysosomes based on its colocalization with the lysosome marker LysoTracker

inside cells (Supplementary Fig. 11). Inside lysosomes, BSA-Alexa647-biotin was fragmented by

proteases before RotSensors were added to cells for internalization (Supplementary Fig. 12). The

working principle of the FRET assay is that once the contents of lysosomes are intermixed with the

phagosome lumen during fusion, the fragmented BSA-Alexa647-biotin from lysosomes will in

close proximity with Alexa568 conjugated RotSensors leading to FRET. The magnitude of the

FRET signal thus indicates the extent of phagosome-lysosome fusion. This assay doesn't

necessarily require biotin-streptavidin binding. We showed in *in vitro* experiments that FRET can

occur when BSA-Alexa647, without biotinylation, was added to FRET-RotSensors in 1xPBS

solution (Supplementary Fig. 13). However, we still used BSA-Alexa647-biotin in this FRET assay

because it was loaded into lysosomes much more efficiently than BSA-Alexa647 alone, which

improved the sensitivity of the FRET assay (Supplementary Fig. 14). The observation that the biotin

moiety promotes uptake of BSA by cells is consistent with previous reports[38, 39].

For a thorough validation of the FRET assay, we first tested the FRET-RotSensors *in vitro* by

mixing them with BSA-Alexa647 in 1xPBS (Supplementary Fig. 13). In this experiment, we could

not measure the acceptor emission in the FRET channel due to the excessive amount of free BSA-

Alexa647 in solution. Instead, we quantified the percentage of donor quenching as an indirect

indicator of FRET efficiency. We found that the percentage of donor quenching, first increased

linearly as a function of BSA-Alexa647 concentration in the lysosome loading solution ([BSA-

Alexa647]$_{loading}$), but reached a plateau at [BSA-Alexa647]$_{loading}$ > 200 µg/ml. This plateau indicates

the maximum FRET efficiency which is determined by the amount of the donor fluorophore

(Alexa568) available on RotSensors. For the FRET assay to accurately detect continuous fusion of

a phagosome with lysosomes, the concentration of acceptor fluorophores (Alexa647) loaded into

lysosomes must be optimized to achieve sufficient FRET efficiency without saturating the donor

on the RotSensors. We did this by optimizing the loading concentration of BSA-Alexa647-biotin

into lysosomes.

We next tested and optimized the FRET assay in live cells. To start with, we loaded lysosomes

using 10 µg/mL BSA-Alexa647-biotin ([BSA-Alexa647-biotin]$_{loading}$ = 10 µg/mL). After

internalization of FRET-RotSensors, lysosomes docked on the phagosomes, and the FRET-

RotSensors inside phagosomes exhibited decreased fluorescence emission in the donor channel

(ex/em: 561/586 nm) and simultaneously increased emission in FRET channel (ex/em: 561/680

nm) (Fig. 2b and c). For quantification, we calculated FRET ratio defined as:

$$FRET\ ratio = \frac{FRET_{em}}{AF568_{em}} = \frac{FRET_0 - \alpha \times AF568_{em} - \beta \times AF647_{em}}{AF568_{em}} \tag{2}$$

In this equation, $FRET_0$ is the measured FRET signal (ex/em 561/680 nm) before correction for

spectral bleed-through, $AF568_{em}$ is the emission in the donor channel (ex/em 561/586 nm), and

$AF647_{em}$ is the emission in the acceptor channel (ex/em 660/680 nm). The coefficient $\alpha$ corrects

the bleed-through contribution from the donor (Alexa568) due to the overlap of the donor emission

in the FRET channel. The coefficient $\beta$ corrects the non-FRET bleed-through from the acceptor

(Alexa647) due to the excitation of the acceptor at the donor excitation wavelength. Correction

factors $\alpha$ and $\beta$ were determined independently for each experiment (see Methods). Simultaneous

with the observation of lysosome docking on phagosomes, the FRET ratio of single phagosomes

increased continuously with time until it reached a plateau, following a sigmoidal relationship (Fig.

2d). This is a general phenomenon confirmed in over 40 phagosomes from 22 cells (Supplementary

Fig. 15a-c). The gradual increase in FRET ratio as a function of time indicates an increasing

concentration of acceptors (Alexa647) in phagosomes resulted from their continuous fusion with

lysosomes. FRET signal was diminished when microtubules were disrupted by nocodazole

treatment, which further confirms that this assay measures phagosome-lysosome fusion

(Supplementary Fig. 16).

We then asked: what caused the plateau in the FRET ratio of single phagosomes? This is unlikely

to be caused by the completion of phagosome-lysosome fusion, because lysosomes were seen to

dock on phagosomes after the FRET ratio reached its plateau. We explored two possible causes for

the plateau seen in the FRET ratio vs. time plots. One is that the concentration of acceptor

fluorophores delivered into phagosomes by lysosome fusion reached the threshold needed to

saturate the donor fluorophores on RotSensors. The second is that the acceptor concentration in the

phagosome had reached equilibrium with that in the lysosomes that were fusing with it. Even as

more lysosomes continued to fuse with the phagosome, the content exchange would no longer

increase the concentration of acceptors in the phagosome. To determine which of these two possible

circumstances caused the FRET ratio to plateau, we varied [BSA-Alexa647-biotin]$_{loading}$ from 2.5

µg/mL to 25 µg/mL (note: concentrations higher than 25 µg/mL resulted in decreased cell viability).

While the exact concentration of acceptors (Alexa647) inside lysosomes is unknown, it is

proportional to [BSA-Alexa647-biotin]$_{loading}$ (Supplementary Fig. 14 and Supplementary Fig. 15a-

c). The rationale here is that if the plateau in FRET ratio was caused by the saturation of donors on

RotSensors, the magnitude of plateau would be determined by the amount of donors on RotSensors

and thus should not change with [BSA-Alexa647-biotin]$_{loading}$. On the contrary, if the plateau of

FRET ratio occurred because the acceptor concentration reached an equilibrium between the

phagosome and fusing lysosomes, the magnitude of plateau would be determined by the acceptor

concentration inside lysosomes and thus should increase with [BSA-Alexa647-biotin]$_{loading}$. In

experiments, we found that the plateau level of FRET ratio increased with [BSA-Alexa647-

biotin]$_{loading}$ from 2.5 to 25 µg/mL (Supplementary Fig. 15a-d). This indicates that the FRET ratio

plateau was due to the establishment of equilibrium between the phagosome lumen and lysosomes,

and not saturation of donors on RotSensors.

We further optimized [BSA-Alexa647-biotin]$_{loading}$ to minimize cell-to-cell variation in loading

BSA-Alexa647-biotin into lysosomes. We found that the higher [BSA-Alexa647-biotin]$_{loading}$, the

more variation in the amount of BSA-Alexa647-biotin loaded into lysosomes between cells

(Supplementary Fig. 14). As a result, the plateau level of FRET ratio had larger variations at [BSA-

Alexa647-biotin]$_{loading}$ > 10 µg/mL (Supplementary Fig. 15d). To estimate how the cell-to-cell

variation affects the measurement of phagosome-lysosome fusion kinetics, we normalized the

FRET ratio vs. time plot of each phagosome to its plateau level and then from the sigmoidal

Boltzmann fitting of each FRET ratio plot, we calculated the slope at half-response point $t_0$ as the

kinetic rate of phagosome-lysosome fusion (referred to as FRET rate) (Fig. 2d and Supplementary

Fig. 15e). In the FRET rate vs. [BSA-Alexa647-biotin]$_{loading}$ plot (Supplementary Fig. 15f), the

standard deviation of FRET rate between phagosomes increased significantly at [BSA-Alexa647-

biotin]$_{loading}$ > 10 µg/ml. All results suggest that an optimal range of [BSA-Alexa647-biotin]$_{loading}$

for the FRET assay is ≤ 10 µg/ml. We therefore chose to use 10 µg/ml, the highest concentration

within this range, to improve the sensitivity of the FRET assay.

**284  FRET-RotSensors reveal correlation between phagosome-lysosome fusion kinetics and**

**285  phagosomes centripetal transport**

After the validation and calibration of the FRET fusion assay, we then investigated whether the

kinetics of phagosome-lysosome fusion (measured as the FRET rate) correlates with the motility of

phagosomes. Much like the heterogeneity in acidification, the FRET rate differed greatly between

phagosomes even within the same cell (Fig. 2e and f). However, all single phagosome FRET ratio

data follow a linear correlation with the translational and rotational velocities of phagosomes, with

a Pearson's coefficient of 0.75 and 0.86, respectively (Fig. 2e and f). This indicates that phagosomes

that moved faster translationally and rotated more along microtubules also fused more rapidly with

lysosomes.

Because lysosomes in macrophages are more concentrated in the perinuclear region (Fig. 3a)[40], we

then asked if, besides velocity, the direction of phagosome transport plays any role in their fusion

with lysosomes. To identify centripetal towards cell nucleus, we analyzed the distance of each

phagosome from the center of cell nucleus as a function of time (Fig. 3b). The effective transport

distance of a phagosome (grey shaded areas in Fig.3b) is the distance from the time when it begins

to fuse with lysosomes to the time when it reaches the nucleus boundary. The starting time of phagosome-lysosome fusion was obtained from the sigmoidal fitting of the FRET ratio vs. time plot (Fig. 2d). We also identified segments of active centripetal motion following a previously reported method (Fig. 3b)[13]. Most phagosomes moved centripetally (Supplementary Fig. 17), but not all of them exhibited active centripetal runs (Fig. 3b). As shown in Fig. 3A, phagosome #1 underwent a few segments of active centripetal runs, but phagosome #2 did not. Concurrently, phagosome #1 also fused with lysosomes more rapidly (Fig. 3c). To confirm the generality of this result, we quantified the centripetal velocity and percentage of active centripetal runs of single phagosomes. Centripetal velocity was calculated as the effective transport distance divided by time. The percentage of active centripetal run was calculated as the sum of time of all active centripetal runs divided by the total duration of the effective transport (N = 40 phagosomes from 22 cells). The results demonstrate that phagosomes with larger centripetal velocity and higher percentage of active centripetal movements fuse faster with lysosomes (Fig. 3d).

**MagSensors for magnetic manipulation and imaging of phagosomes**

To determine whether the centripetal motility of phagosomes determines their fusion with lysosomes and consequently their acidification, we applied magnetic tweezers to manipulate the intracellular transport of single phagosomes. We simultaneously measured changes in phagosome degradation (Supplementary Fig. 18). We designed 1 µm magnetically modulated phagosome sensors (MagSensors) that were also pH-responsive (Fig. 4a). The MagSensors were biotinylated and conjugated with SAv-pHrodo Red (pH indicator) and SAv-CF640 (reference), using the same procedure as for the preparation of RotSensors. The MagSensors exhibited similar pH responses (Supplementary Fig. 19). In all experiments, a solenoid tip was positioned on the opposite side of the cell from the phagosome of interest, so that the phagosome could be pulled by magnetic attraction from the cell periphery towards the center. The magnetic force exerted on individual

MagSensors was calibrated by tracking the movement of particles in a polydimethylsiloxane (PDMS) base calibration medium under magnetic pulling force. The magnetic force changes with the distance between the MagSensor and the solenoid tip following theoretical predictions (Fig. 4b)[41]. On average, the solenoid tip was positioned ≈ 43 μm from the MagSensor of interest. This means that the average force exerted on a MagSensor by the tip was ≈ 21 pN at the beginning of manipulation and reached as high as ≈ 31 pN as it moved closer to the tip at the end of imaging (Supplementary Fig. 20).

We performed the magnetic experiments in resting macrophages stably expressing actin-GFP (Supplementary Fig. 21). The fluorescence from actin-GFP allowed us to identify when phagosome formation completes[42, 43], so that we can apply magnetic pulling force on the MagSensor-containing phagosomes only after their internalization. We used resting cells for these experiments because they are less flat, which allows for better magnetic manipulation of the phagosome. The acidification-motility correlation of phagosomes in resting cells was similar to that in activated cells (Supplementary Fig. 6). In the experiments, we turned on the magnetic force immediately after the MagSensors were internalized into phagosomes, which was indicated by the actin-GFP intensity peak, and pulled the MagSensor-containing phagosomes from the cell periphery towards the center (Fig. 4a). The magnetic force remained on throughout imaging. Under the magnetic pulling force, phagosomes moved towards the magnetic tip with accelerated translational velocity and larger mean square displacements (Fig. 4d and Supplementary Fig. 22). Their trajectories were zigzagged (Fig. 4c), likely because the magnetic force applied on the MagSensors was only slightly larger than the collective forces from the microtubule-based molecular motors. As mentioned above, the magnetic pulling force on single MagSensors was ≈ 21 to 31 pN (Supplementary Fig. 20). For comparison, microtubule-based molecular motors were shown to exert collective forces as high as ≈ 20 pN on vacuoles encapsulating 1 μm particles[44]. Therefore, the zigzagged movements of

phagosomes are plausibly a result of the combined influence of the magnetic force and the forces exerted by molecular motors. As the translational velocity of phagosomes accelerated under magnetic pulling, they also acidified more rapidly (Fig. 5a). Phagosomes that were magnetically pulled acidified at an average rate of $0.79 \pm 0.54$ pH unit/min (N = 38). In contrast, for phagosomes not subject to magnetic pulling, an average rate of $0.48 \pm 0.34$ pH unit/min was observed (N = 33) (Fig. 5b). Surprisingly, magnetic manipulation had no effect on the final pH of the phagosomes, as phagosomes with or without magnetic manipulation reached an average final pH of $4.7 \pm 0.4$ and $4.7 \pm 0.3$, respectively (Fig. 5c). These results evidently demonstrate that faster transport of phagosomes causes their faster acidification.

We next investigated how magnetic pulling affects phagosome-lysosome fusion using the FRET fusion assay shown in Fig. 2. BSA-Alexa647-biotin (10 µg/ml) was used to load lysosomes. As shown in Fig. 5c, the FRET signals from magnetically pulled phagosomes (average from N = 20 each) were noticeably more intense and increased more rapidly, indicating enhanced phagosome fusion with lysosomes. The FRET rate, which indicates the phagosome-lysosome fusion kinetic rate, is $0.28 \pm 0.21$ a.u. (N = 20) for phagosomes under magnetic pulling, in contrast to the average rate of $0.15 \pm 0.07$ a.u. without magnetic forces (N = 20) (Fig. 5e). To investigate whether the movement of phagosomes alone is sufficient to facilitate productive fusion with lysosomes, we next treated cells with nocodazole, which disrupts microtubule-based transport of both phagosomes and lysosomes. We then magnetically moved phagosomes the same way as in non-treated cells. Despite the fast movement of phagosomes under magnetic pulling forces, they exhibited minimal fusion with lysosomes and minimal acidification (Supplementary Fig. 23). This suggests that productive

phagosome-lysosome fusion requires active transport of not only the phagosomes but also the lysosomes that they fuse with.

The magnetic tweezers results, taken together, demonstrate that faster transport of phagosomes on microtubules promotes their encounters with lysosomes, which leads to faster acidification. This explains why phagosome transport velocity positively correlates with phagosome-lysosome fusion kinetics and acidification rate.

**Discussion**

Phagosomes, even those from the same cells, show a significant amount of variation from one another in their degradation and transport. Each phagosome functions as a separate degradative unit with its own distinct degradation kinetics and transport velocity. However, the individuality of phagosomes has been overlooked in previous studies that relied on population-based measurements of phagosomes[8, 30, 45, 46]. There is no known quantitative relationship between the seemingly stochastic rates of degradation of single phagosomes and their equally stochastic transport dynamics. In the current study, we detail the development of a single-phagosome imaging and manipulation toolset that allowed us to probe both degradative processes and dynamics of individual phagosomes. Multimodal particle sensors, called RotSensors, were engineered as phagosome probes to measure acidification within the lumen of an individual phagosome, the kinetics of that same phagosome's fusion with lysosomes, and its overall rotational and translational motion within the cell. The RotSensors were further combined with magnetic tweezers to perturb phagosome transport and observe the consequences for the degradation of single phagosomes. This integrated approach allowed us to demonstrate that the biochemical progression of degradative functions in individual phagosomes is coupled to and determined by the dynamics of their active transport.

By simultaneously tracking the transport dynamics and degradation of single phagosomes, we first

determined that phagosomes that move faster centripetally fuse with lysosomes and acidify faster.

While phagosome degradation is known to require microtubule-dependent transport[19, 21, 33, 47, 48, 49,

50, 51], our results demonstrate that the seemingly stochastic rate of degradation of single phagosomes

is positively correlated with their equally stochastic velocity of transport. In the magnetic tweezers

experiments, we then showed that when phagosomes are accelerated towards nucleus, they speed

up their fusion with lysosomes and their acidification. The velocity of phagosomes directly

regulates where and how fast they fuse with lysosomes, and this, in turn, determines the

acidification. Surprisingly however, phagosome velocity doesn't influence the final pH of the

phagosomes. Our results suggest that centripetal transport of phagosomes functions as a clock that

determines the timing of their sequential biochemical activities during pathogen degradation. The

precise positioning of phagosomes largely relies on microtubules-based motors[52]. It was shown

previously that Rab7-RILP (Rab7-interacting lysosomal protein) complex induces the recruitment

of the dynein-dynactin motor protein complex to phagosomes[53], and the clustering of dynein

generates extra force to mediate the rapid centripetal transport of phagosomes towards the

perinuclear region[50]. Because lysosomes are enriched in perinuclear region (Fig. 3a) and the

perinuclear lysosomes are more acidic than those near cell periphery[54, 55] , it is possible that dynein-

mediated active centripetal transport enhances the probability of physical collision between a

phagosome and lysosomes, hence promoting the degradation of phagosome lumen. We further

confirmed that it is the active microtubule-based centripetal transport, rather than simple diffusion,

regulates phagosome degradation. Because phagosomes failed to fuse with lysosomes or to acidify

in cells with disrupted microtubules, even if phagosomes were moved magnetically. Moreover,

Rab7 not only bridges phagosomes with dynein-dynactin, but also directly recruits the homotypic

fusion and vacuole protein sorting (HOPS) complex and V-ATPase subunit V1G1, which are

involved in membrane fusion and phagosome acidification[56, 57]. Those literature reports, albeit scattered, suggest that phagosome membrane associated molecules, such as Rab7-RILP complexes, may function as the molecular links that relate phagosome motility with their degradation activities.

In addition to studying phagosomes, we envision that the RotSensor toolset presented here could be further applied, with appropriate modifications, to studying intracellular pathogens and endosome degradation. Many pathogens, such as the bacterium *Legionella pneumophila*, hijack the phagosome-lysosome fusion process to evade immune clearance[9, 10, 11]. Our study here was done with synthetic beads, which are far less complex than pathogens. Nevertheless, the results imply that intracellular pathogens might prolong their survival in host cells by disrupting the intracellular transport of phagosomes to slow down fusion with lysosomes and thus delay the degradation. This hypothesis can be tested in future studies where pathogen-extracted bioparticles or intact pathogens, instead of synthetic particles, can be used in the single-particle assays. Similarly, when smaller nanoparticles are used to replace the micron-sized beads in the single-particle assays, it will be possible to study the degradative function of endosomes that are smaller than phagosomes.

**Methods**

**Materials**

Carboxylate-modified yellow-green fluorescent polystyrene nanoparticles (diameter 100 nm), carboxylate-modified superparamagnetic Dynabeads (diameter 1 μm), pentylamine-biotin, Alexa Fluor 568 NHS ester (succinimidyl ester), Alexa Fluor 647 NHS ester (succinimidyl ester), pHrodo iFL Red STP Ester, Streptavidin, LysoTracker Green DND-26, Alexa Fluor 488 anti-tubulin-α antibody, Alexa Fluor Plus 647 phalloidin, and DQ-green BSA were purchased from ThermoFisher (Waltham, MA). Immunoglobulin G from rabbit plasma, albumin from bovine serum (BSA), biotin N-hydroxysuccinimide ester (biotin-NHS), nocodazole, and lipopolysaccharides were purchased

from Sigma-Aldrich (St. Louis, MO). Monodisperse amine-modified silica particles (diameter 1.0 µm) were purchased from Spherotech Inc. (Lake Forest, IL). Sylgard 184 PDMS base was from Dow Corning (Midland, MI). 1-(3-Dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) was purchased from Alfa Aesar (Haverhill, MA). Nigericin sodium salt was purchased from Tocris Bioscience (Minneapolis, MN). Recombinant Murine IFN-γ was purchased from Peprotech (Rocky Hill, NJ). CF640R-amine was purchased from Biotium (Fremont, CA). RAW264.7 macrophage was purchased from ATCC (Manassas, VA). RAW264.7 macrophages stably expressing EGFP-actin have been previously described [42]. Ringer's solution (pH = 7.3, 10 mM HEPES, 10 mM glucose, 155 mM NaCl, 2 mM $NaH_2PO_4$, 5 mM KCl, 2 mM $CaCl_2$, 1 mM $MgCl_2$) was used for live-cell imaging. Potassium-rich solution (135 mM KCl, 2 mM $K_2HPO_2$, 1.2 mM $CaCl_2$, 0.8 mM $MgSO_4$) was used for intracellular pH calibration. Acidic washing solution (135 mM KCl, 2 mM $K_2HPO_2$, 1.2 mM $CaCl_2$, 0.8 mM $MgSO_4$, 5 mM sodium citrate) at pH of 4.5 was used for particle washing in the EDC coupling step of the RotSensor fabrication. Artificial lysosome fluid (55 mM NaCl, 0.5 mM $Na_2HPO_4$, 0.26 mM trisodium citrate dihydrate, 0.79 mM glycine, 150 mM NaOH, 108 mM citric acid, 0.87 mM $CaCl_2 \cdot 2H_2O$, 0.27 mM $Na_2SO_4$, 0.25 mM $MgCl_2 \cdot 6H_2O$, 0.46 mM disodium tartrate, 1.6 mM sodium pyruvate) was prepared following a previously reported protocol [58] and used for washing particles during the protein conjugation step of the RotSensor fabrication.

**Cell Culture, Pharmacological Treatments**

Both RAW264.7 macrophage and stable cell line expressing EGFP-actin were cultured in Dulbecco's Modified Eagle Medium (DMEM) complete medium supplemented with 10% fetal bovine serum (FBS), 100 units/ml penicillin and 100 µg/ml streptomycin at 37°C and 5% $CO_2$. Resting macrophages were activated with a combination of 50 ng/ml LPS and 100 units/ml IFN-γ for 9 h prior to live cell imaging. Microtubule depolymerization was achieved by incubating cells

471   with 10 µM nocodazole for 1 h prior to imaging, and 10 µM nocodazole was presented during live

472   cell imaging.

**Fabrication and Characterization of Phagosome Sensors**

**(a) Fluorescence labeling of Streptavidin and Bovine Serum Albumin (BSA)**

To prepare streptavidin-pHrodo Red conjugates (SAv-pHrodo Red), 0.7 mg of streptavidin, 60 µg

of pHrodo Red STP Ester were mixed in 350 µl $NaHCO_3$ solution (100mM, pH 8.2) for 3 h at room

temperature. Free dyes were removed by centrifugal filtration using Amicon Ultra filters (30K). To

prepare streptavidin-CF640R conjugates (SAv-CF640), 0.7 mg of streptavidin, 80 µg of CF640R-

amine, and 3 mg of EDC were mixed in 350 µl MES buffer (50mM, pH 4.5) for 3 h at room

temperature. Free dyes were removed by centrifugal filtration using Amicon Ultra filters (30K). To

prepare streptavidin-Alexa568 conjugates (SAv-Alexa568), 0.5 mg of streptavidin, 60 µg of Alexa

Fluor 568 NHS ester were mixed in 235 µl $NaHCO_3$ solution (100mM, pH 8.2) for 1 h at room

temperature. Free dyes were removed by centrifugal filtration using Amicon Ultra filters (30K). To

prepare bovine serum albumin-Alexa647-biotin (BSA-Alexa647-biotin) conjugates, a mixture

containing 2 mg/ml BSA and 188 µg/ml (molar ratio of 1: 20) biotin-NHS in $NaHCO_3$ solution

(100 mM, pH 8.2) was incubated at room temperature for 1 h. Unbound biotin-NHS was removed

by centrifugal filtration using Amicon Ultra filters (30K). After washing, a mixture containing 2

mg/ml BSA-biotin and 780 µg/ml Alexa Fluor 647 NHS ester in $NaHCO_3$ solution (100 mM, pH

8.2) was incubated at room temperature for 3 h. Free dyes were removed by centrifugal filtration

using Amicon Ultra filters (30K). The resulting BSA-Alexa647-biotin yielded dye: protein ratios

of 1.2:1 on NanoDrop UV-Vis measurement (ThermoFisher, Waltham, MA). To prepare bovine

serum albumin-Alexa647 (BSA-Alexa647) conjugates, a mixture containing 2 mg/ml BSA and 780

µg/ml Alexa Fluor 647 NHS ester in $NaHCO_3$ solution (100 mM, pH 8.2) was incubated at room

temperature for 1 h. Free dyes were removed by centrifugal filtration using Amicon Ultra filters (30K). The resulting BSA-Alexa647 yielded dye: protein ratios of 1.3:1 based on NanoDrop UV-Vis measurement (ThermoFisher, Waltham, MA).

**(b) Fabrication of Rotational Phagosome Sensors (RotSensors)**

RotSensors: Amine-modified non-fluorescent silica particles (1 µm) were incubated with 150 µg/ml biotin-NHS in $NaHCO_3$ solution (10 mM, pH 8.25) for 2 h at room temperature and then with 500 µg/ml of biotin-NHS for 30 min for a second round of biotinylation. After biotinylation, silica particles were washed in methanol and deionized (DI) water. The "snowman"-like design was introduced by covalently conjugating the carboxylate-modified yellow-green fluorescent nanoparticles (100 nm) to the biotinylated and amine-modified silica particles (1 µm) using EDC coupling. In brief, carboxylate-modified yellow-green fluorescent nanoparticles (100 nm) were mixed with the biotinylated and amine-modified silica particles (1 µm) at a molar ratio of 333:1 in phosphate buffer (10 mM, pH 7.0) containing 1.0 mg/ml EDC for 2 h at room temperature. After the EDC coupling, particles were washed with acidic washing solution and 1×PBS to remove the non-covalently bounded carboxylate-modified yellow-green fluorescent nanoparticles (100 nm). 20% of the particles were found to have the "snowman" shape with 1:1 coupling ratio of the biotinylated and amine-modified silica particles (1 µm) and the carboxylate-modified yellow-green fluorescent nanoparticles (100 nm).

pH-RotSensors: pH-RotSensors were fabricated by labeling RotSensors with SAv-pHrodo Red and SAv-CF640. RotSensors were incubated with 25 µg/ml of SAv-pHrodo Red, 2.5 µg/ml of SAv-CF640, 5 µg/ml of BSA and 1 µg/ml of IgG in 1×PBS for 5 h at room temperature. Unbound proteins were rinsed off with artificial lysosome fluid and 1×PBS. The resulting pH-RotSensors were further opsonized with 30 µg/ml IgG in 1×PBS for additional 2 h before adding to cell samples.

FRET-RotSensors: FRET-RotSensors were fabricated by labeling RotSensors with SAv-Alexa 568. The RotSensors were incubated with 27.5 µg/ml of SAv-Alexa568, 5 µg/ml of BSA, and 1 µg/ml of IgG in 1×PBS for 5 h at room temperature. Unbound proteins were rinsed off with artificial lysosome solution and 1×PBS. The FRET-RotSensors were further opsonized with 30 µg/ml of IgG in 1×PBS for additional 2 h before adding to cell samples.

**(c) Fabrication of Magnetically Modulated Phagosome Sensors (MagSensors)**

pH-MagSensors: The first step is the biotinylation of carboxylate-modified Dynabeads (1 µm). In brief, 10 µl of stock Dynabeads suspension (10 mg/ml) were added to 100 µl of MES buffer (50 mM, pH 6.2) containing 10 mg/ml EDC and 1 mM of biotin pentylamine. After 1-h incubation at room temperature, biotinylated Dynabeads (1 µm) were washed in 1×PBS to remove unbound biotin pentylamine. pH-MagSensors were fabricated by labeling biotinylated Dynabeads (1 µm) with SAv-pHrodo Red and SAv-CF640. Biotinylated Dynabeads were incubated with 50 µg/ml of SAv-pHrodo Red and 50 µg/ml of SAv-CF640 for 1 h at room temperature. The resulting pH-MagSensors were further opsonized with 1 mg/ml of IgG in 1×PBS for additional 1 h before adding to cell samples.

FRET-MagSensors: Biotinylated Dynabeads (1 µm) were prepared following the same procedure described above. FRET-MagSensors were fabricated by labeling biotinylated Dynabeads (1 µm) with SAv-Alexa568. Biotinylated Dynabeads were incubated with 27.5 µg/ml of SAv-Alexa568 for 1 h at room temperature. The resulting FRET-MagSensors were further opsonized with 1 mg/ml of IgG in 1×PBS for additional 1 h before adding to cell samples.

**Magnetic Tweezers Setup and Force Calibration**

Magnetic tweezers were built on an inverted fluorescence microscope system (Nikon Eclipse Ti-U,

Nikon, Tokyo, Japan), as shown in Supplementary Fig. 18. The setup mainly includes a solenoid

and a power supply. The solenoid was assembled by inserting a high permeability HyMu-80 alloy

rod (Carpenter Technology, Reading, PA) into an aluminum bobbin wrapped with 600 turns of

copper coil [59] (Supplementary Fig. 18b). The tip of the rod has a diameter of ≈ 1 µm (Supplementary

Fig. 18c). Position of the solenoid was controlled with a manual micromanipulator (Narishige

NMN-21) to achieve independent control of its position in the x-, y-, and z-direction with a

minimum graduation of 250 nm in the x-y plane and 1 µm in the z-direction. The current going

through the solenoid was generated by a programmable power supply (Tekpower, Montclair, CA)

with a maximum power output of 5 A.

Magnetic force was calibrated as a function of particle-to-tip distance by measuring the movements

of Dynabeads (1 µm) in magnetic field. In brief, CF640R labeled Dynabeads (1 µm) were

suspended in PDMS base at a concentration of ≈ $3.0 \times 10^5$ particles/ml. This low particle

concentration was necessary to avoid particle aggregation and inter-particle magnetic inference [60].

The PDMS base, which has a high viscosity of 5.1 Pa·s, was chosen to slow down particle

movements to make particle tracking feasible. The initial positions of the solenoid tip and of the

magnetic particles were imaged in bright field before time-lapse epi-fluorescence images of

particles were acquired with an interval time of 0.2 s. Working current of the magnetic tweezers

was 1.0 A in the calibration. The magnetic force $F(r)$ exerted on each magnetic particles at

particle-to-tip distance r was converted from particle velocity values using Stokes-Einstein

equation:

566
$$F(r) = 3 \cdot \pi \cdot \eta \cdot d \cdot v, \tag{1}$$

where $\eta$ is the viscosity of calibration solution PDMS base, d represents the diameter of particle, and v is the velocity of particle. The relationship between magnetic force F(r) and particle-to-tip distance $r$ was fitted using equation:

$$F(r) = \frac{F_0}{[\frac{r}{2r_0}+\frac{1}{2}]^{C_F}},$$ (2)

Where $F_0$ is a force constant with the unit of pN, $r_0$ is a distance constant with the unit of μm, $C_F$ is unitless and $F_{(r0)} = F_0$ [59].

**Fluorescence Microscopy**

**(a) Single-phagosome pH Assay**

RAW264.7 macrophage cells were seeded on glass coverslips at $2.0 \times 10^5$ cells/ml in complete medium for 24 h and then activated with LPS and IFN-γ following the procedure described above. For all single-phagosome pH studies, the recognition of pH-Sensors (i.e. pH-RotSensor and pH-MagSensor) by macrophages was synchronized following a previously reported protocol [61]. Briefly, cell samples were cooled on ice for 3 min to postpone phagocytosis. pH-Sensors were then added at a particle-to-cell ratio of ~5:1. Bindings of particles on macrophage were synchronized by centrifuging cell samples at 200×g for 30 s. Live cell imaging was conducted in Ringer's solution at 37°C on a Nikon Eclipse-Ti inverted microscope equipped with a 1.49 N.A. ×100 TIRF objective (Nikon, Tokyo, Japan) and an Andor iXon3 EMCCD camera (Andor Technology, Belfast, U.K.). Fluorescence emissions at three wavelengths (ex: 488, 561, and 660 nm; em: 515, 586 and 680 nm) were acquired for the time-lapse imaging. The acquisition rate was 2 sec/frame. Single-phagosome pH assay in actin-GFP expressing RAW264.7 macrophages was conducted following the same procedure described above.

  **(b) Single-phagosome FRET-fusion Assay**

RAW264.7 macrophage cells were seeded on glass coverslips at $2.0 \times 10^5$ cells/ml in complete

medium for 24 h and then activated with LPS and IFN-γ following the procedure described above.

Loading of either BSA-Alexa647-biotin or BSA-Alexa647 into lysosome compartment was carried

out following a previously reported pulse-chase protocol [62, 63, 64]. In brief, RAW264.7 macrophage

cells were incubated overnight in complete medium containing either BSA-Alexa647-biotin or

BSA-Alexa647 at indicated concentration. 2 h prior to live cell imaging, cells were rinsed twice

with complete medium. Labeled endocytic compartment was then chased at 37°C for 2 h to ensure

complete accumulation and fragmentation of fluorescently labeled BSA in lysosome compartments

(Supplementary Fig. 11 and Supplementary Fig. 12). Synchronized internalization of FRET-

Sensors (i.e. FRET-RotSensor and FRET-MagSensor) was conducted following the same

procedure described above. Time-lapse epi-fluorescence images were acquired to record FRET

emission (FRET$_0$; ex/em 561/680 nm), donor emission (Alexa568$_{em}$; ex/em 561/586 nm), acceptor

emission (Alexa647$_{em}$; ex/em 660/680 nm) and the emission from 100 nm yellow-green

fluorescence particle (ex/em 488/515 nm). The acquisition rate was 4 sec/frame.

The broad emission spectra of Alexa568 (donor) overlap partially with the acceptor emission

channel (680 nm) and the broad excitation spectra of Alexa647 (acceptor) did overlap partially with

donor excitation channel (561 nm). This resulted in two types of spectral cross-talks: (i) the

detection of donor fluorophore (Alexa568) emission at acceptor emission channel (680 nm) under

the donor excitation of 561 nm and (ii) the detection of acceptor fluorophore (Alexa647) emission

at acceptor emission channel (680 nm) under the donor excitation of 561 nm in the absence of FRET

effect. These two cross-talks, however, are constant throughout FRET-fusion assay, and could be

deducted from the measured FRET emission (FRET$_0$) to obtain FRET-generated emission

($FRET_{em}$). Correction factors $\alpha$ and $\beta$ for cross-talks were determined for each independent

experiment based on: (1) the measured FRET emission and donor emission of Alexa568

streptavidin particles outside cells and (2) the measured FRET emission and acceptor emission of acceptor fluorophore (Alexa647) in cells without FRET-sensor. The FRET ratio is obtained as follows:

$$FRET\ ratio = \frac{FRET_{em}}{AF568_{em}} = \frac{FRET_0 - \alpha \times AF568_{em} - \beta \times AF647_{em}}{AF568_{em}}, \qquad (3)$$

where $FRET_0$ is the measured FRET emission (ex/em 561/680 nm) before cross-talk correction, $FRET_{em}$ is the FRET emission, $AF568_{em}$ is the donor emission (ex/em 561/586 nm) and $AF647_{em}$ is the acceptor emission (ex/em 660/680 nm). The FRET increase was gradual, not stepwise, because multiple dye-loaded lysosomes can simultaneously fuse with a single phagosome and the increment in FRET signal from a single phagosome-lysosome fusion is too small to be resolved.

**(c) DQ-green BSA Digestion Assay**

To estimate the fragmentation level of BSA-Alexa647-biotin chasing, we loaded DQ-green BSA into lysosome compartment following the same pulse-chase produced described above, and measured the dequenching kinetics of lysosomal DQ-green BSA as a function of time. A DQ-green BSA concentration at 50 µg/ml was used during pulse. After the overnight pulse, cells were rinsed 3 times with flour-free Ringer's solution and imaged using Epi-fluorescence microscope. The emission of DQ-green (ex/em 488/515 nm) was recorded every 5 min for 2 h.

**(d) Immunofluorescence Staining and Imaging**

RAW 264.7 macrophages were seeded and activated as described above. For immunostaining of α-tubulin and F-actin, cells were first washed with 1× PBS and fixed with 2% PFA at room temperature for 5 min. Next, they were permeabilized with 0.1% Triton X-100 for 5 min at room temperature, and blocked with 2% BSA for 1 h at room temperature. After the BSA blocking, cells were incubated with 2 µg/mL Alexa Fluor 488 anti-tubulin-α antibody and 2 µg/mL Alexa Fluor

647 phalloidin for 1 h at room temperature. Super-resolution structured illumination microscopy

(SIM) images of the labeled cells were acquired using a DeltaVision OMX SR imaging system

equipped with a Olympus Plan Apo 60×/1.42 PSF objective and a sCMOS camera.

### (e) Lysotracker Labeling and Imaging

RAW 264.7 macrophages were seeded and activated as described above. After overnight incubation

in complete medium containing 5 µg/mL BSA-Alexa647-biotin, the labeled endocytic

compartments were chased at 37°C for 1.5 h. Cells were then incubated with 50 nM Lysotracker

Green DND-26 at 37°C for 30 min. After that, cells were rinsed twice with Ringer's solution to

remove the free fluorophore and imaged using Re-scan Confocal Microscopy (RCM) equipped with

a 1.49 N.A. ×100 TIRF objective and ORCA-fusion CMOS camera.

### Image Analysis

### (a) Image Registration

Image registration was done to correct for the optical shift between different imaging channels. 100

nm TetraSpeck fluorescent particles (Ex/Em: 360/430 nm, 505/515 nm, 560/580 nm, and 660/680

nm) were adsorbed on poly-l-lysine-coated glass coverslips at a surface density of $\approx 0.10$ bead/µm$^2$

and used as markers for image registration. After sequential imaging of the marker particles in three

channels (Ex: 488, 561, and 660 nm), an affine transformation was applied to align particle

localization maps from 488 and 660 nm channels (termed target images) to that from the 561 nm

channel (reference image) using ImageJ plugin MultistackReg [65]. A global mapping matrix was

obtained to record all the transformation steps and used to apply the same operations to all images.

### (b) Single-Particle Localization and Intensity Determination

The centroids of single particles in epi-fluorescence images were localized using a Gaussian-based

localization algorithm in the ImageJ plugin Trackmate. To measure the emission of CF640R and

pHrodo Red on pH-Sensors (i.e. pH-RotSensor and pH-MagSensor), pixel intensities within a

diameter of 2 μm from the localized centroid of the particle were integrated and background-

corrected using custom MATLAB algorithms. Same procedure was carried out for determining

$FRET_0$, $AF568_{em}$ and $AF647_{em}$ in single-phagosome FRET-fusion assay. To determine the

localization uncertainties, pH-RotSensors were immobilized on poly-l-lysine-coated glass coverslip

and imaged for 200 consecutive frames in Ringer's solution. Localization uncertainty was defined

as the standard deviation of the tracked particle positions in x- and y-coordinates. The localization

uncertainties of the 1 μm pHrodo Red-coated particle and 100 nm yellow-green fluorescence

nanoparticle were determined to be 20.6 ± 4.05 nm and 13.1 ± 2.32 nm, respectively

(Supplementary Fig. 25).

**(c) Translational and Rotational Tracking Analysis of RotSensors**

The single particle tracking analysis was done as described previously [24, 66]. Briefly, translational

velocity was determined from the centroid location of the 1 μm pHrodo Red- or SAv-Alexa568-

coated particle ($X_{red}, Y_{red}$) as a function of time. For rotational tracking, a vector was drawn from

the centroid of the 1 μm red fluorescent particle ($X_{red}, Y_{red}$) to that of the 100 nm yellow-green

nanoparticle ($X_{green}, Y_{green}$) belonging to the same RotSensor. Orientation of the vector was obtained

as the in-plane angle, $\varphi$, for the single RotSensor. The length of the vector was used to calculate

the out-of-plane angle, $\theta$, using the equation below:

689
$$\theta = arcsin\left(\frac{d}{r}\right), \tag{4}$$

690 where $d$ is the projection inter-particle distance between a pair of green and red dots in each given

691 image, and $r$ is the physical inter-particle distance. Because $r$ varies slightly from one particle to

692 another due to the size distribution, $r$ was obtained as the maximal projected inter-particle distance

$d_{max}$ when a RotSensor samples all possible orientations and the $d_{max}$ was larger than 550 nm. In cases when $d_{max}$ < 550 nm, $r$ was set to 550 nm. Two-dimensional rotational velocity was determined from the rotation matrix derived from $\varphi$ and $\theta$ at consecutive frames. Using this method, φ can be measured within a full range (−180° to +180°), but there is ambiguity in measuring θ, because RotSensors oriented at a polar angle of θ or (180° − θ) are indistinguishable from one another when observed as a two-dimensional projection image. However, this ambiguity does not affect the measurements of Δθ and translational velocity in this study, because Δθ of phagosomes between consecutive image frames (2s/frame) is expected to be < 90°, based on the result that phagosomes rotate an average Δφ of 0.12 ± 0.16 rad (6.9° ± 9.2°) between consecutive frames (Supplementary Fig. 26). Therefore, only Δθ < 90° was used in the tracking analysis for calculating rotational velocity of phagosomes.

**(d) Determination of Active Centripetal Run** The directional transport of phagosomes was analyzed following a previously reported method with slight modification [13]. The centroid of the nucleus was obtained based on bright field image of the cell (Supplementary Fig. 27) [48, 67]. At each time point of a particle trajectory, the distances between the phagosome and the centroid of the nucleus was quantified (referred to as 'distance-to-nucleus') (Fig. 4b). The travelled distance of phagosome towards the nucleus was subdivided into segments with a length of 32 seconds. Each segment was categorized as either a segment of active run or as a segment of passive motion based on the travelled distance with regards to nucleus. The distance threshold for active run was set to 0.5 μm, independent of the direction of motion whether it is towards the nucleus (centripetal) or towards the periphery (centrifugal) [68]. Finally, segments of active runs were classified either as active centripetal run or as active centrifugal run depending on the directionality of the transport.

**(e) Intracellular pH Calibration**

The intracellular pH calibration of pH-RotSensors and pH-MagSensors was done following a previously published protocol [26]. Macrophages seeded on glass coverslips were pretreated with 10 µM concanamycin in Ringer's solution for 10 min before the addition of particles and then incubated at 37°C for 10 min to promote particle phagocytosis. Cell medium was replaced with potassium-rich buffers at different pH values. All buffers contain 20 µM of nigericin, but different buffering agent: 5 mM acetic acid for pH 4.5, 5 mM MES for pH 5.5, and 5mM HEPES for pH 6.5 and 7.3. The pH calibration was done from pH 7.3 to 4.5. For each pH, cells were rinsed twice with the correspondingly buffer and allowed to equilibrate for 5-10 min before image acquisition. Fluorescence emission of the pHrodo Red at 586 nm ($I_{pHrodoRed}$) and reference dye CF640R at 680 nm ($I_{ref}$) was obtained at various pH and background-corrected to obtain ratiometric pH calibration plots (Supplementary Fig. 2).

In live cell imaging, pH calibration was done for individual internalized pH-RotSensors and pH-MagSensors to eliminate the effect of particle-to-particle variation in their pH responses. By plotting $I_{pHrodo}/I_{ref}$ vs. time for single phagosomes, we notice that acidification process started with a standby period during which $I_{pHrodo}/I_{ref}$ ratio remains unchanged denoting that the phagosome pH remained equivalent to extracellular pH of 7.3 (Ringer's solution, pH = 7.3) before the initiation of phagosome acidification (Supplementary Fig. 28a and b). We further confirmed this by imaging phagosome acidification in cells expressing actin-GFP, in which intensity peak of actin-GFP pinpointed the time of particle internalization [42, 43] (Supplementary Fig. 28c-e). After live cell imaging, the same cell sample was incubated in pH 4.5 potassium-rich pH calibration buffer containing 20 µM of nigericin to obtain the ratiometric emission ($I_{pHrodo}/I_{ref}$) of the RotSensor at pH 4.5 (Supplementary Fig. 28a). In the final step of the calibration procedure, a linear function was generated based on the known $I_{pHrodo}/I_{ref}$ ratios at pH 4.5 and 7.3, and the result function

was used to transform the fluorescence measurements of the RotSensor to luminal pH values (Supplementary Fig. 28b and f).

## (f) Wavelet Transform Analysis

A wavelet analysis algorithm reported by Chen *et al*. was used to distinguish active rotation from passive one. Both in-plane ($\theta$) and out-of-plane ($\varphi$) displacement of a RotSensor was convoluted using the 1-dimensional Haar continuous time wavelet transform [32]. The wavelet coefficients after the transformation were plotted as a function of time and for different widths of the wavelet function (Supplementary Fig. 8). The width of the wavelet function used here is the number of image frames and referred to as "scale". Scale 10 was chosen as a universal threshold such that no active rotation of phagosomes was detected in cells that were treated with microtubule-disrupting drug nocodazole.

## (g) Determination of location of cell entry

The boundary of the cell was manually traced in ImageJ according to the fluorescence emission (ex/em 561/586 nm) of the cell in Ringer's solution containing trypan blue (Supplementary Fig. 27a). The boundary and the center of the nucleus was manually traced in ImageJ based on the bright field image of the given cell [48, 69] (Supplementary Fig. 27b).

## (h) Calculation of Mean-Square Displacements (MSD)

Time-resolved MSD shown in Fig. 4d was analyzed using custom MATLAB algorithms. Briefly, the MSD values of each single phagosome trajectories were calculated as a function of lag time $\boldsymbol{\delta t}$ based on the following equation:

$$MSD(\delta t) = \; < \Delta R(t)^2 > \; = \; < (R(t + \delta t) - R(t))^2 > \tag{5}$$

Where $(R(t + \delta t) - R(t))$ is end-to-end displacement of an individual particle from the time point t to $t + \delta t$.

**Transmission Electron Microscopy**

Cell samples for TEM imaging were prepared based on a previously reported protocol with slight

modification [66]. Particle sensors were incubated with cells for 50 min at 37 °C before trypsinization.

After centrifugation, the cell pellet was fixed on ice for 1 h by using a mixture of 2.5% (w/v)

glutaraldehyde and 1% (w/v) osmium tetroxide in 1×PBS buffer. Subsequently, the pellet was

stained in 0.5% (w/v) uranyl acetate aqueous solution for 12 h on ice. Cell pellets were dehydrated

sequentially in a series of ice-cold aqueous solutions containing 30% (v/v), 50%, 75%, 90%, 95%

and 100% ethanol for 5 min each. Dehydration in 100% (v/v) ethanol was repeated three times at

room temperature. The dehydrated cell pellet was immersed sequentially in resin infiltration

solutions that contain ethanol and Spurr's resin at various volume ratio (ethanol: resin, 2:1, 1:1 and

1:2) for 30 min each at room temperature. Cells were cured in 100% resin for 18 h at 65°C prior to

microtome sectioning. Sections were stained with 2% (w/v) uranyl acetate for 10 min before

imaging. Samples were imaged with a JEOL JEM-1010 Transmission Electron Microscope

(Electron Microscopy Center, Indiana University).

**References**

1.    Harrison RE, Grinstein S. Phagocytosis and the microtubule cytoskeleton. *Biochem Cell Biol* **80**, 509-515 (2002).

2.    Al-Haddad AH*, et al.* Myosin va bound to phagosomes binds to F-actin and delays microtubule-dependent motility. *Mol Biol Cell* **12**, 2742-2755 (2001).

3.    Araki N. Role of microtubules and myosins in Fc gamma receptor-mediated phagocytosis. *Front Biosci* **11**, 1479-1490 (2006).

4.    Kelleher JF, Titus MA. Intracellular motility: How can we all work together? *Curr Biol* **8**, R394-R397 (1998).

5.    Oberhofer A*, et al.* Molecular underpinnings of cytoskeletal cross-talk. *P Natl Acad Sci USA* **117**, 3944-3952 (2020).

6.   Steinberg B, Huynh K, Grinstein S. Phagosomal acidification: measurement, manipulation
and functional consequences.). Portland Press Ltd. (2007).

7.   Lennon-Duménil A-M*, et al.* Analysis of protease activity in live antigen-presenting cells
shows regulation of the phagosomal proteolytic contents during dendritic cell activation. *J*
*Exp Med* **196**, 529-540 (2002).

8.   Rybicka JM, Balce DR, Chaudhuri S, Allan ERO, Yates RM. Phagosomal proteolysis in
dendritic cells is modulated by NADPH oxidase in a pH-independent manner. *Embo J* **31**,
932-944 (2012).

9.   Smith LM, May RC. Mechanisms of microbial escape from phagocyte killing. *Biochem*
*Soc Trans* **41**, 475-490 (2013).

10.  Cossart P, Helenius A. Endocytosis of viruses and bacteria. *Cold Spring Harb Perspect*
*Biol* **6**,  (2014).

11.  Kagan JC, Roy CR. Legionella phagosomes intercept vesicular traffic from endoplasmic
reticulum exit sites. *Nat Cell Biol* **4**, 945-954 (2002).

12.  Blocker A, Griffiths G, Olivo JC, Hyman AA, Severin FF. A role for microtubule
dynamics in phagosome movement. *Journal of Cell Science* **111**, 303-312 (1998).

13.  Keller S, Berghoff K, Kress H. Phagosomal transport depends strongly on phagosome
size. *Sci Rep* **7**, 17068 (2017).

14.  Jiang M*, et al.* Microtubule motors transport phagosomes in the RPE, and lack of KLC1
leads to AMD-like pathogenesis. *Journal of Cell Biology* **210**, 595-611 (2015).

15.  Rybicka JM, Balce DR, Khan MF, Krohn RM, Yates RM. NADPH oxidase activity
controls phagosomal proteolysis in macrophages through modulation of the lumenal redox
environment of phagosomes. *Proc Natl Acad Sci U S A* **107**, 10496-10501 (2010).

16.  Guerin I, de Chastellier C. Disruption of the actin filament network affects delivery of
endocytic contents marker to phagosomes with early endosome characteristics: The case
of phagosomes with pathogenic mycobacteria. *Eur J Cell Biol* **79**, 735-749 (2000).

17.  Gibbs D, Kitamoto J, Williams DS. Abnormal phagocytosis by retinal pigmented
epithelium that lacks myosin Vila, the Usher syndrome 1B protein. *P Natl Acad Sci USA*
**100**, 6481-6486 (2003).

18.  Arora PD, Manolson MF, Downey GP, Sodek J, McCulloch CA. A novel model system
for characterization of phagosomal maturation, acidification, and intracellular collagen
degradation in fibroblasts. *J Biol Chem* **275**, 35432-35441 (2000).

19.  Funato K, Beron W, Yang CZ, Mukhopadhyay A, Stahl PD. Reconstitution of
phagosome-lysosome fusion in streptolysin O-permeabilized cells. *Journal of Biological*
*Chemistry* **272**, 16147-16151 (1997).
847

20. Blocker A, Severin FF, Habermann A, Hyman AA, Griffiths G, Burkhardt JK. Microtubule-associated protein-dependent binding of phagosomes to microtubules. *J Biol Chem* **271**, 3803-3811 (1996).

21. Desjardins M, Huber LA, Parton RG, Griffiths G. Biogenesis of Phagolysosomes Proceeds through a Sequential Series of Interactions with the Endocytic Apparatus. *Journal of Cell Biology* **124**, 677-688 (1994).

22. Claus V*, et al.* Lysosomal enzyme trafficking between phagosomes, endosomes, and lysosomes in J774 macrophages enrichment of cathepsin H in early endosomes. *J Biol Chem* **273**, 9842-9851 (1998).

23. Savina A*, et al.* NOX2 controls phagosomal pH to regulate antigen processing during crosspresentation by dendritic cells. *Cell* **126**, 205-218 (2006).

24. Yu YQ, Gao Y, Yu Y. "Waltz" of Cell Membrane-Coated Nanoparticles on Lipid Bilayers: Tracking Single Particle Rotation in Ligand-Receptor Binding. *Acs Nano* **12**, 11871-11880 (2018).

25. Deriy LV*, et al.* Disease-causing Mutations in the Cystic Fibrosis Transmembrane Conductance Regulator Determine the Functional Responses of Alveolar Macrophages. *Journal of Biological Chemistry* **284**, 35926-35938 (2009).

26. Naufer A*, et al.* pH of endophagosomes controls association of their membranes with Vps34 and PtdIns(3)P levels. *Journal of Cell Biology* **217**, 329-346 (2018).

27. Nunes P, Guido D, Demaurex N. Measuring Phagosome pH by Ratiometric Fluorescence Microscopy. *Jove-J Vis Exp*, (2015).

28. Canton J, Khezri R, Glogauer M, Grinstein S. Contrasting phagosome pH regulation and maturation in human M1 and M2 macrophages. *Mol Biol Cell* **25**, 3330-3341 (2014).

29. Ogawa M, Kosaka N, Regino CAS, Mitsunaga M, Choyke PL, Kobayashi H. High sensitivity detection of cancer in vivo using a dual-controlled activation fluorescent imaging probe based on H-dimer formation and pH activation. *Mol Biosyst* **6**, 888-893 (2010).

30. Yates RM, Hermetter A, Russell DG. The kinetics of phagosome maturation as a function of phagosome/lysosome fusion and acquisition of hydrolytic activity. *Traffic* **6**, 413-420 (2005).

31. Gao Y, Anthony SM, Yu YQ, Yi Y, Yu Y. Cargos Rotate at Microtubule Intersections during Intracellular Trafficking. *Biophys J* **114**, 2900-2909 (2018).

32. Chen KJ, Wang B, Guan J, Granick S. Diagnosing Heterogeneous Dynamics in Single-Molecule/Particle Trajectories with Multiscale Wavelets. *Acs Nano* **7**, 8634-8644 (2013).

33. Sun-Wada GH, Tabata H, Kawamura N, Aoyama M, Wada Y. Direct recruitment of H+-ATPase from lysosomes for phagosomal acidification. *J Cell Sci* **122**, 2504-2513 (2009).

34. Pauwels AM, Trost M, Beyaert R, Hoffmann E. Patterns, Receptors, and Signals: Regulation of Phagosome Maturation. *Trends Immunol* **38**, 407-422 (2017).

35. Riazanski V*, et al.* TRPC6 channel translocation into phagosomal membrane augments phagosomal function. *P Natl Acad Sci USA* **112**, E6486-E6495 (2015).

36. Mcneil PL, Tanasugarn L, Meigs JB, Taylor DL. Acidification of Phagosomes Is Initiated before Lysosomal-Enzyme Activity Is Detected. *Journal of Cell Biology* **97**, 692-702 (1983).

37. Duvvuri M, Krise JP. A novel assay reveals that weakly basic model compounds concentrate in lysosomes to an extent greater than pH-partitioning theory would predict. *Mol Pharmaceut* **2**, 440-448 (2005).

38. Fahrer J, Funk J, Lillich M, Barth H. Internalization of biotinylated compounds into cancer cells is promoted by a molecular Trojan horse based upon core streptavidin and clostridial C2 toxin. *N-S Arch Pharmacol* **383**, 263-273 (2011).

39. Horn MA, Heinstein PF, Low PS. Characterization of Parameters Influencing Receptor-Mediated Endocytosis in Cultured Soybean Cells. *Plant Physiol* **98**, 673-679 (1992).

40. Pu J, Guardia CM, Keren-Kaplan T, Bonifacino JS. Mechanisms and functions of lysosome positioning. *J Cell Sci* **129**, 4329-4339 (2016).

41. Jiang C, Lionberger TA, Wiener DM, Meyhofer E. Electromagnetic tweezers with independent force and torque control. *Rev Sci Instrum* **87**, 084304 (2016).

42. Scott CC*, et al.* Phosphatidylinositol-4,5-bisphosphate hydrolysis directs actin remodeling during phagocytosis. *Journal of Cell Biology* **169**, 139-149 (2005).

43. Huang NN*, et al.* Canonical and Noncanonical G-Protein Signaling Helps Coordinate Actin Dynamics To Promote Macrophage Phagocytosis of Zymosan. *Mol Cell Biol* **34**, 4186-4199 (2014).

44. Hendricks AG, Holzbaur EL, Goldman YE. Force measurements on cargoes in living cells reveal collective dynamics of microtubule motors. *Proc Natl Acad Sci U S A* **109**, 18447-18452 (2012).

45. Kaufmann AM, Goldman SD, Krise JP. A fluorescence resonance energy transfer-based approach for investigating late endosome-lysosome retrograde fusion events. *Anal Biochem* **386**, 91-97 (2009).

46. Yates RM, Russell DG. Phagosome maturation proceeds independently of stimulation of toll-like receptors 2 and 4. *Immunity* **23**, 409-417 (2005).

47. Huynh KK, Eskelinen EL, Scott CC, Malevanets A, Saftig P, Grinstein S. LAMP proteins are required for fusion of lysosomes with phagosomes. *EMBO J* **26**, 313-324 (2007).

48. Dumas A*, et al.* The HIV-1 protein Vpr impairs phagosome maturation by controlling microtubule-dependent trafficking. *J Cell Biol* **211**, 359-372 (2015).

49.    Blocker A*, et al.* Molecular requirements for bi-directional movement of phagosomes along microtubules. *J Cell Biol* **137**, 113-129 (1997).

50.    Rai A, Pathak D, Thakur S, Singh S, Dubey AK, Mallik R. Dynein Clusters into Lipid Microdomains on Phagosomes to Drive Rapid Transport toward Lysosomes. *Cell* **164**, 722-734 (2016).

51.    Rai AK, Rai A, Ramaiya AJ, Jha R, Mallik R. Molecular Adaptations Allow Dynein to Generate Large Collective Forces inside Cells. *Cell* **152**, 172-182 (2013).

52.    Ferro LS, Can S, Turner MA, ElShenawy MM, Yildiz A. Kinesin and dynein use distinct mechanisms to bypass obstacles. *Elife* **8**,  (2019).

53.    Harrison RE, Bucci C, Vieira OV, Schroer TA, Grinstein S. Phagosomes fuse with late endosomes and/or lysosomes by extension of membrane protrusions along microtubules: role of Rab7 and RILP. *Mol Cell Biol* **23**, 6494-6506 (2003).

54.    Gowrishankar S*, et al.* Massive accumulation of luminal protease-deficient axonal lysosomes at Alzheimer's disease amyloid plaques. *P Natl Acad Sci USA* **112**, E3699-E3708 (2015).

55.    Johnson DE, Ostrowski P, Jaumouille V, Grinstein S. The position of lysosomes within the cell determines their luminal pH. *Journal of Cell Biology* **212**, 677-692 (2016).

56.    Balderhaar HJK, Ungermann C. CORVET and HOPS tethering complexes - coordinators of endosome and lysosome fusion. *J Cell Sci* **126**, 1307-1316 (2013).

57.    Kissing S*, et al.* Vacuolar ATPase in Phagosome-Lysosome Fusion. *Journal of Biological Chemistry* **290**, 14166-14180 (2015).

58.    Pelfrene A, Cave MR, Wragg J, Douay F. In Vitro Investigations of Human Bioaccessibility from Reference Materials Using Simulated Lung Fluids. *Int J Environ Res Public Health* **14**,  (2017).

59.    Zahn C*, et al.* Measurement of the magnetic moment of single Magnetospirillum gryphiswaldense cells by magnetic tweezers. *Sci Rep* **7**, 3558 (2017).

60.    Etoc F, Lisse D, Bellaiche Y, Piehler J, Coppey M, Dahan M. Subcellular control of Rac-GTPase signalling by magnetogenetic manipulation inside living cells. *Nat Nanotechnol* **8**, 193-198 (2013).

61.    FitzGerald LI, Johnston APR. It's what's on the inside that counts: Techniques for investigating the uptake and recycling of nanoparticles and proteins in cells. *J Colloid Interf Sci* **587**, 64-78 (2021).

62.    Jeschke A, Haas A. Sequential actions of phosphatidylinositol phosphates regulate phagosome-lysosome fusion. *Mol Biol Cell* **29**, 452-465 (2018).

63. Jeschke A*, et al.* Phosphatidylinositol 4-phosphate and phosphatidylinositol 3-phosphate regulate phagolysosome biogenesis. *P Natl Acad Sci USA* **112**, 4636-4641 (2015).

64. Becken U, Jeschke A, Veltman K, Haas A. Cell-free fusion of bacteria-containing phagosomes with endocytic compartments. *Proc Natl Acad Sci U S A* **107**, 20726-20731 (2010).

65. Micheva KD, O'Rourke N, Busse B, Smith SJ. Array tomography: semiautomated image alignment. *Cold Spring Harb Protoc* **2010**, pdb prot5527 (2010).

66. Sanchez L, Yi Y, Yu Y. Effect of partial PEGylation on particle uptake by macrophages. *Nanoscale* **9**, 288-297 (2017).

67. Le-Bury G, Deschamps C, Dumas A, Niedergang F. Phagosome Migration and Velocity Measured in Live Primary Human Macrophages Infected with HIV-1. *Jove-J Vis Exp*, (2016).

68. Gross SP, Tuma MC, Deacon SW, Serpinskaya AS, Reilein AR, Gelfand VI. Interactions and regulation of molecular motors in Xenopus melanophores. *Journal of Cell Biology* **156**, 855-865 (2002).

69. Ba QL, Raghavan G, Kiselyov K, Yang G. Whole-Cell Scale Dynamic Organization of Lysosomes Revealed by Spatial Statistical Analysis. *Cell Reports* **23**, 3591-3606 (2018).

**Acknowledgments:** We thank Prof. Sergio Grinstein at University of Toronto for his helpful discussion. We thank Mr. Andy Alexander at the Electronic Instrument Services of Indiana University for assistance with the setup of magnetic tweezers. We thank Dr. Barry Stein at the IUB Electron Microscopy Imaging Center, Dr. Jim Power at the IUB Light Microscopy Imaging Center (LMIC), Dr. Yi Yi at the Nanoscale Characterization Center for assistance with instrument use.

**Funding:** The development of single-particle rotational tracking method was supported by the National Science Foundation, Division of Chemical, Bioengineering, Environmental, and Transport Systems under Award No. 1554078. All other research reported in this publication was supported by the National Institute of General Medical Sciences of the NIH under Award No. R35GM124918 (Y. Y.). G.F.W.W. is supported by a Vanier Canada Graduate Scholarship (CIHR). The content is

1033 solely the responsibility of the authors and does not necessarily represent the official views of the

1034 NIH.

1035 **Author contributions:** Y.-q. Y., Z. Z., and Y. Y. designed research; Y.-q. Y. and Z. Z. performed

1036 research and analyzed data; G. F. W. W. contributed critical reagents; Y.-q. Y., Z. Z., and Y. Y.

1037 wrote the paper and G.F.W. W. provided critical discussion.

1038

1039 **Competing interests:** The authors declare that they have no competing interests.

1040 **Data and materials availability:** All data needed to evaluate the conclusions in the paper are

1041 present in the paper and/or the Supplementary Materials. Additional data related to this paper may

1042 be requested from the authors.

1043

1044

1045

1046

1047

1048

1049

1050

1051

**Figures**

**Fig. 1 Simultaneous measurement of phagosome dynamics and acidification. a** Schematics showing internalization

of pH-sensitive RotSensors into phagosomes in macrophage cells. **b** Schematic illustration of the RotSensor design.

Each RotSensor contains a 1 μm silica particle tethered with a 100 nm yellow-green fluorescent particle. The silica

particles were coated with streptavidin (SAv)-pHrodo Red (pH reporter) and SAv-CF640R (reference). The azimuthal

(φ) and polar (θ) angles of each pH-RotSensor were analyzed based on its projection fluorescence image. d and r denote

the projection inter-particle distance and the physical inter-particle distance, respectively. **c** A representative trajectory

of a pH-RotSensor-containing phagosome. Vectors indicates the orientation of the pH-RotSensor and are color-coded

based on time. Scale bar, 2 μm. Insets show fluorescence images of the RotSensor (top) and the pH response (bottom).

Scale bar, 1 μm. **d-f** Line plots showing acidification (**d**), translational (**e**) and rotational velocities (**f**) of the pH-

RotSensor-containing phagosome in **c**. The acidification profile is fitted with sigmoidal-Boltzmann function to

determine the initial pH, final pH, the period of rapid acidification ($t_{initial}$ to $t_{final}$), and acidification rate. The period

of rapid acidification is highlighted in grey. Darker lines in **e** and **f** are velocity values after wavelet denoising. **g** and **h**

Scattered plots showing acidification rates against translational (**g**) and rotational velocities (**h**) of single phagosomes.

Each data point represents one single phagosome. Data points from multiple phagosomes within the same cells are

shown in the same solid color. Data points from cells containing only one phagosome are shown as black circles. N =

1068  42 phagosomes from 24 cells for translational tracking, and 17 phagosomes from 12 cells for rotational tracking.

1069  Pearson's coefficients of 0.78 and 0.81 were obtained in **g** and **h**, respectively. **I** and **j** Scatter plots showing acidification

1070  rates against percentage of active rotation (**i**) and maximum rotation amplitude (**j**) of single phagosomes during rapid

1071  acidification period. N = 17 phagosomes from 12 cells. Pearson's coefficients of 0.76 and 0.61 are obtained in **i** and **j**.

1072

1073

**Fig. 2 Simultaneous measurement of phagosome-lysosome fusion and phagosome transport dynamics. a**

Schematic illustration of the Förster resonance energy transfer (FRET)-based phagosome-lysosome fusion assay. The

FRET-RotSensor was coated with SAv-Alexa568 (FRET donor). BSA-Alexa647-biotin (FRET acceptor) was loaded

into lysosomes. Phagosome fusion with lysosomes leads to intermixing between donor fluorophore (Alexa568) and

fluid phase acceptor fluorophore (Alexa647) generating FRET emission (680 nm) under the donor excitation of 561

nm. **b** and **c** Fluorescence images and intensity plots showing the change of FRET emission (ex/em: 561/680 nm) and

donor emission (red, ex/em: 561/586 nm) from a FRET-RotSensor (100 nm yellow-green nanoparticle in green) in a

RAW264.7 macrophage. Scale bar, 2μm. **d** FRET ratio vs. time plot is fitted with sigmoidal-Boltzmann function

(shown as the black solid line) to determine the initial time point of phagosome-lysosome fusion ($t_{initial}$), the time point

where FRET-signal reaches a plateau ($t_{final}$), and the FRET rate, as indicated by the red dotted line. **e** and **f** Scatter

plots showing normalized FRET rate against translational (**e**) and rotational velocities (**f**) of single phagosomes during

the period of its fusion with lysosomes. Each data point represents data from a single phagosome. Data points from

multiple phagosomes within the same cells are shown in the same solid color. Data points from cells containing only

one phagosome are shown as black circles. For translational tracking, N = 40 phagosomes from 22 cells. For rotational

tracking, N = 11 phagosomes from 10 cells. The black lines indicate linear regression with a Pearson's coefficient of

0.75 in **e** and of 0.86 in **f**.

**Fig. 3 Correlation of phagosome-lysosome fusion kinetics with the centripetal motility of phagosomes. a** Fluorescence image showing a cell in which lysosomes were labeled with BSA-Alexa647-biotin. Green line indicates the cell periphery, and the asterisk indicates the centroid of the nucleus. The initial positions of two representative phagosomes were indicated by the two number-labeled white dots and their trajectories are shown in red. Scale bar, 10 μm. **b** Plots showing the phagosome-to-nucleus distance as a function of time for both the phagosomes indicated in **a**. Segments of active centripetal runs are highlighted in green. Grey shaded area indicates the effective transport distance of the phagosome from the time where it begins to fuse with lysosomes to the time when it reaches the nucleus boundary. The starting time of phagosome lysosome fusion was obtained based on sigmoidal fitting the FRET ratio vs. time plot. Phagosome #1 underwent active centripetal runs in 48% of the time during effective transport, whereas phagosome #2 had no active centripetal run. **c** FRET ratio vs. time plots for both phagosomes marked in **a**. Solid lines indicate sigmoidal-Boltzmann fitting to the data. **d** and **e** Scatter plots showing normalized FRET rate of single phagosomes plotted against the percentage of active centripetal motion (**d**) and centripetal velocity (**e**). Each data point represents data from a single phagosome. Data from multiple phagosomes inside the same cell are shown in the same solid color. Data from cells in which only one phagosome was studied are shown as black circles. N = 40 phagosomes

 from 22 cells. The black lines indicate linear regression with a Pearson's coefficient of 0.68 and 0.49, respectively.

1107

1108

**Fig. 4 Phagosome degradation measurements during magnetically manipulated transport. a** Schematic

illustration of the experimental setup. The magnetic gradient is generated by a homemade magnetic tweezers setup

built on an inverted fluorescence microscope system. Magnetic pulling force was applied on MagSensors after their

internalization into phagosomes. The 1 μm MagSensors were coated with SAv-pHrodo Red (pH indicator), SAv-

CF640 (reference), and physically adsorbed IgG. **b** Calibration plot showing the magnetic force exerted on each

MagSensor as a function of its distance to the tip of the magnetic tweezers solenoid (shown in inset). Error bars are

standard deviations from 5 samples. **c** Bright-field image of a cell overlaid with the trajectory of a MagSensor-

containing phagosome under magnetic pulling. The start and end time points of the exertion of magnetic force are

indicated. The blue-colored segment of the trajectory indicates the movement of the MagSensor before cell entry; the

red-colored segment of the trajectory indicates the intracellular movement of a MagSensor-containing phagosome

under magnetic manipulation. Scale bar, 5 μm. **d** Mean-square displacements (MSD) calculated from trajectories of

individual MagSensors under different conditions as indicated. Each line is an average of results from N = 10

phagosomes from 8 cells (no force), 9 cells (magnetic pulling), and 7 cells (nocodazole). Shaded areas indicate

standard deviations.

1123

1124

**Fig. 5 Phagosome acidification and phagosome-lysosome fusion under magnetic pulling. a** Line plots showing the

average phagosome pH as a function of time with or without magnetic pulling as indicated. Each line plot is an average

from 20 phagosomes. Shaded areas represent standard deviations. **b** Box graph showing the average acidification rate

of phagosomes with or without magnetic pulling. The average acidification rate is $0.48 \pm 0.34$ pH unit/min without

magnetic pulling (N = 33 phagosomes in 29 cells from 11 independent experiments) and $0.79 \pm 0.54$ pH unit/min with

magnetic pulling (N = 38 phagosomes in 38 cells from 13 independent experiments). **c** Box graph showing the average

final pH in different experiment conditions as indicated. The average final pH is $4.7 \pm 0.3$ without magnetic

manipulation (N = 33) and $4.7 \pm 0.4$ with magnetic pulling (N = 38). **d** Line plots showing the average normalized

FRET ratio as a function of time with or without magnetic pulling as indicated. The line curves are averaged from 20

individual phagosomes in each experimental condition. Shaded areas represent standard deviations. **e** Box graph

showing the average normalized FRET rate in different experiment conditions as indicated. The average FRET rate is

$0.15 \pm 0.07$ without magnetic manipulation (N = 20 phagosomes in 16 cells from 4 independent experiments) and 0.28

$\pm 0.21$ with magnetic pulling (N = 20 phagosomes in 19 cells from 8 independent experiments). In **b**, **c** and **e**, each box

plot indicates the mean (horizontal line) and the interquartile range from 25% to 75% of the corresponding data set.

Statistical significance is highlighted by p-values (Mann-Whitney Test) as follows: ** $p < 0.01$, *** $p < 0.0001$, NS p

$> 0.05$.