Cell-type specific circadian bioluminescence rhythms recorded

from *Dbp* reporter mice reveal circadian oscillator misalignment

[a,b,1] Ciearra B. Smith, [a,c,1] Vincent van der Vinne, [d] Eleanor McCartney, [e] Adam C. Stowie, [f] Tanya L. Leise, [d,2] Blanca Martin-Burgos, [d] Penny C. Molyneux, [g] Lauren A. Garbutt, [h] Michael H. Brodsky, [e] Alec J. Davidson, [d] Mary E. Harrington, [g] Robert Dallmann, and [a,b,j,3] David R. Weaver

[a]   Department of Neurobiology, University of Massachusetts Medical School, Worcester MA

[b]   Graduate Program in Neuroscience, University of Massachusetts Medical School, Worcester MA

[c]   Department of Biology, Williams College, Williamstown, MA

[d]   Neuroscience Program, Smith College, Northampton MA

[e]   Neuroscience Institute, Morehouse School of Medicine, Atlanta GA

[f]   Department of Mathematics and Statistics, Amherst College, Amherst MA

[g]   Division of Biomedical Sciences, Warwick Medical School, University of Warwick, Coventry, UK

[h]   Department of Molecular, Cell and Cancer Biology, University of Massachusetts Medical School, Worcester MA

[i]   Department of Biochemistry and Molecular Pharmacology, University of Massachusetts Medical School, Worcester MA

[j]   NeuroNexus Institute, University of Massachusetts Medical School, Worcester MA

[1] C.B.S and V.v.d.V contributed equally to this work.

[2] Present address: University of California, San Diego, La Jolla, CA USA

[3] To whom correspondence may be addressed:

   Email: David.weaver@umassmed.edu

   David R. Weaver, Ph.D., Department of Neurobiology, LRB-723, UMass Medical School, 364 Plantation St., Worcester MA 01605

ORCID ID's:
CBS   0000-0003-2999-3387   Ciearra.Smith@umassmed.edu
VvdV 0000-0003-3926-5041   vv5@williams.edu
EM   0000-0002-5806-1995   mccartneyee1921@gmail.com
ACS   0000-0002-7007-9135   astowie@msm.edu
TLL   0000-0002-7458-7604   tleise@amherst.edu
BMB   0000-0002-1388-9123   bmartinb@ucsd.edu
PCM   pmolyneu@smith.edu
LAG   0000-0002-9366-4468   L.A.Garbutt@warwick.ac.uk
MHB   0000-0001-7703-4872   Michael.brodsky@umassmed.edu
AJD   0000-0003-4205-1968   adavidson@msm.edu
MEH   0000-0003-2266-6455   mharring@smith.edu
RD   0000-0002-7490-0218   R.Dallmann@warwick.ac.uk
DRW   0000-0001-7941-6719   david.weaver@umassmed.edu

Classification

43   Major Classification: Biological Sciences

44   SubClassification: Physiology

Keywords (at least three and no more than five).

47   Circadian Rhythms, Bioluminescence, Luciferase, Misalignment, Liver

Conflict of interest statement: The authors declare no conflicts of interest.

This PDF contains

52   Main Text

53   Figures 1 to 7

54   Table 1

Abstract

Circadian rhythms are endogenously generated physiological and molecular rhythms with a cycle length of

about 24 h. Bioluminescent reporters have been exceptionally useful for studying circadian rhythms

in numerous species. Here, we report development of a reporter mouse generated by modification

of a widely expressed and highly rhythmic gene encoding D-site albumin promoter binding protein

(*Dbp*). In this line of mice, firefly luciferase is expressed from the *Dbp* locus in a *Cre*-recombinase-

dependent manner, allowing assessment of bioluminescence rhythms in specific cellular

populations. A mouse line in which luciferase expression was *Cre*-independent was also generated.

The *Dbp* reporter alleles do not alter *Dbp* gene expression rhythms in liver or circadian locomotor

activity rhythms. *In vitro* and *in vivo* studies show the utility of the reporter alleles for monitoring

rhythmicity. Our studies reveal cell-type specific characteristics of rhythms among neuronal

populations within the suprachiasmatic nuclei *in vitro*. *In vivo* studies show stable *Dbp*-driven

bioluminescence rhythms in the liver of *Albumin-Cre;Dbp^{KI/+}* "liver reporter" mice. After a shift

of the lighting schedule, locomotor activity achieved the proper phase relationship with the new

lighting cycle more rapidly than hepatic bioluminescence did. As previously shown, restricting

food access to the daytime altered the phase of hepatic rhythmicity. Our model allowed assessment

of the rate of recovery from misalignment once animals were provided with food *ad libitum*. These

studies provide clear evidence for circadian misalignment following environmental perturbations

and reveal the utility of this model for minimally invasive, longitudinal monitoring of rhythmicity

from specific mouse tissues.

Significance Statement

Disruption of temporal coordination among circadian oscillators and exposure to light at

biologically inappropriate times are important drivers of the increased incidence of adverse health

outcomes observed in shift workers and rodent models of chronic circadian disruption. Here, we

demonstrate the utility of a new mouse line that enables tissue-specific monitoring of circadian

molecular rhythms *in vivo* and *ex vivo*. This reporter mouse provides a major advance in our

capabilities for monitoring rhythms in a variety of tissues under normal and disruptive conditions.

Our studies provide an unprecedented longitudinal assessment of tissue-specific rhythmicity, a key

step in the identification of mechanisms underlying the circadian disruption inherent to life in

modern 24/7 societies.

Introduction

Circadian rhythms are endogenous rhythms with a cycle length of ~24 hours. The mammalian circadian system is hierarchical. The hypothalamic suprachiasmatic nuclei (SCN) serve as the pacemaker[1,2]. The SCN are synchronized by environmental cues, of which the light-dark cycle is the most influential. The SCN are not unique in their capacity for rhythmicity, however. The transcriptional-translational feedback loop regulating molecular oscillations in the SCN is also present in individual cells throughout the body[1,2]. SCN-driven neural, behavioral and hormonal rhythms synchronize these cell-autonomous oscillators, leading to rhythmicity with predictable phase relationships among tissues, genes and physiological processes[1-4]. Repeated disruption of this internal temporal order by inappropriately timed light exposure or food intake leads to adverse health consequences in shift-working humans and animal models[4-15]. Progress in identifying the mechanisms by which chronic circadian disruption leads to adverse health consequences will require long-term monitoring of central and peripheral rhythms[7,8].

Rhythmically expressed reporter genes have been extremely important for demonstrating cell-autonomous circadian clocks in several organisms[16-23], and in screens identifying clock genes and modifiers[24-28]. Circadian reporters have also been used to assess rhythmicity in peripheral tissues and the impact of dynamic alterations in environmental conditions (food availability, lighting cycles) on peripheral oscillators, conducted by measuring bioluminescence rhythms in tissue explants monitored *ex vivo*[21,29-35]. These studies complement work done by assessing population rhythms in gene expression in tissue samples[36-42] indicating altered rhythm amplitude, phase, and phase relationships in and between SCN and peripheral oscillators following resetting[43]. More recent advances include development of methods for monitoring bioluminescence rhythms from the SCN *in vivo*[44-50], and for assessing peripheral rhythms in anesthetized[51-53] and ambulatory[54,55] mice.

Here, we report a new transgenic mouse line in which firefly luciferase is expressed from the mouse *Dbp* locus in a *Cre*-recombinase-dependent manner. *Dbp* is widely and rhythmically expressed[3,57], allowing detection of circadian bioluminescence rhythms in numerous tissues, *in vivo* and *ex vivo*. *Cre*-dependent bioluminescence in specific cell types revealed unexpected differences among SCN neuronal populations.

Furthermore, we observed transient misalignment between behavioral and hepatic bioluminescence

rhythms in freely moving mice subjected to a shift of the light-dark cycle or restricted food access.

Results

<u>Generation of a bifunctional reporter mouse.</u> CRISPR/Cas9 genome editing was used to introduce a

bifunctional reporter into the mouse *Dbp* locus (**Fig. 1**). The reporter consists of a T2A sequence (to allow

expression of separate proteins from a single transcript[56]), a destabilized, enhanced GFP (d2EGFP, hereafter

GFP) sequence flanked by loxP sites, and a codon optimised synthetic firefly *luciferase* (luc2 from

Photinus pyralis, hereafter luc). In the absence of *Cre* expression, DBP and GFP are expressed as separate

proteins. After CRE-mediated recombination, the floxed GFP is removed, and separate DBP and luciferase

proteins are expressed from the *Dbp* locus. Sequencing of genomic DNA confirmed successful generation

of the Dbp^{KI} conditional reporter allele.

<u>GFP expression from the Dbp^{KI} allele.</u> To examine expression of GFP from the conditional allele, $Dbp^{KI/+}$

mice (n=5-6 mice per time-point) were anesthetized and perfused with fixative at 4-h intervals over 24 h

(**Fig. S1**). Liver sections from $Dbp^{KI/+}$ and control (WT) mice were examined by confocal microscopy.

Fluorescence signal intensity did not differ between time-points (ANOVA $F_{5,26} =1.279$, $p = 0.7560$). GFP

signal from $Dbp^{KI/+}$ liver sections was 5-10x higher than from WT sections, but absolute levels were quite

low. The low level of GFP expression may be due to the use of destabilized GFP with a 2-hour half-life,

intended to more accurately track changes on a circadian time-scale. The relatively low level and lack of

detectable rhythmicity in GFP expression was unexpected, especially considering that liver is the tissue

with the highest levels of *Dbp* expression[57] and thus may represent a 'best-case' scenario. As the primary

objective of this project was to generate a mouse model with Cre-dependent expression of bioluminescence

from the *Dbp* locus, however, the absence of robust GFP-driven fluorescence rhythms in *Cre*-negative cells

did not preclude achieving this objective. GFP is effectively serving as a 'floxed stop' to make luciferase

expression from the *Dbp* locus exclusively *Cre*-dependent.

Non-conditional luciferase expression from the Dbp^{Luc} allele. A non-conditional reporter allele was
generated by breeding to combine the conditional Dbp^{KI} allele with *Cre*-recombinase expressed in the
germline, leading to germline excision of GFP. The resulting Dbp^{Luc} allele produces wide-spread, rhythmic
luciferase expression, both and *in vivo* and *ex vivo*. More specifically, explants of lung and anterior pituitary
gland from $Dbp^{Luc/+}$ mice incubated with D-luciferin had robust circadian rhythms in bioluminescence (**Fig.
2**). Furthermore, *in vivo* imaging of $Dbp^{Luc/+}$ mice at 7 time-points over a ~30-h period revealed rhythmic
bioluminescence in the abdomen and throat in ventral views, and in the lower back in dorsal views (**Fig.
3B**), similar to the distribution of bioluminescence signal from $Per2^{Luc}$ mice[51-53] (**Fig. 3A**). The level of
light output was ~2.5-fold greater in ventral views than in dorsal views (p<0.0001, Wilcoxon matched pairs
test, W=151, n=17). In the abdomen, we defined a rostral ("liver") region of interest (ROI) and a more
caudal "lower abdominal" ROI. The liver ROI accounted for 46.6 ± 3.0% (Mean ± SEM; n=17) of
bioluminescence from the ventral view, while the lower abdomen contributed another 38.4 ± 3.5%.
Bioluminescence rhythms from the throat region have previously been shown to originate in the
submandibular gland[51]. Bioluminescence was absent in mice with wild-type *Dbp* alleles or with the
conditional Dbp^{KI} allele (in the absence of *Cre*).

Post-mortem dissection and imaging revealed that tissues of the gastrointestinal tract (intestine,
cecum and colon), pancreas and mesenteric fat, perigonadal fat and uterus were major contributors to
overall light output, with liver and kidney emitting lower levels (**Fig. S2**). Organs contributing minor
amounts to total bioluminescence in dissected mice were esophagus, heart, lung, thymus, spleen, and testes.

Previous reports have shown that in a number of tissues, *Dbp* RNA levels peak earlier than *Per2*
RNA levels[3]. Consistent with this literature, the time of peak of bioluminescence rhythms from $Dbp^{Luc/+}$
tissues preceded the time of peak of bioluminescence rhythms from $Per2^{Luc/+}$ tissues by ~6 hours, both *in
vitro* (**Fig. 2C, 2F)** and *in vivo* (**Fig. 3G-3I).** Unexpectedly, bioluminescence rhythms from $Per2^{Luc/+}$ tissue
explants had significantly greater period length than explants from $Dbp^{Luc/+}$ mice (Lung: 25.29 ± 0.13 vs

23.93 $\pm$ 0.11 h; $F_{1,27.7}$ = 95.55, p < 0.0001; Anterior Pituitary: 25.27 $\pm$ 0.08 vs 23.73 $\pm$ 0.112 h; $F_{1,24.53}$ =

66.12, p < 0.0001).

Molecular and Behavioral Rhythms in Mice with *Dbp* Reporter Alleles. To confirm that the

introduction of the reporter construct into the *Dbp* locus did not alter circadian clock function, molecular

and behavioral rhythms were assessed. Male mice used for these analyses had either one or two copies of

the GFP-containing conditional allele (*Dbp*$^{KI/+}$ and *Dbp*$^{KI/KI}$, respectively), one or two copies of the

luciferase-expressing allele (*Dbp*$^{Luc/+}$ and *Dbp*$^{Luc/Luc}$, respectively), or were wild-type (WT) littermate

controls.

RNA was isolated from livers collected at 4-h intervals over 24-h. Northern blots were prepared

and probed for *Dbp* and *Actin* (loading control). As expected, the transcripts from *Dbp*^{KI} and *Dbp*^{Luc} alleles

migrated more slowly than the wild-type transcript (**Fig. 4A**), due to inclusion of GFP and luciferase coding

sequence in these transcripts, respectively, as verified by probing for reporter sequences in a replicate blot.

Peak levels of *Dbp* expression in liver occurred at ZT10 in all genotypes (**Fig. 4B**), as expected based on

previous studies[3,42,57]. For each transcript type, the *Dbp*/*Actin* ratios were ranked within each series of 6

timepoints. These ranks differed significantly among the timepoints for each transcript (Friedman's One-

Way analysis of variance, p < 0.002), and post-hoc testing indicated significantly higher rankings at ZT10

than at ZT2, ZT18 and ZT22 (Dunn's test, p < 0.05). These data indicate that the temporal profile of

transcript expression from the *Dbp* locus was unaffected by the inclusion of reporter sequences.

Heterozygous mice expressed both *Dbp* and *Dbp-plus-reporter* transcripts. The two transcript types

did not differ in abundance: optical density over film background of the *Dbp*^{KI} transcript was 100.5 $\pm$ 5.3 %

of the Dbp$^+$ transcript in *Dbp*$^{KI/+}$ mice (t=0.084, df=7, p= 0.94, one-sample t-test vs 100%), while the *Dbp*^{Luc}

transcript was 102.3 $\pm$ 5.0 % of *Dbp*$^+$ transcript in *Dbp*$^{Luc/+}$ mice (t=0.446, df=7, p=0.669). The equivalent

expression level of the two transcript types in heterozygous animals strongly suggests that transcript

regulation and stability were not altered by inclusion of reporter-encoding sequences.

Potential influences of the *Dbp* reporter alleles on locomotor activity rhythms were assessed in

constant darkness. Mice of the same five genotypes and both sexes were examined (**Table 1; Fig. S3**). This

assessment was complicated by a significant sex-by-genotype interaction ($F_{4,102} = 2.904$, $p = 0.0254$) that

post-hoc tests indicated was the result of an unexpected sex difference in the $Dbp^{Luc/Luc}$ mice. Indeed, when

this genotype was excluded from the analysis, no significant sex-by-genotype interaction was observed

($F_{3,88} = 1.349$; $p = 0.2636$) and one-way ANOVA demonstrated the absence of a significant main effect of

genotype ($F_{3,91} = 1.174$; $p = 0.3242$). One-way ANOVA within each sex with all five genotypes included

revealed no genotype effect in males ($F_{4,50} = 1.299$, $p = 0.283$). While there was a significant genotype

effect in females ($F_{4,52} = 2.716$, $p = 0.040$), Tukey HSD post-hoc tests did not find a significant result among

any of the pairwise genotype comparisons (all p values > 0.05). Similarly, an alternative post-hoc analysis

revealed that none of the other female genotypes differed from WT females in their free-running period in

constant darkness (Dunnett's test, p > 0.5 in each case). To further examine the effect of sex on free-running

period, males and females of each genotype were compared directly. In both homozygous reporter lines

($Dbp^{Luc/Luc}$ and $Dbp^{KI/KI}$), males had significantly longer period lengths than females ($p < 0.01$), while this

was not seen in wild-type controls or heterozygous reporters ($p > 0.46$). Together, these assessments of

molecular and behavioral rhythms indicate that the reporter alleles do not alter circadian function or change

Dbp expression.

***Cre*-dependent Luciferase Expression in Liver.** The main use we envision for the *Dbp* reporter alleles

involve *Cre* recombinase-mediated excision of GFP, leading to expression of *luciferase* in cells expressing

Cre. The effectiveness of this approach was first assessed in the liver. Hepatocytes were targeted using an

Albumin-Cre-driver line. *In vivo* bioluminescence imaging of intact *Albumin-Cre$^+$*; $Dbp^{KI/+}$ "liver reporter"

mice at the time of expected maximal bioluminescence revealed that $96.6 \pm 0.48\%$ of light originated in the

"liver" ROI (relative to total ventral-view bioluminescence; p<0.0001 versus $46.6 \pm 3.0\%$ in Dbp^{Luc} mice,

U-test, U=0, n=19 and 17, respectively). Light output from the ventral side was 5.14 ± 0.53 times greater

than from the dorsal view (p<0.0001, Wilcoxon matched pairs test, W=190, n=19). Notably, post-mortem

imaging of dissected parts confirmed that the bioluminescence signal originated exclusively from the liver

in these mice (97.4% of light from liver; n=12).

In a separate cohort of liver reporter mice, bioluminescence was assessed around the clock by IVIS

imaging. The cosinor-fitted time of peak of *Dbp*-driven bioluminescence rhythms from the liver 'region of

interest' of these mice (ZT11) was indistinguishable from the peak time of the liver ROI analyzed in whole-

body *Dbp^Luc* mice (**Fig. 3I**).

<u>Cell-type Specific Bioluminescence Rhythms in SCN Slices</u>. The heterogeneity of SCN neurons has

complicated our understanding of central clock function[58]. Neuromedin S (NMS) is expressed in ~40% of

SCN neurons, while Arginine Vasopressin (AVP) is expressed in ~10% of SCN neurons and is contained

within the NMS-expressing population[59]. The utility of our conditional reporter line was demonstrated by

monitoring bioluminescence rhythms within specific subpopulations of SCN neurons (**Fig. 5**). *NMS-iCre;*

Dbp^{KI/+} mice and *AVP-IRES2-Cre; Dbp^{KI/+}* mice were generated, and single-cell bioluminescence rhythms

were compared to those from non-conditional *Dbp^{Luc/+}* mice in SCN slices *in vitro*. For the conditional

mice, bioluminescence was apparent in subsets of cells within the SCN (**Fig. 5**). The anatomical pattern of

bioluminescence in the SCN differed based on the *Cre* line used, consistent with the expected distribution

for each neuronal subtype.

The cell-type specificity of bioluminescence signals from the different genotypes enabled the

assessment of rhythm quality in the different neural populations. This assessment revealed a significantly

shorter period in AVP$^+$ neurons compared to NMS$^+$ cells (**Fig. 5D**; $F_{2,14.64}$ = 4.259, p = 0.0345). Although

the time of peak of *Dbp*-driven bioluminescence did not differ significantly between the different cellular

populations examined (**Fig. 5E**; $F_{2,18.31}$ = 0.6570, p = 0.5302), a reduction in rhythm robustness was

observed in AVP$^+$ neurons compared to rhythms of NMS$^+$ neurons as well as compared to all neurons (**Fig.**

5F; $F_{2,18.11}$= 14.34, p = 0.0002). In line with this reduced robustness of individual cellular oscillators, the

distribution of peak times was also more dispersed in AVP$^+$ cells compared to NMS$^+$ neurons (**Fig. 5G**).

These results complement the recent report from Shan *et al* using a Cre-dependent Color Switch

PER2::LUC reporter mouse demonstrating period and phase differences among sub-populations of SCN

neurons (AVP[+] and VIP[+]), relative to the rest of the SCN[60]. Our *Dbp[KI]* mice and the recently reported Color-

Switch PER2::LUC[60] mouse line will be important additions to our molecular/genetic armamentarium for

unravelling the complicated relationships among the cellular components of the central circadian

pacemaker in the SCN[58-64].

Continuous, Non-invasive Detection of Bioluminescence Rhythms from Liver in Ambulatory Mice.

Addressing issues of internal desynchrony and misalignment of oscillators requires monitoring the

dynamics of tissue resetting over time after a phase-shifting stimulus. The use of *in vivo* bioluminescence

imaging for repeated assessments of organ-level regions of interest over multiple days is feasible[52] but

requires repeated, potentially disruptive anesthesia sessions[65] per circadian cycle for several days, and

intensive effort by investigators. As a result, *in vivo* bioluminescence imaging has generally been relegated

to assessing phase of reporter gene oscillations on relatively few occasions after a shifting stimulus, with

rare exception[52]. An attractive alternative is to perform long-term, non-invasive bioluminescence

recordings, as pioneered by Saini et al.[54] who administered a virally encoded luciferase reporter by tail vein

injection to transduce liver, allowing bioluminescence recording from awake, behaving mice using a

specialized detector unit. This virally mediated reporter method is only appropriate for assessing rhythms

in liver, however. Other studies have attached recording devices directly to mice or used fiber optics to

collect light from specific brain regions in broadly luminescent *Per1-luc* or *Per2-luc* mouse reporter lines[45-

50,66-68], which allow monitoring rhythms from tethered but 'freely moving' mice. An elegant but more

invasive approach involves administering *Cre*-dependent viral reporters to mice with cell- or tissue-specific

Cre expression, allowing specific cellular populations to be monitored *in vivo*[48]. A less invasive approach

that allows long-term assessment of rhythms in a variety of specific tissues is desirable.

In vivo bioluminescence imaging also suffers from a lack of anatomical resolution, with the light

from several abdominal organs potentially merging. Tissues of the gastrointestinal tract are a major source

of abdominal bioluminescence in "whole-body" *(Dbp^(Luc/+))* reporter mice, and these tissues likely

overshadow (or, more literally, out-glow) surrounding tissues, making it impossible to specifically assess

rhythmicity in smaller abdominal structures *in vivo*. Bioluminescence from even relatively large organs like

liver and kidney is likely 'contaminated' by light from the gastrointestinal tract in non-conditional reporter

mice.

To overcome these difficulties with assessing the origin of bioluminescence in "whole-body"

reporter mice, and to refine recently developed methods for long-term monitoring of peripheral rhythms in

ambulatory mice[55], we generated *Albumin-Cre*;*Dbp^(KI/+)* ("liver reporter") mice. First, we examined the

potential impact of route of substrate administration on rhythms using a Lumicycle *In Vivo* system[55]

(Actimetrics, Wilmette IL). Mice were entrained to 12L:12D followed by a skeleton photoperiod consisting

of 4 1-h pulses of light every 24 hr (1L:1D:1L:6D:1L:1D:1L:12D) with the 12-h dark phase coinciding

with 12-h dark phase of the preceding lighting cycle. A skeleton photoperiod was used because detection

of bioluminescence requires the absence of ambient light, while studies of light-induced phase shifting

obviously require light; a skeleton photoperiod is a compromise between these conflicting constraints. After

7 days in the skeleton photoperiod, mice were anesthetized for subcutaneous implantation of a primed

osmotic minipump (Alzet, Model #1002 (0.25µl per hour)) containing either D-luciferin (100 mM) or

phosphate buffered saline (PBS). Mice with PBS-containing pumps received D-luciferin in the drinking

water (2 mM). The time of peak of bioluminescence rhythms was determined 5 days after pump

implantation on the first day of exposure to constant darkness. Time of peak was determined by discrete

wavelet transform (DWT) analysis. There was no difference in time of peak for these routes of D-luciferin

administration (drinking water: mean peak time ($\pm$ SEM) CT 8.75 ± 0.20 (n = 7); osmotic minipumps: mean

peak time CT 8.76 ± 0.19; unpaired t-test, t = 0.0342, df =12, $p = 0.9733$). Thus, the presumed rhythm of

substrate intake, secondary to the rhythm of water intake, does not change the time of peak of the

bioluminescence rhythm from liver reporter mice. This is consistent with recent results from Sinturel et al.

2021[77]. Our subsequent studies used D-luciferin (2 mM) administered in the drinking water.

<u>Circadian Misalignment Following a Phase Shift of the Lighting Cycle</u>. The approach described above

provides an unparalleled system for assessing the timing of rhythmicity in a specific tissue over long periods

of time. Thus, hepatic bioluminescence rhythms were monitored in *Albumin-Cre*; *Dbp*^*KI/+* (liver reporter)

mice before and after a 6-hr phase advance of the skeleton lighting cycle described above. Control mice

remaining in the original (non-shifted) skeleton lighting regimen had a stable phase of hepatic

bioluminescence (**Fig. 6C**). In contrast, mice exposed to a phase-advance of the skeleton photoperiod

displayed a gradual phase-advance in both locomotor activity and hepatic bioluminescence rhythms (**Fig.**

6A, B). Notably, locomotor rhythms shifted more rapidly than hepatic bioluminescence rhythms: the liver

lagged behind (**Fig. 6B**). To compare the re-entrainment of bioluminescence and locomotor activity

rhythms, peak time for each rhythm each day was normalized to the time of peak on the last day before

shifting the lighting cycle in the shifted group (e.g., Day 2 in Fig. 6) for each animal. Data from each

lighting group were analyzed separately using a general linear model with Animal ID as a random variable

(allowing comparison of the two rhythms within individuals) and the main effects of the endpoint

(locomotor activity or bioluminescence) and Day number. In animals not undergoing a phase shift, the

phase relationship of these endpoints was unchanged over time (F < 1.1, p > 0.39). In contrast, in animals

exposed to a 6-hr phase advance of the skeleton photoperiod, the phase relationship of the locomotor

activity and bioluminescence rhythms differed significantly (Measure*Day interaction, $F_{9,54.98}$ = 3.358, p =

0.0024). Post-hoc testing revealed a significant difference in phase between the two measures on day 9

(Tukey HSD, p<0.05). A separate analysis to compare phase (relative to Day 2 baseline) between

bioluminescence and locomotor activity rhythms revealed significant differences between the two measures

on days 5,6,7,8,9 and 10 (t-tests on each day, p < 0.05). Thus, both locomotor activity and hepatic

bioluminescence rhythms shifted following a phase shift of the lighting cycle, but the rhythms differ in their

kinetics of re-adjustment, with the liver lagging behind. These data provide clear evidence for misalignment

of SCN-driven behavioral rhythms and rhythmicity in the liver.

Recovery from Circadian Misalignment Induced by Temporally Restricted Feeding. We next conducted a study to examine misalignment induced by restricted feeding, as previous studies have shown that food availability limited to daytime significantly alters phase of peripheral oscillators[35,54]. Due to our desire to study bioluminescence rhythms without interference from the light-dark cycle, our experiment assessed the timing of the *Dbp*-driven liver bioluminescence rhythms in constant darkness after different feeding regimens were administered in a light-dark cycle. This allowed us to determine the time of peak bioluminescence of the liver after restricted feeding, and the unprecedented opportunity to observe its return toward a normal phase relationship with SCN-driven behavioral rhythms over time in constant darkness with *ad libitum* food access.

Alb-Cre;Dbp[KI/+] liver reporter mice were exposed to one of three feeding regimes (*ad libitum*, nighttime, or daytime food availability; **Fig. 7A**) for ten days preceding bioluminescence recording in constant darkness under *ad libitum* feeding conditions. A previously described automated feeder system[69] was used to restrict food availability. This system limits total daily consumption (to prevent hoarding). With the setting used, this system restored daily food allotments to *ad libitum* fed and night-fed mice daily at 0000h (ZT18), and restricted food pellet delivery for day-fed mice to 0600-1800 h (ZT0-ZT12), and for night-fed mice to 1800-0600 h ZT12 – ZT24/0). This midnight food replenishment resulted in unusual temporal profiles of food intake in *ad libitum* and night-fed mice. Nevertheless, *ad libitum* and nighttime food access both resulted in food intake being concentrated in the night while daytime food availability resulted in the midpoint of daily food intake occurring during the first half of the light phase (**Fig. 7A, 7C**). Within-group variability in the timing of food intake was low except for three clear outliers (**Fig. 7C**) that were excluded from subsequent analyses.

Ad libitum fed mice showed consistently phased rhythms in bioluminescence after transfer

to constant darkness from 12h L:12h D, as did night-fed animals (**Fig. 7A, 7D**). In contrast, mice

fed only during the light period for 10 days prior to housing in DD with *ad libitum* food had an

earlier time of peak of the hepatic bioluminescence rhythm. Daytime feeding resulted in a

significantly advanced peak time compared to both night-fed and *ad libitum* fed mice, while these

latter groups were statistically indistinguishable ($F_{2,259.6} = 76.66$, $p < 0.0001$; **Fig. 7D**). Subsequent

exposure to constant darkness with *ad libitum* feeding allowed the hepatic clock of day-fed mice

to return toward the appropriate phase relationship with the locomotor activity rhythm regulated

by the SCN circadian pacemaker.

Although daytime feeding resulted in an advanced time of peak bioluminescence, the

timing of the liver bioluminescence rhythm was not solely controlled by the timing of food intake.

First, no significant correlations between the timing of food intake and time of peak

bioluminescence were observed within any of the three feeding regimes ($F < 1.13$, $p > 0.32$; **Fig.**

7C). Second, the relationship between the timing of liver bioluminescence rhythms relative to the

midpoint of food intake was significantly different between the different groups ($F_{2,17} = 313.2$, p

< 0.0001; **Fig. 7E**). While *Dbp*-driven hepatic bioluminescence rhythms were roughly in anti-

phase with the midpoint of feeding in *ad libitum* and night-fed mice, daytime feeding resulted in

near synchrony between these different rhythms (**Fig. 7E**). Furthermore, although the average

midpoint of feeding was significantly earlier in night-fed compared to *ad libitum* fed mice ($t_{10} =$

6.21, $p < 0.0001$; **Fig. 7C**), no significant difference was observed in bioluminescence phase

relative to the preceding light-dark cycle (**Fig. 7D**), with the timing of liver bioluminescence

rhythms relative to the midpoint of food intake being significantly delayed in night-fed compared

to *ad libitum* fed mice (**Fig. 7E**). Overall, these results demonstrate that although the timing of

food intake strongly influences liver rhythms, the timing of bioluminescence rhythmicity in liver

reporter mice is not solely driven by the timing of food intake (with food intake regulated for this

duration and in this way).

Discussion

Numerous studies have made use of rhythmically expressed bioluminescent reporter genes

to monitor circadian rhythms. The *Per2^Luc* mouse has been especially useful as it generates robust

bioluminescence rhythms from numerous tissues *in vitro*. The widespread expression of

PER2::LUC (and other 'non-conditional' bioluminescence reporters) comes at a cost, however, as

it is not possible to assess rhythmicity in specific cell populations within a larger tissue without

dissection. Tissue explant preparation can cause phase-resetting, however, especially after

exposure to phase shifting stimuli[70,71]. Furthermore, *ex vivo* culturing of tissues does not allow

assessment of rhythmicity in the context of the hierarchical circadian system or dynamic changes

during environmentally-induced resetting.

We chose to modify the *Dbp* gene to generate a conditional reporter for several reasons.

Dbp is widely and rhythmically expressed at readily detectable levels in numerous tissues[3,57]. This

feature ensures that the reporter mouse would be useful for detecting rhythmicity in numerous

tissues. In addition, individual clock genes are responsive to different signaling pathways. This

differential regulation can lead to circadian misalignment *within* the circadian clock[39,43]. As an

output gene, *Dbp* rhythmicity is likely a good proxy for the integrated output of the molecular

clockwork in total. Finally, concern that the targeting event could disrupt function of the modified

gene led us to steer away from core clock genes. For example, the GFP-expressing *Dbp* transcript

lacks the native 3′ UTR and uses an exogenous polyadenylation sequence, which could affect *Dbp*

gene expression and regulation. Notably, however, our Northern blot analysis suggests little or no

alteration in expression level or dynamics of the reporter transcripts; an observation in line with

the previous finding that mice homozygous for a targeted allele of *Dbp* have only a modest

circadian phenotype[72].

Shan et al.[60] recently reported development of a Color-Switch PER2::LUC line which they

used to demonstrate the utility of a Cre-dependent reporter approach for interrogating SCN

circuitry. The Color-Switch PER2::Luc line has the advantage of reporting on both Cre-positive

and Cre-negative cells in different colors. This strength of the Color-Switch PER2::LUC reporter

can simultaneously be a weakness, in that detection of bioluminescence requires segmentation of

the bioluminescence signal between wavelengths. Our 'simpler' approach of only inducing a

bioluminescence signal in *Cre*-positive cells of $Dbp^{KI/+}$ mice enables recording of bioluminescence

rhythms without the need for wavelength segmentation. In addition, the *Dbp* reporter can easily

be used in *Per2* mutant mice. Like the Color-Switch PER2::LUC line, our *Dbp* conditional reporter

line is useful for *ex vivo* studies, allowing specific cellular populations to be monitored by crossing

to the appropriate *Cre*-expressing lines. Our studies reveal subtle differences among the population

of oscillators defined by AVP-*Cre*, NMS-*Cre*, and the entire SCN cohort. More specifically, AVP

cells had a shorter period, reduced rhythmicity index, and larger within-slice dispersal of peak

times than the NMS cell population with which it overlaps. Our results suggest that AVP cells are

coordinated less well and are less robust, in contrast to the typical view of AVP cells as high-

amplitude 'output' neurons that also contribute to determination of period and rhythm amplitude[58].

We envision this line being very useful for monitoring additional neuronal subpopulations

in the SCN in wild-type and mutant animals. Additional technical development may allow *in vivo*

detection of bioluminescence rhythms from neuronal populations in awake behaving mice.

Approaches to optimize the signal detected from brain include use of highly efficient and cell- and

brain-penetrant substrates[73,74], cranial windows[75] and hairless or albino mice[55,74]. These

approaches may allow interrogation of the SCN circuit *in vivo*, extending the elegant studies being

performed with SCN slices *in vitro*. Bioluminescence rhythms can also be examined in neuronal

populations outside the SCN, by using an appropriate *Cre* driver and/or viral delivery of *Cre*

recombinase.

In vivo bioluminescent imaging allows assessment of bioluminescence from several organs

in vivo, but the signal from these areas likely includes light emitted from nearby organs (e.g.,

intestinal tract and abdominal fat likely contribute to the signal attributed to liver and kidney).

Indeed, the size and shape of the "liver" ROI seen by IVIS imaging differs between *Dbp* liver

reporter mice and whole-body reporter *Dbp^{Luc}* mice. Furthermore, bioluminescence from nearby

tissues can be obscured by the high level of light output coming from liver, kidney and intestines.

Cre-mediated recombination of the conditional *Dbp* reporter allele thus enables assessment of

bioluminescence from other tissues.

Cre-mediated recombination of the *Dbp^{KI}* allele in liver enabled us to perform continuous,

in vivo bioluminescence monitoring of liver in freely moving mice. These studies demonstrate

transient misalignment between the liver oscillator and SCN-regulated behavioral rhythms.

Repeated misalignment among oscillators is thought to contribute to adverse metabolic and health

consequences of chronic circadian disruption. Up until now, technical and practical limitations

have restricted our ability to monitor the behavior of circadian rhythms in different peripheral

tissues during and following environmental disruption of circadian homeostasis. Our *Cre*-

conditional reporter line and the approaches described here for assessing bioluminescence rhythms

in vivo will allow longitudinal and tissue-specific characterization of misalignment and recovery

after a variety of circadian-disruptive lighting and food availability paradigms, allowing more

extensive testing of the consequences of repeated misalignment of peripheral clocks.

434   Previous studies have shown misalignment between central and peripheral clocks induced

by altering the time of food access to daytime, by assessing oscillator phase at various time-points

after a phase shift of the lighting cycle, or by exposure to non-24hr light-dark schedules. The vast

majority of these studies monitored bioluminescence rhythms *ex vivo* or assessed transcript levels

following tissue collection at various times after a shift[29-36,37-41,69]. Notably, *ex vivo*

bioluminescence rhythm timing may be affected by prior lighting conditions[70,71]. Few studies have

followed bioluminescence rhythms *in vivo* over time after a light-induced phase shift or after a

food manipulation that phase-shifts peripheral oscillators[52,76]. Our current data leverage the

unprecedented ability to non-invasively monitor rhythmicity from a peripheral oscillator in

individual animals over many days to show the time course of internal misalignment and recovery

after a phase shift. Other studies with minimally invasive monitoring of bioluminescence rhythms

have relied upon viral introduction of the reporter into liver, and thus are limited to studies of

liver[54,77]. Moreover, efficient expression of virally delivered reporter constructs is limited by the

promoter size and specificity. Future studies of additional tissues in *Cre*-conditional reporter mice

will enable a full deciphering of how other components within the hierarchical, multi-oscillatory

circadian system respond to disruptive stimuli. Several studies suggest that organs differ in their

response to resetting stimuli. For example, the *Dbp* mRNA rhythm in liver is more fully reset than

the rhythm in heart and kidneys 3 days after restricting food availability to daytime[38], and several

studies indicate the SCN (and the locomotor rhythms it regulates) reset more rapidly than

peripheral tissues[21,30-33,38,43,50,52,54].

A further advance in studying the behavior of peripheral oscillators is provided by the

ability to temporally regulate the timing of feeding. A recent study[76] used a feeding device similar

to the one used here to recapitulate food intake patterns in mice with restricted food access that

were more naturalistic than the severe 'all or none' patterns typically used in studies with time-

restricted access to food. The authors found that peripheral oscillators of *Per2*[Luc] mice were not

effectively entrained by these imposed 'natural' feeding patterns. Similarly, our restricted food

access study produced a smaller and more variable phase shift of the hepatic circadian clock (as

indicated by the initial time-of-peak of *Dbp*-driven bioluminescence) than expected based on

published results using presence / absence food availability cycles. A longer period of adjustment

to the restricted feeding schedule or more complete and abrupt transitions between food presence

and absence would likely produce a stronger entraining signal for the liver. Use of a variety of

different *Cre* drivers will allow assessment of whether different peripheral organs respond

similarly to food restriction paradigms. In addition, tissue-specific reporter models will be very

useful in assessing how more naturalistic food ingestion paradigms influence peripheral circadian

clocks in several tissues. These approaches will characterize circadian organization in normal and

disruption paradigms, helping to identify factors contributing to the adverse consequences of

circadian disruption.

Materials and Methods

Animals and Housing Conditions

All animal procedures were reviewed and approved by the Institutional Animal Care and Use

Committees of the University of Massachusetts Medical School, Morehouse School of Medicine, the

University of Warwick, and/or Smith College.

Unless otherwise noted, animals were maintained in a 12h light: 12h dark (LD) lighting cycle with

access to food (Prolab Isopro RMH3000; LabDiet) and water available *ad libitum*. Zeitgeber Time (ZT)

refers to time relative to the lighting cycle. ZT 0-12h is the light phase and ZT 12-24h is the dark phase.

Cre recombinase-expressing lines were obtained from the Jackson Labs and were crossed to mice

bearing the conditional (*Dbp^KI*) reporter allele to generate mice expressing luciferase in specific cells or

tissues. The following tissue-specific *Cre* lines were used:

Albumin-Cre (B6.Cg-*Speer6-ps1^Tg(Alb-Cre)21Mgn*/J ; JAX 003574), *AVP*-IRES2-*Cre* (B6.Cg-*Avp^tm1.1(Cre)Hze*/J;

JAX 0023530), and *NMS-Cre* (Tg(Nms-iCre)[20Ywa], JAX 027205). These lines direct Cre recombinase

expression to hepatocytes, neurons expressing arginine vasopressin (AVP), and neurons expressing

neuromedin S (NMS), respectively. In addition, a *Prrx1-Cre* female (B6.Cg-Tg(Prrx1-Cre[1Cjt]/J), JAX

005584) was used for germline deletion of the conditional allele (see below). All *Cre*-expressing lines were

on the C57BL/6J background.

Founder *Per2^Luc/+* mice with an in-frame fusion of firefly luciferase to PER2, and an SV40

polyadenylation signal[19,78] were generously provided by Dr. Joseph Takahashi, University of Texas

Southwestern Medical School, Dallas. This line was maintained by backcrossing to C57BL/6J mice (JAX

000664). All *Per2^Luc* reporter mice used for experiments here were heterozygous (e.g., *Per2^LucSV/+*).

CRISPR/Cas9 targeting the *Dbp* locus

The mutant allele was generated by CRISPR/Cas9 mediated engineering of the *Dbp* locus. The targeting

construct consisted of a 5′ homology arm terminating just 5′ of the *Dbp* stop codon followed by in-frame

sequences encoding a T2A linker, LoxP, GFP with the bovine growth hormone polyadenylation signal,

LoxP, and Luc2 followed by the 3′-UTR of *Dbp* (3′ homology arm). Two sets of injections were done. In

the first injection set, which led to 17 mice, C57BL/6J blastocysts were microinjected with the donor

construct,   and   20   ng/ul   of   each   of   two   guide   RNAs   (MmDBPki_gR49f   5′

GCCCAGCAUGGGACACUGUG 3′ and MmDBPki_gR69f 5′ AGGCCACCUCCACCCUGCCA 3′).

This set of injections did not lead to any putative founders. For the second set of microinjections, blastocysts

were injected with 40 ng/ul guide RNA MmDBPki_gR49f, 50 ng/ul *Cas9* mRNA (synthesized from a Cas9

PCR product using mMessage mMachine T7 Ultra Kit from Life Technologies) and 20 ng/ul CAS9 protein

(IDT). From 34 mice generated in this set, two putative founders were identified using a primer pair internal

to the construct (primer pair C, see **Table S1**). Primer pairs consisting of a primer in flanking DNA, external

to the construct, and a primer within the construct were used to confirm that one of these animals had the

desired targeting event (primer pairs F and H, which spanned the 5′ and 3′ ends, respectively). Genomic

DNA from this mouse was then amplified using a primer pair flanking the entire construct; sequencing the

product confirmed the construct was inserted properly, *in vivo*. The founder carrying the targeted (knock-

in or Dbp^{KI}) allele and its offspring were backcrossed to C57BL/6J mice (JAX 000664) for three generations

before any intercrossing to reduce off-target mutations.

To generate mice with germline deletion of GFP (and thus leading to expression of luciferase

throughout the body), a male $Dbp^{KI/+}$ was bred to a *Prrx1-Cre* female (JAX 005584), which we had on hand

and which produces germline deletion of floxed alleles at high frequency. Several mice bearing the newly

generated Dbp^{Luc} allele were identified and backcrossed to C57BL/6J, selecting against *Prrx1-Cre*.

Genotyping

Genotyping was performed by PCR amplification of DNA extracted from ear punches.

Amplification products were separated by agarose gel electrophoresis. Genotyping protocols for *Cre*

recombinase and $Per2^{Luc}$ have been published previously[53, 79]. A primer set ("4A") capable of detecting all

possible *Dbp* allele combinations was used for colony genotyping; the three possible alleles (Dbp^+, Dbp^{KI},

Dbp^{Luc}) generate amplicons of different size with this primer set (299, 399 and 490 bp, respectively). Primer

sets are listed in **Table S1**.

Generation of Digoxigenin (DIG) DNA Probes and Northern Blot Assay

DIG-labeled DNA probes were generated by PCR in reactions containing 28 μM of DIG-labeled

UTP following the manufacturer's protocol (Roche). Primer sets are listed in **Table S1**.

WT, $Dbp^{KI/+}$, $Dbp^{KI/KI}$, $Dbp^{Luc/+}$, and $Dbp^{Luc/Luc}$ mice were euthanized by Euthasol injection and liver

tissue was collected at 4-h intervals (ZT 2, 6, 10, 14, 18, 22). RNA was isolated from the liver tissue by

Trizol extraction (Ambion). RNA was quantitated by Nanodrop. Five micrograms per lane were separated

by electrophoresis on a 1.2% formaldehyde gel. RNA was transferred to a nylon membrane and cross-

linked by UV exposure. Blots were prehybridized, probed and detected following the manufacturer's

protocol (Roche), bagged and exposed to X-ray film.

Locomotor Activity Rhythms

Male and female mice of five genotypes (WT, $Dbp^{KI/+}$, $Dbp^{KI/KI}$, $Dbp^{Luc/+}$, and $Dbp^{Luc/Luc}$) were

transferred to the experimental room and single-housed with a running wheel. Animals had access to food

and water *ad libitum*. Running-wheel activity was monitored using ClockLab collection software

(Actimetrics). Mice were entrained to a 12-h light/12-h dark cycle for 18 days, then were placed into

constant darkness (dim red light) for 15 days. The free-running period for each animal was determined on

days 4-15 in DD by periodogram analysis using ClockLab analysis software.

Bioluminescence Recordings from Tissue Explants

Tissue explants were prepared late in the afternoon from $Per2^{Luc/+}$ and $Dbp^{Luc/+}$ mice housed on a

12-h light/12-h dark lighting cycle. Mice were deeply anesthetized with Euthasol and decapitated. Tissues

were dissected and immediately placed in ice-cold 1X HBSS (Gibco). Pituitary gland was subdivided into

4 sections (~2mm^3) with a scalpel and each piece was cultured separately. Lung explants were placed three

per dish. Up to three replicate dishes were studied per tissue per animal. Explants were placed on sterile

35-mm Millicell culture plate inserts (Millipore) in a sealed petri dish containing air-buffered

bioluminescence medium [53,80] plus D-luciferin (100 μM) (Gold Biotechnology) and incubated at 32 °C.

Bioluminescence was measured from each dish for 1 minute every 15 minutes using a Hamamatsu LM-

2400 luminometer.

Bioluminescence records were analyzed to determine period and peak time. The first 12-h were

discarded to exclude acute responses to explant preparation. Photon counts were smoothed to a 3-h running

average and baseline subtracted using a 24-h running average. Circadian period was determined from the

average of the period between each peak, trough, upward crossing and downward crossing between 24 and

88 hr of recording for each record. Peak time was calculated as the clock time of the first peak in the

background-subtracted data. Tissues from mice of the two genotypes were studied together in each run.

Imaging of *In Vivo* Bioluminescence Rhythms

In vivo imaging was performed in the UMass Medical School Small Animal Imaging Core Facility

using an In Vivo Imaging System (IVIS-100, Caliper, now Perkin Elmer) as previously described[52,53]. *Alb-*

Cre$^+$; Dbp$^{KI/+}$, *Dbp$^{Luc/+}$*, and *Per2$^{Luc/+}$* mice were anesthetized with 2% isoflurane (Zoetis Inc.) and skin

covering the liver, kidneys and submandibular glands was shaved. Mice were injected with D-luciferin (i.p.,

100 µl at 7.7 mM, Gold Biotechnology) and Dorsal (9 min post-injection) and Ventral (10.5 min post-

injection) images were captured from each animal. To assess bioluminescence rhythms, anesthesia, D-

luciferin injection and imaging was repeated at 4- to 8-hour intervals over approximately 30 hours.

Experiments to localize the source of bioluminescence utilized a single injection of D-luciferin at the time

of the expected peak bioluminescence, followed by euthanasia, dissection, and *ex vivo* imaging.

IVIS images were analyzed using Caliper Life Sciences' Living Image software (version 4.4).

Bioluminescence within Regions of Interest (ROI) of fixed size for all time-points in each mouse was

determined and analyzed as previously described[53].

Assessing Distribution of Bioluminescence *In Vivo* and *Ex Vivo*

Whole-body reporters (*Dbp$^{Luc/+}$*) and liver reporters (*Alb-Cre+;Dbp$^{KI/+}$*) were used to assess the

distribution of bioluminescence by IVIS imaging. Mice were anesthetized with isoflurane, shaved, and

injected with D-luciferin (100 microliters at 7-10 mM, i.p.) at times of peak expression (ZT 11-16). Images

were captured of ventral and dorsal views at 9-12 minutes after injection.

Bioluminescent counts within regions of interest (ROIs) were calculated using Living Image 4.4

software. ROIs identified on the ventral surface were the whole rectangular region containing the mouse,

and sub-ROI's where a region in the throat (submandibular gland), upper abdomen, and lower abdomen,

and any other hot-spots observed. Dorsal ROI's were the rectangle containing the entire mouse and a sub-

ROI over the lower back, corresponding to the abdomen on the dorsal side. Subsequent calculations were

performed in Microsoft Excel.

Animals were euthanized while under anesthesia, and organs were dissected and imaged to assess

the distribution of light. Due to the time required for sequential dissection and image capture of multiple

animals, some images of dissected tissues were captured as long as 60 minutes after D-luciferin injection,

and up to 10 minutes after euthanasia. Preliminary studies conducted by capturing images at various time

after euthanasia suggested there is no change in the distribution of bioluminescence with time, although

absolute levels fall with time. These studies indicated significant contribution of bioluminescence from

small intestine, colon, cecum, pancreas and mesenteric fat, uterus and perigonadal fat, with liver and kidney

emitting less light. Thymus, heart, lung and spleen contributed very little light output under the imaging

conditions used.

Bioluminescence Imaging of SCN Explants

Coronal sections containing SCN from *NMS-Cre;Dbp$^{KI/+}$* , *AVP-IRES-CRE;Dbp$^{KI/+}$*, and *Dbp$^{Luc/+}$*

mice were dissected, cultured, and imaged as previously described[81,82]. Briefly, sections containing SCN

(150 μm) were collected from adult mice, cultured on a membrane (Millicell CM; Millipore) in 1.2 mL of

air-buffered media containing 100 μM D-luciferin (Gold Biotechnology), and imaged for 5 days using a

Stanford Photonics XR/MEGA-10Z cooled intensified charge-coupled device camera.

Rhythmic parameters of luciferase expression were calculated for each slice and for cell-like

regions of interest (ROIs) within each slice using computational analyses in MATLAB (R2018a,

MathWorks) as described previously[82,83]. Briefly, to locate and extract data from cell-like ROIs, we

employed an iterative process identifying clusters of at least 20 bright pixels after background and local

noise subtraction (through application of a 2D wavelet transform using Wavelab 850,

(https://statweb.stanford.edu/~wavelab/) of a slice image summed across 24 h of bioluminescence. To

extract time series for the ROI's, each image in the sequence was smoothed via convolution with a Gaussian

kernel applied to 12x12-pixel regions and reduced from 512x640 resolution to 256x320. A discrete wavelet

transform (DWT) was applied to each time series to remove the trend and to extract the circadian and noise

components using the *wmtsa* toolbox for MATLAB (https://atmos.uw.edu/~wmtsa/). The criteria for

circadian rhythmicity in the ROI time series were a peak autocorrelation coefficient of at least 0.2, a

circadian component peak-to-peak time between 18 and 30 h, an amplitude above baseline noise (standard

deviation of noise component), and a cross-correlation coefficient of at least 0.4 with an aligned sine wave

over a 48h window. Peaks of the DWT circadian component were used to estimate peak time of each ROI.

Rhythmicity index (RI) is the peak in the autocorrelation of the DWT-detrended time series,

corresponding to a lag between 16 and 36 hrs, as previously described[84,85]. The time of peak

bioluminescence, rhythmicity index and the scatter of peak times within each slice for each ROI was

assessed on the first day *in vitro*. Period of rhythmicity in each ROI was determined as the average peak-

to-peak interval in the second and third cycles. These measures were compared between genotypes by a

general linear model, with slice ID included as a random variable to account for multiple cells being

measured on each slice. Where applicable, post-hoc comparisons were performed using Tukey's HSD

pairwise comparisons.

Bioluminescence Rhythms in Ambulatory Liver Reporter Mice

Bioluminescence was measured in freely moving *Alb-Cre^{+} ; Dbp$^{KI/+}$* reporter mice with the

"Lumicycle *In Vivo*" system (Actimetrics, Wilmette, IL) using methods as recently described[55]. Each unit

contained two PMTs (Hamamatsu H8259-01), and programmable LED lights. A programmable shutter

blocked the PMTs during periods of light exposure and to measure 'dark counts'. Each 1-minute dark-count

value was subtracted from the counts recorded during the subsequent 14 minutes to obtain the background-

corrected count values, to compensate for the effect of temperature fluctuations on PMT signal. Locomotor

activity was recorded using passive infrared motion sensors (Visonic, K940) and Clocklab software

(RRID:SCR_014309). Animals were checked daily at varied times using an infrared viewer (Carson

OPMOD DNV 1.0), or goggles (Pulsar Edge Night Vision Goggles PL75095).

Analysis of Ambulatory Bioluminescence Rhythms. Ambulatory bioluminescence data were

analyzed using RStudio. A discrete wavelet transform (DWT) was applied to each time series to detrend

and to calculate the time of peaks using the wmtsa R package (https://cran.r-

project.org/web/packages/wmtsa/index.html), as described[83-85]. The S12 filter was applied on 15-min

median binned data; medians were used (instead of means) to reduce the effect of large outliers. Data before

the first trough and after the last trough were discarded to avoid edge effects.

Assessing Routes of Administration of Luciferin. To determine whether rhythmic substrate

intake influences the pattern of bioluminescence, we compared the time of peak bioluminescence between

animals receiving continuous administration of substrate (from a subcutaneous osmotic minipump) with

trials in which mice received D-luciferin in the drinking water (2 mM) and implantation of a PBS-filled

osmotic pump.

Liver reporter mice previously housed in 12L:12D were entrained to a skeleton photoperiod (SPP)

consisting of four 1-hour light pulses. A skeleton photoperiod provides additional periods of darkness in

which to record bioluminescence. In the SPP, illumination occurred in four 1-hour blocks within the light

phase in the preceding lighting cycle (e.g., lights were on from ZT 0-1, 2-3, 9-10, and 11-12, so the first

and last hours of light in SPP coincided with light onset and offset, respectively, in the full photocycle with

lights on ZT0-12 and lights off ZT12-24/0). On the seventh day of SPP entrainment, mice were given

analgesics (0.05 mg/kg Buprenorphine and 2.0 mg/kg Meloxicam), anesthetized with 3% isoflurane, shaved

from hips to shoulders, and a primed osmotic minipump (Alzet Model #1002, 0.25µl per hour, 14 day)

containing D-luciferin (100 mM dissolved in PBS) or PBS vehicle was implanted subcutaneously. Mice

were returned to their cages with a warming disc and were provided soft food during the first 24 hours of

recovery. Animals were placed into the LumiCycle *In Vivo* unit 2.5 days after surgery. Bioluminescence

was recorded in SPP lighting for 2.5 days, then lights were disabled at the time of lights-out. The time of

peak bioluminescence was determined by wavelet analysis on the first day in constant darkness. No

difference in peak time of bioluminescence was found (see Results); in subsequent studies we administered

D -luciferin (2 mM) in the drinking water.

Re-entrainment following a Phase Shift of the Skeleton Photoperiod. Additional studies were

conducted to assess re-entrainment of the bioluminescence rhythms after a 6-hr advance of the light-dark

cycle. Animals previously entrained to a full 12L:12D lighting cycle were transferred to the skeleton

photoperiod described above for several days before study. Mice were anesthetized with isoflurane and

shaved 2.5 days prior to placement in the LumiCycle *In Vivo* units. D-Luciferin (2 mM) was provided in

the drinking water. Skeleton photoperiod lighting conditions were either maintained at the initial pattern or

advanced by 6 hr after the second day of recording. Locomotor activity was detected by passive infrared

motion sensors.

Analysis. The circadian time of peak bioluminescence and the mid-point of locomotor activity

were determined by wavelet analysis on each day of recording. For studies of phase shifts in response to

shifting the skeleton photoperiod, the timing of bioluminescence rhythms and locomotor activity rhythms

were normalized relative to the timing of these rhythms on Day 2 (e.g., the last day before shifting the

lighting cycle in the shifted group) for each animal. Data are expressed as mean ± SEM for each genotype

and endpoint on each day. Data from each lighting group were analyzed separately using a general lineal

model with Animal ID as a random variable (allowing comparison of the two rhythms within individuals)

and the main effects of the endpoint measure (locomotor activity or bioluminescence) and Day number,

and the 2-way interaction Measure*Day. In animals not undergoing a phase shift, potential changes in the

timing of the locomotor or bioluminescence rhythm were assessed separately for either measure by testing

the influence of Day number.

Food Restriction Followed by Bioluminescence Recording. Liver reporter mice (*Albumin-Cre;*

Dbp$^{KI/+}$) were fed pellets (300 mg, Dustless Precision Pellets, Rodent, Grain-Based, F0170, BioServ,

Flemington, NJ, USA) through the Actimetrics timed feeding apparatus designed by Phenome

Technologies, Skokie, IL, USA. Pellets were spaced by a minimum of 10 minutes to prevent hoarding

behavior[69]. Three groups were studied: those with *ad libitum* access to food, those with feeding restricted

to the light phase of the LD cycle (daytime feeding), and mice with access to food restricted to the dark

phase of the LD cycle (nighttime feeding). Mice were held under their feeding regime for 10 days prior to

bioluminescence recording. They were weighed regularly to ensure body weight did not decrease below

95% of initial weight. All mice were kept on a 12L:12D lighting schedule during the period of food

manipulation, and then were released into constant darkness for bioluminescence recording. During the LD

period, data were collected on feeding, light levels, and locomotor behavior (using motion sensors). Three

days before entering the Lumicycle In Vivo units, cage bottoms were changed at dark onset. *Ad libitum* and

night-fed mice were placed into the LumiCycle In Vivo units at dark onset with food immediately available.

Day-fed mice were placed into the LumiCycle In Vivo units at dark onset but were provided food after 12

hours (at the time of light onset in the previous LD cycle) to continue the daytime feeding regime during

the first day of the recording period. Bioluminescence was recorded for 7 days.

Liver reporter animals were randomly assigned to treatment groups and recording boxes.

Experimental groups and controls ran in parallel over five cohorts lasting 3 months. 24 hours prior to

placement in the recording boxes, mice were shaved from hips to shoulders on their front and back under

3% isoflurane and returned to their cages. 6 hours prior to placement into the in vivo boxes, mice were

provided with D-luciferin (2mM) in the drinking water to enable instantaneous bioluminescence upon

recording onset.

Analysis. For each animal the center of gravity (COG) of food intake was calculated for the last 5

days of the feeding regimen. Food intake patterns were also independently assessed qualitatively by four

observers. These assessments led to identification of three cohorts of mice, based on food intake patterns.

Three mice were identified as clear outliers compared to these three cohorts based on visual inspection of

the food intake timing. In line with this qualitative assessment, the feeding COG of each of these 3 animals

was >2 h removed from the other animals in their cohort (**Fig. 7C**). These three animals were excluded

from cohort-based assessments. Peak of bioluminescence on each day was calculated by DWT analysis as

above. Missing data resulted from inability to define a time of peak on some days. Hair regrowth contributed

to loss of signal and loss of rhythm amplitude, and thus to missing data in some cases.

Acknowledgments

We thank Christopher Lambert and Jamie Black for technical assistance, and Steven A. Brown (University of Zurich) for discussions and encouragement in the development of this project.

Use of UMass Medical School core facilities (Mutagenesis Core, Mouse Modeling Core, and Small Animal Imaging Core) is gratefully acknowledged.

Research reported in this publication was supported by the National Institute for Neurological Diseases and Stroke and the National Institute of General Medical Sciences of the National Institutes of Health under award numbers R21NS103180 (DRW), SC1GM112567 (AJD), and NIGMS R15GM126545 (MEH), the Hartmann Müller Stiftung (RD), MRC MC_PC_15070 (RD) and BSN (RD and LAG). CBS was a participant in the UMass Medical School Initiative for Maximizing Student Development, supported by NIH grant R25GM113686. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or the other funding agencies.

Author Contributions

R.D and D.R.W. conceived the project

C.B.S., V.v.d.V., E.M., M.H.B., A.J.D., M.E.H., R.D. and D.R.W. designed research

C.B.S., V.v.d.V., E.M., A.C.S., B.M.B., P.C.M., L.A.G., R.D., and D.R.W. performed research

C.B.S., V.v.d.V., E.M., T.L.L., B.M.B., M.E.H., R.D. and D.R.W. analyzed data

C.B.S., V.v.d.V., and D.R.W. wrote the paper

All authors have approved this version of the manuscript.

References

1. S. M. Reppert, D. R. Weaver, Coordination of circadian timing in mammals. *Nature* **418**, 935–941

(2002).

2. J. A. Mohawk, C. B. Green, J. S. Takahashi, Central and peripheral circadian clocks in

mammals. *Annu Rev Neurosci* **35**, 445–462 (2012).

3. R. Zhang, N. F. Lahens, H. I. Ballance, M. E. Hughes, J. B. Hogenesch, A circadian gene expression

atlas in mammals: implications for biology and medicine. *Proc Natl Acad Sci U S A* **111**, 16219–

16224 (2014).

4. A. Patke, M.W. Young, S. Axelrod, Molecular mechanisms and physiological importance of

circadian rhythms. *Nat Rev Mol Cell Biol* 21(2): 67-84 (2020).

5. J. A. Evans, A. J. Davidson, Health consequences of circadian disruption in humans and animal

models. *Prog Mol Biol Transl Sci* **119**, 283–323 (2013).

6. D.M. Arble*, et al*., Impact of sleep and circadian disruption on energy balance and diabetes: A

summary of workshop discussions. *Sleep* **38**, 1849-1860 (2015).

7. A. C. West, D. A. Bechtold, The cost of circadian desynchrony: Evidence, insights and open

questions. *Bioessays* **37**, 777–788 (2015).

8. T. Roenneberg, M. Merrow, The circadian clock and human health. *Curr Biol* **26**, R432-43 (2016).

9. J.D. Johnston, J.M. Ordovás, F.A. Scheer, F.W. Turek, Circadian rhythms, metabolism, and

chrononutrition in rodents and humans. *Adv Nutr* **7**, 399-406 (2016).

10. E. N. C. Manoogian, A. Chaix, S. Panda, When to eat: The importance of eating patterns in health

and disease. *J Biol Rhythms* **34**, 579–581 (2019).

11. L. R. Wegrzyn, *et al.*, Rotating night-shift work and the risk of breast cancer in the Nurses' Health

Studies. *Am J Epidemiol* **186**, 532–540 (2017).

12. A.M. Ramsey, A. Stowie, O. Castanon-Cervantes, A.J. Davidson, Environmental circadian

disruption increases stroke severity and dysregulates immune response. *J Biol Rhythms* **35**, 368-376

(2020).

13. W.H. Walker,2nd, J. C. Walton, A. C. DeVries, R. J. Nelson, Circadian rhythm disruption and

mental health. *Transl Psychiatry* **10**, 28 (2020).

14. T. Papagiannakopoulos, *et al.*, Circadian rhythm disruption promotes lung tumorigenesis. *Cell*

Metab **24**, 324–331 (2016).

15. E. Hadadi, *et al.*, Chronic circadian disruption modulates breast cancer stemness and immune

microenvironment to drive metastasis in mice. *Nat Commun* **11**, 3193-020-16890–6 (2020).

16. A. J. Millar, S. R. Short, N. H. Chua, S. A. Kay, A novel circadian phenotype based on firefly
luciferase expression in transgenic plants. *Plant Cell* **4**, 1075–1087 (1992).

17. T. Kondo, *et al.*, Circadian rhythms in prokaryotes: luciferase as a reporter of circadian gene
expression in cyanobacteria. *Proc Natl Acad Sci U S A* **90**, 5672–5676 (1993).

18. C. Brandes*, et al.,* Novel features of *Drosophila period* transcription revealed by real-time luciferase
reporting. *Neuron* **16,** 687-692 (1996).

19. D.K. Welsh, S.H. Yoo, A.C. Liu, J.S. Takahashi, S.A. Kay, Bioluminescence imaging of individual
fibroblasts reveals persistent, independently phased circadian rhythms of clock gene expression.
Curr Biol **14**, 2289-2295 (2004).

20. E. Nagoshi, *et al.*, Circadian gene expression in individual fibroblasts: cell-autonomous and self-
sustained oscillators pass time to daughter cells. *Cell* **119**, 693–705 (2004).

21. S. Yamazaki, *et al.*, Resetting central and peripheral circadian oscillators in transgenic rats. *Science*
288, 682–685 (2000).

22. M. Abe, *et al.*, Circadian rhythms in isolated brain regions. *J Neurosci* **22**, 350–356 (2002).

23. S. H. Yoo, *et al.*, PERIOD2::LUCIFERASE real-time reporting of circadian dynamics reveals
persistent circadian oscillations in mouse peripheral tissues. *Proc Natl Acad Sci U S A* **101**, 5339–
5346 (2004).

24. R. Stanewsky, *et al.*, The *cry^b* mutation identifies cryptochrome as a circadian photoreceptor in
Drosophila. *Cell* **95**, 681–692 (1998).

25. E. E. Zhang, *et al.,* A genome-wide RNAi screen for modulators of the circadian clock in human
cells. *Cell* **139**, 199-210 (2009).

26. T. Hirota T, *et al.,* High-throughput chemical screen identified a novel potent modulator of cellular
circadian rhythms and reveals CK1 as a clock regulatory kinase. *PLoS Biol* **8**, e1000559 (2010).
https://doi.org/10.1371/journal.pbio.1000559.

27. T. Hirota T, *et al.,* Identification of small molecule activators of cryptochrome. *Science* **337**, 1094-
1097 (2012).

28. Z. Chen, *et al.*, Identification of diverse modulators of central and peripheral circadian clocks by
high-throughput chemical screening. *Proc Natl Acad Sci U S A* **109**, 101–106 (2012).

29. W. Nakamura, S. Yamazaki , N.N. Takasu, K. Mishima, G.D. Block GD, Differential response of
period 1 expression within the suprachiasmatic nucleus. *J Neurosci* **25**, 5481-5487 (2005).

30. Y. Yamanaka, S. Honma, K-I Honma, Scheduled exposures to a novel environment with a running-
wheel differentially accelerate re-entrainment of mice peripheral clocks to new light-dark cycles.
Genes Cells **13**, 497-507(2008).

31. A. J. Davidson, S. Yamazaki, D. M. Arble, M. Menaker, G. D. Block, Resetting of central and
peripheral circadian oscillators in aged rats. *Neurobiol Aging* **29**, 471–477 (2008).

32. A. J. Davidson, O. Castanon-Cervantes, T. L. Leise, P. C. Molyneux, M. E. Harrington, Visualizing
jet lag in the mouse suprachiasmatic nucleus and peripheral circadian timing system. *Eur J*
Neurosci **29**, 171–180 (2009).

33. M. T. Sellix, *et al.*, Aging differentially affects the re-entrainment response of central and peripheral
circadian oscillators. *J Neurosci* **32**, 16193–16202 (2012).

34. P. Pezuk, J. A. Mohawk, L. A. Wang, M. Menaker, Glucocorticoids as entraining signals for
peripheral circadian oscillators. *Endocrinology* **153**, 4775–4783 (2012).

35. K. A. Stokkan, S. Yamazaki, H. Tei, Y. Sakaki, M. Menaker, Entrainment of the circadian clock in
the liver by feeding. *Science* **291**, 490–493 (2001).

36. A. Balsalobre, *et al.*, Resetting of circadian time in peripheral tissues by glucocorticoid
signaling. **289**, 2344–2347 (2000).

37. A. Balsalobre, F. Damiola, U. Schibler, A serum shock induces circadian gene expression in
mammalian tissue culture cells. *Cell* 93(6): 929-937(1998).

38. F. Damiola, *et al.*, Restricted feeding uncouples circadian oscillators in peripheral tissues from the
central pacemaker in the suprachiasmatic nucleus. *Genes Dev* **14**, 2950–2961 (2000).

39. A. B. Reddy, M. D. Field, E. S. Maywood, M. H. Hastings, Differential resynchronisation of
circadian clock gene expression within the suprachiasmatic nuclei of mice subjected to experimental
jet lag. *J Neurosci* **22**, 7326–7330 (2002).

40. M. Nagano, *et al.*, An abrupt shift in the day/night cycle causes desynchrony in the mammalian
circadian center. *J Neurosci* **23**, 6141–6151 (2003).

41. Y. Yamaguchi, *et al.*, Mice genetically deficient in vasopressin V1a and V1b receptors are resistant
to jet lag. *Science* **342**, 85-90 (2013).

42. E. Destici, E.H. Jacobs, F. Tamanini, M. Loos, G.T.J. van der Horst, M. Oklejewicz, Altered phase-
relationship between peripheral oscillators and environmental time in *Cry1* or *Cry2* deficient mouse
models for early and late chronotypes. *PLoS ONE* **8***, e83802 (2013).*

43. S. K. Nicholls, L. P. Casiraghi, W. Wang, E. T. Weber, M. E. Harrington, Evidence for internal
desynchrony caused by circadian clock resetting. *Yale J Biol Med* **92**, 259–270 (2019).

44. W. Nakamura, *et al.*, *In vivo* monitoring of circadian timing in freely moving mice. **18**, 381–385
(2008).

45. D. Ono, K. Honma, S. Honma, Circadian and ultradian rhythms of clock gene expression in the
suprachiasmatic nucleus of freely moving mice. *Sci Rep* **5**, 12310 (2015).

46. D. Ono, S. Honma, K. Honma, Circadian PER2::LUC rhythms in the olfactory bulb of freely

moving mice depend on the suprachiasmatic nucleus but not on behaviour rhythms. *Eur J*

Neurosci **42**, 3128–3137 (2015).

47. D. Ono, *et al.*, Dissociation of *Per1* and *Bmal1* circadian rhythms in the suprachiasmatic nucleus in

parallel with behavioral outputs. *Proc Natl Acad Sci U S A* **114**, E3699–E3708 (2017).

48. Y. Yamaguchi, *et al.*, Real-time recording of circadian *Per1* and *Per2* expression in the

suprachiasmatic nucleus of freely moving rats. *J Biol Rhythms* **31**,108-111 (2016).

49. L. Mei, *et al.*, Long-term in vivo recording of circadian rhythms in brains of freely moving

mice. *Proc Natl Acad Sci U S A* **115**, 4276–4281 (2018).

50. T. Hamada, *et al.*, *In vivo* imaging of clock gene expression in multiple tissues of freely moving

mice. *Nat Commun* **7**, 11705 (2016).

51. Y. Tahara, *et al.*, *In vivo* monitoring of peripheral circadian clocks in the mouse. *Curr Biol* **22**, 1029–

1034 (2012).

52. V. van der Vinne, B. Martin Burgos, M. E. Harrington, D. R. Weaver, Deconstructing circadian

disruption: Assessing the contribution of reduced peripheral oscillator amplitude on obesity and

glucose intolerance in mice. *J Pineal Res* e12654 (2020). https://doi.org/10.1111/jpi.12654

53. V. van der Vinne, S. J. Swoap, T. J. Vajtay, D. R. Weaver, Desynchrony between brain and

peripheral clocks caused by CK1δ/ε disruption in GABA neurons does not lead to adverse metabolic

outcomes. *Proc Natl Acad Sci U S A* **115**, E2437–E2446 (2018).

54. C. Saini, *et al.*, Real-time recording of circadian liver gene expression in freely moving mice reveals

the phase-setting behavior of hepatocyte clocks. *Genes Dev* **27**, 1526–1536 (2013).

55. B. Martin-Burgos, W. Wang, I. William, S. Tir, I. Mohammad, R. Javed, S. Smith, Y. Cui, C.B.

Smith, V. van der Vinne, P.C. Molyneux, S.C. Miller, D. R. Weaver, T.L. Leise, M.E. Harrington,

Methods for detecting PER2::LUCIFERASE bioluminescence rhythms in freely moving mice.

BioRxiv 2020 https://doi.org/10.1038/s41467-017-00462-2

56. J.H. Kim, *et al.*, High cleavage efficiency of a 2A peptide derived from porcine Teschovirus-1 in

human cell lines, zebrafish and mice. PLoS ONE **6**, e18556 (2011).

doi:10.1371/journal.pone.0018556

57. P. Fonjallaz, V. Ossipow, G. Wanner and U. Schibler, The two PAR leucine zipper proteins,

TEF and DBP, display similar circadian and tissue-specific expression, but have different

target promoter preferences. *EMBO J* **15**, 351-362 (1996).

58. E.D. Herzog, T. Hermanstyne, N.J. Smyllie, M.H. Hastings, Regulating the suprachiasmatic nucleus

(SCN) Clockwork: Interplay between cell-autonomous and circuit-level mechanisms. *Cold Spring*

*Harb Perspect Bio*l **9**, a027706 (2017).

59.   I. T. Lee, *et al.*, Neuromedin S-producing neurons act as essential pacemakers in the suprachiasmatic
nucleus to couple clock neurons and dictate circadian rhythms. *Neuron* **85**, 1086–1102 (2015).

60.   Y. Shan, *et al.*, Dual-color single-cell imaging of the suprachiasmatic nucleus reveals a circadian
role in network synchrony. *Neuron* **108**, 164-179 (2020).

61.   M. Mieda, *et al.*, Cellular clocks in AVP neurons of the SCN are critical for interneuronal coupling
regulating circadian behavior rhythm. *Curr Biol* **85**, 1103–1116 (2015).

62.   M. Mieda, H. Okamoto, T. Sakurai, Manipulating the cellular circadian period of arginine
vasopressin neurons alters the behavioral circadian period. *Neuron* **26**, 2535–2542 (2016).

63.   N. J. Smyllie, J. E. Chesham, R. Hamnett, E. S. Maywood, M. H. Hastings, Temporally chimeric
mice reveal flexibility of circadian period-setting in the suprachiasmatic nucleus. *Proc Natl Acad Sci*
U S A **113**, 3657–3662 (2016).

64.   M. Brancaccio, *et al.*, Cell-autonomous clock of astrocytes drives circadian behavior in
mammals. *Science* **363**, 187–192 (2019).

65.   R.C. Poulsen, *et al.,* How does general anaesthesia affect the circadian clock? *Sleep Med Rev.* **37**,
35-44 (2018).

66.   Y. Sawai, *et al.*, *In vivo* evaluation of the effect of lithium on peripheral circadian clocks by real-
time monitoring of clock gene expression in near-freely moving mice. *Sci Comm* **9**, 10909 (2019).

67.   K. Hamada K, *et al.*, Double recording system of *Period1* gene expression rhythm in the olfactory
bulb and liver of freely moving mouse. *Biochem Biophys Res Comm* **529**, 898-903 (2020).

68.   S. Yamaguchi, *et al.*, View of a mouse clock gene ticking. *Nature* **409**, 684 (2001).

69.   Acosta-Rodriguez, M.H.M. de Groot, F. Rijo-Ferreira, C.B. Green, J. S. Takahashi, Mice under
caloric restriction self-impose a temporal restriction of food intake as revealed by an automated
feeder system. *Cell Metab* **26**, 267-277 (2017).

70.   T. Noguchi, *et al.*, Circadian rhythm bifurcation induces flexible phase resetting by reducing
circadian amplitude. *Eur J Neurosci* **51**, 2329–2342 (2020).

71.   T. L. Leise, *et al.*, Recurring circadian disruption alters circadian clock sensitivity to resetting. *Eur J*
Neurosci **51**, 2343–2354 (2020).

72.   L. Lopez-Molina *et al.*, The DBP gene is expressed according to a circadian rhythm in the
suprachiasmatic nucleus and influences circadian behavior. *EMBO J* **16**, 6762-6771 (1997).

73.   M.S. Evans, *et al.*, A synthetic luciferin improves bioluminescence imaging in live mice. *Nat*
Methods **11**, 393-395 (2014).

74.   S. Iwano, *et al.*, Single-cell bioluminescence imaging of deep tissue in freely moving animals.
Science **359**, 935-939 (2018).

75. J.E. Miller*, et al*., Vasoactive intestinal polypeptide mediates circadian rhythms in mammalian
olfactory bulb and olfaction*. J Neurosci* **34**, 6040-6046 (2014).

76. X. Xie, et al., Natural food intake patterns have little synchronizing effect on peripheral circadian
clocks. *BMC Biology* **18**, 160 (2020). https://doi.org/10.1186/s12915-020-00872-7

77. F. Sinturel, *et al*., Circadian hepatocyte clocks keep synchrony in the absence of a master pacemaker
in the suprachiasmatic nucleus or other extrahepatic clocks. *Genes Dev* **35,** (e-pub ahead of print)
(2021). http://www.genesdev.org/cgi/doi/10.1101/gad.346460.120

78. S. H. Yoo, *et al.*, *Period2* 3′-UTR and microRNA-24 regulate circadian rhythms by repressing
PERIOD2 protein accumulation. *Proc Natl Acad Sci U S A* **114**, E8855–E8864 (2017).

79. D. R. Weaver, *et al.*, Functionally complete excision of conditional alleles in the mouse
suprachiasmatic nucleus by *Vgat-ires-Cre*. *J Biol Rhythms* **33**, 179–191 (2018).

80. S. Yamazaki, J. S. Takahashi, Real-time luminescence reporting of circadian gene expression in
mammals. *Methods* **393**, 288–301 (2005).

81. J. A. Evans, T. L. Leise, O. Castanon-Cervantes, A. J. Davidson, Intrinsic regulation of
spatiotemporal organization within the suprachiasmatic nucleus. *PLoS ONE* **6**, e15869 (2011).

82. J. A. Evans, T. L. Leise, O. Castanon-Cervantes, A. J. Davidson, Dynamic interactions mediated by
nonredundant signaling mechanisms couple circadian clock neurons. *Neuron* **80**, 973–983 (2013).

83. T. L. Leise, M. E. Harrington, Wavelet-based time series analysis of circadian rhythms. *J Biol
Rhythms* **26**, 454–463 (2011).

84. T. L. Leise, Analysis of nonstationary time series for biological rhythms research. *J Biol Rhythms*
32, 187–194 (2017).

85. T. L. Leise, *et al.*, Voluntary exercise can strengthen the circadian system in aged mice. *Age (Dordr)*
35, 2137–2152 (2013).

Figure 1. Generation of a bifunctional reporter from the mouse *Dbp* locus.

A. The mouse *Dbp* locus was modified by CRISPR-mediated insertion of the donor construct shown. The

construct contained homology arms from the *Dbp* locus (gray and black) and inserted the reporter sequences

with a T2A-encoding sequence (orange) between DBP and the reporter. Destabilized EGFP (d2EGFP) with

a bovine growth hormone polyadenylation site (PA) was flanked by *loxP* sites (red). Downstream of *GFP*

is a luciferase (*Luc2*) reporter gene. Without recombination *Dbp* and *GFP* are expressed as a single

transcript from the conditional (*Dbp^KI* allele).

B. With *Cre*-mediated recombination, GFP-encoding sequences are excised and *Dbp* and *luciferase* are

expressed as a single transcript. The T2A sequence generates separate proteins from these bifunctional

transcripts. *Cre*-mediated germline recombination led to mice expressing luciferase non-conditionally from

the *Dbp^Luc* allele.

File: Fig 1 Construct figure copy_drw121120.jpg

Figure 2. Tissue explants from *Dbp^{Luc/+}* mice have an earlier time of peak bioluminescence than explants from *Per2^{Luc/+}* mice *in vitro*.

A-C, Anterior Pituitary gland explants. **D-F**, Lung explants.

A., B., D., and E. are representative bioluminescence rhythms from triplicate tissue explants from *Per2^{Luc/+}* (**A, D**) and *Dbp^{Luc/+}* mice (**B, E**). 'Days' refers to time in culture, not projected ZT. Values are 24-h background-subtracted, 3-h smoothed values.

C,F. Time of peak bioluminescence *in vitro*. The large circles represent a 24-h day for each organ. ZT's refer to the lighting cycle to which the mice were exposed prior to sample collection, with ZT0-12 being the light phase. Colored points at the perimeter of the large circle indicate the timing of peak bioluminescence of individual *Per2^{Luc/+}* (dark blue) or *Dbp^{Luc/+}* (teal) tissue explants (n=12-14 mice). Within each tissue/genotype combination, there was significant clustering of times of peak bioluminescence. Radial lines represent the mean peak time, which differed significantly between genotypes for each tissue (Watson-Williams test, p<0.001).

File: Fig 2 in vitro-032821.jpg

Figure 3. Bioluminescence rhythms measured *in vivo*

A-C. Bioluminescence images captured at 4-6 hr intervals from a representative mouse of each genotype.
A. *Per2^{Luc/+}*, **B**. *Dbp^{Luc/+}* **C**. *Alb-Cre+ ; Dbp^{KI/+}*. Ventral (V) and dorsal (D) views are shown for each
mouse. All images for each mouse are set to the same luminescence scale.

D-F. Cosinor-fitting of bioluminescence signal over time for the animals shown in Panels A-C to determine
peak time. Bioluminescence rhythms were assessed in submandibular gland, liver, and kidneys of (**D.**)
Per2^{Luc/+} and (**E.**) *Dbp^{Luc/+}* reporter mice, and from liver of *Alb-Cre+ ; Dbp^{KI/+}* mice (**F.**).

G-I. Time of peak bioluminescence *in vivo* as assessed by IVIS imaging. **G.** Submandibular gland, **H.**
Kidneys, and **I.** Liver. Data plotted as in Fig. 2C and 2F. Dark blue symbols are *Per2^{Luc/+}* tissues (n=10),
teal symbols are *Dbp^{Luc/+}* tissues (n=7). In Panel H, open symbols represent the right kidney and filled
symbols represent the left kidney. In Panel I, purple circles represent livers from *Alb-Cre+ ; Dbp^{KI/+}* mice

(n=8). Radial lines represent the mean peak time for each genotype and tissue. Radial lines from the two

kidneys of a genotype are nearly overlapping. For liver, radial lines for the two *Dbp* reporter lines are

overlapping and appear as a single line. Time of peak for each *Per2$^{Luc/+}$* organ examined differed

significantly from time of peak of the corresponding organs from *Dbp$^{Luc/+}$* and *Alb-Cre+ ; Dbp$^{KI/+}$* mice

(p=0.002, Watson-Williams test). There was no significant difference in peak time between *DbpLuc/+* and

Alb-Cre+ ; Dbp$^{KI/+}$ liver tissues (P >0.05).

File: Fig 3 in vivo IVIS_032821scaled.jpg

Figure 4. *Dbp* mRNA rhythms are not altered in reporter mice

A-C. Representative Northern Blots probed to detect *Dbp* and *Actin* mRNA. **A.** From each of five genotypes, RNA samples were extracted from livers collected at ZT 2 and 10. For each genotype, there are two samples at ZT10 and one sample at ZT2 on this blot. **B**. and **C**. Representative Northern Blots of RNA samples collected from WT and reporter mouse livers at each of six Zeitgeber times (ZT).

D-F. Quantification of *Dbp* mRNA rhythms for each allele in time-series experiments (6 time-points each). Results are expressed as mean (± SEM) percent of the peak *Dbp*/Actin ratio, which occurred at ZT 10 on every blot. **D.** Wild-type *Dbp* transcript (n=12 sample sets). **E.** *Dbp^{KI}* transcript (n= 6). **F.** *Dbp^{Luc}* transcript (n= 6). For each transcript, there was a significant rhythm (Friedman's One-way ANOVA, Q > 19, *p* < 0.002). Asterisks indicate time-points that differed significantly from ZT10 (Dunn's test, * *p* < 0.05, ** *p*

980   < 0.01, *** $p < 0.001$, **** $p \leq 0.0001$). Significant differences among some other time-points are not

981   shown for clarity.

983   *File: Fig 4_Northern figure with bar graphs_010321.jpg*

Figure 5. Cell-type-specific imaging of LUCIFERASE expression in SCN slices

A) 24h summed bioluminescence overlaid onto bright field images of a section through the SCN from

DBP^{Luc/+} (global reporter expression, left), and in mice expressing luciferase from specific subsets of SCN

neurons (NMS$^+$ cells, center; AVP$^+$ cells, right).

B. Representative bioluminescence traces from single neuron-like ROIs in slices from each genotype.

C. Circular plots indicate the peak time of bioluminescence rhythms from each genotype. Time is expressed

relative to the light-dark cycle the mice were housed in prior to sacrifice; numbers >24 are used to indicate

that these measures are recorded on the first day *in vitro* and are plotted relative to previous *in vivo* lighting

conditions. Each slice is represented by a small dot. Placement of the dot relative to outer circle indicates

average peak time (±SD), while the distance from the center corresponds to the number of cells incorporated

in the average ($\sqrt{\text{cell\#}}$).

D. Mean period (± SEM) by genotype. The number of slices per genotype is indicated at the base of each

bar (D-G).

E. Circular mean peak time (± SEM) by genotype.

F. Mean rhythmicity index score (± SEM) by genotype.

G. Mean peak time dispersal (quantified by circular SD of peak times within each slice) by genotype.

1002

1003 *File: Fig 5_Smith et al_032821.jpg*

Figure 6. Light-induced resetting produces misalignment between liver bioluminescence rhythms and locomotor activity rhythms.

A. Representative double-plotted actogram showing locomotor activity (dark gray) and bioluminescence (dark red) of an *Alb-Cre; Dbp*$^{KI/+}$ liver reporter mouse before and after a 6-h advance of the skeleton photoperiod consisting of four 1-h periods of light per 24-h day, as indicated by white. The skeleton photoperiod was advanced by 6 h by shortening the dark phase after the last light pulse on Day 2. Red squares represent the peak of the bioluminescence rhythm, while black circles represent the midpoint of locomotor activity each day. These values were determined by discrete wavelet transform analysis. Six hours of each cycle are double-plotted to aid visualization of the data. Light and dark are indicated by white and light-gray backgrounds, respectively.

B. Mean (± SEM) midpoint of locomotor activity (black) and peak of liver bioluminescence (red) rhythms are shown, relative to their initial value, in a group of 4 mice exposed to a shifted skeleton photoperiod. Both rhythms reset gradually after a 6-h phase advance of the skeleton photoperiod, but the locomotor activity rhythm re-sets more rapidly than the bioluminescence rhythm within animal (Significant Measure * Day interaction, and significant phase difference between the rhythms on Day 9 (Tukey HSD, $p < 0.05$).

C. Mean (± SEM) time of midpoint of locomotor activity (black) and peak liver bioluminescence (red) rhythms are shown, relative to their initial phase, in a group of 4 mice not subjected to a phase shift of the skeleton photoperiod.

1023
1024   *File: Fig 6_Liver Reporter SPP phase shift.JPG*
1025

1026

Figure 7. Time-restricted feeding alters the timing of liver bioluminescence rhythms.

A. Representative actograms of three *Alb-Cre; Dbp^{KI/+}* liver reporter mice exposed to the different feeding regimes as indicated above each panel. Mice were housed in a 12h:12h light-dark cycle and exposed to the specified feeding regime for ten days (-10 to 0) preceding bioluminescence recording. The timing of food intake (blue triangles) and general locomotor activity (dark gray) was recorded continuously. The midpoint of food intake over the last five days preceding bioluminescence recording is indicated by a cyan diamond on day 0. Mice were transferred to the bioluminescence recording setup at the start of the dark phase and housed in constant darkness with *ad libitum* food access starting 12 hr later. Liver bioluminescence levels are depicted in dark red. Red squares represent the time of peak of the bioluminescence rhythm, determined by discrete wavelet transform. Six hours of each cycle are double-plotted and the y-axis has been stretched during the last 6 days to aid visualization of the data. Light and dark are indicated by white and light gray backgrounds, respectively.

B. Individual and mean (± SEM) phase of liver bioluminescence rhythms relative to clock time for three feeding groups. Mice previously exposed to *ad libitum*, nighttime and daytime feeding are plotted in grey/black, blue/cyan and magenta, respectively (key in Panel C). Prior to recording

bioluminescence, mice were entrained to a 12L:12D lighting cycle with lights on at 0600. Mice

previously exposed to daytime feeding show an advanced peak phase of liver bioluminescence that

reverts over time in constant darkness with *ad libitum* food.

C. Relationship between preceding feeding phase and peak liver bioluminescence phase for individual

animals on the first day under constant conditions. *Ad libitum* and night-fed groups had similar

midpoint of food intake; three "outliers" with respect to midpoint of food intake (shown by open

symbols) were not included in further analyses (Panels B, D and E).

D. Mean (± SEM) peak liver bioluminescence phase on the first day under constant conditions, relative

to clock time for the three feeding regimens. The low variability within groups resulted in error bars

that were nearly or completely contained within the symbols.

E. Mean (±SEM) peak liver bioluminescence phase on the first day under constant conditions, relative

to the midpoint of preceding food intake for the three feeding regimens. The low variability within

groups resulted in error bars that were nearly or completely contained within the symbols.

1056

1057   *File: Fig 7_Smith et al_032921.JPG*

1058

1059   **Table 1:** Period length of locomotor activity rhythms in constant darkness, by sex and genotype

1060

1061	Genotype	Sex	N	tau_{DD} (Mean +/- SEM), h
1062	$Dbp^{+/+}$	Male	15	23.88 ± 0.027
1063	$Dbp^{KI/+}$	Male	10	23.91 ± 0.057
1064	$Dbp^{KI/KI}$	Male	11	23.92 ± 0.036
1065	$Dbp^{Luc/+}$	Male	11	23.86 ± 0.025
1066	$Dbp^{Luc/Luc}$	Male	8	23.97 ± 0.029
1067				
1068	$Dbp^{+/+}$	Female	21	23.87 ± 0.021
1069	$Dbp^{KI/+}$	Female	9	23.89 ± 0.036
1070	$Dbp^{KI/KI}$	Female	11	23.79 ± 0.030
1071	$Dbp^{Luc/+}$	Female	8	23.82 ± 0.053
1072	$Dbp^{Luc/Luc}$	Female	8	23.75 ± 0.042