Intragenomic sequence variations in the second internal transcribed spacer (ITS2) ribosomal DNA of the malaria vector *Anopheles stephensi*

Shobhna Mishra[1], Gunjan Sharma[1], Manoj K. Das[2], Veena Pande[3], Om P. Singh[1]*,

[1]National Institute of Malaria Research, Sector 8, Dwarka, New Delhi-110077, India

[2]National Institute of Malaria Research, Field Unit, Itki, Ranchi-835301, India

[3]Department of Biotechnology, Kumaun University, Nainital, Uttarakhand-263001, India

*Corresponding author
Email: dr.opsingh@gmail.com
ORCID ID: 0000-0001-5135-6654

Abstract

Second Internal Transcribed Spacer (ITS2) ribosomal DNA (rDNA) sequence is a widely used molecular marker for the species-identification or -delimitation due to observed concerted evolution which is believed to homogenize rDNA copies in an interbreeding population. However, intra-specific differences in ITS2 of *Anopheles stephensi* have been reported. This study reports the presence of intragenomic variation in the ITS2 region of *An. stephensi* and hypothesized that observed intra-specific differences in this species may have been resulted due to ambiguous DNA-chromatogram resulting from intragenomic heterogeneity. *Anopheles stephensi* collected from different parts of India were sequenced for complete ITS2 and the variable region of 28S rDNA (d1-d3 domains). Intragenomic variations at fixed loci were found in ITS2 of all *An. stephensi* sequenced, but no such variation was observed in d1 to d3 domains of 28S rDNA. Cloning and sequencing of ITS2 through the d3 domain of the 28S region of rDNA from representative samples from northern, central and southern India confirmed the presence of intragenomic variation in ITS2. The variations are due to three transitions and two bp indel in a di-nucleotide microsatellite locus at fixed loci. Multiple haplotypes were observed in ITS2 raised from such variations. Due to the absence of detectable intragenomic sequence variation in the d1 to d3 domain of 28S rDNA of *An. stephensi*, this region can serve as an ideal reference sequence for taxonomic and phylogenetic studies. The presence of intragenomic variation in rDNA should be carefully examined before using this as a molecular marker for species delimitation or phylogenetic analyses.

Key words: *Anopheles stephensi*, intragenomic variation, ITS2, ribosomal DNA, malaria

Introduction

Anopheles stephensi is a major malaria vector in India, especially in an urban setting. This vector is predominantly found in urban areas due to their preference to breed in clean water especially in cement tanks, containers, overhead tanks, cisterns, fountain tanks, and building construction sites [1]—most abundantly found in urban areas. Perhaps due to the adaptation to such breeding habitats, this species is invading several neighbouring territories to their native. Because of the high infectivity rate of *An. stephensi* to human-*Plasmodium* species [2-4] and its invasive nature, this malaria vector has now become a global health concern.

Historically, *An. stephensi* was initially reported from the Middle-East and South Asia region (Afghanistan, Bahrain, Bangladesh, China, Egypt, India, Iran, Iraq, Oman, Pakistan, Saudi Arabia, Thailand) [1]. According to Sharma (2012) [5] *An. stephensi* is an invasive species that appeared first in port cities in India and invaded the riparian town and then in towns with excessive wells. During the last few decades, this species had invaded areas where this species was never reported before. Invasions have been reported in two islands near India, i.e., Lakshadweep (India) [6] and Sri Lanka [7-9], Republic of Djibouti [10], Ethiopia [11-12] and Sudan [13]. *Anopheles stephensi* is now established in the Horns of Africa and contributing to local malaria transmission [14]. As a consequence of invasion, Word Health Organization [13] alerted WHO-member-states and their implementing partners of affected countries to take immediate action.

For the identification of *An. stephensi,* the second internal transcribed spacer (ITS2) ribosomal DNA (rDNA) has been frequently used as a molecular marker. Ribosomal DNA is comprised of tandemly arranged units of ETS, 18S, ITS1, 5.8S, ITS2, and 28S, in several hundreds of copies in insects. DNA sequences of variable regions of 28S-rDNA, IGS, and ITS2 in particular, have been frequently used for species identification, species delimitation, and phylogenetic analyses. 28S rDNA is a relatively conserved region, however, some domains (d2 and d3) are variable and have been used as markers for species differentiation [15-20]. The second internal transcribed spacer region, on the other hand, which is relatively highly variable among different species because of its high evolution rate, is the most extensively used taxonomic marker for species identification and phylogenetic studies. The rDNA sequences are reported to be highly conserved in the interbreeding population due to the homogenization of copies which is thought to be achieved due to concerted evolution [21]. However, in some species, intragenomic variations have been reported with the presence of multiple haplotypes. Such intragenomic variation leads to challenges in DNA sequencing especially when indel is present in some copies. The presence of indel leads to ambiguous sequencing-result starting from the point of indel onwards which can be resolved through cloning [21]. The presence of such indels has resulted in the submission of incorrect sequences in GenBank or published reports. Such a phenomenon has recently been observed in the case of *An. subpictus* [22] and it was noted that all the reported intra-specific differences in a molecular form (Form A) of *An. subpictus* was in fact due to ambiguous DNA sequence chromatogram resulting from the

presence of indel in one of the two haplotypes of ITS2 present in each individual. The selection of the molecular region of rDNA, which can serve as reference sequences for the species identification, is therefore important and a region exhibiting intragenomic heterogeneity should be avoided. Moreover, recording of intragenomic sequence variation in rDNA becomes important while reporting sequence data to avoid ambiguities in taxonomic and phylogenetic studies.

In the case of *An. stephensi,* numerous ITS2 sequences are available in the GenBank and research publications [2, 8, 10, 23-25]. However, polymorphism in length and nucleotide sequence has been observed in sequences reported from Iran [23], India [25], and in several other GenBank sequences. Such intraspecific polymorphism is intriguing. In some other reports from India [24], Saudi Arabia [26], Ethiopia [11] and Sri Lanka [8], however, no intraspecific differences were noted. One plausible reason which explains reports of polymorphism in some reports is the presence of intragenomic sequence variation which has not been taken care of during DNA sequence analysis.

In this study, we characterized partial 5.8S, complete ITS2, and partial 28S (d1 to d3 domains) of *An. stephensi* to investigate intragenomic variations and delineate the region of rDNA suitable for molecular taxonomy that can be correctly sequenced without cloning.

Material and methods

Mosquito samples

Anopheles stephensi mosquitoes were collected from different parts of India, i.e., Gurugram (Haryana), Nuh (Haryana), Alwar (Rajasthan), and New Delhi from northern India, Ranchi (Jharkhand), Raipur (Chattishgarh), and Gadhchiroli (Maharashtra) from central India, Goa, Bangalore (Karnataka) Mangalore (Karnataka), Chennai (Tamil Nadu) and Mysuru (Karnataka) from southern India (**Figure 1**). Adult mosquitoes were identified using keys by Christophers [28].

DNA isolation and direct sequencing

DNA was isolated from individual female mosquitoes using the method by Livak [27]. A part of rDNA spanning part of 5.8S rDNA, complete ITS2, and part of 28S rDNA (d1 to d3 domains) was amplified by using primers ITS2A (5'-TGT GAA CTG CAG GAC ACA T-3') [29] and D3B

(5'-TCG GAA GGA ACC AGT TAC TA-3') [30]. The PCR reaction mixture (20 µL) contained 1X buffer, 1.5 mM MgCl$_2$, 200 µM of each dNTP, 0.25 µM of each primer, 0.50 unit of Taq polymerase (GoTaq®, Promega Corporation Inc), and 0.5 µL of DNA template. The PCR conditions were: an initial denaturation step at 95 °C for 3 min, followed by 35 cycles each with a denaturation step at 95 °C for 30 sec, annealing step at 55 °C for 30 sec and extension step at 72 °C for 1.5 min, followed by a final extension at 72 °C for 7 min. The PCR product quality was checked on 1.5% agarose gel. The PCR products were subjected to DNA sequencing using Sanger's method. The PCR products were treated with Exo-Sap (ExoSAP-IT™, Thermo Fisher, USA) to remove unutilized primers and dNTPs, and subjected to cycle sequencing reaction using BigDye Terminator v3.2 (Invitrogen Inc., USA) following vendor's protocol. The primers used for sequencing termination reaction were ITS2A, ITS2D (5'-TAT GCT TAA ATT CTG AGG GT-3'), D2A (5'-AGT CGT GTT GCT TGA TAG TGC AG-3') [31], D2B (5'-TTG GTC CGT GTT TCA AGA CGG G-3') [31], D3A (5'-GAC CCG TCT TGA AAC ACG GA-3') [30] and D3B. The sequence termination reaction products were cleaned up using ethanol precipitation and were electrophoresed in ABI Prism 3730xl. The numbers of samples sequenced from different parts of India has been shown in **Table 1**.

Cloning and sequencing

For cloning, the rDNA was first amplified from individual female mosquitoes from Nuh (representing northern India), Gadchiroli (central India) and Chennai (representing southern India) using a high-fidelity DNA Taq polymerase to minimize PCR-errors. The PCR reaction mixture (25 µL) contained 0.5 µM of each forward and reverse primer (ITS2A and D3B), 0.50 unit of Phusion® High-Fidelity DNA Polymerase, 5 µL of 5X Phusion HF reaction buffer (all from New England Biolabs, USA), 0.5 µL of 10 mM premixed dNTPs and 0.5 µL of gDNA. PCR conditions were: initial denaturation at 98˚C for 30 sec, 35 cycles of each of denaturation at 98˚C for 10 sec, annealing at 65 °C for 30 sec and extension at 72 °C for 45 sec, followed by a cycle of final extension at 72 °C for 2 min. Five µL of PCR products were visualized on 2% agarose gel and the remaining were purified using QIAquick PCR purification kit (Qiagen Inc, USA) following the manufacturer's protocol. A-tail was incorporated at 3' end of the purified PCR product by incubating 50 ng of PCR product at 70 °C for 20 min in a reaction mixture (20 µL) containing 1X PCR buffer, 1.5 mM MgCl$_2$, 0.5 unit of Taq DNA polymerase, and 200 µM

of dATP. Four µL of purified A-tailed PCR products were ligated to pGEM-T easy vector (Promega Corporation) and 5 µL of this ligated mixture was then transformed into DH5α competent cells (New England Biolab, USA). Transformed cultures were plated on Luria-Bertani (LB) agar plates containing 5-bromo-4-chloro-3-indolyl-beta-D-galactopyranoside (X-gal), isopropyl-beta-D-thiogalactopyranoside (IPTG), and 50 µg/mL ampicillin. White transformant colonies originating from each mosquito were picked up and DNA was isolated by boiling them in TE buffer at 95 °C. The plasmid DNA was subjected to PCR using primers ITS2 and D3B and the amplified products with successful insert, as evident from the size of PCR product on agarose gel, were treated with Exo-sap and sequenced as described above.

Results

Sequencing of cloned PCR products revealed the presence of a total of ten haplotypes (**Figure 2**) resulting from SNPs at three fixed loci in the ITS2 region, i.e., C>T transition at nucleotide base positions 172 and 383, G>A transition at 401, and two bp indel (CA) at base positions 182-183 (base position numbering as per **Figure 2**). No intragenomic variation was found in the sequenced region of 5.8S and d1 to d3 domains of 28S rDNA. To avoid inclusion of an SNP in cloned samples resulting from possible PCR error, any SNP which was present only in one clone, were suspected to be raised from PCR error and were excluded from the analysis. The number of clones sequenced and distribution of haplotypes in representative samples from three populations have been shown in **Table 2**. The Hap-1 was the dominant haplotype. The sequences of all haplotypes have been submitted to GenBank (accession numbers: MW676288—MW676292; MW732930—MW732931). Since NCBI no more accept annotation of rDNA sequences submitted, due to recent policy change, annotation details have been provided in supplementary file S1.

Direct sequencing of *An. stephensi* from different localities, representing northern, central, and southern India, was performed for ITS2 and d1-d3 domains of 28S-rDNA (**Table 1**). Due to the presence of indel at base positions 182-183 of ITS2, the sequences were ambiguous in sequence reads beyond this point (**Figure 3**). Therefore, forward sequences were read before the indel point and reverse sequences were read after the indel. Mixed bases were seen at all polymorphic loci, as seen in clonal sequencing, in all the samples. However, it was difficult to recognize transition 383C>T in some samples and differentiate from noise due to the low peak

area of T-allele. The result shows that intragenomic variation is universal, at least in the Indian population. The direct sequence chromatogram also showed that CA-microsatellite present in ITS2 (base positions 172-182) was dimorphic (with six or seven repeats) and not highly polymorphic like other microsatellites. No polymorphism was seen in the 5.8S and 28S rDNA regions sequenced.

To investigate if mixed haplotypes in *An. stephensi* is due to the hybridization of type form and *var. mysorensis*, two individuals of each form were also sequenced for ITS2 and 28S rDNA. No difference was found in the sequence and pattern of intragenomic variations.

Discussion

Ribosomal DNA sequence is a widely used molecular taxonomic marker for the species-identification/delimitation and phylogenetic studies due to homogenization of copies of rDNA in an interbreeding population through observed concerted evolution, which leads to a remarkably low or complete absence of intra-specific variation. Among rDNA regions, the ITS2-rDNA is the most frequently used molecular marker that exhibits remarkably high inter-specific differentiation due to high evolution rate, although it remains under evolutionary constraints to maintain the specific secondary structures that provide functionality [32] of rDNA. However, the use of rDNA sometimes becomes problematic due to intragenomic variation, particularly indel. Intragenomic variations in rDNA, particularly in ITS2, are not uncommon and have been observed in several anophelines [33-36]. The indel in some copies of rDNA sequence significantly affects the DNA sequence quality which leads to the collapse of the DNA sequence chromatogram [21] downstream to the point of indel. Instances exist where such intragenomic variations are overlooked and potentially inaccurate sequences are submitted to GenBank or documented in the research articles. A recent report by Sindhania et al. [22] revealed that most of the ITS2-sequences of a molecular form of *An. subpictus* (prevalent mainly in Indian mainland, Form A) available in the public domain are incorrect due to the presence of indel in one haplotype. Therefore, the generation of quality sequence and documentation of intragenomic variation in rDNA is essential for the correct molecular characterization of a species.

The ITS2 sequences of *An. stephensi* have been reported by several authors but intragenomic variations have not been reported. However, significant intra-specific variations in ITS2 have been reported in some reports. It is suspected that the observed intraspecific

polymorphism in ITS2 is raised due to compromised sequence quality. Such intraspecific variations have been exploited for population genetic analysis [23] which may be misleading. In a study carried out in Sri Lanka [8], only a part of sequences, representing approximately half of the sequence generated (using primers from 5.8S and 28S rDNA), were submitted. Similarly, partial sequences were submitted from the Horn of Africa [2]. Trimming of significant portions of DNA sequence in these studies is probably due to ambiguities encountered in sequence read due to intragenomic variation particularly indel at the microsatellite locus. However, none of the reports documented intragenomic variation except in one GenBank entry from the Horn of Africa [2] (accession number: MN826065) where mixed bases (G/A) have been shown at base position 401. It is most likely that intragenomic variation is present beyond India which is evident from the fact that the sequences available from the public domain from other countries are either divergent or only fragments of the sequences have been reported from. However, the presence of intragenomic variation in other countries needs confirmation. This study also revealed that polymorphic sites are fixed in all Indian populations studied. A similar observation has been documented in the case of *An. subpictus* (molecular Form A) where two ITS2 haplotypes were found present in all populations from northern India to Sri Lanka [22].

Due to limited numbers of clones sequenced, the possibility of missing some rare variants, if any, may exist; although, such variations, present in very low frequency, will not be affecting the quality of direct sequencing chromatogram. Extremely rare variants will not affect the quality of DNA sequences and utility of sequence in species delimitation or phylogenetic studies. It has been shown [37] that DNA copies that are present in less than one-tenth proportion do not have obvious signals in sequence chromatogram. All the variations we observed through clonal sequencing in *An. stephensi* can also be detected to a greater extent through a direct-sequencing chromatogram. However, the quality of DNA sequence with minimal noise at base of chromatogram is crucial. The noise in the DNA sequence can be significantly reduced by choosing a sequencing primer different than the primers that were used for PCR amplification. This eliminates parallel sequencing of PCR artifacts such as non-specific PCR products present in very low quantity and primer-dimers.

The second internal transcribed spacer region has been an extensively used molecular marker for species characterization, however, coding regions of rDNA have also been used for this purpose. The 28S-rDNA, which is believed to be highly conserved, has less frequently been

used in molecular taxonomy. However, the variable regions of 28S-rDNA, i.e., D2 and D3 domain [15-20] have exhibited the taxonomic significance and have been used for developing molecular differentiation of closely related sibling species/groups. This study provides the first-ever reference sequence of *An. stephensi* for d1 to d3 domains of 28S rDNA. This region didn't exhibit obvious intragenomic sequence variation. Therefore, this sequence can be used as a reference sequence for *An. stephensi* for molecular taxonomy.

Conclusions

Due to the presence of intragenomic sequence variations in the ITS2-rDNA of *An. stephensi,* this region is not a suitable molecular marker for the identification of *An. stephensi* and molecular phylogenetic analysis. The 28S-rDNA (d1 through d3 domains) sequence can serve as a superior molecular marker for this purpose due to the absence of intragenomic variation.

Acknowledgements

The authors acknowledge Dr. Robin Marwal for providing specimens of type form and *var mysorensis*, Dr. V.P. Ojha for providing mosquitoes from Gadchiroli and Mr. Shri Bhagwan for collecting mosquitoes from Chennai. SM and GS were supported by ICMR-Senior Research Fellowship.

References

1. Dash AP, Adak T, Raghavendra K, Singh OP. The biology and control of malaria vectors in India. 2007; Curr Sci: 92, 1571-1578.
2. Tadesse FG, Ashine T, Teka H, Esayas E, Messenger LA, Chali W, et al. *Anopheles stephensi* mosquitoes as vectors of *Plasmodium vivax* and *falciparum*, Horn of Africa, 2019. Emerg Infect Dis. 2021; 27(2):603-607. doi: 10.3201/eid2702.200019.
3. Adak T, Kaur S, Singh OP. Comparative susceptibility of different members of the *Anopheles culicifacies* complex to *Plasmodium vivax*. Trans R Soc Trop Med Hyg. 1999; 93(6):573-7. doi: 10.1016/s0035-9203(99)90052-4.
4. Kaur S, Singh OP, Adak T. Susceptibility of species A, B, and C of *Anopheles culicifacies* complex to *Plasmodium yoelii yoelii* and *Plasmodium vinckei petteri*

infections. J Parasitol. 2000; 86(6):1345-8. doi: 10.1645/0022-
3395(2000)086[1345:SOSABA]2.0.CO;2.

5. Sharma VP. Continuing challenge of malaria in India. 2012; Curr Sci, 102: 678–682

6. Sharma SK, Hamzakoya KK. Geographical spread of *Anopheles stephensi* vector of
urban Malaria, and Aedes aegypti, vector of dengue/ DHF, in the Arabian Sea Islands of
Lakshadweep, India. Dengue Bulletin. 2001; 25: 88–91.
https://apps.who.int/iris/handle/10665/148798

7. Gayan Dharmasiri AG, Perera AY, Harishchandra J, Herath H, Aravindan K, Jayasooriya
HTR, et al. First record of *Anopheles stephensi* in Sri Lanka: a potential challenge for
prevention of malaria reintroduction. Malar J. 2017;16(1):326. doi: 10.1186/s12936-017-
1977-7.

8. Surendran SN, Sivabalakrishnan K, Gajapathy K, Arthiyan S, Jayadas TTP, Karvannan
K, et al. Genotype and biotype of invasive *Anopheles stephensi* in Mannar Island of Sri
Lanka. Parasit Vectors. 2018; 11(1):3. doi: 10.1186/s13071-017-2601-y.

9. Surendran SN, Sivabalakrishnan K, Sivasingham A, Jayadas TTP, Karvannan K,
Santhirasegaram S, et al. Anthropogenic factors driving recent range expansion of the
malaria vector *Anopheles stephensi*. Front Public Health. 2019; 7:53. doi:
10.3389/fpubh.2019.00053.

10. Faulde MK, Rueda LM, Khaireh BA. First record of the Asian malaria vector *Anopheles
stephensi* and its possible role in the resurgence of malaria in Djibouti, Horn of Africa.
Acta Trop. 2014;139:39-43. doi: 10.1016/j.actatropica.2014.06.016.

11. Carter TE, Yared S, Gebresilassie A, Bonnell V, Damodaran L, Lopez K, et al. First
detection of *Anopheles stephensi* Liston, 1901 (Diptera: culicidae) in Ethiopia using
molecular and morphological approaches. Acta Trop. 2018;188:180-186. doi:
10.1016/j.actatropica.2018.09.001.

12. Balkew M, Mumba P, Dengela D, Yohannes G, Getachew D, Yared S, et al.
Geographical distribution of *Anopheles stephensi* in eastern Ethiopia. Parasit Vectors.
2020;13(1):35. doi: 10.1186/s13071-020-3904-y..

13. WHO. Vector alert: *Anopheles stephensi* invasion and spread. 26 August 2019.
https://www.who.int/news/item/26-08-2019-vector-alert-anopheles-stephensi-invasion-
and-spread (retrieved: 1 March, 2021)

14. Seyfarth M, Khaireh BA, Abdi AA, Bouh SM, Faulde MK. Five years following first detection of *Anopheles stephensi* (Diptera: Culicidae) in Djibouti, Horn of Africa: populations established-malaria emerging. Parasitol Res. 2019;118(3):725-732. doi: 10.1007/s00436-019-06213-0.

15. Sharpe RG, Hims MM, Harbach RE, Butlin RK. PCR-based methods for identification of species of the *Anopheles minimus* group: allele-specific amplification and single-strand conformation polymorphism. Med Vet Entomol. 1999; 13(3):265-73. doi: 10.1046/j.1365-2915.1999.00178.x.

16. Singh OP, Chandra D, Nanda N, Raghavendra K, Sunil S, Sharma SK, et al. Differentiation of members of the *Anopheles fluviatilis* species complex by an allele-specific polymerase chain reaction based on 28S ribosomal DNA sequences. Am J Trop Med Hyg. 2004;70(1):27-32.

17. Singh OP, Goswami G, Nanda N, Raghavendra K, Chandra D, Subbarao SK. An allele-specific polymerase chain reaction assay for the differentiation of members of the *Anopheles culicifacies* complex. J Biosci. 2004;29(3):275-80. doi: 10.1007/BF02702609.

18. Alam MT, Das MK, Dev V, Ansari MA, Sharma YD. PCR-RFLP method for the identification of four members of the *Anopheles annularis* group of mosquitoes (Diptera: Culicidae). Trans R Soc Trop Med Hyg. 2007;101(3):239-44. doi: 10.1016/j.trstmh.2006.03.007.

19. Singh OP, Nanda N, Chandra D, Jha D, Adak T, Dua VK, et al. Modified PCR-based assay for the differentiation of members of *Anopheles fluviatilis* complex in consequence of the discovery of a new cryptic species (species V). Malar J. 2020;19(1):96. doi: 10.1186/s12936-020-03172-1.

20. Raghavendra K, Cornele AJ, Reddy BP, Collins FH, Nanda N, Chandra D, et al. Multiplex PCR assay and phylogenetic analysis of sequences derived from D2 domain of 28S rDNA distinguished members of the *Anopheles culicifacies* complex into two groups, A/D and B/C/E. Infect Genet Evol. 2009;9(2):271-7. doi: 10.1016/j.meegid.2008.12.007.

21. Beebe NW. DNA barcoding mosquitoes: advice for potential prospectors. Parasitology. 2018;145(5):622-633. doi: 10.1017/S0031182018000343.

22. Sindhania A, Das MK, Sharma G, Surendran SN, Kaushal BR, Lohani HP, et al. Molecular forms of *Anopheles subpictus* and *Anopheles sundaicus* in the Indian subcontinent. Malar J. 2020;19(1):417. doi: 10.1186/s12936-020-03492-2.

23. Djadid ND, Gholizadeh S, Aghajari M, Zehi AH, Raeisi A, Zakeri S. Genetic analysis of rDNA-ITS2 and RAPD loci in field populations of the malaria vector, *Anopheles stephensi* (Diptera: Culicidae): implications for the control program in Iran. Acta Trop. 2006; 97(1):65-74. doi: 10.1016/j.actatropica.2005.08.003.

24. Alam MT, Bora H, Das MK, Sharma YD. The type and mysorensis forms of the *Anopheles stephensi* (Diptera: Culicidae) in India exhibit identical ribosomal DNA ITS2 and domain-3 sequences. Parasitol Res. 2008;103(1):75-80. doi: 10.1007/s00436-008-0930-7.

25. Bhinder P, Chaudhry A, Barna B, Kaur S. Imidacloprid and Thiamethoxam Induced Mutations in Internal Transcribed Spacer 2 (ITS2) of *Anopheles stephensi*. Toxicol Int. 2012;19(2):201-6. doi: 10.4103/0971-6580.97223

26. Munawar K, Saleh A, Afzal M, Qasim M, Khan KA, Zafar MI, et al. Molecular characterization and phylogenetic analysis of anopheline (Anophelinae: Culicidae) mosquitoes of the Oriental and Afrotropical Zoogeographic zones in Saudi Arabia. Acta Trop. 2020;207:105494. doi: 10.1016/j.actatropica.2020.105494.

27. Livak KJ. Organization and mapping of a sequence on the *Drosophila melanogaster* X and Y chromosomes that is transcribed during spermatogenesis. Genetics. 1984; 107(4):611-34.

28. Christophers SR. The fauna of British India, including Ceylon and Burma. 1933, Vol IV., Diptera, Family Culicidae, Tribe Anophelini; London: Taylor and Francis. 371 pp.

29. Beebe NW, Saul A. Discrimination of all members of the *Anopheles punctulatus* complex by polymerase chain reaction--restriction fragment length polymorphism analysis. Am J Trop Med Hyg. 1995;53(5):478-81. doi: 10.4269/ajtmh.1995.53.478.

30. Litvaitis MK, Nunn G, Thomas WK, Koche TD. A molecular approach for the identification of *Meiofaunal turbellarians* (Platyhelminthes, Turbellaria). Marine Biol. 1994; 120, 437–42.

31. Campbell BC, Steffen-Campbell JD, Werren JH. Phylogeny of the Nasonia species complex (Hymenoptera: Pteromalidae) inferred from an internal transcribed spacer

(ITS2) and 28S rDNA sequences. Insect Mol Biol. 1993; 2(4):225-37. doi: 10.1111/j.1365-2583.1994.tb00142.x.

32. Zhang W, Tian W, Gao Z, Wang G, Zhao H. Phylogenetic Utility of rRNA ITS2 Sequence-Structure under Functional Constraint. Int J Mol Sci. 2020 3; 21(17):6395. doi: 10.3390/ijms21176395.

33. Bezzhonova OV, Goryacheva II. Intragenomic heterogeneity of rDNA internal transcribed spacer 2 in *Anopheles messeae* (Diptera: Culicidae). J Med Entomol. 2008; 45(3):337-41. doi: 10.1603/0022-2585(2008)45[337:ihorit]2.0.co;2.

34. Fairley TL, Kilpatrick CW, Conn JE. Intragenomic heterogeneity of internal transcribed spacer rDNA in neotropical malaria vector *Anopheles aquasalis* (Diptera: Culicidae). J Med Entomol. 2005; 42(5):795-800. doi: 10.1093/jmedent/42.5.795.

35. Li C, Wilkerson RC. Intragenomic rDNA ITS2 variation in the neotropical *Anopheles* (Nyssorhynchus) *albitarsis* complex (Diptera: Culicidae). J Hered. 2007; 98(1):51-9. doi: 10.1093/jhered/esl037.

36. Onyabe DY, Conn JE. Intragenomic heterogeneity of a ribosomal DNA spacer (ITS2) varies regionally in the neotropical malaria vector *Anopheles nuneztovari* (Diptera:Culicidae). Insect Mol Biol. 1999; 8(4):435-42. doi: 10.1046/j.1365-2583.1999.00134.x.

37. Sharma D, Lather M, Dykes CL, Dang AS, Adak T, Singh OP. Disagreement in genotyping results of drug resistance alleles of the *Plasmodium falciparum* dihydrofolate reductase (Pfdhfr) gene by allele-specific PCR (ASPCR) assays and Sanger sequencing. Parasitol Res. 2016;115(1):323-8. doi: 10.1007/s00436-015-4750-2.

Table 1: Number of samples subjected to direct sequencing of rDNA regions

Population	Geographical coordinates	rDNA region	
		ITS2	28S (d1-d3)
1. Delhi	28.63, 77.15	2	0
2. Gurugram (Haryana)	28.43, 76.97	4	4
3. Nuh (Haryana)	28.10, 76.99	4	4
4. Alwar (Rajasthan)	27.55, 76.62	4	4
5. Ranchi (Jharkhand)	23.36, 85.32	9	9
6. Raipur (Chhatishgarh)	21.26, 81.63	1	1
7. Gadchiroli (Maharashtra)	20.18, 79.99	2	2
8. Goa (UT)	15.36, 74.03	4	4
9. Bengaluru (Karnataka)	13.01, 77.61	2	2
10. Mangalore (Karnataka)	12.92, 74.85	4	4
11. Mysuru (Karnataka)	12.31, 76.64	3	3
12. Chennai (Tamil Nadu)	13.06, 80.23	5	5
13. Laboratory colony: "type form" (Jodhpur, Rajasthan)		2	2
14. Laboratory colony: "*var mysorensis*" (Jodhpur, Rajasthan)		2	2
		48	46

Table 2: Distribution of haplotypes in clonal DNA samples (nucleotide base position numbering based on the alignment displayed in Figure 2)

Haplotypes	Base positions					Number of clones			
	1 7 2	1 8 2	1 8 3	3 8 3	4 0 1	Nuh (northern India)	Gadchiroli (central India)	Bengaluru (southern India)	Total
Hap-1	C	C	A	C	G	12	11	7	30
Hap-2	T	-	-	C	G	4	2	3	9
Hap-3	T	-	-	C	A	3	1	3	7
Hap-4	C	-	-	T	G	3	2	1	6
Hap-5	C	C	A	T	G	2	1	2	5
Hap-6	C	C	A	C	A	1	0	2	3
Hap-7	C	C	A	T	A	0	0	1	1
Hap-8	T	C	A	T	G	0	0	1	1
Hap-9	C	-	-	C	A	0	1	0	1
Hap-10	C	-	-	C	G	0	1	0	1
Total						25	19	20	64

Figure 1 Geographical location of mosquito sampling

Figure 2: Alignment of haplotypes of complete ITS2 sequence of *An. stephensi* (the flanking 5.8S and 28S rDNA sequences are not shown here). The polymorphic loci have been highlighted (SNPs at three base positions, i.e., 172, 383 and 401; two bp indel at base positions 182-183). Dot represents similarity with Hap-1 sequence and dash represents gap in nucleotide sequence.

Figure 3 Snapshots of portions of ITS2 chromatogram (a) forward sequence, and (b) reverse sequence chromatogram (reverse compliment) showing polymorphic sites. Polymorphism 172C>T and 182_183del can be seen in forward sequence and 372C>T and 401G>A can be seen in reverse sequence chromatogram. The ambiguity (mixed bases) in sequences can be seen from the point of indel onwards (a).

Supplementary information

S1: Annotation features of GenBank entries

```
>Feature gb|MW676288
<1    52          5.8S ribosomal RNA
53    522         internal transcribed spacer 2
523   >1679       28S ribosomal RNA
>Feature gb|MW676289
<1    52          5.8S ribosomal RNA
53    520         internal transcribed spacer 2
521   >1677       28S ribosomal RNA
>Feature gb|MW676290
<1    52          5.8S ribosomal RNA
53    520         internal transcribed spacer 2
521   >1677       28S ribosomal RNA
>Feature gb|MW676291
<1    52          5.8S ribosomal RNA
53    520         internal transcribed spacer 2
521   >1677       28S ribosomal RNA
>Feature gb|MW676292
<1    52          5.8S ribosomal RNA
53    522         internal transcribed spacer 2
523   >1679       28S ribosomal RNA
>Feature gb|MW676293
<1    52          5.8S ribosomal RNA
53    522         internal transcribed spacer 2
523   >1679       28S ribosomal RNA
>Feature gb|MW676294
<1    52          5.8S ribosomal RNA
53    522         internal transcribed spacer 2
523   >1679       28S ribosomal RNA
>Feature gb|MW676295
<1    52          5.8S ribosomal RNA
53    522         internal transcribed spacer 2
523   >1679       28S ribosomal RNA
>Feature gb|MW732930
<1    52          5.8S ribosomal RNA
53    520         internal transcribed spacer 2
521   >1677       28S ribosomal RNA
>Feature gb|MW732931
<1    52          5.8S ribosomal RNA
53    520         internal transcribed spacer 2
521   >1677       28S ribosomal RNA
```