**Identification of arginine phosphorylation in *Mycolicibacterium smegmatis***

Emmanuel C. Ogbonna[1], Henry R. Anderson[2] and Karl R. Schmitz[1,2]*

Department of Biological Sciences[1]
Department of Chemistry & Biochemistry[2]
University of Delaware
Newark DE, 19716

*corresponding author: schmitzk@udel.edu

running title: Identification of arginine phosphorylation in *M. smegmatis*

**Abstract**

Tuberculosis is a leading cause of worldwide infectious mortality. The prevalence of multidrug-resistant *Mycobacterium tuberculosis* (*Mtb*) infections drives an urgent need to exploit new drug targets. One such target is the ATP-dependent protease ClpC1P1P2, which is strictly essential for viability. However, few proteolytic substrates of mycobacterial ClpC1P1P2 have been identified to date. Recent studies in *Bacillus subtilis* have shown that the orthologous ClpCP protease recognizes proteolytic substrates bearing post-translational arginine phosphorylation. While several lines of evidence suggest that ClpC1P1P2 is similarly capable of recognizing phosphoarginine-bearing proteins, the existence of phosphoarginine modifications in mycobacteria has remained in question. Here, we confirm the presence of post-translational phosphoarginine modifications in *Mycolicibacterium smegmatis* (*Msm*), a nonpathogenic surrogate of *Mtb.* Using a phosphopeptide enrichment workflow coupled with shotgun phosphoproteomics, we identify arginine phosphosites on several functionally diverse targets within the *Msm* proteome. Interestingly, phosphoarginine modifications are not upregulated by heat stress, suggesting divergent roles in mycobacteria and *Bacillus*. Our findings provide new evidence supporting the existence of phosphoarginine-mediated proteolysis by ClpC1P1P2 in mycobacteria and other actinobacterial species.

**Introduction**

Tuberculosis is a leading cause of worldwide infectious mortality, ranking above HIV/AIDS and Ebola, and is one of the top ten leading causes of death overall (WHO, 2021)**.** Advances in diagnosis, vaccinations and therapeutics have reduced tuberculosis morbidity and mortality. However, the prevalence of multidrug-resistance in the causative bacterium, *Mycobacterium tuberculosis* (*Mtb*), remains high (WHO, 2021). These statistics underscore the urgent need to discover new drugs and exploit new molecular targets. One promising target is the mycobacterial Clp protease. Several studies show that Clp protease components are strictly essential for *Mtb* viability (Sassetti et al., 2003; Griffin et al., 2011; Raju et al., 2012; Raju et al., 2014; DeJesus et al., 2017) and are viable targets for anti-*Mtb* therapeutics (Compton et al., 2013; Gavrish et al, 2014; Gao et al, 2015; Moreira et al., 2015; Famulla et al., 2016; Moreno-Cinos et al., 2019).

Clp proteases mechanically unfold and destroy native cytosolic proteins (Sauer and Baker; 2011; Baker and Sauer 2012). These large enzymatic complexes consist of a core peptidase and an associated hexameric ATP-dependent unfoldase. The mycobacterial Clp peptidase, ClpP1P2, is a heteromer composed of distinct ClpP1 and ClpP2 rings that stack face-to-face to create a barrel-shaped tetradecamer (Akopian et al., 2012; Schmitz et al., 2014; Schmitz et al., 2014b; Li et al., 2016; Vahidi et al., 2020). ClpC1 is one of two mycobacterial unfoldases (the other is ClpX) that can assemble with ClpP1P2 to form a functional protease. ClpC1 is an 848-residue protein with a globular N-terminal domain (NTD) and two AAA+ (ATPases associated with various cellular activities) modules. Ring-shaped hexamers of ClpC1 dock coaxially on the surface of the ClpP2 face of the peptidase (Leodolter et al., 2015; Nagpal et al., 2019). Protein substrates destined for destruction by ClpC1P1P2 must first be recognized by ClpC1. The unfoldase therefore serves a critical regulatory role in proteolysis.

Clp proteases have become major targets of novel drug development against *Mtb* and other pathogenic bacteria. Several classes of antimicrobials specifically target the Clp peptidase,

including dysregulators (e.g., acyldepsipeptides (Brötz-Oesterhelt et al., 2005; Kirstein et al,

2009)), activators (e.g., sclerotiamide (Lavey et al., 2016)), and catalytic inhibitors (e.g., β-

lactones and boronate compounds (Compton et al., 2013; Akopian et al., 2015)). Importantly,

multiple cyclic peptides have been discovered that exhibit anti-*Mtb* activity through dysregulation

of ClpC1, including cyclomarin A, lassomycin, ecumicin, rufomycin and metamarin (Schmitt et al.,

2011; Gavrish et al., 2014; Gao et al., 2015; Wolf et al., 2019; Li et al., 2021). These compounds

bind to the ClpC1 NTD and disrupt proteolysis by uncoupling unfoldase activity from proteolysis

or by causing uncontrolled degradation of cellular proteins. The antimicrobial activity of ClpC1-

targeting compounds underscores the importance of ClpC1P1P2 in mycobacteria. However, the

specific proteolytic functions responsible for its essentiality remain obscure, and only a handful of

protein substrates have been identified (Barik et al., 2010; Gopal et al., 2020; Lunge et al., 2020;

Ziemski et al., 2021).

An expanded understanding of the physiological roles played by ClpC1P1P2 would bolster

efforts to develop antimicrobial compounds. In other bacteria, Clp proteases participate in cellular

processes ranging from targeted pathway regulation to cell-wide protein quality control (Gerth et

al., 2008; Trentini et al., 2016). Multiple mechanisms of substrate recognition have been

described, including direct interactions with the unfoldase (Gopal et al., 2020; Lunge et al., 2020)

and indirect recognition with aid of adaptors (Marsee et al., 2018; Ziemski et al., 2021). Recent

studies in *Bacillus subtilis* and related Firmicutes have demonstrated that post-translational

arginine phosphorylation marks some proteins for destruction by ClpCP (Schmidt et al., 2014;

Trentini et al., 2016; Junker et al., 2018)*. Dual phosphoarginine (pArg) binding sites on the *B.*

*subtilis* ClpC NTD allow ClpCP to recognize phosphoarginylated proteins as proteolytic substrates

(Trentini et al., 2016). Phosphoarginine-mediated proteolysis in these bacteria is upregulated

during stress through activation of the arginine kinase McsB (Fuhrmann et al., 2009).

Phosphoproteomic studies have uncovered widespread arginine phosphorylation during heat

stress, implicating ClpCP in turnover of misfolded proteins (Schmidt et al., 2014; Trentini et al.,

2016; Trentini et al., 2018). Additionally, McsB regulates the global stress response through

targeted phosphorylation of the negative transcriptional regulators CtsR and HrcA (Fuhrmann et

al., 2009; Schmidt et al., 2014; Suskiewicz et al., 2019). Proteolysis of these targets by ClpCP

allows transcriptional activation of stress response genes (Fuhrmann et al., 2009). Interestingly,

ClpC and ClpP are themselves targets of McsB, and specific pArg sites on ClpC are required for

its activation by McsB (Schmidt et al., 2014, Elsholz et al., 2012). These studies underline the role

of arginine phosphorylation in Firmicutes as both a degradation signal and a regulatory

mechanism.

Several lines of evidence suggest that an analogous pArg-mediated proteolytic pathway

exists in mycobacteria. Sequence and structural data reveal overall homology between the NTDs

of *Mtb* ClpC1 and *B. subtilis* ClpC, as well as strong conservation of the residues surrounding the

pArg-binding sites (Trentini et al., 2016; Weinhäupl et al., 2018). Moreover, *in vitro* experiments

confirm that the *Mtb* ClpC1 NTD does indeed interact with both free pArg and with arginine-

phosphorylated model substrates (Weinhäupl et al., 2018). However, the existence of pArg-

mediated proteolysis in mycobacteria remains in question, as no mycobacterial McsB homologs

are known and phosphoarginine modifications have not yet been described in these bacteria.

Here, we confirm the existence of post-translational phosphoarginine modifications in

*Mycolicibacterium smegmatis* (*Msm*), a nonpathogenic surrogate of *Mtb.* Using a phosphopeptide

enrichment workflow coupled to shotgun phosphoproteomics, we identify arginine phosphosites

on several protein targets within the *Msm* proteome. Our findings suggest that these modifications

are widespread among actinobacterial species.

**Materials and Methods**

*ClpC1 Sequence Analysis* — Actinobacterial homologs of *Msm* ClpC1 were identified using

HMMER (Eddy, 1995) and aligned using Clustal Omega (Sievers et al., 2011). Fragmentary

sequences were omitted from analysis. To reduce overrepresentation of similar taxa (e.g.,

multiple *Mtb* strains), alignments were pruned such that no two sequences had greater than 90%

sequence identity. Positional conservation scores were exported from Jalview (Waterhouse et al.,

2009) and plotted as a heatmap in Prism (Graphpad). The sequence logo of the NTD region was

generated using WebLogo (Crooks et al., 2004).

*Protein Purification and Binding Assays* — The nucleotide sequence encoding the NTD of ClpC1

(codons 1 – 147; referred to as ClpC1$^{NTD}$) was amplified from *M. smegmatis* MC$^2$155 gDNA

(ATCC) and cloned into a pET22b-derived vector (EMD Millipore) in frame with a C-terminal

LPETGG sortase recognition sequence (Glasgow et al., 2016) and 6xHis tag. ClpC1$^{NTD}$ was

overexpressed in *E. coli* strain ER2566 (NEB) by induction with 0.5 mM isopropyl β-D-1-

thiogalactopyranoside at 30°C for 4 h. Cells were harvested by centrifugation, resuspended in

Lysis Buffer (25 mM HEPES, 500 mM NaCl, 10 mM imidazole, 10% glycerol, pH 7.5), and lysed

by sonication. ClpC1$^{NTD}$ was purified from clarified lysate by Ni-NTA (Marvelgent Biosciences)

and anion exchange (Source Q, Cytiva) chromatography. ClpC1$^{NTD}$ was fluorescently labeled via

sortase transpeptidation (Glasgow et al., 2016), by 2 h incubation with sortase A (~1:30 molar

ratio) and a 2-fold molar excess of a Gly-Gly-Asn-Lys-(fluorescein-isothiocyanate) peptide

(Biomatik) in PBS Buffer (10 mM Na$_2$HPO$_4$, 1.8 mM KH$_2$PO$_4$, 137 mM NaCl, 2.7 mM KCl, 10%

glycerol, pH 7.4). Excess peptide was removed by size exclusion chromatography (Superdex 75,

Cytiva). Purified ClpC1$^{NTD-FITC}$ was concentrated and stored in CPD Buffer (25 mM HEPES, 200

mM NaCl, 10 mM MgCl$_2$, 0.1 mM EDTA, pH 7.0). Binding of phosphoarginine (MilliporeSigma) to

0.1 µM purified ClpC1$^{NTD-FITC}$ was assayed in CPD supplemented with 0.05% Tween 20 and 8

mg/ml BSA by microscale thermophoresis using a Monolith NT.115 (NanoTemper) (Jerabek-

Willemsen et al., 2014). Thermophoretic data were fit to a quadratic single site binding equation

in Prism (Graphpad).

*Cell Culture Conditions* — Liquid starter cultures of *M. smegmatis* MC$^2$155 were prepared in 20

mL Middlebrook broth base (HiMedia) containing 0.2% v/v glycerol (Fisher Scientific), 0.05% v/v

Tween 80, and grown for 48 hours at 37 °C with 250 rpm orbital shaking. Saturated starter cultures

were sub-cultured into 900 mL of fresh media at a starting $A_{600}$ ~0.02, and grown at 37 °C until

$A_{600}$ ~1.0. 500 mL of culture was collected and added to 400 mL of fresh media. Unstressed

control samples were further grown at 37 °C, while samples for heat stress were grown at 50 °C

for 21 hours. Cells were harvested by centrifugation at 9,000 × g for 20 min at 4 °C, and pellets

were resuspended in 5 mL of ice-cold lysis buffer (25 mM HEPES, 200 mM KCl, 10 mM MgCl$_2$,

0.1 mM EDTA, pH 7.5) containing 10 mM ATP (Fisher), 200 µL phosphatase inhibitor cocktail

(MilliporeSigma), 30 mM sodium pervanadate (Acros Organics), 3 mM NaF and 3 mM sodium

pyrophosphate (both from Fisher). Cells were lysed at high pressure using a microfluidizer

(Microfluidics). Lysates were clarified at 15,000 × g for 30 min at 4 °C, and supernatant was stored

at -80 °C prior to further processing. Total protein content was estimated by Bradford assay (Bio-

Rad).

*Filter-Aided Sample Preparation for Mass Spectrometry* — Replicate samples of clarified cell

lysates (approximately 10 mg total protein) were prepared for mass spectrometry using filter-aided

sample preparation (FASP) (Wiśniewski et al., 2009; Schmidt et al., 2014). Reduction of disulfide

bonds was achieved by the addition of 200 mM dithiothreitol (DTT; Fisher), followed by incubation

at 56 °C for 50 min.  Afterwards, samples were diluted in 7 mL of 0.1 M Tris (Gold Biotechnology),

pH 7.5 containing 8 M urea and 25 mM 2-iodoacetamide (both from Acros Organics) in 15 mL

tubes and incubated in the dark at room temperature for 45 min to carbamidomethylate cysteines.

After alkylation, samples were transferred to an Amicon filter (10,000 MWCO, MilliporeSigma),

washed twice with 5 mL 0.1 M Tris, pH 7.5, containing 8 M urea by centrifugation at 4,000 × g,

then washed twice with 5 mL 0.1 M Tris, pH 7.5, containing 4 M urea, and finally twice with 5 mL

100 mM ammonium bicarbonate (Honeywell). After the last wash step, retentate was reduced to

less than 1 mL by spin concentration.

*In-Solution Trypsin Proteolytic Cleavage* — MS Grade Trypsin Protease (Pierce) was dissolved

to obtain a 20 µg/mL stock in 100 mM ammonium bicarbonate. Trypsin digestion of processed *M.*

*smegmatis* lysates was performed in an Amicon filter unit at an approximately 1500:1

protein:trypsin ratio overnight at 37 °C. Upon digestion, peptides were recovered by centrifugation

at 4,000 × g for 10 minutes, followed by the addition of 0.5 M NaCl. Eluted samples were dried

under vacuum in a SpeedVac centrifuge (Thermo Fisher).

*TiO$_2$ Enrichment of Phosphopeptides* — Phosphopeptides were enriched from tryptic peptide

samples using Titansphere TiO$_2$ beads (5 µm; GL Sciences) in buffer conditions that limited acid

hydrolysis of phosphoarginine (Thingholm et al., 2006; Schmidt et al., 2014; Trentini et al., 2016).

5 mg TiO$_2$ resin was resuspended in 1 mL Binding Buffer (300 mg/mL lactic acid, 12.5% acetic

acid, 60% acetonitrile, 0.2% heptafluorobutyric acid, pH 4 with NH$_4$OH). Lyophilized peptide

samples were re-dissolved in the TiO$_2$ suspension, incubated for 35 min at 20 °C with gentle

agitation, then transferred to graphite spin columns (Thermo Fisher). Unbound peptides were

removed by a first wash step with 150 µL Binding Buffer and spun at 2000 × g for 1 min, followed

by three wash steps using 400 µL of Wash Solution A (200 mg/ml lactic acid, 75% acetonitrile,

2% trifluoroacetic acid, 2% heptafluorobutyric acid), Wash Solution B (200 mg/mL lactic acid, 75%

acetonitrile, 10% acetic acid, 0.1% heptafluorobutyric acid, pH 4 with NH$_4$OH), Wash Solution C

(80% acetonitrile, 10% acetic acid), respectively. The resin was then incubated with 100 µL

Elution Solution 1 (1% NH$_4$OH, 30 mM ammonium phosphate) and Elution Solution 2 (1.25%

NH$_4$OH in 50% acetonitrile) for 15 minutes each. The eluate containing phosphopeptides was

collected by centrifugation after each incubation. To remove salts, samples were desalted using

a HyperSep C18 column (Thermo Scientific). Samples were lyophilized and stored at -80 °C prior

to mass spectrometry.

*Liquid Chromatography - Tandem Mass Spectrometry (LC-MS/MS)* — Lyophilized and desalted

tryptic digests were resuspended in 20 µL 0.5% acetic acid (pH 4.5). An Orbitrap Eclipse MS

(Thermo Scientific) coupled with an Ultimate 3000 nano-LC system and a FAIMS Pro Interface

(Thermo Scientific) was used for the LC-MS/MS analysis. Peptide samples were first loaded onto

a trap column (PepMap C18; 2 cm x 100 µm I.D.) and afterwards separated at a flow rate of 300

nL/min on an analytical column (PepMap C18, 3.0 µm; 10 cm x 75 mm I.D.; Thermo Scientific). A

binary buffer system (buffer A, 0.1% formic acid in water; buffer B, 0.1% formic acid in acetonitrile)

with a 165-min gradient (1% to 25% buffer B over 125 min; 25% to 32% buffer B in 10 min, then

to 95% buffer B over 3 min; back to 1% B in 5 min, and stay equilibration at 1% B for 20 min) was

utilized. To achieve field asymmetric ion mobility spectrometry (FAIMS) separation, multiple CVs

(-45, -60 and -80) were applied. For all experiments, the survey scans (MS1) were acquired over

a mass range of 375-1500 m/z at a resolution of 60,000 in the Orbitrap. Isolation of precursors

was done with a width of 1.6 m/z for MS/MS acquisition. They were subsequently fragmented with

higher energy collisional dissociation (HCD) using 30% collision energy with a maximum injection

time of 100 ms, and collected in Orbitrap at 15,000 resolution. The dynamic exclusion was set to

60 s, and was shared across different FAIMS experiments. LS-MS/MS data was collected in

independent biological triplicates.

*Mass Spectrometry Data Analysis* — Proteomic analysis was performed in the Proteome

Discoverer software suite (version 2.2, Thermo Fisher). Raw data was searched against the *M.*

*smegmatis* (strain ATCC 700084 / MC[2] 155) UniProt Reference Proteome (Proteome ID

UP000000757; 6,602 entries in total) using Sequest HT (University of Washington and Thermo

Fisher) (Eng et al., 1994). Iodoacetamide-mediated cysteine carbamidomethylation was set as a

static modification, while methionine oxidation and phosphorylation of arginine, serine, threonine,

tyrosine and histidine residues were entered as dynamic modifications. Complete trypsinization

with a maximum of two missed cleavages was allowed. Precursor mass tolerance was set at 10

ppm, while allowing fragment ions to have a mass deviation of 0.02 Da for the HCD data.

Validation of peptide-spectrum matches (PSM) based on q-value was done using Percolator -

with target false discovery rates (FDR) of 1% and 5% for stringent and relaxed validation,

respectively. The false discovery rate of high confidence protein and peptide identification was at

1%. Localization probability of phosphopeptide hits were analyzed using the PhosphoRS (ptmRS)

node of the Proteome Discoverer software (Taus et al., 2011). Only modifications with a

PhosphoRS (ptmRS) score ≥75% were selected.

*Gene Ontology Annotation Analysis* — Gene Ontology (GO) annotation was performed using the

OmicsBox software suite (Conesa et al., 2005). Homologs were identified by BLAST against the

SwissProt/UniProt database (Bairoch et al., 1997), with a minimum expectation value $10^{-3}$.

Homolog annotations were compiled from the InterPro database (Hunter et al., 2009). BLAST and

InterPro results were used to generate GO terms in terms of biological process. For proteins still

unannotated, direct Uniprot BLAST was performed, and tentative assignment of functional group

was based on those of obtained homologs.

*Sequence and Structural Analysis of Arginine Phosphosites* — For alignment analysis, a 21-mer

sequence was obtained containing 10 residues flanking each side of phosphoarginine. All 21-

mers were then aligned in BioEdit (Hall, 1999). Structural information, if available, was obtained

from Protein Data Bank (PDB) (Berman et al, 2000). Predicted structures were retrieved using

neural network-based machine learning model, AlphaFold (version 2, Jumper et al., 2021) and

molecular images were prepared in PyMOL (version 2.5.2, Schrödinger). Protein BLAST (Basic

Local Alignment Search Tool) of arginine-phosphorylated proteins was performed in the NCBI

suite (Gish and States, 1994). Analysis was restricted to the Corynebacterineae suborder.

Sequences were aligned using Clustal Omega (Sievers et al., 2011), and conservation and

consensus values obtained in Jalview (Waterhouse et al., 2009).

*Data Availability* — The mass spectrometry data from this publication have been submitted to the

ProteomeXchange Consortium (http://proteomecentral.proteomexchange.org) via the PRIDE

partner repository (Perez-Riverol et al., 2019), and assigned the identifier PXD032083.

**Results**

***Phosphoarginine Binding Sites in the ClpC1 NTD Are Conserved Across Actinobacteria –***

ClpC1 is an essential enzyme in *Mtb* and *Msm* (Sassetti et al., 2003; DeJesus et al., 2017), yet

few specific cellular roles or proteolytic substrates of the ClpC1P1P2 protease are known. To gain

insight into its cellular function, we began by examining ClpC1 sequence conservation across the

phylum Actinobacteria by constructing an alignment of 1,195 actinobacterial orthologs (**Fig. 1A**).

Like most other type-II Clp unfoldases, ClpC1 possesses a ~150 amino acid N-terminal domain

(NTD), which likely participates in regulation and substrate recognition, but lacks mechanical

function or a direct role in catalysis (Schlothauer et al, 2003; Marsee et al., 2018). Surprisingly,

the NTD is as well or better conserved than the D1 and D2 AAA+ ATPase rings, suggesting that

the NTD has important and conserved functions across Actinobacteria. This may help explain

why naturally occurring antibiotics that target and dysregulate ClpC1 have evolved to specifically

bind the NTD (Kirstein et al., 2009; Schmitt et al., 2011; Vasudevan et al., 2013; Gavrish et al.,

2014; Bürstner et al., 2015; Gao et al., 2015; Jung et al., 2017).

In *Bacillus subtilis* and other Firmicutes, ClpCP recognizes phosphorylated arginine

residues as degradation signals via twin binding sites on opposite ends of the ClpC NTD (Trentini

et al., 2016). Examination of actinobacterial ClpC1 sequences revealed that the residues known

to be important for phosphoarginine binding in *B. subtilis* are conserved across actinobacterial

orthologs (**Fig. 1B**). We found that we could readily model bound phosphoarginines in an existing

structure of the *M. tuberculosis* ClpC1 NTD (Wolf et al., 2020), with minor side chains

rearrangements, based on the binding mode observed to the *B. subtilis* ClpC NTD (Trentini et al.,

2016) (**Fig. 1C,D**). Finally, we directly assessed binding of pArg to purified *Msm* ClpC1$^{NTD}$ by

microscale thermophoresis (**Fig. 1E**). Phosphoarginine bound with an affinity of 1.4 µM, similar to

the ~5 µM affinity previously reported to the *Mtb* ClpC1 NTD (Weinhäupl et al., 2018). The ability

of the ClpC1$^{NTD}$ to bind pArg and the conservation of pArg binding modules across Actinobacteria

provides strong indirect evidence for the existence of this post-translational modification in this

phylum.

***Identification of Phosphoarginine Modifications in Mycolicibacterium smegmatis –*** To test

for the physiological existence of arginine phosphorylation, we probed dot blots of several

bacterial lysates with a phosphoarginine-specific antibody (**Fig. 1F**) (Fuhrmann et al., 2015). In

line with prior studies, signal was detected in *B. subtilis* lysate (Fuhrmann et al., 2015), but not in

lysate from *E. coli*, which lacks arginine phosphorylation (Potel et al., 2018, Fu et al., 2020). We

also observed a positive immunoblot reaction in lysates from *Msm* and a second actinobacterium,

*Corynebacterium jeikeium*. These observations suggest that pArg modifications occur at least

within    the    suborder    Corynebacterineae,    which    encompasses    *Corynebacterium*,

*Mycolicibacterium*, and *Mycobacterium*.

To determine which specific cellular proteins carry pArg modifications in *Msm*, we

employed an unbiased shotgun proteomics approach. Phosphoarginine is acid-labile and has a

short half-life in the acidic TFA-containing solvent systems typically used for proteomic sample

preparation and LS-MS/MS (Schmidt et al., 2014). Optimized protocols for arginine

phosphoproteomics have been reported that utilize alternative solvent systems at pH ≥ 4 for most

steps (Schmidt et al., 2014; Trentini et al., 2016; Trentini et al., 2018). We adapted these methods

to our workflow for sample preparation, phosphopeptide enrichment, and LC-MS/MS, thereby

minimizing phosphoarginine hydrolysis (**Fig. 2A**). Since stress conditions such as heat shock

upregulated the occurrence of this modification in *B. subtilis* (Schmidt et al., 2014), we analyzed

lysates from *Msm* cultures grown at either normal growth temperature (37 °C) or 50 °C for 21

hours.

A similar number (~3000) of proteins were observed in samples from normal growth (3147,

3004, and 3145 proteins in the three replicates, respectively) and heat stress conditions (2969,

2724 and 2899 proteins) (**Fig. 2B**). This suggests that at this level of stress, compensatory stress

responses do not involve a complete shutdown of translational machinery or dramatic

upregulation of proteolytic activity. Typically, a major part of stress response to heat shock is the

increased expression of heat shock proteins or chaperones. We evaluated the differential levels

of observed proteins based on Sequest HT scores. As shown in **Fig. 2C,** several proteins were

observed at significantly higher levels ($p$-value ≤ 0.05) in heat shock than in normal growth

conditions. These included Spore protein (Msmeg_5611)*,* which belongs to the Hsp20 small heat

shock protein family and was enriched over 2-fold upon heat shock. Other chaperone proteins

enriched in heat stressed samples included DnaJ1, DnaK, and the SecB-like chaperone SmegB.

We also performed comparative GO annotation analysis of strongly enriched/depleted proteins,

whose levels differed significantly and by a magnitude of at least two-fold between conditions

(**Supplemental Figure S1; Supplemental Table S1**). Proteins involved in amino acid, lipid and

noncanonical metabolism, along with redox proteins, were strongly depleted during heat stress.

Spore protein was the only stress response protein strongly enriched under heat shock; no protein

involved in stress response was strongly depleted. Taken together, this differential expression

suggests the activation of canonical heat shock response in these samples.

Using Proteome Discoverer we identified arginine-phosphorylated sites in proteins from

these samples. Only peptides with a phosphosite localization probability (PhosphoRS/ptmRS

score) ≥75% were selected (Bäsell et al., 2014). We unambiguously localized six phosphoarginine

sites in six different proteins (**Fig. 3A; Supplemental Table S2; Supplemental Figure S2**). Most

pArg sites were observed in multiple replicates, in both stressed and unstressed samples (**Fig.**

**3A**). The existence of phosphoarginine modifications in our samples implies the existence of an

unidentified *Msm* pArg kinase.

***Functional Characterization of Arginine-Phosphorylated Proteins –*** We examined the known

or predicted functions of the six proteins observed to carry pArg modifications and found that they

have diverse functions with no obvious common physiological role (**Fig. 3A**). PhoU1 is a central

regulator of the SenX3-RegX3 two-component system responsible for uptake of inorganic

phosphate ($P_i$) during phosphate starvation (Brokaw et al., 2017). This raises the possibility that

pArg modifications are linked to $P_i$ availability. Two other pArg-bearing targets were metabolic

enzymes: the multifunctional 2-oxoglutarate metabolism enzyme Kgd, involved in the tricarboxylic

acid cycle (Wagner et al., 2011), and the glycerol dehydratase large subunit (Msmeg_1547),

which potentially contributes to the catabolic pathway involving the glycerol dehydration reaction

which yields 3-hydroxypropanal in the presence of adenosylcobalamin coenzyme (**Fig. 3A;**

**Supplemental Figure S2**) (Liu et al., 2010; Nasir et al., 2020). pArg may thus play a role in

regulating cellular metabolic pathways. Another phosphosite was detected in the transmembrane

xanthine/uracil permease (Msmeg_1293), a nucleobase transporter (**Figs. 3A and 3C**). A site

was found on the hemerythrin domain-containing protein Msmeg_0907, which belongs to a class

of $O_2$-binding proteins involved in signal transduction, response to $H_2O_2$, oxygen sensing and nitric

oxide reduction (Isaza et al., 2006; Li et al. 2015; Lo et al., 2016; Ma et al., 2020). Finally, a site

was observed on Msmeg_0879, a small (48 aa) uncharacterized protein predicted to be

predominantly disordered (**Figs. 3A and B**). Homologs of Msmeg_0879 were detected in close

relatives of *Msm*, but are not widely conserved in actinobacteria.

***Structural and Physicochemical Analysis of Arginine Phosphorylation Sites –*** We assessed

whether arginine phosphorylation occurred at sites with particular physicochemical properties.

We aligned 21-residue sequence segments centered on each unique phosphoarginine site (**Fig.**

**4A**) but observe no clear consensus motif. Ala, Leu, Val and Gly appear to be common in flanking

positions, although we note that these are the four most abundant amino acids in the *M.*

*smegmatis* proteome (13%, 10%, 9%, 9% of total, respectively) (**Supplemental Table S3**). The

sequence diversity surrounding pArg positions suggests that target discrimination is guided by

characteristics other than primary sequence.

We next examined structural characteristics of phosphorylated positions, based on

predicted structures generated by AlphaFold2 (Jumper et al., 2021), except for the small protein

Msmeg_0879 for which structure prediction failed. (Notably, the AlphaFold2 prediction for kgd

was virtually identical to its reported X-ray structure (PDB ID: 2XT6) with RMSD ~0.5 A in the

region of the phosphosite. (Wagner et al., 2011)) As expected for this charged residue, arginine

phosphosites were solvent exposed (**Supplemental Figure S3**), but were located on structured

elements rather than loops. Interestingly, in all structures, arginine phosphosites occurred

proximal to the beginning or the end of an alpha helix (**Figs. 4B-F**), which may reflect a structural

constraint important for recognition by an arginine kinase.

Finally, we assessed the sequence conservation of arginine at these positions. We aligned

the arginine-phosphorylated *Msm* proteins with homologs within the Corynebacterineae suborder,

and analyzed the positional conservation of arginine at the respective positions. As shown in **Fig.**

**5**, conservation varies. The phosphosite arginine was well conserved in Kgd (99.8%),

Msmeg_0907 (79%), and Msmeg_0879 (62%) (**Figs 5A-C**). Conservation was lower in

Msmeg_1547 (34.1%) and PhoU1 (32.63%) (**Figs. 5D and 5E**). Arg was rarely present in this

position in homologs of Msmeg_1293 (0.23%) (**Fig. 5F**). In cases where the phosphorylated

arginine was well conserved, it may indicate that phosphorylation at this position is a conserved

phenomenon with functional or regulatory significance.

**Discussion**

Protein phosphorylation is a ubiquitous mechanism of signal propagation and pathway

regulation in bacteria (Macek et al., 2019). While examples of His, Asp, Ser, Thr, and Tyr

phosphorylation are widespread (Chow et al., 1994; Canova et al., 2008; Prisic et al., 2010;

Nakedi et al., 2015), the importance of arginine phosphorylation has become apparent only

recently. A growing body of data links pArg modifications to protein quality control pathways and

spore germination in Firmicutes, including *B. subtilis* (Fuhrmann et al., 2009; Elsholz et al., 2012;

Schmidt et al., 2014; Fuhrmann et al., 2016; Trentini et al., 2016; Junker et al., 2018; Weinhäupl

et al., 2018; Zhou et al., 2019). However, it has remained unclear how prevalent pArg

modifications are in other bacterial phyla. Here we provide direct evidence for the existence of

arginine phosphorylation in *Mycolicibacterium smegmatis*, an actinobacterium.

In *B. subtilis* and *S. aureus*, this modification is upregulated by stress (Fuhrmann et al.,

2009; Elsholz et al., 2012; Schmidt et al., 2014; Fuhrmann et al., 2016; Trentini et al., 2016; Junker

et al., 2018; Weinhäupl et al., 2018; Zhou et al., 2019). Hence, we performed our study by

examining pArg levels in cells grown under heat stress and normal conditions. While we saw

expected increases in levels of some heat shock proteins overall, arginine phosphorylation

abundance was not altered by heat shock, as occurs in firmicutes. It is possible that mycobacterial

pArg levels are linked to different stress stimuli that were not tested here. Alternatively, pArg may

have no relationship to the mycobacterial stress response, and may instead play a targeted

regulatory role.

The six arginine phosphosites localized here were far fewer than those identified in other

bacterial studies. Two independent studies in *B. subtilis* observed 121 and 217 sites in 87 and

134 proteins, respectively (Elsholz et al., 2012; Schmidt et al., 2014), while in *S. aureus* 207 sites

were identified in 126 proteins (Junker et al., 2018). On its face, the small number of pArg

modifications found in *Msm* argues against a proteome-wide quality control role, and appears

more consistent with targeted regulation of select proteins and processes. However, we note that

phosphoproteomic studies in Firmicutes utilized deletion strains lacking known arginine

phosphatases ([Bs]YwlE and [Sa]PtpB), which elevate levels of pArg-bearing proteins. No arginine

phosphatase has yet been identified in *Msm* or *Mtb*, thus our approach utilized wild-type *Msm*. If

a mycobacterial arginine phosphatase is eventually identified, it would be interesting to observe

the effect of its knockout or knockdown on pArg levels.

For arginine phosphorylation to be useful to the cell, it must presumably be applied

selectively through the regulated activity of arginine kinases. Many protein kinases recognize

substrates through characteristic sequence motifs surrounding the phosphosite. By contrast, our

examination of mycobacterial arginine phosphosites revealed no clearly enriched consensus

sequence around pArg. Prior analysis of Arg phosphosites in *B. subtilis* similarly revealed no

evident consensus sequence (Schmidt et al., 2014). This degeneracy suggests either that distal

interactions guide *Msm* arginine kinases to phosphosites or that other characteristics of target

proteins guide substrate selection. Supporting this latter possibility, we noted that all

phosphorylated arginine residues (for which structural models could be obtained) were found near

one end of an alpha helix, which may indicate a structural feature recognized by an arginine

kinase.

What role does arginine phosphorylation play in mycobacterial cells? In *B. subtilis*, pArg

tagging helps enforce protein quality control by marking misfolded proteins for destruction by

ClpCP (Elsholz et al., 2012; Schmidt et al., 2014; Trentini et al., 2016). Prior studies have

demonstrated that mycobacterial ClpC1P1P2 can degrade model substrates *in vitro* bearing pArg

modifications (Weinhäupl et al., 2018). The pArg-bearing proteins identified here may therefore

be recognized as proteolytic substrates by ClpC1P1P2. Alternatively, pArg may modulate some

aspect of the targets' function or interactions with other proteins. We observed pArg on PhoU1,

which helps regulate $P_i$ uptake by inhibiting the activity of the SenX3-RegX3 two-component

system when $P_i$ is readily available (Brokaw et al., 2017). Additionally, the signal transduction

protein Msmeg_0907, metabolic proteins (Kgd and Msmeg_1547), a transmembrane

xanthine/uracil transport protein (Msmeg_1293) and an uncharacterized protein Msmeg_0879

were found to be phosphorylated. In terms of essentiality, a prior study showed that the *Mtb*

homolog of Kgd is essential for viability (Sassetti et al., 2003); while another report showed that

PhoU1 is jointly essential with the PhoU2 protein (Msmeg_1605) for *in vitro Msm* growth (Brokaw

et al., 2017). More work is required to determine whether pArg modifications regulate these

pathways, and whether this modification plays an essential role in mycobacterial physiology.

This study lays the groundwork for future efforts to expound on the roles of

phosphoarginine modifications in mycobacteria. The immediate impediment to further

understanding this system is the absence of identifiable orthologs of known arginine kinases or

phosphatases. Nevertheless, the lack of consensus motif in the identified phosphosites predicts

a promiscuous kinase that targets cellular proteins, in a manner independent of a specific primary

sequence. Future work will be required to decipher how arginine phosphorylation is regulated,

and how it contributes to the physiology of mycobacteria and other actinobacterial species.

**SUPPLEMENTAL DATA**

This article contains supplemental data.

**ACKNOWLEDGEMENTS**

We thank members of the Schmitz lab for advice; VP, AJL, PCB, MP, AD, PAO and OU for

comments on the manuscript; and the Parashar lab for use of instrumentation. For technical help

with mass spectrometry, we thank Dr Yanbao Yu and the University of Delaware Mass

Spectrometry Facility. Sortase A was a kind gift from S Novo and JM Fox. The anti-pArg antibody

was a kind gift from Dr. Paul Thompson, UMass Medical School.

**Funding and Additional Information**

KRS was supported by NIH NIGMS Award P20GM104316. The UD Department of Chemistry &

Biochemistry Mass Spectrometry core was additionally supported by NIH NIGMS award

P30GM110758-02. This content is solely the responsibility of the authors and does not

necessarily represent the official views of the National Institutes of Health.

**Author Contributions**

E. O. and K. S. designed research; E. O. and H. A. performed research; E. O. and K. S. analyzed

data; E. O. and K. S. wrote the paper.

**References**

Akopian, T., Kandror, O., Raju, R. M., Unnikrishnan, M., Rubin, E. J., and Goldberg, A. L. (2012)

The active ClpP protease from M. tuberculosis is a complex composed of a heptameric

ClpP1 and a ClpP2 ring. *The EMBO journal* **31(6):** 1529–1541.

Akopian, T., Kandror, O., Tsu, C., Lai, J. H., Wu, W., Liu, Y., Zhao, P., Park, A., Wolf, L., Dick, L.

R., Rubin, E. J., Bachovchin, W., and Goldberg, A. L. (2015) Cleavage Specificity of

*Mycobacterium tuberculosis* ClpP1P2 Protease and Identification of Novel Peptide

Substrates and Boronate Inhibitors with Anti-bacterial Activity. *The Journal of biological*

*chemistry* **290(17):** 11008–11020.

Bairoch, A., and Apweiler, R. (1997) The SWISS-PROT protein sequence data bank and its

supplement TrEMBL. *Nucleic acids research* **25(1):** 31–36.

Baker, T. A., and Sauer, R. T. (2012) ClpXP, an ATP-powered unfolding and protein-degradation

machine. *Biochimica et biophysica acta* **1823(1):** 15–28.

Barik, S., Sureka, K., Mukherjee, P., Basu, J., and Kundu, M. (2010) RseA, the SigE specific anti-

sigma factor of *Mycobacterium tuberculosis*, is inactivated by phosphorylation-dependent

ClpC1P2 proteolysis. *Molecular microbiology* **75(3):** 592–606.

Bäsell, K., Otto, A., Junker, S., Zühlke, D., Rappen, G. M., Schmidt, S., Hentschker, C., Macek,
B., Ohlsen, K., Hecker, M., and Becher, D. (2014) The phosphoproteome and its
physiological dynamics in Staphylococcus aureus. *International journal of medical*
*microbiology: IJMM* **304(2):** 121–132.

Berman, H. M., Westbrook, J., Feng, Z., Gilliland, G., Bhat, T. N., Weissig, H., Shindyalov, I. N.,
& Bourne, P. E. (2000) The Protein Data Bank. *Nucleic acids research* **28(1):** 235–242.

Brokaw, A. M., Eide, B. J., Muradian, M., Boster, J. M., & Tischler, A. D. (2017) *Mycobacterium*
*smegmatis* PhoU Proteins Have Overlapping Functions in Phosphate Signaling and Are
Essential. *Frontiers in microbiology* **8:** 2523.

Brötz-Oesterhelt, H., Beyer, D., Kroll, H. P., Endermann, R., Ladel, C., Schroeder, W., Hinzen,
B., Raddatz, S., Paulsen, H., Henninger, K., Bandow, J. E., Sahl, H. G., and Labischinski,
H. (2005) Dysregulation of bacterial proteolytic machinery by a new class of
antibiotics. *Nature medicine* **11(10):** 1082–1087.

Bürstner, N., Roggo, S., Ostermann, N., Blank, J., Delmas, C., Freuler, F., Gerhartz, B., Hinniger,
A., Hoepfner, D., Liechty, B., Mihalic, M., Murphy, J., Pistorius, D., Rottmann, M., Thomas,
J. R., Schirle, M., and Schmitt, E. K. (2015) Gift from Nature: Cyclomarin A Kills
Mycobacteria and Malaria Parasites by Distinct Modes of Action. *Chembiochem : a*
*European journal of chemical biology* **16(17):** 2433–2436.

Canova, M. J., Veyron-Churlet, R., Zanella-Cleon, I., Cohen-Gonsaud, M., Cozzone, A. J., Becchi,
M., Kremer, L., and Molle, V. (2008) The *Mycobacterium tuberculosis* serine/threonine
kinase PknL phosphorylates Rv2175c: mass spectrometric profiling of the activation loop
phosphorylation sites and their role in the recruitment of Rv2175c. *Proteomics* **8(3):** 521–
533.

Chow, K., Ng, D., Stokes, R., and Johnson, P. (1994) Protein tyrosine phosphorylation in
*Mycobacterium tuberculosis*. *FEMS microbiology letters* **124(2):** 203–207.

Compton, C. L., Schmitz, K. R., Sauer, R. T., and Sello, J. K. (2013) Antibacterial activity of and resistance to small molecule inhibitors of the ClpP peptidase. *ACS chemical biology* **8(12):** 2669–2677.

Conesa, A., Götz, S., García-Gómez, J. M., Terol, J., Talón, M., and Robles, M. (2005) Blast2GO: a universal tool for annotation, visualization and analysis in functional genomics research. *Bioinformatics (Oxford, England)* **21(18):** 3674–3676.

Crooks, G. E., Hon, G., Chandonia, J. M., and Brenner, S. E. (2004) WebLogo: a sequence logo generator. *Genome research* **14(6):** 1188–1190.

DeJesus, M.A., Gerrick, E.R., Xu, W., Park, S.W., Long, J.E.  Boutte, C.C., Rubin, E.J., Schnappinger, D., Ehrt, S., Fortune, S.M., Sassetti, C.M., and Ioerger, T.R. (2017) Comprehensive Essentiality Analysis of the *Mycobacterium tuberculosis* Genome via Saturating Transposon Mutagenesis. *mBio* **8(1):** e02133-16.

Eddy S. R. (1995) Multiple alignment using hidden Markov models. *Proceedings. International Conference on Intelligent Systems for Molecular Biology* **3:** 114–120.

Elsholz, A. K., Turgay, K., Michalik, S., Hessling, B., Gronau, K., Oertel, D., Mäder, U., Bernhardt, J., Becher, D., Hecker, M., and Gerth, U. (2012) Global impact of protein arginine phosphorylation on the physiology of *Bacillus subtilis*. *Proceedings of the National Academy of Sciences of the United States of America* **109(19):** 7451–7456.

Eng, J. K., McCormack, A. L., and Yates, J. R. (1994) An approach to correlate tandem mass spectral data of peptides with amino acid sequences in a protein database. *Journal of the American Society for Mass Spectrometry* **5(11):** 976–989.

Famulla, K., Sass, P., Malik, I., Akopian, T., Kandror, O., Alber, M., Hinzen, B., Ruebsamen-Schaeff, H., Kalscheuer, R., Goldberg, A. L., and Brötz-Oesterhelt, H. (2016) Acyldepsipeptide antibiotics kill mycobacteria by preventing the physiological functions of the ClpP1P2 protease. *Molecular microbiology* **101(2)**: 194–209.

Fu, S., Fu, C., Zhou, Q., Lin, R., Han, O., Wang, M., Sun, Y., Liu, Y., and Zhao, Y. (2020) Widespread arginine phosphorylation in human cells—a novel protein PTM revealed by mass spectrometry. *Sci. China Chem.* **63:** 341–346.

Fuhrmann, J., Schmidt, A., Spiess, S., Lehner, A., Turgay, K., Mechtler, K., Charpentier, E., and Clausen, T. (2009) McsB is a protein arginine kinase that phosphorylates and inhibits the heat-shock regulator CtsR. *Science (New York, N.Y.)* **324(5932):** 1323–1327.

Fuhrmann, J., Subramanian, V., and Thompson, P. R. (2015) Synthesis and Use of a Phosphonate Amidine to Generate an Anti-Phosphoarginine-Specific Antibody. *Angewandte Chemie (International ed. in English)* **54(49):** 14715–14718.

Fuhrmann, J., Subramanian, V., Kojetin, D. J., and Thompson, P. R. (2016) Activity-Based Profiling Reveals a Regulatory Link between Oxidative Stress and Protein Arginine Phosphorylation. *Cell chemical biology* **23(8):** 967–977.

Gao, W., Kim, J. Y., Anderson, J. R., Akopian, T., Hong, S., Jin, Y. Y., Kandror, O., Kim, J. W., Lee, I. A., Lee, S. Y., McAlpine, J. B., Mulugeta, S., Sunoqrot, S., Wang, Y., Yang, S. H., Yoon, T. M., Goldberg, A. L., Pauli, G. F., Suh, J. W., Franzblau, S. G., and Cho, S. (2015) The cyclic peptide ecumicin targeting ClpC1 is active against *Mycobacterium tuberculosis* in vivo. *Antimicrobial agents and chemotherapy* **59(2):** 880–889.

Gavrish, E., Sit, C. S., Cao, S., Kandror, O., Spoering, A., Peoples, A., Ling, L., Fetterman, A., Hughes, D., Bissell, A., Torrey, H., Akopian, T., Mueller, A., Epstein, S., Goldberg, A., Clardy, J., and Lewis, K. (2014) Lassomycin, a ribosomally synthesized cyclic peptide, kills *Mycobacterium tuberculosis* by targeting the ATP-dependent protease ClpC1P1P2. *Chemistry & biology* **21(4):** 509–518.

Gerth, U., Kock, H., Kusters, I., Michalik, S., Switzer, R. L., & Hecker, M. (2008) Clp-dependent proteolysis down-regulates central metabolic pathways in glucose-starved Bacillus subtilis. *Journal of bacteriology* **190(1):** 321–331.

Glasgow, J. E., Salit, M. L., and Cochran, J. R. (2016) *In vivo* site-specific protein tagging with

diverse amines using an engineered sortase variant. *Journal of the American Chemical*

*Society* **138 (24):** 7496-7499

Gopal, P., Sarathy, J. P., Yee, M., Ragunathan, P., Shin, J., Bhushan, S., Zhu, J., Akopian, T.,

Kandror, O., Lim, T. K., Gengenbacher, M., Lin, Q., Rubin, E. J., Grüber, G., and Dick, T.

(2020) Pyrazinamide triggers degradation of its target aspartate decarboxylase. *Nature*

*communications* **11(1):** 1661.

Griffin, J.E., Gawronski, J.D., Dejesus, M.A., Ioerger ,T.R., Akerley, B.J., and Sassetti, C.M.

(2011) High-resolution phenotypic profiling defines genes essential for mycobacterial

growth and cholesterol catabolism. *PLoS Pathog* **7(9):** e1002251.

Hall, T.A. (1999) BioEdit: A User-Friendly Biological Sequence Alignment Editor and Analysis

Program for Windows 95/98/NT. *Nucleic Acids Symposium Series* **41:** 95-98.

Hunter, S., Apweiler, R., Attwood, T. K., Bairoch, A., Bateman, A., Binns, D., Bork, P., Das, U.,

Daugherty, L., Duquenne, L., Finn, R. D., Gough, J., Haft, D., Hulo, N., Kahn, D., Kelly,

E., Laugraud, A., Letunic, I., Lonsdale, D., Lopez, R., and Yeats, C. (2009) InterPro: the

integrative protein signature database. *Nucleic acids research* **37(Database issue):**

D211–D215.

Iakoucheva, L. M., Radivojac, P., Brown, C. J., O'Connor, T. R., Sikes, J. G., Obradovic, Z., &

Dunker, A. K. (2004) The importance of intrinsic disorder for protein

phosphorylation. *Nucleic acids research* **32(3):** 1037–1049.

Isaza, C. E., Silaghi-Dumitrescu, R., Iyer, R. B., Kurtz, D. M., Jr, & Chan, M. K. (2006) Structural

basis for $O_2$ sensing by the hemerythrin-like domain of a bacterial chemotaxis protein:

substrate tunnel and fluxional N terminus. *Biochemistry* **45(30):** 9023–9031.

Jerabek-Willemsen, M., André, T., Wanner, R., Roth, H.M., Duhr, S., Baaske, P., & Breitsprecher,

D. (2014) MicroScale Thermophoresis: Interaction analysis and beyond. *Journal of*

*Molecular Structure* **1077:** 101-113.

Jumper, J., Evans, R., Pritzel, A., Green, T., Figurnov, M., Ronneberger, O.,Tunyasuvunakool,

K., Bates, R., Žídek, A., Potapenko, A., Bridgland, A., Meyer, C., Kohl, S., Ballard, A. J.,

Cowie, A., Romera-Paredes, B., Nikolov, S., Jain, R., Adler, J., Back, T., Hassabis, D.

(2021) Highly accurate protein structure prediction with AlphaFold. *Nature* **596(7873):**

583–589.

Jung, I. P., Ha, N. R., Kim, A. R., Kim, S. H., and Yoon, M. Y. (2017) Mutation analysis of the

interactions between *Mycobacterium tuberculosis* caseinolytic protease C1 (ClpC1) and

ecumicin. *International journal of biological macromolecules* **101:** 348–357.

Junker, S., Maaβ, S., Otto, A., Michalik, S., Morgenroth, F., Gerth, U., Hecker, M., and Becher,

D. (2018) Spectral Library Based Analysis of Arginine Phosphorylations

in *Staphylococcus aureus*. *Molecular & cellular proteomics: MCP* **17(2):** 335–348.

Kirstein, J., Hoffmann, A., Lilie, H., Schmidt, R., Rübsamen-Waigmann, H., Brötz-Oesterhelt, H.,

Mogk, A., and Turgay, K. (2009) The antibiotic ADEP reprogrammes ClpP, switching it

from a regulated to an uncontrolled protease. *EMBO molecular medicine* **1(1):** 37–49.

Lavey, N. P., Coker, J. A., Ruben, E. A., and Duerfeldt, A. S. (2016) Sclerotiamide: The First Non-

Peptide-Based Natural Product Activator of Bacterial Caseinolytic Protease P. *Journal of*

*natural products* **79(4):** 1193–1197.

Leodolter, J., Warweg, J., and Weber-Ban, E. (2015) The *Mycobacterium tuberculosis* ClpP1P2

Protease Interacts Asymmetrically with Its ATPase Partners ClpX and ClpC1. *PloS*

*one* **10(5):** e0125345.

Li, L., MacIntyre, L. W., Ali, T., Russo, R., Koirala, B., Hernandez, Y., & Brady, S. F. (2021)

Biosynthetic Interrogation of Soil Metagenomes Reveals Metamarin, an Uncommon

Cyclomarin Congener with Activity against *Mycobacterium tuberculosis*. *Journal of natural*

*products* **84(4):** 1056–1066.

Li, M., Kandror, O., Akopian, T., Dharkar, P., Wlodawer, A., Maurizi, M. R., and Goldberg, A. L.

(2016) Structure and Functional Properties of the Active Form of the Proteolytic Complex,

ClpP1P2, from *Mycobacterium tuberculosis*. *The Journal of biological chemistry* **291(14):** 7465–7476.

Li, X., Li, J., Hu, X. *et al.* (2015) Differential roles of the hemerythrin-like proteins of *Mycobacterium smegmatis* in hydrogen peroxide and erythromycin susceptibility. *Sci Rep* **5:** 16130

Liu, Y., Gallo, A. A., Bajpai, R. K., Chistoserdov, A., Nelson, A., Segura, L., & Xu, W. (2010) The diversity and molecular modelling analysis of $B_{12}$ and $B_{12}$-independent glycerol dehydratases. *International journal of bioinformatics research and applications* **6(5):** 484–507.

Lunge, A., Gupta, R., Choudhary, E., and Agarwal, N. (2020) The unfoldase ClpC1 of *Mycobacterium tuberculosis* regulates the expression of a distinct subset of proteins having intrinsically disordered termini. *The Journal of biological chemistry* **295(28):** 9455–9473.

Ma, Z., Abendroth, J., Buchko, G. W., Rohde, K. H., & Davidson, V. L. (2020) Crystal structure of a hemerythrin-like protein from *Mycobacterium kansasii* and homology model of the orthologous Rv2633c protein of *M. tuberculosis*. *The Biochemical journal* **477(2**): 567–581.

Macek, B., Forchhammer, K., Hardouin, J., Weber-Ban, E., Grangeasse, C., and Mijakovic, I. (2019) Protein post-translational modifications in bacteria. *Nature reviews. Microbiology* **17(11):** 651–664.

Marsee, J. D., Ridings, A., Yu, T., and Miller, J. M. (2018) *Mycobacterium tuberculosis* ClpC1 N-Terminal Domain Is Dispensable for Adaptor Protein-Dependent Allosteric Regulation. *International journal of molecular sciences* **19(11):** 3651.

Moreira, W., Ngan, G. J., Low, J. L., Poulsen, A., Chia, B. C., Ang, M. J., Yap, A., Fulwood, J., Lakshmanan, U., Lim, J., Khoo, A. Y., Flotow, H., Hill, J., Raju, R. M., Rubin, E. J., and

639        Dick, T. (2015) Target mechanism-based whole-cell screening identifies bortezomib as an

640           inhibitor of caseinolytic protease in mycobacteria. *mBio* **6(3**): e00253-15.

641 Moreno-Cinos, C., Goossens, K., Salado, I. G., Van Der Veken, P., De Winter, H., and Augustyns,

642           K. (2019) ClpP Protease, a Promising Antimicrobial Target. *International journal of*

643           *molecular sciences* **20(9):** 2232.

644 Nagpal, J., Paxman, J. J., Zammit, J. E., Alhuwaider, A., Truscott, K. N., Heras, B., and Dougan,

645           D. A. (2019) Molecular and structural insights into an asymmetric proteolytic

646           complex (ClpP1P2) from *Mycobacterium smegmatis. Scientific reports* **9(1):** 18019.

647 Nakedi, K. C., Nel, A. J., Garnett, S., Blackburn, J. M., and Soares, N. C. (2015) Comparative

648           Ser/Thr/Tyr phosphoproteomics between two mycobacterial species: the fast growing

649           *Mycobacterium smegmatis* and the slow growing *Mycobacterium bovis* BCG. *Frontiers in*

650           *microbiology* **6:** 237.

651 Nasir, A., Ashok, S., Shim, J. Y., Park, S., & Yoo, T. H. (2020) Recent Progress in the

652           Understanding and Engineering of Coenzyme $B_{12}$-Dependent Glycerol

653           Dehydratase. *Frontiers in bioengineering and biotechnology* **8**: 500867.

654 Perez-Riverol, Y., Csordas, A., Bai, J., Bernal-Llinares, M., Hewapathirana, S., Kundu, D. J.,

655           Inuganti, A., Griss, J., Mayer, G., Eisenacher, M., Pérez, E., Uszkoreit, J., Pfeuffer, J.,

656           Sachsenberg, T., Yilmaz, S., Tiwary, S., Cox, J., Audain, E., Walzer, M., Jarnuczak, A. F.,

657           and Vizcaíno, J. A. (2019) The PRIDE database and related tools and resources in 2019:

658           improving support for quantification data. *Nucleic acids research* **47(D1):** D442–D450.

659 Potel, C. M., Lin, M. H., Heck, A., and Lemeer, S. (2018) Widespread bacterial protein histidine

660           phosphorylation revealed by mass spectrometry-based proteomics. *Nature*

661           *methods* **15(3):** 187–190.

662 Prisic, S., Dankwa, S., Schwartz, D., Chou, M. F., Locasale, J. W., Kang, C. M., Bemis, G.,

663           Church, G. M., Steen, H., and Husson, R. N. (2010) Extensive phosphorylation with

664           overlapping specificity by *Mycobacterium tuberculosis* serine/threonine protein

kinases. *Proceedings of the National Academy of Sciences of the United States of America* **107(16):** 7521–7526.

Raju, R. M., Unnikrishnan, M., Rubin, D. H., Krishnamoorthy, V., Kandror, O., Akopian, T. N., Goldberg, A. L., and Rubin, E. J. (2012) *Mycobacterium tuberculosis* ClpP1 and ClpP2 function together in protein degradation and are required for viability in vitro and during infection. *PLoS pathogens* **8(2):** e1002511.

Raju, R.M., Jedrychowski, M.P., Wei, J.R., Pinkham, J.T., Park, A.S., O'Brien, K., Rehren, G., Schnappinger, D., Gygi S.P., and Rubin, E.J. (2014) Post-translational regulation via Clp protease is critical for survival of *Mycobacterium tuberculosis*. *PLoS Pathog* **10(3):** e1003994.

Sassetti, C.M., Boyd, D.H., and Rubin E.J. (2003) Genes required for mycobacterial growth defined by high density mutagenesis. *Molecular microbiology* **48 (1):** 77-84.

Sauer, R. T., and Baker, T. A. (2011) AAA+ proteases: ATP-fueled machines of protein destruction. *Annual review of biochemistry* **80:** 587–612.

Schlothauer, T., Mogk, A., Dougan, D. A., Bukau, B., and Turgay, K. (2003) MecA, an adaptor protein necessary for ClpC chaperone activity. *Proceedings of the National Academy of Sciences of the United States of America* **100(5):** 2306–2311.

Schmidt, A., Trentini, D. B., Spiess, S., Fuhrmann, J., Ammerer, G., Mechtler, K., and Clausen, T. (2014) Quantitative phosphoproteomics reveals the role of protein arginine phosphorylation in the bacterial stress response. *Molecular & cellular proteomics: MCP* **13(2):** 537–550.

Schmitt, E. K., Riwanto, M., Sambandamurthy, V., Roggo, S., Miault, C., Zwingelstein, C., Krastel, P., Noble, C., Beer, D., Rao, S. P., Au, M., Niyomrattanakit, P., Lim, V., Zheng, J., Jeffery, D., Pethe, K., & Camacho, L. R. (2011) The natural product cyclomarin kills *Mycobacterium tuberculosis* by targeting the ClpC1 subunit of the caseinolytic protease. *Angewandte Chemie (International ed. in English)* **50(26)**: 5889–5891.

Schmitz, K. R., and Sauer, R. T. (2014) Substrate delivery by the AAA+ ClpX and ClpC1

unfoldases activates the mycobacterial ClpP1P2 peptidase. *Molecular*

*microbiology* **93(4):** 617–628.

Schmitz, K. R., Carney, D. W., Sello, J. K., and Sauer, R. T. (2014) Crystal structure of

*Mycobacterium tuberculosis* ClpP1P2 suggests a model for peptidase activation by AAA+

partner binding and substrate delivery. *Proceedings of the National Academy of Sciences*

*of the United States of America* **111(43):** E4587–E4595.

Sievers, F., Wilm, A., Dineen, D., Gibson, T. J., Karplus, K., Li, W., Lopez, R., McWilliam, H.,

Remmert, M., Söding, J., Thompson, J. D., and Higgins, D. G. (2011) Fast, scalable

generation of high-quality protein multiple sequence alignments using Clustal

Omega. *Molecular systems biology* **7:** 539.

Singh, G. P. (2015) Association between intrinsic disorder and serine/threonine phosphorylation

in *Mycobacterium tuberculosis*. *PeerJ* **3:** e724.

States, D. J., & Gish, W. (1994) Combined use of sequence similarity and codon bias for coding

region identification. *Journal of computational biology: a journal of computational*

*molecular cell biology* **1(1):** 39–50.

Suskiewicz, M. J., Hajdusits, B., Beveridge, R., Heuck, A., Vu, L. D., Kurzbauer, R., Hauer, K.,

Thoeny, V., Rumpel, K., Mechtler, K., Meinhart, A., and Clausen, T. (2019) Structure of

McsB, a protein kinase for regulated arginine phosphorylation. *Nature chemical*

*biology* **15(5):** 510–518.

Taus, T., Köcher, T., Pichler, P., Paschke, C., Schmidt, A., Henrich, C., and Mechtler, K. (2011)

Universal and confident phosphorylation site localization using phosphoRS. *Journal of*

*proteome research* **10(12):** 5354–5362.

Thingholm, T. E., Jørgensen, T. J., Jensen, O. N., and Larsen, M. R. (2006) Highly selective

enrichment of phosphorylated peptides using titanium dioxide. *Nature protocols* **1(4):**

1929–1935.

Trentini, D. B., Fuhrmann, J., Mechtler, K., & Clausen, T. (2014) Chasing Phosphoarginine Proteins: Development of a Selective Enrichment Method Using a Phosphatase Trap. *Molecular & cellular proteomics: MCP* **13(8):** 1953–1964.

Trentini, D. B., Suskiewicz, M. J., Heuck, A., Kurzbauer, R., Deszcz, L., Mechtler, K., and Clausen, T. (2016) Arginine phosphorylation marks proteins for degradation by a Clp protease. *Nature* **539(7627):** 48–53.

Vahidi, S., Ripstein, Z. A., Juravsky, J. B., Rennella, E., Goldberg, A. L., Mittermaier, A. K., Rubinstein, J. L., and Kay, L. E. (2020) An allosteric switch regulates *Mycobacterium tuberculosis* ClpP1P2 protease function as established by cryo-EM and methyl-TROSY NMR. *Proceedings of the National Academy of Sciences of the United States of America* **117(11):** 5895–5906.

Vasudevan, D., Rao, S. P., and Noble, C. G. (2013) Structural basis of mycobacterial inhibition by cyclomarin A. *The Journal of biological chemistry* **288(43):** 30883–30891.

Wagner, T., Bellinzoni, M., Wehenkel, A., O'Hare, H. M., & Alzari, P. M. (2011) Functional plasticity and allosteric regulation of α-ketoglutarate decarboxylase in central mycobacterial metabolism. *Chemistry & biology* **18(8):** 1011–1020.

Waterhouse, A. M., Procter, J. B., Martin, D. M., Clamp, M., and Barton, G. J. (2009) Jalview Version 2--a multiple sequence alignment editor and analysis workbench. *Bioinformatics (Oxford, England)* **25(9):** 1189–1191.

Weinhäupl, K., Brennich, M., Kazmaier, U., Lelievre, J., Ballell, L., Goldberg, A., Schanda, P., and Fraga, H. (2018) The antibiotic cyclomarin blocks arginine-phosphate-induced millisecond dynamics in the N-terminal domain of ClpC1 from *Mycobacterium tuberculosis*. *The Journal of biological chemistry* **293(22):** 8379–8393.

Wiśniewski, J. R., Zougman, A., Nagaraj, N., and Mann, M. (2009) Universal sample preparation method for proteome analysis. *Nature methods* **6(5):** 359–362.

Wolf, N. M., Lee, H., Choules, M. P., Pauli, G. F., Phansalkar, R., Anderson, J. R., Gao, W., Ren,
J., Santarsiero, B. D., Lee, H., Cheng, J., Jin, Y. Y., Ho, N. A., Duc, N. M., Suh, J. W.,
Abad-Zapatero, C., and Cho, S. (2019) High-Resolution Structure of ClpC1-Rufomycin
and Ligand Binding Studies Provide a Framework to Design and Optimize Anti-
Tuberculosis Leads. *ACS infectious diseases* **5(6):** 829–840.

Wolf, N. M., Lee, H., Zagal, D., Nam, J. W., Oh, D. C., Lee, H., Suh, J. W., Pauli, G. F., Cho, S.,
and Abad-Zapatero, C. (2020) Structure of the N-terminal domain of ClpC1 in complex
with the antituberculosis natural product ecumicin reveals unique binding
interactions. *Acta crystallographica. Section D, Structural biology* **76(Pt 5):** 458–471.

World Health Organization (2021) *Global Tuberculosis Report* (World Health Organization,
Geneva)

Zhou, B., Semanjski, M., Orlovetskie, N., Bhattacharya, S., Alon, S., Argaman, L., Jarrous, N.,
Zhang, Y., Macek, B., Sinai, L., and Ben-Yehuda, S. (2019) Arginine dephosphorylation
propels spore germination in bacteria. *Proceedings of the National Academy of Sciences
of the United States of America* **116(28):** 14228–14237.

Ziemski, M., Leodolter, J., Taylor, G., Kerschenmeyer, A., & Weber-Ban, E. (2021) Genome-wide
interaction screen for *Mycobacterium tuberculosis* ClpCP protease reveals toxin-antitoxin
systems as a major substrate class. *The FEBS journal*, **288(1):** 111–126

**Figure 1**. **Indirect evidence for phosphoarginine in actinobacteria**. **A**) The domain organization of ClpC1 is shown above a plot of amino acid conservation among actinobacterial ClpC1 orthologs, where conservation at each alignment position is plotted as a purple vertical strip. The apparent poor conservation at the beginning of the NTD (marked by *) likely reflects mis-annotation of the start site in some entries. **B**) The sequence of the *Bacillus subtilis* ClpC NTD is shown above equivalent regions from several actinobacterial ClpC orthologs. The sequence logo below shows amino acid conservation in actinobacterial ClpC1 orthologs. Arrows mark positions reported to be important for phosphoarginine binding to site 1 (orange) or site 2 (yellow) (Weinhäupl et al., 2018). Phosphoarginines from the crystal structure of *B. subtilis* ClpC NTD (5HBN) were modeled on the *M. tuberculosis* ClpC1 NTD (6PBQ) putative phosphoarginine-binding sites 1 (**C**) and 2 (**D**) (Wolf et al., 2020). **E**) Binding of phosphoarginine to *Msm* ClpC1$^{NTD}$ was measured by microscale thermophoresis. Data were fit to a noncooperative binding model (gray curve), yielding a $K_D$ of 1.4 ± 0.6 µM. Values are averages of three technical replicates (N = 3) ± one SD. **F**) Dot blots of *Mycolicibacterium smegmatis*, *Corynebacterium jeikeium*, *Bacillus subtilis*, and *Escherichia coli* cell lysates, probed with anti-phosphoarginine antibody (Fuhrmann et al., 2015) or stained with Ponceau S.

**Figure 2. Comparative analysis of mass spectrometry output. A**) Workflow for arginine

phosphoproteomics in *Mycolicibacterium smegmatis*. **B**) Number of proteins obtained across biological

replicates from cells grown in heat stress (50°C) and normal conditions (37°C). **C**) Volcano plot showing

enrichment of proteins observed by LC-MS/MS in heat stress versus normal growth conditions. Plotted on

the *x*-axis is the $\log_2$ of the ratio of average Sequest HT score of stressed to unstressed samples. The *y*-

axis shows $-\log_{10}$ of p-values obtained by Student's t-test. The horizontal green line indicates a cut-off p-

value = 0.05; the vertical blue line indicates score ratio = 1. Highlighted in red are notable chaperone

proteins with stressed:unstressed ratio > 1.

| Protein | Accession | Gene | Position | Observations | | | | | |
|---|---|---|---|---|---|---|---|---|---|
| | | | | Unstressed | | | Stressed | | |
| | | | | 1 | 2 | 3 | 1 | 2 | 3 |
| Uncharacterized protein | A0QQU4 | MSMEG_0879 | R31 | • | • | • | • | • | • |
| Xanthine/uracil permeases family protein | A0QRZ5 | MSMEG_1293 | R475 | | • | | | | |
| Phosphate-specific transport system accessory protein PhoU1 | A0R4B7 | phoU | R149 | | • | • | • | • | • |
| Multifunctional 2-oxoglutarate metabolism enzyme | A0R2B1 | kgd | R433 | | • | | | | |
| Hemerythrin domain-containing protein | A0QQX2 | MSMEG_0907 | R34 | • | • | • | | • | • |
| Glycerol dehydratase large subunit | A0QSP1 | MSMEG_1547 | R326 | • | • | | • | • | • |

**Figure 3. Identification of arginine phosphosites in *Msm* proteome**. (**A**) Localized arginine phosphosites in six *Msm* proteins. Localization probability is reported as PhosphoRS/ptmRS score. Representative secondary fragmentation spectra show arginine-phosphorylated peptides from (**B**) Phou1 and (**C**) hemerythrin domain-containing protein. Additional spectra are shown in **Supplemental Figure S2.**

**Figure 4. Structural characteristics of phosphoarginine sites**. (**A**) Sequence alignment of 20 residues

flanking the arginine phosphosites. Structural models of (**B**) Msmeg_1293, (**C**) Msmeg_1547, (**D**) PhoU1,

(**E**) Kgd and (**F**) Msmeg_0907 are shown with the phosphorylated arginine colored magenta and marked

with an arrow. All models were generated by AlphaFold2 (Jumper *et al.*, 2021).

**Figure 5. Sequence conservation of phosphorylated arginine residues.** Sequence logos show

conservation near phosphorylated arginines in (A) Kgd R433, (B) Msmeg_0907 R34, (C) Msmeg_0879

R31, (D) Msmeg_1547 R326, (E) PhoU1 R149, (F) Msmeg_1293 R475. Percentages indicates positional

conservation of the phosphorylated arginine; "n" indicates the number of aligned orthologous sequences

used to generate the sequence logo.

**Supplemental Figure 1. Gene ontology (GO) annotation of proteins strongly enriched or depleted**

**upon heat shock.** Proteins with significantly (p ≤ 0.05) and strongly different levels (two-fold or more) in

heat shock and control conditions were sorted into GO annotation categories based on biological process.

**Supplemental Figure 2. Phosphosite secondary spectra.** Secondary fragmentation spectra showing

arginine-phosphorylated peptides in (**A**) 2-oxoglutarate, (**B**) glycerol dehydratase, (**C**) xanthine uracyl

permease, and (**D**) MSMEG_0879.

**Supplemental Figure 3. Solvent accessibility of phosphorylated arginines.** Surface representations of

proteins (**A**) MSMEG_1293 (**B**) PhoU1 (**C**) Kgd (**D**) MSMEG_0907 (**E**) MSMEG_1547 based on AlphaFold2

prediction. Magenta patches represent specific arginine residues observed to be arginine-phosphorylated

located on protein surface.

821