1   Differential effects of cerebellar degeneration on feedforward versus feedback control across
2   speech and reaching movements
3

4   Benjamin Parrell[1*], Hyosub Kim[2,3], Assaf Breska[2], Arohi Saxena[2], & Rich Ivry[2]
5
6   [1] University of Wisconsin–Madison, Department of Communication Sciences & Disorders,
7   Madison, WI, 53706, USA
8   [2] University of California, Berkeley, Department of Psychology, Berkeley, CD, 94720
9   [3] University of Delaware, Department of Physical Therapy, Newark, DE, 19713
10   [*] corresponding author, bparrell@wisc.edu

11
Number of pages: 24
Number of figures: 6
Number of tables: 0
Number of multimedia items: 0
Number of 3D models: 0
Number of words for abstract: 247
Number of words for significance: 116
Number of words for introduction: 647
Number of words for discussion: 1299
21
22   Conflict of interests: None of the authors has a conflict of interest to disclose.
23

24   Acknowledgments: This work was funded by grant R01 DC017091 from the NIDCD and R35
25   NS116883 from the NINDS.

Abstract

Errors that result from a mismatch between predicted movement outcomes and sensory afference are used to correct ongoing movements through feedback control and to adapt feedforward control of future movements. The cerebellum has been identified as a critical part of the neural circuit underlying implicit adaptation across a wide variety of movements (reaching, gait, eye movements, and speech). The contribution of this structure to feedback control is less well understood: although it has recently been shown in the speech domain that individuals with cerebellar degeneration produce even larger online corrections for sensory perturbations than control participants, similar behavior has not been observed in other motor domains. Currently, comparisons across domains are limited by different population samples and potential ceiling effects in existing tasks. To assess the relationship between changes in feedforward and feedback control associated with cerebellar degeneration across motor domains, we evaluated adaptive (feedforward) and compensatory (feedback) responses to sensory perturbations in reaching and speech production in individuals with cerebellar degeneration and neurobiologically healthy controls. As expected, the cerebellar group demonstrated impaired adaptation in both reaching and speech. In contrast, the groups did not differ in their compensatory response in either domain. Moreover, compensatory and adaptive responses in the cerebellar group were not correlated within or across motor domains. Together, these results point to a general impairment in feedforward control with spared feedback control in cerebellar degeneration. However, the magnitude of feedforward impairments and potential changes in feedback-based control manifest in a domain-specific manner across individuals.

Significance Statement
The cerebellum contributes to feedforward updating of movement in response to sensory
errors, but its role in feedback control is less understood. Here, we tested individuals with
cerebellar degeneration (CD), using sensory perturbations to assess adaptation of feedforward
control and feedback gains during reaching and speech production tasks. The results confirmed
that CD leads to reduced adaption in both domains. However, feedback gains were unaffected
by CD in either domain. Interestingly, measures of feedforward and feedback control were not
correlated across individuals within or across motor domains. Together, these results indicate a
general impairment in feedforward control with spared feedback control in CD. However, the
magnitude of feedforward impairments manifests in a domain-specific manner across
individuals.
62

Introduction

Coordinated movement relies on a combination of feedback control and anticipatory mechanisms. A mismatch between the predicted and actual feedback resulting from a motor command can lead to online corrections as well adaption of feedforward control for future movements (Shadmehr and Krakauer, 2008). The cerebellum plays a critical role in this latter process, helping ensure that the predictive system is optimally calibrated. One line of supportive evidence comes from the substantial literature showing markedly impaired performance of individuals with cerebellar degeneration (CD) during sensorimotor adaptation tasks involving upper limb movement (Martin et al., 1996; Smith and Shadmehr, 2005; Tseng et al., 2007; Donchin et al., 2012; Schlerf et al., 2013), gait (Morton and Bastian, 2006), eye movements (Xu-Wilson et al., 2009), and speech (Parrell et al., 2017).

If, and how, the cerebellum contributes to feedback control is less clear. One clue comes from intentional tremor, a prominent feature of CD where low-frequency oscillations occur around the movement endpoint. Intentional tremor is reduced when movement is produced without visual feedback (Day et al., 1998). Such behavior is broadly consistent with a control system that relies on visual feedback (Beppu et al., 1987), and could occur if the gains on sensory feedback errors are larger in individuals with CD. Larger gains could lead to overcorrections for errors and the need for additional counter-corrections. This hypothesis is supported by evidence showing that, relative to controls, individuals with CD produce larger compensatory responses to auditory perturbations of speech (Parrell et al., 2017; Houde et al., 2019; Li et al., 2019).

There is mixed evidence concerning feedback gains in other types of movement. Individuals with CD produce a smaller long-latency muscle response to mechanical perturbations (Kurtzer et al., 2013), consistent with a decreased gain in the response to proprioceptive feedback. However, their feedback-based corrections during split-belt treadmill walking are relatively normal (Morton and Bastian, 2006), suggesting proprioceptive gains are unaffected for at least some tasks. Moreover, individuals with CD have normal feedback gains in a continuous visual tracking task, but with a substantial phase lag (Zimmet et al., 2020). Computationally, this is consistent with an increased reliance on (delayed) feedback in the absence of a predictive function for state estimation (Wolpert et al., 1998; Miall et al., 2007).

It is possible that potential increases in feedback gains in non-speech tasks may be obscured by a ceiling effect. While the auditory feedback response in speech only partially compensates for the perturbation, the responses to perturbations in tasks such as visually-guided movements or walking typically provide nearly complete compensation to the perturbation (Morton and Bastian, 2006; Tseng et al., 2007). As such, reductions in feedback gains could be readily measured but increases in gain might be hard to detect. Alternatively (or additionally), modification of feedback gains in CD may be task-dependent, with variable changes across movement types. This would stand in contrast to feedforward control, where the cerebellum appears to play a similar role in implicit adaptation across domains.

The preceding hypotheses are based on inferences drawn from disparate studies that distinct methods and typically focus on a single motor domain. In the present study, we used a 2 x 2 design to evaluate feedback and feedforward control in two motor tasks, one involving visually-guided reaching and the other speech production. To avoid potential ceiling effects in the former, we used a task shown previously to induce only partial corrections to the visual

perturbation (Körding and Wolpert, 2004). Moreover, by testing the same individuals on all four
tasks, we can perform a correlational analysis, focusing on two key questions related to
individual differences associated with CD. First, are patterns of impairment similar across the
two motor domains and/or forms of control? Second, is the degree of impairment in
feedforward control (adaptation) predictive of feedback gains, a signature that would be
consistent with the hypothesis that enhanced feedback gains arise as a compensatory
mechanism.

Materials and Methods
<u>Participants</u>
23 individuals with cerebellar degeneration (CD: 19 female, 37-89 years, mean age 62 years)
and 15 age-matched neurobiologically healthy controls (CO: 8 female, 43-85 years, mean age 61
years) were recruited for the study. The participants had normal or corrected-to-normal vision,
and none reported any history of speech or hearing impairments, or significant neurological
issues (other than ataxia in the CD group). Participants provided informed consent and received
financial compensation for their time. The protocols were approved by the institutional review
boards at the University of Wisconsin–Madison and the University of California, Berkeley.
All of the participants were administered the Montreal Cognitive Assessment (MoCA) as
a gross measure of cognitive impairment. The CD group was additionally administered the
Standard Ataxia Rating Scale (SARA) to assess the severity of their ataxia. SARA subscores were
calculated for both upper limb control and speech. One individual with CD was excluded due a
MOCA score < 18, indicative of moderate cognitive impairment. See the Appendix for a full
characterization of the CD group. Data from one control participant was excluded from Exp 4
due to equipment failure during data collection. The final sample size was based on typical
sample sizes in previous studies and, with sample sizes of 22 and 15 for the CD and control
groups, gives a power of 0.75 to detect a large between group effect (d = 0.8) as well as a
power of 0.75 detect a medium effect (d = 0.5) for correlations within the CD group.
133
<u>Experimental Design</u>
Each participant completed four conditions in a single experimental session that lasted
approximately one hour, including breaks of approximately 5 min between each condition.
Feedforward and feedback control during reaching were assessed in Conditions 1 and 2,
respectively; similarly, feedforward and feedback control during speech were assessed in
Conditions 3 and 4, respectively. A schematic of each of the four conditions is shown in **Figure**
2. The same order of the four conditions was used for each participant. While this approach
introduces order confounds, we opted to keep the order fixed given that a) this is preferable for
correlational analyses and b) the sample size was insufficient to properly assess order effects.
Stimulus presentation and data collection was controlled with Matlab for all conditions.
144
145

FEEDFORWARD CONTROL   *FEEDBACK CONTROL*

Figure 1: Experimental design. *The top row of each panel shows the structure of the experimental*
session and the bottom part depicts the perturbation. **A:** *Condition 1, feedforward control in reaching:*
Participants made 8 cm center-out reaches to targets (blue dot) with endpoint feedback. During the
perturbation phase, the location of the feedback cursor was rotated by 20° from actual hand position
(direction cross-balanced across participants). The dashed line represents the unseen hand movement. **B:**
Condition 2, feedback control in reaching: Participants made 16 cm reaches to a target (blue dot). During
the perturbation phase, visual feedback about hand position (black dot) was given at reach midpoint
with the feedback shifted by -1, 0, or 1 cm, with the shift on each trial determined in a pseudo-random
manner. The thin gray line depicts location of reach midpoint, but was not visible to the participant. **C:**
Condition 3, feedforward control in speech: Participants spoke a single word on each trial, hearing a
playback of their speech over headphones with minimal delay. During the perturbation phase, feedback
of the first formant during the vocalic portion of the utterance was perturbed by imposition of a − 125
mel shift (cyan trace superimposed on the spectrogram). **D:** *Condition 4, feedback control in speech:*
Participants spoke a single word on each trial. During the perturbation phase, auditory perturbations
(cyan trace) of -125, 0, or 125 mels were pseudo-randomly applied to the first formant during the vocalic
portion of the utterance.

Condition 1: Feedforward control in reaching.
Participants were seated in front of a 53.2 x 20 cm LCD screen (ASUS) that was horizontally
encased in a table frame mounted 27 cm above a 49.3 x 32.7 cm digitizing tablet (Intuos 4XL,
Wacom, Vancouver, WA). Participants held a modified air hockey paddle and made reaching
movements by sliding the paddle across the table. The position of a stylus embedded in the
paddle was recorded by the tablet at 200 Hz. Feedback, when available, was presented in the
form of a cursor on the LCD screen. Participants' view of their hand was blocked by the LCD

screen. To further limit vision of the upper arm, the experiment was conducted in a darkened
room. The experiment was controlled with Psychtoolbox (Brainard, 1997; Pelli, 1997; Kleiner et
al., 2007).
Participants made center-out reaches, moving to targets located at a radial distance of 8
cm from the center of the workspace. The start location was marked by a white ring (6 mm
diameter). On each trial, the participant moved the digitizing stylus to position the hand within
the start location. Visual feedback about the stylus position was provided by a small (3.5 mm
diameter) white circle. After the participant had maintained the stylus position within the start
location for 500 ms, one of 3 equally-spaced targets (6 mm diameter) appeared (0°, 120°, or
240°). The participant was instructed to reach, attempting to "slice through" the target. The
instructions emphasized that the movements should be made quickly and, to minimize
demands on endpoint (radial) accuracy, should terminate beyond the target location. RT was
not emphasized; the movement could be initiated at any time after the presentation of the
target.
Visual feedback about the movement was limited to endpoint feedback, eliminating the
opportunity for visually guided corrections. The feedback cursor disappeared at movement
onset and reappeared when the radial distance of the hand movement reached 8 cm. The
cursor remained visible for 50 ms and then was blanked. If participants reached the 8 cm
threshold in ≤ 500 ms, a knocking sound was played, indicating that the movement was "fast
enough and far enough." If the movement duration was > 500 ms, participants heard a
recording of the words "too slow". During the return movement to the start location, visual
feedback was withheld until the stylus was within 3 cm of the start position.
The condition consisted of five phases. 1) A *no feedback baseline* phase of 30 trials to
measure movement variability in the absence of visual feedback. 2) A *baseline phase* of 30 trials
with veridical endpoint feedback of the cursor. 3) A *perturbation phase* of 90 trials in which the
location of the cursor at movement endpoint was rotated 20° from the true position. The 20°
rotation was chosen to limit awareness of the perturbation and minimize the use of explicit re-
aiming (Bond and Taylor, 2015; Morehead et al., 2015; Werner et al., 2015). The perturbation
was either clockwise or counter-clockwise, counterbalanced across participants. 4) An
aftereffect phase of 15 trials in which there was no visual feedback. 5) A *washout* phase of 15
trials with veridical endpoint feedback.
The hand angle for each trial was measured as the angle between a line connecting the
start location and the hand position at the time of peak radial velocity, relative to the line
connecting the start and target locations. To remove intrinsic biases in reaching, the mean hand
angle during the baseline phase was subtracted from the heading angle for each trial. We focus
on two measures of adaptation, the mean of the last 15 trials of the perturbation phase
(*asymptote*) and the mean of the 15 trials in the aftereffect phase. We additionally measured
reaction time (time from target appearance to movement onset) and movement time (time
from movement onset to target), as well as the percentage of trials where any part of the
cursor representing hand position overlapped with the target (*hits*).
211
Condition 2: Feedback control in reaching.
The experimental apparatus was identical to Exp 1. On each trial, the white ring
indicating the start location appeared near the bottom edge of the screen at the horizontal

meridian. After this positioned was maintained for 500 ms, the target appeared at a fixed
position, 16 cm in front of the start location. The longer distance was used so that feedback
could be presented at the midpoint of the movement, providing sufficient distance (and time)
for an online correction.
There were five phases to Condition 2. 1) An *acclimation phase* of 20 trials with
continuous veridical feedback. 2) A *baseline phase* of 30 trials in which there was no visual
feedback, either during the movement or at the target distance. 3) A *practice* phase of 20 trials
to introduce the method for providing limited visual feedback. The cursor appeared twice on
these trials, first when the radial amplitude of the stylus was 8 cm from the start location
(midpoint, duration = 100 ms), and second when the radial amplitude reach 16 cm (endpoint,
duration = 50 ms). Participants were instructed to "use what you see midway through the
reach to get as close as possible to the target". 4) A *perturbation phase* of 120 trials with
midpoint feedback. On 50% of the trials, the midpoint feedback was given at the true location
of the stylus, and on the remaining 50% of the trials, the midpoint feedback was shifted 1 cm to
the left of the stylus position or 1 cm to the right (25% each). Each cluster of 4 reaches
consisted of 2 unperturbed trials, and 1 trial with the leftward and rightward perturbation, in a
randomized order. Endpoint feedback was not provided on perturbed trials to minimize
learning effects (i.e., anticipatory effects based on prior responses to the perturbation).
Endpoint feedback was provided on the unperturbed trials to help participants remain
calibrated to the target reaching location. 5) A *final phase* of 20 trials with no visual feedback.
Data from this last phase was not analyzed further.
As in Condition 1, participants were instructed to slice through the target. The
movement time criterion was increased to 1200 ms. The extra time was employed to
compensate for the larger amplitude movements and to encourage participants to adopt a
movement speed that allowed them time to use the midpoint feedback to make an online
correction (if warranted). The knocking sound was played if the target amplitude was reached
within 1200 ms, and if the movement duration was between 400 and 1200 ms, the target circle
turned green. If the movement was < 400 ms, the words "too fast" appeared on the screen. If
the movement duration was longer than 1200 ms, the target circle turned red and a recording
of the words "too slow" was played.
The focus here was on the online corrections made in response to the horizontally
displaced midpoint feedback; that is, the horizontal position of the hand relative to the target
at the target distance. However, the raw horizontal displacement would also reflect the
horizontal position of the hand at reach midpoint, which may vary between trials. To account
for this, we fit the trial-by-trial data of each participant with a linear model that predicted the
final horizontal position of the hand on each trial from the horizontal hand position at reach
midpoint, the visual perturbation (treated as a categorical variable and coded using separate
dummy predictors for each direction), and the interaction between midpoint hand position and
the perturbation:

$x_{target} \sim \beta_0 + \beta_1 x_{midpoint} + \beta_2 perturbation(L) + \beta_3 perturbation(R) + \beta_4 x_{midpoint}:perturbation(L) +$
$\beta_5 x_{midpoint}:perturbation(R)$

This model allows us to estimate two dependent measures of feedback-based corrections: 1)
the magnitude of the correction for the perturbation (direction effect estimates, β_2, β_3) and 2)
corrections for self-produced variability (the midpoint position estimate, β_1, and its adjustment
in the presence of perturbations, β_4, β_5). For statistical analysis, the sign of β_2 values were
flipped such that negative values always reflected a compensatory response to the
perturbation, regardless of perturbation direction. Additionally, this method allows us to
estimate the effect of the correction for the visual perturbation while accounting for any
individual differences in the bias of the overall reach trajectory, an effect captured by the
intercept term (β_0), which reflects the horizontal displacement relative to the target position for
unperturbed reaches. We measured movement time, reaction time, and proportion of trials
where the cursor hit the target as for Condition 1. Hit proportion was calculated only for
unperturbed trials.
270
Condition 3: Feedforward control in speech.
Participants were again seated in front of the horizontally-aligned monitor. Participants
wore a head-mounted microphone (AKG C520) and close-backed, over-the-ear headphones
(Beyerdynamic DT 770). On each trial, participants spoke the word "head" after it appeared on
the monitor. The utterance was digitized through a Scarlett 2i2 sound card and processed with
Audapter (Cai et al., 2008; Tourville et al., 2013) to synthesize a sound for playback over the
headphones. The recording, processing, and playback occurred in near real time (~20 ms delay).
The intensity of the synthesized feedback was set to roughly 80 dB based on each participants
comfortable speaking volume, mixed with speech-shaped noise at ~60 dB to mask air- or bone-
conducted direct feedback of the utterance. The actual intensity level of the feedback varied
with changes in the intensity of participants' speech.
There were four phases in Condition 3. For each phase, the word "head" remained
visible for 2.5 s on each trial, and the trials were separated by 1.1-1.3 s (randomly jittered). 1) A
baseline phase of 60 trials with speech feedback resynthesized through Audapter with no
auditory perturbation applied. 2) A *perturbation phase* of 100 trials during which a constant
shift of -125 mels was applied to the first vowel formant (F1) throughout the utterance, shifting
the F1 value of "head" towards that of "hid". 3) A *washout phase* of 30 trials in which there was
no auditory perturbation.
Vowel onset and offset was labelled for each trial using a semi-automated procedure.
First, automatic labels were generated by identifying on the waveforms where the speech
amplitude first crossed (onset) or fell below (offset) a participant-specific amplitude value. The
automatic labels were then visually inspected and corrected when the waveform and
spectrogram indicated that the automatic markings were inappropriate. A semi-automated
procedure was used to track the formants during the vocalic phase of the utterance using
participant-specific vales for LPC order and pre-emphasis. Formant tracking was performed
with Praat (Boersma and Weenink, 2019). Using the Wave Runner software package (Niziolek
and Houde, 2015), the tracks were visually inspected and, when they did not align with visible
formants on the speech spectrogram, the formant tracking parameters (pre-emphasis, LPC
order) were modified.
The primary outcome measure was adaptation of the vocalic portion of the utterances
across trials. For each trial, we calculated the average F1 value from 50-100 ms after vowel

onset. This window avoids the transitional phase of the formant during the word-initial
consonant (initial 50 ms) as well as any online compensatory response to the perturbation,
which typically begins later than 100 ms after vowel onset (Tourville et al., 2008; Cai et al.,
2012; Parrell et al., 2017). To facilitate comparisons across participants, the F1 values were
normalized with respect to the average F1 value taken over the 50-100 ms window during the
second half of the baseline phase (trials 31-60). We chose to use the second half of the baseline
phase to allow participants time to acclimate to the processed auditory feedback.
309
Condition 4: Feedback control in speech.
The stimulus set consisted of four words, "dead", "fed", "said", and "shed", selected
because they share the same vowel /ɛ/. The condition began with a calibration phase designed
to shape the participant's speaking rate such that the produced vowel duration would be
between 300 and 500 ms. This criterion was important to ensure that there would be sufficient
time for feedback-based corrections, shown in previous work to have a latency of ~150 ms
(Tourville et al., 2008; Cai et al., 2012; Parrell et al., 2017). On each trial, one of the four words
was displayed. After each utterance, the automated estimate of the vowel duration in
milliseconds was displayed on the monitor. This procedure was repeated for 10 trials. If the
duration fell outside the 300-500 ms window on more than 2 of these 10 trials, the procedure
was repeated for 10 more trials. When the utterances fell within the criterion window for at
least 8 of the 10 trials, the main experiment began.
Condition 4 had two phases. 1) A *practice phase* of 10 trials with veridical auditory
feedback. 2) A *perturbation phase* of 120 trials. During the perturbation phase, participants
heard veridical feedback on 50% of trials, a + 125 mel shift of F1 (moving the vowel towards
that in "had") on 25% of trials, and a -125 mel shift of F1 (moving the vowel towards that in
"hid") on the remaining 25% of trials. Each group of 4 trials consisted of 2 unperturbed trials,
and 1 trial each of positive and negative F1 perturbations, randomly ordered. Stimuli were
presented in a random order (selection with replacement), with the constraint that the total
number of each stimulus word be as equal as possible across the experimental condition.
Vowels were tracked as in Exp 3. The primary outcome measure was the online
correction to the perturbation, calculated following standard approaches (Cai et al., 2012;
Niziolek and Guenther, 2013; Parrell et al., 2017; Daliri et al., 2020). First, an average baseline
F1 trajectory was calculated for each stimulus word from the unperturbed productions of that
word. Second, the F1 trajectory from each perturbed trial was normalized by subtracting the
appropriate word-specific average baseline F1 trajectory, giving a normalized F1 response for
each perturbed trial. Trajectories from trials with upward and downward F1 perturbations were
separately averaged to generate an average F1 response in each direction. To generate a
composite feedback response, the sign of the average response to the upward perturbation
was flipped, such that positive values always reflected a compensatory response to the
perturbation, regardless of perturbation direction. Note that because of the normalization
process (and lack of an explicit target), this differs slightly from the approach in Exp 2, where
midpoint variability had to be accounted for. To quantify the magnitude of the compensatory
response, the mean value of each average F1 response trajectory between 200-300 ms after
vowel onset was calculated. This window begins well after the expected 150 ms latency of the
compensatory response is ~150 ms (Tourville et al., 2008; Cai et al., 2012; Parrell et al., 2017).

Statistical Analysis

Statistical analysis for all conditions was conducted in R (R Core Team, 2013) using
mixed ANOVAs or Welches' two-sample t-tests. When necessary, post-hoc tests were
conducted with a Tukey HSD correction. Where appropriate, one-sample t-tests with Holm-
Bonferroni corrections were used to determine whether the responses for each group differed
significantly from 0. Means are reported with standard deviations. Effect sizes are given as
Hedges *g* or partial eta-squared.

For conditions 1 and 3, a mixed ANOVA was used that included group (ataxic vs control)
and phase (end of perturbation and aftereffects), as well as the interaction between the two
factors. Similarly, for condition 4, the mixed ANOVA included group (ataxic vs control) and
perturbation direction (up and down), as well as the interaction between the two.

For condition 2, separate ANOVAs were used to evaluate differences between the ataxic
and control group in 1) the magnitude of the correction in response to the perturbation and 2)
the magnitude of correction in response to self-produced variability. These were conducted as
second-level analyses, on the beta coefficients estimated in the first-level analyses that were
conducted within each individual (see above). The dependent variable for the perturbation
correction model was the direction effect coefficient, estimated separately for each
perturbation direction (β_2, β_3). In addition to the perturbation direction factor (left vs right), this
model included a group factor as well as the interaction between perturbation direction and
group. For the variability correction, the dependent variable was the midpoint coefficient
representing the magnitude of correction for variability at reach midpoint in unperturbed (β_1),
left ($\beta_1+\beta_4$), and right ($\beta_1+\beta_5$) conditions. In addition to the main variable of interest, group, this
model also included perturbation condition (no perturbation, left, right). Welches' t-tests were
used to compare the ataxic and control groups on the reaction time, movement time, and hit
percentage in unperturbed trials.

In order to assess the relationship between feedback and feedforward control, we
conducted a set of planned correlational analyses using data from pairs of conditions. One set
of correlations compared the measures of feedback and feedforward control within each motor
domain (i.e., one correlation between the two reach tasks and a second between the two
speech tasks). We also performed a second set of correlations of similar forms of control across
the motor domains (i.e., one correlation of the feedforward measures from the reach and
speech tasks, and a second correlation of the feedback measures from the reach and speech
tasks). Because the primary aim of these analyses is to investigate the possibility that changes
in motor behavior due to cerebellar degeneration are correlated across sensory domains and
control systems, we limited these correlational analyses to the data from the CD group. We also
examined the extent to which our experimental measures of feedforward and feedback control
in the ataxic group were related to ataxia symptom severity, as measured by SARA speech and
upper limb subscores.

Results

Condition 1: Feedforward control in reaching.

In the baseline phase with visual feedback, the CD group was substantially more variable
than controls (standard deviation of 6.2 ± 2.0° vs 4.0 ± 0.8°, t(31.6) = 4.6 p < 0.0001, g = -0.89)
and less accurate (percentage of targets hit, 67 ± 25% vs 87 ± 15%, t(35.8) = -3.0, p = 0.005, g =
1.28). The CD group also produced slower movements than controls (347 ± 148 ms vs 264 ± 61
ms, t(31.6) = 2.4, p = 0.02, g = 0.67) and had longer reaction times (544 ± 116 ms vs 442 ± 58
ms, t(34.0) = 3.6, p = 0.001, g = 1.02).
Our principle outcome measure, adaptation, was operationalized as the change in
heading angle following the introduction of a 20° rotation of the feedback cursor. As can be
seen in Figure 2, the perturbation induced a gradual change in heading angle, with the
functions appearing to reach or approach asymptote by the end of the perturbation trials.
Adaptation in the control participants compensated for approximately 83% of the perturbation;
the comparable figure for the CD participants was only 45%. A decline in adaptation is visible
throughout the aftereffect phase in which there was no visual feedback and the following
washout phase in which veridical feedback was reintroduced.
Statistically, we first confirmed that the sign-dependent changes in heading angle were
significantly different than zero, the signature of adaptation. When measured during the final
trials of the perturbation block, the heading angle values were different than zero for both the
control (16.6 ± 2.7°, $t(14) = 24.1$, $p < 0.0001$, $g = 5.90$) and CD groups (9.2 ± 5.7°, $t(22) = 7.8$, $p <$
0.0001, $g = 1.56$). The adapted response persisted into the aftereffect block (controls: 15.1 ±
2.9°, $t(14) = 20.0$, $p < 0.0001$, $g = 4.87$; CD: 7.3 ± 7.5°, $t(22) = 4.7$, $p < 0.0001$, $g = 0.95$), although
the value was smaller than observed at the end of the perturbation block for both groups
($F(1,36) = 7.4$, $p = 0.01$, $\eta^2 = 0.17$). Across both the perturbation and aftereffect phases,
adaptation was significantly greater in the control group ($F(1, 36) = 20.1$, $p < 0.0001$, $\eta^2 = 0.36$)
and the Group x Phase interaction was not significant ($F(1,36) = 0.05$, $p = 0.82$, $\eta^2 = 0.001$).
These results are in accord with previous results (for a review, see Krakauer et al., 2019) in
showing that adaptation of feedforward control for reaching is impaired in individuals with CD.

Figure 2: In the visuomotor rotation task, the CD group reached a lower level of adaptation
and exhibited a smaller aftereffect relative to the control group. A: Hand angle over the course
of the experimental condition. Means and standard errors are shown for control participants
(blue) and individuals with cerebellar degeneration (red). Visual feedback is withheld in the
shaded phases. The perturbation is shown with a dashed black line. B,C: Individual data (semi-
transparent dots) and group means ± standard error for the CD and control participants during
the last 15 reaches of the perturbation phase (B) and 15-trial aftereffect phase in which the
visual feedback was withheld (C).

<u>Condition 2: Feedback control during reaching</u>
As in Condition 1, there were some kinematic differences between the two groups. The
most salient group difference was in movement accuracy: control participants hit the target
more often than the CD participants on the unperturbed trials (control: 85 ± 10 %, CD: 70± 24%,
t(31.9) = 2.7, *p* = 0.01, *g* = 0.75). There were small differences between the groups in reach
curvature, with controls showing a slight leftward shift from midpoint to target on unperturbed
trials and the CD group showing a slight rightward shift on these trials (control: -0.17 ± 0.193
cm, CD: 0.13 ± 0.52 cm, *t*(30.2) = 2.5, *p* = 0.02, *g* = 0.69, **Figure 3**F). Reaction times were also
slower in the CD group (control: 452 ± 79 ms, CD: 525 ± 138 ms, *t*(35.5) = 2.08, *p* = 0.045, *g* =
0.61) whereas movement time was similar between groups (control: 717 ± 114 ms, CD: 722 ±
114 ms, *t*(30.1) = 0.14, *p* = 0.89, *g* =0.05).
Our principle dependent measure, online corrective responses, was operationalized as a
lateral shift in the trajectory in response to perturbed feedback that was presented at the
midpoint of the movement, with the direction of the perturbation randomized across trials. As
can be seen in the data from a representative control participant (Fig 3A), the hand trajectory
deviated in the opposite direction of the perturbation, a signature of a feedback-based
response. The lateral shift accounted for roughly 33% of the perturbation magnitude in the

control group and 21% in the ataxic group (Fig 3B, D). These corrective feedback responses
were significantly different from 0 in both groups in response to the leftward perturbation (CD:
-0.21 ± 0.34 cm, $t(22) = 3.3$, $p = 0.003$, $g = 0.67$; control: -0.35 ± 0.30 cm, $t(14) = 4.6$, $p = 0.0005$,
$g = 1.12$) and rightward perturbation (CD: -0.20 ± 0.32 cm, $t(22) = 2.9$, $p = 0.007$ $g = 0.60$;
control: -0.34 ± 0.28 cm, $t(14) = 4.7$, $p = 0.0003$, $g = 1.15$). While the magnitude of the feedback-
based response was smaller for the CD group, this difference was not significant ($F(1,36) = 2.8$,
$p = 0.1$, $\eta^2 = 0.07$). There was no difference between the responses to the two perturbation
directions ($F(1,36) = 0.16$, $p = 0.69$, $\eta^2 = 0.004$) and the Group x Direction interaction was not
significant ($F(1,36) = 0.009$, $p = 0.93$, $\eta^2 = 0.0002$). These results indicate that feedback-based
corrective responses to visual perturbations are not enhanced in individuals with CD; indeed, as
a group the trend was for attenuated feedback responses.
In a second measure of feedback control, we looked at adjustments in the reach
trajectory in response to motor variability (estimated as the change in horizontal position from
reach midpoint to endpoint unrelated to the visual perturbation; model parameters β_1, $\beta_1+\beta_4$,
and $\beta_1+\beta_5$). Although both groups showed a slight shift in trajectory in the compensatory
direction in response to the position of the cursor at reach midpoint (**Figure 3**C,E,G), this shift
was not significantly different from 0 after correction for multiple comparisons (CD: -0.010 ±
0.035, $p = 0.05$, $g = -0.30$; control: -0.009 ± 0.042, $p = 0.24$, $g = -0.21$). There was no difference
between the two groups ($F(1,36) = 0.01$, $p = 0.91$, $\eta^2 = 0.0004$), nor any interaction between
group and direction ($F(2,72) = 2.4$, $p = 0.10$, , $\eta^2 = 0.06$). However, there was a significant effect
of perturbation direction ($F(2,72) = 7.3$, $p = 0.001$, , $\eta^2 = 0.17$) such that the shift in trajectory
was greater in the presence of leftward perturbations (-0.021 ± 0.0332) compared to trials with
no perturbation (0.001 ± 0.038; $p = 0.008$, $g = 0.62$). There was a similar trend in for trials with
rightward perturbations (-0.010 ± 0.036), but this did not reach significance ($p = 0.15$, $g = 0.33$).
In sum, there is little evidence that either group corrected for self-produced variability at reach
midpoint.
467

468
Figure 3: In the reaching feedback control task, both the CD and control groups showed a
robust on-line correction to perturbed visual feedback presented at the midpoint of the reach,
with no group differences. *A: Example data from one control participant showing average*
reach trajectories for unperturbed reaches (black), reaches with a 1 cm rightward visual
perturbation (teal), and reaches with a 1 cm leftward visual perturbation (purple). Dots in
corresponding colors show the location of the visual feedback. The corrective response only
partially corrects for the perturbation. *B, D*: *Compensatory response to the feedback*
perturbation. *F: Estimate of the change in horizontal hand position from reach midpoint on*
unperturbed trials. *C, E, G: Estimates of the effect of true hand position at reach midpoint on the*
compensatory response for leftward (C), rightward (E), and unperturbed (G) trials. Across all
plots, compensatory responses have negative values. Semi-transparent dots represent
individuals.

481
<u>Condition 3: Feedforward control in speech</u>
Adaptation was operationalized as the change in F1 in response to a -125 mel F1 shift in
the feedback heard by participants, introduced via the real-time resynthesis of their utterances.
As can be seen in Figure 4A, this auditory perturbation caused a gradual change in F1 that
opposed the perturbation. At asymptote, the adaptive response had reached approximately
40% of the perturbation in the control group and 25% in the CD group. The lower degree of
compensation relative to reaching is consistent with previous studies (Houde and Jordan, 1998;
Purcell and Munhall, 2006; Parrell et al., 2017). Note that, unlike Condition 1, post-perturbation
trials (washout phase) always included veridical feedback; we did not include a no-feedback
phase as masking auditory feedback with noise leads to substantial changes in speech
(Lombard, 1911; Summers et al., 1988).

We first confirmed that both groups adapted during the perturbation phase (CD: 31 ± 39 mels, *t*(1,22) = 3.9, *p* = 0.0007, *g* = 0.79; control: 53 ± 31 mels, *t*(1,14) = 6.6, *p* < 0.0001, *g* = 1.61; Figure 4B) and maintained their adapted state during the initial part of the washout phase (CD: 21 ± 32 mels, *t*(1,22) = 3.1, *p* = 0.005, *g* = 0.64; control: 40 ± 23 mels, *t*(1,14) = 6.6, *p* < 0.0001, *g* = 1.62; Figure 4C). The CD group exhibited less adaptation than the control participants (*F*(1,36) = 4.7, *p* = 0.036, , η^2 = 0.12), and this difference was similar in both phases (Group x Phase interaction: *F*(1,36) = 0.06, *p* = 0.80, η^2 = 0.002). In sum, we find that sensorimotor adaptation in speech is impaired in individuals with CD, similar to the impairment observed for reach adaptation in Condition 1 and consistent with previous findings (Parrell et al., 2017),

***Figure 4: In the speech adaptation task, the CD group adapted less than controls. A:** Change in first formant over the course of the experimental condition. Means and standard errors are shown for control (blue) and CD participants (red). For each participant, the change was calculated relative to their mean F1 value during the second half of the baseline phase, with positive values corresponding to changes in the opposite direction of perturbation. The perturbation is shown with a dashed black line (sign flipped).*
***B,C:** Group means ± standard error (solid dots) and individual data points (semi-transparent dots) of adaptive response during the last 10 trials of the perturbation phase and first 10 trials of the washout phase.*

Condition 4: : Feedback control in speech

Online corrective responses for speech were operationalized as the change in F1 during the time window from 200-300 ms after vowel onset in response to an upward or downward perturbation of F1, randomized across trials. The data are plotted relative to F1 values measured on unperturbed trials. As can be seen in Figure 5A, the non-predictable auditory perturbations resulted in compensatory responses that opposed the perturbation in both groups. The magnitude of the corrective response (7.6% in controls, 9.2% in the ataxic group) was similar to that typically observed in response to auditory perturbations of speech and much lower than observed for the non-predictive perturbations during reaching used in Condition 2.

520   The change on perturbed trials was significantly larger than 0 in response to upward and
521 downward perturbations in the CD group (up: 10.2 ± 13.6 mels, $t(1,22)$ = 3.6, p = 0.002, g =
522 0.73; down: 11.8 ± 16.0 mels, $t(1,22)$ = 3.5, p = 0.002, g = 0.71) and control group (up: 11.0 ±
523 14.4 mels, $t(1,13)$ = 2.9, p = 0.02, g = 0.72; down: 8.2 ± 12.4 mels, $t(1,13)$ = 2.9, p = 0.02, g =
524 0.72) (Figure 5B, C). While the mean values were larger in the CD group and individuals in this
525 group showed the largest compensatory response, the difference between the two groups was
526 not significant ($F(1,35)$ = 0.35, p = 0.56, η^2 = 0.009). There were no differences between the
527 responses to the two perturbation directions ($F(1,35)$ = 0.1, p = 0.74, , η^2 = 0.003) and the
528 Group x Direction interaction was not significant ($F(1,35)$ = 0.07, p = 0.80, η^2 = 0.002). These
529 results suggest feedback gains for auditory perturbations in speech are similar in individuals
530 with CD and healthy controls, consistent with the reaching results in Exp 2. Of note, the null
531 effects here are inconsistent with the results from previous studies involving speech
532 articulation (Parrell et al., 2017) and vocal pitch production (Houde et al., 2019; Li et al., 2019)
533 in which individuals with CD were found to show an enhanced feedback response.

534
535 ***Figure 5: In the speech feedback control task, both*** *the CD group and controls corrected for formant*
536 **perturbations in both directions, with no difference between the two groups.** *A: Normalized formant*
537 *trajectories in response to a downward F1 perturbation (solid lines) or upward perturbation (dotted lines)*
538 *in the control (blue) and CD (red) groups. Shaded area indicates standard error. Data are normalized to*
539 *the formant trajectories produced in trials with no perturbation. B, C: Magnitude of F1 values in trials*
540 *with a downward (**B**) or upward (**C**) perturbation, averaged over the time window spanning from 150-*
541 *300 ms following onset of vocalic portion of the utterances. Compensatory responses are expressed as*
542 *positive values in both cases. Group means ± standard error are shown as solid dots;* s*emi-transparent*
543 *dots represent individuals.*

544 <u>Feedforward and feedback control within and across motor domains</u>
545   By testing each participant in all four conditions, we can compare the measures of
546 feedback and feedforward control within and across task domains. Because our focus in this

analysis is how deficits in these domains may be correlated in individuals with cerebellar
degeneration, we limit this analysis to the 22 individuals with CD.
The between-domain comparisons assess the similarity of impairment (or lack thereof)
between reaching and speech. Although the ataxic group adapted less than the controls at the
group level, there was no significant correlation within the ataxic group between the magnitude
of adaptation between speech and reaching (Figure 6A). Similarly, the magnitude of feedback-
based compensation (Figure 6B) was unrelated across motor domains.
The within domain correlations provide a test of the hypothesis that feedback gains may
increase in response to impaired feedforward control: in this case, we would expect a negative
correlation between these measures such that individuals who are more impaired in
feedforward control should be more likely to exhibit a greater reliance on feedback control.
This relationship might hold even if there is no overall increase in compensatory responses
when comparing the CD and control groups. Contrary to this prediction, the relationship
between feedforward adaptation and feedback-based compensation was not significant in
either reaching (Figure 6C) or speech (Figure 6D).
We additionally tested whether any of our behavioral measures correlated with ataxia
severity as assessed with the SARA. Neither a summary measure of overall upper limit ataxia
nor intentional tremor severity were correlated with reach adaptation or compensation (all $p >$
0.2). Similarly, overall speech impairment was not correlated with speech compensation ($r = -$
0.24, $p = 0.26$). We did observe a positive correlation between the SARA measure of speech
impairment and speech adaptation, with lower levels of adaptation to the auditory
perturbation associated with greater speech impairment ($r = 0.45$, $p = 0.03$).
569

570
Figure 6: Correlational analyses of feedforward adaptation and feedback-based compensation in
individuals with cerebellar degeneration. Within the CD group, there were no correlations in behavioral
*measures obtained across (**A-B**) or within (**C-D**) motor domain. Plots show individual CD participants as*

red dots and the mean of the control participants as a blue dot. To simplify interpretation, the sign of
compensation in reaching has been flipped such that larger positive values reflect greater compensation,
*as for all other measures. **A:** Adaptation in reaching and speech. **B:** Compensation in reaching and*
*speech. **C:** Adaptation and compensation in reaching. **D:** Adaptation and compensation in speech.*

Discussion
We conducted a set of experiments to evaluate the impact of cerebellar degeneration on
feedforward and feedback control in two motor domains, reaching and speech. Individuals with
cerebellar degeneration showed a marked impairment in feedback control relative to controls,
manifest as reduced adaptation in response to a sensory perturbation that remained constant
from trial to trial. Feedback control, measured in terms of the on-line response to a variable
perturbation, was intact in both motor domains.
586
Multi-modal impairment in feedforward control in individuals with cerebellar degeneration
Our principle positive result, that individuals with cerebellar degeneration are impaired
in adapting their motor behavior in the presence of sensory prediction errors, is consistent with
prior neuropsychological studies involving upper limb control and speech. These results are
consistent with the hypothesis that the cerebellum provides a domain-general mechanism for
generating sensory predictions and using error information to keep this predictive system well-
calibrated. In particular, these results agree with evidence that the cerebellum is critical for
implicitly updating motor behavior. Adaptation in speech is highly likely to be an implicit
process (Kim and Max, 2020; Lametti et al., 2020): the majority of participants are unaware of
the auditory perturbation (Parrell and Niziolek, 2020), and adaptation is similar whether
participants are unaware of the perturbation or made aware of the perturbation and explicitly
told to ignore the feedback (Munhall et al., 2009). The design for the reaching task in the
current study, one in which the perturbation was only 20°, was chosen to also primarily engage
implicit processes (Bond and Taylor, 2015; Morehead et al., 2015; Werner et al., 2015).
While adaptation was impaired in both reaching and speech at the group level, there
was no correlation between the magnitude of adaptation in reaching and speech in the CD
group. That is, individuals who were more impaired in reaching were not more likely to be
impaired in speech. It is possible that the failure to find such a relationship is due either to
unreliable estimates of adaptation and compensation in each domain (given that we only
tested each condition a single time, and no previous work, to our knowledge, has established
the reliability of these measures) or to a lack of power given our sample size of 22 individuals
with CD. However, previous work has identified a similar dissociation in individuals with CD
between adaptation to dynamic (force-field) and kinematic (visuomotor rotation) perturbations
during reaching, and deficits in adapting reach dynamics and reach kinematics have been
localized to different regions in the cerebellum in both individuals with CD (Rabe et al., 2009)
and acute cerebellar lesions (Donchin et al., 2012). Our results further specify this dissociation
by showing that cerebellar mechanisms of adapting kinematic perturbations to different motor
effectors do not overlap. Given somatotopic representations across subregions of the
cerebellar cortex (e.g., Marvel and Desmond, 2010; Mottolese et al., 2013), we would

anticipate that deficits in speech adaptation would be associated with more medial cerebellar
regions, relative to deficits in limb adaptation, though future work examining patterns of
cerebellar damage in patients will be needed to confirm this hypothesis.
619
Intact, but not enhanced feedback control in individuals with cerebellar degeneration
Feedback-based corrections for errors were similar in magnitude between individuals
with cerebellar degeneration and controls for both speech and reaching. Although this result
agrees with estimates of feedback gains in a continuous visuomotor tracking task (Zimmet et
al., 2020), the absence of a difference between the CD and control groups on the speech task
fails to replicate a previous finding from our labs showing an enhanced feedback response to a
similar perturbation (Parrell et al., 2017). Moreover, a much larger increase in the feedback
response, relative to controls, has been observed in response to auditory perturbations of pitch
(Houde et al., 2019; Li et al., 2019). Our previous results had led to the hypothesis that the
enhanced feedback response reflected a compensatory mechanism, one to help offset the
disruptive effects of impairment in feedforward control. The failure to observe enhanced
feedback in the current study in both domains argues against this compensatory hypothesis.
The absence of a correlation between the feedforward and feedback measures within both
motor domains also argues against a compensatory hypothesis.
There are a few issues to consider in terms of the discrepancy between our results and
previous studies as well as the interpretation of the null results regarding the compensatory
hypothesis. First, we may have failed to find any changes in feedback control in individuals with
CD simply due to sampling issues. Our sample of this population (n = 23) is consistent with, or
larger than, many previous studies (Morton and Bastian, 2006; Parrell et al., 2017; e.g., Houde
et al., 2019; Li et al., 2019; Zimmet et al., 2020), and we were adequately powered to detect
relatively large between-group effects (.75 power to detect *d* of .8). However, it is possible the
effect size of the expected increase in feedback gains in certain domains, as in our previous
work on oral articulation, may be relatively small. Thus, we may simply have been under-
powered to detect any changes.
Second, it may be that increased feedback gains are a secondary, and somewhat
sporadic, effect of cerebellar degeneration. The consistent and striking deficit in feedforward
control points to inaccurate or attenuated predictive signals. In this case, some individuals may
learn to rely more on sensory feedback to help with movement accuracy as a compensatory
mechanism for impairments in feedforward control, though at some loss of movement speed
and fluidity. It may be that our sample did not include enough participants with altered
feedback gains to show an effect at the group level. If this is the case, we might still expect to
find some evidence that individuals with larger feedback gains have greater impairments in
feedforward control, even if there were no group differences. However, we found no evidence
of this negative correlation. While this does not eliminate this explanation, it may make it an
unlikely account.
A final potential explanation is that increases in feedback gains are domain-specific
while impairments in feedforward control are more general, at least at the group level. This
may have to do with domain-specific use of feedback for sensorimotor control. The strongest
example of higher feedback gains in individuals with cerebellar degeneration comes from work
on pitch control, where the on-line response to a pitch perturbation was roughly twice as large

in a CD group compared to a control group (Houde et al., 2019; Li et al., 2019). This increase
may result from an inherent reliance on auditory feedback for pitch control, compared to
higher reliance on feedforward control for speech and reaching. There is some evidence that
pitch control is indeed heavily weighted towards online feedback control even in healthy
individuals: pitch control rapidly degenerates after post-lingual hearing loss while oral
articulation is better maintained (Cowie and Douglas-Cowie, 1983; Lane and Webster, 1991).
Thus, cerebellar degeneration may cause increased feedback gains only in those motor domains
which are already primarily reliant on sensory-feedback-based control. The lack of any
correlation between the magnitude of feedback-based corrections across motor domains is
consistent with the idea that changes in feedback use may be domain-specific.
670
Conclusions
Adaptation of feedforward control based on sensory errors is impaired in individuals
with cerebellar degeneration in both reach and speech. Interestingly, these individuals
exhibited intact feedback control in both domains. Contrary to our initial hypothesis and data
from vocal pitch control, we found no evidence for increased feedback gains in either domain.
However, these results, together with those from a recent study of upper extremity control
(Zimmet et al., 2020), motivate further investigation into how feedback gains may be
differentially affected by the specific demands of different motor tasks, as well as to determine
the variability in feedback control associated with cerebellar dysfunction.
680

References
Beppu H, Nagaoka M, Tanaka R (1987) Analysis of Cerebellar Motor Disorders By Visually-
Guided Elbow Tracking Movement 2. Contribution of the Visual Cues on Slow Ramp
Pursuit. Brain 110:1–18.

Boersma P, Weenink D (2019) Praat: doing phonetics by computer. Available at:
ttp://www.praat.org/.

Bond KM, Taylor JA (2015) Flexible explicit but rigid implicit learning in a visuomotor adaptation
task. J Neurophysiol 113:3836–3849.

Brainard DH (1997) The Psychophysics Toolbox. Spatial Vision 10:433–436.

Cai S, Beal DS, Ghosh SS, Tiede MK, Guenther FH, Perkell JS (2012) Weak responses to auditory
feedback perturbation during articulation in persons who stutter: evidence for abnormal
auditory-motor transformation. PLoS One 7:e41830.

Cai S, Boucek M, Ghosh S, Guenther FH, Perkell J (2008) A System for Online Dynamic
Perturbation of Formant Trajectories and Results from Perturbations of the Mandarin
Triphthong /iau/. In: Proceedings of the 8th International Seminar on Speech
Production, pp 65–68. Strasbourg, France.

Cowie RID, Douglas-Cowie E (1983) 6 - Speech Production in Profound Postlingual Deafness. In:
Hearing Science and Hearing Disorders, pp 183–230. London: Academic Press. Available

699   at: https://www.sciencedirect.com/science/article/pii/B9780124604407500119
700   [Accessed January 16, 2018].

701 Daliri A, Chao S-C, Fitzgerald LC (2020) Compensatory Responses to Formant Perturbations
702   Proportionally Decrease as Perturbations Increase. Journal of Speech, Language, and
703   Hearing Research 63:3392–3407.

704 Day BL, Thompson PD, Harding AE, Marsden CD (1998) Influence of vision on upper limb
705   reaching movements in patients with cerebellar ataxia. Brain 121 (Pt 2):357–372.

706 Donchin O, Rabe K, Diedrichsen J, Lally N, Schoch B, Gizewski ER, Timmann D (2012) Cerebellar
707   regions involved in adaptation to force field and visuomotor perturbation. J
708   Neurophysiol 107:134–147.

709 Houde JF, Gill JS, Agnew Z, Kothare H, Hickok G, Parrell B, Ivry RB, Nagarajan SS (2019)
710   Abnormally increased vocal responses to pitch feedback perturbations in patients with
711   cerebellar degeneration. The Journal of the Acoustical Society of America 145:EL372–
712   EL378.

713 Houde JF, Jordan IM (1998) Sensorimotor adaptation in speech production. Science 279:1213–
714   1216.

715 Kim KS, Max L (2020) Speech auditory-motor adaptation lacks an explicit component: reduced
716   adaptation in adults who stutter reflects limitations in implicit sensorimotor learning.
717   bioRxiv:2020.09.06.284638.

718 Kleiner M, Brainard D, Pelli D, Ingling A, Murray R, Broussard C (2007) What's new in
719   psychtoolbox-3. Perception 36:1–16.

720 Körding KP, Wolpert DM (2004) Bayesian integration in sensorimotor learning. Nature 427:244–
721   247.

722 Krakauer JW, Hadjiosif AM, Xu J, Wong AL, Haith AM (2019) Motor Learning. Compr Physiol
723   9:613–663.

724 Kurtzer I, Trautman P, Rasquinha RJ, Bhanpuri NH, Scott SH, Bastian AJ (2013) Cerebellar
725   damage diminishes long-latency responses to multijoint perturbations. Journal of
726   Neurophysiology 109:2228–2241.

727 Lametti DR, Quek MYM, Prescott CB, Brittain J-S, Watkins KE (2020) The perils of learning to
728   move while speaking: One-sided interference between speech and visuomotor
729   adaptation. Psychon Bull Rev 27:544–552.

730 Lane H, Webster JW (1991) Speech deterioration in postlingually deafened adults. The Journal
731   of the Acoustical Society of America 89:859–866.

Li W, Zhuang J, Guo Z, Jones JA, Xu Z, Liu H (2019) Cerebellar contribution to auditory feedback
control of speech production: Evidence from patients with spinocerebellar ataxia. Hum
Brain Mapp 40:4748–4758.

Lombard E (1911) Le signe de l'elevation de la voix. Annale des Maladies de L'Oreille et du
Larynx 37:2.

Martin TA, Keating JG, Goodkin HP, Bastian AJ, Thach WT (1996) Throwing while looking
through prisms. I. Focal olivocerebellar lesions impair adaptation. Brain 119 (Pt
4):1183–1198.

Marvel CL, Desmond JE (2010) Functional Topography of the Cerebellum in Verbal Working
Memory. Neuropsychol Rev 20:271–279.

Miall RC, Christensen LOD, Cain O, Stanley J (2007) Disruption of state estimation in the human
lateral cerebellum. PLoS Biol 5:e316.

Morehead JR, Qasim SE, Crossley MJ, Ivry R (2015) Savings upon Re-Aiming in Visuomotor
Adaptation. J Neurosci 35:14386–14396.

Morton SM, Bastian AJ (2006) Cerebellar contributions to locomotor adaptations during
splitbelt treadmill walking. J Neurosci 26:9107–9116.

Mottolese C, Richard N, Harquel S, Szathmari A, Sirigu A, Desmurget M (2013) Mapping motor
representations in the human cerebellum. Brain 136:330–342.

Munhall GK, MacDonald EN, Byrne SK, Johnsrude I (2009) Talkers alter vowel production in
response to real-time formant perturbation even when instructed not to compensate.
Journal of the Acoustical Society of America 125:384–390.

Niziolek CA, Guenther FH (2013) Vowel category boundaries enhance cortical and behavioral
responses to speech feedback alterations. Journal of Neuroscience 33:12090–12098.

Niziolek CA, Houde J (2015) Wave_Viewer: First Release. Available at:
https://zenodo.org/record/13839 [Accessed January 30, 2020].

Parrell B, Agnew Z, Nagarajan S, Houde JF, Ivry RB (2017) Impaired Feedforward Control and
Enhanced Feedback Control of Speech in Patients with Cerebellar Degeneration. Journal
of Neuroscience 37:9249–9258.

Parrell B, Niziolek C (2020) Increased speech contrast induced by sensorimotor adaptation to a
non-uniform auditory perturbation. Journal of Neurophysiology Available at:
https://journals.physiology.org/doi/abs/10.1152/jn.00466.2020 [Accessed January 13,
2021].

Pelli DG (1997) The VideoToolbox software for visual psychophysics: transforming numbers into
movies. Spat Vis 10:437–442.

Purcell DW, Munhall KG (2006) Adaptive control of vowel formant frequency: evidence from
real-time formant manipulation. Journal of Acoustical Society of America 120:966–977.

R Core Team (2013) R: A Language and Environment for Statistical Computing. Vienna, Austria:
R Foundation for Statistical Computing. Available at: http://www.R-project.org/.

Rabe K, Livne O, Gizewski ER, Aurich V, Beck A, Timmann D, Donchin O (2009) Adaptation to
visuomotor rotation and force field perturbation is correlated to different brain areas in
patients with cerebellar degeneration. J Neurophysiol 101:1961–1971.

Schlerf JE, Xu J, Klemfuss NM, Griffiths TL, Ivry RB (2013) Individuals with cerebellar
degeneration show similar adaptation deficits with large and small visuomotor errors. J
Neurophysiol 109:1164–1173.

Shadmehr R, Krakauer JW (2008) A computational neuroanatomy for motor control.
Experimental Brain Research 185:359–381.

Smith MA, Shadmehr R (2005) Intact ability to learn internal models of arm dynamics in
Huntington's disease but not cerebellar degeneration. J Neurophysiol 93:2809–2821.

Summers WV, Pisoni DB, Bernacki RH, Pedlow RI, Stokes MA (1988) Effects of noise on speech
production: acoustic and perceptual analyses. Journal of Acoustical Society of America
84:917–928.

Tourville JA, Cai S, Guenther F (2013) Exploring auditory-motor interactions in normal and
disordered speech. In: Proceedings of Meetings on Acoustics, pp 060180 Available at:
http://asa.scitation.org/doi/abs/10.1121/1.4800684 [Accessed April 9, 2020].

Tourville JA, Reilly KJ, Guenther FH (2008) Neural mechanisms underlying auditory feedback
control of speech. Neuroimage 39:1429–1443.

Tseng Y-WW, Diedrichsen J, Krakauer JW, Shadmehr R, Bastian AJ (2007) Sensory prediction
errors drive cerebellum-dependent adaptation of reaching. Journal of Neurophysiology
98:54–62.

Werner S, Aken BC van, Hulst T, Frens MA, Geest JN van der, Strüder HK, Donchin O (2015)
Awareness of Sensorimotor Adaptation to Visual Rotations of Different Size. PLOS ONE
10:e0123321.

Wolpert DM, Miall RC, Kawato M (1998) Internal models in the cerebellum. Trends Cogn Sci
2:338–347.

Xu-Wilson M, Chen-Harris H, Zee DS, Shadmehr R (2009) Cerebellar contributions to adaptive
control of saccades in humans. J Neurosci 29:12930–12939.

Zimmet AM, Cao D, Bastian AJ, Cowan NJ (2020) Cerebellar patients have intact feedback
control that can be leveraged to improve reaching Ivry RB, ed. eLife 9:e53246.

Appendix

Table A1: Characteristics of individuals with cerebellar degeneration. One participant, shown in
gray, was excluded due to a MOCA score indicative of moderate cognitive impairment.

Diagnosis	Age	MOCA	SARA overall score	SARA upper limb subscore	SARA intention tremor subscore	SARA speech subscore
AOA2	42	27	22	6	2.5	2
SCA3	61	28	16	5	1.5	3
SCA3	75	26	16.5	5.5	1.5	0
SCA3	65	25	24	7	2	1
SCA6	77	22	8	2	1	1
SCA6	70	23	7.75	3.25	0.5	0
SCA6	60	26	3	0.5	0	0
SCA6	49	26	1	1	0	0
SCA6	65	24	9	3.5	0.5	0
SCA6	57	27	6	3	0	0
SCA8	57	27	11	3.5	1	0
SCA8	53	30	3.5	2	0	0
SCA, sporadic	57	22	10	4	1	2
SCA, unknown	65	17	25	6	2	3
SCA, unknown	89	26	9.5	2	0.5	4
SCA, unknown	68	27	23	5	2	1
SCA, unknown	64	25	8	1.5	0.5	1
SCA, unknown	37	27	14	3	1	2
SCA, unknown	72	27	18.5	4	1.5	1
SCA, unknown	64	28	11	3.5	0	1.5
SCA, unknown	54	28	8.5	4.5	0.5	3
SCA, unknown	65	25	9.5	2.5	1	0
SCA, unknown	62	30	12	5.5	1	1

806