TOOLS AND RESOURCES

Novel human cell expression method reveals the role and prevalence of posttranslational modification in non-muscle tropomyosins

Peter J. Carman[1,2], Kyle R. Barrie[1,2], Roberto Dominguez[1,*]

[1] Department of Physiology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, 19104, USA

[2] Biochemistry and Molecular Biophysics Graduate Group, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, 19104, USA

* Correspondence: droberto@pennmedicine.upenn.edu

Abstract

Biochemical studies require large protein quantities, which are typically obtained using bacterial expression. However, the folding machinery of bacteria is inadequate for many mammalian proteins, which additionally undergo posttranslational modifications (PTMs) that bacteria, yeast, or insect cells cannot perform. Many proteins also require native N- and C-termini and cannot tolerate extra tag amino acids for function. Tropomyosin, a coiled coil that decorates most actin filaments in cells, requires both native N- and C-termini and PTMs, specifically N-terminal acetylation, to polymerize along actin filaments. Here, we describe a new method that combines native protein expression in human cells with an intein-based purification tag that can be precisely removed after purification. Using this method, we expressed several non-muscle tropomyosin isoforms. Mammalian cell-expressed tropomyosins are functionally different from their *E. coli*-expressed counterparts, display multiple types of PTMs, and can form heterodimers. This method can be extended to other proteins, as demonstrated here for α-synuclein.

Introduction

Biochemical and structural studies require large amounts of pure proteins and protein complexes. When these cannot be purified from natural sources, scientists typically opt for protein expression in bacteria, and N- or C-terminal affinity tags are commonly used for protein purification. These tags are often left on proteins during subsequent studies and, when they are enzymatically removed, extra amino acids often remain (1). While these approaches have proven powerful and will likely continue to be used in the future, there are many circumstances in which they fail to produce accurate results. For example, most human proteins undergo multiple types of posttranslational modifications (PTMs), including acetylation, phosphorylation, and methylation (2). Such PTMs play crucial roles in the regulation of protein function, including protein stability, protein-protein interactions, and enzymatic activity (3-6). Most PTMs present in human proteins cannot be processed by expression systems such as bacteria, yeast, or insect cells (7-12). These expression systems may also lack the adequate machinery to fold certain mammalian proteins (13-16). Many proteins also require native N- and C-termini and cannot tolerate the presence of extra amino acids from purification tags for proper function (17-21). We were confronted with these problems while studying tropomyosin (Tpm), a protein of central importance in the actin cytoskeleton (22).

52
Tpm is a coiled coil dimer that polymerizes head-to-tail to form two contiguous chains on
opposite sides of the actin filament (F-actin). Through alternative splicing of four different genes
(*TPM1-4*), up to 40 distinct Tpm isoforms can be produced, of which ~30 have been detected as
proteins (23). Most isoforms are ~280-amino acids long (called long isoforms), comprising
seven pseudo-repeats of ~40 amino acids, with each pseudo-repeat spanning the length of one
actin subunit along the long-pitch helix of the actin filament. Long Tpm isoforms decorate the
contractile actin filaments of muscle cells. In non-muscle cells, the majority of actin filaments are
also decorated with Tpm (24). Some of the non-muscle Tpm isoforms lack exon 2 and as a
consequence are one pseudo-repeat shorter (called short isoforms). Because the six human
actin isoforms are very similar, sharing 93–99% sequence identity, it has been proposed that
the decoration of actin filaments with distinct Tpm isoforms provides a mechanism for filaments
to assume different identities and become functionally segregated in cells (25). Tpm can move
azimuthally on actin, helping to regulate the interactions of most actin-binding proteins, including
myosin (26,27).
67
Although there is high sequence similarity among Tpm isoforms, gene knockout and isoform
overexpression studies have shown that they play non-redundant roles (22). For example,
Tpm4.2 knockout causes low platelet count (28), whereas Tpm3.5 knockout causes softer, less
mechanically resilient lenses in the eye (29), indicating that other Tpm isoforms cannot
compensate for the lack of Tpm4.2 or Tpm3.5. Similarly, different Tpm isoforms have been
implicated in processes such as muscle contraction (30,31), cytokinesis (32,33), embryogenesis
(34,35), and endocytosis (36). The temporal expression of Tpm isoforms is also tightly
controlled. Thus, sixteen isoforms are up- or down-regulated at different stages of brain
development in rodents (37,38). Tpm3.1 and Tpm3.2 are found in the axon in developing
neurons, whereas in mature neurons they are expressed only in cell bodies and replaced in
axons by Tpm1.12 and Tpm4.2 (39).
79
PTMs, and in particular N-terminal acetylation (Nt-acetylation) (40-42) and phosphorylation (43-
46), have been shown to critically regulate Tpm function. Proper head-to-tail assembly of Tpm
coiled coils along actin filaments requires native N- and C-termini, i.e. free of extra amino acids
from tags and Nt-acetylated (40,47), whereas the phosphorylation state of Tpm can also
modulate the stiffness and affinity of the head-to-tail interaction as well as the F-actin binding
affinity (43,44,46). Despite the importance of these factors, previous biochemical studies have
focused on one or two Tpm isoforms purified from muscle or *E. coli*-expressed full-length and
peptide fragments of Tpm isoforms that lack PTMs and native N and C-termini. While extra
amino acids have been added at the N-terminus to substitute for Nt-acetylation (40,41,48,49), it
is unclear how precisely this approach mimics acetylation. Tpm coexpression with the yeast
acetyltransferase NatB in *E. coli* has been also used to acetylate the N-terminus (50). However,
while yeast NatB can acetylate the N-terminus, the extent of this modification varies for different
Tpm isoforms, being as low as 30% for certain mammalian Tpm isoforms, whereas other types
of PTMs are not accounted for (8). Furthermore, as we show here, the initiator methionine of
many Tpm isoforms must be removed by mammalian N-terminal aminopeptidases, and Nt-
acetylation occurs on the second residue, which this approach can also not account for (51).
96
Here, we describe a method that combines expression in human Expi293F cells, which grow in
suspension to high density and contain the native PTM machinery, with an intein-based affinity
tag that can be precisely removed through self-cleavage after purification. We applied this
method to the expression of six Tpm isoforms. We demonstrate the general applicability of this
method to other proteins by also showing expression of α-synuclein, a protein implicated in
several neurodegenerative disorders whose function is critically regulated by numerous PTMs

(52,53). We characterize the expressed proteins, including a functional analysis of the
expressed Tpm isoforms in comparison with their *E. coli*-expressed counterparts, and a
comprehensive analysis of PTMs using proteomics analysis. We finally show that non-muscle
Tpm isoforms can form heterodimers *in vitro* and in cells, as previously shown for muscle
isoforms purified from tissues.
108
109
110 **Results**
111
112 **Human cell expression and intein-mediated purification method and its application to**
113 **Tpm and α-synuclein**
114
The expression method developed here combines mammalian cell expression with a self-
cleavable affinity tag. The cells used are Expi293F, commercialized by Thermo Fisher Scientific,
which are based on human embryonic kidney (HEK) cells and have been adapted to grow in
suspension to high density (54,55). These cells can be transiently transfected, display high
protein expression levels, and have the endogenous protein folding and PTM machineries
necessary for native expression and modification of human proteins. The self-cleavable affinity
tag consists of a modified intein element followed by a chitin-binding domain (CBD), and is
based on the IMPACT (intein-mediated purification with an affinity chitin-binding tag) system
marketed by New England Biolabs (56,57). Both, N- and C-terminal intein-CBD tags are
commercially available for bacterial protein expression, but we opted here to use a C-terminal
intein-CBD tag for expression in mammalian cells to allow for N-terminal processing and
acetylation, which affects ~90% of human proteins (58). After purification on the high-affinity
chitin resin, self-cleavage of the intein takes place precisely after the last endogenous amino
acid of the expressed target protein, leaving no extra amino acids from the tag. While both
mammalian cell expression (59-61) and intein-CBD purification tags (62) have been previously
used, to our knowledge this is the first time that the two are combined in a single method. Self-
cleavage of the intein is thiol-catalyzed, meaning that a portion of the intein of bacterial origin is
prematurely activated in the reducing environment of the cytoplasm of eukaryotic cells, likely
explaining why they have not been used for mammalian cell expression.
134
We built a new vector (named pJCX4), which incorporates the *Mxe* GyrA intein and CBD into a
mammalian-cell expression vector (*Figure 1A*). The *Mxe* GyrA intein is a 198-amino acid self-
splicing protein from *Mycobacterium xenopi* (63). In its native form, the intein catalyzes its own
excision from the precursor protein, forming a new peptide bond between the N- and C-terminal
fragments that flank the intein (64,65). To harness this process for expression and purification of
a target protein, the N- and C-terminal fragments of the intein precursor protein are substituted
respectively by the target protein and the CBD affinity tag, and two amino acids at the C-
terminus of the intein moiety (Asn-Cys) are mutated to Ala-Thr, which turns the intein splicing
activity into precise cleavage at the junction between the target protein and the intein (66).
144
In addition to the intein-CBD element, the expression vector includes the cytomegalovirus
promoter (CMV) and the simian virus 40 (SV40) polyadenylation (polyA) signal, as well as the
woodchuck hepatitis virus posttranscriptional regulatory element (WPRE) for enhanced
expression (*Figure 1A*). It also includes pUC and f1 origins of replication for propagation in *E.*
coli and the SV40 origin of replication for propagation in mammalian cells. Finally, the vector
contains the kanamycin resistance gene for selection in bacteria. These elements collectively
allow for plasmid cloning and amplification in *E. coli* and expression in Expi293F cells. A number
of restriction sites are available to clone the 5' end of a target gene, while the SapI or SpeI sites
must be used to clone the 3' end, to ensure that no vector- or tag-derived amino acids remain

Figure 1. Cloning, expression and purification of Tpm isoforms and α-synuclein in Expi293F cells. (A) Schematic representation of the mammalian-cell expression vector designed here, featuring an intein-CBD affinity-purification and self-cleavage element. Cloning into this vector results in the expression of the target protein as a fusion protein with intein-CBD (see also *Figure S1*). (B) Schematic representation of the expression and purification steps, including purification on a chitin resin of the protein expressed in Expi293F cells, thiol-catalyzed self-cleavage of the intein (using β-mercaptoethanol or βME), and elution of the target protein. As an example, SDS-PAGE analysis at the bottom of the panel illustrates the progression of these steps for Tpm3.1 (see also *Figure S1*). (C and D) SDS-PAGE analysis and Coomassie Brilliant Blue R-250 (BIO-RAD) staining of the six Tpm isoforms (panel C) and α-synuclein (panel D) expressed and purified using this strategy, and further purified using ion exchange chromatography (see Materials and methods).

Figure S1. Instructions for cloning of a target gene into vector pJCX4. (A) Schematic representation of the expression cassette of vector pJCX4 for mammalian cell expression, including a restriction enzyme map of the multicloning site for the insertion of a target gene (see also Figure 1A). (B) Two examples of forward (5') and reverse (3') primer design to amplify a target gene for insertion into vector pJCX4, showing the general strategy (above) and a specific example for Tpm3.1 (below) (C) SDS-PAGE characterization and Coomassie Brilliant Blue R-250 (BIO-RAD) staining of the purification steps for Tpm3.1.

after self-cleavage of the intein (*Figure 1A, Figure S1A and B*). Growth and transfection of
Expi293F cells is carried out according to the manufacturer's protocol, with minor changes (see
Materials and methods). After harvesting and lysing cells, the fusion precursor protein is loaded
onto a chitin affinity resin. The high affinity of the CBD-chitin interaction allows for extensive
washing, and the resin can be analyzed during washing by SDS-PAGE to monitor the removal
of contaminants (*Figure 1B, Figure S1C*). The intein is then activated with the addition of a
reducing agent which allows the target protein to be released from the column.
161
We used this method here to express and purify the six most studied Tpm isoforms and α-
synuclein (*Figure 1C and D*). As an example, after extensive washing on the chitin resin, the
fusion protein Tpm3.1-intein-CBD is nearly contaminant free (step 1 in *Figure 1B*). Self-cleavage
of the intein, induced with β-mercaptoethanol for 24 h (step 2), is followed by elution of the
target protein (step 3). Although the proteins were mostly pure after the intein-affinity purification
step, most proteins here were additionally purified by ion exchange chromatography. Final
yields varied from protein to protein, ranging from 0.25 to 2 mg of purified proteins from 250 mL
of Expi293F culture.
170
171 **Post-translational modification of Expi293F-expressed Tpm and α-synuclein**
172
To evaluate the presence of PTMs and native N- and C-termini in the proteins expressed here
we used two approaches, tandem mass spectrometry (MS)-based proteomics and Pro-Q
Diamond phosphoprotein gel staining (Thermo Fisher Scientific). Fluorescent Pro-Q Diamond
staining is used to directly detect the presence of phosphate groups on tyrosine, serine, or
threonine residues in acrylamide gels. It is important to note that signaling cascades that get
activated during cell lysis can lead to increased dephosphorylation (67-69). Therefore,
phosphatase inhibitors can be used during cell lysis if necessary to preserve native
phosphorylation sites (70,71). We used this approach here in the Pro-Q Diamond staining
analysis (*Figure 2B and D*), but not for PTM proteomics analysis.
182
To ensure full peptide coverage during MS analysis, the expressed proteins were subjected to
digestions using three different enzymes: trypsin, chymotrypsin, and Glu-C. To establish the
identity of each peptide in the observed MS spectra, their masses were matched to the peptides
in a Tpm sequence database, using a false discovery rate (FDR) threshold of less than 1%
(*Table S1*). Full peptide coverage was obtained for all the proteins expressed, including Tpm
isoforms and α-synuclein, with one exception, Tpm4.2, for which coverage was 99% (*Figure 2A*
and C, Figure S2). This analysis revealed several interesting findings. All the proteins, including
α-synuclein, were Nt-acetylated, which as mentioned above is a precondition for optimal Tpm
polymerization along actin filaments (40,41). The three short Tpm isoforms analyzed (Tpm3.1,
Tpm3.2, Tpm4.2) were acetylated on the second residue. In other words, for these three
isoforms the initiator methionine is first removed during N-terminal processing and Nt-
acetylation then takes place on the second residue. Different N-terminal acetyl transferases
(NATs) likely acetylate Tpm isoforms at the initiator methionine and at position two. Based on
the known sequence specificity of the various human NATs (72), we anticipate isoforms
Tpm3.1, Tpm3.2, and Tpm4.2 with N-terminal sequence Met-Ala-Gly have their initiator
methionine removed by MetAP (methionine aminopeptidase) and then become acetylated by
NatA on Ala-2, whereas isoforms Tpm1.6, Tpm1.7, and Tpm2.1 with N-terminal sequence Met-
Asp-Ala are acetylated by NatB on the initiator methionine. α-Synuclein, with N-terminal
sequence Met-Asp-Val is also acetylated on the initiator methionine by NatB (73) (*Figure 2C*).
For all the proteins expressed here, intact and native C-termini peptides were identified (*Figure*
2A and C, Figure S2), confirming that self-cleavage of the intein proceeded as expected, without
the removal of any endogenous amino acids of the target protein or the presence of any extra

Figure 2. Post-translational modification analysis of Expi293F-expressed Tpm3.1 and α-synuclein. (A and C) Mass spectrometry peptide coverage maps of Tpm3.1 and α-synuclein, shown relative to exon and protein domain diagrams, respectively. Green, magenta and blue lines indicate peptides obtained by digestions with trypsin, chymotrypsin, and Glu-C, respectively, and whose masses were identified by mass spectrometry using a false discovery rate (FDR) threshold of < 1% (see also *Table S1* and *Figure S2*). Orange, yellow, red, purple, and green circles indicate Nt-acetylation, Ser/Thr phosphorylation, Tyr phosphorylation, Lys acetylation, and Met oxidation, respectively. The mass spectrometry spectra (MS2) of three representative peptides (highlighted by black dashed lines and gray background) are shown as examples, including N-terminal peptides of Tpm3.1 and α-synuclein, and a central peptide of α-synuclein. These peptides show different types of PTMs (Nt-acetylation, Ser/Thr phosphorylation, and Met oxidation). The MS2 spectra show the b (N-terminal) and y (C-terminal) fragment ions detected by mass spectrometry for each peptide. Sequence diagrams corresponding to the three spectra illustrate all the theoretical fragment ions colored black (observed experimentally) or gray (non-observed). (B and D) SDS-PAGE analysis of purified samples of Tpm3.1 and α-synuclein, using phosphatase inhibitors to reduce sample dephosphorylation during purification, and stained using either Coomassie Brilliant Blue R-250 (BIO-RAD) or Pro-Q Diamond Phosphoprotein Gel Stain (Thermo Fisher Scientific).

Table 1. PTMs identified in the human cell-expressed proteins (FDR < 1%)

Protein	Nt-acetylation	Ser/Thr phosphorylation	Tyr phosphorylation	Lys acetylation
Tpm1.6	Met-1	Ser-87, Ser-123, Ser-174, Thr-199, Ser-252		Lys-112, Lys-189
Tpm1.7	Met-1	Ser-87, Ser-174, Ser-245, Ser-252	Tyr-221	Lys-48
Tpm2.1	Met-1	Ser-87		Lys-48, Lys-77
Tpm3.1	Ala-2	Thr-5, Thr-6, Ser-51, Thr-234,	Tyr-126	Lys-13, Lys-104
Tpm3.2	Ala-2	Ser-51, Ser-155, Thr-216		Lys-169, Lys-190
Tpm4.2	Ala-2	Ser-6		Lys-169
α-synuclein	Met-1	Thr-33, Ser-42, Ser-87		Lys-12, Lys-60

Figure S2. Post-translational modification analysis of Expi293F-expressed Tpm isoforms. Mass spectrometry peptide coverage maps of Tpm isoforms 1.6, 1.7, 2.1, 3.2, and 4.2, shown relative to the exon diagram of each isoform. Green, magenta and blue lines indicate peptides obtained by digestions with trypsin, chymotrypsin, and Glu-C, respectively, and whose masses were identified by mass spectrometry using a false discovery rate (FDR) threshold of less than 1% (see also Figure 2, Table S1, and Materials and methods). Orange, yellow, red, and purple circles indicate Nt-acetylation, Ser/Thr phosphorylation, Tyr phosphorylation, and Lys acetylation, respectively.

amino acids from the tag. Numerous phosphorylation sites, including serine/threonine and
tyrosine phosphorylation, were identified in the expressed proteins with an FDR < 1% (*Table 1,*
Figure 2, Figure S2). The identity of these phosphorylation sites is mostly consistent with
previous phosphoproteomics studies curated by PhosphoSitePlus (74). Other forms of PTMs
were also identified using the stringent FDR cutoff of < 1%, including most notably several sites
of lysine acetylation (*Table 1, Figure 2, Figure S2, Table S1*). Methionine oxidation was also
observed, including for instance α-synuclein Met-5 (*Figure 2C, Table S1*), however this is not a
native modification but rather one that occurs during sample preparation for mass spectrometry
(75). Table S1 contains a complete list of all the peptides and PTMs observed.
214
215 **Comparison of the activities of Expi293F- vs. *E. coli*-expressed Tpm isoforms**
216
The defining biochemical activity of Tpm is to decorate actin filaments, which by combinatorial
association of different isoforms of both actin and Tpm can give rise to a large number of Tpm-
actin copolymers with distinct properties and activities in cells (25). Accordingly, we used here
actin cosedimentation assays to compare the actin-binding affinities of a representative group of
three Expi293F-expressed Tpm isoforms, corresponding to three different genes and
comprising both short (Tpm3.1) and long (Tpm1.6 and Tpm2.1) isoforms with differently
processed N-termini (see Materials and methods). Quantification of multiple such experiments,
using SDS-PAGE and densitometric analysis, and fitting of the data to a Hill equation revealed
the parameters of the interactions of Tpm isoforms with F-actin, including the apparent
dissociation constant (K_{app}) and Hill coefficient (a measure of the cooperativity of the interaction)
(*Figure 3A, B and E, Figure S3*). This analysis confirmed binding of all three of the Tpm
isoforms to F-actin, but with significantly higher affinity for Tpm1.6 (K_{app} = 0.32 ± 0.06 μM) than
Tpm2.1 or Tpm3.1 (K_{app} = 1.78 ± 0.10 μM and 1.72 ± 0.09 μM, respectively). Interestingly, the
latter two isoforms had very similar affinities for F-actin, despite one being one pseudo-repeat
longer than the other and having differently processed N-termini.
232
We then compared the two most studied non-muscle isoforms (Tpm1.6 and Tpm3.1) with their
E. coli-expressed counterparts, both with and without the addition at the N-terminus of three
amino acids (Met-Ala-Ser) that have been widely used to mimic acetylation (41,76). This
analysis revealed at least three interesting results. First, for the long isoform Tpm1.6, the
Expi293F- and *E. coli*-expressed proteins had very similar affinities for F-actin when Met-Ala-
Ser was added at the N-terminus (*Figure 3C and E*). However, the affinity of *E. coli*-expressed
Tpm1.6 dropped ~15-fold when Met-Ala-Ser was not added. This result suggests that for
Tpm1.6 the addition of Met-Ala-Ser adequately mimics N-terminal acetylation for binding to F-
actin, although it remains to be shown whether this modification can fully account for other
regulatory activities of Tpm1.6 in controlling the interactions of actin-binding proteins (ABPs)
with the actin filament. Second, for the short isoform Tpm3.1 the two *E. coli*-expressed proteins,
with and without Met-Ala-Ser at the N-terminus, had nearly identical affinities for F-actin, but
lower than the affinity of Expi293F-expressed Tpm3.1 (*Figure 3D and E*). Therefore, for Tpm3.1
the addition of extra amino acids at the N-terminus cannot substitute for native Nt-acetylation,
indicating that native PTMs added by Expi293F cells are necessary for function. We suspect
this behavior may affect other Tpm isoforms, which is an important consideration since many
studies have used extra N-terminal amino acids to substitute for Nt-acetylation irrespective of
isoform. Third, it is known that most Tpm isoforms bind cooperatively to F-actin (77-79). Here,
the isoforms that bound F-actin with lower affinity displayed higher cooperativity, and this was
true among different isoforms as well as among variants of the same isoform expressed in
different cell types. Thus, Tpm2.1 and Tpm3.1 exhibited higher cooperativity than Tpm1.6,
whereas the *E. coli*-expressed Tpm isoforms that bound F-actin with lower affinity also showed
higher cooperativity (*Figure 3E*).

Figure 3. Binding of Tpm isoforms to F-actin using cosedimentation analysis. (A) Representative SDS-PAGE cosedimentation analysis of Expi293F-expressed Tpm3.1 (0.75-10 µM) with F-actin (7 µM). Controls include actin alone, found mostly in the pellet fraction (P), and Tpm3.1 alone, found exclusively in the supernatant fraction (S). (B) Between three and four cosedimentation experiments were performed for each of the three Tpm isoforms analyzed (Tpm1.6, Tpm2.1, Tpm3.1; color-coded), the density of the bands in the gels were quantified and data were plotted as the fraction of maximal binding *vs.* the free concentration of each Tpm isoform (see *Figure S3* and Materials and methods). The data were fit to a Hill equation to obtain the dissociation constant (K_{app}) and Hill coefficient of each interaction. (C and D) experiments similar to those shown in panels *A* and *B* were performed to compare the F-actin-binding affinities of Expi293F-expressed Tpm1.6 and Tpm3.1 with their *E. coli*-expressed counterparts, with and without the N-terminal addition of Met-Ala-Ser (MAS), a modification thought to mimic acetylation {Gateva, 2017 #12568;Palm, 2003 #12954}. (E) Parameters of the fits shown in panels B to D, including K_{app} and Hill coefficient.

Figure S3. Cosedimentation of Tpm isoforms with F-actin. Cosedimentation analysis of the interactions of Tpm isoforms with F-actin using 12% SDS-PAGE gels, showing Expi293F-expressed Tpm1.6, Tpm2.1, and Tpm3.1 and *E. coli*-expressed Tpm1.6 and Tpm3.1 with and without the addition of Met-Ala-Ser (MAS) at the N-terminus to mimic Nt-acetylation. The actin concentration in all the experiments was 7 µM, whereas the concentration of Tpm varied for each experiment and is shown on the top of the gels. Controls include actin alone, found mostly in the pellet fraction (P), and Tpm alone, found exclusively in the supernatant fraction (S) for all the proteins analyzed. This data was used to produce the plots and data fits shown in Figure 3.

256
257 **Non-muscle Tpm isoforms can form heterodimers**

258
259 The actin filaments of the contractile apparatus of muscle cells are decorated with two or more
260 Tpm isoforms, which often associate as heterodimers (80-83). For instance, in skeletal muscle
261 Tpm1.1/Tpm2.2 (also known as isoforms α and β) heterodimers form more favorably than
262 Tpm1.1/Tpm1.1 homodimers (81). It is unknown whether non-muscle Tpm isoforms can also
263 form heterodimers in cells.

264
To explore this question, we modified the mammalian cell expression system described here to
coexpress pairs of Tpm isoforms. We designed a new mammalian cell expression vector,
containing an internal ribosomal entry site (IRES) inserted in between two multi-cloning sites
(MCS), one for each Tpm isoform being coexpressed (*Figure 4A*). The use of the coexpression
IRES vector, instead of cotransfection with two separate vectors, ensures that similar amounts
of the two Tpm isoforms are expressed together in time and space. Adjacent to MCS1 and
MCS2 the vector contains a V5 and a FLAG affinity tag, respectively. In this way, the
coexpressed Tpm isoforms have native N-termini, but either V5 or FLAG affinity tags added at
the C-terminus. These tags are used for tandem affinity purification and detection via Western
blot analysis (*Figure 4B*), a method we previously used to study wild-type/mutant heterodimers
of IRSp53 (84). The presence of these tags likely disrupts the head-to-tail interaction of Tpm
coiled coils along F-actin, but because coiled coil formation does not depend on Tpm's ability to
bind F-actin, these tags were not removed for the experiments described here. When two Tpm
isoforms are coexpressed using this strategy, they can form either two independent
homodimers, heterodimers or a combination of homodimers and heterodimers. To discriminate
among these possibilities, we purify the mixture on a FLAG resin, wash extensively, and the
eluted sample is then analyzed by anti-FLAG and anti-V5 Western blotting (*Figure 4B*). The
presence of an anti-V5 signal in the eluted sample (but not in the last wash) is a strong
indication of heterodimer formation. This procedure is illustrated here for the coexpressed
isoform pair Tpm3.1-V5/Tpm4.2-FLAG (*Figure 4C*). Note that the anti-FLAG and anti-V5 signals
disappear almost entirely in the last wash, but both signals show up strongly in the eluted
sample, indicating that any Tpm3.1-V5 that was retained in the FLAG resin was forming
heterodimers with Tpm4.2-FLAG.

288
Because heterodimerization of long muscle isoforms has been demonstrated (80,81,85), the
analysis here focused on four short, non-muscle Tpm isoforms: Tpm3.1, Tpm3.2, Tpm3.5, and
Tpm4.2. Note that Tpm isoforms of different lengths are unlikely to heterodimerize since this
would lead to parts of the hydrophobic core of the coiled coil being exposed to the solvent. We
co-expressed the following pairs: Tpm3.1-V5/Tpm3.2-FLAG, Tpm3.1-V5/Tpm4.2-FLAG, and
Tpm3.5-V5/Tpm4.2-FLAG, which all formed heterodimers (*Figure 4D*). To control for potential
differences in the expression levels of the proteins cloned into MCS1 and MCS2, we switched
the order of isoforms and coexpressed the pair Tpm3.2-V5/Tpm3.1-FLAG, which also formed a
heterodimer. Another control showed that V5- and FLAG-tagged isoforms Tpm3.1 and Tpm3.2
form homodimers despite the different tags. Finally, Tpm3.1-FLAG cloned alone into MCS2
(MCS1 empty) also eluted as a homodimer and no anti-V5 signal was detected in the eluted
sample (*Figure 4D*).

301
302 These results strongly suggest that Tpm3.1, Tpm3.2, Tpm3.5, and Tpm4.2, and likely other non-
303 muscle isoforms can form heterodimers as well as homodimers in cells when coexpressed
304 together. We then asked whether they would preferentially reassociate as homodimers or
305 heterodimers after boiling *in vitro*, which unfolds the coiled coil, as shown for muscle isoforms
306 (86-88). For this, we modified the procedure described above introducing two additional steps;

Figure 4. Heterodimerization of non-muscle Tpm isoforms. (A) Schematic representation of the MCS1-IRES-MCS2 elements of vector pJC8 used for coexpression of pairs of Tpm isoforms, including a depiction of the restriction sites and affinity tags (FLAG and V5) associated with each MCS. (B) Illustration of the potential association of coexpressed Tpm pairs as homodimers or heterodimers and strategy for the FLAG tag-based purification and Western blot-based detection of such species using anti-FLAG and anti-V5 antibodies. (C) Illustration of this strategy for the coexpressed pair of Tpm3.1-V5 and Tpm4.2-FLAG, showing the presence of both isoforms, i.e. heterodimers reactive to both anti-FLAG and anti-V5 antibodies, in the elution sample after extensive washing on the FLAG resin. (D) Western blot analysis of the FLAG elution samples from seven such experiments, including six Tpm pairs and a control experiment with Tpm3.1-FLAG cloned into MCS2 and no Tpm isoform cloned into MCS1. (E) Experimental design for the detection of Tpm heterodimers after unfolding and refolding. Following the FLAG elution, samples are boiled and cooled and then purified on a V5 resin. (F) Illustration of this strategy for the coexpressed pair of Tpm3.1-V5 and Tpm3.2-FLAG. After extensive washing, the Western blot detection of anti-FLAG and anti-V5 signals in the sample from the V5 resin demonstrates that Tpm3.1/Tpm3.2 heterodimers formed spontaneously during refolding.

after purification on the FLAG resin, the eluted samples were boiled, then allowed to slowly
refold during cooling, repurified on a V5 resin, and then analyzed by anti-FLAG and anti-V5
Western blotting (*Figure 4E*). We analyzed in this manner the pair Tpm3.1-V5/Tpm3.2-FLAG
and monitored each step of this procedure by Western blotting (*Figure 4F*). The initial steps of
the procedure (i.e. purification on the FLAG resin) confirmed the formation of heterodimers, as
described above (*Figure 4D*). After boiling/cooling the sample was loaded onto the V5 resin.
The flowthrough from the V5 resin reacted only with the anti-FLAG antibody, indicating that
some of the heterodimers had reassociated as homodimers. Yet, the V5 elution showed strong
anti-FLAG and anti-V5 signals. Any Tpm3.2-FLAG retained in the V5 resin was likely forming
heterodimers with Tpm3.1-V5. Together, the data presented here show that non-muscle Tpm
isoforms can form both homodimer and heterodimers in cells and *in vitro*.
318
319 **Discussion**
320
321 **Main features and general applicability of the expression-purification method**
322
We have developed a novel method for protein expression and purification that integrates the
advantages of mammalian cell expression with those of an intein-CBD purification tag. The
mammalian cells used, Expi293F, grow to high density in suspension and are adapted for high-
yield protein expression. More importantly, these cells naturally contain the necessary folding
and PTM machineries to produce fully native human proteins. The intein-CBD tag, on the other
hand, provides three key advantages. First, it is a high-affinity tag that allows for extensive on-
column washing, such that high protein purity can be achieved in a single purification step.
Second, this is a self-cleavable tag, which can be activated on-column, bypassing the necessity
for enzymatic cleavage after purification and additional purification associated with other tags
that can in some cases produce non-specific cleavage and introduces two new contaminants,
i.e. the cleavage enzyme and the released affinity tag. Third, through proper design of the
cloning primers (as shown in *Figure S1*) self-cleavage of the intein can take place precisely after
the last C-terminal amino acid of the expressed protein, without the deletion of endogenous
amino acids or the presence of extra amino acids from the tag. One potential drawback of this
system is that a certain amount of premature cleavage may take place in the reducing
environment of the cytoplasm of Expi293F cells, which ultimately reduces protein yields. Yet, for
all the proteins expressed here yields of ~1 mg of highly pure proteins were obtained from
relatively small Expi293F cultures of 250 mL. Such protein amounts are clearly sufficient for
most biochemical and structural studies and the technology is cost permissive for most
laboratories possessing a mammalian cell incubator. The combined benefits and relative
simplicity of this method make it easily adaptable for the expression of most human proteins,
and particularly those whose activities depend upon key PTMs. Thus, while the initial motivation
for this work was the expression of several cytoskeletal proteins of interest to our laboratory,
such as Tpm, we extended it here to the expression of α-synuclein, a protein of intensive
interest and a notable target of many PTMs (52,53). To facilitate the adoption of this system, we
are depositing the two vectors developed here with the Addgene plasmid repository
(www.addgene.org).
350
351 **PTMs of human cell-expressed non-muscle Tpm isoforms**
352
The use of the human cell-expression system described here allowed us to make several
discoveries. One of these discoveries concerns the presence of numerous forms of PTMs,
including Nt-acetylation, Ser/Thr and Tyr phosphorylation and Lys acetylation in non-muscle
Tpm isoforms (*Figure 2A*, *Figure S2*, *Table 1*). We have limited the list of PTMs reported here to
sites detected with very high fidelity (FDR < 1%), but many other sites were detected with

relatively high confidence (FDR < 5%) and could possibly play functional roles (*Table S1*). The
importance of Ser/Thr phosphorylation in muscle Tpm isoforms has been recognized since the
seventies (89,90), and a more recent study suggests that phosphorylation of Ser-283 near the
C-terminus of striated muscle Tpm (Tpm1.1) enhances the stiffness of the head-to-tail
interaction (43) . Virtually nothing, however, is known about the role of Ser/Thr phosphorylation
in non-muscle Tpm isoforms. Curiously, among the phosphorylation sites observed here,
several occur near the N- and/or C-terminal ends of Tpm3.1, Tpm1.6, Tpm1.7, Tpm3.2, and
Tpm4.2, and could possibly have a similar role as in muscle Tpm1.1, by modulating the stiffness
of the head-to-tail interaction. To our knowledge, the role Tyr phosphorylation in Tpm isoforms
remains unexplored, and our finding here of Tyr phosphorylation sites in Tpm3.1 and Tpm1.7
opens an interesting new line of inquire to understand the role of this modification. Lys
acetylation of Tpm isoforms is also virtually unexplored. One of the Lys acetylation sites
detected here occurs near the N-terminus of Tpm3.1 and could also modulate the
polymerizability of this isoform along actin filaments, as *in vitro* studies have suggested for
muscle Tpm (91). A recent study also finds that Lys acetylation of F-actin decreases Tpm-
dependent inhibition of actomyosin activity (92). Lys acetylation of non-muscle Tpm isoforms
could have a similar effect, by modulating the interactions of ABPs. The presence and extent of
Nt-acetylation among non-muscle Tpm isoforms is to this day questioned (93). Since ~90% of
human proteins are Nt-acetylated (58) and Nt-acetylation plays a crucial role in the
polymerization of muscle Tpm isoforms (41), it is not surprising that we found here that all the
non-muscle isoforms analyzed were Nt-acetylated. What is more, we did not detect a single N-
terminal peptide that was not acetylated, suggesting that N-terminal acetylation is a constitutive
modification for all Tpm isoforms. More surprising, but consistent with the known sequence
specificity of human NATs (72), was the observation that the short Tpm isoforms (Tpm3.1,
Tpm3.2, Tpm4.2) undergo Nt-acetylation on the second residue, after removal of the initiator
Met by MetAP, which the PTM machinery of Expi293F cells was also able to achieve for these
overexpressed proteins. These three isoforms are likely Nt-acetylated by NatA, whereas the
long isoforms (Tpm1.6, Tpm1.7, Tpm2.1) are likely Nt-acetylated by NatB on the initiator Met
(72).
387
388 **Different activities of Tpm isoforms expressed in human and *E. coli* cells**
389
The vast majority of *in vitro* studies on Tpm isoforms have used *E. coli*-expressed proteins. For
years, these studies have used the addition of extra amino acids at the N-terminus of Tpm as a
way to mimic Nt-acetylation (40,41,48,49) and have assumed that this modification fully
replicates the natural capacity of Tpm to polymerize along actin filaments. By comparing for the
first time side-by-side the F-actin-binding activity of naturally acetylated and *E. coli*-expressed
Tpm isoforms ± Nt-acetylation-mimics we have shown here that this assumption is not always
true. Thus, while the N-terminal addition of Met-Ala-Ser mostly restores the F-actin-binding
affinity of *E. coli*-expressed Tpm1.6, it fails to do the same for Tpm3.1 (*Figure 3*). Tpm3.1 is a
short isoform, whereas Tpm1.6 is a long isoform and interacts with one more actin protomer
along F-actin. While this difference could possibly explain their different behaviors in these
experiments, it appears prudent to test this effect individually for each Tpm isoform, particularly
that this work provides a solution to the problem of Nt-acetylation. Indeed, Tpm2.1 is also a long
isoform, but we found its affinity for F-actin to be very similar to that of Tpm3.1 (*Figure 3A*). We
also noticed an interesting correlation between affinity and cooperativity; lower Tpm binding
affinity for F-actin correlates with higher cooperativity of the interaction, both among naturally
occurring isoforms or within a single isoform with different head-to-tail interaction sequences
(*Figure 3E*). Although we could not find a previous reference to this correlation, it is intuitively
clear that as the affinity of single Tpm dimers for F-actin becomes lower, binding to F-actin
would become increasingly dependent on the pre-polymerization of Tpm into mini-filaments of

ever-growing length through head-to-tail interaction. This is also why a strong head-to-tail
interaction, which Nt-acetylation favors, enhances binding to F-actin.
411
412 **Non-muscle Tpm isoforms can form heterodimers *in vitro* and in cells**
413
It has long been known that muscle Tpm isoforms function as a mixture of homodimers and
heterodimers (81). Similarly, we have shown here that non-muscle Tpm isoforms purify from
cells as both homodimers and heterodimers, and that their ability to heterodimerize is retained
after boiling and cooling *in vitro,* with no apparent preference for one form of association vs.
another. This is an important finding, since Tpm heterodimerization can further expand the
range and functional diversity of Tpm-actin copolymers with segregated localization and function
in cells. The prevalence of heterodimers in cells could be dictated by the time, location and
amount of expression of one isoform vs. another.
422
423 In summary, we have developed a novel method for the expression of fully native proteins in
424 human cells, we have demonstrated its applicability to two types of proteins of general interest,
425 we have used this method to make new discoveries about non-muscle Tpm that had so far
426 resisted investigation, and we have provided the scientific community access to our newly-
427 developed vectors to facilitate the adoption of this method.
428
429
430 **Materials and methods**
431
432 **Human cell expression vectors**
433
Vector pJCX4: The QuickChange site-directed mutagenesis kit (Agilent Technologies, Santa
Clara, CA) was used to remove two SapI restriction sites from vector pEGFP-C1 (Takara Bio
USA, Mountain View, CA). A Woodchuck hepatitis virus (WHP) posttranscriptional regulatory
element (WPRE) was added between the XbaI and MfeI sites of the multi-cloning site (MCS).
The MCS-intein-chitin binding domain of vector pTXB1 (New England Biolabs, Ipswitch, MA)
was then inserted between sites NheI and Kpn2I, and the MCS of vector pEGFP-C1 was
removed by annealed oligos which were ligated between sites Kpn2I and XbaI. The vector is
depicted in figure 1*A*.
442
Vector pJC8: The intein-CBD cassette was removed from vector pJCX4 and replaced with an
internal ribosome entry site (IRES) element between sites BamHI and BspE1. Two new
multicloning sites, one containing a V5- and the other a FLAG-affinity tag, were introduced on
either side of the IRES using annealing oligos. The V5 and FLAG epitope tags provide clear
signals by Western blot analysis and are used here to recognize the expressed Tpm isoforms.
The MCSs and IRES region of the vector are depicted in figure 4*A*.
449
450 **Cloning of Tpm isoforms and α-synuclein**
451
The following cDNAs were used in cloning (NCBI codes): Tpm1.6 (NM_001018004), Tpm1.7
(NM_001018006), Tpm2.1 (NM_213674), Tpm3.1 (NM_153649), Tpm3.2 (NM_001043351),
Tpm4.2 (NM_003290), α-synuclein (NM_000345.4). The cDNAs were cloned into vector pJCX4
by PCR-amplification with primers that included the NotI (5') and SapI (3') restriction enzyme
sites (*Figure 1, Figure S1*). The genes encoding isoforms Tpm1.6 and Tpm3.1 with added N-
terminal Met-Ala-Ser in vector pBAT4 were a gift from Dr. Pekka Lappalainen. The two isoforms
were also cloned without N-terminal extensions into vector pBAT4 by PCR-amplification with
primers that included the NcoI (5') and either BamHI or XhoI (3') restriction enzyme sites. Four

of the Tpm isoforms (Tpm3.1, Tpm3.2, Tpm3.5, Tpm4.2) were separately cloned as pairs into
the IRES-containing vector pJC8 (see above) using PCR primers which included XhoI and
BspEI (MCS1) or NotI and NheI (MCS2) on their overhangs. For this, the cDNA encoding
isoform Tpm3.5 was generated starting from Tpm3.1 (amino acids 1-222) and the nucleotides
encoding for the C-terminal 26 amino acids that differ among these two isoforms were added by
overlap extension PCR. All the primers used in this study are listed in Table S2.
466
467   **Protein expression in Expi293F cells and intein-based purification**
468
Expression was carried out with minor changes to the manufacture's protocol for the Expi293
Expression System Kit (Thermo Fisher Scientific, Waltham, MA). Expi293F cells were grown in
Expi293 media to a density of 2.5×10^6 cells/mL. Cultures were incubated on an orbital shaker
(125 rpm) at 37°C using an Isotemp incubator (Thermo Fisher Scientific, Waltham, MA) with a
humidified atmosphere of 8% CO_2. Pure plasmid DNA (1 µg per mL of Expi293F culture) and
polyethylenimine (PEI, 3 µL per mL of Expi293F culture) were added to separate 5 mL aliquots
of 37°C Opti-MEM (Thermo Fisher Scientific), mixed after 5 min, and incubated for 30 min at
25°C and then added to the Expi293F cell culture. Cultures were grown for 48-72 h, optimized
for each protein. Cells were harvested by centrifugation (4,000 x g) at 4°C for 10 min.
478
For purification, cell pellets were resuspended in 20-35 mL of lysis buffer (20 mM HEPES pH
7.5, 500 mM NaCl, 1 mM DTT). Cells were lysed using a Dounce glass piston homogenizer
(Kontes Glass, Vineland, NJ) and the supernatants were clarified by centrifugation at 48,000 x g
at 4°C for 20 min. The clarified supernatants were loaded onto 15 mL chitin resin (packed into a
column) pre-equilibrated with lysis buffer and washed extensively. Self-cleavage of the intein
was activated with the addition of 50 mM β-mercaptoethanol at 25°C for 24 h. The expressed
proteins were then eluted with lysis buffer, dialyzed overnight into A-buffer (20 mM HEPES pH
7.5, 50 mM NaCl) and further purified on a Source Q column (MilliporeSigma, St. Louis, MO)
using a 50-300 mM NaCl gradient. Peak fractions were dialyzed into storage buffer (20 mM
HEPES pH 7.5, 50 mM NaCl), flash frozen in liquid nitrogen and stored at -80°C.
489
490   *E. coli* **expression and purification of Tpm isoforms**
491
Tpm isoforms Tpm1.6 and Tpm3.1 (with and without Met-Ala-Ser at the N-terminus) were
expressed in ArcticExpress(DE3) RIL cells (Agilent Technologies), grown in Terrific Broth (TB)
medium for 6 h at 37°C to an optical density of ~1.5-2 at 600 nm (OD_{600}), followed by 24 h at
10°C in with 0.4 mM isopropyl-β-D-thiogalactoside (IPTG). Cells were harvested by
centrifugation (4,000 x g) and resuspended in 20 mM HEPES (pH 7.5), 100 mM NaCl, 1 mM
EDTA, and 1 mM phenylmethanesulfonyl fluoride (PMSF). Cells were boiled in a water bath for
10 min, cooled on ice for 10 min, and clarified by centrifugation at 48,000 x g for 20 min at 4°C.
Sodium acetate (1 M, pH 4.5) was added to the clarified supernatant containing the Tpm
isoforms (that are heat-stable) until the solution reached the isoelectric point for precipitation of
each isoform (4.5-4.7). The solutions were then centrifuged at 48,000 x g to pellet the Tpm
isoforms. Pellets were dissolved in A-buffer (20 mM HEPES pH 7.5, 100 mM NaCl) and
dialyzed in the same buffer overnight. Proteins were loaded on Source Q ion exchange
(MilliporeSigma) and eluted with a 100-500 mM NaCl gradient. Peak fractions containing pure
proteins were dialyzed in storage buffer (20 mM HEPES pH 7.5, 50 mM NaCl), flash frozen in
liquid nitrogen and stored at -80°C.
507
508   **Preparation of samples for proteomics analysis**
509

Gel bands of the purified proteins were cut and destained with 100 mM ammonium bicarbonate/acetonitrile (50:50). The bands were reduced in 10 mM dithiothreitol and 100 mM ammonium bicarbonate for 60 min at 52°C. The bands were then alkylated with 50 mM iodoacetamide and 100 mM ammonium bicarbonate at 25°C for 1 h in the dark. The proteins were digested with enzymes (trypsin, chymotrypsin, or Glu-C) at 37°C for 12 h. Supernatants were removed and transferred to fresh tubes. Additional peptides were extracted from the gel by adding 50% acetonitrile and 1% TFA and shaking for 10 min. The supernatants were combined and dried. The dried samples were reconstituted in 0.1% formic acid for mass spectrometry analysis.

Proteomics analysis by nano-LC-MS/MS

Peptides were analyzed on a Q-Exactive Orbitrap Mass Spectrometer attached to an Easy-nLC (Thermo Fisher Scientific) at 400 nL/min. Peptides were eluted with a 25 min gradient of 5-32% ACN followed by 5 min 90% ACN and 0.1% formic acid. The data-dependent acquisition mode with a dynamic exclusion of 45 sec was used for analysis. A full MS scan range of 350-1200 m/z was collected, with a resolution of 70 K, maximum injection time of 50 ms, and automatic gain control (AGC) of 1×10^6. A series of MS2 scans were then acquired for the top 12 most abundant ions of the MS1 scans. Ions were filtered with charge 2–4. An isolation window of 2 m/z was used with the quadruple isolation mode. Ions were fragmented using higher-energy collisional dissociation (HCD), with a collision energy of 27%. Orbitrap detection was used with a scan range of 140-2000 m/z, resolution of 30 K, maximum injection time of 54 ms and AGC of 50,000. All the samples were analyzed using a multiplexed parallel reaction monitoring (PRM) method based on a scheduled inclusion list containing the target precursor ions. Full MS scans were acquired on the Orbitrap from 350–1200 m/z at a resolution of 60,000, using an AGC of 50,000. The minimum threshold was set to 100,000 ion counts. Precursor ions were fragmented with the quadrupole using an isolation width of 2 m/z units, a maximum injection time of 50 ms and AGC of 10,000.

Proteomics MS Data Analysis Including Peptide Identification and Quantification

The mass spectrometry raw spectra were processed using Proteome Discoverer version 2.3 (Thermo Fisher Scientific). The spectra were searched against a database of target protein sequences using default setting: precursor mass tolerance of 10 ppm, fragment mass tolerance of 0.02 Da, enzymes specific cleavage sites, and up to two missed cleavages. Cysteine carbamidomethylation was set as a fixed modification, while methionine oxidation and N-terminal or lysine acetylation were set as variable modifications. The search results were filtered using the target-decoy approach with the false discovery rate (FRD) cutoff of < 1% at the peptide and protein levels. To identify PTMs, the raw data were re-processed with the program MetaMorepheus version 0.0.316 optimized for PTM analysis (94). The calibrate, G-PTM-D, and search tasks of the program were used with default parameters, including default fixed and variable modifications, precursor mass tolerance of 5 ppm, enzyme specific cleavage, up to two missed cleavages, and an FRD cutoff of < 1% at the peptide and protein levels.

Cosedimentation assays

Actin filaments at a fixed concentration of 7 µM were incubated for 30 min at 25°C with increasing Tpm concentrations in cosedimentation buffer (20 mM HEPES pH 7.5, 200 mM NaCl, 5 mM MgCl$_2$) and a total volume of 100 µL. The reactions were centrifuged for 30 min at 80,000 rpm (280,000 rcf) to pellet F-actin and any bound Tpm (*Figure 3A, Figure S3*). For each Tpm protein, the cosedimentation experiments were repeated three or four times (*Figure S3*).

After centrifugation, 100 µL of the supernatants were mixed with 25 µL of 4x Laemmli SDS-
PAGE loading buffer (S fraction). Pellets were gently washed with 100 µL of cosedimentation
buffer, to remove any remaining soluble protein, and resuspended in 100 µL of cosedimentation
buffer and mixed with 25 µL of 4x Laemmli SDS-PAGE loading buffer (P fraction). Identical
volumes (7 µL) of the P and S fractions were analyzed on SDS-PAGE gels, Coomassie-stained,
and the bands were densitometrically quantified with the program Image Lab (BIO-RAD). In the
absence of actin filaments, all the Tpm isoforms were found in the S fraction, whereas most of
the actin was found in the P fraction with or without Tpm (*Figure 3A, Figure S3*). In the presence
of actin filaments, Tpm cosedimented with actin to different degrees. The free Tpm
concentration (plotted on the x-axis) was determined by dividing the intensity of the Tpm band in
the S fraction by the sum of the intensities of the Tpm bands in the S and P fractions and
multiplied by the total Tpm concentration. The fraction of Tpm bound in the pellet was
determined by dividing the intensities of the Tpm and actin bands in the pellet fraction.
According to a commonly used convention (95,96), we plotted on the y-axis the fraction of
maximal Tpm binding to F-actin, determined by normalizing the ratio of the intensities of the
Tpm and actin bands in the pellet fraction to the plateau value for each Tpm sample (i.e. the
maximum fraction of Tpm/actin in the pellet fraction). This allows to compare side-by-side Tpm
isoforms with different F-actin-binding affinities by plotting the fraction of maximal binding vs. the
free Tpm concentration and fitting the data to a Hill equation for cooperative binding. We used
the program Prism 9.0.0 (GraphPad) for fitting with a "specific binding with Hill slope" model,
which yielded the apparent dissociation constant (K_{app}) and Hill coefficient for each interaction
(*Figure 3E*).
583
584 **Detection of non-muscle Tpm heterodimers**
585
Expi293F cell cultures (25 mL) expressing pairs of Tpm isoforms using the IRES-containing
vector pJC8 (see above) were resuspended in FLAG lysis buffer (phosphate buffered saline, 1
mM EDTA, 1 mM PMFS, and 0.5% Triton X-100) and lysed by three cycles of freeze-thaw in
liquid nitrogen. Lysates were clarified at 4^{o}C by centrifugation at 48,000 x g for 20 min, and
incubated with 100 µL of anti-DYKDDDDK (FLAG) resin (Thermo Fisher Scientific) for 2 h at
4^{o}C. The resin was washed five times with 200 µL lysis buffer. FLAG-tagged proteins were
released from the resin by incubation with buffer supplemented with 50 mM FLAG peptide
(Genscript, Piscataway, NJ) for 30 min.
594
For boiling and refolding experiments, the proteins purified from the FLAG resin were heated for
10 min at 90^{o}C and then allowed to cool to 25^{o}C over 30 min. The proteins were incubated for 2
h at 4^{o}C with 100 µL of V5 resin, washed five times with 200 µL lysis buffer, and directly used
for analysis. Equal volumes (100 µL) of lysate, flowthrough, 1st wash, last wash, FLAG elution,
and V5 resin-bound proteins were mixed with 25 µL of 4x Laemmli SDS-PAGE loading buffer
and analyzed by Western blotting using primary FLAG (Santa Cruz Biotechnology, Dallas, TX)
and V5 antibodies (Thermo Fisher Scientific). Secondary ECL-linked anti-mouse and anti-rabbit
antibodies (Cytiva Life Sciences) were applied and the blots were imaged using a Pierce ECL
Plus Western Blotting Substrate (Thermo Fisher Scientific) on a G:BOX Chemi XR5 imager
(Syngene, Frederick, MD).
605
606
607 **Acknowledgements**
608

609 This work was supported by National Institutes of Health Grants R01 GM073791 and R01
610 MH087950 to RD and T32 AR053461 to PJC. We thank Pekka Lappalainen for the gift of
611 vectors for the bacterial expression of tropomyosin isoforms. We thank the Quantitative

Proteomics Resource Core at the Perelman School of Medicine (University of Pennsylvania), Hyoungjoo Lee, Joseph Cesare, and Michael Gilbert for help with the proteomics data acquisition and analysis. We thank Michael Shortreed for help with the use of the program MetaMorpheus.

Author contributions

Peter J. Carman: conceptualization, resources, data analysis, validation, investigation, visualization, methodology, project administration, writing and review/editing
Kyle R. Barrie: methodology, review/editing
Roberto Dominguez: conceptualization, supervision, project administration, funding acquisition, writing and review/editing

Author ORCIDs

Peter J. Carman   http://orcid.org/0000-0002-6120-5507
Kyle R. Barrie   https://orcid.org/0000-0003-0860-1639
Roberto Dominguez   https://orcid.org/0000-0003-3186-5229

References

1.   Waugh, D. S. (2011) An overview of enzymatic reagents for the removal of affinity tags. *Protein Expr Purif* **80**, 283-293
2.   Khoury, G. A., Baliban, R. C., and Floudas, C. A. (2011) Proteome-wide post-translational modification statistics: frequency analysis and curation of the swiss-prot database. *Sci Rep* **1**
3.   Bah, A., and Forman-Kay, J. D. (2016) Modulation of Intrinsically Disordered Protein Function by Post-translational Modifications. *J Biol Chem* **291**, 6696-6705
4.   Hunter, T. (2009) Tyrosine phosphorylation: thirty years and counting. *Curr Opin Cell Biol* **21**, 140-146
5.   Johnson, L. N., and Lewis, R. J. (2001) Structural basis for control by phosphorylation. *Chem Rev* **101**, 2209-2242
6.   Walsh, C. T., Garneau-Tsodikova, S., and Gatto, G. J., Jr. (2005) Protein posttranslational modifications: the chemistry of proteome diversifications. *Angew Chem Int Ed Engl* **44**, 7342-7372
7.   Amann, T., Schmieder, V., Faustrup Kildegaard, H., Borth, N., and Andersen, M. R. (2019) Genetic engineering approaches to improve posttranslational modification of biopharmaceuticals in different production platforms. *Biotechnol Bioeng* **116**, 2778-2796
8.   Brown, C. W., Sridhara, V., Boutz, D. R., Person, M. D., Marcotte, E. M., Barrick, J. E., and Wilke, C. O. (2017) Large-scale analysis of post-translational modifications in E. coli under glucose-limiting conditions. *BMC Genomics* **18**, 301
9.   Higel, F., Seidl, A., Sorgel, F., and Friess, W. (2016) N-glycosylation heterogeneity and the influence on structure, function and pharmacokinetics of monoclonal antibodies and Fc fusion proteins. *Eur J Pharm Biopharm* **100**, 94-100
10.   McKenzie, E. A., and Abbott, W. M. (2018) Expression of recombinant proteins in insect and mammalian cells. *Methods* **147**, 40-49
11.   Tang, H., Wang, S., Wang, J., Song, M., Xu, M., Zhang, M., Shen, Y., Hou, J., and Bao, X. (2016) N-hypermannose glycosylation disruption enhances recombinant protein

production by regulating secretory pathway and cell wall integrity in Saccharomyces
cerevisiae. *Sci Rep* **6**, 25654

12. Weis, B. L., Guth, N., Fischer, S., Wissing, S., Fradin, S., Holzmann, K. H., Handrick, R.,
and Otte, K. (2018) Stable miRNA overexpression in human CAP cells: Engineering
alternative production systems for advanced manufacturing of biologics using miR-136
and miR-3074. *Biotechnol Bioeng* **115**, 2027-2038

13. Dunn, A. Y., Melville, M. W., and Frydman, J. (2001) Review: cellular substrates of the
eukaryotic chaperonin TRiC/CCT. *J Struct Biol* **135**, 176-184

14. Liu, Z., Tyo, K. E., Martinez, J. L., Petranovic, D., and Nielsen, J. (2012) Different
expression systems for production of recombinant proteins in Saccharomyces
cerevisiae. *Biotechnol Bioeng* **109**, 1259-1268

15. Resnicow, D. I., Deacon, J. C., Warrick, H. M., Spudich, J. A., and Leinwand, L. A.
(2010) Functional diversity among a family of human skeletal muscle myosin motors.
Proc Natl Acad Sci U S A **107**, 1053-1058

16. Thomas, J. A., and Tate, C. G. (2014) Quality control in eukaryotic membrane protein
overproduction. *J Mol Biol* **426**, 4139-4154

17. Brault, V., Reedy, M. C., Sauder, U., Kammerer, R. A., Aebi, U., and Schoenenberger,
C. (1999) Substitution of flight muscle-specific actin by human (beta)-cytoplasmic actin in
the indirect flight muscle of Drosophila. *J Cell Sci* **112 (Pt 21)**, 3627-3639

18. Sliwinska, M., Skorzewski, R., and Moraczewska, J. (2008) Role of actin C-terminus in
regulation of striated muscle thin filament. *Biophys J* **94**, 1341-1347

19. Chant, A., Kraemer-Pecore, C. M., Watkin, R., and Kneale, G. G. (2005) Attachment of a
histidine tag to the minimal zinc finger protein of the Aspergillus nidulans gene regulatory
protein AreA causes a conformational change at the DNA-binding site. *Protein Expr Purif*
39, 152-159

20. Suh-Lailam, B. B., and Hevel, J. M. (2009) Efficient cleavage of problematic tobacco
etch virus (TEV)-protein arginine methyltransferase constructs. *Anal Biochem* **387**, 130-
132

21. Wu, J., and Filutowicz, M. (1999) Hexahistidine (His6)-tag dependent protein
dimerization: a cautionary tale. *Acta Biochim Pol* **46**, 591-599

22. Gunning, P. W., Hardeman, E. C., Lappalainen, P., and Mulvihill, D. P. (2015)
Tropomyosin - master regulator of actin filament function in the cytoskeleton. *Journal of*
cell science **128**, 2965-2974

23. Geeves, M. A., Hitchcock-DeGregori, S. E., and Gunning, P. W. (2015) A systematic
nomenclature for mammalian tropomyosin isoforms. *J Muscle Res Cell Motil* **36**, 147-
153

24. Meiring, J. C. M., Bryce, N. S., Wang, Y., Taft, M. H., Manstein, D. J., Liu Lau, S., Stear,
J., Hardeman, E. C., and Gunning, P. W. (2018) Co-polymers of Actin and Tropomyosin
Account for a Major Fraction of the Human Actin Cytoskeleton. *Curr Biol* **28**, 2331-2337
e2335

25. Gunning, P. W., Ghoshdastider, U., Whitaker, S., Popp, D., and Robinson, R. C. (2015)
The evolution of compositionally and functionally distinct actin filaments. *J Cell Sci* **128**,
2009-2019

26. Lehman, W., Hatch, V., Korman, V., Rosol, M., Thomas, L., Maytum, R., Geeves, M. A.,
Van Eyk, J. E., Tobacman, L. S., and Craig, R. (2000) Tropomyosin and actin isoforms
modulate the localization of tropomyosin strands on actin filaments. *J Mol Biol* **302**, 593-
606

27. McKillop, D. F., and Geeves, M. A. (1993) Regulation of the interaction between actin
and myosin subfragment 1: evidence for three states of the thin filament. *Biophys J* **65**,
693-701

28. Cheng, C., Nowak, R. B., Amadeo, M. B., Biswas, S. K., Lo, W.-K., and Fowler, V. M.
(2018) Tropomyosin 3.5 protects the F-actin networks required for tissue biomechanical
properties. *Journal of cell science* **131**, jcs.222042-jcs.222042
29. Pleines, I., Woods, J., Chappaz, S., Kew, V., Foad, N., Ballester-Beltrán, J., Aurbach, K.,
Lincetto, C., Lane, R. M., Schevzov, G., Alexander, W. S., Hilton, D. J., Astle, W. J.,
Downes, K., Nurden, P., Westbury, S. K., Mumford, A. D., Obaji, S. G., Collins, P. W.,
Delerue, F., Ittner, L. M., Bryce, N. S., Holliday, M., Lucas, C. A., Hardeman, E. C.,
Ouwehand, W. H., Gunning, P. W., Turro, E., Tijssen, M. R., Kile, B. T., and Kile, B. T.
(2017) Mutations in tropomyosin 4 underlie a rare form of human
macrothrombocytopenia. *The Journal of clinical investigation* **127**, 814-829
30. Kee, A. J., Gunning, P. W., and Hardeman, E. C. (2009) A cytoskeletal tropomyosin can
compromise the structural integrity of skeletal muscle. *Cell Motil Cytoskeleton* **66**, 710-
720
31. Vlahovich, N., Kee, A. J., Van der Poel, C., Kettle, E., Hernandez-Deviez, D., Lucas, C.,
Lynch, G. S., Parton, R. G., Gunning, P. W., and Hardeman, E. C. (2009) Cytoskeletal
tropomyosin Tm5NM1 is required for normal excitation-contraction coupling in skeletal
muscle. *Mol Biol Cell* **20**, 400-409
32. Eppinga, R. D., Li, Y., Lin, J. L., and Lin, J. J. (2006) Tropomyosin and caldesmon
regulate cytokinesis speed and membrane stability during cell division. *Arch Biochem
Biophys* **456**, 161-174
33. Thoms, J. A., Loch, H. M., Bamburg, J. R., Gunning, P. W., and Weinberger, R. P.
(2008) A tropomyosin 1 induced defect in cytokinesis can be rescued by elevated
expression of cofilin. *Cell Motil Cytoskeleton* **65**, 979-990
34. Hook, J., Lemckert, F., Qin, H., Schevzov, G., and Gunning, P. (2004) Gamma
Tropomyosin Gene Products Are Required for Embryonic Development. *Molecular and
Cellular Biology* **24**, 2318-2323
35. Hook, J., Lemckert, F., Schevzov, G., Fath, T., and Gunning, P. (2011) Functional
identity of the gamma tropomyosin gene: Implications for embryonic development,
reproduction and cell viability. *Bioarchitecture* **1**, 49-59
36. Dalby-Payne, J. R., O'Loughlin, E. V., and Gunning, P. (2003) Polarization of specific
tropomyosin isoforms in gastrointestinal epithelial cells and their impact on CFTR at the
apical surface. *Mol Biol Cell* **14**, 4365-4375
37. Dufour, C., Weinberger, R. P., and Gunning, P. (1998) Tropomyosin isoform diversity
and neuronal morphogenesis. *Immunology and Cell Biology* **76**, 424-429
38. Had, L., Faivre-Sarrailh, C., Legrand, C., and Rabie, A. (1993) The expression of
tropomyosin genes in pure cultures of rat neurons, astrocytes and oligodendrocytes is
highly cell-type specific and strongly regulated during development. *Brain Res Mol Brain
Res* **18**, 77-86
39. Hannan, A. J., Gunning, P., Jeffrey, P. L., and Weinberger, R. P. (1998) Structural
Compartments within Neurons: Developmentally Regulated Organization of
Microfilament Isoform mRNA and Protein. *Molecular and Cellular Neuroscience* **11**, 289-
304
40. Heald, R. W., and Hitchcock-DeGregori, S. E. (1988) The structure of the amino
terminus of tropomyosin is critical for binding to actin in the absence and presence of
troponin. *J Biol Chem* **263**, 5254-5259
41. Palm, T., Greenfield, N. J., and Hitchcock-DeGregori, S. E. (2003) Tropomyosin ends
determine the stability and functionality of overlap and troponin T complexes. *Biophys J*
84, 3181-3189
42. Maytum, R., Geeves, M. A., and Konrad, M. (2000) Actomyosin regulatory properties of
yeast tropomyosin are dependent upon N-terminal modification. *Biochemistry* **39**, 11913-
11920

43. Lehman, W., Medlock, G., Li, X. E., Suphamungmee, W., Tu, A. Y., Schmidtmann, A., Ujfalusi, Z., Fischer, S., Moore, J. R., Geeves, M. A., and Regnier, M. (2015) Phosphorylation of Ser283 enhances the stiffness of the tropomyosin head-to-tail overlap domain. *Arch Biochem Biophys* **571**, 10-15

44. Palani, S., Köster, D. V., Hatano, T., Kamnev, A., Kanamaru, T., Brooker, H. R., Hernandez-Fernaud, J. R., Jones, A. M. E., Millar, J. B. A., Mulvihill, D. P., and Balasubramanian, M. K. (2019) Phosphoregulation of tropomyosin is crucial for actin cable turnover and division site placement. *Journal of Cell Biology* **218**, 3548-3559

45. Rajan, S., Jagatheesan, G., Petrashevskaya, N., Biesiadecki, B. J., Warren, C. M., Riddle, T., Liggett, S., Wolska, B. M., Solaro, R. J., and Wieczorek, D. F. (2019) Tropomyosin pseudo-phosphorylation results in dilated cardiomyopathy. *J Biol Chem* **294**, 2913-2923

46. Rao, V. S., Marongelli, E. N., and Guilford, W. H. (2009) Phosphorylation of tropomyosin extends cooperative binding of myosin beyond a single regulatory unit. *Cell Motil Cytoskeleton* **66**, 10-23

47. Brooker, H. R., Geeves, M. A., and Mulvihill, D. P. (2016) Analysis of biophysical and functional consequences of tropomyosin-fluorescent protein fusions. *FEBS Lett* **590**, 3111-3121

48. Frye, J., Klenchin, V. A., and Rayment, I. (2010) Structure of the tropomyosin overlap complex from chicken smooth muscle: insight into the diversity of N-terminal recognition. *Biochemistry* **49**, 4908-4920

49. Monteiro, P. B., Lataro, R. C., Ferro, J. A., and Reinach Fde, C. (1994) Functional alpha-tropomyosin produced in Escherichia coli. A dipeptide extension can substitute the amino-terminal acetyl group. *J Biol Chem* **269**, 10461-10466

50. Johnson, M., Coulton, A. T., Geeves, M. A., and Mulvihill, D. P. (2010) Targeted amino-terminal acetylation of recombinant proteins in E. coli. *PLoS One* **5**, e15801

51. Wingfield, P. T. (2017) N-Terminal Methionine Processing. *Curr Protoc Protein Sci* **88**, 6 14 11-16 14 13

52. Schmid, A. W., Fauvet, B., Moniatte, M., and Lashuel, H. A. (2013) Alpha-synuclein post-translational modifications as potential biomarkers for Parkinson disease and other synucleinopathies. *Mol Cell Proteomics* **12**, 3543-3558

53. Zhang, J., Li, X., and Li, J. D. (2019) The Roles of Post-translational Modifications on alpha-Synuclein in the Pathogenesis of Parkinson's Diseases. *Front Neurosci* **13**, 381

54. Backliwal, G., Hildinger, M., Hasija, V., and Wurm, F. M. (2008) High-density transfection with HEK-293 cells allows doubling of transient titers and removes need for a priori DNA complex formation with PEI. *Biotechnol Bioeng* **99**, 721-727

55. Liu, C., Dalby, B., Chen, W., Kilzer, J. M., and Chiou, H. C. (2008) Transient transfection factors for high-level recombinant protein production in suspension cultured mammalian cells. *Mol Biotechnol* **39**, 141-153

56. Shah, N. H., and Muir, T. W. (2014) Inteins: Nature's Gift to Protein Chemists. *Chem Sci* **5**, 446-461

57. Southworth, M. W., Amaya, K., Evans, T. C., Xu, M. Q., and Perler, F. B. (1999) Purification of proteins fused to either the amino or carboxy terminus of the Mycobacterium xenopi gyrase A intein. *Biotechniques* **27**, 110-114, 116, 118-120

58. Aksnes, H., Drazic, A., Marie, M., and Arnesen, T. (2016) First Things First: Vital Protein Marks by N-Terminal Acetyltransferases. *Trends Biochem Sci* **41**, 746-760

59. Gray, D. (2001) Overview of protein expression by mammalian cells. *Curr Protoc Protein Sci* **Chapter 5**, Unit5 9

60. Khan, K. H. (2013) Gene expression in Mammalian cells and its applications. *Adv Pharm Bull* **3**, 257-263

61. Tokmakov, A. A., Kurotani, A., Takagi, T., Toyama, M., Shirouzu, M., Fukami, Y., and
Yokoyama, S. (2012) Multiple post-translational modifications affect heterologous protein
synthesis. *J Biol Chem* **287**, 27106-27116
62. Wood, D. W., and Camarero, J. A. (2014) Intein applications: from protein purification
and labeling to metabolic control methods. *J Biol Chem* **289**, 14512-14519
63. Telenti, A., Southworth, M., Alcaide, F., Daugelat, S., Jacobs, W. R., Jr., and Perler, F.
B. (1997) The Mycobacterium xenopi GyrA protein splicing element: characterization of
a minimal intein. *J Bacteriol* **179**, 6378-6382
64. Hirata, R., Ohsumk, Y., Nakano, A., Kawasaki, H., Suzuki, K., and Anraku, Y. (1990)
Molecular structure of a gene, VMA1, encoding the catalytic subunit of H(+)-translocating
adenosine triphosphatase from vacuolar membranes of Saccharomyces cerevisiae. *J
Biol Chem* **265**, 6726-6733
65. Kane, P. M., Yamashiro, C. T., Wolczyk, D. F., Neff, N., Goebl, M., and Stevens, T. H.
(1990) Protein splicing converts the yeast TFP1 gene product to the 69-kD subunit of the
vacuolar H(+)-adenosine triphosphatase. *Science* **250**, 651-657
66. Chong, S., Shao, Y., Paulus, H., Benner, J., Perler, F. B., and Xu, M. Q. (1996) Protein
splicing involving the Saccharomyces cerevisiae VMA intein. The steps in the splicing
pathway, side reactions leading to protein cleavage, and establishment of an in vitro
splicing system. *J Biol Chem* **271**, 22159-22168
67. Garcia, A., Cayla, X., Guergnon, J., Dessauge, F., Hospital, V., Rebollo, M. P.,
Fleischer, A., and Rebollo, A. (2003) Serine/threonine protein phosphatases PP1 and
PP2A are key players in apoptosis. *Biochimie* **85**, 721-726
68. Sun, H., and Wang, Y. (2012) Novel Ser/Thr protein phosphatases in cell death
regulation. *Physiology (Bethesda)* **27**, 43-52
69. Van Hoof, C., and Goris, J. (2003) Phosphatases in apoptosis: to be or not to be, PP2A
is in the heart of the question. *Biochim Biophys Acta* **1640**, 97-104
70. Huyer, G., Liu, S., Kelly, J., Moffat, J., Payette, P., Kennedy, B., Tsaprailis, G., Gresser,
M. J., and Ramachandran, C. (1997) Mechanism of inhibition of protein-tyrosine
phosphatases by vanadate and pervanadate. *J Biol Chem* **272**, 843-851
71. Ishihara, H., Martin, B. L., Brautigan, D. L., Karaki, H., Ozaki, H., Kato, Y., Fusetani, N.,
Watabe, S., Hashimoto, K., Uemura, D., and et al. (1989) Calyculin A and okadaic acid:
inhibitors of protein phosphatase activity. *Biochem Biophys Res Commun* **159**, 871-877
72. Drazic, A., Myklebust, L. M., Ree, R., and Arnesen, T. (2016) The world of protein
acetylation. *Biochim Biophys Acta* **1864**, 1372-1401
73. Anderson, J. P., Walker, D. E., Goldstein, J. M., de Laat, R., Banducci, K., Caccavello,
R. J., Barbour, R., Huang, J., Kling, K., Lee, M., Diep, L., Keim, P. S., Shen, X.,
Chataway, T., Schlossmacher, M. G., Seubert, P., Schenk, D., Sinha, S., Gai, W. P., and
Chilcote, T. J. (2006) Phosphorylation of Ser-129 is the dominant pathological
modification of alpha-synuclein in familial and sporadic Lewy body disease. *J Biol Chem*
281, 29739-29752
74. Hornbeck, P. V., Zhang, B., Murray, B., Kornhauser, J. M., Latham, V., and Skrzypek, E.
(2015) PhosphoSitePlus, 2014: mutations, PTMs and recalibrations. *Nucleic Acids Res*
43, D512-520
75. Schey, K. L., and Finley, E. L. (2000) Identification of peptide oxidation by tandem mass
spectrometry. *Acc Chem Res* **33**, 299-306
76. Gateva, G., Kremneva, E., Reindl, T., Kotila, T., Kogan, K., Gressin, L., Gunning, P. W.,
Manstein, D. J., Michelot, A., and Lappalainen, P. (2017) Tropomyosin Isoforms Specify
Functionally Distinct Actin Filament Populations In Vitro. *Curr Biol* **27**, 705-713
77. Tobacman, L. S. (2008) Cooperative binding of tropomyosin to actin. *Adv Exp Med Biol*
644, 85-94

78. Wegner, A. (1979) Equilibrium of the actin-tropomyosin interaction. *J Mol Biol* **131**, 839-
853
79. Yang, Y. Z., Korn, E. D., and Eisenberg, E. (1979) Cooperative binding of tropomyosin to
muscle and Acanthamoeba actin. *J Biol Chem* **254**, 7137-7140
80. Bicer, S., and Reiser, P. J. (2013) Complex tropomyosin and troponin T isoform
expression patterns in orbital and global fibers of adult dog and rat extraocular muscles.
J Muscle Res Cell Motil **34**, 211-231
81. Bronson, D. D., and Schachat, F. H. (1982) Heterogeneity of contractile proteins.
Differences in tropomyosin in fast, mixed, and slow skeletal muscles of the rabbit. *J Biol*
Chem **257**, 3937-3944
82. Sanders, C., Burtnick, L. D., and Smillie, L. B. (1986) Native chicken gizzard
tropomyosin is predominantly a beta gamma-heterodimer. *J Biol Chem* **261**, 12774-
12778
83. Muthuchamy, M., Grupp, I. L., Grupp, G., O'Toole, B. A., Kier, A. B., Boivin, G. P.,
Neumann, J., and Wieczorek, D. F. (1995) Molecular and physiological effects of
overexpressing striated muscle beta-tropomyosin in the adult murine heart. *J Biol Chem*
270, 30593-30603
84. Kast, D. J., and Dominguez, R. (2019) Mechanism of IRSp53 inhibition by 14-3-3. *Nat*
Commun **10**, 483
85. Coulton, A., Lehrer, S. S., and Geeves, M. A. (2006) Functional homodimers and
heterodimers of recombinant smooth muscle tropomyosin. *Biochemistry* **45**, 12853-
12858
86. Jancso, A., and Graceffa, P. (1991) Smooth muscle tropomyosin coiled-coil dimers.
Subunit composition, assembly, and end-to-end interaction. *J Biol Chem* **266**, 5891-5897
87. Graceffa, P. (1992) Heat-treated smooth muscle tropomyosin. *Biochim Biophys Acta*
1120, 205-207
88. Lehrer, S. S., and Qian, Y. (1990) Unfolding/refolding studies of smooth muscle
tropomyosin. Evidence for a chain exchange mechanism in the preferential assembly of
the native heterodimer. *J Biol Chem* **265**, 1134-1138
89. Heeley, D. H. (2013) Phosphorylation of tropomyosin in striated muscle. *J Muscle Res*
Cell Motil **34**, 233-237
90. Ribolow, H., and Barany, M. (1977) Phosphorylation of tropomyosin in live frog muscle.
Arch Biochem Biophys **179**, 718-720
91. Johnson, P., and Smillie, L. B. (1977) Polymerizability of rabbit skeletal tropomyosin:
effects of enzymic and chemical modifications. *Biochemistry* **16**, 2264-2269
92. Schmidt, W., Madan, A., Foster, D. B., and Cammarato, A. (2020) Lysine acetylation of
F-actin decreases tropomyosin-based inhibition of actomyosin activity. *J Biol Chem* **295**,
15527-15539
93. Jansen, S., and Goode, B. L. (2019) Tropomyosin isoforms differentially tune actin
filament length and disassembly. *Mol Biol Cell* **30**, 671-679
94. Solntsev, S. K., Shortreed, M. R., Frey, B. L., and Smith, L. M. (2018) Enhanced Global
Post-translational Modification Discovery with MetaMorpheus. *J Proteome Res* **17**, 1844-
1851
95. Moraczewska, J., Greenfield, N. J., Liu, Y., and Hitchcock-DeGregori, S. E. (2000)
Alteration of tropomyosin function and folding by a nemaline myopathy-causing
mutation. *Biophysical journal* **79**, 3217-3225
96. Rynkiewicz, M. J., Prum, T., Hollenberg, S., Kiani, F. A., Fagnant, P. M., Marston, S. B.,
Trybus, K. M., Fischer, S., Moore, J. R., and Lehman, W. (2017) Tropomyosin Must
Interact Weakly with Actin to Effectively Regulate Thin Filament Function. *Biophys J* **113**,
2444-2451
914

Table S2. Primers used in this study

Primer name	Primer sequence	Purpose
Primers used to construct vector pJCX4		
a3106g_for	gagcccctgatgctcctcgtccagatcatcc	Removal of SapI site from
a3106g_rev	ggatgatctggacgaggagcatcaggggctc	pEGFPC1
a3316g_for	tcgccgccaagctcctcagcaatatcacggg	Removal of SapI site from
a3316g_rev	cccgtgatattgctgaggagcttggcggcga	pEGFPC1
WPRE_for_xbaI	ttctctagataatcaacctctggattacaaaatttgtgaaagattga	PCR amplify WPRE
WPRE_rev_mfeI	ttccaattgatgcggggaggcggc	
MCS_KO_top	ccggaggagacaataccggaat	Removing the pEGFP-C1
MCS_KO_bottom	ctagattccggtattgtctcct	MCS
Primers used to construct vector pJC8		
IRES_for_BamHI	ttcggatccaattcgcccctctccctccc	PCR amplify IRES
IRES_rev_BspEI	ttctccggaggttgtggccatattatcatcgtgttttc	
MCS2_top	tcgagtcttgatcactaactagtaataccggtactctgca	Adding MCS1 into pJC8
MCS2_bottom	gagtaccggtattactagttagtgatcaagac	
MCS2_top	ccggatatcccgggatcagatctatcgagctcattt	Adding MCS2 into pJC8
MCS2_bottom	ctagaaatgagctcgatagatctgatcccgggatat	
Primers used for cloning of Tpm and α-synuclein		
TPM1_for_NotI	ttcgcggccgcgccaccatggacgccatcaagaagaagatg	Tpm1.6, Tpm1.7
TPM1_rev_SapI	ggtggttgctcttccgcacatgttgtttaactccagtaaagtctgatcc	into pJCX4
TPM1_for_NotI	ttcgcggccgcgccaccatggacgccatcaagaagaagatg	Tpm2.1
TPM2_rev_SapI	ggtgctcttccgcacaggttgttgagttccagcagg	into pJCX4
TPM3_for_NotI	ttcgcggccgcgccaccatggctgggatcaccaccatcg	Tpm3.1, Tpm3.2
TPM3_rev_SapI	ggttgctcttccgcacatctcattcaggtcaagcagggtctg	into pJCX4
TPM4_for_NotI	ttcgcggccgcgccaccatggccggcctcaactcc	Tpm4.2
TPM4_rev_SapI	ggttgctcttccgcatatacagttaagttcgtttagtgtctgatccag	into pJCX4
αSyn_for_NotI	ttcgcggccgcatggatgtattcatgaaaggactttcaaag	α-synuclein
αSyn_rev_SapI	ggttgctcttccgcaggcttcaggttcgtagtcttgatac	into pJCX4
1.6_for_NcoI	ttcccatggacgccatcaagaagaagatg	Tpm1.6
1.6_rev_BamHI	ttcggatcctcacatgttgtttaactccagtaaagtctgatccag	into pBAT4
3.1_for_NcoI	ttcccatggctgggatcaccaccatc	Tpm3.1
3.1_rev_XhoI	ttcctcgagtcacatctcattcaggtcaagcagggtctg	into pBAT4
TPM3_for_XhoI	ttcctcgagatggctgggatcaccaccatc	Tpm3.1, Tpm3.2
TPM3_rev_BspEI	ttctccggacatctcattcaggtcaagcaggg	into pJC8 MCS1
TPM3_for_NotI	ttcgcggccgcgccaccatggctgggatcaccaccatcg	Tpm3.1, Tpm3.2
TPM3_rev_NheI	ttcgctagccatctcattcaggtcaagcaggg	into pJC8 MCS2
TPM3_for_XhoI	ttcctcgagatggctgggatcaccaccatc	Tpm3.5
Tpm3.5_rev_BspE1	ttctccggatatagaggtcatgtcattgagggcgtggtccagctcctcgctaatg	into pJC8 MCS1
TPM4_for_NotI	ttcgcggccgcgccaccatggccggcctcaactcc	Tpm4.2
TPM4_rev_NheI	ttcgctagctatacagttaagttcgtttagtgtctgatccag	into pJC8 MCS2
TPM3_for_XhoI	ttcctcgagatggctgggatcaccaccatc	Tpm3.5 overlap extension
Tpm3.5_rev_1	acttcagtttctgggcatagagctcatcttccaggtcatcaattgtcttttccag	PCR to generate isoform
Tpm3.5_rev_2	gtggtccagctcctcgctaatggccttgtacttcagtttctgggcatagagc	
Tpm3.5_rev_BspEI	ttctccggatatagaggtcatgtcattgagggcgtggtccagctcctcgctaatg	