1

Main Manuscript for

Bioinformatic and mechanistic analysis of the palmerolide PKS-NRPS biosynthetic pathway from the microbiome of an Antarctic ascidian

Nicole E. Avalon;[1] Alison E. Murray;[2] Hajnalka E. Daligault;[3] Chien-Chi Lo;[3] Karen W. Davenport;[3] Armand E.K. Dichosa;[3] Patrick S.G. Chain;[3] Bill J. Baker[1]

[1]Department of Chemistry, University of South Florida, Tampa, FL 33620

[2]Division of Earth and Ecosystem Sciences, Desert Research Institute, Reno, NV 89512

[3]Los Alamos National Laboratory, Los Alamos, NM 87545

Corresponding author:

Bill J. Baker, Department of Chemistry, University of South Florida, Tampa, FL 33620, USA. +1-813-974-1967, bjbaker@usf.edu

Author Contributions: This work was the result of a team effort in which the following contributions are recognized: conceptualization, A.E.M., P.S.G.C., and B.J.B.; methodology and experimentation, N.E.A., A.E.M., H.E.D., C.L., P.S.G.C., B.J.B; data analysis, N.E.A., A.E.M., H.E.D, C.L., K.W.D., A.E.K.D., P.S.G.C, B.J.B; data curation, N.E.A., A.E.M., H.E.D, C.L., P.S.G.C, B.J.B; original draft preparation, N.E.A., B.J.B., A.E.M., H.E.D.; review and editing, N.E.A., B.J.B., A.E.M., C.L., H.E.D., P.S.G.C., A.E.K.D; funding acquisition, A.E.M., P.S.G.C., and B.J.B. All authors have read and agreed to the published version of the manuscript.

Competing Interest Statement: The authors have no competing interests to disclose.

Classification: Biological sciences, environmental sciences and ecology

30 **Keywords:** microbial biosynthesis, secondary metabolites, Antarctic microbiology, *trans*-AT Type
31 I polyketide synthase, marine natural products

33 **This PDF file includes:**

34   Main Text.
35   Figures 1 to 4.
36
Abstract

Polyketides are a complex family of natural products that often serve competitive or pro-
survival purposes but can also demonstrate bioactivity in human diseases as, for example
cholesterol lowering agents, anti-infectives, or anti-tumor agents. Marine invertebrates and
microbes are a rich source of polyketides. Palmerolide A, a polyketide isolated from the Antarctic
ascidian *Synoicum adareanum,* is a vacuolar-ATPase inhibitor with potent bioactivity against
melanoma cell lines. The biosynthetic gene clusters (BGC) responsible for production of
secondary metabolites are encoded in the genomes of the producers as discrete genomic
elements. A putative palmerolide BGC was identified from a *S. adareanum* metagenome based
on a high degree of congruence with a chemical structure-based retrobiosynthetic prediction.
Protein family homology analysis, conserved domain searches, and active site and motif
identification were used to identify and propose the function of the 75 kb *trans*-acyltransferase
(AT) polyketide synthase-non-ribosomal synthase (PKS-NRPS) domains responsible for the
synthesis of palmerolide A. Though PKS systems often act in a predictable co-linear sequence,
this BGC includes multiple *trans*-acting enzymatic domains, a non-canonical condensation
termination domain, a bacterial luciferase-like monooxygenase (LLM), and multiple copies found
within the metagenome-assembled genome (MAG) of *Candidatus* Synoicohabitans
palmerolidicus. Detailed inspection of the five highly similar pal BGC copies suggests the
potential for biosynthesis of other members of the palmerolide chemical family. This is the first
delineation of a biosynthetic gene cluster from an Antarctic species. These findings have
relevance for fundamental knowledge of PKS combinatorial biosynthesis and could enhance drug
development efforts of palmerolide A through heterologous gene expression.

Significance Statement

Complex interactions exist between microbiomes and their hosts. Increasingly, defensive
metabolites that have been attributed to host biosynthetic capability are now being recognized as
products of associated microbes. These unique metabolites often have bioactivity in targets of
human disease and can be purposed as pharmaceuticals. The molecular machinery for
production of palmerolide A, a macrolide that is potent and selective against melanoma, was
discovered as a genomic cluster in the microbiome of an Antarctic ascidian. Multiple non-identical
copies of this genomic information provide clues to differences in specific enzymatic domains and
point to Nature's ability to perform combinatorial biosynthesis *in situ*. Harnessing this genetic
information may pave a path for development of a palmerolide-based drug.

69
70 **Introduction**
71
Marine invertebrates such as corals, sponges, mollusks, and ascidians, are known to be
a rich source of bioactive compounds (1). Due to their sessile or sluggish nature, chemical
defenses such as secondary metabolites are often key to their survival. Many compound classes
are represented among benthic invertebrates including terpenes, nonribosomal peptide synthase
(NRPS) products, ribosomally synthesized and post-translationally modified peptides (RiPPs),
and polyketides. It is estimated that that over 11,000 secondary metabolites from marine and

terrestrial environments understood to be of polyketide synthase (PKS) and NRPS origin have
been isolated and described (2). Biosynthetic gene clusters (BGCs) exist as a series of genomic
elements that encode for megaenzymes responsible for production of these secondary
metabolites. BGCs can have distinct nucleotide composition properties such as codon usage and
guanine-cytosine content that do not match the remainder of the genome (3, 4), suggesting a
mechanism of transfer from organisms that are only distantly related, perhaps even from
interkingdom horizontal gene transfer (5, 6). Interestingly, the BGCs for many natural products
isolated from marine invertebrates are found in the host-associated microbiota, reflecting the role
of these compounds in symbiosis (7).
Polyketides are a complex family of natural products produced by a variety of PKS
enzymes that are related to, but evolutionarily divergent from, fatty acid synthases (8). They often
possess long carbon chains with varied degrees of oxidation, can contain aromatic components,
and may be either cyclic or linear. It is estimated that of the polyketides that have been isolated
and characterized, 1% have potential biological activity against human diseases, making this
class of compounds particularly appealing from a drug discovery and development standpoint (9).
This potential for use as pharmaceuticals is approximately five times greater than for compounds
of all other natural product classes (9). Many polyketides are classified as macrolides, which are
large-ring lactones that are pharmaceutically relevant due to a number of biological actions,
including, for example, targeting the cytoskeleton, ribosomal protein biosynthesis, and vacuolar
type V-ATPases (10–13). V-ATPases are responsible for acidification of cells and organelles via
proton transport across membranes, including those of lysosomes, vacuoles, and endosomes.
These enzymes appear to have an impact on angiogenesis, apoptosis, cell proliferation, and
tumor metatastisis (10). A number of marine macrolides inhibit V-ATPases, including
lobatamides, chondropsins, iejimalides, and several of the palmerolides (14–18).
There are three types of PKS systems. Type I PKS systems in bacteria are primarily
comprised of non-iteratively acting multimodular enzymes that lead to progressive elongation of a
polyketide chain, though these megaenzymes can also include "stuttering" modules that may act
iteratively (19–21). In addition, some bacterial Type I PKS systems are comprised solely of
iteratively acting monomodular enzymes that catalyze a series of chain elongation steps for
polyketide formation (22). Type II PKS systems typically contain separate, iteratively acting
enzymes that biosynthesize polycyclic aromatic polyketides, while Type III PKS systems possess
iteratively-acting homodimeric enzymes that often result in monocyclic or bicyclic aromatic
polyketides (19). Type I PKS systems can be subdivided into two groups, depending upon
whether the acyl transferase (AT) modules are encoded within each module at the site that is
parallel to the functional role of the ATs, as in *cis*-AT Type I PKS, or physically distinct from the
megaenzyme, as in *trans*-AT Type I PKS. In both cases, there are often parallel relationships
between the genome order, the action of enzymatic modules, and the functional groups present
in the growing polyketide chain, though in *trans*-AT systems deviations from these parallel
relationships is more likely to be observed (23). In *trans*-AT systems, AT domains may be
incorporated in a mosaic fashion through horizontal gene transfer (23). This introduces greater
molecular architectural diversity over evolutionary time, as one clade of *trans*-ATs may select for
a malonyl-CoA derivative, while the *trans*-AT domains in another clade may select for unusual or
functionalized subunits (24, 25). Additionally, recombination, gene duplication, and conversion
events can lead to further diversification of the resultant biosynthetic machinery (26). Predictions
regarding the intrinsic relationship between a secondary metabolite of interest, the biosynthetic
megaenzyme, and the biosynthetic gene cluster (BGC) can be harnessed for natural product
discovery and development (27–29).
In the search for new and bioactive chemotypes as inspiration for the next generation of
drugs, underexplored ecosystems hold promise as biological and chemical hotspots (30). The
vast Southern Ocean comprises one-tenth of the total area of Earth's oceans and is largely
unstudied for its chemodiversity. The coastal marine environment of Antarctica experiences
seasonal extremes in, for example, ice cover, light field, and food resources. Taken with the
barrier to migration imposed by the Antarctic Circumpolar Current and the effects of repeated

glaciation events on speciation, a rich and endemic biodiversity has evolved, with consequent
potential for new chemodiversity (30–32).
Palmerolide A (**Fig. 1**) is the principal secondary metabolite isolated from *Synoicum*
adareanum, an ascidian which can be found in abundance at depths of 10 to 40 m in the coastal
waters near Palmer Station, Antarctica (17). Palmerolide A is a macrolide polyketide that
possesses potent bioactivity against malignant melanoma cell lines while demonstrating minimal
cytotoxicity against other cell lines (17). The NCI's COMPARE algorithm was used to correlate
experimental findings with a database for prediction of the biochemical mechanism by identifying
the mechanism of action of palmerolide A as a vacuolar-ATPase (V-ATPase) inhibitor (33).
Downstream effects of V-ATPase inhibition include an increase in both hypoxia induction factor-
1α and autophagy (17, 34). Increased expression of V-ATPase on the surface of metastatic
melanoma cells (34) perhaps explain palmerolide A's selectivity for UACC-62 cell lines over all
cell types (17). Despite the relatively high concentrations of palmerolide A in the host tissue
(0.49–4.06 mg palmerolide A x g^{-1} host dry weight) (35), isolation of palmerolide A from its
Antarctic source in mass sufficient for drug development it is not ecologically nor logistically
feasible. Synthetic strategies for the synthesis of palmerolide A have been reported (6–13),
though a clear pathway to quantities needed for drug development has been elusive. Therefore,
there is substantial interest in identifying the biosynthetic gene cluster (BGC) responsible for
palmerolide A production.
Our approach to identify the palmerolide BGC (*pal* BGC) began with the characterization
of the host-associated microbiome (43). The core microbiome for *Synoicum adareanum*, a
persistent cohort of bacteria present in many individual ascidians, was established through
analysis of occurrence of distinct amplicon sequence variants (ASV) from the iTag sequencing of
the Variable 3-4 regions of the bacterial 16S rRNA (35). This ultimately led to the evaluation of
the metagenome and then the assembly of the metagenome assembled genome (MAG) of
Candidatus Synoicohabitans palmerolidicus, a verrucomicrobium in the family *Opitutaceae* (44).
Contained within the genome are five non-identical copies of the *pal* BGC, of which, three
correlate with palmerolides, the structures of which have been previously discovered in this
macrolide family. Here we report a detailed analysis of the *pal* BGCs.
160
161   **Results and Discussion**
162
163   **A. Retrobiosynthetic Scheme for Palmerolide A**
164
A retrobiosynthetic scheme of the *pal* BGC was developed based on the chemical structure
for palmerolide A, including modules consistent with a hybrid PKS-NRPS with tailoring enzymes
for key functional groups (**Fig. 1**). We hypothesized that the initial module would be PKS-like in
nature to utilize 3-methylcrotonic acid as the starter unit followed by a NRPS domain for the
incorporation of glycine. PKS elongation was predicted to be an 11-step sequence resulting in 22
contiguous carbons. Modifying enzymes that are encoded co-linearly were predicted to create the
architectural diversity with olefin placement, reduction of certain carbonyl groups to secondary
alcohols, and full reduction of other subunits. In addition, incorporation of methylmalonyl CoA or
enzymatic activity of carbon methyltransferases (cMTs) were predicted to be responsible for the
placement of methyl groups C-26 and C-27 from *S*-adenosylmethionine (SAM).
Several key structural features proposed to result from the action of *trans*-acting enzymes
are present. As seen in the kalimantacins (45), the carbamate on C-11 was hypothesized to be
installed by a carbamoyl transferase (CT). The C-25 methyl group is located on C-17, in the β-
position. This structural feature suggests the origin of this branch from hydroxymethylglutaryl-CoA
Synthase (HCS) catalysis, rather than the SAM-mediated methylation occurring at the α-position,
which appears to be the origin of the C-26 and C-27 methyl groups. Methylation in this acetate
carboxyl position is unusual, but represented in a number of notable BGC's, such as those of the
jamaicamides, bryostatins, curacin A, oocydin, pederin, and psymberin, among others; in
biochemically characterized Type I PKS BGCs, HCS-mediated β-branch formation is the common
mechanism (46–50). The hydroxy group on C-10, in the α-position of the acetate subunit, was

hypothesized to result from elongation resulting from hydroxymalonyl-CoA incorporation or from
the action of a hydroxylase at a later stage of biosynthesis.
187
188 **B. Proposed Biosynthesis and Architecture of the Putative *pal* Biosynthetic Gene Cluster**
189
Synoicum adareanum collected near Palmer Station, Antarctica was processed to separate
eukaryotic from prokaryotic cells of the host microbiome. Iterations of metagenomic analysis (44)
yielded a large number of BGCs when analyzed via the antiSMASH annotation platform (51). A
candidate cluster with multiple copy numbers was identified and has excellent congruence with
retrobiosynthetic predictions, including the key genomic markers outlined above that are likely
involved in biosynthesis of palmerolide A. Integration of BGC annotations with information from
protein family homology analysis, conserved domain searches, active site and motif identification,
supported the hypothesis that this putative 75 kb BGC is responsible for palmerolide A
production. The core scaffold was explained by the NRPS and *trans*-AT PKS hybrid system. In
addition, each of the tailoring enzymes that are expected for the distinct chemical features are
encoded in the genome. The proposed BGC for palmerolide A is comprised of 14 core
biosynthetic modules and 25 genes in a single operon of 74,655 bases (**Fig. 2**). Fourteen
modules are co-linear and two *trans*-AT domains (modules 15 & 16) follow the core biosynthetic
genes. Additional *trans*-acting genes contribute to backbone modifications with at least one gene
contributing to post-translational tailoring (**Fig. 3**).
205
An unusual starter unit and NRPS domains of *palA*. Bioinformatic analysis of the gene
sequence suggests that the initial core biosynthetic domains of *palA* (modules 1 and 2) encode
for the requisite acyl carrier proteins (ACP) (**Fig. 2**). Encoded in module 1, there are three ACPs
in tandem, which could serve to promote an increase in metabolite production (52). The second in
series is an ACP-β containing the conserved domain sequence GXDS (53) and is likely the
acceptor of a starter unit containing a β-branch, consistent with the proposed starter unit for
palmerolide A, 3-methylcrotonic acid. While both *trans*-acting ATs, PalE and PalF, (**Fig. 2**)
possess the catalytic active site serine which is key for the proper positioning of subunit within the
hydrophobic cleft of the active site (8, 54), only the first AT, PalE, has a characteristic motif that
includes an active site phenylalanine, conferring specificity for malonate selection (55). The AT
selecting the methylcrotonic acid starter unit is likely the second of the two *trans*-AT domains
(PalF), which lacks definitive specificity for malonyl-CoA. In support, some *trans*-acting ATs have
demonstrated affinity for a wider range of substrates than their *cis*-acting counterparts (26, 56). 3-
Methylcrotonyl-CoA is an intermediate of branched-chain amino acid catabolism in leucine
degradation; intermediates of which can be diverted to secondary metabolite production (57).
Interestingly, an upregulation of leucine catabolic pathways can be seen as a mechanism of cold
adaption to maintain cell wall fluidity via production of branched chain fatty acids, a factor which
may be at play in this Antarctic microorganism (57, 58). The subsequent NRPS module
containing condensation (C) and adenylation (A) domains as well as a carrier protein comprise
module 2. Signature sequence information and NRPSPredictor2 method (51, 59) of the A domain
are consistent with selection of a glycine residue. These domains incorporate the amino acid
residue, resulting in the addition of a nitrogen and two carbons in this step of the biosynthesis of
palmerolide A.
In a non-canonical fashion, the carrier proteins flanking the NRPS modules do not appear
to be the expected ACP and PCP for module 1 and 2, respectively. The carrier protein following
the KS domain in module 1 was initially annotated as a non-β-branching ACP; however,
phylogenetic analysis with the amino acid sequences of carrier proteins from other hybrid PKS-
NRPS systems, demonstrates that this carrier protein is in the same clade as peptidyl-carrier
proteins (*SI Appendix*, **Fig. S1**). The carrier protein associated with module 2 which was initially
annotated simply as a phosphopantetheine attachment site (Pfam00550.24) is found to be more
phylogenetically-related to ACPs within PKS-NRPS systems (*SI Appendix,* **Fig. S1**). Notably, it
possesses the (D/E)xGxDSL motif for phosphopantetheine arm attachment (60) with the
exception of an isoleucine rather than leucine in the final position of the motif, a residue seen in

this position in other ACPs from hybrid PKS-NRPS systems (**SI Appendix, Fig. S2**). Typically, a
PCP would follow the domains in NRPS-like modules; however, there are exceptions in the
literature. For example, the BGCs for both corallopyronin and oxazolamycin contain ACPs
following an A domain (61, 62). This non-canonical finding could point to the acquisition of these
domains over time, as the carrier protein for module 1 is encoded in *palA*, the same gene as the
proteins for both modules 1 and 2, whereas the carrier protein for module 2 is encoded at the
beginning of *palB*, a gene which encodes for only PKS domains (**Fig. 2**).
246
Contiguous PKS chain and *trans*-acting enzymes at site of action for *palB – palD*. The
contiguous carbon backbone of palmerolide A arises from 11 cycles of elongation from a series of
modules with a variety of enzymatic domains that include an ACP, KS, and associated genes that
establish the oxidation state of each subunit (**Fig 2**). The first module of *palB* (module 3) includes
a dehydratase (DH) and cMT domains, a sequence which results in a chain extension
modification to an α,β-unsaturated thioester, a result of the action of the encoded DH. The
expected KR domain that would be responsible for the Δ^{22} olefin (**Fig. 2**) is not present. The
BGCs for bryostatin 1, corallopyronin, and sorangicin also lack an accompanying KR domain to
work in concert with an encoded DH. The unaccompanied DH in the first two systems are
deemed inactive; however, an olefin results from the DH in the absence of an accompanying KR
in both modules 9 and 11 of the sorangicin BGC (46, 61, 63). The subsequent cMT methylation is
consistent with an *S*-adenosylmethionine (SAM)-derived methyl group, as expected for C-27 in
palmerolide A. Module 4, spanning the end of *palB* and beginning of *palC*, includes a DH, a
ketoreductase (KR), and another cMT domain. This cluster of domains is predicted to result in the
methyl-substituted conjugated diene of the macrolide tail (C-19 through C-24, C-26 on
palmerolide A).
The substrate critical for macrolactonization of the polyketide is the C-19 hydroxy group, a
result of the KS and KR domains encoded in module 5 (**Fig. 2**). Additionally, a domain initially
annotated as a dehydratase (DHt) at this location may contribute to the final cyclization and
release of the molecule from the megaenzyme by assisting the terminal C domain with ring
closure (53). This sequence shows some homology with condensation domains and does not
possess the hotdog fold that is indicative of canonical dehydratases, and therefore, may not truly
represent a DH (64). Alternatively, this domain could be responsible for the olefin shifts to the β,γ-
positions, as seen in bacillaene and ambruticin biosynthesis (65, 66).
Only an enoyl-CoA hydratase (ECH) in addition to standard ACP and KS is encoded in
module 6 which would lead to a ketone function; based on our retrobiosynthetic analysis, the
resulting ketone at C-17 is the necessary substrate for HCS-catalyzed β-branch formation,
resulting in the C-25 methyl group on C-17. The HCS cassette (PalK through PalO) is comprised
of a series of *trans*-acting domains, including an ACP, an HCS, a free KS, and 2 additional ECH
modules (**Fig. 2**). The HCS cassette can act while the elongating chain is tethered to an ACP
module, rather than after cyclization and release (67, 68). The two ECHs in the HCS cassette
along with the ECH encoded in-line with the core biosynthetic genes would be responsible for
isomerization of a terminal methylene to the observed internal olefin. An HCS cassette formed by
the combination of a *trans*-KS and at least one ECH module with an HCS domain is reported in
several other bacterial BGCs such as bryostatin 1, calyculin A, jamaicamide, mandelalide,
phormidolide, and psymberin (47, 50, 53, 69–71). The domain structure for the HCS cassettes
has a remarkably high degree of synteny across these diverse BGCs (72).
There is substantial similarity in the domain structure of module 7, module 10, and module
13, which each include a KS, DH, and KR (**Fig. 2**). The olefin that arises from the action of
module 7, concomitant with carbon chain elongation, is conjugated with the Δ^{16} olefin adjacent
the C-17 β-branch. Modules 10 and 13 have similar enzymatic composition to 7 and are likely
responsible for Δ^8 and Δ^2 olefins. The combination of KR and DH domains are also found in
modules 8 and 12; however, in concert with an unidentified *trans*-acting enoyl reductase domain
(ER), these olefins would be reduced to fully saturated monomeric subunits. There are some
examples of *trans*-acting ER domains carrying out this function, including OocU in oocydin, SorN
in sorangicin, and MndM in mandelalide (49, 63, 71), while in other systems, such as

corallopyronin and leinamycin, the reductions of the olefins are largely unexplained (61, 73). The
reduction by a *trans*-acting enzyme often occurs while the elongating polyketide is tethered to the
megaenzyme, as evidenced by the downstream specificity of the KS module for Claisen-type
condensation with subunits containing single or double bonds (63).
The genetic architecture for the biosynthesis of two functional groups, essential for bioactivity,
is encoded in module 9 (**Fig. 2**); structure-activity relationship studies demonstrate the
importance of the C-10 hydroxyl group and the C-11 carbamate (74). The KR domain that is
present, predicting the C-11 alcohol function, serves as the substrate for the carbamoyl
transferase (*palQ*) in a post-translational modification (75–77). Intriguingly, a domain annotated
as a luciferase-like monooxygenase (LLM) initially seems out of place; however, palmerolide A
has a hydroxy group at C-10, which is the α-position of the acetate subunit inserted in module 9.
LLMs associated with BGCs may not serve as true luciferases, but instead demonstrate oxidizing
effects on polyketides and peptides without evidence of corresponding bioluminescence (78, 79).
For example, there is an overrepresentation of LLMs in *Candidatus* Entotheonella BGCs without
known bioluminescence (80). As demonstrated through individual inactivation of the LLM in the
BGC of mensacarin, a Type II PKS system, Msn02, Msn04, and Msn08 have key activity as
epoxidases and hydroxlases (79). There are several examples of LLMs in modular Type I PKS
systems. OnnC from onnamide and NazB from nazumamide are two LLMs in *Candidatus*
Endotheonella that are proposed to serve biosynthetically as hydroxylases (80). In calyculin and
mandelalide, the CalD and MndB LLMs catalyze chain shortening reactions through α-
hydroxylation and Baeyer-Villiger-type oxidation reactions (69, 71). Phormidolide has a LLM that
adds a hydroxy group which is then hypothesized to attack an olefin through a Michael-type
addition for cyclization with enzymatic assistance from a pyran synthase (53). The hydroxylation
that is key in cyclization of oocydin A is likely installed by OocK or OocM, flavin-dependent
monooxygenases that are contiguous to the PKS genes and are thought to act while the
substrate is bound to a portion of the PKS megaenzyme (49). It is this hydroxylase activity that
we propose for the LLM in module 9. Since the producing bacteria is yet to be cultured, it is not
established whether this LLM may also serve a role in bioluminescence and/or quorum sensing.
Further evidence for the role of the LLM is provided through alignment against other LLM's. The
pal BGC contains both LuxR family transcriptional regulator as well as the DNA-binding response
regulator. In addition to the annotation within Pfam00296, which includes the bacterial LLMs, the
sequence is homologous with the multiple sequence alignments and the hidden Markov models
of the TIGR subfamily 04020 of the conserved protein domain family cl19096, which is noted for
natural product biosynthesis LLMs (80). The subfamily occurs in both NRPS and PKS systems as
well as small proteins with binding of either flavin mononucleotide or coenzyme F420. Alignment
of the LLMs from multiple PKS systems, including palmerolide A, shows homology with model
sequences from the TIGR subfamily 04020 (*SI Appendix*, **Fig. S3**).
The addition of C-5 and C-6 and the reduction of the β-carbonyl to form the C-7 hydroxy
group of palmerolide A, is due to module 11, which possesses a KR domain in addition to
elongating KS (**Fig. 2**). In the structure of palmerolide A this is followed by the fully reduced
subunit from module 12 as discussed above. The final elongation results from module 13, which
includes DH and KR domains that contribute to the conjugated ester found as palmerolide A's C-
1 through C-3, completing the palmerolide A C_{24} carbon skeleton.
336
Noncanonical termination condensation domain in *palD* for product cyclization and
release. Typically, PKS systems terminate with a thioesterase (TE) domain, leading to release of
the polyketide from the megaenzyme (71, 81–83). This canonical domain is not present in the *pal*
cluster. Instead, the final module in the *cis*-acting biosynthetic gene cluster includes a truncated
condensation domain comprised of 133 amino acid residues; compared to the approximately 450
residues that comprise a standard condensation domain (84) (**Fig. 2**). Condensation domains
catalyze cyclization through ester formation in free-standing condensation domains that act in
trans as well as in NRPS systems (85, 86). In addition, this non-canonical termination domain is
not without precedent in hybrid PKS-NRPS and in PKS systems, as basiliskamide and
phormidolide both include condensation domains for product release (53, 87). Though the

terminal condensation domain in the *pal* BGC is truncated, it maintains much of the HHXXDDG
motif (*SI Appendix,* **Fig. S3**), most notably the second histidine, which serves as the catalytic
histidine in the condensation reaction (84).
350
Stereochemical and structural confirmation based on sequence information. KR domains
are NADPH-dependent enzymes that belong to the short-chain dehydrogenase superfamily, with
Rossman-like folds for co-factor binding (60, 88). Enzymatically, the two KR subtypes, A-Type
and B-type, are responsible for stereoselective reduction of β-keto groups and can also
determine the stereochemistry of α-substituents. C-type KRs, however, lack reductase activity
and often serve as epimerases. A-Type KRs have a key tryptophan residue in the active site, do
not possess the LDD amino acid motif, and result in the reduction of β-carbonyls to ʟ-configured
hydroxy groups (60). B-Type, which are identified by the presence of an LDD amino acid motif,
result in formation of ᴅ-configured hydroxy groups (60). The stereochemistry observed in
palmerolide A is reflected in the active site sequence information for the ᴅ-configured hydroxyl
groups from module 5 and module 11 (**Fig. 1** and **Fig. 2**). When an enzymatically active DH
domain is within the same module, the stereochemistry of the *cis-* versus *trans*-olefin can be
predicted, as the combination of an A-Type KR with a DH results in a *cis*-olefin formation and the
combination of a B-Type KR with a DH results in *trans*-olefin formation. The *trans*-α,β-olefins
arising from module 7 (Δ^{14}), module 10 (Δ^8), and module 13 (Δ^2) arise from B-Type KRs and
active DHs. The other three olefins present in the structure of palmerolide A, as noted above,
likely have positional and stereochemical influence during the enzymatic shifts to the β,γ-
positions (Δ^{21} and Δ^{23}) or from the ECH domain (module 6).
Additional structural information was obtained through defining the specificity of KS
domains using phylogenetic analysis and the *trans*-AT PKS Polyketide Predictor (*trans*ATor)
bioinformatic tool (89). KS domains catalyze the sequential two-carbon elongation steps through
a Claisen-like condensation with a resulting β-keto feature (90). Additional domains within a given
module can modify the β-carbonyl or add functionality to the adjacent α- or γ-positions (60).
Specificity of KSs, based the types of modification located on the upstream acetate subunit were
determined and were found to be mostly consistent with the retrobiosynthetic predictions (*SI*
Appendix, **Fig. S4,** *SI Appendix,* **Table S1**). For example, the first KS, KS1 (module 1), is
predicted to receive a subunit containing a β-branch. KS3 (module 4) and KS4 (module 5) are
predicted to receive an upstream monomeric unit with α-methylation and an olefinic shift,
consistent with the structure of palmerolide A and with the enzymatic transformations resulting
from module 3 and module 4, respectively. Interestingly, the KS associated with the HCS
cassette is in a clade of its own in the phylogenetic tree. *Trans*ATor also aided in confirming the
stereochemical outcomes of the hydroxy groups and olefins, which occur through reduction of the
β-carbonyls. The predictions for the ᴅ-configured hydroxy groups were consistent with not only
the presence of the LDD motif, indicative of type B-type KR as outlined above, but also with
stereochemical determination based on the clades of the KS domains of the receiving modules,
KS5 (module 6) and KS11 (module 12). They are also consistent with the structure of palmerolide
A. The KS predictions, however, did not aid in confirming reduction of the upstream olefins for
KS8 (module 9) and KS12 (module 13).
389
Additional *trans*-acting domains and domains between genes responsible for
biosynthesis. A glycosyl transferase (*palP*) and lactone oxidase (*palH*) that are associated with
glycosylation of polyketides are encoded in the palmerolide A BGC (**Fig. 2**), though glycosylated
palmerolides have not been observed. Glycosylation as a means of self-resistance has been
described for oleandomycin, which encodes not only two glycosyl transferases, one with
substrate specificity toward oleandomycin and the other relaxed specificity, but also a glycosyl
hydrolase (91, 92). The glycosylation of the macrolide confers internal resistance to ribosomal
inhibition, making the glycosylated macrolide a pro-drug, which is activated by the glycosyl
hydrolase that is excreted extracellularly (91, 92). Glycosyl transferases are found in other
macrolide- and non-macrolide-producing organisms as a means to inactivate hydroxylated
polyketides (93, 94). Though prokaryotic V-ATPases tend to be simpler than those of eukaryotes,

there is homology in the active sites making the pro-drug hypothesis for self-resistance
reasonable in palmerolide A biosynthesis (95). The D-arabinono-1,4-lactone oxidase (*palH*) is a
FAD-dependent oxidoreductase that likely works in concert with the glycosyltransferase. An ATP-
binding cassette (ABC) transporter is encoded between the core biosynthetic genes and the
genes for the *trans*-acting enzymes. This transporter, which has homology to SryD and contains
the key nucleotide-binding domain GGNGSGKST, may be responsible for the translocation of the
macrolide out of the cell. Since it is housed within the BGC, it is likely under the same regulatory
control. Additionally, a Band7 protein is encoded. Band7 proteins belong to a ubiquitous family of
slipin or stomatin-like integral membrane proteins that are found in all kingdoms, yet lack
assignment of function (96). A domain of unknown function (DUF) is annotated; however, this
particular DUF, DUF179, has no superfamily relatives in protein homology searches. Together,
these *trans*-acting domains decorate the carbon backbone with features that give palmerolide A
its unique chemical structure and contribute to its bioactivity.

C. Multiple copies of the *pal* Biosynthetic Gene Cluster Explain Structural Variants in the Palmerolide Family

Careful assembly of the metagenome assembled genome (MAG) of Ca. *S. palmerolidicus*
revealed the BGC was present in multiple copies (**Fig. 4** and ***SI Appendix,* Fig. S5-S7**) (44),
evidenced by their independent anchoring loci within the MAG and supported by a five-fold
increase in depth of coverage relative to the rest of the genome. The structural complexity of the
multicopy BGCs represents a biosynthetic system that is similar to that which is found in *Ca.*
Didemnitutus mandela, another ascidian-associated verrucomicrobium in the family *Opitutaceae*
(71). A total of 5 distinct BGCs consistent with Type-I PKS systems that have much overlap with
one another and are likely responsible for much of the structural diversity in the family of
palmerolides (17, 18) (**Fig. 4**). Palmerolide A is hypothesized to arise from the BGC designated
as *pal* BGC 4 with additional compounds also from this same cluster. The clusters designated as
pal BGC 1, *pal* BGC 2, *pal* BGC 3, and *pal* BGC 5 have been identified in *Ca.* S. palmerolidicus
and the potential products of each are described below. It is hypothesized that there are three
levels of diversity introduced to create the family of palmerolides: (1) differences in the site of
action for the *trans*-acting domains (with additional *trans*-acting domains at play as well), (2)
promiscuity of the initial selection of the starter subunit, and (3) differences in the core
biosynthetic genes with additional PKS domains or stereochemical propensities within a module.
There are several palmerolides that likely arise from the same BGC encoding the
megaenzyme responsible for palmerolide A (*pal* BGC 4). We hypothesize that the *trans*-acting
domains have different sites of action than what is seen in palmerolide A biosynthesis. For
example, the chemical scaffold of palmerolide B (**Fig. 4**) is similar to palmerolide A, though the
carbamate transfer occurs on the C-7 hydroxy group. Palmerolide B instead bears a sulfate group
on the C-11 hydroxy group; proteins with homology to multiple types of sulfatases from the
UniProtKB database (P51691, P15289, O69787, Q8ZQJ2) are found in the genome of *Ca.* S.
palmerolidicus (44), but are not encoded within the BGCs. One of these *trans*-acting sulfatases
likely modifies the molecule post-translationally. Other structural differences including the
hydroxylation on C-8 instead of C-10 (as observed in palmerolide A) and the Δ^9 olefin that differs
from palmerolide A's Δ^8 olefin, are either due to a difference of the site of action of the LLM
(module 9) or a *trans*-acting hydroxylase. Another member of the compound family, palmerolide
C, has structural differences attributable to *trans*-acting enzymes as well. Again, a *trans*-acting
hydroxylase or the LLM is proposed to be responsible for hydroxylation on C-8. A hydroxy group
on C-9 occurs through reduction of the carbonyl. The carbamate installation occurs on C-11 after
trans-acting hydroxylation or LLM hydroxylation. In addition, the Δ^8 olefin in palmerolide A is not
observed, but rather a Δ^6 olefin.
Additional levels of structural variation are seen at the site of the starter unit, likely due to
a level of enzymatic promiscuity of the second AT (PalF). This, combined with differences in the
sites of action for the *trans*-acting domains, is likely responsible for the structural differences
observed in palmerolide F (**Fig. 4**). The terminal olefin on the tail of the macrolide which perhaps

is a product of promiscuity of the selection of the starter unit, the isomeric 3-methyl-3-butenoic
acid, is consistent with the aforementioned lack of consensus for malonate selection by the AT. In
addition, the KS that receives the starter unit is phylogenetically distinct from the other KS in the
pal clusters (***SI Appendix,*** **Fig. S4**).

The retrobiosynthetic hypothesis for palmerolide G (**Fig. 4**) has much similarity to what is
present in *pal* BGC 4; however, the presence of a *cis*-olefin rather than a *trans*-olefin could arise
from a difference in the enzymatic activity of module 4. This olefin is the result of a shift from an
epimerase and, therefore, the stereochemistry is not solely reliant upon the action of the
associated KR. Although this difference has not been identified in the BGCs in the samples
sequenced, this could be present in other environmental samples that have been batched for
processing and compound isolation. Currently, the biosynthetic mechanism is unknown.

The core biosynthetic genes of two palmerolide BGCs (*pal* BGC 1 and *pal* BGC 3) are
identical to one another (***SI Appendix,*** **Fig. S5**) and possess an additional elongation module
when compared to *pal* BGC 4. Palmerolide D (**Fig. 4**) is structurally very similar to palmerolide A
with the exception of elongation in the carboxylate tail of the macrolide by an isopropyl group.
This could arise from one additional round of starter unit elongation via a KS and methylation.
These two identical BGCs are consistent with this structural difference. The overall architecture
and stereochemistry are otherwise maintained. Palmerolide H (**Fig. 4**) includes the structural
differences of both palmerolide B and palmerolide D. It contains the extended carboxylate tail with
a terminal olefin and incorporates hydroxylation on C-8 rather than C-10. Again, there is no
genomic evidence that this hydroxylation in the α-position is due to incorporation of
hydroxymalonate to explain this but is instead likely due to a *trans*-hydroxylase. The carbamate
installation occurs on C-7, while sulfonation occurs on C-11 and α-hydroxy placement is on C-8.

The final two palmerolide BGCs both have truncation of the core biosynthetic genes. The
gene structure of *pal* BGC 5 (***SI Appendix,*** **Fig. S6**) shows preservation of genes upstream to the
core biosynthetic genes; however, there are no pre-NRP PKS modules noted in the BGC. The
HCS cassette, glycosyl transferase, and CT are all present downstream. The predicted product of
this cluster does not align with a known palmerolide, though post-translational hydrolysis of the C-
24 amide may result in a structure similar to palmerolide E (**Fig. 4**), which maintains much of the
structure of palmerolide A; however, is missing the initial polyketide starter unit and the glycine
subunit. The final *pal* BGC in *Ca.* S. palmerolidicus, *pal* BGC 2 (***SI Appendix,*** **Fig. S7**), includes
only 5 elongating modules, which would result in a 10-carbon structure that has not been
observed. Interestingly, despite the shortened BGC, the HCS cassette, glycosyl transferase, and
CT are all present downstream. There would only be a single hydroxy group serving as a
substrate for the CT, glycosyl transferase, and sulfatase to act. The 2-carbon site of action for the
β-branch introduced in the palmerolide A structures would not be present. The structure-based
retrobiosynthesis of the eight known palmerolides (A-F) can be hypothesized to arise from
differences in the core biosynthetic genes of these non-identical copies of the *pal* BGC, starter
unit promiscuity, and differing sites of action in the *trans*-acting enzymes.

494
495

Conclusion

The putative *pal* BGC has been identified and represents the first BGC elucidated from an
Antarctic organism. As outlined in a retrobiosynthetic strategy, the *pal* BGC represents a *trans*-AT
Type I PKS/NRPS hybrid system with compelling alignment to the predicted biosynthetic steps for
palmerolide A. The *pal* BGC is proposed to begin with PKS modules resulting in the incorporation
of an isovaleric acid derivative, 3-methylcrotonic acid, as a starter unit, followed by incorporation
of a glycine residue with NRPS-type modules. Then, eleven rounds of progressive polyketide
elongation likely occur leading to varying degrees of oxidation introduced with each module.
There are several interesting non-canonical domains encoded within the BGC, such as an HCS,
CT, LLM, and a truncated condensation termination domain. Additionally, a glycosylation domain
may be responsible for reversible, pro-drug formation to produce self-resistance to the V-ATPase

activity of palmerolide A. There are additional domains, the function of which have yet to be
determined.
A combination of modular alterations, starter unit differences, and activity of *trans*-acting
enzymes contributes to Nature's production of a suite of palmerolide analogues. There are a total
of five distinct *pal* BGCs in the MAG of *Ca.* S. palmerolidicus, yielding the known eight
palmerolides, with genetic differences that explain some of the structural variety seen within this
family of compounds. These include differences in modules that comprise the core biosynthetic
genes. Additionally, it is proposed that some of the architectural diversity of palmerolides arises
from different sites of action of the *trans*-acting, or non-colinear, modules. Starter unit promiscuity
is another potential source of the structural differences observed in the compounds. Analysis of
the *pal* BGC not only provides insight into the architecture of this Type I PKS/NRPS hybrid BGC
with unique features, but also lays the foundational groundwork for drug development studies of
palmerolide A via heterologous expression.
521
522
523
524 **Materials and Methods**
525
The details of field collections, sample processing and genomic methods (sequencing,
assembly, and analysis) used to identify the putative biosynthetic gene clusters are described in
detail in two related publications (35, 44). In brief, *Synoicum adareanum* samples were collected
by SCUBA from the Antarctic Peninsula in the Anvers Island Archipelago and flash frozen.
Microbial cells were separated from host tissue using a homogenization protocol established for
this holobiont followed by differential centrifugation to separate the host cellular debris and
microbial cells (31). The metagenome assembled genome (MAG) sequence analyzed in this
study was generated from two *S. adareanum* samples (Bon-1c-2011 and Del-2b-2011) with high
copy numbers of the putative biosynthetic gene cluster sequenced using PacBio technology. The
resulting assembly of produced a nearly complete 4.3 MB genome, with five unique contigs and
five varying copies of the *pal* BGC (referred to as *pal* BGC 1 through *pal* BGC 5) (44). Given the
phylogenomic novelty of the MAG, the name *Candidatus* Synoicihabitans palmerolidicus was
proposed for this new genus in the *Opitutaceae* family (Verrucomicrobia phylum).
The methods employed in this study used bioinformatic tools to develop predictive models
of palmerolide biosynthesis. Enzymatic reactions and organic synthetic interpretations were
based on homology analyses. Automated annotation and manual bioinformatic tools were used to
discern the details of palmerolide A biosynthesis in addition to generating predictions for the other
BGCs. The *Ca*. S. palmerolidicus MAG was annotated using antiSMASH (v. 5.0) (51) using the
full complement of annotation options available. Then we predicted the gene cluster responsible
for palmerolide A biosynthesis using retrobiosynthetic predictions focused on the 5' end of the
BGCs (Fig. 1). Only one of the five BGCs met the criteria for a non-ribosomal peptide glycine
starter unit. The annotation predictions were integrated and validated with results of additional
protein family homology analysis, conserved domain searches, active site and motif identification
to predict the step-wise biosynthesis of palmerolide A. Manual annotation of the *pal* BGC
sequences included BLASTP searches to confirm enzymatic identities, then protein family
alignments were used to identify active site residues key for stereochemical outcomes, confirm
substrate affinities, and other biochemical synthesis details.
Additional manual bioinformatic efforts included obtaining BGCs from public NCBI
databases for basiliskamide, bryostatin 1, calyculin, corallopyronin, mandelalide, onnamide,
oxazolamycin, pederin, phormidolide, psymberin, sorangicin, and myxoviricin (***SI Appendix***,
Table S2). ClustalO alignment tool in the CLC Genome Workbench (QIAGEN aarhau A/S v.
20.0.3) was used for multiple sequence alignments of enzymatic domains with HMM Pfam Seeds
obtained from EMBL-EBI and the amino acid sequences from the other PKS BGCs. MIBiG (97)
was used to acquire the KS amino acid sequence from the type III PKS BGC responsible for 3-
(2'-hydroxy-3'-oxo-4'-methylpentyl)-indole biosynthesis from *Xenorhabdus bovienii SS-2004*
(GenBank Accession: FN667741.1), which was used for an outgroup. The *pal* BGC ACPs and

PCPs were numbered according to their position in the proposed biosynthesis of palmerolide A (***SI Appendix*, Fig. S1 and S2**). The BGC KSs were numbered according to their position in their proposed biosynthesis in the literature. Prior to the construction of the phylogenetic tree for the KS domains (***SI Appendix*, Fig. S4**), the sequences in the alignment were manually inspected and trimmed. Phylogenetic trees were created in the CLC Genome Workbench (QIAGEN aarhau A/S v. 20.0.3) with Neighbour Joining (NJ) as a distance method and Bayesian estimation for ACP and PCP comparisons as well as for KS analysis. Jukes-Cantor was selected for the genetic distance model and bootstrapping with performed with 100 replicates. Additionally, the sequence of each KS in the *pal* BGCs was queried using the *trans*-AT PKS Polyketide Predictor (*trans*ATor) to help define the specificity of KS domains. The software is based on phylogenetic analyses of fifty-four *trans*-AT type I PKS systems with 655 KS sequences and the resulting clades are referenced to help predict the KS specificity for the upstream unit (89).

Data Deposition

Data from this project has been deposited to the NCBI under BioProject Accession Number PRJNA662631. The GenBank Accession Number for the MAG of *Ca.* Synoicohabitans palmerolidicus is JAGGDC000000000. The BGCs can be accessed in the MiBIG database with the associated Accession Numbers: BGC0002118 (for *pal* BGC 4) and BGC0002119 (for *pal* BGC 3).

Acknowledgments

Support for this research was provided in part by the National Institutes of Health award (CA205932) to A.E.M., B.J.B., and P.S.G.C., with additional support from National Science Foundation awards (ANT-0838776, and PLR-1341339 to B.J.B., ANT-0632389 to A.E.M.) The authors acknowledge the assistance of field team members, including William Dent, Charles D. Amsler, James B. McClintock, Margaret O. Amsler, and Katherine Schoenrock. This work would not have been possible without the outstanding logistical support of the United States Antarctic Program. Lucas Bishop, Robert Read, and Mary L. Higham are also recognized for their contributions.

References

1. A. R. Carroll, B. R. Copp, R. A. Davis, R. A. Keyzers, M. R. Prinsep, Marine natural products. *Nat. Prod. Rep.* **36**, 122–173 (2019).
2. C. A. Dejong, *et al.*, Polyketide and nonribosomal peptide retro-biosynthesis and global gene cluster matching. *Nat. Chem. Biol.* **12**, 1007–1014 (2016).
3. J. G. Lawrence, H. Ochman, M. A. Ragan, Reconciling the many faces of lateral gene transfer. *Trends Microbiol.* **10**, 1–4 (2002).
4. M. Ravenhall, N. Škunca, F. Lassalle, C. Dessimoz, Inferring Horizontal Gene Transfer. *PLoS Comput. Biol.* **11**, 1–16 (2015).
5. I. Schmitt, H. T. Lumbsch, Ancient horizontal gene transfer from bacteria enhances biosynthetic capabilities of fungi. *PLoS One* **4**, 1–8 (2009).
6. E. W. Schmidt, Trading molecules and tracking targets in symbiotic interactions. *Nat. Chem. Biol.* **4**, 466–473 (2008).
7. E. W. Schmidt, The secret to a successful relationship: Lasting chemistry between ascidians and their symbiotic bacteria. *Invertebr. Biol.* **134**, 88–102 (2015).
8. E. J. N. Helfrich, J. Piel, Biosynthesis of polyketides by *trans*-AT polyketide synthases. *Nat. Prod. Rep.* **33**, 231–316 (2016).
9. A. M. P. Koskinen, K. Karisalmi, Polyketide stereotetrads in natural products. *Chem. Soc. Rev.* **34**, 677–690 (2005).

10. J. Napolitano, A. Daranas, M. Norte, J. Fernandez, Marine Macrolides, a Promising
Source of Antitumor Compounds. *Anticancer. Agents Med. Chem.* **9**, 122–137 (2012).
11. R. Ueoka, *et al.*, Metabolic and evolutionary origin of actin-binding polyketides from
diverse organisms. *Nat. Chem. Biol.* **11**, 705–712 (2015).
12. M. E. Bordeleau, *et al.*, Stimulation of mammalian translation initiation factor eIF4A activity
by a small molecule inhibitor of eukaryotic translation. *Proc. Natl. Acad. Sci. U. S. A.* **102**,
10460–10465 (2005).
13. S. Nishimura, *et al.*, Structure-activity relationship study on 13-deoxytedanolide, a highly
antitumor macrolide from the marine sponge *Mycale adhaerens*. *Bioorganic Med. Chem.*
13, 455–462 (2005).
14. R. Shen, C. T. Lin, E. J. Bowman, B. J. Bowman, J. A. Porco, Lobatamide C: Total
synthesis, stereochemical assignment, preparation of simplified analogues, and V-ATPase
inhibition studies. *J. Am. Chem. Soc.* **125**, 7889–7901 (2003).
15. E. J. Bowman, K. R. Gustafson, B. J. Bowman, M. R. Boyd, Identification of a New
Chondropsin Class of Antitumor Compound that Selectively Inhibits V-ATPases. *J. Biol.*
Chem. **278**, 44147–44152 (2003).
16. S. Kazami, *et al.*, Iejimalides show anti-osteoclast activity via V-ATPase inhibition. *Biosci.*
Biotechnol. Biochem. **70**, 1364–1370 (2006).
17. T. Diyabalanage, C. D. Amsler, J. B. McClintock, B. J. Baker, Palmerolide A, a cytotoxic
macrolide from the antarctic tunicate *Synoicum adareanum*. *J. Am. Chem. Soc.* **128**,
5630–5631 (2006).
18. J. H. Noguez, *et al.*, Palmerolide macrolides from the Antarctic tunicate *Synoicum*
adareanum. *Bioorganic Med. Chem.* **19**, 6608–6614 (2011).
19. B. Shen, *et al.*, Polyketide biosynthesis beyond the type I, II, and III polyketide synthase
paradigms: A progress report. *ACS Symp. Ser.* **955**, 154–166 (2007).
20. B. Wilkinson, *et al.*, Novel octaketide macrolides related to 6-deoxyerythronolide B provide
evidence for iterative operation of the erythromycin polyketide synthase. *Chem. Biol.* **7**,
111–117 (2000).
21. S. Tatsuno, K. Arakawa, H. Kinashi, Analysis of Modular-iterative Mixed Biosynthesis of
Lankacidin by Heterologous Expression and Gene Fusion. *J. Antibiot. (Tokyo).* **60**, 700–
708 (2007).
22. B. Wang, F. Guo, C. Huang, H. Zhao, Unraveling the iterative type I polyketide synthases
hidden in *Streptomyces*. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 8449–8454 (2020).
23. T. A. Nguyen, *et al.*, Exploiting the mosaic structure of *trans*-acyltransferase polyketide
synthases for natural product discovery and pathway dissection. *Nat. Biotechnol.* **26**, 225–
233 (2008).
24. S. F. Haydock, *et al.*, Divergent sequence motifs correlated with the substrate specificity of
(methyl)malonyl-CoA:acyl carrier protein transacylase domains in modular polyketide
synthases. *FEBS Lett.* **374**, 246–248 (1995).
25. H. Jenke-Kodama, A. Sandmann, R. Müller, E. Dittmann, Evolutionary implications of
bacterial polyketide synthases. *Mol. Biol. Evol.* **22**, 2027–2039 (2005).
26. A. Nivina, K. P. Yuet, J. Hsu, C. Khosla, Evolution and Diversity of Assembly-Line
Polyketide Synthases. *Chem. Rev.* **119**, 12524–12547 (2019).
27. P. Videau, K. N. Wells, A. J. Singh, W. H. Gerwick, B. Philmus, Assessment of *Anabaena*
sp. Strain PCC 7120 as a Heterologous Expression Host for Cyanobacterial Natural
Products: Production of Lyngbyatoxin A. *ACS Synth. Biol.* **5**, 978–988 (2016).
28. E. J. Kim, *et al.*, Heterologous Production of 4-O-Demethylbarbamide, a Marine
Cyanobacterial Natural Product. *Org. Lett.* **14**, 5824–5827 (2012).
29. C. Greunke, *et al.*, Direct Pathway Cloning (DiPaC) to unlock natural product biosynthetic
potential. *Metab. Eng.* **47**, 334–345 (2018).
30. J. B. McClintock, C. D. Amsler, B. J. Baker, R. W. M. Van Soest, Ecology of antarctic
marine sponges: An overview. *Integr. Comp. Biol.* **45**, 359–368 (2005).
31. A. Clarke, J. A. Crame, Evolutionary dynamics at high latitudes: Speciation and extinction
in polar marine faunas. *Philos. Trans. R. Soc. B Biol. Sci.* **365**, 3655–3666 (2010).

32. R. M. Young, *et al.*, Site-specific variability in the chemical diversity of the Antarctic red
671    alga *Plocamium cartilagineum. Mar. Drugs* **11**, 2126–2139 (2013).
672 33. K. D. Paull, E. Hamel, L. Malspeis, Prediction of Biochemical Mechanism of Action from
673    the In Vitro Antitumor Screen of the National Cancer Institute. *Cancer Chemother. Agents*,
674    9–45 (1995).
675 34. K. Von Schwarzenberg, *et al.*, Mode of cell death induction by pharmacological vacuolar H
676    $^+$-ATPase (V-ATPase) inhibition. *J. Biol. Chem.* **288**, 1385–1396 (2013).
677 35. A. E. Murray, *et al.*, Uncovering the Core Microbiome and Distribution of Palmerolide in
678    *Synoicum adareanum* Across the Anvers Island Archipelago, Antarctica. *Mar. Drugs* **18**,
679    298 (2020).
680 36. K. P. Kaliappan, P. Gowrisankar, Synthetic studies on a marine natural product,
681    palmerolide A: Synthesis of C1-C9 and C15-C21 fragments. *Synlett*, 1537–1540 (2007).
682 37. M. P. Lisboa, D. M. Jones, G. B. Dudley, Formal synthesis of palmerolide A, featuring
683    alkynogenic fragmentation and syn-selective vinylogous aldol chemistry. *Org. Lett.* **15**,
684    886–889 (2013).
685 38. M. D. Lebar, B. J. Baker, Synthesis of the C3-14 fragment of palmerolide A using a chiral
686    pool based strategy. *Tetrahedron* **66**, 1557–1562 (2010).
687 39. Z. K. Wen, Y. H. Xu, T. P. Loh, Palladium-catalyzed cross-coupling of unactivated alkenes
688    with acrylates: Application to the synthesis of the C13-C21 fragment of palmerolide A.
689    *Chem. - A Eur. J.* **18**, 13284–13287 (2012).
690 40. S. A. Pujari, P. Gowrisankar, K. P. Kaliappan, A shimizu non-aldol approach to the formal
691    total synthesis of palmerolide A. *Chem. - An Asian J.* **6**, 3137–3151 (2011).
692 41. K. R. Prasad, A. B. Pawar, Stereoselective synthesis of C1-C18 region of palmerolide A
693    from tartaric acid. *Synlett* **4**, 1093–1095 (2010).
694 42. A. B. Pawar, K. R. Prasad, Formal total synthesis of palmerolide A. *Chem. - A Eur. J.* **18**,
695    15202–15206 (2012).
696 43. C. S. Riesenfeld, A. E. Murray, B. J. Baker, Characterization of the microbial community
697    and polyketide biosynthetic potential in the palmerolide-producing tunicate *Synoicum
698    adareanum. J. Nat. Prod.* **71**, 1812–1818 (2008).
699 44. A. E. Murray, *et al.*, Discovery of an Antarctic ascidian-associated uncultivated
700    Verrucomicrobia that encodes antimelanoma palmerolide biosynthetic capacity. *BioRxiv.*
701    (Preprint: 10.1101/2021.05.05.442870) (2021).
702 45. W. Mattheus, *et al.*, Isolation and Purification of a New Kalimantacin/Batumin-Related
703    Polyketide Antibiotic and Elucidation of Its Biosynthesis Gene Cluster. *Chem. Biol.* **17**,
704    149–159 (2010).
705 46. S. Sudek, *et al.*, Identification of the putative bryostatin polyketide synthase gene cluster
706    from "*Candidatus* Endobugula sertula", the uncultivated microbial symbiont of the marine
707    bryozoan *Bugula neritina. J. Nat. Prod.* **70**, 67–74 (2007).
708 47. D. J. Edwards, *et al.*, Structure and Biosynthesis of the Jamaicamides, New Mixed
709    Polyketide-Peptide Neurotoxins from the Marine Cyanobacterium *Lyngbya majuscula.*
710    *Chem. Biol.* **11**, 817–833 (2004).
711 48. Z. Chang, *et al.*, Biosynthetic Pathway and Gene Cluster Analysis of Curacin A, an
712    Antitubulin Natural Product from the Tropical Marine Cyanobacterium *Lyngbya majuscula*
713    †. *J. Nat. Prod.* **67**, 1356–1367 (2004).
714 49. M. A. Matilla, H. Stöckmann, F. J. Leeper, G. P. C. Salmond, Bacterial biosynthetic gene
715    clusters encoding the anti-cancer haterumalide class of molecules: Biogenesis of the
716    broad spectrum antifungal and anti-oomycete compound, oocydin A. *J. Biol. Chem.* **287**,
717    39125–39138 (2012).
718 50. K. M. Fisch, *et al.*, Polyketide assembly lines of uncultivated sponge symbionts from
719    structure-based gene targeting. *Nat. Chem. Biol.* **5**, 494–501 (2009).
720 51. K. Blin, *et al.*, antiSMASH 5.0: updates to the secondary metabolite genome mining
721    pipeline. *Nucleic Acids Res.* **47**, W81–W87 (2019).
722 52. T. A. M. Gulder, M. F. Freeman, J. Piel, The Catalytic Diversity of Multimodular Polyketide
723    Synthases: Natural Product Biosynthesis Beyond Textbook Assembly Rules (2011)

https:/doi.org/10.1007/128_2010_113.

53.   M. J. Bertin, *et al.*, The Phormidolide Biosynthetic Gene Cluster: A *trans*-AT PKS Pathway
Encoding a Toxic Macrocyclic Polyketide. *ChemBioChem* **17**, 164–173 (2016).

54.   C. D. Reeves, *et al.*, Alteration of the substrate specificity of a modular polyketide
synthase acyltransferase domain through site-specific mutations. *Biochemistry* **40**, 15464–
15470 (2001).

55.   G. Yadav, R. S. Gokhale, D. Mohanty, Computational approach for prediction of domain
organization and substrate specificity of modular polyketide synthases. *J. Mol. Biol.* **328**,
335–363 (2003).

56.   B. J. Dunn, K. R. Watts, T. Robbins, D. E. Cane, C. Khosla, Comparative analysis of the
substrate specificity of *trans*- versus *cis*-acyltransferases of assembly line polyketide
synthases. *Biochemistry* **53**, 3796–3806 (2014).

57.   A. L. Díaz-Pérez, C. Díaz-Pérez, J. Campos-García, Bacterial l-leucine catabolism as a
source of secondary metabolites. *Rev. Environ. Sci. Bio/Technology* **15**, 1–29 (2016).

58.   T. Mahmud, *et al.*, A novel biosynthetic pathway providing precursors for fatty acid
biosynthesis and secondary metabolite formation in myxobacteria. *J. Biol. Chem.* **277**,
32768–32774 (2002).

59.   M. Röttig, *et al.*, NRPSpredictor2 - A web server for predicting NRPS adenylation domain
specificity. *Nucleic Acids Res.* **39**, 362–367 (2011).

60.   A. T. Keatinge-Clay, The structures of type I polyketide synthases. *Nat. Prod. Rep.* **29**,
1050 (2012).

61.   Ö. Erol, *et al.*, Biosynthesis of the myxobacterial antibiotic corallopyronin A.
ChemBioChem **11**, 1253–1265 (2010).

62.   C. Zhao, *et al.*, Oxazolomycin biosynthesis in *Streptomyces albus* JA3453 featuring an
"acyltransferase-less" type I polyketide synthase that incorporates two distinct extender
units. *J. Biol. Chem.* **285**, 20097–20108 (2010).

63.   H. Irschik, *et al.*, Analysis of the sorangicin gene cluster reinforces the utility of a combined
phylogenetic/retrobiosynthetic analysis for deciphering natural product assembly by *trans*-
AT PKS. *ChemBioChem* **11**, 1840–1849 (2010).

64.   D. C. Cantu, Y. Chen, P. J. Reilly, Thioesterases: A new perspective based on their
primary and tertiary structures. *Protein Sci.* **19**, 1281–1295 (2010).

65.   J. Moldenhauer, *et al.*, The Final Steps of Bacillaene Biosynthesis in *Bacillus*
amyloliquefaciens FZB42: Direct Evidence for β,γ Dehydration by a *trans*-Acyltransferase
Polyketide Synthase. *Angew. Chemie* **122**, 1507–1509 (2010).

66.   G. Berkhan, C. Merten, C. Holec, F. Hahn, The Interplay between a Multifunctional
Dehydratase Domain and a C-Methyltransferase Effects Olefin Shift in Ambruticin
Biosynthesis. *Angew. Chemie - Int. Ed.* **55**, 13589–13592 (2016).

67.   C. Hertweck, The biosynthetic logic of polyketide diversity. *Angew. Chemie - Int. Ed.* **48**,
4688–4716 (2009).

68.   J. Moldenhauer, X. H. Chen, R. Borriss, J. Piel, Biosynthesis of the antibiotic bacillaene,
the product of a giant polyketide synthase complex of the *trans*-AT family. *Angew. Chemie*
- Int. Ed. **46**, 8195–8197 (2007).

69.   T. Wakimoto, *et al.*, Calyculin biogenesis from a pyrophosphate protoxin produced by a
sponge symbiont. *Nat. Chem. Biol.* **10**, 648–655 (2014).

70.   E. Esquenazi, *et al.*, Visualizing the spatial distribution of secondary metabolites produced
by marine cyanobacteria and sponges via MALDI-TOF imaging. *Mol. Biosyst.* **4**, 562–570
(2008).

71.   J. Lopera, I. J. Miller, K. L. McPhail, J. C. Kwan, Increased Biosynthetic Gene Dosage in a
Genome-Reduced Defensive Bacterial Symbiont. *mSystems* **2** (2017).

72.   T. J. Buchholz, *et al.*, Polyketide β-Branching in Bryostatin Biosynthesis: Identification of
Surrogate Acetyl-ACP Donors for BryR, an HMG-ACP Synthase. *Chem. Biol.* **17**, 1092–
1100 (2010).

73.   Y. Q. Cheng, G. L. Tang, B. Shen, Type I polyketide synthase requiring a discrete
acyltransferase for polyketide biosynthesis. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 3149–

3154 (2003).

74.   K. C. Nicolaou, *et al.*, Chemical synthesis and biological evaluation of palmerolide A
analogues. *J. Am. Chem. Soc.* **130**, 10019–10023 (2008).

75.   S. F. Haydock, *et al.*, Organization of the biosynthetic gene cluster for the macrolide
concanamycin A in *Streptomyces neyagawaensis* ATCC 27449. *Microbiology* **151**, 3161–
3169 (2005).

76.   W. Chen, *et al.*, Characterization of the Polyoxin Biosynthetic Gene Cluster from
Streptomyces cacaoi and Engineered Production of Polyoxin H. *J. Biol. Chem.* **284**,
10627–10638 (2009).

77.   T. K. Mihali, W. W. Carmichael, B. A. Neilan, A putative gene cluster from a *Lyngbya*
wollei bloom that encodes paralytic shellfish toxin biosynthesis. *PLoS One* **6** (2011).

78.   A. K. El-Sayed, J. Hothersall, C. M. Thomas, Quorum-sensing-dependent regulation of
biosynthesis of the polyketide antibiotic mupirocin in *Pseudomonas fluorescens* NCIMB
10586. *Microbiology* **147**, 2127–2139 (2001).

79.   S. Maier, *et al.*, Functional characterization of different ORFs including luciferase-like
monooxygenase genes from the mensacarcin gene cluster. *ChemBioChem* **16**, 1175–
1182 (2015).

80.   G. Lackner, E. E. Peters, E. J. N. Helfrich, J. Piel, Insights into the lifestyle of uncultured
bacterial natural product factories associated with marine sponges. *Proc. Natl. Acad. Sci.*
114, E347–E356 (2017).

81.   J. J. Gehret, *et al.*, Terminal alkene formation by the thioesterase of curacin A
biosynthesis: Structure of a decarboxylating thioesterase. *J. Biol. Chem.* **286**, 14445–
14454 (2011).

82.   L. Gu, *et al.*, Metamorphic enzyme assembly in polyketide diversification. *Nature* **459**,
731–735 (2009).

83.   J. Piel, A polyketide synthase-peptide synthetase gene cluster from an uncultured
bacterial symbiont of Paederus beetles. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 14002–14007
(2002).

84.   T. Stachelhaus, H. D. Mootz, V. Bergendahl, M. A. Marahiel, Peptide Bond Formation in
Nonribosomal Peptide Biosynthesis. *J. Biol. Chem.* **273**, 22773–22781 (1998).

85.   S. Lin, S. G. Van Lanen, B. Shen, A free-standing condensation enzyme catalyzing ester
bond formation in C-1027 biosynthesis. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 4183–4188
(2009).

86.   K. Zaleta-Rivera, *et al.*, A bidomain nonribosomal peptide synthetase encoded by FUM14
catalyzes the formation of tricarballylic esters in the biosynthesis of fumonisins.
Biochemistry **45**, 2561–2569 (2006).

87.   C. M. Theodore, *et al.*, Genomic and metabolomic insights into the natural product
biosynthetic diversity of a feral-hog-associated *Brevibacillus laterosporus* strain. *PLoS*
One **9**, 3–12 (2014).

88.   A. T. Keatinge-Clay, R. M. Stroud, The Structure of a Ketoreductase Determines the
Organization of the β-Carbon Processing Enzymes of Modular Polyketide Synthases.
Structure **14**, 737–748 (2006).

89.   E. J. N. Helfrich, *et al.*, Automated structure prediction of *trans*-acyltransferase polyketide
synthase products. *Nat. Chem. Biol.* **15**, 813–821 (2019).

90.   C. Khosla, Y. Tang, A. Y. Chen, N. A. Schnarr, D. E. Cane, Structure and Mechanism of
the 6-Deoxyerythronolide B Synthase. *Annu. Rev. Biochem.* **76**, 195–221 (2007).

91.   L. M. Quirós, I. Aguirrezabalaga, C. Olano, C. Méndez, J. A. Salas, Two
glycosyltransferases and a glycosidase are involved in oleandomycin modification during
its biosynthesis by *Streptomyces antibioticus*. *Mol. Microbiol.* **28**, 1177–1185 (1998).

92.   T. A. Wencewicz, Crossroads of Antibiotic Resistance and Biosynthesis. *J. Mol. Biol.* **431**,
3370–3399 (2019).

93.   G. Jenkins, E. Cundliffe, Cloning and characterization of two genes from *Streptomyces*
lividans that confer inducible resistance to lincomycin and macrolide antibiotics. *Gene* **108**,
55–62 (1991).

94.   A. Gourmelen, M. H. Blondelet-Rouault, J. L. Pernodet, Characterization of a glycosyl
transferase inactivating macrolides, encoded by gimA from *Streptomyces ambofaciens*.
Antimicrob. Agents Chemother. **42**, 2612–2619 (1998).
95.   K. Yokoyama, H. Imamura, Rotation, structure, and classification of prokaryotic V-
ATPase. *J. Bioenerg. Biomembr.* **37**, 405–410 (2005).
96.   M. Boehm, *et al.*, Structural and mutational analysis of band 7 proteins in the
cyanobacterium *Synechocystis* sp. strain PCC 6803. *J. Bacteriol.* **191**, 6425–6435 (2009).
97.   S. A. Kautsar, *et al.*, MIBiG 2.0: A repository for biosynthetic gene clusters of known
function. *Nucleic Acids Res.* **48**, D454–D458 (2020).

Figures

843
844
845

846
847
Fig. 1. Structure of palmerolide A with notations for the proposed retrobiosynthesis. Backbone
synthesis is a result of incorporation of the starter unit, a glycine residue, and acetate subunits
(C1 indicated by black squares). Structural features from *trans*-acting tailoring enzymes (indicated
by grey ovals) utilize additional substrates: methyl transfers from SAM (purple dots), installation of
C-25 methyl from acetate (blue dot) via an HCS cassette, and carbamoyl transfer to the
secondary alcohol on C-11. The α-hydroxy group on C-10 is predicted to arise from incorporation
of hydroxymalonic acid or a *trans*-acting hydroxylase.
855

856
857
Fig. 2. The proposed BGC for palmerolide A, showing the hybrid PKS-NRPS system. KS:
ketosynthase domain, C: condensation domain, gly: adenylation domain for glycine incorporation,
DH: dehydratase domain, cMT: carbon methyl transferase domain, KR: ketoreductase domain,
DHt: dehydratase variant; ECH: enoyl-CoA hydratase, LLM: luciferase-like monooxygenase, AT:
acyl transferase; polysacc synt_2: polysaccharide biosynthesis protein, LO: lactone oxidase, ABC

trans: ATP-binding cassette transporter, Band7: stomatin-like integral membrane, PPTase:
phosphopantetheinyl transferase, NMO: nitronate monooxygenase, HCS: hydroxymethylglutaryl-
CoA synthase, GTF: glycosyl transferase ER: enoyl reductase, CT: carbamoyl transferase, small
blue circles represent acyl- or peptidyl-carrier proteins. Ppant arms are symbolized by wavy lines.
The grey domains indicate domains that would be expected to perform an enzymatic
transformation; however, are not encoded in the BGC. Blue arrows indicate biosynthetic genes.
Green arrows indicate genes that encode for non-biosynthetic proteins. White arrows reflect
hypothetical genes. The BGC is displayed in reverse compliment.
871

873
874 **Fig. 3.** Comparison of BGC organization of select *trans*-AT systems. There is significant
875 variability in the order of the core modules, AT modules, and modules which contain *trans*-acting
876 tailoring enzymes. There is also variability in the number of encoded AT modules, though the AT
877 modules are typically encoded on separate, but tandem genes if more than one is present.
878
879

880
881
Fig. 4 (a) Comparison of the modular structure of the 5 *pal* BGCs. **(b)** Family of palmerolides.
Much of the structural diversity can be explained by differences due to starter unit promiscuity,
sites of action for the *trans*-acting tailoring enzymes, and differences in the core modules of the
multiple *pal* BGCs. It is proposed that *pal* BGC 4 is responsible for not only palmerolide A, but
also palmerolide B, palmerolide C, palmerolide F, and palmerolide G. It is interesting to note that
the modular structure of the domains responsible for biosynthesis are equivalent for *pal* BGC 1
and *pal* BGC 3. These two BGCs contain an additional KS domain as compared to *pal* BGC 4
and are likely responsible for the biosynthesis of palmerolide D and palmerolide H.