Palmerolide PKS-NRPS gene clusters characterized from the microbiome of an Antarctic ascidian

Nicole E. Avalon;[1] Alison E. Murray;[2] Hajnalka E. Daligault;[3] Chien-Chi Lo;[3] Karen W. Davenport;[3] Armand E.K. Dichosa;[3] Patrick S.G. Chain;[3] Bill J. Baker[1]

[1]Department of Chemistry, University of South Florida, Tampa, FL 33620

[2]Division of Earth and Ecosystem Sciences, Desert Research Institute, Reno, NV 89512

[3]Los Alamos National Laboratory, Los Alamos, NM 87545

Corresponding authors:

Bill J. Baker, Department of Chemistry, University of South Florida, Tampa, FL 33620, USA. +1-813-974-1967, bjbaker@usf.edu

Alison E. Murray, Division of Earth and Ecosystem Sciences, Desert Research Institute, Reno, NV 89512, USA. +1 775-673-7361. alison.murray@dri.edu.

Competing Interest Statement: The authors have no competing interests to disclose.

Classification: Biological sciences, environmental sciences and ecology

Keywords: microbial biosynthesis, secondary metabolites, Antarctic microbiology, *trans*-AT Type I polyketide synthase, marine natural products

Abstract

Polyketides are a complex family of natural products that often serve competitive or pro-survival purposes but can also demonstrate bioactivity in human diseases as, for example cholesterol lowering agents, anti-infectives, or anti-tumor agents. Marine invertebrates and microbes are a rich source of polyketides. Palmerolide A, a polyketide isolated from the Antarctic ascidian *Synoicum adareanum,* is a vacuolar-ATPase inhibitor with potent bioactivity against melanoma cell lines. The biosynthetic gene clusters (BGC) responsible for production of secondary metabolites are encoded in the genomes of the producers as discrete genomic elements. A putative palmerolide BGC was identified from a *S. adareanum* metagenome based on a high degree of congruence with a chemical structure-based retrobiosynthetic prediction. Protein family homology analysis, conserved domain searches, and active site and motif identification were used to confirm the identity and propose the function of the 75 kb *trans*-acyltransferase (AT) polyketide synthase-non-ribosomal synthase (PKS-NRPS) domains responsible for the synthesis of palmerolide A. Though PKS systems often act in a predictable co-linear sequence, this BGC includes multiple *trans*-acting enzymatic domains, a non-canonical condensation termination domain, a bacterial luciferase-like monooxygenase (LLM), and multiple

copies found within the metagenome-assembled genome (MAG) of *Candidatus* Synoicohabitans palmerolidicus. Detailed inspection of the five highly similar pal BGC copies suggests the potential for biosynthesis of other members of the palmerolide chemical family. This is the first delineation of a biosynthetic gene cluster from an Antarctic species. These findings have relevance for fundamental knowledge of PKS combinatorial biosynthesis and could enhance drug development efforts of palmerolide A through heterologous gene expression.

Significance Statement

Complex interactions exist between microbiomes and their hosts. Increasingly, defensive metabolites that have been attributed to host biosynthetic capability are now being recognized as products of associated microbes. These unique metabolites often have bioactivity in targets of human disease and can be purposed as pharmaceuticals. The molecular machinery for production of palmerolide A, a macrolide that is potent and selective against melanoma, was discovered as a genomic cluster in the microbiome of an Antarctic ascidian. Multiple non-identical copies of this genomic information provide clues to differences in specific enzymatic domains and point to Nature's ability to perform combinatorial biosynthesis *in situ*. Harnessing this genetic information may pave a path for development of a palmerolide-based drug.

Introduction

Marine invertebrates such as corals, sponges, mollusks, and ascidians, are known to be a rich source of bioactive compounds (1). Due to their sessile or sluggish nature, chemical defenses such as secondary metabolites are often key to their survival. Many compound classes are represented among benthic invertebrates including terpenes, nonribosomal peptide synthase (NRPS) products, ribosomally synthesized and post-translationally modified peptides (RiPPs), and polyketides. It is estimated that that over 11,000 secondary metabolites from marine and terrestrial environments understood to be of polyketide synthase (PKS) and NRPS origin have been isolated and described (2). Biosynthetic gene clusters (BGCs) exist as a series of genomic elements that encode for megaenzymes responsible for production of these secondary metabolites. BGCs can have distinct nucleotide composition properties such as codon usage and guanine-cytosine content that do not match the remainder of the genome (3, 4), suggesting a mechanism of transfer from organisms that are only distantly related, perhaps even from interkingdom horizontal gene transfer (5, 6). Interestingly, the BGCs for many natural products isolated from marine invertebrates are found in the host-associated microbiota, reflecting the role of these compounds in symbiosis (7).

Polyketides are a complex family of natural products produced by multi-modular PKS enzymes that are related to, but evolutionarily divergent from, fatty acid synthases (8). They often possess long carbon chains with varied degrees of oxidation, can contain aromatic components, and may be either cyclic or linear. It is estimated that of the polyketides that have been isolated and characterized, 1% have potential biological activity against human diseases, making this class of compounds particularly appealing from a drug discovery and development standpoint (9). This potential for use as pharmaceuticals is approximately five times greater than for compounds of all other natural product classes (9). Many polyketides are classified as macrolides, which are large-ring lactones that are pharmaceutically relevant due to a number of biological actions, including, for example, targeting the cytoskeleton, ribosomal protein biosynthesis, and vacuolar type V-ATPases (10–13). V-ATPases are responsible for acidification of cells and organelles via proton transport across membranes, including those of lysosomes, vacuoles, and endosomes. These enzymes appear to have an impact on angiogenesis, apoptosis, cell proliferation, and tumor metatastisis (10). A number of marine macrolides inhibit V-ATPases, including lobatamides, chondropsins, iejimalides, and several of the palmerolides (14–18).

There are three types of PKS systems: Type I, which are comprised of non-iteratively acting modular enzymes that lead to progressive elongation of a polyketide chain, Type II, which contain iteratively acting enzymes that biosynthesize polycyclic aromatic polyketides, and Type III

which possess iteratively-acting homodimeric enzymes that often result in monocyclic or bicyclic aromatic polyketides (19). Type I PKS systems can be subdivided into two groups, depending upon whether the acyl transferase (AT) modules are encoded co-linearly, as in *cis*-AT Type I PKS, or physically distinct from the megaenzyme, as in *trans*-AT Type I PKS. The concept of co-linearity refers to the parallel relationship between the order of enzymatic modules and the functional groups present in the growing polyketide chain (20). In *trans*-AT systems, AT domains may be incorporated in a mosaic fashion through horizontal gene transfer (20). This introduces greater molecular architectural diversity over evolutionary time, as one clade of *trans*-ATs may select for a malonyl-CoA derivative, while the *trans*-AT domains in another clade may select for unusual or functionalized subunits (21). Additionally, recombination, gene duplication, and conversion events can lead to further diversification of the resultant biosynthetic machinery (21). Predictions regarding the intrinsic relationship between a secondary metabolite of interest, the biosynthetic megaenzyme, and the biosynthetic gene cluster (BGC) can be harnessed for natural product discovery and development (22–24).

In the search for new and bioactive chemotypes as inspiration for the next generation of drugs, underexplored ecosystems hold promise as biological and chemical hotspots (25). The vast Southern Ocean comprises one-tenth of the total area of Earth's oceans and is largely unstudied for its chemodiversity. The coastal marine environment of Antarctica experiences seasonal extremes in, for example, ice cover, light field, and food resources. Taken with the barrier to migration imposed by the Antarctic Circumpolar Current and the effects of repeated glaciation events on speciation, a rich and endemic biodiversity has evolved, with consequent potential for new chemodiversity (25–27).

Palmerolide A (**Fig. 1**) is the principal secondary metabolite isolated from *Synoicum adareanum*, an ascidian which can be found in abundance at depths of 10 to 40 m in the coastal waters near Palmer Station, Antarctica (17). Palmerolide A is a macrolide polyketide that possesses potent bioactivity against malignant melanoma cell lines while demonstrating minimal cytotoxicity against other cell lines (17). The NCI's COMPARE algorithm was used to correlate experimental findings with a database for prediction of the biochemical mechanism by identifying the mechanism of action of palmerolide A as a vacuolar-ATPase (V-ATPase) inhibitor (28). Downstream effects of V-ATPase inhibition include an induction of both hypoxia induction factor-1α and autophagy (17, 29). Increased expression of V-ATPase on the surface of metastatic melanoma cells perhaps explain palmerolide A's selectivity for UACC-62 cell lines over all cell types (30). Despite the relatively high concentrations of palmerolide A in the host tissue (0.49–4.06 mg palmerolide A x g^{-1} host dry weight) (31), isolation of palmerolide A from its Antarctic source in mass sufficient for drug development it is not ecologically nor logistically feasible. Synthetic strategies for the synthesis of palmerolide A have been reported (6–13), though a clear pathway to quantities needed for drug development has been elusive. Therefore, there is substantial interest in identifying the biosynthetic gene cluster (BGC) responsible for palmerolide A production.

Our approach to identify the palmerolide BGC (*pal* BGC) began with the characterization of the host-associated microbiome (39). The core microbiome for *Synoicum adareanum*, a persistent cohort of bacteria present in many individual ascidians, was established through analysis of occurrence of distinct amplicon sequence variants (ASV) from the iTag sequencing of the Variable 3-4 regions of the bacterial 16S rRNA (31). This ultimately led to the evaluation of the metagenome and then the assembly of the metagenome assembled genome (MAG) of *Candidatus* Synoicohabitans palmerolidicus, a verrucomicrobium in the family *Opitutaceae* (40). Contained within the genome are five non-identical copies of the *pal* BGC, of which, three correlate with palmerolides, the structures of which have been previously discovered in this macrolide family (**Fig. 1**). Here we report a detailed analysis of the *pal* BGCs.

Results and Discussion

A. Retrobiosynthetic Scheme for Palmerolide A

A retrobiosynthetic scheme of the *pal* BGC was developed based on the chemical structure for palmerolide A, including modules consistent with a hybrid PKS-NRPS with tailoring enzymes for key functional groups (**Fig. 1**). We hypothesized that the initial module would be PKS-like in nature to utilize 3-methylcrotonic acid as the starter unit followed by a NRPS domain for the incorporation of glycine. PKS elongation was predicted to be an 11-step sequence resulting in 22 contiguous carbons. Modifying enzymes that are encoded co-linearly were predicted to create the architectural diversity with olefin placement, reduction of certain carbonyl groups to secondary alcohols, and full reduction of other subunits. In addition, carbon methyltransferases (cMTs) were predicted to catalyze the placement of methyl groups C-26 and C-27 from *S*-adenosylmethionine (SAM).

Several key structural features proposed to result from the action of *trans*-acting enzymes are present. As seen in the kalimantacins (41), the carbamate on C-11 was hypothesized to be installed by a carbamoyl transferase (CT). The C-25 methyl group is located on C-17, a carbon atom in α-position of the acetate subunit. This structural feature suggests its origin from hydroxymethylglutamoyl-CoA Synthetase (HCS) catalysis, rather than the more common SAM-mediated methylation at the acetate β-position, which appears to be the origin of the C-26 and C-27 methyl groups. Precedence for HCS-mediated β-branch formation is found in the biosynthetic pathways for of the jamaicamides, bryostatins, curacin A, oocydin, pederin, and psymberin, among others (42–46). The hydroxy group on C-10, in the α-position of the acetate subunit, was hypothesized to result from incorporation of hydroxymalonic acid or the action of a hydroxylase on a malonyl-CoA monomer.

B. Proposed Biosynthesis and Architecture of the Putative *pal* Biosynthetic Gene Cluster

Synoicum adareanum collected near Palmer Station, Antarctica was processed to separate eukaryotic from prokaryotic cells of the host microbiome. Iterations of metagenomic analysis (40) yielded a large number of BGCs when analyzed via the antiSMASH annotation platform. A candidate cluster with multiple copy numbers was identified and has excellent congruence with retrobiosynthetic predictions, including the key genomic markers outlined above that are likely involved in biosynthesis of palmerolide A. Integration of BGC annotations with information from protein family homology analysis, conserved domain searches, active site and motif identification, supported the hypothesis that this putative 75 kb BGC is responsible for palmerolide A production. The core scaffold was explained by the NRPS and *trans*-AT PKS hybrid system. In addition, each of the tailoring enzymes that are expected for the distinct chemical features are encoded in the genome. The BGC for palmerolide A is comprised of 14 core biosynthetic modules and 25 genes in a single operon of 74,655 bases (**Fig. 2**). Fourteen modules are co-linear and two *trans*-AT domains (modules 15 & 16) follow the core biosynthetic genes. Additional *trans*-acting genes contribute to backbone modifications with at least one gene contributing to post-translational tailoring (**Fig. 3**).

An unusual starter unit and NRPS domains of *palA*. The initial core biosynthetic domains of *palA* (modules 1 and 2) encode for the requisite acyl carrier proteins (ACP) (**Fig. 2**). Encoded in module 1, there are three ACPs in tandem, which could serve to promote an increase in metabolite production (47). The second in series is an ACP-β containing the conserved domain sequence GXDS (48) and is likely the acceptor of a starter unit containing a β-branch, consistent with the proposed starter unit for palmerolide A, 3-methylcrotonic acid. While both *trans*-acting ATs, PalE and PalF, (**Fig. 2**) possess the catalytic active site serine which is key for the proper

positioning of subunit within the hydrophobic cleft of the active site (8, 49), only the first AT, PalE, has a characteristic motif that includes an active site phenylalanine, conferring specificity for malonate selection (50). The AT selecting the methylcrotonic acid starter unit is likely the second of the two *trans*-AT domains (PalF), which lacks definitive specificity for malonyl-CoA. In support, *trans*-ATs have demonstrated affinity for a wider range of substrates than their *cis*-acting counterparts (21, 51). 3-Methylcrotonyl-CoA is an intermediate of branched-chain amino acid catabolism in leucine degradation; intermediates of which can be diverted to secondary metabolite production (52). Interestingly, an upregulation of leucine catabolic pathways can be seen as a mechanism of cold adaption to maintain cell wall fluidity via production of branched chain fatty acids, a factor which may be at play in this Antarctic microorganism (52, 53). Condensation (C) and adenylation (A) domains accompany the ACP in module 2. Signature sequence information and NRPSPredictor2 method (54, 55) of the A domain are consistent with selection of a glycine residue. These domains incorporate the amino acid residue, resulting in the addition of a nitrogen and two carbons in this step of the biosynthesis of palmerolide A.

Contiguous PKS chain and *trans*-acting enzymes at site of action for *palB – palD*. The contiguous carbon backbone of palmerolide A arises from 11 cycles of elongation from a series of modules with a variety of enzymatic domains that include an ACP, KS, and associated genes that establish the oxidation state of each subunit (**Fig 2**). The first module of *palB* (module 3) includes a dehydratase (DH) and cMT domains, a sequence which results in a chain extension modification to an α,β-unsaturated thioester, a result of the action of the encoded DH. The subsequent cMT methylation is consistent with an *S*-adenosylmethionine (SAM)-derived methyl group, as expected for C-27 in palmerolide A. Module 4, spanning the end of *palB* and beginning of *palC*, includes a DH, a ketoreductase (KR), an epimerization (E), and another cMT domain. This cluster of domains is predicted to result in the methyl-substituted conjugated diene of the macrolide tail (C-19 through C-24, C-26 on palmerolide A). The epimerization domain is hypothesized to set the stereochemistry at C-20.

The substrate critical for macrolactonization of the polyketide is the C-19 hydroxy group, a result of the KS, E, and KR domains encoded in module 5 (**Fig. 2**). Additionally, a domain initially annotated as a dehydratase (DHt) at this location may contribute to the final cyclization and release of the molecule from the megaenzyme by assisting the terminal C domain with ring closure (48). This sequence shows some homology with condensation domains and does not possess the hotdog fold that is indicative of canonical dehydratases, and therefore, may not truly represent a DH (56). Alternatively, this domain could be responsible for the olefin shift to the β,γ-position, as seen in bacillaene and ambruticin biosynthesis (57, 58). Finally, the E domain is hypothesized to establish the stereochemistry at C-19.

Only an enoyl-CoA hydratase (ECH) in addition to standard ACP and KS is encoded in module 6 which would lead to a ketone function; based on our retrobiosynthetic analysis, the resulting ketone at C-17 is the necessary substrate for HCS-catalyzed β-branch formation, resulting in the C-25 methyl group on C-17. The HCS cassette (PalK through PalO) is comprised of a series of *trans*-acting domains, including an ACP, an HCS, a free KS, and 2 additional ECH modules (**Fig. 2**). The HCS cassette can act while the elongating chain is tethered to an ACP module, rather than after cyclization and release (59, 60). The two ECHs in the HCS cassette along with the ECH encoded in-line with the core biosynthetic genes would be responsible for isomerization of a terminal methylene to the observed internal olefin. An HCS cassette formed by the combination of a *trans*-KS and at least one ECH module with an HCS domain is reported in several other bacterial BGCs such as bryostatin 1, calyculin A, jamaicamide, mandelalide, phormidolide, and psymberin (43, 46, 48, 61–63). The domain structure for the HCS cassettes has a remarkably high degree of syntony (64).

There is significant similarity in the domain structure of module 7, module 10, and module 13, which each include a KS, DH, and KR (**Fig. 2**). The olefin that arises from the action of module 7, concomitant with carbon chain elongation, is conjugated with the C16-C17 olefin adjacent the C-17 β-branch. Modules 10 and 13 have similar enzymatic composition to 7 and are likely responsible for Δ^8 and Δ^2 olefins. The combination of KR and DH domains are also found in

modules 8 and 12; however, in concert with an unidentified *trans*-acting ER domain, these olefins would be reduced to fully saturated monomeric subunits. There are some examples of *trans*-acting ER domains carrying out this function, including OocU in oocydin, SorN in sorangicin, and MndM in mandelalide (45, 63, 65), while in other systems, such as corallopyronin and leinamycin, the reductions of the olefins are largely unexplained (66, 67). The reduction by a *trans*-acting enzyme often occurs while the elongating polyketide is tethered to the megaenzyme, as evidenced by the downstream specificity of the KS module for Claisen-type condensation with subunits containing single or double bonds (65).

The genetic architecture for the biosynthesis of two functional groups, essential for bioactivity, is encoded in module 9 (**Fig. 2**); structure-activity relationship studies demonstrate the importance of the C-10 hydroxyl group and the C-11 carbamate (68). The KR domain that is present, predicting the C-11 alcohol function, serves as the substrate for the carbamoyl transferase (*palQ*) in a post-translational modification (69–71). Intriguingly, a domain annotated as a luciferase-like monooxygenase (LLM) initially seems out of place; however, palmerolide A has a hydroxy group at C-10, which is the α-position of the acetate subunit inserted in module 9. LLMs associated with BGCs may not serve as true luciferases, but instead demonstrate oxidizing effects on polyketides and peptides without evidence of corresponding bioluminescence (72, 73). For example, there is an overrepresentation of LLMs in *Candidatus* Entotheonella BGCs without known bioluminescence (74). As demonstrated through individual inactivation of the LLM in the BGC of mensacarin, a Type II PKS system, Msn02, Msn04, and Msn08 have key activity as epoxidases and hydroxlases (73). There are several examples of LLMs in modular Type I PKS systems. OnnC from onnamide and NazB from nazumamide are two LLMs in *Candidatus* Endotheonella that are proposed to serve biosynthetically as hydroxylases (74). In calyculin and mandelalide, the CalD and MndB LLMs catalyze chain shortening reactions through α-hydroxylation and Baeyer-Villiger-type oxidation reactions (61, 63). Phormidolide has a LMM that adds a hydroxy group which is then hypothesized to attack an olefin through a Michael-type addition for cyclization with enzymatic assistance from a pyran synthase (48). The hydroxylation that is key in cyclization of oocydin A is likely installed by OocK or OocM, flavin-dependent monooxygenases that are contiguous to the PKS genes and are thought to act while the substrate is bound to a portion of the PKS megaenzyme (45). It is this hydroxylase activity that we propose for the LLM in module 9. Since the producing bacteria is yet to be cultured, it is not established whether this LLM may also serve a role in bioluminescence and/or quorum sensing. Further evidence for the role of the LLM is provided through alignment against other LLM's. The *pal* BGC contains both LuxR family transcriptional regulator as well as the DNA-binding response regulator. In addition to the annotation within Pfam00296, which includes the bacterial LLMs, the sequence is homologous with the multiple sequence alignments and the hidden Markov models of the TIGR subfamily 04020 of the conserved protein domain family cl19096, which is noted for natural product biosynthesis LLMs (74). The subfamily occurs in both NRPS and PKS systems as well as small proteins with binding of either flavin mononucleotide or coenzyme F420. Alignment of the LLMs from multiple PKS systems, including palmerolide A, shows homology with model sequences from the TIGR subfamily 04020 (**Fig. 4**).

The addition of C-5 and C-6 and the reduction of the β-carbonyl to form the C-7 hydroxy group of palmerolide A, is due to module 11, which possesses a KR domain in addition to elongating KS (**Fig. 2**). In the structure of palmerolide A this is followed by the fully reduced subunit from module 12 as discussed above. The final elongation results from module 13, which includes DH and KR domains that contribute to the conjugated ester found as palmerolide A's C-1 through C-3, completing the palmerolide A C_{24} carbon skeleton.

Noncanonical termination condensation domain in *palD* for product cyclization and release. Typically, PKS systems terminate with a thioesterase (TE) domain, leading to release of the polyketide from the megaenzyme (63, 75–77). This canonical domain is not present in the *pal* cluster. Instead, the final module in the *cis*-acting biosynthetic gene cluster includes a truncated condensation domain comprised of 133 amino acid residues; compared to the approximately 450 residues that comprise a standard condensation domain (78) (**Fig. 2**). Condensation domains

catalyze cyclization through ester formation in free-standing condensation domains that act in *trans* as well as in NRPS systems (79, 80). In addition, this non-canonical termination domain is not without precedent in hybrid PKS-NRPS and in PKS systems, as basiliskamide and phormidolide both include condensation domains for product release (48, 81). Though the terminal condensation domain in the *pal* BGC is truncated, it maintains much of the HHXXDDG motif (**Fig. 4**), most notably the second histidine, which serves as the catalytic histidine in the condensation reaction (78).

Stereochemical and structural confirmation based on sequence information. KR domains are NADPH-dependent enzymes that belong to the short-chain dehydrogenase superfamily, with Rossman-like folds for co-factor binding (82, 83). Enzymatically, the two KR subtypes, A-Type and B-type, are responsible for stereoselective reduction of β-keto groups and can also determine the stereochemistry of α-substituents. C-type KRs, however, lack reductase activity and often serve as epimerases. A-Type KRs have a key tryptophan residue in the active site, do not possess the LDD amino acid motif, and result in the reduction of β-carbonyls to $_L$-configured hydroxy groups (83). B-Type, which are identified by the presence of an LDD amino acid motif, result in formation of $_D$-configured hydroxy groups (83). The stereochemistry observed in palmerolide A is reflected in the active site sequence information for the $_D$-configured hydroxyl groups from module 5 and module 11 (**Fig. 1** and **Fig. 2**). When an enzymatically active DH domain is within the same module, the stereochemistry of the *cis*- versus *trans*-olefin can be predicted, as the combination of an A-Type KR with a DH results in a *cis*-olefin formation and the combination of a B-Type KR with a DH results in *trans*-olefin formation. The *trans*-α,β-olefins arising from module 7 (C14-C15), module 10 (C8-C9), and module 13 (C2-C3) are each consistent with the presence of a B-Type KR and an active DH. The other three olefins present in the structure of palmerolide A, as noted above, likely have positional and stereochemical influence from E domains (module 4 and module 5) or an ECH domain (module 6).

Additional structural confirmation was obtained through defining the specificity of KS domains using phylogenetic analysis and the *Trans*-AT PKS Polyketide Predictor (TransATor) bioinformatic tool (84). KS domains catalyze the sequential two-carbon elongation steps through a Claisen-like condensation with a resulting β-keto feature (85). Additional domains within a given module can modify the β-carbonyl or add functionality to the adjacent α- or γ-positions (83). Specificity of KSs, based the types of modification located on the upstream acetate subunit were determined and were found to be mostly consistent with the retrobiosynthetic predictions (***SI Appendix*, Fig. S1,** *SI Appendix*, **Table S1**). For example, the first KS, KS1 (module 1), is predicted to receive a subunit containing a β-branch. KS3 (module 4) and KS4 (module 5) are predicted to receive an upstream monomeric unit with α-methylation and an olefinic shift, consistent with the structure of palmerolide A and with the enzymatic transformations resulting from module 3 and module 4, respectively. Interestingly, the KS associated with the HCS cassette is in a clade of its own in the phylogenetic tree. TransATor also aided in confirming the stereochemical outcomes of the hydroxy groups and olefins, which occur through reduction of the β-carbonyls. The predictions for the $_D$-configured hydroxy groups were consistent with not only the the presence of the LDD motif, indicative of type B-type KR as outlined above, but also with stereochemical determination based on the clades of the KS domains of the receiving modules, KS5 (module 6) and KS11 (module 12). They are also consistent with the structure of palmerolide A. The KS predictions, however, did not aid in confirming reduction of the upstream olefins for KS8 (module 9) and KS12 (module 13).

Additional *trans*-acting domains and domains between genes responsible for biosynthesis. A glycosyl transferase (*pal*P) and lactone oxidase (*palH*) that are associated with glycosylation of polyketides are encoded in the palmerolide A BGC (**Fig. 2**), though glycosylated palmerolides have not been observed. Glycosylation as a means of self-resistance has been described for oleandomycin, which encodes not only two glycosyl transferases, one with substrate specificity toward oleandomycin and the other relaxed specificity, but also a glycosyl hydrolase (86, 87). The glycosylation of the macrolide confers internal resistance to ribosomal

inhibition, making the glycosylated macrolide a pro-drug, which is activated by the glycosyl hydrolase that is excreted extracellularly (86, 87). Glycosyl transferases are found in other macrolide- and non-macrolide-producing organisms as a means to inactivate hydroxylated polyketides (88, 89). Though prokaryotic V-ATPases tend to be simpler than those of eukaryotes, there is homology in the active sites making the pro-drug hypothesis for self-resistance reasonable in palmerolide A biosynthesis (90). The D-arabinono-1,4-lactone oxidase (*palH*) is a FAD-dependent oxidoreductase that likely works in concert with the glycosyltransferase. Additionally, a Band7 protein is encoded. Band7 proteins belong to a ubiquitous family of slipin or stomatin-like integral membrane proteins that are found in all kingdoms, yet lack assignment of function (91). A domain of unknown function (DUF) is annotated; however, this particular DUF, DUF179, has no superfamily relatives in protein homology searches. Together, these *trans*-acting domains decorate the carbon backbone with features that give palmerolide A its unique chemical structure and contribute to its bioactivity.

C. Multiple copies of the *pal* Biosynthetic Gene Cluster Explain Structural Variants in the Palmerolide Family

Careful assembly of the metagenome assembled genome (MAG) of Ca. *S. palmerolidicus* revealed the BGC was present in multiple copies (**Fig. 1** and ***SI Appendix*, Fig. S2-S4**) (40). The structural complexity of the multicopy BGCs represents a biosynthetic system that is similar to that which is found in *Ca.* Didemnitutus mandela, another ascidian-associated verrucomicrobium in the family *Opitutaceae* (63). A total of 5 distinct BGCs consistent with Type-I PKS systems that have much overlap with one another and are likely responsible for much of the structural diversity in the family of palmerolides (17, 18) (**Fig. 1**). Palmerolide A is hypothesized to arise from the BGC designated as *pal* BGC 4 with additional compounds also from this same cluster. The clusters designated as *pal* BGC 1, *pal* BGC 2, *pal* BGC 3, and *pal* BGC 5 have been identified in *Ca.* S. palmerolidicus and the potential products of each are described below. It is hypothesized that there are three levels of diversity introduced to create the family of palmerolides: (1) differences in the site of action for the *trans*-acting domains (with additional *trans*-acting domains at play as well), (2) promiscuity of the initial selection of the starter subunit, and (3) differences in the core biosynthetic genes with additional PKS domains or stereochemical propensities within a module.

There are several palmerolides that likely arise from the same BGC encoding the megaenzyme responsible for palmerolide A (*pal* BGC 4). We hypothesize that the *trans*-acting domains have different sites of action than what is seen in palmerolide A biosynthesis. For example, the chemical scaffold of palmerolide B (**Fig. 1**) is similar to palmerolide A, though the carbamate transfer occurs on the C-7 hydroxy group. Palmerolide B instead bears a sulfate group on the C-11 hydroxy group; proteins with homology to multiple types of sulfatases from the UniProtKB database, (P51691, P15289, O69787, Q8ZQJ2) found in the genome of *Ca.* S. palmerolidicus (40), but are not encoded within the BGCs. One of these *trans*-acting sulfatases likely modifies the molecule post-translationally. Other structural differences including the hydroxylation on C-8 instead of C-10 (as observed in palmerolide A) and the Δ^9 olefin that differs from palmerolide A's Δ^8 olefin, are either due to a difference of the site of action of the LLM (module 9) or a *trans*-acting hydroxylase. Another member of the compound family, palmerolide C, has structural differences attributable to *trans*-acting enzymes as well. Again, a *trans*-acting hydroxylase or the LLM is proposed to be responsible for hydroxylation on C-8. A hydroxy group on C-9 occurs through reduction of the carbonyl. The carbamate installation occurs on C-11 after *trans*-acting hydroxylation or LLM hydroxylation. In addition, the Δ^8 olefin in palmerolide A is not observed, but rather a Δ^6 olefin.

Additional levels of structural variation are seen at the site of the starter unit, likely due to a level of enzymatic promiscuity of the second AT (PalF). This, combined with differences in the sites of action for the trans-acting domains, is likely responsible for the structural differences

observed in palmerolide F (**Fig. 1**). The terminal olefin on the tail of the macrolide which perhaps is a product of promiscuity of the selection of the starter unit, the isomeric 3-methyl-3-butenoic acid, is consistent with the aforementioned lack of consensus for malonate selection by the AT. In addition, the KS that receives the starter unit is phylogenetically distinct from the other KS in the *pal* clusters (**SI Appendix, Fig. S1**).

The retrobiosynthetic hypothesis for Palmerolide G (**Fig. 1**) has much similarity to what is present in *pal* BGC 4; however, the presence of a *cis*-olefin rather than a *trans*-olefin could arise from a difference in the enzymatic activity of module 4. This olefin is the result of a shift from an epimerase and, therefore, the stereochemistry is not solely reliant upon the action of the associated KR. Although this difference has not been identified in the BGCs in the samples sequenced, this could be present in other environmental samples that have been batched for processing and compound isolation. Currently, the biosynthetic mechanism is unknown.

The core biosynthetic genes of two identical palmerolide BGCs (*pal* BGC 1 and *pal* BGC 3) (**SI Appendix, Fig. S2**) possess an additional elongation module when compared to *pal* BGC 4. Palmerolide D (**Fig. 1**) is structurally very similar to palmerolide A with the exception of elongation in the carboxylate tail of the macrolide by an isopropyl group. This could arise from one additional round of starter unit elongation via a KS and methylation. These two identical BGCs are consistent with this structural difference. The overall architecture and stereochemistry are otherwise maintained. Palmerolide H (**Fig. 1**) includes the structural differences of both palmerolide B and palmerolide D. It contains the extended carboxylate tail with a terminal olefin and incorporates hydroxylation on C-8 rather than C-10. Again, there is no genomic evidence that this hydroxylation in the α-position of the acetate subunit is due to incorporation of hydroxy-malonate to explain this but is instead likely due to a *trans*-hydroxylase. The carbamate installation occurs on C-7, while sulfonation occurs on C-11 and α-hydroxy placement is on C-8.

The final two palmerolide BGCs both have truncation of the core biosynthetic genes. The gene structure of *pal* BGC 5 (**SI Appendix, Fig. S3**) shows preservation of genes upstream to the core biosynthetic genes; however, there are no pre-NRP PKS modules noted in the BGC. The HCS cassette, glycosyl transferase, and CT are all present downstream. The predicted product of this cluster does not align with a known palmerolide, though posttranslational hydrolysis of the C-24 amide may result in a structure similar to palmerolide E (**Fig. 1**), which maintains much of the structure of palmerolide A; however, is missing the initial polyketide starter unit and the glycine subunit. The final *pal* BGC in *Ca.* S. palmerolidicus, *pal* BGC 2 (**SI Appendix, Fig. S4**), includes only 5 elongating modules, which would result in a 10-carbon structure that has not been observed. Interestingly, despite the shortened BGC, the HCS cassette, glycosyl transferase, and CT are all present downstream. There would only be a single hydroxy group serving as a substrate for the CT, glycosyl transferase, and sulfatase to act. The 2-carbon site of action for the β-branch introduced in the palmerolide A structures would not be present. The structure-based retrobiosynthesis the eight known palmerolides (A-F) can be hypothesized to arise from differences in the core biosynthetic genes of these non-identical copies of the *pal* BGC, starter unit promiscuity, and differing sites of action in the *trans*-acting enzymes.

Conclusion

The putative *pal* BGC has been identified and represents the first BGC elucidated from an Antarctic organism. As outlined in a retrobiosynthetic strategy, the *pal* BGC represents a *trans*-AT Type I PKS/NRPS hybrid system with compelling alignment to the predicted biosynthetic steps for palmerolide A. The *pal* BGC is proposed to begin with PKS modules resulting in the incorporation of an isovaleric acid derivative, 3-methylcrotonic acid, as a starter unit, followed by incorporation of a glycine residue with NRPS-type modules. Then, eleven rounds of progressive polyketide elongation likely occur leading to varying degrees of oxidation introduced with each module. There are several interesting non-canonical domains encoded within the BGC, such as an HCS, CT, LLM, and a truncated condensation termination domain. Additionally, a glycosylation domain

may be responsible for reversible, pro-drug formation to produce self-resistance to the V-ATPase activity of palmerolide A. There are additional domains, the function of which have yet to be determined.

A combination of modular alterations, starter unit differences, and activity of *trans*-acting enzymes contributes to Nature's production of a suite of palmerolide analogues. There are a total of five distinct *pal* BGCs in the MAG of *Ca.* S. palmerolidicus, yielding the known eight palmerolides, with genetic differences that explain some of the structural variety seen within this family of compounds. These include differences in modules that comprise the core biosynthetic genes. Additionally, it is proposed that some of the architectural diversity of palmerolides arises from different sites of action of the *trans*-acting, or non-colinear, modules. Starter unit promiscuity is another potential source of the structural differences observed in the compounds. Analysis of the *pal* BGC not only provides insight into the architecture of this Type I PKS/NRPS hybrid BGC with unique features, but also lays the foundational groundwork for drug development studies of palmerolide A via heterologous expression.

Materials and Methods

Specimens of *Synoicum adareanum* were collected by SCUBA from the Antarctic Peninsula in the Anvers Island Archipelago and flash frozen. The methods for processing as well as metagenome sequencing and assembly are outlined in detail in two related publications (31, 40). In brief, the first step of data analysis was the result of metagenome sequencing of two Antarctic *S. adareanum* samples (Sample IDs: Nor-2c-2007 and Nor-2a-2007). Ion Proton and 454, two different high throughput DNA sequencing technologies, were utilized. The sequence data was independently assembled using both Newbler (v. 2.9)(92) and SPAdes (v. 3.5) (93). The results were merged via MeGAmerge (v. 1.2) (94). Assembly 1 resulted in 86,387 contigs with a maximum contig size 153,680 and total contig size 144,953,904 bases. The second step in the metagenome sequencing effort was based on two additional *S. adareanum* samples (Bon-1c-2011 and Del-2b-2011) that were also collected by SCUBA and then frozen. These samples were rich in domains for the *pal* BGCs. Six cells of data were obtained from PacBio sequencing, two cells from Bon1C and four cells from Del2B. We generated Circular Consensus Sequences (CCSs) using SMRT Link. The CCS reads were assembled on EDGE Bioinformatics (95) using wtdbg2 (96) to create Assembly 2. The assembled contigs were binned using MaxBin2 (v. 2.2.6) and the binned contigs from different assemblies with 58% GC content were used to re-assemble them via phrap (97), an overlap based assembler. Manual editing of the assembled genome was performed, and CCS reads were used to close gaps and verify repeat regions. We obtained a 4.3 MB genome of palmerolide A producing bacteria in 5 contigs with unique sequences and 5 varying copies of the BGC repeats (referred to as *pal* BGC 1 through *pal* BGC 5).

The assembled metagenome was annotated using antiSMASH (v. 5.0). These annotations were integrated with information from protein family homology analysis, conserved domain searches, active site and motif identification to delineate the biosynthetic steps. Manual interrogations of sequence information of the *pal* BGCs was used to perform BLASTP searches, confirm enzymatic identities, and to identify active site residues key for stereochemical outcomes, substrate affinities, and other granular details. BGCs were obtained from public NCBI databases for basiliskamide, bryostatin 1, calyculin, corallopyronin, mandelalide, onnamide, oxazolamycin, pederin, phormidolide, psymberin, sorangicin, and myxoviricin. ClustalO alignment tool in the CLC Genome Workbench (QIAGEN aarhau A/S v. 20.0.3) was used for multiple sequence alignments of enzymatic domains with the HMM Pfam Seed obtained from EMBL-EBI and the amino acid sequences from the other PKS BGCs. MIBiG (98) was used to acquire the KS amino acid sequence from the type III PKS BGC responsible for 3-(2'-hydroxy-3'-oxo-4'-methylpentyl)-indole biosynthesis from *Xenorhabdus bovienii SS-2004* (GenBank Accession: FN667741.1), which was used for an outgroup. The BGC KSs were numbered according to their position in their proposed biosynthesis in the literature. Prior to the construction of the phylogenetic tree for the

KS domains (***SI Appendix,* Fig. S1**), the sequences in the alignment were manually inspected and trimmed. Phylogenetic trees were created in the CLC Genome Workbench (QUIAGEN aarhau A/S v. 20.0.3) with Neighbour Joining (NJ) as a distance method and Bayesian estimation. Jukes-Cantor was selected for the genetic distance model and bootstrapping with performed with 100 replicates. Additionally, the sequence of each KS in the *pal* BGCs was queried using the *Trans*-AT PKS Polyketide Predictor (TransATor) to help define the specificity of KS domains. The software is based on phylogenetic analyses of fifty-four *trans*-AT type I PKS systems with 655 KS sequences and the resulting clades are referenced to help predict the KS specificity for the upstream unit (84).

Data Desposition

Data from this project has been deposited to the NCBI under BioProject Accession Number PRJNA662631. The GenBank Accession Number for the MAG of *Ca.* Synoicohabitans palmerolidicus is JAGGDC000000000. The BGCs can be accessed in the MiBIG database with the associated Accession Numbers: BGC0002118 (for *pal* BGC_4) and BGC0002119 (for *pal* BGC_3).

Acknowledgments

Support for this research was provided in part by the National Institutes of Health award (CA205932) to A.E.M., B.J.B., and P.S.G.C., with additional support from National Science Foundation awards (ANT-0838776, and PLR-1341339 to B.J.B., ANT-0632389 to A.E.M.) The authors acknowledge the assistance of field team members, including William Dent, Charles D. Amsler, James B. McClintock, Margaret O. Amsler, and Katherine Schoenrock. This work would not have been possible without the outstanding logistical support in Antarctica from the employees and subcontractors of Raytheon Polar Services. Lucas Bishop and Mary L. Higham are also recognized for their contributions.

Author Contributions:

This work was the result of a team effort in which the following contributions are recognized: conceptualization, A.E.M., P.S.G.C., and B.J.B.; methodology and experimentation, N.E.A., A.E.M., H.E.D., C.L., P.S.G.C., B.J.B; data analysis, N.E.A., A.E.M., H.E.D, C.L., K.W.D., A.E.K.D., P.S.G.C, B.J.B; data curation, N.E.A., A.E.M., H.E.D, C.L., P.S.G.C, B.J.B; original draft preparation, N.E.A., B.J.B., A.E.M., H.E.D.; review and editing, N.E.A., B.J.B., A.E.M., C.L., H.E.D., P.S.G.C., A.E.K.D; funding acquisition, A.E.M., P.S.G.C., and B.J.B. All authors have read and agreed to the published version of the manuscript.

References

1.   N. G. Wilson, J. A. Maschek, B. J. Baker, A species flock driven by predation? Secondary metabolites support diversification of slugs in Antarctica. *PLoS One* **8**, 1–7 (2013).
2.   C. A. Dejong, *et al.*, Polyketide and nonribosomal peptide retro-biosynthesis and global gene cluster matching. *Nat. Chem. Biol.* **12**, 1007–1014 (2016).
3.   J. G. Lawrence, H. Ochman, M. A. Ragan, Reconciling the many faces of lateral gene transfer. *Trends Microbiol.* **10**, 1–4 (2002).

4.   M. Ravenhall, N. Škunca, F. Lassalle, C. Dessimoz, Inferring Horizontal Gene Transfer. *PLoS Comput. Biol.* **11**, 1–16 (2015).

5.   I. Schmitt, H. T. Lumbsch, Ancient horizontal gene transfer from bacteria enhances biosynthetic capabilities of fungi. *PLoS One* **4**, 1–8 (2009).

6.   E. W. Schmidt, Trading molecules and tracking targets in symbiotic interactions. *Nat. Chem. Biol.* **4**, 466–473 (2008).

7.   E. W. Schmidt, The secret to a successful relationship: Lasting chemistry between ascidians and their symbiotic bacteria. *Invertebr. Biol.* **134**, 88–102 (2015).

8.   E. J. N. Helfrich, J. Piel, Biosynthesis of polyketides by trans-AT polyketide synthases. *Nat. Prod. Rep.* **33**, 231–316 (2016).

9.   A. M. P. Koskinen, K. Karisalmi, Polyketide stereotetrads in natural products. *Chem. Soc. Rev.* **34**, 677–690 (2005).

10.   J. Napolitano, A. Daranas, M. Norte, J. Fernandez, Marine Macrolides, a Promising Source of Antitumor Compounds. *Anticancer. Agents Med. Chem.* **9**, 122–137 (2012).

11.   R. Ueoka, *et al.*, Metabolic and evolutionary origin of actin-binding polyketides from diverse organisms. *Nat. Chem. Biol.* **11**, 705–712 (2015).

12.   M. E. Bordeleau, *et al.*, Stimulation of mammalian translation initiation factor eIF4A activity by a small molecule inhibitor of eukaryotic translation. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 10460–10465 (2005).

13.   S. Nishimura, *et al.*, Structure-activity relationship study on 13-deoxytedanolide, a highly antitumor macrolide from the marine sponge Mycale adhaerens. *Bioorganic Med. Chem.* **13**, 455–462 (2005).

14.   R. Shen, C. T. Lin, E. J. Bowman, B. J. Bowman, J. A. Porco, Lobatamide C: Total synthesis, stereochemical assignment, preparation of simplified analogues, and V-ATPase inhibition studies. *J. Am. Chem. Soc.* **125**, 7889–7901 (2003).

15.   E. J. Bowman, K. R. Gustafson, B. J. Bowman, M. R. Boyd, Identification of a New Chondropsin Class of Antitumor Compound that Selectively Inhibits V-ATPases. *J. Biol. Chem.* **278**, 44147–44152 (2003).

16.   S. Kazami, *et al.*, Iejimalides show anti-osteoclast activity via V-ATPase inhibition. *Biosci. Biotechnol. Biochem.* **70**, 1364–1370 (2006).

17.   T. Diyabalanage, C. D. Amsler, J. B. McClintock, B. J. Baker, Palmerolide A, a cytotoxic macrolide from the antarctic tunicate Synoicum adareanum. *J. Am. Chem. Soc.* **128**, 5630–5631 (2006).

18.   J. H. Noguez, *et al.*, Palmerolide macrolides from the Antarctic tunicate Synoicum adareanum. *Bioorganic Med. Chem.* **19**, 6608–6614 (2011).

19.   B. Shen, *et al.*, Polyketide biosynthesis beyond the type I, II, and III polyketide synthase paradigms: A progress report. *ACS Symp. Ser.* **955**, 154–166 (2007).

20.   T. A. Nguyen, *et al.*, Exploiting the mosaic structure of trans-acyltransferase polyketide synthases for natural product discovery and pathway dissection. *Nat. Biotechnol.* **26**, 225–233 (2008).

21.   A. Nivina, K. P. Yuet, J. Hsu, C. Khosla, Evolution and Diversity of Assembly-Line Polyketide Synthases. *Chem. Rev.* **119**, 12524–12547 (2019).

22.   P. Videau, K. N. Wells, A. J. Singh, W. H. Gerwick, B. Philmus, Assessment of Anabaena sp. Strain PCC 7120 as a Heterologous Expression Host for Cyanobacterial Natural Products: Production of Lyngbyatoxin A. *ACS Synth. Biol.* **5**, 978–988 (2016).

23.   E. J. Kim, *et al.*, Heterologous Production of 4-O-Demethylbarbamide, a Marine Cyanobacterial Natural Product. *Org. Lett.* **14**, 5824–5827 (2012).

24.   C. Greunke, *et al.*, Direct Pathway Cloning (DiPaC) to unlock natural product biosynthetic potential. *Metab. Eng.* **47**, 334–345 (2018).

25.   J. B. McClintock, C. D. Amsler, B. J. Baker, R. W. M. Van Soest, Ecology of antarctic marine sponges: An overview. *Integr. Comp. Biol.* **45**, 359–368 (2005).

26.   A. Clarke, J. A. Crame, Evolutionary dynamics at high latitudes: Speciation and extinction in polar marine faunas. *Philos. Trans. R. Soc. B Biol. Sci.* **365**, 3655–3666 (2010).

27.   R. M. Young, *et al.*, Site-specific variability in the chemical diversity of the Antarctic red

alga Plocamium cartilagineum. *Mar. Drugs* **11**, 2126–2139 (2013).

28. K. D. Paull, E. Hamel, L. Malspeis, Prediction of Biochemical Mechanism of Action from the In Vitro Antitumor Screen of the National Cancer Institute. *Cancer Chemother. Agents*, 9–45 (1995).

29. K. Von Schwarzenberg, *et al.*, Mode of cell death induction by pharmacological vacuolar H +-ATPase (V-ATPase) inhibition. *J. Biol. Chem.* **288**, 1385–1396 (2013).

30. T. Diyabalanage, Chemical Investigation of two Antarctic Invertebrates, Synoicum adareanum (Chordata: Ascidiaceae; Enterogona; Polyclinidae) and Austrodoris kergulenensis (Molusca; Gastropoda; Nudibranchia; Dorididae). 211 (2006).

31. A. E. Murray, *et al.*, Uncovering the Core Microbiome and Distribution of Palmerolide in Synoicum adareanum Across the Anvers Island Archipelago, Antarctica. *Mar. Drugs* **18**, 298 (2020).

32. K. P. Kaliappan, P. Gowrisankar, Synthetic studies on a marine natural product, palmerolide A: Synthesis of C1-C9 and C15-C21 fragments. *Synlett*, 1537–1540 (2007).

33. M. P. Lisboa, D. M. Jones, G. B. Dudley, Formal synthesis of palmerolide A, featuring alkynogenic fragmentation and syn-selective vinylogous aldol chemistry. *Org. Lett.* **15**, 886–889 (2013).

34. M. D. Lebar, B. J. Baker, Synthesis of the C3-14 fragment of palmerolide A using a chiral pool based strategy. *Tetrahedron* **66**, 1557–1562 (2010).

35. Z. K. Wen, Y. H. Xu, T. P. Loh, Palladium-catalyzed cross-coupling of unactivated alkenes with acrylates: Application to the synthesis of the C13-C21 fragment of palmerolide A. *Chem. - A Eur. J.* **18**, 13284–13287 (2012).

36. S. A. Pujari, P. Gowrisankar, K. P. Kaliappan, A shimizu non-aldol approach to the formal total synthesis of palmerolide A. *Chem. - An Asian J.* **6**, 3137–3151 (2011).

37. K. R. Prasad, A. B. Pawar, Stereoselective synthesis of C1-C18 region of palmerolide A from tartaric acid. *Synlett* **4**, 1093–1095 (2010).

38. A. B. Pawar, K. R. Prasad, Formal total synthesis of palmerolide A. *Chem. - A Eur. J.* **18**, 15202–15206 (2012).

39. C. S. Riesenfeld, A. E. Murray, B. J. Baker, Characterization of the microbial community and polyketide biosynthetic potential in the palmerolide-producing tunicate Synoicum adareanum. *J. Nat. Prod.* **71**, 1812–1818 (2008).

40. A. E. Murray, *et al.*, Discovery of an Antarctic ascidian-associated uncultivated verrucomicrobium that encodes the biosynthetic capability for antimelanoma palmerolides. **(In prep.)** (2021).

41. W. Mattheus, *et al.*, Isolation and Purification of a New Kalimantacin/Batumin-Related Polyketide Antibiotic and Elucidation of Its Biosynthesis Gene Cluster. *Chem. Biol.* **17**, 149–159 (2010).

42. S. Sudek, *et al.*, Identification of the putative bryostatin polyketide synthase gene cluster from "Candidatus Endobugula sertula", the uncultivated microbial symbiont of the marine bryozoan Bugula neritina. *J. Nat. Prod.* **70**, 67–74 (2007).

43. D. J. Edwards, *et al.*, Structure and Biosynthesis of the Jamaicamides, New Mixed Polyketide-Peptide Neurotoxins from the Marine Cyanobacterium Lyngbya majuscula. *Chem. Biol.* **11**, 817–833 (2004).

44. Z. Chang, *et al.*, Biosynthetic Pathway and Gene Cluster Analysis of Curacin A, an Antitubulin Natural Product from the Tropical Marine Cyanobacterium Lyngbya m ajuscula †. *J. Nat. Prod.* **67**, 1356–1367 (2004).

45. M. A. Matilla, H. Stöckmann, F. J. Leeper, G. P. C. Salmond, Bacterial biosynthetic gene clusters encoding the anti-cancer haterumalide class of molecules: Biogenesis of the broad spectrum antifungal and anti-oomycete compound, oocydin A. *J. Biol. Chem.* **287**, 39125–39138 (2012).

46. K. M. Fisch, *et al.*, Polyketide assembly lines of uncultivated sponge symbionts from structure-based gene targeting. *Nat. Chem. Biol.* **5**, 494–501 (2009).

47. T. A. M. Gulder, M. F. Freeman, J. Piel, "The Catalytic Diversity of Multimodular Polyketide Synthases: Natural Product Biosynthesis Beyond Textbook Assembly Rules" in

(2011) https:/doi.org/10.1007/128_2010_113.

48. M. J. Bertin, *et al.*, The Phormidolide Biosynthetic Gene Cluster: A trans -AT PKS Pathway Encoding a Toxic Macrocyclic Polyketide. *ChemBioChem* **17**, 164–173 (2016).

49. C. D. Reeves, *et al.*, Alteration of the substrate specificity of a modular polyketide synthase acyltransferase domain through site-specific mutations. *Biochemistry* **40**, 15464–15470 (2001).

50. G. Yadav, R. S. Gokhale, D. Mohanty, Computational approach for prediction of domain organization and substrate specificity of modular polyketide synthases. *J. Mol. Biol.* **328**, 335–363 (2003).

51. B. J. Dunn, K. R. Watts, T. Robbins, D. E. Cane, C. Khosla, Comparative analysis of the substrate specificity of trans - Versus cis- acyltransferases of assembly line polyketide synthases. *Biochemistry* **53**, 3796–3806 (2014).

52. A. L. Díaz-Pérez, C. Díaz-Pérez, J. Campos-García, Bacterial l-leucine catabolism as a source of secondary metabolites. *Rev. Environ. Sci. Bio/Technology* **15**, 1–29 (2016).

53. T. Mahmud, *et al.*, A novel biosynthetic pathway providing precursors for fatty acid biosynthesis and secondary metabolite formation in myxobacteria. *J. Biol. Chem.* **277**, 32768–32774 (2002).

54. M. Röttig, *et al.*, NRPSpredictor2 - A web server for predicting NRPS adenylation domain specificity. *Nucleic Acids Res.* **39**, 362–367 (2011).

55. K. Blin, *et al.*, antiSMASH 5.0: updates to the secondary metabolite genome mining pipeline. *Nucleic Acids Res.* **47**, W81–W87 (2019).

56. D. C. Cantu, Y. Chen, P. J. Reilly, Thioesterases: A new perspective based on their primary and tertiary structures. *Protein Sci.* **19**, 1281–1295 (2010).

57. J. Moldenhauer, *et al.*, The Final Steps of Bacillaene Biosynthesis in Bacillus amyloliquefaciens FZB42: Direct Evidence for β,γ Dehydration by a trans-Acyltransferase Polyketide Synthase. *Angew. Chemie* **122**, 1507–1509 (2010).

58. G. Berkhan, C. Merten, C. Holec, F. Hahn, The Interplay between a Multifunctional Dehydratase Domain and a C-Methyltransferase Effects Olefin Shift in Ambruticin Biosynthesis. *Angew. Chemie - Int. Ed.* **55**, 13589–13592 (2016).

59. C. Hertweck, The biosynthetic logic of polyketide diversity. *Angew. Chemie - Int. Ed.* **48**, 4688–4716 (2009).

60. J. Moldenhauer, X. H. Chen, R. Borriss, J. Piel, Biosynthesis of the antibiotic bacillaene, the product of a giant polyketide synthase complex of the trans-AT family. *Angew. Chemie - Int. Ed.* **46**, 8195–8197 (2007).

61. T. Wakimoto, *et al.*, Calyculin biogenesis from a pyrophosphate protoxin produced by a sponge symbiont. *Nat. Chem. Biol.* **10**, 648–655 (2014).

62. E. Esquenazi, *et al.*, Visualizing the spatial distribution of secondary metabolites produced by marine cyanobacteria and sponges via MALDI-TOF imaging. *Mol. Biosyst.* **4**, 562–570 (2008).

63. J. Lopera, I. J. Miller, K. L. McPhail, J. C. Kwan, Increased Biosynthetic Gene Dosage in a Genome-Reduced Defensive Bacterial Symbiont. *mSystems* **2** (2017).

64. T. J. Buchholz, *et al.*, Polyketide β-Branching in Bryostatin Biosynthesis: Identification of Surrogate Acetyl-ACP Donors for BryR, an HMG-ACP Synthase. *Chem. Biol.* **17**, 1092–1100 (2010).

65. H. Irschik, *et al.*, Analysis of the sorangicin gene cluster reinforces the utility of a combined phylogenetic/retrobiosynthetic analysis for deciphering natural product assembly by trans-AT PKS. *ChemBioChem* **11**, 1840–1849 (2010).

66. Ö. Erol, *et al.*, Biosynthesis of the myxobacterial antibiotic corallopyronin A. *ChemBioChem* **11**, 1253–1265 (2010).

67. Y. Q. Cheng, G. L. Tang, B. Shen, Type I polyketide synthase requiring a discrete acyltransferase for polyketide biosynthesis. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 3149–3154 (2003).

68. K. C. Nicolaou, *et al.*, Chemical synthesis and biological evaluation of palmerolide A analogues. *J. Am. Chem. Soc.* **130**, 10019–10023 (2008).

69. S. F. Haydock, *et al.*, Organization of the biosynthetic gene cluster for the macrolide concanamycin A in Streptomyces neyagawaensis ATCC 27449. *Microbiology* **151**, 3161–3169 (2005).

70. W. Chen, *et al.*, Characterization of the Polyoxin Biosynthetic Gene Cluster from Streptomyces cacaoi and Engineered Production of Polyoxin H. *J. Biol. Chem.* **284**, 10627–10638 (2009).

71. T. K. Mihali, W. W. Carmichael, B. A. Neilan, A putative gene cluster from a Lyngbya wollei bloom that encodes paralytic shellfish toxin biosynthesis. *PLoS One* **6** (2011).

72. A. K. El-Sayed, J. Hothersall, C. M. Thomas, Quorum-sensing-dependent regulation of biosynthesis of the polyketide antiobiotic mupirocin in Pseudomonas fluorescens NCIMB 10586. *Microbiology* **147**, 2127–2139 (2001).

73. S. Maier, *et al.*, Functional characterization of different ORFs including luciferase-like monooxygenase genes from the mensacarcin gene cluster. *ChemBioChem* **16**, 1175–1182 (2015).

74. G. Lackner, E. E. Peters, E. J. N. Helfrich, J. Piel, Insights into the lifestyle of uncultured bacterial natural product factories associated with marine sponges. *Proc. Natl. Acad. Sci.* **114**, E347–E356 (2017).

75. J. J. Gehret, *et al.*, Terminal alkene formation by the thioesterase of curacin A biosynthesis: Structure of a decarboxylating thioesterase. *J. Biol. Chem.* **286**, 14445–14454 (2011).

76. L. Gu, *et al.*, Metamorphic enzyme assembly in polyketide diversification. *Nature* **459**, 731–735 (2009).

77. J. Piel, A polyketide synthase-peptide synthetase gene cluster from an uncultured bacterial symbiont of Paederus beetles. *Proc. Natl. Acad. Sci. U. S. A.* **99**, 14002–14007 (2002).

78. T. Stachelhaus, H. D. Mootz, V. Bergendahl, M. A. Marahiel, Peptide Bond Formation in Nonribosomal Peptide Biosynthesis. *J. Biol. Chem.* **273**, 22773–22781 (1998).

79. S. Lin, S. G. Van Lanen, B. Shen, A free-standing condensation enzyme catalyzing ester bond formation in C-1027 biosynthesis. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 4183–4188 (2009).

80. K. Zaleta-Rivera, *et al.*, A bidomain nonribosomal peptide synthetase encoded by FUM14 catalyzes the formation of tricarballylic esters in the biosynthesis of fumonisins. *Biochemistry* **45**, 2561–2569 (2006).

81. C. M. Theodore, *et al.*, Genomic and metabolomic insights into the natural product biosynthetic diversity of a feral-hog-associated Brevibacillus laterosporus strain. *PLoS One* **9**, 3–12 (2014).

82. A. T. Keatinge-Clay, R. M. Stroud, The Structure of a Ketoreductase Determines the Organization of the β-Carbon Processing Enzymes of Modular Polyketide Synthases. *Structure* **14**, 737–748 (2006).

83. A. T. Keatinge-Clay, The structures of type I polyketide synthases. *Nat. Prod. Rep.* **29**, 1050 (2012).

84. E. J. N. Helfrich, *et al.*, Automated structure prediction of trans-acyltransferase polyketide synthase products. *Nat. Chem. Biol.* **15**, 813–821 (2019).

85. C. Khosla, Y. Tang, A. Y. Chen, N. A. Schnarr, D. E. Cane, Structure and Mechanism of the 6-Deoxyerythronolide B Synthase. *Annu. Rev. Biochem.* **76**, 195–221 (2007).

86. L. M. Quirós, I. Aguirrezabalaga, C. Olano, C. Méndez, J. A. Salas, Two glycosyltransferases and a glycosidase are involved in oleandomycin modification during its biosynthesis by Streptomyces antibioticus. *Mol. Microbiol.* **28**, 1177–1185 (1998).

87. T. A. Wencewicz, Crossroads of Antibiotic Resistance and Biosynthesis. *J. Mol. Biol.* **431**, 3370–3399 (2019).

88. G. Jenkins, E. Cundliffe, Cloning and characterization of two genes from Streptomyces lividans that confer inducible resistance to lincomycin and macrolide antibiotics. *Gene* **108**, 55–62 (1991).

89. A. Gourmelen, M. H. Blondelet-Rouault, J. L. Pernodet, Characterization of a glycosyl

transferase inactivating macrolides, encoded by gimA from Streptomyces ambofaciens. *Antimicrob. Agents Chemother.* **42**, 2612–2619 (1998).

90. K. Yokoyama, H. Imamura, Rotation, structure, and classification of prokaryotic V-ATPase. *J. Bioenerg. Biomembr.* **37**, 405–410 (2005).
91. M. Boehm, *et al.*, Structural and mutational analysis of band 7 proteins in the cyanobacterium Synechocystis sp. strain PCC 6803. *J. Bacteriol.* **191**, 6425–6435 (2009).
92. M. J. Chaisson, P. A. Pevzner, Short read fragment assembly of bacterial genomes. *Genome Res.* **18**, 324–330 (2008).
93. S. Nurk, *et al.*, Assembling single-cell genomes and mini-metagenomes from chimeric MDA products. *J. Comput. Biol.* **20**, 714–737 (2013).
94. M. Scholz, C. C. Lo, P. S. G. Chain, Improved assemblies using a source-agnostic pipeline for metagenomic assembly by merging (MeGAMerge) of contigs. *Sci. Rep.* **4**, 1–9 (2014).
95. P. E. Li, *et al.*, Enabling the democratization of the genomics revolution with a fully integrated web-based bioinformatics platform. *Nucleic Acids Res.* **45**, 67–80 (2017).
96. J. Ruan, H. Li, Fast and accurate long-read assembly with wtdbg2. *Nat. Methods* **17**, 155–158 (2020).
97. B. Ewing, P. Green, Base-calling of automated sequencer traces using phred. II. Error probabilities. *Genome Res.* **8**, 186–194 (1998).
98. S. A. Kautsar, *et al.*, MIBiG 2.0: A repository for biosynthetic gene clusters of known function. *Nucleic Acids Res.* **48**, D454–D458 (2020).

Fig. 1. (a) Structure of palmerolide A with notations for the proposed retrobiosynthesis. Backbone synthesis is a result of incorporation of the starter unit, a glycine residue, and acetate subunits (C1 indicated by black squares). Structural features from *trans*-acting tailoring enzymes (indicated by grey ovals) utilize additional substrates: methyl transfers from SAM (purple dots), installation of C-25 methyl from acetate (blue dot) via HCS cassette, and carbamoyl transfer to the secondary alcohol on C-11. The α-hydroxy group on C-10 from incorporation of hydroxymalonic acid or a *trans*-acting hydroxylase. **(b)** Family of palmerolides. Much of the structural diversity can be explained by differences due to starter unit promiscuity, sites of action for the *trans*-acting tailoring enzymes, and differences in the core modules of the multiple pal BGCs.

Fig. 2. The proposed BGC for palmerolide A, showing the hybrid PKS-NRPS system. KS: ketosynthase domain, C: condensation domain, gly: adenylation domain for glycine incorporation, DH: dehydratase domain, cMT: carbon methyl transferase domain, KR: ketoreductase domain, E: epimerization domain, DHt: dehydratase variant; ECH: enoyl-CoA hydratase, LLM: luciferase-like monooxygenase, AT: acyl transferase; polysacc synt_2: polysaccharide biosynthesis protein, LO: lactone oxidase, Band7: stomatin-like integral membrane, PPTase: phosphopantetheinyl transferase, NMO: nitronate monooxygenase, HCS: hydroxymethylglutaryl-CoA synthase, GTF: glycosyl transferase ER: enoyl reductase, CT: carbamoyl transferase, small blue circles represent acyl- or peptidyl-carrier proteins. Blue arrows indicate biosynthetic genes. White arrows reflect hypothetical genes. The BGC is displayed in reverse compliment.

Fig. 3. Comparison of BGC organization of select *trans*-AT systems. There is significant variability in the order of the core modules, AT modules, and modules which contain *trans*-acting tailoring enzymes. There is also variability in the number of encoded AT modules, though the AT modules are typically encoded on separate, but tandem genes if more than one is present.

Fig. 4. (a) Alignment of the proposed palmerolide hydroxylase luciferase-like monooxygenase (LLM) with sequences from the TIGR subfamily 04020. **(b)** Alignment of the proposed termination condensation domain of the *pal* BGC_4 with that of basiliskamide and phormidolide as well as the HMM seed sequences for PF00668 (condensation domains); conservation of the HHXXXDG motif can be seen.