Construction of continuously expandable single-cell atlases through integration of heterogeneous datasets in a generalized cell-embedding space

Lei Xiong[1,2,4], Kang Tian[1,2,4], Yuzhe Li[1,3], Qiangfeng Cliff Zhang[1,2,*]

[1] MOE Key Laboratory of Bioinformatics, Beijing Advanced Innovation Center for Structural Biology & Frontier Research Center for Biological Structure, Center for Synthetic and Systems Biology, School of Life Sciences, Tsinghua University, Beijing, China 100084

[2] Tsinghua-Peking Center for Life Sciences, Beijing, China 100084

[3] Academy for Advanced Interdisciplinary Studies, Peking University, Beijing, China 100871

[4] Co-first authorship

[*] Correspondence: qczhang@tsinghua.edu.cn (Q.C.Z.)

ABSTRACT

Single-cell RNA-seq and ATAC-seq analyses have been widely applied to decipher cell-type and regulation complexities. However, experimental conditions often confound biological variations when comparing data from different samples. For integrative single-cell data analysis, we have developed SCALEX, a deep generative framework that maps cells into a generalized, batch-invariant cell-embedding space. We demonstrate that SCALEX accurately and efficiently integrates heterogenous single-cell data using multiple benchmarks. It outperforms competing methods, especially for datasets with partial overlaps, accurately aligning similar cell populations while retaining true biological differences. We demonstrate the advantages of SCALEX by constructing continuously expandable single-cell atlases for human, mouse, and COVID-19, which were assembled from multiple data sources and can keep growing through the inclusion of new incoming data. Analyses based on these atlases revealed the complex cellular

landscapes of human and mouse tissues and identified multiple peripheral immune

subtypes associated with COVID-19 disease severity.

INTRODUCTION

Single-cell RNA sequencing (scRNA-seq) and assay for transposase-accessible

chromatin using sequencing (scATAC-seq) technologies enable decomposition of

diverse cell-types and states to elucidate their function and regulation in tissues and

heterogeneous systems[1-4]. Efforts like the Human Cell Atlas project[5] and Tabula Muris

Consortium[6] are constructing a single-cell reference landscape for a new era of highly

resolved cell research. With the explosive accumulation of single-cell studies,

integrative analysis of data from experiments of different contexts is essential for

characterizing heterogenous cell populations[7]. However, potentially informative

biological insights are often confounded by batch effects that reflect different donors,

conditions, and/or analytical platforms[8,9].

Integration methods have been developed to remove batch effects in single-cell

datasets[10-16]. One common strategy is to identify similar cells or cell populations across

batches. This includes the mutual nearest neighborhood (MNN) method[10] which

identifies correspondent pairs of cells between two batches by searching for mutual

nearest neighbors in gene expression. Scanorama[11] generalizes the process of neighbor

searching from within two batches to a multiple-batch manner. Seurat v2[13] applies

canonical correlation analysis (CCA) to identify common cell populations in

low-dimensional embeddings across data batches, while Seurat v3[14] introduces "cell

anchors" to mitigate the problem of mixing non-overlapping populations, an issue

experienced in Seurat v2. Harmony[16] also applies population matching across batches,

specifically through a fuzzy clustering algorithm.

It is notable that all of these cell similarity-based methods are local-based,

wherein cell-correspondence across batches are identified through the similarity of

individual cells or cell anchors/clusters. Accordingly, these methods all suffer from

two common limitations. First, they are prone to mixing cell populations that only exist

in some batches. This becomes a severe problem for the integration of datasets that

contain non-overlapping cell populations in each batch (*i.e.*, partially-overlapping data).

Second, these methods can only remove batch effects from the current batches being

assessed but cannot manage batch effects from additional, subsequently obtained

batches. So each time a new batch is added, it requires an entirely new integration

process that again examines the previous batches. This severely limits the capacity to

integrate new single-cell sequencing datasets.

As an alternative to the cell similarity-based local methods, scVI[17] applies a

conditional variational autoencoder (VAE)[18] framework to model the inherent

distribution/structure of the input single-cell data. VAE is a deep generative method

that comprises an encoder and a decoder, wherein the encoder projects all

high-dimensional input data into a low-dimensional embedding, and the decoder

recovers them back to the original data space. The VAE framework can maintain the

same global internal data structure between the high- and low-dimensional spaces[19].

However, scVI includes a set of batch-conditioned parameters into its encoder that

restrains the encoder from learning a batch-invariant embedding space, limiting its

generalizability with new batches.

We previously applied VAE and designed SCALE (Single-Cell ATAC-seq

Analysis via Latent feature Extraction) to model and analyze single-cell ATAC-seq

data[20]. We found that the VAE framework in SCALE can disentangle cell-type-related

and batch-related features in a low-dimensional embedding space. Here, having

redesigned the VAE framework, we introduce SCALEX as a method for integration of

heterogeneous single-cell data. We demonstrate that SCALEX integration is accurate,

scalable, and computationally efficient for multiple benchmark datasets from

scRNA-seq and scATAC-seq studies. As a specific advantage, SCALEX accomplishes

data integration through projecting all single-cell data into a generalized

cell-embedding space using a batch-free encoder and a batch-specific decoder. Since

the encoder is trained to only preserve batch-invariant biological variations, the

resulting cell-embedding space is a generalized one, *i.e.*, common to all projected data.

SCALEX is therefore able to accurately integrate partially-overlapping datasets

without mixing of non-overlapping cell populations. By design, SCALEX runs very

efficiently on huge datasets. These two advantages make SCALEX especially useful

for the construction and research utilization of large-scale single-cell atlas studies,

based on integrating data from heterogeneous sources. New data can be projected to

augment an existing atlas, enabling continuous expansion and improvement of an atlas.

We demonstrated these functionalities of SCALEX in the construction and analyses of

atlases for human, mouse, and COVID-19 PBMCs.

RESULTS

Projecting single-cell data into a generalized cell-embedding space

The central goal of single-cell data integration is to identify and align similar cells

across different batches, while retaining true biological variations within and across

cell-types. The fundamental concept underlying SCALEX is disentangling

batch-related components away from batch-invariant components of single-cell data

and projecting the batch-invariant components into a generalized, batch-invariant

cell-embedding space. To accomplish this, SCALEX implements a batch-free encoder

and a batch-specific decoder in an asymmetric VAE framework[18] (Fig. 1a. Methods).

While the batch-free encoder extracts only biological-related latent features (z) from

input single-cell data (x), the batch-specific decoder is responsible for reconstructing

the original data from z by incorporating batch information back during data

reconstruction.

Supplying batch information to the decoder in data reconstruction allows the

encoder to learn a batch-invariant data representation for each individual cell during

model training, which, as a whole, defines a generalized low-dimensional

cell-embedding space. This learning is also facilitated by random slicing of all input

single cells from different batches into mini-batches. Each mini-batch is forced into

alignment with the same data distribution under the restriction of KL-divergence in the

same cell-embedding space[21]. SCALEX also implements Domain-Specific Batch

Normalization (DSBN)[22] (Methods), a multi-branch Batch Normalization[23], in its

decoder to support incorporation of batch-specific variations to reconstruct single-cell

data.

The design underlying SCALEX renders the encoder to function as a data projector

that projects single cells of different batches into a generalized, batch-invariant

cell-embedding space. SCALEX thus removes batch-related variations present in

single-cell data while preserving batch-invariant biological signals in cell-embedding,

making it an enabling tool for integration analyses of diverse single cell datasets,

without relying on searching for cell similarities.

SCALEX integration is accurate, scalable, and accommodates diverse data types

We first evaluated the data integration performance of SCALEX on multiple

well-curated scRNA-seq datasets, including human *pancreas* (eight batches of five

studies)[24-28], *heart* (two batches of one study)[29] and *liver* (two studies)[30,31]; as well as

human non-small-cell lung cancer (*NSCLC*, four studies)[32-35] and peripheral blood

mononuclear cell (*PBMC*; two batches assayed by two different protocols)[13]. For

comparison, we included several other methods in the analyses, including Seurat v3,

Harmony, Conos, BBKNN, MNN, Scanorama, and scVI (Methods).

We used Uniform Manifold Approximation and Projection (UMAP)[36] embeddings
to visualize the integration performance of all methods (Methods). Note that all of the
raw datasets displayed strong batch effects: cell-types that were common in different
batches were separately distributed. Overall, SCALEX, Seurat v3, and Harmony
achieved the best integration performance for most of the datasets by merging common
cell-types across batches while keeping disparate cell-types apart (Fig. S1). MNN and
Conos integrated many datasets but left some common cell populations not well
aligned. BBKNN, Scanorama, and scVI often had unmerged common cell-types, and
sometimes incorrectly mixed distinct cell-types together. For example, in the *PMBC*
dataset (Fig. 1b), considering the T cell populations between the two batches, while
SCALEX, Seurat v3, Harmony, and MMN integrations were effective, Scanorama
showed both a larger misalignment and mixed all cell-types together without
maintaining clear boundaries.

We quantified single-cell data integration performance using a silhouette score[37]
and a batch entropy mixing score[10] (Methods). Briefly, the silhouette score assesses the
separation of biological distinctions, and the batch entropy mixing score evaluates the
extent of mixing of cells across batches. Overall, SCALEX outperformed all of the
other methods as assessed by the silhouette score, and tied with Seurat and Harmony as
the best-performing methods based on the batch entropy mixing score (Fig. 1c). We
note that SCALEX obtained a slightly lower batch entropy mixing score, compared to
Seurat v3 and Harmony on the *liver* dataset, which contains batch-specific cell-types
and thus is a partially-overlapping dataset. However, Seurat v3 and Harmony may
have obtained a high batch entropy mixing score because of misaligning different
cell-types together. Indeed, by only considering the degree of batch mixing but
ignoring cell-type differences, the batch entropy mixing score is not ideally suited for
assessing batch mixing for partially-overlapping datasets.

We also tested the scalability and computation efficiency of SCALEX on

large-scale datasets by applying it to 1,369,619 cells from the *human fetal atlas* dataset

(two data batches, Methods)[38,39]. SCALEX accurately integrated these two batches,

showing good alignment of the same cell-types (Fig. S2, Fig. 1d). We then compared

the computational efficiency of different methods using down-sampled datasets (of 10

K, 50 K, 250 K, 1 M) from the *human fetal atlas* dataset. SCALEX consumed almost

constant runtime and memory that increased only linearly with data size, whereas MNN,

Seurat v3, and Conos consumed runtime and memory that increased exponentially, thus

did not scale well beyond 250 K cells. Harmony consumed over 400 gigabytes (GB) of

memory in analyzing the 1 M dataset, rendering it unsuitable for integration of datasets

at this scale (Fig. 1e). Notably, the deep learning framework of SCALEX enables it to

run very efficiently on GPU devices, requiring much reduced runtime (took about 10

minutes and 16 GB of memory on the 1 M dataset).

Finally, SCALEX can be used to integrate scATAC-seq data as well as

cross-modality data (*e.g.* scRNA-seq and scATAC-seq) (Methods). For example,

SCALEX integrated the mouse brain scATAC-seq dataset (two batches assayed by

snATAC and 10X)[40] very well, aligning common cell subpopulations and separate

distinct ones (Fig. 1f). We also integrated the cross-modality PBMC data between

scRNA-seq and scATAC-seq[41,42], and found that SCALEX could correctly integrate

the two types of data, and could distinguish rare cells that are specific to scRNA-seq

data, including pDC and platelet cells (Fig. 1g). Thus, SCALEX has broad integration

capacity across various types of single-cell data.

SCALEX integrates partially-overlapping datasets

Partially-overlapping datasets present a major challenge for single-cell data integration

for local cell similarity-based methods[13,14], often leading to over-correction (*i.e.*,

mixing of distinct cell-types). As a global integration method that project cells into a

generalized cell-embedding space, SCALEX is expected to be immune to this problem.

For example, the *liver* dataset is a partially-overlapping dataset where the hepatocyte

population contains multiple subtypes specific to different batches: three subtypes are

specific to LIVER_GSE124395, and two other subtypes only appear in

LIVER_GSE115469 (Fig. S3). We noticed that SCALEX maintained the five

hepatocyte subtypes apart, whereas Seurat v3 mixed all five and Harmony mixed the

hepatocyte-SCD and hepatocyte-TAT-AS1 cells (Fig. 2a).

To characterize the performance of SCALEX on partially-overlapping datasets, we

constructed test datasets with a range of common cell-types, down-sampled from the

six major cell-types in the *pancreas* dataset (Methods). SCALEX integration was

accurate for all cases, aligning the same cell-types without over-correction, whereas

both Seurat v3 and Harmony frequently mixed the cell-types, particularly for the

low-overlapping cases (Fig. 2b, Fig. S4). When there was none common cell-type, both

Seurat v3 and Harmony collapsed the six cell-types to three, mixing alpha with gamma

cells, beta with delta cells, and acinar with ductal cells in various extent. We repeated

the cell-type down-sampling analysis from the 12 cell-types in the *PBMC* dataset as a

more complex partial-overlapping example and observed similar results (Fig. S5),

demonstrating that SCALEX is robust in retaining informative biological variations for

partially-overlapping datasets.

Projection of unseen data into an existing cell-embedding space

The accurate, scalable, and efficient integration performance of SCALEX depends on

its encoder's capacity to project cells from various sources into a generalized,

batch-invariant cell-embedding space. We speculate that once a cell-embedding space

has been constructed after integration of existing data, SCALEX should be able to use

the same encoder to project additional (*i.e.*, previously unseen) data onto the same

embedding space. To test this hypothesis, we used the *pancreas* dataset. SCALEX

integration removed the strong batch effect in the raw data and aligned the same

cell-types together and kept different cell-types were clearly distinguished (Fig. 3a, Fig.

S6a). Cell-types were validated by the expression of their canonical markers, including

rare cells such as Schwann cells, epsilon cells (Fig. S6b).

We projected three new batches[43-45] for pancreas tissues (Fig. 3b) into this

"pancreas cell space" using the same encoder trained on the *pancreas* dataset. After

projection, most of the cells in the new batches were accurately aligned to the correct

cell-types in the pancreas cell space, enabling their accurate annotation by cell-type

label transfer (Fig. 3c, Method). We benchmarked annotation accuracy by calculating

the adjusted Rand Index (ARI)[46], the Normalized Mutual Information (NMI)[47], and the

F1 score using the cell-type information in the original studies as a gold standard

(Methods). The SCALEX annotations achieved the highest accuracy in comparisons

with annotations using three other methods (Seurat v3, Conos, and scmap).

Expanding an existing cell space by including new data

The ability to project new single-cell data into a generalized cell-embedding space

allows SCALEX to readily extend this cell space. To verify this, we projected two

additional melanoma data batches (SKCM_GSE72056, SKCM_GSE123139)[48,49] onto

the previously constructed PBMC space. The common cell-types were correctly

projected onto the same locations in the PBMC cell space (Fig. 3d). For the tumor and

plasma cells only present in the melanoma data batches, SCALEX did not project these

cells onto any existing cell populations in the PBMC space; rather, it projected them

onto new locations close to similar cells, with the plasma cells projected to a location

near B cells, and the tumor cells projected to a location near HSC cells (Fig. 3e).

SCALEX projection enables *post hoc* annotation of unknown cell-types in the

existing cell space using new data. We noted a group of cells previously

uncharacterized in the *pancreas* dataset (Fig. 3a). We found that these cells displayed

high expression levels for known epithelial genes (Methods). We therefore assembled a

collection of epithelial cells from the *bronchial epithelium* dataset[50]. We then projected

these epithelial cells onto the pancreas cell space and found that a group of

antigen-presenting airway epithelial (SLC16A7+ epithelial) cells were projected onto

the same location of the uncharacterized cells (Fig. 3f). This, together with the

observation that both cell populations showed similar marker gene expression (Fig. 3g),

indicates that these uncharacterized cells are also SLC16A7+ epithelial cells. SCALEX

thus enables discovery science in cell biology by supporting exploratory analysis with

large numbers of diverse datasets.

SCALEX supports construction of expandable single-cell atlases

The ability to combine partially-overlapping data onto a generalized cell-embedding

space makes SCALEX a powerful tool to construct a single-cell atlas from a collection

of diverse and large datasets. We applied SCALEX integration to two large and

complex datasets—the *mouse atlas* dataset (comprising multiple organs from two

studies assayed by 10X, Smart-seq2, and Microwell-seq[6,51]) (Fig. 4a) and the *human*

atlas dataset (comprising multiple organs from two studies assayed by 10X and

Microwell-seq[39,52]).

Despite the strong batch effects in the raw data, SCALEX integrated the three

batches of the *mouse atlas* dataset into a unified cell-embedding space (Fig. 4b,c, Fig.

S7a). Common cell-types (including both B, T, and endothelial cells in all tissues and

proximal tubule, urothelial, and hepatocytic cells in certain tissues) were well-aligned

together at the same position in the cell space. Non-overlapping cell-types (such as

sperm, Leydig, and small intestine cells from the Microwell-seq data, keratinocyte stem

cells and large intestine cells in the Smart-seq2 data, and oligodendrocytes in the

Smart-seq2 and Microwell-seq data) were located separately in the space, indicating

that biological variations were preserved well (Fig S7b).

Importantly, atlases generated with SCALEX can be used and further expanded by

projecting new single-cell data to support comparative studies of cells both in the

original atlas and in the new data. Illustrating this, we projected two additional data

batches of aged mouse tissues from *Tabula Muris Senis* (Smart-seq2 and 10X) [53] and

two single tissue datasets (lung and kidney)[54] onto the SCALEX mouse atlas space. We

found that the same cell-types in the new data batches were correctly projected onto the

same locations on the cell-embedding space of the initial mouse atlas (Fig. 4d), which

was also confirmed by the accurate cell-type annotations for the new data by label

transfer from the corresponding cell-types in the initial atlas (Fig. 4e. Methods). On one

way, this mouse atlas then can be used to accurately identify/characterize the cells in

the new data based on their projected locations in the cell space; and on the other way,

projection of new data enables ongoing (and informative) expansion of an existing

atlas.

Following the same strategy, we also constructed a human atlas by SCALEX

integration of multiple tissues from two studies (GSE134255, GSE159929) (Fig. S8a,b).

SCALEX, effectively eliminated the batch effects in the original data and integrated the

two datasets in a unified cell-embedding space (Fig. S8c,d). Again, we were able to

correctly project two additional human skin datasets (GSE130973, GSE147424)[55,56]

onto the human atlas cell-embedding space (Fig. S8e), and again accurately annotated

these projected skin cells (Fig. S8f. Methods). These results illustrate that: i) SCALEX

enables researchers to evaluate their project-specific single cell datasets by leveraging

existing information in large-scale (and ostensibly well annotated) cell atlases; and ii) it

also enables atlas creators to informatively integrate new datasets and attendant

biological insights from many research programs.

An integrative SCALEX COVID-19 PBMC atlas

Many single-cell studies have been conducted to analyze COVID-19 patient immune

responses[57-64]. However, these studies often suffer from small sample size and/or

limited sampling of various disease states[58,64]. For a comprehensive study, we collected

data from multiple COVID-19 PBMC studies, involving 860,746 single cells, and 10

batches from 9 studies[57-63] (Fig. 5a, Fig. S9a), and used SCALEX to generate a

COVID-19 PBMC atlas, identifying 22 cell-types, each of which were supported by

canonical marker gene expression (Fig. 5b,c, Fig. S9b,c. Methods). Cells across

different studies were integrated accurately with the same cell-types aligned together,

confirming integration performance of SCALEX (Fig. 5c, Fig. S9d).

We observed that some cell subpopulations were differentially associated with

patient status (Fig 5d). A subpopulation of CD14 monocytes (CD14-ISG15-Mono),

specifically associated with COIVD-19 patients, was characterized by its high

expression of Type I interferon-stimulated genes (ISGs) and genes associated with

immune-response-related GO terms (Fig 5e,f). The frequency of CD14-ISG15-Mono

cells increased significantly from healthy donors to mild/moderate and severe patients

(Fig. 6g, Fig. S9e. Methods). Within the COVID-19 patients, we observed a significant

decrease in ISG gene expression in CD14-ISG15-Mono cells between the

mild/moderate and severe cases, indicating apparently dysfunctional anti-viral immune

response in severe COVID-19 patients (Fig. 5e). Specifically enriched in severe verse

mild/moderate patients, a neutrophil subpopulation (NCF1-Immature_Neutrophil)

lacked expression of the genes responsible for neutrophil activation but showed

elevated expression of genes associated with viral-process-related GO terms (Fig.

S10a,b). Also enriched in severe patients, a plasma cell subpopulation (MZB1-Plasma)

cells displayed decreased expression for antibody production and were enriched for GO

terms of immune and inflammatory responses (Fig. S10c,d). Thus, the SCALEX

COVID-19 PBMC atlas, generated by integrating a highly diverse collection of

single-cell data from individual studies, identified multiple immune cells-types

showing dysregulations during COVID-19 disease progression. Note that these trends

could not have been detected in the small-scale, individual studies that served as the

basis for our SCALEX COVID-19 PBMC atlas.

Comparative analysis of the SCALEX COVID-19 PBMC atlas and the SC4

consortium study

Recently, a large-scale effort of the Single Cell Consortium for COVID-19 in China

(SC4) has generated a single-cell atlas that contains over 1 million cells (including

PBMCs and other tissues) from 171 COVID-19 patients and 25 healthy controls[65] (Fig.

S11a). We projected the consortium dataset into the cell-embedding space of the

SCALEX COVID-19 PBMC atlas, and found that the cell-types of two atlases were

well-aligned in the embedding space (Fig. 5h,i, Fig. S11b,c).

Our analysis, based on the SCALEX COVID-19 PBMC atlas, yielded findings

consistent with two conclusions from the SC4 study[65]. First, in both analyses diverse

immune subpopulations displayed differential associations with COVID-19 severity.

The proportions of CD14 monocytes, megakaryocytes, plasma cells, and pro T cells

were elevated with increasing disease severity, while the proportion of pDC and mDC

cells decreased (Fig. 5g). Second, we confirmed that the megakaryocytes and monocyte

populations are associated with cytokine storms triggered by SARS-Cov2 infection and

are further elevated in severe patients[66], based on calculating the same cytokine score

and inflammatory score (defined in the SC4 study) for the cells of our SCALEX

COVID-19 PBMC atlas (Fig. 5j. Methods).

Integration of the SC4 data further substantially improved both the scope and

resolution of the SCALEX COVID-19 PBMC atlas. First, this data added macrophages

and epithelial cells to the cell space, enabling investigation of their potential

involvement in COVID-19. The integration also supported more precise

characterization of specific cell subpopulations. For example, the megakaryocyte

population, not distinguished in either single atlas, could be divided into two

subpopulations in the combined atlas (Fig. 5h). An exploratory functional analysis of

the differentially expressed genes in these two newly delineated megakaryocyte

subpopulations (TUBA8-Mega and IGKC-Mega, Fig. S11d,e) revealed enrichment for

the GO terms "humoral immune response" for IGKC-Mega cells yet enrichment for

"negative regulation of platelet activation" for TUBA8-Mega cells (Fig. 5k). These

results illustrate how the continuously expandable single-cell atlases generated using

SCALEX capitalize on existing large-scale data resources and also facilitate discovery

of biological and biomedical insights.

DISCUSSION

SCALEX provides a VAE framework for integration of heterogeneous single-cell data

by disentangling batch-invariant components from batch-related variations and

projecting the batch-invariant components into a generalized, low-dimensional

cell-embedding space. By design, SCALEX models the inherent batch-invariant

patterns of single-cell data, distinguishing it from previously reported integration

methods based on cell similarities. SCALEX does not rely on the identification of

common cell-types across batches, and therefore avoids the problem of cell-type

over-correction, a severe problem for partially-overlapping datasets. SCALEX thus

also overcomes issues of computational complexity in cell similarity-based methods;

that is, the computational time required to identify similar cells may increase

exponentially as the cell number increases.

These two features make SCALEX particularly useful for construction and

integrative analysis of large-scale single-cell atlases based on very heterogenous data

(*i.e.*, datasets acquired by different labs and using different single-cell analysis

platforms). Our construction of human, mouse, and COVID-19 patient single-cell

atlases—which aligned well with previously reported atlases generated from

coordinated large-scale consortium efforts—demonstrates the particular ability of

SCALEX to producing large-scale atlases from extant small-scale datasets. SCALEX

achieves data integration by projecting all single cells into a generalized

cell-embedding space using a universal data projector (*i.e.*, the encoder). This data

projector only needs to be trained once, and then can be used without retraining to

continuously integrate new incoming data into an existing single-cell atlas. This

continuous growth ability makes a SCALEX atlas an elastic resource, allowing the

integration of many single-cell studies to support ongoing, very large-scale research

programs throughout the life sciences and biomedicine.

While the number of single-cell studies is increasing enormously each year, best

practices for experimental design and sample processing are not established, and there

is no obviously dominant data-acquisition platform. SCALEX's ability to

informatively combine data from heterogenous studies and platforms makes it

particularly suitable for the current era of single-cell biological research. Finally, the

ability to conduct exploratory analysis within a generalized cell space supports that

SCALEX should be particularly useful for large-scale integrative (*e.g.*, pan-cancer)

studies. We speculate that use of SCALEX to project single-cell datasets (including

for example scATAC-seq and scRNA-seq) from highly diverse cancer types to

construct a pan-cancer single-cell atlas may lead to the discovery of previously

unknown cell types that are common to divergent carcinomas and that function in

pathogenesis, malignant progression, and/or metastasis.

REFERENCES

1   Kolodziejczyk, A. A., Kim, J. K., Svensson, V., Marioni, J. C. & Teichmann,
S. A. The technology and biology of single-cell RNA sequencing. *Mol Cell* **58**,
610-620, doi:10.1016/j.molcel.2015.04.005 (2015).
2   Papalexi, E. & Satija, R. Single-cell RNA sequencing to explore immune cell
heterogeneity. *Nat Rev Immunol* **18**, 35-45, doi:10.1038/nri.2017.76 (2018).
3   Tanay, A. & Regev, A. Scaling single-cell genomics from phenomenology to
mechanism. *Nature* **541**, 331-338, doi:10.1038/nature21350 (2017).

4   Satpathy, A. T. *et al.* Massively parallel single-cell chromatin landscapes of
human immune cell development and intratumoral T cell exhaustion. *Nature*
Biotechnology **37**, 925-936, doi:10.1038/s41587-019-0206-z (2019).
5   Regev, A. *et al.* The Human Cell Atlas. *Elife* **6**, doi:10.7554/eLife.27041
(2017).
6   Tabula Muris, C. *et al.* Single-cell transcriptomics of 20 mouse organs creates
a Tabula Muris. *Nature* **562**, 367-372, doi:10.1038/s41586-018-0590-4 (2018).
7   Lahnemann, D. *et al.* Eleven grand challenges in single-cell data science.
Genome Biol **21**, 31, doi:10.1186/s13059-020-1926-6 (2020).
8   Leek, J. T. *et al.* Tackling the widespread and critical impact of batch effects
in high-throughput data. *Nat Rev Genet* **11**, 733-739, doi:10.1038/nrg2825
(2010).
9   Hicks, S. C., Townes, F. W., Teng, M. & Irizarry, R. A. Missing data and
technical variability in single-cell RNA-sequencing experiments. *Biostatistics*
19, 562-578, doi:10.1093/biostatistics/kxx053 (2018).
10   Haghverdi, L., Lun, A. T. L., Morgan, M. D. & Marioni, J. C. Batch effects in
single-cell RNA-sequencing data are corrected by matching mutual nearest
neighbors. *Nat Biotechnol* **36**, 421-427, doi:10.1038/nbt.4091 (2018).
11   Hie, B., Bryson, B. & Berger, B. Efficient integration of heterogeneous
single-cell transcriptomes using Scanorama. *Nat Biotechnol*,
doi:10.1038/s41587-019-0113-3 (2019).
12   Polanski, K. *et al.* BBKNN: fast batch alignment of single cell transcriptomes.
Bioinformatics **36**, 964-965, doi:10.1093/bioinformatics/btz625 (2020).
13   Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating
single-cell transcriptomic data across different conditions, technologies, and
species. *Nat Biotechnol* **36**, 411-420, doi:10.1038/nbt.4096 (2018).
14   Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**,
1888-1902 e1821, doi:10.1016/j.cell.2019.05.031 (2019).
15   Barkas, N. *et al.* Joint analysis of heterogeneous single-cell RNA-seq dataset
collections. *Nat Methods* **16**, 695-698, doi:10.1038/s41592-019-0466-z
(2019).
16   Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data
with Harmony. *Nat Methods*, doi:10.1038/s41592-019-0619-0 (2019).
17   Lopez, R., Regier, J., Cole, M. B., Jordan, M. I. & Yosef, N. Deep generative
modeling for single-cell transcriptomics. *Nat Methods* **15**, 1053-1058,
doi:10.1038/s41592-018-0229-2 (2018).
18   Kingma, D. P. & Welling, M. Auto-Encoding Variational Bayes.
arXiv:1312.6114 (2013).
19   Rezende, D. J., Mohamed, S. & Wierstra, D. Stochastic Backpropagation and
Approximate Inference in Deep Generative Models. doi:abs/ (2014).

435	20	Xiong, L. *et al.* SCALE method for single-cell ATAC-seq analysis via latent
436		feature extraction. *Nat Commun* **10**, 4576, doi:10.1038/s41467-019-12630-7
437		(2019).

20 Xiong, L. *et al.* SCALE method for single-cell ATAC-seq analysis via latent
feature extraction. *Nat Commun* **10**, 4576, doi:10.1038/s41467-019-12630-7
(2019).

21 Kullback, S. & Leibler, R. A. On Information and Sufficiency. *Ann. Math.*
Statist. **22**, 79-86, doi:10.1214/aoms/1177729694 (1951).

22 Chang, W.-G., You, T., Seo, S., Kwak, S. & Han, B. Domain-Specific Batch
Normalization for Unsupervised Domain Adaptation. *arXiv:1906.03950*
(2019).

23 Ioffe, S. & Szegedy, C. Batch Normalization: Accelerating Deep Network
Training by Reducing Internal Covariate Shift. *arXiv:1502.03167* (2015).

24 Lawlor, N. *et al.* Single-cell transcriptomes identify human islet cell signatures
and reveal cell-type-specific expression changes in type 2 diabetes. *Genome*
Res **27**, 208-222, doi:10.1101/gr.212720.116 (2017).

25 Segerstolpe, A. *et al.* Single-Cell Transcriptome Profiling of Human
Pancreatic Islets in Health and Type 2 Diabetes. *Cell Metab* **24**, 593-607,
doi:10.1016/j.cmet.2016.08.020 (2016).

26 Muraro, M. J. *et al.* A Single-Cell Transcriptome Atlas of the Human Pancreas.
Cell Syst **3**, 385-394 e383, doi:10.1016/j.cels.2016.09.002 (2016).

27 Grun, D. *et al.* De Novo Prediction of Stem Cell Identity using Single-Cell
Transcriptome Data. *Cell Stem Cell* **19**, 266-277,
doi:10.1016/j.stem.2016.05.010 (2016).

28 Baron, M. *et al.* A Single-Cell Transcriptomic Map of the Human and Mouse
Pancreas Reveals Inter- and Intra-cell Population Structure. *Cell Syst* **3**,
346-360 e344, doi:10.1016/j.cels.2016.08.011 (2016).

29 Litvinukova, M. *et al.* Cells of the adult human heart. *Nature* **588**, 466-472,
doi:10.1038/s41586-020-2797-4 (2020).

30 Aizarani, N. *et al.* A human liver cell atlas reveals heterogeneity and epithelial
progenitors. *Nature* **572**, 199-204, doi:10.1038/s41586-019-1373-2 (2019).

31 MacParland, S. A. *et al.* Single cell RNA sequencing of human liver reveals
distinct intrahepatic macrophage populations. *Nat Commun* **9**, 4383,
doi:10.1038/s41467-018-06318-7 (2018).

32 Lambrechts, D. *et al.* Phenotype molding of stromal cells in the lung tumor
microenvironment. *Nat Med* **24**, 1277-1289, doi:10.1038/s41591-018-0096-5
(2018).

33 Song, Q. *et al.* Dissecting intratumoral myeloid cell plasticity by single cell
RNA-seq. *Cancer Med* **8**, 3072-3085, doi:10.1002/cam4.2113 (2019).

34 Zilionis, R. *et al.* Single-Cell Transcriptomics of Human and Mouse Lung
Cancers Reveals Conserved Myeloid Populations across Individuals and
Species. *Immunity* **50**, 1317-1334 e1310, doi:10.1016/j.immuni.2019.03.009
(2019).

35   Kim, N. *et al.* Single-cell RNA sequencing demonstrates the molecular and
cellular reprogramming of metastatic lung adenocarcinoma. *Nat Commun* **11**,
2285, doi:10.1038/s41467-020-16164-1 (2020).
36   McInnes, L., Healy, J. & Melville, J. UMAP: Uniform Manifold
Approximation and Projection for Dimension Reduction. *arXiv:1802.03426*
(2018).
37   Rousseeuw, P. J. Silhouettes: A graphical aid to the interpretation and
validation of cluster analysis. *Journal of Computational and Applied*
Mathematics **20**, 53-65, doi:https://doi.org/10.1016/0377-0427(87)90125-7
(1987).
38   Cao, J. *et al.* A human cell atlas of fetal gene expression. *Science* **370**,
doi:10.1126/science.aba7721 (2020).
39   Han, X. *et al.* Construction of a human cell landscape at single-cell level.
Nature **581**, 303-309, doi:10.1038/s41586-020-2157-4 (2020).
40   Fang, R. *et al.* Comprehensive analysis of single cell ATAC-seq data with
SnapATAC. *Nat Commun* **12**, 1337, doi:10.1038/s41467-021-21583-9 (2021).
41   Genomics, X. 10k Peripheral blood mononuclear cells (PBMCs) from a
healthy donor, Single Cell ATAC Dataset by Cell Ranger 1.0.1.   (2018).
42   Genomics, X. 10k PBMCs from a Healthy Donor (v3 chemistry), Single Cell
Gene Expression Dataset by Cell Ranger 3.0.0.   (2018).
43   Wang, Y. J. *et al.* Single-Cell Transcriptomics of the Human Endocrine
Pancreas. *Diabetes* **65**, 3028-3038, doi:10.2337/db16-0405 (2016).
44   Enge, M. *et al.* Single-Cell Analysis of Human Pancreas Reveals
Transcriptional Signatures of Aging and Somatic Mutation Patterns. *Cell* **171**,
321-330 e314, doi:10.1016/j.cell.2017.09.004 (2017).
45   Xin, Y. *et al.* Pseudotime Ordering of Single Human beta-Cells Reveals States
of Insulin Production and Unfolded Protein Response. *Diabetes* **67**, 1783-1794,
doi:10.2337/db18-0365 (2018).
46   Hubert, L. & Arabie, P. Comparing partitions. *Journal of Classification* **2**,
193-218, doi:10.1007/BF01908075 (1985).
47   Amelio, A. & Pizzuti, C. in *Proceedings of the 2015 IEEE/ACM International*
Conference on Advances in Social Networks Analysis and Mining 2015
1584–1585 (Association for Computing Machinery, Paris, France, 2015).
48   Tirosh, I. *et al.* Dissecting the multicellular ecosystem of metastatic melanoma
by single-cell RNA-seq. *Science* **352**, 189-196, doi:10.1126/science.aad0501
(2016).
49   Li, H. *et al.* Dysfunctional CD8 T Cells Form a Proliferative, Dynamically
Regulated Compartment within Human Melanoma. *Cell* **176**, 775-789.e718,
doi:https://doi.org/10.1016/j.cell.2018.11.043 (2019).
50   Plasschaert, L. W. *et al.* A single-cell atlas of the airway epithelium reveals
the CFTR-rich pulmonary ionocyte. *Nature* **560**, 377-381,
doi:10.1038/s41586-018-0394-6 (2018).

51   Han, X. *et al.* Mapping the Mouse Cell Atlas by Microwell-Seq. *Cell* **172**,
1091-1107 e1017, doi:10.1016/j.cell.2018.02.001 (2018).

52   He, S. *et al.* Single-cell transcriptome profiling of an adult human cell atlas of
15 major organs. *Genome Biol* **21**, 294, doi:10.1186/s13059-020-02210-0
(2020).

53   Tabula Muris, C. A single-cell transcriptomic atlas characterizes ageing
tissues in the mouse. *Nature* **583**, 590-595, doi:10.1038/s41586-020-2496-1
(2020).

54   Kimmel, J. C. *et al.* Murine single-cell RNA-seq reveals cell-identity- and
tissue-specific trajectories of aging. *Genome Res* **29**, 2088-2103,
doi:10.1101/gr.253880.119 (2019).

55   Sole-Boldo, L. *et al.* Single-cell transcriptomes of the human skin reveal
age-related loss of fibroblast priming. *Commun Biol* **3**, 188,
doi:10.1038/s42003-020-0922-4 (2020).

56   He, H. *et al.* Single-cell transcriptome analysis of human skin identifies novel
fibroblast subpopulation and enrichment of immune subsets in atopic
dermatitis. *J Allergy Clin Immunol* **145**, 1615-1628,
doi:10.1016/j.jaci.2020.01.042 (2020).

57   Schulte-Schrepping, J. *et al.* Severe COVID-19 Is Marked by a Dysregulated
Myeloid Cell Compartment. *Cell* **182**, 1419-1440.e1423,
doi:10.1016/j.cell.2020.08.001 (2020).

58   Lee, J. S. *et al.* Immunophenotyping of COVID-19 and influenza highlights
the role of type I interferons in development of severe COVID-19. *Sci*
Immunol **5**, doi:10.1126/sciimmunol.abd1554 (2020).

59   Wilk, A. J. *et al.* A single-cell atlas of the peripheral immune response in
patients with severe COVID-19. *Nat Med* **26**, 1070-1076,
doi:10.1038/s41591-020-0944-y (2020).

60   Guo, C. *et al.* Single-cell analysis of two severe COVID-19 patients reveals a
monocyte-associated and tocilizumab-responding cytokine storm. *Nat*
Commun **11**, 3924, doi:10.1038/s41467-020-17834-w (2020).

61   Yao, C. *et al.* Cell-Type-Specific Immune Dysregulation in Severely Ill
COVID-19 Patients. *Cell Rep* **34**, 108590, doi:10.1016/j.celrep.2020.108590
(2021).

62   Zhang, J. Y. *et al.* Single-cell landscape of immunological responses in
patients with COVID-19. *Nat Immunol* **21**, 1107-1118,
doi:10.1038/s41590-020-0762-x (2020).

63   Ballestar, E. *et al.* Single cell profiling of COVID-19 patients: an international
data resource from multiple tissues. *medRxiv*, 2020.2011.2020.20227355,
doi:10.1101/2020.11.20.20227355 (2020).

64   Bernardes, J. P. *et al.* Longitudinal Multi-omics Analyses Identify Responses
of Megakaryocytes, Erythroid Cells, and Plasmablasts as Hallmarks of Severe

558   COVID-19. *Immunity* **53**, 1296-1314 e1299,
559   doi:10.1016/j.immuni.2020.11.017 (2020).

560   65   Ren, X. *et al.* COVID-19 immune features revealed by a large-scale single-cell
561   transcriptome atlas. *Cell*, doi:10.1016/j.cell.2021.01.053 (2021).

562   66   Chen, G. *et al.* Clinical and immunological features of severe and moderate
563   coronavirus disease 2019. *J Clin Invest* **130**, 2620-2629,
564   doi:10.1172/JCI137244 (2020).

565   67   Kingma, D. P. & Ba, J. Adam: A Method for Stochastic Optimization.
566   *arXiv:1412.6980* (2014).

567   68   Danese, A., Richter, M. L., Fischer, D. S., Theis, F. J. & Colomé-Tatché, M.
568   EpiScanpy: integrated single-cell epigenomic analysis. *bioRxiv*,
569   doi:10.1101/648097 (2019).

570   69   Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene
571   expression data analysis. *Genome Biol* **19**, 15,
572   doi:10.1186/s13059-017-1382-0 (2018).

573   70   Stuart, T., Srivastava, A., Lareau, C. & Satija, R. Multimodal single-cell
574   chromatin analysis with Signac. *bioRxiv*, doi:10.1101/2020.11.09.373613
575   (2020).

576   71   Abdi, H. & Williams, L. J. Principal component analysis. *WIREs Comput. Stat.*
577   **2**, 433-459, doi:10.1002/wics.101 (2010).

578   72   Traag, V. A., Waltman, L. & van Eck, N. J. From Louvain to Leiden:
579   guaranteeing well-connected communities. *Scientific Reports* **9**, 5233,
580   doi:10.1038/s41598-019-41695-z (2019).

581
582

 Figures

Fig. 1 | The design and performance of SCALEX for single-cell data integration.

a, SCALEX models the global structure of single-cell data using a variational

autoencoder (VAE) framework. **b**, UMAP embeddings of the *PBMC* dataset before

and after integration using SCALEX, Seurat v3, Harmony, Conos, or Scanorama

integration, colored by batch and cell-type. **c**, Scatter plot showing a quantitative

comparison of the silhouette score (y-axis) and the batch entropy mixing score (x-axis)

on the benchmark datasets. **d**, UMAP embeddings of the SCALEX integration of the

human fetal atlas dataset, colored by batch and cell-type. **e**, Comparison of

computation efficiency on datasets of different sizes sampled from the whole *human*

fetal atlas dataset) including runtime (left) and memory usage (right). **f**, UMAP

embeddings of the mouse brain scATAC-seq dataset before (left) and after integration

(middle, right); colored by data batch or Leiden clustering. **g**, UMAP embeddings of

the PBMC cross-modality dataset before (left) and after integration (middle, right);

colored by batch or cell-type.

Fig. 2 | Comparison of integration performance over partially-overlapping

datasets by different methods. a, Comparison over the *liver* dataset. **b,** Comparison

over simulated datasets with different numbers of common cell-types (obtained by

down-sampling the *pancreas* dataset). Misalignments are highlighted with red circles.

Fig. 3 | Projecting heterogenous data into a generalized cell-embedding space. a,

UMAP embeddings of the *pancreas* dataset after integration by SCALEX, colored by

cell-type. **b**, UMAP embeddings of three projected pancreas data batches projected

onto the pancreas space, colored by cell-types; the light gray shadows represent the

original *pancreas* dataset. **c**, Confusion matrix between ground truth cell-types and

those annotated by different methods. ARI, NMI and F1 scores (top) measure the

annotation accuracy. **d**, UMAP embeddings of the *PBMC* dataset after integration and

the two projected melanoma data batches onto the PBMC space, colored by cell-types

with light gray shadows represent the original *PBMC* dataset. **e**, The PBMC space that

includes the original *PBMC* dataset and the two projected melanoma data batches. **f**,

Annotating an uncharacterized small cell population in the *pancreas* dataset by

projection of the bronchial epithelium data batches into the pancreas cell space. Only

the uncharacterized cells in the *pancreas* dataset (left) and the SLC16A7+ epithelial

cells in the bronchial epithelium data batches (right) are colored. **g**, Heatmap showing

the normalized expression of the top-10 ranking specific genes for the uncharacterized

cell population in different cell-types.

Fig. 4 | Construction of an expandable mouse single-cell atlas. a, Datasets acquired

using different technologies (Smart-seq2, 10X, and Microwell-seq) covering various

tissues used for construction of the mouse atlas. **b**, UMAP embeddings of the *mouse*

atlas dataset colored by batch and tissue. **c**, UMAP embeddings of the *mouse atlas*

after SCLAEX integration, labeld with and colored by cell-type. **d**, Two *Tabula Muris*

Senis data batches and two mouse tissues (lung and kidney) data are projected onto

the cell space of the mouse atlas, with the same cell-type color as in **c**. **e**, Confusion

matrix of the cell-type annotations by SCALEX and those in the original studies.

Color bar represents the percentage of cells in confusion matrix C_{ij} known to be

cell-type i and predicted to be cell-type j.

Fig. 5 | Construction and expansion of a COVID-19 single-cell atlas. a, COVID-19

dataset composition, including healthy controls and influenza patients, as well as

mild/moderate, severe, and convalescent COVID-19 patients. **b**,**c** UMAP embeddings

of COVID-19 PBMC atlas after SCLAEX integration colored by batch (**b**), and by

cell-types (**c**). **d**, UMAP embeddings of the COVID-19 PBMC atlas separated by

disease state. **e**, Stacked violinplot of differentially-expressed ISGs among CD14

monocytes across disease states. **f**, GO terms enriched in the differentially-expressed

genes for CD14-IL1B-Mono and CD14-ISG15-Mono cells. **g**, Cell-type frequency

across healthy and influenza controls, and among mild/moderate, severe, and convalescent COVID-19 patients. Dirichlet-multinomial regression was used for pairwise comparisons, ***p<0.001, **p<0.01, *p<0.05. **h**, United UMAP embeddings of the SCALEX COVID-19 PBMC atlas and the SS4 atlas (from the Single Cell Consortium for COVID-19 in China, projected onto the cell space of the SCALEX COVID-19 PBMC atlas). Left: the SCALEX COVID-19 PBMC atlas, middle: SC4 colored by cell clusters in the original study, right: Expanded atlas combining the SCALEX COVID-19 PBMC atlas and the SC4 atlas. **i**, Similarity matrix of meta-cell representations for cell-types between the SCALEX COVID-19 PBMC atlas and SC4 in the generalized cell-embedding space after SCALEX integration. Color bar represents the Pearson correlation coefficient between the average meta-cell representation of two cell-types from a respective data batch. **j**, UMAP embeddings of the SCALEX COVID-19 PBMC atlas colored by the cytokine score and the inflammatory score. **k**, GO terms enriched in the differentially-expressed genes for TUBA8-Mega and IGKC-Mega cells.

Supplementary figures

Fig. S1 | Comparison of integration performance on benchmark datasets. UMAP embeddings for benchmark datasets grouped by batches and cell-types, before and after integration by different methods. Misalignments are highlighted with red circles.

Fig. S2 | The human fetal atlas. a, UMAP embeddings of the *human fetal atlas* dataset colored by batch before integration. **b,** Similarity matrix of meta-cell representations for different cell-types in the two data batches in the generalized cell-embedding space. Color bar represents the Pearson correlation coefficient between the average meta-cell representation of two cell-types from a respective data batch. **c,** Comparison of computation efficiency on datasets of different sizes

(sampled from the whole *human fetal atlas* dataset), including runtime (left) and
memory usage (right), in log scale.

Fig. S3 | Canonical marker genes of different cell-types and UMAP embeddings
of the *liver* dataset. a, Dotplot of canonical marker genes for each cell-type. Dot
color represents average expression level, while dot size represents the proportion of
cells in the group expressing the marker. **b,** UMAP embeddings of the *liver* dataset,
colored by batch (left) and cell-type (right) after SCALEX integration. **c**, Normalized
marker gene expression on the UMAP embeddings of the five hepatocyte subtypes.
Color bar represents the expression level.

Fig. S4 | Integration over partially-overlapping datasets down-sampled from the
***pancreas* dataset.** Partially-overlapping datasets were generated by down-sampling
the *pancreas* dataset, consisted of common cell-types with a decreased overlapping
number (ranging from 0 to 6). Integration results for SCALEX, Seurat, and Harmony
are shown in the UMAP embeddings colored by batches (left) and cell-types (right)
respectively (overlapping number decreases from 6 to 0). Misalignments are
highlighted with red circles.

Fig. S5 | Integration over partially-overlapping datasets down-sampled from the
***PBMC* dataset.** Partially-overlapping datasets were generated by down-sampling the
PBMC dataset, consisted of common cell-types with a decreased overlapping number
(ranging from 0 to 6). Integration results for SCALEX, Seurat and Harmony are
shown in the UMAP embeddings colored by batches (left) and cell-types (right)
respectively (overlapping number decreases from 6 to 0). Misalignments are
highlighted with red circles.

Fig. S6 | The *pancreas* dataset and the additional data batches. a, UMAP
embeddings of the *pancreas* dataset, the three additional pancreas data batches and
the bronchial epithelium data batches (data from three donors), grouped by batch. **b**,

Dot plot of canonical markers of cell-types of reference *pancreas* dataset; dot color

represents average expression level, while dot size represents the proportion of cells

in the group expressing the marker.

Fig. S7 | The SCALEX mouse atlas. a, UMAP embeddings of the mouse atlas data

before integration, colored by batch. **b**, UMAP embeddings of three mouse atlas data

batches (Smart-seq2, 10X, and Microwell-seq) after integration, colored by cell-type;

the light gray shadows represent the original *mouse atlas* dataset. **c**, Dotplot of the top

5 cell-type-specific genes for each cell-type in the *mouse atlas* dataset. Dot color

represents average expression level, while dot size represents the proportion of cells

in the group expressing the marker.

Fig. S8 | The SCALEX human atlas. a, The *human atlas* dataset acquired using

different technologies (Smart-seq2, 10X, and Microwell-seq) covering various tissues

used for construction of the human atlas. **b-c**, UMAP embeddings of the *human atlas*

dataset colored by batch and cell-type, before (**b**) and after integration (**c**). **d**,

Similarity matrix of meta-cell representations for cell-types in the two data batches in

the generalized cell-embedding space after SCALEX integration between two batches.

Color bar represents the Pearson correlation coefficient between the average meta-cell

representation of two cell-types from a respective data batch. **e**, UMAP embeddings

of the human atlas and two additional projected data batches colored by cell-type. **f**,

Confusion matrix of the cell-type annotations by SCALEX and those in the original

study. Color bar represents the percentage of cells in confusion matrix C_{ij} known to be

in cell-type *i* and predicted to be in cell-type j.

Fig. S9 | COVID-19 immune landscape. a, UMAP embeddings of the raw

COVID-19 PBMC dataset before integration. **b**, UMAP embeddings of the

COVID-19 PBMC atlas colored by condition and Leiden clustering after SCALEX

integration. **c**, Dotplot of canonical marker genes for each cell-type. Dot color

represents average expression level, while dot size represents the proportion of cells

in the group expressing the marker. **d**, UMAP embeddings of the COVID-19 PBMC

atlas in individual batches after SCALEX integration, colored by cell-type; the light

gray shadows represent the other batches of COVID-19 PBMC atlas. **e**, Frequency of

cell distributions across healthy people and influenza patient controls, and among

mild/moderate, severe, and convalescent COVID-19 patients. Dirichlet-multinomial

regression was used for pairwise comparisons, ***$p<0.001$, **$p<0.01$, *$p<0.05$.

Fig. S10 | COVID-19 heterogeneous dysfunctional immune response. a, Stacked

violin plot of differentially-expressed genes between PNPLA2-Immature_Neutrophil

and NCF1-Immature_Neutrophil cells. **b**, GO terms enriched in the

differentially-expressed genes for PNPLA2-Immature_Neutrophil and

NCF1-Immature_Neutrophil cells. **c**, Stacked violinplot of differentially-expressed

genes between PRDM1-Plasma and MZB1-Plasma. **d**, GO terms enriched in the

differentially-expressed genes for PRDM1-Plasma and MZB1-Plasma cells.

Fig S11 | Projection of the SC4 dataset onto the SCLAEX COVID-19 PBMC

atlas. a-b, UMAP embeddings of the SC4 dataset before integration (**a**) and after

projection onto the SCLAEX COVID-19 PBMC space (**b**). **c**, Separate UMAP

embeddings of each SC4 data batch, after being projected onto the SCALEX

COVID-19 PBMC space, colored by cell-type. **d**, UMAP embeddings of the

TUBA8-Mega and IGKC-Mega cells. **e**, UMAP embeddings of the

differentially-expressed genes of TUBA8-Mega and IGKC-Mega cells.

Fig 1

Fig 2

Fig 3

Fig 4

Fig 5

Supplementary Fig 1

Supplementary Fig 2

Supplementary Fig 3

a

Supplementary Fig 4

Supplementary Fig 5

Supplementary Fig 6

a

b

Supplementary Fig 7

Supplementary Fig 8

Supplementary Fig 9

Supplementary Fig 10

Supplementary Fig 11