Host factors drive the within-host competition between highly and
low pathogenic H5N8 avian influenza viruses

Pierre Bessière[1][¶], Thomas Figueroa[1][¶], Amelia Coggon[1], Charlotte Foret-
Lucas[1], Alexandre Houffschmitt[1], Maxime Fusade-Boyer[1], Gabriel Dupré[1],
Jean-Luc Guérin[1], Maxence Delverdier[1], and Romain Volmer[1][*]

[1] Ecole nationale vétérinaire de Toulouse, Université de Toulouse, ENVT,
INRAE, IHAP, UMR 1225, Toulouse, France.

* Correspondence:

Dr Romain Volmer

romain.volmer@envt.fr

¶ These authors contributed equally to this work.

ABSTRACT

Highly pathogenic avian influenza viruses (HPAIV) emerge from low pathogenic avian influenza viruses (LPAIV) through the introduction of basic amino acids at the hemagglutinin (HA) cleavage site. Following viral evolution, the newly formed HPAIV likely represents a minority variant within the index host, predominantly infected with the LPAIV precursor. Using reverse-genetics engineered H5N8 viruses differing solely at the HA cleavage, we tested the hypothesis that the interaction between the minority HPAIV and the majority LPAIV could modulate the risk of HPAIV emergence and that the nature of the interaction could depend on the host species. In chickens, we observed that the $H5N8_{LP}$ increased $H5N8_{HP}$ replication and pathogenesis. By contrast, the $H5N8_{LP}$ antagonized $H5N8_{HP}$ replication and pathogenesis in ducks. Ducks mounted a more potent antiviral innate immune response than chickens against the $H5N8_{LP}$, which correlated with $H5N8_{HP}$ inhibition. These data provide experimental evidence that HPAIV may be more likely to emerge in chickens than in ducks and underscore the importance of within-host viral variants interactions in viral evolution.

INTRODUCTION

Highly pathogenic avian influenza virus (HPAIV) outbreaks have a major impact on animal health, food security and economy, as well as on public health (EFSA Panel on Animal Health and Welfare (AHAW) et al., 2017; Lee et al., 2021). A better understanding of the factors leading to HPAIV emergence is therefore of paramount importance. Avian influenza viruses of H5 and H7 subtypes can become highly pathogenic through the introduction of multiple basic amino acids within the hemagglutinin (HA) cleavage site (Böttcher-Friebertshäuser et al., 2014). Several mechanisms account for these introductions: nucleotide substitutions and insertions or non-homologous recombination with viral or cellular RNAs (Abdelwhab et al., 2013; Lee et al., 2021; Richard et al., 2017). The acquisition of a multi-basic cleavage site (MBCS) is a virus-dependent event that occurs in a bird infected with a parental low pathogenic avian influenza virus (LPAIV). To emerge successfully, the newly formed HPAIV must become a predominant variant in order to overcome the transmission bottleneck between individuals (Hughes et al., 2012; McCrone and Lauring, 2018; Varble et al., 2014). In order to achieve this, the newly formed HPAIV must therefore outcompete its LPAIV precursor within the individual in which it has arisen. We therefore suggest that HPAIV emergence is a two-step process: firstly, the acquisition of a MBCS, followed by the ability to become a predominant variant of the viral quasi-species within an individual.

Over the past decades, the vast majority of HPAIV emergences have been linked to H5 and H7 LPAIV circulation in *Galliformes* (such as chickens and turkeys) (Abdelwhab et al., 2013; EFSA Panel on Animal Health and Welfare (AHAW) et al., 2017; Lee et al., 2021). By contrast, *Anseriformes* (such as ducks and geese) are considered as reservoirs for LPAIV precursors, rather than a species in which HPAIV emerge (Bodewes and Kuiken, 2018; Munster et al., 2007; van Dijk et al., 2018; Wahlgren, 2011). These observations suggest that host factors could modulate HPAIV emergence. To our knowledge, this has never been investigated experimentally.

To model the intra-host competition between a newly formed HPAIV and its parental LPAIV, we co-infected embryonated eggs and chickens and ducks *in vivo* with a H5N8 HPAIV as a minority variant and a reverse-genetics engineered LPAIV that differed from the HPAIV only at the level of the HA cleavage site, as a majority variant. Our results demonstrate that chickens and ducks have opposite effects on the interaction between the H5N8 HPAIV and LPAIV and that the HPAIV has a stronger selective advantage in chickens than in ducks. To our knowledge this is the first experimental evidence that HPAIV emergence could be more likely in chickens than in ducks.

RESULTS

Characterization of H5N8$_{HP}$ and H5N8$_{LP}$ viruses in cell culture

We used reverse-genetics to generate the wild-type H5N8$_{HP}$ from a HPAIV isolated during the 2016 H5N8 epizootics in France (Guinat et al., 2018). This virus belonged to clade 2.3.4.4 group B, which caused high levels of mortality in *Galliformes* and in wild and domestic *Anseriformes* during the 2016-2017 HPAIV outbreak in Europe (Grund et al., 2018; Guinat et al., 2018; Kleyheeg et al., 2017; More et al., 2017). Using site-directed mutagenesis and reverse-genetics, we mutated the H5N8$_{HP}$ HA polybasic cleavage site PLRELRRLR/G to a monobasic cleavage site PQRETR/G to obtain the H5N8$_{LP}$ virus with a typical LPAIV HA sequence (NCBI Genbank accession number: AB261853.1) (Hiono et al., 2016). The H5N8$_{HP}$ and the H5N8$_{LP}$ differed solely at the level of the HA cleavage site, as verified by whole genome sequencing following virus amplification in chicken embryonated eggs (Fig. 1A).

Viral growth kinetics was analyzed in Madin-Darby canine kidney (MDCK) cells, in the DF-1 chicken fibroblast cell line and in the CCL-141 duck fibroblast cell line infected at a multiplicity of infection (MOI) of 10^{-5} tissue culture infectious dose 50 (TCID$_{50}$). H5N8$_{HP}$ replicated with similar kinetics in the absence or presence of exogenous trypsin, demonstrating that proteolytic processing of the HA MBCS was independent of trypsin-like

proteases, as expected for a HPAIV (Fig. 1B). By contrast, H5N8$_{LP}$

replication was severely impaired in the absence of trypsin, while its

replication kinetics was indistinguishable from that of the H5N8$_{HP}$ in the

presence of trypsin (Fig. 1B). These results indicate that mutations

introduced in the H5N8$_{LP}$ HA have no negative impact on viral replication

kinetics in cell culture beyond the expected trypsin-like proteases

requirement for H5N8$_{LP}$ HA proteolytic processing, as expected for a LPAIV.

H5N8$_{HP}$ and H5N8$_{LP}$ viruses co-infections *in ovo*

Next, we evaluated H5N8$_{HP}$ and H5N8$_{LP}$ replication in embryonated chicken

(*Gallus gallus*) and Pekin duck (*Anas platyrhynchos domesticus*) eggs, as

they have a complex cellular architecture and provide an intermediate

model between monolayer cell culture and *in vivo* studies. To investigate

how the host species modulated the interaction between a HPAIV and a

LPAIV, we inoculated embryonated chicken and duck eggs via the allantoic

cavity either with 10^2 egg infectious dose 50 (EID$_{50}$) H5N8$_{HP}$ alone, with

increasing doses of H5N8$_{LP}$ alone, or with 10^2 EID$_{50}$ H5N8$_{HP}$ in combination

with increasing doses of H5N8$_{LP}$ (Fig. 2A). We inoculated chicken eggs at

day 10 and duck eggs at day 11 in order to work at similar development

stages (Li et al., 2019). After 24 hours of incubation at 37°C, we harvested

the allantoic fluid and quantified the level of virus replication by RT-qPCR

using primers specific for H5N8$_{HP}$ or H5N8$_{LP}$ HA. In chicken eggs, we

detected equivalent levels of viral RNA following inoculation with 10^2EID$_{50}$

H5N8$_{HP}$ alone or with H5N8$_{LP}$ alone regardless of the inoculum dose (Fig.

2B). Upon co-infection with H5N8$_{HP}$ and H5N8$_{LP}$ in chicken eggs, the level

of H5N8$_{HP}$ and H5N8$_{LP}$-specific viral RNA did not differ from mono-

infections regardless of the quantity of H5N8$_{LP}$ (Fig. 2B). Thus, the

replication of H5N8$_{HP}$ is not affected by H5N8$_{LP}$, regardless of the amount of

H5N8$_{LP}$ co-inoculated in chicken eggs. In duck eggs, we detected similar

levels of viral RNA following inoculation with 10^2 EID$_{50}$ H5N8$_{HP}$ alone or

with 10^2 EID$_{50}$ H5N8$_{LP}$ alone (Fig. 2C). When duck eggs were infected

simultaneously with H5N8$_{HP}$ and H5N8$_{LP}$, we observed a decrease of

H5N8$_{HP}$ viral RNA levels that correlated with the amount of H5N8$_{LP}$ co-

inoculated, indicating that the H5N8$_{LP}$ antagonized H5N8$_{HP}$ replication in duck eggs (Fig. 2C). The *in ovo* experiments provide evidence that a ≥100-fold excess of H5N8$_{LP}$ significantly antagonized H5N8$_{HP}$ replication in duck eggs, while H5N8$_{LP}$ did not affect H5N8$_{HP}$ replication in chicken eggs regardless of the amount of H5N8$_{LP}$. This observation thus indicates that the H5N8$_{HP}$ had a stronger selective advantage over the H5N8$_{LP}$ in chicken embryonated eggs than in duck embryonated eggs.

Potentiation of HPAIV replication and pathogenesis by LPAIV in chickens

In a preliminary experiment (data not shown), we observed transient low-level oropharyngeal viral RNA shedding in 6/8 of the 10^6 EID$_{50}$ H5N8$_{LP}$ infected chickens (*Gallus gallus*), whereas all chickens infected with 10^7 EID$_{50}$ H5N8$_{LP}$ shed high levels of viral RNA for a prolonged time. No clinical signs were observed in chickens inoculated with H5N8$_{LP}$. Furthermore, by using increasing doses of H5N8$_{HP}$, we evaluated the H5N8$_{HP}$ chicken infectious dose 50 at 6×10^3 EID$_{50}$ and noticed that mortality reached 100% within four days with doses higher than 10^5 EID$_{50}$. H5N8$_{HP}$-infected sick birds presented dyspnea that quickly progressed to severe dyspnea, associated with anorexia and lethargy. These results provided evidence that the H5N8$_{HP}$ and H5N8$_{LP}$ caused, respectively, commonly observed HPAIV and LPAIV-associated infection patterns in chickens (EFSA Panel on Animal Health and Welfare (AHAW) et al., 2017).

Based on these results, we investigated the interaction between a HPAIV and a LPAIV in chickens. Four-week old chickens were assigned one of six groups: L7, H3, H4, L7H3, L7H4 or NI (Fig. 3A). L7 animals were infected with 10^7 EID$_{50}$ H5N8$_{LP}$. H3 animals were infected with 10^3 EID$_{50}$ H5N8$_{HP}$. H4 animals were infected with 10^4 EID$_{50}$ H5N8$_{HP}$. L7H3 animals were infected with a mixture of 10^7 EID$_{50}$ H5N8$_{LP}$ and 10^3 EID$_{50}$ H5N8$_{HP}$. L7H4 animals were infected with a mixture of 10^7 EID$_{50}$ H5N8$_{LP}$ and 10^4 EID$_{50}$ H5N8$_{HP}$. Finally, non-infected animals from group NI were administered vehicle only. Neither mortality nor clinical signs were observed in chickens from group L7 and group H3 (Fig. 3B). Chickens in the H4 group reached

45% mortality (Fig. 3B). When chickens were infected with a mixture of H5N8$_{LP}$ and H5N8$_{HP}$, we observed an increase in mortality compared to H5N8$_{HP}$ mono-infected chickens: mortality reached 18% in the L7H3 group, compared to 0% in the H3 group; and 72% in the L7H4 group, compared to 45% in the H4 group (Fig. 3B). Although the increase in mortality did not reach statistical significance, these results indicate that H5N8$_{LP}$ and H5N8$_{HP}$ synergized, resulting in increased pathogenesis in co-infected chickens compared to H5N8$_{HP}$ mono-infected chickens.

We measured viral shedding from oropharyngeal and cloacal swabs by quantifying viral RNA using RT-qPCR with primers detecting specifically H5N8$_{LP}$ or H5N8$_{HP}$ HA sequences. Regardless of the experimental group, all H5N8$_{HP}$ swab-positive chickens eventually developed severe dyspnea and either died or were euthanized when they reached humane endpoints, as previously observed (Leyson et al., 2019). In line with the mortality rate (Fig 3B), no H5N8$_{HP}$ swab-positive chicken was found in the H3 group, while only two chickens tested positive for H5N8$_{HP}$ in the L7H3 group. Because of the low rate of infection in the H3 and L7H3 groups, these animals were excluded from further analyses. H5N8$_{HP}$ oropharyngeal and cloacal shedding was higher in group L7H4 compared to group H4 at 2 days post-infection (dpi), but the difference was not statistically significant (Fig. 3C&D). Average H5N8$_{HP}$ oropharyngeal and cloacal shedding remained otherwise similar between the L7H4 and the H4 groups and was not associated with different H5N8$_{HP}$ transmission rates to contact chickens introduced in the poultry isolators at 1 dpi (data not shown). H5N8$_{LP}$ oropharyngeal shedding did not differ between the L7 and L7H4 group (Fig. 3E).

Antagonism of HPAIV replication and pathogenesis by LPAIV in ducks

We inoculated groups of 4-week old Pekin ducks (*Anas platyrhynchos domesticus*) with H5N8$_{HP}$ alone, H5N8$_{LP}$ alone, or a mixture containing both H5N8$_{LP}$ and H5N8$_{HP}$. Ducks were assigned one of four groups: L7, H4, L7H4 or NI (Fig. 7A). L7 animals were infected with 10^7 EID$_{50}$ H5N8$_{LP}$. H4 animals were infected with 10^4 EID$_{50}$ H5N8$_{HP}$. L7H4 animals were infected with a mixture of 10^7 EID$_{50}$ H5N8$_{LP}$ and 10^4 EID$_{50}$ H5N8$_{HP}$. Non-infected animals

from group NI were administered vehicle only. Neither mortality nor clinical signs were observed in any duck of group L7 (Fig. 7B). Mortality reached 87% in the H4 group and was preceded by predominantly neurological clinical signs, in accordance with the pronounced neurotropism of clade 2.3.4.4 group B H5N8 viruses in ducks (Grund et al., 2018; Kleyheeg et al., 2017; More et al., 2017). Mortality was significantly reduced to 37% in the L7H4 group. Thus, $H5N8_{LP}$ antagonized $H5N8_{HP}$ pathogenesis in co-infected ducks, in contrast to what we observed in chickens.

To determine how this related to the levels of virus replication, we analyzed $H5N8_{LP}$ and $H5N8_{HP}$ viral RNA shedding from oropharyngeal and cloacal swabs using RT-qPCR. Oropharyngeal $H5N8_{HP}$ shedding was significantly reduced in group L7H4 compared to group H4 in the first days post-infection (dpi), with a hundred-fold difference at 1 dpi ($p<0.001$) and a ten-fold difference at 2 and 3 dpi ($p<0.01$) (Fig. 7C). Cloacal $H5N8_{HP}$ shedding was also decreased in group L7H4, but the difference only reached statistical difference at day 3, with a ten-fold reduction ($p<0.05$) (Fig. 7D). There was no difference in $H5N8_{LP}$ shedding between groups L7H4 and L7 (Fig. 7E).

Next, we analyzed viral RNA levels in the lungs and brain. $H5N8_{LP}$ was not detected in the lungs or brain, indicating that $H5N8_{LP}$ replicated mostly in the upper respiratory tract of ducks. In contrast to chickens, we detected $H5N8_{HP}$ nucleic viral RNA from the lungs and brain of most animals at 1 and 3 dpi (Fig. 7F&G). $H5N8_{HP}$ viral RNA load was decreased in the lungs and brain of $H5N8_{LP}$ and $H5N8_{HP}$ co-infected birds (group L7H4), compared to $H5N8_{HP}$ mono-infected birds (group H4). High levels of $H5N8_{HP}$ viral RNA were also detected in the lungs and brain of moribund ducks, which developed predominantly neurological signs.

Extensive viral antigen staining was detected by immunohistochemistry in the brain of moribund animals, while more modest viral antigen staining was observed in the lungs, with no difference between groups L7H4 and H4 (Fig. 8A&B). We then evaluated the expression of host immune response markers in the lungs and brain using RT-qPCR. Mx mRNA expression was

significantly increased in the lungs of infected ducks belonging to the L7H4 and H4 groups at 1 and 3 dpi, as well as in moribund animals (Fig. 9A). Compared to NI duck, we also observed an upregulation of Mx mRNA expression in the lungs of H5N8$_{LP}$ mono-infected ducks (L7 group) at 1 and 3 dpi, although the difference did not reach statistical significance. IFN-γ mRNAs levels were significantly higher in the lungs of H5N8$_{HP}$ mono-infected ducks (group H4) at 1 dpi compared to NI, L7H4 and L7 ducks (Fig. 9B). Similar results were observed in the brain (data not shown). To evaluate the antiviral innate immune response in the upper respiratory tract, we measured Mx mRNA expression by RT-qPCR from RNA extracted from oropharyngeal swabs. Compared to NI ducks, Mx mRNA was increased in all infected groups (Fig. 9C). In contrast to chickens, Mx mRNA expression level remained high in H5N8$_{LP}$ mono-infected ducks from the L7 group (Fig. 9C).

To further compare the antiviral innate immune response in ducks and chickens, we plotted the levels of Mx mRNA as a function of the level of viral RNA in the lungs (Fig. 10A). We observed a correlation between the level of Mx mRNA and the level of viral RNA in ducks (Pearson r=0.64; p<0.001), confirming that the intensity of the antiviral innate immune response is proportional to the level of viral RNA, as previously reported (Baccam et al., 2006; Hagenaars et al., 2016; Soubies et al., 2013; Volmer et al., 2011). This correlation was less visible in chickens, possibly because viral RNA was only found in the lungs of a limited number of chickens (Pearson r=0.50; p=0.21). The Mx/HA ratios were distributed to higher values in ducks compared to chickens (Fig. 10A), resulting in mean Mx/HA ratios that were significantly higher in ducks than in chickens (Fig. 10B). This result demonstrates that ducks mounted a more potent antiviral innate immune response against influenza virus infection than chickens.

DISCUSSION

To model the intra-host competition between a newly formed HPAIV and its parental LPAIV, we inoculated chickens and ducks with an H5N8$_{HP}$ virus and an H5N8$_{LP}$ virus differing solely at the level of the HA cleavage site. When inoculated alone, H5N8$_{HP}$ replication and pathogenesis were equivalent in chickens and ducks. However, when chickens and ducks were co-inoculated with this pair of viruses, we found that the H5N8$_{HP}$ had a stronger selective advantage over the H5N8$_{LP}$ in chickens than in ducks. Surprisingly, we observed that the H5N8$_{LP}$ increased H5N8$_{HP}$ replication and pathogenesis in chickens. By contrast, we observed that the H5N8$_{LP}$ antagonized H5N8$_{HP}$ replication and pathogenesis in ducks. Thus, the nature of the interaction between the HPAIV and the parental LPAIV may be a critical determinant of HPAIV emergence, which we show here to depend on the host species.

The interaction between co-infecting viruses ranges from synergy to antagonism depending on the virus pairs and the host (Andreu-Moreno et al., 2020; Bou et al., 2019; DaPalma et al., 2010; Domingo-Calap et al., 2019; Kumar et al., 2018; Opatowski et al., 2018; Phipps et al., 2020). Functional complementation occurs when one virus provides a function that is absent or less efficient in the other virus. As the genomes of the H5N8$_{HP}$ and H5N8$_{LP}$ differ solely at the level of the HA cleavage site, it is unlikely that the H5N8$_{LP}$ is able to provide a function missing in the H5N8$_{HP}$. We therefore speculate that the H5N8$_{LP}$-mediated increase in H5N8$_{HP}$ replication and pathogenesis observed in chickens could be due to complementation of semi-infectious H5N8$_{HP}$ particles by the H5N8$_{LP}$ inoculated at a higher dosage (Brooke, 2017). Such complementation is likely to occur both in chickens and in ducks. However, this complementation may be undetectable in ducks due to the dominant effect of host-mediated antagonism between the H5N8$_{LP}$ and the H5N8$_{HP}$. The antagonism between the H5N8$_{LP}$ and the H5N8$_{HP}$ is unlikely to be due to viral interference due to receptor or cell machinery usage, as no competition between the H5N8$_{LP}$ and the H5N8$_{HP}$ was observed in

chickens, although the H5N8$_{LP}$ replicated to higher levels and persisted longer in chickens than in ducks. In contrast to chickens, ducks have a functional RIG-I receptor, which has been proposed to contribute to the more efficient type I interferon mediated-antiviral innate immune response against influenza viruses observed in ducks compared to chickens (Barber et al., 2010; Burggraaf et al., 2014; Cornelissen et al., 2013, 2012; Kuchipudi et al., 2014; Magor et al., 2013; Soubies et al., 2013). The H5N8$_{LP}$ and the H5N8$_{HP}$ are likely to be intrinsically equally sensitive to the antiviral effects of type I interferon. However, a stronger antiviral innate immune response could impose a stronger selective pressure on the minority variant (Zwart and Elena, 2015). In line with this hypothesis, type I interferon antiviral innate immune response was shown to constitute a strong bottleneck shaping poliovirus population in mice (Kuss et al., 2008). Similarly, treatment of hepatitis C virus-infected patients with type I IFN resulted in a dramatic reduction of viral diversity (Farci et al., 2002). Furthermore, a delayed type I IFN response in influenza A virus-infected obese mice promoted minority variant emergence, while type I interferon treatment reduced viral diversity (Honce et al., 2020). Altogether, these results indicate that the innate immunity can shape viral populations. By decreasing viral replication and increasing the within-host competition between variants, a strong innate immune response may inhibit the selection of variants that arise through *de novo* mutations during infection of an individual, unless these variants have a strong selective advantage within this individual. We suggest that the more pronounced antiviral innate immune response observed in ducks compared to chickens may impact the H5N8$_{HP}$ minority variant more severely in ducks than in chickens. Admittedly, the differential outcome of H5N8$_{HP}$ and H5N8$_{LP}$ co-infections in chickens and ducks may also depend on other host factors, including microbiota composition or receptor distribution (Figueroa et al., 2020; Hiono et al., 2016).

Recent work investigating the interaction between an H7N7 HPAIV and a closely related H7N7 LPAIV field isolate demonstrated that the LPAIV inhibited HPAIV replication in chickens when the LPAIV was inoculated at a

≥ 100-fold higher dose (Graaf et al., 2018). These results indicate that the
nature of the interaction between the HPAIV and the LPAIV may depend on
the viral strain. If competition between a LPAIV and a HPAIV occurred in
chickens, we speculate that the competition could even be stronger in
ducks, because of the stronger antiviral innate immune-mediated selective
pressure imposed on the minority HPAIV. However, further work is needed
to properly address this question. These studies should ideally be
performed using different pairs of reverse-genetics engineered HPAIV and
LPAIV differing solely at the level of the HA cleavage site to specifically
investigate how the selective advantage conferred by the MBCS depends on
the host species.

Epidemiological investigations of HPAIV outbreaks indicate that the vast
majority of HPAIV emerge upon replication and inter-individual
transmission of a LPAIV in *Galliformes*, such as chickens and turkeys,
although *Anseriformes*, such as ducks and geese, are considered as the main
reservoirs for LPAIV (Abdelwhab et al., 2013; EFSA Panel on Animal Health
and Welfare (AHAW) et al., 2017; Lee et al., 2021; Richard et al., 2017). In
this study, we focused on the two main representatives of the *Galliformes*
and *Anseriformes* orders: chickens, which are with current global
production of 22 billion chickens the main source of animal protein for
human consumption worldwide and Pekin ducks, which are also
intensively raised in farms and represent the best characterized
Anseriformes in respect to the interaction with influenza viruses (Lee et al.,
2021; "Poultry species | Gateway to poultry production and products | Food
and Agriculture Organization of the United Nations," n.d.). How the LPAIV
and the newly formed HPAIV would interact in other bird species, such as
turkeys, or in wild dabbling or diving ducks, which have a prominent role in
the ecology of avian influenza viruses will be the subject of future studies
(Bodewes and Kuiken, 2018; EFSA Panel on Animal Health and Welfare
(AHAW) et al., 2017; Lee et al., 2021). In addition to host factors, anthropic
factors are also likely to play a major role in the emergence of HPAIV and
may contribute to the higher frequency of HPAIV emergence observed in
chickens that are usually raised at higher densities than ducks (Capua and

Marangon, 2000; Dhingra et al., 2018; M.c et al., 2009; Richard et al., 2017).

Estimating to which extent farming processes contribute to the emergence

of HPAIV is however a difficult task and therefore their contribution to

HPAIV emergence remains currently unclear.

To minimize the risk of HPAIV evolution, present European regulation and

national regulations in many countries require the culling of all birds in

flocks infected with H5 and H7 LPAIV (EFSA Panel on Animal Health and

Welfare (AHAW) et al., 2017). Culling of H5 or H7 LPAIV infected flocks

aims at eliminating LPAIV from the susceptible poultry population before

they have a chance to evolve into HPAIV, and is therefore considered a

preventive biosecurity measure. However, the broader public, breeders and

professional breeder societies question the need for preventive culling for

economical, animal welfare and breeder welfare issues (Organization,

2008). Indeed, preventive culling of animals, especially healthy animals,

which is commonly the case in LPAIV infected *Anseriformes*, is poorly

accepted. Identifying how host factors modulate HPAIV emergence

therefore provides important scientific data that may be taken into account

by animal health policy makers in the future.

MATERIALS AND METHODS

Viruses

The eight segments from the HPAIV field isolate A/mulard duck/France/171201g/2017 (H5N8) were cloned in a pHW2000 plasmid vector using a set of universal primers in order to generate a reverse-genetics engineered wild-type HPAIV H5N8 (H5N8$_{HP}$) (Hoffmann et al., 2001, 2000). Using site-directed mutagenesis, a 9 nucleotides deletion was performed on the HA cleavage site sequence, along with two single nucleotide polymorphism on both sides of the deletion to obtain a reverse-genetics engineered LPAIV H5N8 (H5N8$_{LP}$) according to the manufacturer's instructions (In-Fusion, Takara Bio, France). 2.5×10^5 HEK 293T cells cultured in 6-well plates were transfected in Opti-MEM medium, using the lipophilic transfection reagent LTX with Plus reagent (Invitrogen, Thermo Fisher Scientific Inc., Canada), either with 0.5µg of each seven common H5N8$_{HP/LP}$ pHW2000 plasmids (PB2, PB1, PA, NA, NP, M and NS) and with 0.5µg of HA$_{HP}$ or HA$_{LP}$. L-(tosylamido-2-phenyl)ethyl chloromethyl ketone (TPCK) treated trypsin (TPCK-treated) was added 24 hours post-transfection at a 0.5µg/ml final concentration. 48 hours post-transfection, scraped cells and culture medium were transferred on MDCK cells grown with Opti-MEM supplemented with 0.5µg/ml TPCK-treated trypsin. 72 hours later, cell supernatant was collected. To produce viral stock, both viruses were then propagated in 10-day-old chicken specific-pathogen-free (SPF) embryonated chicken eggs (INRAE, PFIE, Nouzilly, France), by inoculation in the allantoic cavity of 100µL of 1:100 dilutions of infected MDCK cells supernatant. Infectious allantoic fluid was harvested at 72 hours postinoculation and titrated in 10-day-old SPF embryonated chicken eggs to determine the 50% egg infective dose (EID$_{50}$)/mL using the Reed-Muench method. The identity of amplified viruses was verified by Sanger sequencing of each viral gene segment (H5N8$_{HP}$: accession numbers MK859904 to MK859911; H5N8$_{LP}$: accession numbers MK859926 to MK859933). These viruses were referenced by the French biotechnology ethics committee (Haut Conseil des Biotechnologies) and were manipulated exclusively in biosafety level 3 laboratories.

In vitro infections

MDCK cells, DF-1 cells and CCL-141 cells were cultured in DMEM containing 1% antibiotics (penicillin/streptomycin) and 10% fetal bovine serum. When cells reach 90% confluency, they were washed with PBS and infected either with H5N8$_{HP}$ or with H5N8$_{LP}$ at a multiplicity of infection (MOI) of 10^{-5} TCID$_{50}$. After one hour, the inoculum was removed, cells were washed twice with PBS and Opti-MEM supplemented with 0.5µg/mL TPCK-treated trypsin (Pierce, Thermo Fisher Scientific Inc., Canada) was added. Viral RNA extraction was performed on 140µL supernatant collected at 1, 24, 48 and 72 hours post-infection according to the manufacturer's instructions (QIAamp viral RNA; Qiagen, Toronto, Canada). Influenza nucleic acid load was determined by RT-qPCR using primers targeting both H5N8$_{LP}$ and H5N8$_{HP}$ HA gene in a final volume of 10µL. Mixes were prepared according to the manufacturer's instructions (iTaq Universal SYBR green One-Step kit, BioRad) with 1µL of RNA and a final concentration of 0.3µM of each primer.

In ovo co-infection experiments

Specific pathogen-free (SPF) White Leghorn (PA12) embryonated chicken eggs (PFIE, INRAE, Nouzilly, France) and Pekin Duck (ST5 Heavy) embryonated eggs (ORVIA-Couvoir de la Seigneurtière, Vieillevigne, France) were respectively incubated for 10 and 11 days at 37°C to work at similar stages of development (Li et al., 2019). They were then inoculated in the allantoic cavity with 200µL virus diluted in PBS containing 1% antibiotics (penicillin/streptomycin). Eggs from group H2 were infected with 10^2 EID$_{50}$ H5N8$_{HP}$. Eggs from groups L2, L3, L4 and L5 were infected with 10^2 EID$_{50}$, 10^3 EID$_{50}$, 10^4 EID$_{50}$, and 10^5 EID$_{50}$ H5N8$_{LP}$ respectively. Eggs from groups L2H2, L3H2, L4H2 and L5H2 were infected with a mixture of 10^2 EID$_{50}$ H5N8$_{HP}$ and increasing doses of H5N8$_{LP}$: 10^2, 10^3, 10^4 and 10^5 EID$_{50}$ H5N8$_{LP}$ respectively. Each group contained 9 to 29 eggs from two to three independent experiments. 24 hours post-infection eggs were

incubated at 4°C overnight and allantoic fluid was harvested for RNA extraction.

Animals and groups

Chickens and ducks experiment were conducted separately. One-day-old Pekin ducklings (*Anas platyrhyncos domesticus*, ST5 heavy) were obtained from a commercial hatchery (ORVIA-Couvoir de la Seigneurtière, Vieillevigne, France) and one-day-old White Leghorn chicks (*Gallus gallus domesticus*, PA12) from a research hatchery (PFIE, INRAE, Nouzilly, France). Animals were fed *ad libitum* with a starter diet and housed in biosafety level II facilities for 2 weeks in a litter-covered floor pen at the National Veterinary School of Toulouse, France. They were then transferred into a biosafety level III facility, equipped with bird isolators (I-Box; Noroit, Nantes, France) ventilated under negative pressure with HEPA-filtered air. Chicken preliminary experiment: 39 chickens were randomly assigned to four groups: 10 animals were assigned to group H3 and were infected with 10^3 EID_{50} $H5N8_{HP}$. 10 animals were assigned to group H4 and were infected with 10^4 EID_{50} $H5N8_{HP}$. 8 animals were assigned to group L6 and were infected with 10^6 EID_{50} $H5N8_{LP}$. 11 animals were assigned to group L7 and were infected with 10^7 EID_{50} $H5N8_{LP}$.

Chickens co-infection experiment: A total of 130 chickens were randomly assigned to 4 groups: 26 animals (including 5 contact birds introduced 24 hours post-infection) were assigned to group H3 and were infected with 10^3 EID_{50} $H5N8_{HP}$. 26 animals (including 5 contact birds) were assigned to group H4 and were infected with 10^4 EID_{50} $H5N8_{HP}$. 21 animals were assigned to group L7 and were infected with 10^7 EID_{50} $H5N8_{LP}$. 26 animals (including 5 contact birds) were assigned to group L7H3 and were simultaneously infected with a mixture of 10^7 EID_{50} $H5N8_{LP}$ and 10^3 EID_{50} $H5N8_{HP}$. 26 animals (including 5 contact birds) were assigned to group L7H4 and were simultaneously infected with a mixture of 10^7 EID_{50} $H5N8_{LP}$ and 10^4 EID_{50} $H5N8_{HP}$. 5 animals were assigned to the non-infected control group (NI).

Ducks co-infection experiment: A total of 64 ducks were randomly assigned
to 4 groups: 21 animals (including 3 contact birds) were assigned to group
H4 and were infected with 10^4 EID$_{50}$ H5N8$_{HP}$. 16 animals were assigned to
group L7 and were infected with 10^7 EID$_{50}$ H5N8$_{LP}$. 21 animals (including 3
contact birds) were assigned to group L7H4 and were simultaneously
infected with a mixture of 10^7 EID$_{50}$ H5N8$_{LP}$ and 10^4 EID$_{50}$ H5N8$_{HP}$. 5
animals were assigned to the non-infected control group (NI).
479
***In vivo* experimental infections**
Serum was collected from all animals pre-infection to ensure that they
were serologically negative to influenza virus by using a commercial
influenza A NP antibody competition enzyme-linked immunosorbent assay
kit (ID Screen; ID-Vet, Montpellier, France) according to the manufacturer's
instructions. When they were 4-week-old, animals were infected through
the choanal route using an inoculum volume of 100μL. Non-infected groups
received the equivalent volume of allantoic fluid collected from non-
infected SPF embryonated chicken eggs. Contact birds were introduced in
the poultry isolators 1 day post-infection. Clinical signs were recorded over
8 to 9 days. Oropharyngeal and cloacal swabs were performed on all
animals daily. 5 animals from each group were humanely euthanized and
necropsied at 1 and 3 dpi. Moribund animals reaching humane termination
criteria (as dyspnea, convulsions, severe lethargy) were humanely killed
and also necropsied. For each necropsied animal, brain and lungs were
collected and stored frozen in TRIzol reagent (Invitrogen, Carlsbad, CA) or
stored in 10% neutral formalin.
497
H5N8$_{LP}$ and H5N8$_{HP}$ quantification
Cloacal and oropharyngeal swabs from *in vivo* experiments were briefly
vortexed in 500μL of sterile PBS and viral RNA was extracted from 200μL
using a QiaCube automated platform according to the manufacturer's
instructions (Cador Pathogen QIAcube HT kit; Qiagen, Toronto, Canada).
Allantoic fluids from *in ovo* experiments were collected and viral RNA was
extracted from 140μL according to the manufacturer's instructions

(QIAamp viral RNA; Qiagen, Toronto, Canada). cDNA was synthesized by reverse transcription of 5μL of RNA, using a HA-specific primer (5'-GTCCTTGCGACTG-3') and a RevertAid first-strand cDNA synthesis kit (Invitrogen, Thermo Fisher Scientific) according to the manufacturer's instructions. Influenza nucleic acid load was determined by qPCR using primers targeting either the polybasic or the monobasic HA cleavage site. qPCR was performed in 384-well plates in a final volume of 10μL using a Bravo automated liquid handling platform (Agilent Technologies, Palo Alto, CA) and a ViaaA 7 real-time PCR system (Applied Biosystems, Foster City, CA) at the GeT-TRiX platform (GénoToul, Génopole, Toulouse, France). Mixes were prepared according to the manufacturer's instructions (iTaq SYBR green PCR, BioRad) with 2μL of cDNA and a final concentration of 0.5μM of each primer. Absolute quantification was performed using a standard curve based on 10-fold serial dilutions of plasmid containing either H5N8$_{LP}$ or H5N8$_{HP}$ HA gene.

RNA extraction from tissue samples and cDNA synthesis

For each organ, 30mg portions of tissue were placed in tubes with beads (Precellys lysis kit; Stretton Scientific, Ltd., Stretton, United Kingdom) filled with 1mL of TRIzol reagent (Invitrogen, Carlsbad, CA) and mixed for 5s at 6,000rpm three times in a bead beater (Precellys 24; Bertin Technologies, Montigny-le-Bretonneux, France). After TRIzol extraction, the aqueous phase was transferred to a RNA extraction column and processed according to the manufacturer's instructions (NucleoSpin RNA; Macherey-Nagel GmbH & Co, Germany). cDNA was synthesized by reverse transcription of 500ng of total RNA using either both oligo(dT)18 (0.25μg) and random hexamer (0.1μg) primers or HA-specific primer and a RevertAid first-strand cDNA synthesis kit (Invitrogen, Thermo Fisher Scientific) according to the manufacturer's instructions.

Quantitative PCR from tissue samples

Quantitative PCR for the analysis of host genes expression was performed in 384-well plates in a final volume of 10μL using a Bravo automated liquid

handling platform (Agilent Technologies, Palo Alto, CA) and a ViiA 7 real-time PCR system (Applied Biosystems, Foster City, CA) at the GeT-TRiX platform (GenoToul, Genopole, Toulouse, France). Mixes were prepared according to the manufacturer's instructions (iTaq SYBR green PCR, BioRad) with 2µl of 1:20 diluted cDNA and a final 0.5µM concentration of each primer (Table 1). Quantification of influenza virus nucleic acid load in tissues was performed in 96-well plates with a 10µL final volume according to the manufacturer's instructions (iTaq SYBR green PCR, BioRad), along with 2µL of cDNA and a final 0.5µM concentration of $H5N8_{HP}$ or $H5N8_{LP}$ specific primers. qPCR was performed on a LightCycler 96 (Roche). Relative quantification was carried out by using the $2^{-\Delta Ct}$ method. Chickens: RNA levels were normalized with the geometric mean of GAPDH/HPRT1 mRNA levels. Ducks: mRNA levels were normalized with the geometrics means of RPL4/RPL30 mRNA levels in the lungs and RPL4/GAPDH in the brain. Oropharyngeal Mx mRNA expression was performed in 96-well plates with a final volume of 10µL according to the manufacturer's instructions (iTaq SYBR green one-step, BioRad), with 2µL RNA and a final 0.3µM concentration of each primer. Mx mRNA levels were normalized with GAPDH mRNA levels in chickens and ducks samples using the $2^{-\Delta Ct}$ method.

Immunohistochemistry

Tissue samples of the lungs and brain were taken and stored in 10% neutral formalin. After fixation, tissues were processed in paraffin blocks, sectioned at 4µm, and immunohistochemistry was performed on paraffin-embedded sections with a monoclonal mouse anti-nucleoprotein influenza A virus antibody (Argene 11-030; pronase 0.05% retrieval solution, 10 min at 37°C: antibody dilution 1/50, incubation overnight at 4°C). The immunohistochemical staining was revealed with a biotinylated polyclonal goat anti-mouse immunoglobulin conjugated with horseradish peroxidase (HRP; Dako, LSAB2 system-HRP, K0675) and the diaminobenzidine HRP chromogen (Thermo Scientific, TA-125-HDX). Negative controls comprised sections incubated either without specific primary antibody or with another monoclonal antibody of the same isotype (IgG2).

572   Table 1: Primers used for qPCR

Gene name	Primers sequences (5' to 3')	NCBI accession number or Reference
HA H5N8$_{HP}$	CCCAAACAGCCCTCTTT	
	GGCTCAGAAATAGTCCTCT	
HA H5N8$_{LP}$	GCCATTCATCCACAGTTGACA	
	CATGGTGCTGACAGTGGAGTCT	
HA H5N8$_{LP}$ and $_{HP}$	GACCTCTGTTACCCAGGGAGCCT	
	GGACAAGCTGCGCTTACCCCT	
Chicken HPRT1	TGGTGGGGATGACCTCTCAA	(Garrido et al., 2017)
	GGCCGATATCCCACACTTCG	
Chicken GADPH	TCCTCTCTGGCAAAGTCCAAG	(Figueroa et al., 2016)
	CACAACATACTCAGCACCTGC	
Duck GAPDH	CCACTTCCGGGGCACTGTCA	AY436595.1
	AGCACCAGCATCTGCCCACT	
Chicken IFN-γ	ACACTGACAAGTCAAAGCCGC	(Brisbin et al., 2010)
	AGTCGTTCATCGGGAGCTTG	
Duck IFN-γ	ACCTCGTGGAACTGTCAAACCT	AF087134
	CAGCTCACTCACAGCCTTGC	
Chicken Mx	CACTGCAACAAGCAAAGAAGGA	(James et al., 2019)
	TGATCAACCCCACAAGGAAAA	
Duck Mx	TCACACGAAGGCCTATTTTACTGG	Z21549
	GTCGCCGAAGTCATGAAGGA	
Duck RPL4	AAGCTGAACCCATACGCCAA	XM_005015528
	CCTGGGCCTTAGCTGTAACC	

Duck RPL30	GCAAAGCCAAGCTGGTCATC	XM_027452502.1
	CTCAATGTTGTTGCCGCTGT	

573 GAPDH, glyceraldehyde-3-phosphate dehydrogenase; HPRT1, hypoxanthine

574 phosphoribosyltransferase 1; IFN-γ, interferon gamma; Mx, Mx Dynamin Like

575 GTPase; RPL4/30, ribosomal protein L4/L30; F, forward; R, reverse

REFERENCES

Abdelwhab E-SM, Veits J, Mettenleiter TC. 2013. Genetic changes that accompanied shifts of low pathogenic avian influenza viruses toward higher pathogenicity in poultry. *Virulence* **4**:441–452. doi:10.4161/viru.25710

Andreu-Moreno I, Bou J-V, Sanjuán R. 2020. Cooperative nature of viral replication. *Sci Adv* **6**. doi:10.1126/sciadv.abd4942

Baccam P, Beauchemin C, Macken CA, Hayden FG, Perelson AS. 2006. Kinetics of Influenza A Virus Infection in Humans. *Journal of Virology* **80**:7590–7599. doi:10.1128/JVI.01623-05

Barber MRW, Aldridge JR, Webster RG, Magor KE. 2010. Association of RIG-I with innate immunity of ducks to influenza. *Proc Natl Acad Sci USA* **107**:5913–5918. doi:10.1073/pnas.1001755107

Bodewes R, Kuiken T. 2018. Changing Role of Wild Birds in the Epidemiology of Avian Influenza A Viruses. *Adv Virus Res* **100**:279–307. doi:10.1016/bs.aivir.2017.10.007

Böttcher-Friebertshäuser E, Garten W, Matrosovich M, Klenk HD. 2014. The Hemagglutinin: A Determinant of Pathogenicity In: Compans RW, Oldstone MBA, editors. Influenza Pathogenesis and Control - Volume I, Current Topics in Microbiology and Immunology. Cham: Springer International Publishing. pp. 3–34. doi:10.1007/82_2014_384

Bou J-V, Geller R, Sanjuán R. 2019. Membrane-Associated Enteroviruses Undergo Intercellular Transmission as Pools of Sibling Viral Genomes. *Cell Rep* **29**:714-723.e4. doi:10.1016/j.celrep.2019.09.014

Brisbin JT, Gong J, Parvizi P, Sharif S. 2010. Effects of lactobacilli on cytokine expression by chicken spleen and cecal tonsil cells. *Clin Vaccine Immunol* **17**:1337–1343. doi:10.1128/CVI.00143-10

Brooke CB. 2017. Population Diversity and Collective Interactions during Influenza Virus Infection. *Journal of Virology* **91**. doi:10.1128/JVI.01164-17

Burggraaf S, Karpala AJ, Bingham J, Lowther S, Selleck P, Kimpton W, Bean AGD. 2014. H5N1 infection causes rapid mortality and high cytokine levels in chickens compared to ducks. *Virus Res* **185**:23–31. doi:10.1016/j.virusres.2014.03.012

Capua I, Marangon S. 2000. The avian influenza epidemic in Italy, 1999-2000: a review. *Avian Pathol* **29**:289–294. doi:10.1080/03079450050118403

Cornelissen JBWJ, Post J, Peeters B, Vervelde L, Rebel JMJ. 2012. Differential innate responses of chickens and ducks to low-pathogenic avian influenza. *Avian Pathol* **41**:519–529. doi:10.1080/03079457.2012.732691

Cornelissen JBWJ, Vervelde L, Post J, Rebel JMJ. 2013. Differences in highly pathogenic avian influenza viral pathogenesis and associated early inflammatory response in chickens and ducks. *Avian Pathol* **42**:347–364. doi:10.1080/03079457.2013.807325

DaPalma T, Doonan BP, Trager NM, Kasman LM. 2010. A systematic approach to virus–virus interactions. *Virus Research* **149**:1–9. doi:10.1016/j.virusres.2010.01.002

Dhingra MS, Artois J, Dellicour S, Lemey P, Dauphin G, Von Dobschuetz S, Van Boeckel TP, Castellan DM, Morzaria S, Gilbert M. 2018. Geographical and Historical Patterns in the Emergences of Novel Highly Pathogenic Avian Influenza (HPAI) H5 and H7 Viruses in Poultry. *Front Vet Sci* **5**. doi:10.3389/fvets.2018.00084

Domingo-Calap P, Segredo-Otero E, Durán-Moreno M, Sanjuán R. 2019. Social evolution of innate immunity evasion in a virus. *Nat Microbiol* **4**:1006–1013. doi:10.1038/s41564-019-0379-8

EFSA Panel on Animal Health and Welfare (AHAW), More S, Bicout D, Bøtner A, Butterworth
A, Calistri P, Depner K, Edwards S, Garin-Bastuji B, Good M, Gortázar Schmidt C,
Michel V, Miranda MA, Nielsen SS, Raj M, Sihvonen L, Spoolder H, Thulke H-H,
Velarde A, Willeberg P, Winckler C, Breed A, Brouwer A, Guillemain M, Harder T,
Monne I, Roberts H, Baldinelli F, Barrucci F, Fabris C, Martino L, Mosbach-Schulz O,
Verdonck F, Morgado J, Stegeman JA. 2017. Avian influenza. *EFSA J* **15**:e04991.
doi:10.2903/j.efsa.2017.4991
Farci P, Strazzera R, Alter HJ, Farci S, Degioannis D, Coiana A, Peddis G, Usai F, Serra G,
Chessa L, Diaz G, Balestrieri A, Purcell RH. 2002. Early changes in hepatitis C viral
quasispecies during interferon therapy predict the therapeutic outcome. *Proc Natl*
Acad Sci U S A **99**:3081–3086. doi:10.1073/pnas.052712599
Figueroa T, Bessière P, Coggon A, Bouwman KM, van der Woude R, Delverdier M, Verheije
MH, de Vries RP, Volmer R. 2020. The Microbiota Contributes to the Control of
Highly Pathogenic H5N9 Influenza Virus Replication in Ducks. *J Virol* **94**.
doi:10.1128/JVI.00289-20
Figueroa T, Boumart I, Coupeau D, Rasschaert D. 2016. Hyperediting by ADAR1 of a new
herpesvirus lncRNA during the lytic phase of the oncogenic Marek's disease virus.
Journal of General Virology, **97**:2973–2988. doi:10.1099/jgv.0.000606
Garrido D, Chanteloup NK, Trotereau A, Lion A, Bailleul G, Esnault E, Trapp S, Quéré P,
Schouler C, Guabiraba R. 2017. Characterization of the Phospholipid Platelet-
Activating Factor As a Mediator of Inflammation in Chickens. *Front Vet Sci* **4**.
doi:10.3389/fvets.2017.00226
Graaf A, Ulrich R, Maksimov P, Scheibner D, Koethe S, Abdelwhab EM, Mettenleiter TC, Beer
M, Harder T. 2018. A viral race for primacy: co-infection of a natural pair of low and
highly pathogenic H7N7 avian influenza viruses in chickens and embryonated
chicken eggs. *Emerg Microbes Infect* **7**:204. doi:10.1038/s41426-018-0204-0
Grund C, Hoffmann D, Ulrich R, Naguib M, Schinköthe J, Hoffmann B, Harder T, Saenger S,
Zscheppang K, Tönnies M, Hippenstiel S, Hocke A, Wolff T, Beer M. 2018. A novel
European H5N8 influenza A virus has increased virulence in ducks but low zoonotic
potential. *Emerg Microbes Infect* **7**:132. doi:10.1038/s41426-018-0130-1
Guinat C, Nicolas G, Vergne T, Bronner A, Durand B, Courcoul A, Gilbert M, Guérin J-L, Paul
MC. 2018. Spatio-temporal patterns of highly pathogenic avian influenza virus
subtype H5N8 spread, France, 2016 to 2017. *Eurosurveillance* **23**:1700791.
doi:10.2807/1560-7917.ES.2018.23.26.1700791
Hagenaars TJ, Fischer E a. J, Jansen CA, Rebel JMJ, Spekreijse D, Vervelde L, Backer JA, Jong
MCM de, Koets AP. 2016. Modelling the Innate Immune Response against Avian
Influenza Virus in Chicken. *PLOS ONE* **11**:e0157816.
doi:10.1371/journal.pone.0157816
Hiono T, Okamatsu M, Igarashi M, McBride R, de Vries RP, Peng W, Paulson JC, Sakoda Y,
Kida H. 2016. Amino acid residues at positions 222 and 227 of the hemagglutinin
together with the neuraminidase determine binding of H5 avian influenza viruses to
sialyl Lewis X. *Arch Virol* **161**:307–316. doi:10.1007/s00705-015-2660-3
Hoffmann E, Neumann G, Kawaoka Y, Hobom G, Webster RG. 2000. A DNA transfection
system for generation of influenza A virus from eight plasmids. *Proc Natl Acad Sci*
USA **97**:6108–6113. doi:10.1073/pnas.100133697
Hoffmann E, Stech J, Guan Y, Webster RG, Perez DR. 2001. Universal primer set for the full-
length amplification of all influenza A viruses. *Arch Virol* **146**:2275–2289.
Holzinger D, Jorns C, Stertz S, Boisson-Dupuis S, Thimme R, Weidmann M, Casanova J-L,
Haller O, Kochs G. 2007. Induction of MxA Gene Expression by Influenza A Virus
Requires Type I or Type III Interferon Signaling. *Journal of Virology* **81**:7776–7785.
doi:10.1128/JVI.00546-06

Honce R, Karlsson EA, Wohlgemuth N, Estrada LD, Meliopoulos VA, Yao J, Schultz-Cherry S.
2020. Obesity-Related Microenvironment Promotes Emergence of Virulent Influenza
Virus Strains. *mBio* **11**. doi:10.1128/mBio.03341-19
Hughes J, Allen RC, Baguelin M, Hampson K, Baillie GJ, Elton D, Newton JR, Kellam P, Wood
JLN, Holmes EC, Murcia PR. 2012. Transmission of Equine Influenza Virus during an
Outbreak Is Characterized by Frequent Mixed Infections and Loose Transmission
Bottlenecks. *PLOS Pathogens* **8**:e1003081. doi:10.1371/journal.ppat.1003081
James J, Smith N, Ross C, Iqbal M, Goodbourn S, Digard P, Barclay WS, Shelton H. 2019. The
cellular localization of avian influenza virus PB1-F2 protein alters the magnitude of
IFN2 promoter and NFκB-dependent promoter antagonism in chicken cells. *Journal*
of General Virology, **100**:414–430. doi:10.1099/jgv.0.001220
Kleyheeg E, Slaterus R, Bodewes R, Rijks JM, Spierenburg MAH, Beerens N, Kelder L, Poen
MJ, Stegeman JA, Fouchier RAM, Kuiken T, van der Jeugd HP. 2017. Deaths among
Wild Birds during Highly Pathogenic Avian Influenza A(H5N8) Virus Outbreak, the
Netherlands. *Emerg Infect Dis* **23**:2050–2054. doi:10.3201/eid2312.171086
Kuchipudi SV, Tellabati M, Sebastian S, Londt BZ, Jansen C, Vervelde L, Brookes SM, Brown
IH, Dunham SP, Chang K-C. 2014. Highly pathogenic avian influenza virus infection in
chickens but not ducks is associated with elevated host immune and pro-
inflammatory responses. *Vet Res* **45**:118. doi:10.1186/s13567-014-0118-3
Kumar N, Sharma S, Barua S, Tripathi BN, Rouse BT. 2018. Virological and Immunological
Outcomes of Coinfections. *Clinical Microbiology Reviews* **31**.
doi:10.1128/CMR.00111-17
Kuss SK, Etheredge CA, Pfeiffer JK. 2008. Multiple host barriers restrict poliovirus trafficking
in mice. *PLoS Pathog* **4**:e1000082. doi:10.1371/journal.ppat.1000082
Lee D-H, Criado MF, Swayne DE. 2021. Pathobiological Origins and Evolutionary History of
Highly Pathogenic Avian Influenza Viruses. *Cold Spring Harb Perspect Med* **11**.
doi:10.1101/cshperspect.a038679
Leyson C, Youk S-S, Smith D, Dimitrov K, Lee D-H, Larsen LE, Swayne DE, Pantin-Jackwood
MJ. 2019. Pathogenicity and genomic changes of a 2016 European H5N8 highly
pathogenic avian influenza virus (clade 2.3.4.4) in experimentally infected mallards
and chickens. *Virology* **537**:172–185. doi:10.1016/j.virol.2019.08.020
Li S, Bai S, Qin X, Zhang J, Irwin DM, Zhang S, Wang Z. 2019. Comparison of whole embryonic
development in the duck (Anas platyrhynchos) and goose (Anser cygnoides) with the
chicken (Gallus gallus). *Poultry Science* **98**:3278–3291. doi:10.3382/ps/pez133
Magor KE, Miranzo Navarro D, Barber MRW, Petkau K, Fleming-Canepa X, Blyth GAD, Blaine
AH. 2013. Defense genes missing from the flight division. *Developmental &*
Comparative Immunology, Chicken Immunity **41**:377–388.
doi:10.1016/j.dci.2013.04.010
M.c JD, A S, Der GJV, G K. 2009. Intra- and interspecies transmission of H7N7 highly
pathogenic avian influenza virus during the avian influenza epidemic in the
Netherlands in 2003 **28**:333–340. doi:10.20506/rst.28.1.1859
McCrone JT, Lauring AS. 2018. Genetic bottlenecks in intraspecies virus transmission. *Current*
Opinion in Virology, Emerging viruses: intraspecies transmission • Viral Immunology
28:20–25. doi:10.1016/j.coviro.2017.10.008
More S, Bicout D, Bøtner A, Butterworth A, Calistri P, Depner K, Edwards S, Garin-Bastuji B,
Good M, Schmidt CG, Michel V, Miranda MA, Nielsen SS, Raj M, Sihvonen L, Spoolder
H, Thulke HH, Velarde A, Willeberg P, Winckler C, Adlhoch C, Baldinelli F, Breed A,
Brouwer A, Guillemain M, Harder T, Monne I, Roberts H, Abrahantes JC, Mosbach-
Schulz O, Verdonck F, Morgado J, Stegeman A. 2017. Urgent request on avian
influenza. *EFSA Journal* **15**:e04687. doi:https://doi.org/10.2903/j.efsa.2016.4687

Munster VJ, Baas C, Lexmond P, Waldenström J, Wallensten A, Fransson T, Rimmelzwaan GF,
Beyer WEP, Schutten M, Olsen B, Osterhaus ADME, Fouchier RAM. 2007. Spatial,
Temporal, and Species Variation in Prevalence of Influenza A Viruses in Wild
Migratory Birds. *PLOS Pathog* **3**:e61. doi:10.1371/journal.ppat.0030061
Opatowski L, Baguelin M, Eggo RM. 2018. Influenza interaction with cocirculating pathogens
and its impact on surveillance, pathogenesis, and epidemic profile: A key role for
mathematical modelling. *PLOS Pathogens* **14**:e1006770.
doi:10.1371/journal.ppat.1006770
Organization WH. 2008. Addressing ethical issues in pandemic influenza planning: discussion
papers. World Health Organization.
Phipps KL, Ganti K, Jacobs NT, Lee C-Y, Carnaccini S, White MC, Manandhar M, Pickett BE,
Tan GS, Ferreri LM, Perez DR, Lowen AC. 2020. Collective interactions augment
influenza A virus replication in a host-dependent manner. *Nat Microbiol* **5**:1158–
1169. doi:10.1038/s41564-020-0749-2
Poultry species | Gateway to poultry production and products | Food and Agriculture
Organization of the United Nations. n.d. http://www.fao.org/poultry-production-
products/production/poultry-species/en/
Richard M, Fouchier R, Monne I, Kuiken T. 2017. Mechanisms and risk factors for mutation
from low to highly pathogenic avian influenza virus. *EFSA Supporting Publications*
14:1287E. doi:10.2903/sp.efsa.2017.EN-1287
Soubies SM, Hoffmann TW, Croville G, Larcher T, Ledevin M, Soubieux D, Quéré P, Guérin J-L,
Marc D, Volmer R. 2013. Deletion of the C-terminal ESEV domain of NS1 does not
affect the replication of a low-pathogenic avian influenza virus H7N1 in ducks and
chickens. *J Gen Virol* **94**:50–58. doi:10.1099/vir.0.045153-0
van Dijk JG, Verhagen JH, Wille M, Waldenström J. 2018. Host and virus ecology as
determinants of influenza A virus transmission in wild birds. *Curr Opin Virol* **28**:26–
36. doi:10.1016/j.coviro.2017.10.006
Varble A, Albrecht RA, Backes S, Crumiller M, Bouvier NM, Sachs D, García-Sastre A,
tenOever BR. 2014. Influenza A Virus Transmission Bottlenecks Are Defined by
Infection Route and Recipient Host. *Cell Host & Microbe* **16**:691–700.
doi:10.1016/j.chom.2014.09.020
Volmer C, Soubies SM, Grenier B, Guérin J-L, Volmer R. 2011. Immune response in the duck
intestine following infection with low-pathogenic avian influenza viruses or
stimulation with a Toll-like receptor 7 agonist administered orally. *J Gen Virol*
92:534–543. doi:10.1099/vir.0.026443-0
Wahlgren J. 2011. Influenza A viruses: an ecology review. *Infect Ecol Epidemiol* **1**.
doi:10.3402/iee.v1i0.6004
Zwart MP, Elena SF. 2015. Matters of Size: Genetic Bottlenecks in Virus Infection and Their
Potential Impact on Evolution. *Annu Rev Virol* **2**:161–179. doi:10.1146/annurev-
virology-100114-055135

ACKNOWLEDGEMENTS

This work was funded by a grant from the Agence Nationale de la Recherche (ANR-16-CE35-0005-01) to Romain Volmer. Pierre Bessière was supported by a Ph.D. scholarship funded by the Region Occitanie (France) and by the Chaire de Biosécurité at the École Nationale Vétérinaire de Toulouse (French Ministry of Agriculture). Maxime Fusade-Boyer and Gabriel Dupré are supported by PhD scholarships funded by the French Ministry of Research and Education.

AUTHOR CONTRIBUTIONS

Pierre Bessière, Conceptualization, Formal analysis, Investigation, Visualization, Methodology, Writing – original draft; Thomas Figueroa, Conceptualization, Formal analysis, Investigation, Visualization, Methodology, Writing – review and editing; Amelia Coggon, Charlotte Foret-Lucas, Alexandre Houffschmitt, Maxime Fusade-Boyer, Gabriel Dupré, Jean-Luc Guérin, Maxence Delverdier, Investigation, Writing – review and editing; Romain Volmer, Conceptualization, Formal analysis, Investigation, Visualization, Methodology, Funding acquisition, Writing – original draft, Writing – review and editing, Funding acquisition, Project administration.

COMPETING INTERESTS

The authors declare no competing interests.

ETHICS

This study was carried out in compliance with European animal welfare regulation. The protocols were approved by the Animal Care and Use Committee (Comité d'Ethique en Science et Santés Animales – 115) under protocol 13025-2018012311319709.

MATERIALS & CORRESPONDANCE

Correspondence and material requests should be addressed to Romain

Volmer (romain.volmer@envt.fr).

FIGURES

A

B

Fig. 1. **Characterization of H5N8$_{HP}$ and H5N8$_{LP}$ viruses in cell culture. (A)** Sequence alignment of H5N8$_{HP}$ and H5N8$_{LP}$ HA cleavage sites. **(B)** H5N8$_{HP}$ and H5N8$_{LP}$ growth kinetics in the absence or presence of exogenous trypsin in Madin-Darby canine kidney (MDCK) cells, DF-1 cells and CCL-141 cells. Cells were infected at a MOI of 10^{-5} TCID$_{50}$. HA RNA load was analyzed by RT-qPCR using primers common to H5N8$_{HP}$ and H5N8$_{LP}$. The dotted line represents the limit of detection for each experiment. Results are representative of an experiment performed three times.

A

	H2	L2H2	L3H2	L4H2	L5H2	L2	L3	L4	L5
$H5N8_{HP}$	10^2 EID_{50}	10^2 EID_{50}	10^2 EID_{50}	10^2 EID_{50}	10^4 EID_{50}	-	-	-	-
$H5N8_{LP}$	-	10^2 EID_{50}	10^3 EID_{50}	10^4 EID_{50}	10^5 EID_{50}	10^2 EID_{50}	10^3 EID_{50}	10^4 EID_{50}	10^5 EID_{50}

Fig. 2. $H5N8_{HP}$ has a stronger advantage over $H5N8_{LP}$ in embryonated chicken eggs than in embryonated duck eggs. (A) EID_{50} doses used in the *in ovo* experiments. **(B-C)** Chicken **(B)** and duck **(C)** embryonated eggs were infected in the allantoic cavity either with $H5N8_{HP}$ or $H5N8_{LP}$ alone, or with a combination of both. Following a 24 hours incubation at 37°C, viral RNA was extracted from the allantoic fluid and HA levels were determined by RT-qPCR using $H5N8_{HP}$ or $H5N8_{LP}$ specific primers. Statistical analysis: chicken embryonated eggs: n=9 (experiment performed twice); duck embryonated eggs: n=26 (experiment performed three times); one-way ANOVA with Tukey's multiple comparisons test. Results are expressed as means ± SEM. The dotted line represents the limit of detection for each experiment.

Fig. 3. H5N8$_{HP}$ and H5N8$_{LP}$ coinfection in chickens results in increased pathogenesis. Chickens were inoculated via the choanal route with H5N8$_{LP}$ alone (L7), H5N8$_{HP}$ alone (H3 and H4) or with a combination of H5N8$_{LP}$ and H5N8$_{HP}$ (L7H3 and L7H4). **(A)** EID$_{50}$ doses used in the *in vivo* chicken experiments. **(B)** Survival curves. Statistical analysis: Log-rank (Mantel-Cox) test. **(C-E)** Viral shedding was analyzed by quantifying HA RNA levels by RT-qPCR using H5N8$_{HP}$ specific primers from RNA extracted from oropharyngeal swabs **(C)** and cloacal swabs **(D)** or by RT-qPCR using H5N8$_{LP}$ specific primers from RNA extracted from oropharyngeal swabs **(E)**. Numbers of H5N8$_{HP}$ or H5N8$_{HP}$ swab-positive animals are indicated below each time point. Statistical analysis: two-tailed Mann-Whitney test. Results are expressed as means ± SEM. The dotted line represents the limit of detection. **(F-G)** H5N8$_{HP}$ load was analyzed from total RNA extracted from the lungs **(F)** and the brain **(G)** using H5N8$_{HP}$ specific primers. HA RNA levels were normalized using the 2$^{-\Delta Ct}$ method. Statistical analysis: one-way ANOVA with Tukey's multiple comparisons test. Results are expressed as means ± SEM. The dotted line represents the limit of detection. dpi, days post-infection.

Fig. 4. **Immunohistochemical anti-NP staining of hematoxylin-counterstained chickens lung or brain sections.** Analysis was performed on non-infected (NI) and moribund chickens. Scale bar, 50 μm.

Fig. 5. Mx and IFN-γ mRNA expression following chickens infection. (A-B) mRNA expression levels of Mx **(A)** and IFN-γ **(B)** determined by RT-qPCR performed on lung total RNA. **(C)** mRNA expression levels of Mx determined by RT-qPCR performed on RNA extracted from oropharyngeal swabs. mRNA levels were normalized using the $2^{-\Delta Ct}$ method. Statistical analysis: one-way ANOVA with Tukey's multiple comparisons test. Results are expressed as means ± SEM. #, $p < 0.05$ compared to non-infected (NI) animals.

Fig. 6. H5N8_HP and H5N8_LP coinfection in chickens results in higher H5N8_HP growth. (A) Individual H5N8_HP oropharyngeal shedding curves in H4 chickens (blue lines) and L7H4 chickens (red lines). Each curve corresponds to a single animal from the onset of H5N8_HP excretion to death. **(B)** The slopes of individual H5N8_HP excretion curves were calculated. **(C)** H5N8_HP shedding duration: number of days where chickens were H5N8_HP oropharyngeal swab-positive. Statistical analysis: two-tailed Mann-Whitney test. Results are expressed as means ± SEM. The dotted line corresponds to the limit of detection. dpi, days post-infection.

Fig. 7. H5N8$_{HP}$ and H5N8$_{LP}$ coinfection in ducks results in decreased pathogenesis. Ducks were inoculated via the choanal route with H5N8$_{LP}$ alone (L7), H5N8$_{HP}$ alone (H4) or with a combination of H5N8$_{LP}$ and H5N8$_{HP}$ (L7H4). **(A)** EID$_{50}$ doses used in the *in vivo* duck experiments**. (B)** Survival curves. Statistical analysis: Log-rank (Mantel-Cox) test. **(C-E)** Viral shedding was analyzed by quantifying HA RNA levels by RT-qPCR using H5N8$_{HP}$ specific primers from RNA extracted from oropharyngeal swabs **(C)** and cloacal swabs **(D)** or by RT-qPCR using H5N8$_{LP}$ specific primers from RNA extracted from oropharyngeal swabs **(E)**. Numbers of H5N8$_{HP}$ or H5N8$_{HP}$ swab-positive animals are indicated below each time point. Statistical analysis: two-tailed Mann-Whitney test. Results are expressed as means ± SEM. The dotted line represents the limit of detection. **(F-G)** H5N8$_{HP}$ load was analyzed from total RNA extracted from the lungs **(F)** and the brain **(G)** using H5N8$_{HP}$ specific primers. HA RNA levels were normalized using the 2$^{-\Delta Ct}$ method. Statistical analysis: one-way ANOVA with Tukey's multiple comparisons test. Results are expressed as means ± SEM. The dotted line represents the limit of detection. dpi, days post-infection.

Fig. 8. Immunohistochemical anti-NP staining of hematoxylin-counterstained ducks lung or brain sections. Analysis was performed on ducks necropsied at 1 (D1) and 3 days (D3) post infection, moribund ducks and non-infected (NI) ducks. **(A)** Lung. **(B)** Brain. Scale bar, 50 µm.

Fig. 9. **Mx and IFN-γ mRNA expression following infection in ducks. (A-B)** mRNA expression levels of Mx **(A)** and IFN-γ **(B)** determined by RT-qPCR performed on lung total RNA. **(C)** mRNA expression levels of Mx determined by RT-qPCR performed on RNA extracted from oropharyngeal swabs. mRNA levels were normalized using the $2^{-\Delta Ct}$ method. Statistical analysis: one-way ANOVA with Tukey's multiple comparisons test. Results are expressed as means ± SEM. #, p < 0.05 compared to non-infected (NI) animals.

Fig. 10. **Ducks mount a more potent antiviral innate immune response compared to chickens. (A)** Correlation between Mx mRNA levels and viral RNA levels in the lungs. **(B)** Mean Mx/HA ratios. Mx mRNA and HA RNA levels were normalized using the $2^{-\Delta Ct}$ method. Statistical analysis: two-tailed Mann-Whitney test. Results are expressed as means ± SEM.

816