1   **Fungal community turnover along steep environmental gradients from geothermal soils in**
2   **Yellowstone National Park**
3

4   Anna L. Bazzicalupo†*[1], Sonya Erlandson*[2], Margaret Branine[3], Megan Ratz[2], Lauren Ruffing[2],
5   Nhu H. Nguyen[4], Sara Branco[5]
6
[1]Department of Zoology, University of British Columbia, Vancouver, BC
[2] Department of Microbiology and Immunology, Montana State University, Bozeman, MT
[3] Graduate Field of Microbiology, Cornell University, Ithaca, NY
[4] Department of Tropical Plant and Soil Sciences, University of Hawaii at Manoa, HI
[5] Department of Integrative Biology, University of Colorado Denver, Denver, CO
*Authors equally contributed to the study
†Corresponding author: bazzicalupo@zoology.ubc.ca

14
Abstract

Geothermal soils offer unique insight into the way extreme environmental factors shape
communities of organisms. However, little is known about the fungi growing in these
environments and in particular how localized steep abiotic gradients affect fungal diversity. We
used metabarcoding to characterize soil fungi surrounding a hot spring-fed thermal creek with
water up to 84 ℃ and pH 10 in Yellowstone National Park. We found a significant association
between fungal communities and soil variable principal components, and we identify the key
trends in co-varying soil variables that explain the variation in fungal community. Saprotrophic
and ectomycorrhizal fungi community profiles followed, and were significantly associated with,
different soil variable principal components, highlighting potential differences in the factors that
structure these different fungal trophic guilds. In addition, *in vitro* growth experiments in four
target fungal species revealed a wide range of tolerances to pH levels but not to heat. Overall,
our results documenting turnover in fungal species within a few hundred meters suggest many
co-varying environmental factors structure the diverse fungal communities found in the soils of
Yellowstone National Park.

30
31
32 **Keywords**
33 Soil, pH, hot spring, geothermal activity, mycorrhizal, saprobe, *Pisolithus*, *Agaricus*, *Fusarium*
34
35
Declarations

The authors declare no funding.
The authors have no conflicts of interest to declare.
The research did not require ethical approval, no consent to participate or publication was
needed.
All sequences are in GenBank and will be made public after publication: MW471687 -
MW472279.
Code for analyses is available at: https://github.com/abazzical/YellowstoneFungi
All authors have approved this draft of the paper.

Introduction

Soil abiotic factors are known to strongly influence global fungal diversity [1], with soil moisture and chemistry as key drivers of large-scale fungal richness and composition [2]. However, little is known about whether these same drivers function to structure fungal communities at small scales, especially in environments with dramatic small-scale soil abiotic gradients. Geothermal areas are ideal systems to investigate such effects. Thermal sites are geologic units characterized by hot geothermal water, steam, and gases such as carbon dioxide, sulfur dioxide, and hydrogen chloride [3]. They are also notable for increased ambient soil temperature and landscape level thermal features such as hot springs and fumaroles. Thermal areas are by definition hot but can vary greatly in temperature ranging from mildly hot (~30˚C) to boiling point. In addition, geothermal water can cover the range of the pH scale [3]. Thermal soils can therefore display strong edaphic gradients within localized areas, with high variation in soil chemistry, moisture, and temperature.

Thermal areas and hot springs in particular are known for hosting diverse and specialized bacterial and archaeal communities, composed of thermophilic lineages, including novel ones, that evolved to tolerate extreme environmental conditions [4-7]. However, much less is known about microbial eukaryotic communities from thermal areas [8]. For the most part, members of Eukarya lack thermally stable membranes and are much less resilient to high temperatures compared to Archaea and Bacteria [9]. While the vast majority of Fungi fit this pattern, a very small number of thermophilic fungi have been documented. Thermophily evolved independently multiple times across the fungal tree of life [10] and a few species are able to withstand temperatures up to ~60 ºC [11]. Thermophilic fungi have been found in a wide variety of habitats

responding to a range of different environmental pressures and can also occur outside of thermal

areas [12]. Additionally, fungi are well known to withstand other extreme abiotic factors

including broad pH ranges [13]. Several fungal species are able to grow within 5-9 pH unit

differences even when originating from non-extreme habitats [14-16], suggesting that at the local

scale, pH does not contribute strongly to structuring fungal communities. Thermophily in fungi

is rare, so it is likely that in thermal areas, soil temperature may be the main factor driving fungal

richness and composition, while other parameters such as pH are expected to be less relevant for

structuring fungal diversity.

Little is known about the soil fungi from Yellowstone National Park (YNP)'s thermal

habitats that are famous for extremophile research. Available studies have been either based on

fungal culturing from soil (known to detect only a small portion of fungal diversity) or used low-

resolution molecular approaches that limit the ability to fully detect fungal species diversity [17-

20]. Only one very recent metagenomic study describes the mycobiome diversity of two hot

springs in North Sikkim in India where they found the environments dominated by genera of

Ascomycota and detected environmental yeast species [21]. Here we report on the effects of

steep, localized soil abiotic gradients on fungi from Rabbit Creek, an alkaline thermal area in

YNP. We expected geothermal water to form strong abiotic gradients forming in surrounding

soils, directly impacting soil fungal diversity. Specifically, we hypothesized that more extreme

soil conditions (i.e. higher temperature and pH) will host different and species-depauperate

fungal communities relative to less extreme conditions. Given the low fungal tolerance to high

temperatures, we predicted temperature to be the main factor affecting fungal diversity in this

site. We used amplicon sequencing of the fungal ITS rRNA gene to characterize the fungal

communities in soils across a gradient surrounding a hot spring-fed thermal creek with water up

to ~85 ºC and pH ~10. We found soil temperature was much lower than that of nearby boiling

thermal water and covaried with moisture and pH. We found a significant association between

fungal community and environmental soil variables. Although no single variable was a

determining factor driving fungal community composition, soil variables were highly colinear

allowing us to identify major soil chemistry trends that explained turnover in fungal

communities. However, we found saprotrophic fungi and mycorrhizal (tree-associated) fungi

were significantly associated with different sets of soil variables. In addition, we conducted *in*

vitro growth assays testing for temperature and pH tolerance in four target fungal species (three

saprobes and one mycorrhizal species) from Rabbit Creek. Although none of our target fungi

were found to be thermophilic, they strongly differed in the ability to tolerate high or low soil

pH.

Methods

Sampling site and sample processing

We sampled soils around Rabbit Creek, located in the Yellowstone National Park Lower Geyser

Basin, an area characterized by numerous thermal features. Rabbit Creek flows directly out of a

set of hot springs (Table S1, Figure S1). Water temperatures in Rabbit Creek start at

approximately 84 °C at the main hot spring source and cool to 30 °C downstream. The water

stays consistently alkaline at approximately pH 10. Site vegetation consisted of *Pinus contorta*

forest with herbaceous plants including the heat tolerant hot springs panic grass, *Dichanthelium*

lanuginosum (see Stout and Al☐Niemi [22] for a complete geothermal plant survey of

Yellowstone National Park).

In September 2018, we collected a total of 70 soil cores (2.5 × 10 cm) along 14 transects in the

Rabbit Creek area. For each 20 m transect we sampled a total of five cores with one core every 5

m. We sampled eight transects perpendicular to Rabbit Creek (T2-T8) that were 150 m apart

along the creek. We also collected six transects perpendicular to three nearby hot springs (two

transects per hot spring, T1, T9-T13) and an additional transect (T14) away from surface water

and in the pine forest (Table S1).

We measured the soil temperature 10 cm deep next to each soil core and homogenized the soil in

each core before flash freezing and storing approximately 2 g of soil. We saved the remaining

soil for chemical analyses. All cores were processed within 24 h of collection. Soil chemistry and

moisture analyses were conducted at the Environmental Analytical Laboratory (Land Resources

and Environmental Sciences, Montana State University). See Tables S1 and S2 for complete list

of measurements.

DNA extraction and library preparation for sequencing

We extracted genomic DNA from soil with the QIAGEN DNA Power Soil Pro kit using 0.5 g of

soil instead of the recommended 0.25 g. For library preparation, we amplified the ITS1 region as

it detects more taxa than ITS2 and recovers the same community patterns [23]. We used indexed

primers ITS1F [24] and ITS2 [25] in the first round of PCR , followed by a second PCR to attach

the 8 bp barcodes and Illumina adapters, using Hi-Fidelity Phusion DNA Polymerase for both

reactions. Between each amplification step the product was cleaned from reagents and primers

using the NGS SPRI Bead clean-up kit (ABM). Negative PCR controls were sequenced along

with the experimental samples [26]. We included a synthetic mock community for ITS1 as a

positive control to recover tag-switching among samples [27]. Amplicons of equimolar

concentrations were pooled from each of 71 samples. The libraries were sent for paired 250-

nucleotide reads on the Illumina MiSeq sequencing platform at the UC Davis DNA Sequencing

Facility (Davis, California).

Sequence processing

Raw sequence reads were demultiplexed at the UC Davis DNA Sequencing Facility (Davis,

California). We used the QIIME2 software package [28], and a modified QIIME2 pipeline for

fungal ITS by Nguyen [29] with some additional modifications. We quality filtered the

demultiplexed raw sequences (q > 30) setting the truncation of sequences 240 bp for forward and

200 bp for reverse sequences, and denoised the raw reads with DADA2 [30]. After quality

filtering, we *de novo* clustered sequences into operational taxonomic units (OTU) with

VSEARCH at a 97% sequence similarity followed by a re-clustering step with UCLUST at 97%

to ensure the best recovery of our OTUs according to our input mock community [26]. We then

excluded all sequences that were ≤ 85% coverage in the BLAST search since this primer set

amplifies sequences that only partially match to a small part of the ITS gene. We used the

UNITE database v8, [31, 32] for chimera checking, OTU clustering, and assigning taxonomy.

Initial ordination analyses of the OTU community matrix led us to exclude 30 samples as they

clustered due to partial, very short sequences spanning only ~20 nucleotides at the beginning of

the gene making them unidentifiable. In short, an initial non-metric dimensional scaling (NMDS)

ordination including all remaining samples clustered in three groups (Figure S2). A species

indicator analysis using *indicspecies* package (ver. 1.7.8) [33] in R with 9999 permutations

showed the group labelled '3' (Figure S2) clustered due to shared unidentifiable sequences and

in fact, the ITS1 sequences for the OTUs clustered in this group because they could not be

identified. We confirmed this by performing additional BLAST searches. We therefore excluded

cores S5, S6, S7, S8, S9, S60, S63, S64, S67, and S69 from subsequent analyses. These cores

were sampled near the three pools and the forest transect away from surface water (Table S1 and

S2). After filtering, we used 40 cores to perform all further analyses. We produced a species

accumulation curve using the *specaccum* function estimating the mean curve and its standard

deviation from 100 random permutations of the species accumulation data. All scripts and data

tables are available online at https://github.com/abazzical/YellowstoneFungi.

Soil Chemistry and Fungal Community Analysis

We performed Principal Components Analysis (PCA) to assess chemistry similarities across soil

cores. To investigate how the fungal community correlated with soil variables we performed a

Redundancy analysis (RDA) on the OTU table and the first two PCs of the PCA using the vegan

package [34, 35]. For the RDA, we transformed the OTU table by a centered log ratio (CLR).

We first added 1 to the OTU table to make the log transformation possible, then calculated the

CLR transform with the *compositions* R package [36]. We categorized fungi into different

trophic guilds using FUNGuild [37]. We further used the ectomycorrhizal (EM) OTUs and

saprotrophic OTUs for additional RDAs to untangle which variables are more important for each

fungal guild. To test for significant differences in the RDA we performed an ANOVA and

controlled for transects with 1000 permutations within blocks.

To summarize species richness, we calculated Shannon Index in the *vegan* R package [34] and

we report OTU numbers by rarefying the raw OTU table to a sampling depth of 2831. We tested

if the species richness changed with distance from the edge of Rabbit Creek with a linear

regression using the core number as the predictive variable.

Fungal culture isolation and growth assays

We obtained pure cultures for four species growing at the Rabbit Creek site along the transect

sampling sites. We cultured the saprotrophic *Agaricus campestris* and ectomycorrhizal *Pisolithus*

tinctorius from fruitbodies on Potato Dextrose Agar (PDA) and Modified Melin-Norkans

(MMN, [38]) media respectively. We also isolated cultures of potential plant pathogens

Fusarium oxysporum and *F. avenae* from soil by plating serial soil dilutions in PDA. For all four

species, we measured mycelial radial expansion at 30, 35, and 40°C and pH 4, 7, and 9

separately. Each culture was grown in triplicate and measured at 2-4 day intervals. *Fusarium* and

Agaricus cultures were grown on PDA and *Pisolithus* was grown on MMN media, more suitable

for mycorrhizal species. The growth in pH was calculated as the average of four radius

measurements per colony for each time point. The temperature measurements were made by

calculating the area of the fungal colony on the agar plate by importing the drawn outline of the

fungus into ImageJ [39, 40].

We obtained culture identifications by sequencing the ITS rRNA gene. In short, we extracted

genomic DNA using the Extract-N-Amp Sigma kit and amplified the ITS region with primers

ITS1F [24] and ITS4 [25]. The PCR product was Sanger sequenced at GenScript and the

sequences were checked and trimmed for low quality bases in the software FinchTV

(http://www.geospiza.com/finchtv/). To identify the fungi, we used the BLAST tool in the NCBI

database. Sequences are deposited in GenBank (MW471687 - MW472279).

Results

Soil abiotic variation

Soils in the Rabbit Creek area showed wide variation in temperature, moisture content, and

chemistry (Fig. 1, S3). Despite the creek's very high temperature water sources (84ºC), soil

temperature was relatively low and varied between 10 ºC and 31ºC. Soil pH and moisture content

varied broadly, with pH ranging from 4 to 10 and moisture from 10% to 100%. pH and moisture

contributed more than expected to PCA axis 1 and 2, respectively. Soil chemical parameters

varied considerably (Table S1 and S2). Lead (Pb), iron (Fe), total carbon (TC), manganese (Mn),

cobalt (Co), cadmium (Cd) were inversely proportional to alkaline values of pH and were

contributing more than expected to PC1. Phosphorous (PO_4), total nitrogen (TN), moisture

content, copper (Cu), total carbon (TC), and zinc (Zn) were orthogonal to the first set of

parameters and were contributing more than expected to PC2.

Figure 1 – Rabbit Creek soil chemistry variation. (A) Principal components analysis on abiotic parameters measured across soil samples. Colors reflect the soil core number in each transect, core 1 is the closest to Rabbit Creek and core 5 is the furthest away (20 m from the shore). (B) Contribution of variables for PC1 and (C) PC2. Variables with contribution proportion above the red line contribute more than expected to a principal component.

Overall soil fungal community composition

We found a total of 593 fungal OTUs at Rabbit Creek and were able to obtain taxonomic

identifications at a lower taxonomic level than "fungi" for close to 65% of them. This thermal

site was heavily dominated by members of Ascomycota (38.4%) and Basidiomycota (18.7%) but

also included Chytridiomycota, Rozellomycota, Mucoromycota, Glomeromycotina,

Kickxellomycota, Mortierellomycota, and Calcarisporiellomycota. We detected a few fungi

known for thriving at high temperatures (>30 ℃), but not resilient enough to be considered heat

tolerant fungi (with the ability to grow at 40 ˚C [41]). These included *Exophiala opportunistica*,

a black yeast that also withstands high moisture and alkalinity and is commonly found in

dishwashers [42, 43], *Waitea circinata*, a grass pathogen with optimal growth at 25-30 ℃ [44],

Umbelopsis vinacea, a soil fungus that remains active up to 37 ℃ [45], *Gaertneriomyces*

semiglobifer, a chytrid that also growth up to 37 ℃ [46], and *Fusarium kerasplasticarum*, an

opportunistic animal pathogen that can also grow up to 37 ℃. In addition, we found one

unidentified species of *Talaromyces*, a genus that includes thermophilic and thermotolerant

species [47]. Notably, we recovered a number of cold-adapted fungi in our survey, including

Extremus aquaticus, *Elasticomyces elasticus*, *Endophoma alongata*, *Alternaria*

chlamydosporigena, *Mrakiella aquatica*, *Mrakia frigida*, *Tausonia pamirica*, *T. pulullans*,

Naganishia adeliensis, and *Udeniomyces megalosporus*. Many of these species occur in Polar

regions and have documented optimal growth temperatures <20 ℃ [48-53].

Our species accumulation curve based on samples does not plateau (Figure S4), suggesting that

community diversity is not saturated with our sampling.

Redundancy analysis

We performed an RDA between the OTU matrix and the first two PCs of the environmental

variables PCA and we found fungal communities were significantly different and correlated with

the first two PCs of the environmental variables PCA (Figure 2, Table 1). The adjusted R^2 for

this model was 0.5 (Table 1).

Figure 2 – Redundancy analysis of the fungal community and first two PCs of the soil
environmental variable PCA. Both PCs contributed to RDA axes and significantly explained
community composition. The variance explained by each RDA axis is for fitted (constrained)

values. Colors reflect the soil core number in each transect, core 1 is the closest to Rabbit Creek
and core 5 is the furthest away (20 m).

Two hundred and thirty-four (out of 593) OTUs were classified into trophic guilds with

FUNGuild. Twenty two percent (52) OTUs were mycorrhizal (including 16.7% (39) EM fungi),

34.6% (81) saprotrophic, 9.8% (23) plant pathogens, 15% (35) animal pathogens, and 6.8% (16)

dung-associated fungi (Table S3). We performed RDAs on OTU tables composed only of

saprotrophic and EM fungal taxa (Figure 3). Both guilds were found to be significantly

associated with soil variables PCs (Table 1), however, saprotrophic fungi were only significantly

associated with PC2 while EM fungi were significantly associated only with PC1. The total

variance explained (adj. R^2) in the RDA was 9.4% for saprotrophic fungi and 5.7% for EM fungi

(Table 1).

270
Figure 3 – Ectomycorrhizal (EM) and saprotrophic fungal community redundancy analyses. (A)
Saprotrophic fungi are significantly associated with PC2. (B) EM fungi are significantly
associated with PC1. Variables contributing more than expected to PC1 were: pH, Lead (Pb),

iron (Fe), total carbon (TN), manganese (Mn), cobalt (Co), and cadmium (Cd). Variables

contributing more than expected to PC2 were: Phosphorous (PO_4), total nitrogen (TN), moisture

content, copper (Cu), total carbon (TC), and zinc (Zn).

Table 1. Summary of RDA full model analysis of variances (1000 permutations), proportion of

fitted variance explained by RDA axes 1 and 2 for each model, and analysis of variance of

individual predictors (soil principal components) in each RDA model.

RDAs summary statistics						
	All OTUs	residuals	Saprotrophic	residuals	EM	residuals
Adjusted R2	0.05	--	0.094	--	0.057	--
DF	2	45	2	35	2	35
Variance	25.70	269.26	8.70	51.76	5.10	41.80
F	2.15	--	2.94	--	2.14	--
Pf(>F)	**0.000999 *****	--	**0.02597 ***	--	**0.004995 ****	--

RDA1 and RDA2 contributions						
	All OTUs		Saprotrophic		EM	
	RDA1	RDA2	RDA1	RDA2	RDA1	RDA2
Eigenvalue	18.12	7.57	6.18	2.52	3.39	1.72
Proportion Explained	0.71	0.29	0.71	0.29	0.66	0.34
Cumulative Proportion	0.71	1	0.71	1	0.66	1

Individual significance of Principal Components						
	All OTUs		Saprotrophic		EM	
	PC1	PC2	PC1	PC2	PC1	PC2
DF	1	1	1	1	1	1
Variance	16.00	9.97	5.54	3.29	3.39	1.73
F	2.67	1.67	3.75	2.22	2.83	1.45
Pf(>F)	**0.000999 *****	**0.023976 ***	0.07	**0.000999 ***	**0.004995 ***	0.63

Species richness measures

We report OTU numbers and Shannon Index measurements per core. The cores of each transect

represent a soil environmental gradient moving away from the creek shore that is supported by

the soil PCA. We tested if the distance from the creek edge affected the fungal species richness.

We performed linear regressions for the Shannon Index and the OTU numbers against the core

number which reflects the distance from the creek edge. However, we do not find a changing

pattern of species richness away from the creek shore suggesting a change in community

turnover rather than change in number or richness of species (Figure 4). None of the linear

models showed overall significance, however, we found two significant differences for a single

predictive variable in two groups: in OTU numbers for core 2 when we tested all the OTUs

(Estimate=15.929 Std. Error=6.332 t value=2.516 Pr(>|t|)=0.0169) and the saprotrophic OTUs

296
297
the shore, core 3=10m, core 4=15m, core 5=20m. (A) Shannon Index of all fungal OTUs; (B)
Shannon Index of saprotrophic fungal OTUs; (C) Shannon Index of EM fungal OTUs; (D)
number of all fungal OTUs; (E) number of saprotrophic fungal OTUs; (E) number of EM fungal
OTUs. Note the scale bar differences in D-F. Asterisks (*) indicate significance.

Effect of temperature and pH on four fungi from Yellowstone National Park soils

The four fungal species tested for temperature and pH tolerance lacked thermal tolerance but had

the ability to tolerate a wide range of pH (Figures 5 and 6). Notably, all four species performed

well at pH 7. Both *Fusarium oxysporum* and *F. avenae* grew equally well at pH 9 and pH 7

showing less growth at pH 4. Both *Agaricus campestris* and *Pisolithus tinctorius* grew best at pH

7, however, *P. tinctorius* was the only fungus with no growth at pH 9 (Figure 5). All four species

grew well at 30 ℃ but very poorly at 40 ℃ (Figure 6). *P. tinctorius* and *F. oxysporum* grew

equally well at 30 ˚C and 35 ˚C, while *A. campestris* and *F. avenae* grew better at 30 ˚C than 35

˚C. Unfortunately, we did not detect any of the taxa we cultured in our metabarcoding data. This

is not very surprising given the species accumulation curves do not plateau (Figure S4)

indicating our environmental sequencing only detected part of the fungi inhabiting the Rabbit

Creek soil.

Figure 5 – Mycelial growth (average of 4 radius measurement per fungal thallus) at pH 4 (dark blue), pH 7 (green), and pH 9 (yellow) in *Agaricus campestris* (A), *Pisolithus tinctorius* (B), *Fusarium oxysporum* (C), and *Fusarium avenae* (D).

Figure 6 – Mycelial growth (fungus growth area mm^3) at 30ºC (peach), 35ºC (pink), and 40ºC (burgundy) in *Agaricus campestris* (A), *Pisolithus tinctorius* (B), *Fusarium oxysporum* (C), and *Fusarium avenae* (D).

Discussion

We investigated soil fungal communities from a geothermal area in Yellowstone National Park

characterized by a steep abiotic gradient and found a significant association between fungal

communities and soil environmental variables. We summarized the highly co-varied soil

environmental data with PCA and the first two environmental variable PCs significantly

explained variation for the whole fungal community. In addition, saprotrophic and EM fungi

were significantly associated with different components of soil variables. It is known that global

and continental patterns of fungal diversity are shaped by edaphic factors [1, 54]. We present

evidence that the same co-varying edaphic factors that structure fungal diversity at the global

scale also structure fungal communities at a scale of only a few hundred meters. In our study,

these edaphic factors included gradients of co-varied pH, moisture, heavy metals, phosphate,

total carbon and nitrogen. We also found no decline in number or richness measure for OTUs

suggesting that species turnover rather than a reduction in diversity are responsible for the

patterns we reported.

We were surprised by the absence of thermophilic fungi and the presence of cold-adapted

lineages in Rabbit Creek. On the contrary, we detected several cold-adapted species previously

recorded in the region. Specifically, we found the psychrophilic basidiomycete yeast *Tausonia*

pulullans [55] and a few cold adapted Ascomycete yeasts identified in Vu *et al.* (2016) [56],

including *Naganishia adeliensis* and *Udenomyces megalosporus* (=*U. pyricola* in Vu et al 2016).

We also found the psychrophiles *Elasticomyces elasticus* and *Mrakia sp.*, that had been detected

in previous soil surveys in both West Yellowstone and Hyalite near Cooke City, Montana [57].

Additionally, our *Tausonia pamirica* OTU matched 100% to the sequence of the type specimen

(Genbank # NR_154490), a known psychrotolerant yeast from Antarctica [58]. The occurrence

of these species is most likely unrelated to the thermal nature of Rabbit Creek and rather may

reflect the regional climate characterized by severe cold winters. We were also surprised that

none of the fungi we isolated in culture were able to grow at temperatures higher than 35˚C. Soil

temperature is thought to be the most significant factor influencing bacterial and archaeal species

composition in geothermally heated environments [22, 59, 60]. Thermophilic fungi are mostly

known from studies on plant-associated fungi from geothermal environments [61] and surveys

such as the YNP metagenome project [62]. There is also experimental evidence documenting

previously unknown thermophilic and acidophilic fungal species from hot spring habitats [63].

However, the soil temperature of our sampled cores was relatively low, ranging from 10 ºC to

31ºC, which is well within the range of known temperature tolerance of most fungi including the

detected psychrophilic species [55, 56] and the fungi tested in this study (Fig. 6). It is important

to note that our findings do not imply the absence of heat-adapted fungi in Yellowstone, as other

thermal areas characterized by warmer soils may well host thermophilic fungi of unknown

taxonomic identity.

Conditions in highly acidic or alkaline environments often co-vary with other environmentally

extreme variables such as elevated salt concentrations or temperatures[64], and in our case, soil

moisture content. It is known that edaphic parameters structure communities of soil fungi, and

variation in chemistry (including pH) affects fungal diversity on global, continental and local

scales [1, 2]. Individual fungi appear to have a wider pH tolerance range compared to bacteria

[65], however there are several examples of fungi (including *P. tinctorus*, which was studied

here) being able to withstand both low and high pH levels [14]. Studies on acid- or alkaline-

loving fungi have recorded new species and have expanded our knowledge of the possible

habitat types for commonly found fungal species [13]. In addition, research on fungi associated

with hot spring sediment across the globe has documented the phylum Ascomycota as dominant

in these habitats. These included newly discovered species [63] and common members of

widespread genera, such as *Aspergillus*, *Penicillium*, and *Cladosporium*, consistent with taxa

found in hot spring environments in North Sikkim, India [21]. Mycorrhizal species belonging to

lineage Ascomycota have also been detected soda lake habitats [66]. We found the Rabbit Creek

soils with higher pH are inhabited by many members of the Ascomycota lineage, these include

several alkalitolerant fungi such as *Acremonium* from the Emericellopsis clade [67].

We found differences on the effect of edaphic parameters across fungal guilds. EM fungal composition was significantly associated with the axis influenced by heavy metals, pH values, and carbon, while saprotrophic fungi were significantly associated with an axis influenced by phosphorous, nitrogen, moisture, carbon (to a lesser extent than the first axis), and copper and zinc. Additionally, the number of saprotrophic OTUs significantly changed with distance from the creek shore, with an uptick at an intermediate distance while there was no difference found in EM fungi. Interestingly, mycorrhizal fungi have been predicted to display more restricted ranges compared to saprotrophs [68], although global empirical data failed to recover this pattern [69]. This would lead to a prediction of edaphic factors affecting mycorrhizal fungi disproportionately, which is the opposite of what we found. Instead, our study suggests either that the ability to associate with a host allows mycorrhizal fungi to inhabit any micro-habitat where the host is present or that saprotrophic fungi might be more sensitive to environmental variables when resource competition and successional factors are removed.

In conclusion, we found statistical support suggesting soil variables structured fungal community composition at the scale of only a few hundred meters. We also found thermophilic fungi to be absent from the soils surrounding thermal water in Rabbit Creek. Further research on thermal areas characterized by extreme soil conditions is however needed improving our understanding of the effects of extreme edaphic factors on the ecology and evolution of fungi.

Acknowledgements

We thank Maddie Trent, Rio Wofford, Colin Kennedy, Seamus Hoolahan, and Kathryn Gannon for assistance in the field and laboratory, and the Quandt lab for suggestions on the manuscript. Cathy Zabinski helped obtaining the collecting permit, Dan Colman assisted with sampling

design, and Kabir Peay provided raw data from another study that allowed fungal community

comparisons. In addition, we thank three anonymous reviewers for crucial input on previous

versions of this study. We also thank Yellowstone National Park for allowing the collection of

samples in Rabbit Creek (YELL-2018-SCI8062).

Cited Literature

1. Tedersoo L, Bahram M, Põlme S, Kõljalg U, Yorou NS, Wijesundera R, Ruiz LV, Vasco-Palacios AM, Thu PQ, Suija A (2014) Global diversity and geography of soil fungi. Science 346: 1256688.
2. Peay KG, Kennedy PG, Talbot JM (2016) Dimensions of biodiversity in the Earth mycobiome. Nature Reviews Microbiology 14: 434-447. doi: 10.1038/nrmicro.2016.59
3. Amenábar M, Urschel M, Boyd E (2015) Metabolic and taxonomic diversification in continental magmatic hydrothermal systems. In: Bakermans, C (ed.) Microbial evolution under extreme conditions. De Gruyter, Berlin, Boston, pp. 57-96
4. Brock TD, Freeze H (1969) *Thermus aquaticus* gen. n. and sp. n., a Nonsporulating Extreme Thermophile. J Bacteriol 98: 289.
5. Meyer-Dombard DR, Shock EL, Amend JP (2005) Archaeal and bacterial communities in geochemically diverse hot springs of Yellowstone National Park, USA. Geobiology 3: 211-227. doi: https://doi.org/10.1111/j.1472-4669.2005.00052.x
6. Colman DR, Jay ZJ, Inskeep WP, Jennings Rd, Maas KR, Rusch DB, Takacs-Vesbach CD (2016) Novel, Deep-Branching Heterotrophic Bacterial Populations Recovered from Thermal Spring Metagenomes7. doi: 10.3389/fmicb.2016.00304
7. Colman DR, Lindsay MR, Boyd ES (2019) Mixing of meteoric and geothermal fluids supports hyperdiverse chemosynthetic hydrothermal communities. Nature Communications 10: 681. doi: 10.1038/s41467-019-08499-1
8. Brown SR, Fritz SC (2019) Eukaryotic organisms of continental hydrothermal systems. Extremophiles 23: 367-376. doi: 10.1007/s00792-019-01101-y
9. Tansey MR, Brock TD (1972) The Upper Temperature Limit for Eukaryotic Organisms. Proceedings of the National Academy of Sciences 69: 2426. doi: 10.1073/pnas.69.9.2426
10. Morgenstern I, Powlowski J, Ishmael N, Darmond C, Marqueteau S, Moisan M-C, Quenneville G, Tsang A (2012) A molecular phylogeny of thermophilic fungi. Fungal Biology 116: 489-502. doi: https://doi.org/10.1016/j.funbio.2012.01.010
11. Hutchinson MI, Powell AJ, Herrera J, Natvig DO (2019) New Perspectives on the Distribution and Roles of Thermophilic Fungi. In: Tiquia-Arashiro, SM, Grube, M (eds.) Fungi in Extreme Environments: Ecological Role and Biotechnological Significance. Springer International Publishing, Cham, pp. 59-80
12. Powell AJ, Parchert KJ, Bustamante JM, Ricken JB, Hutchinson MI, Natvig DO (2012) Thermophilic fungi in an aridland ecosystem. Mycologia 104: 813-825. doi: 10.3852/11-298

441   13.   Aguilera A, González-Toril E (2019) Eukaryotic Life in Extreme Environments:
442   Acidophilic Fungi. In: Tiquia-Arashiro, SM, Grube, M (eds.) Fungi in Extreme
443   Environments: Ecological Role and Biotechnological Significance. Springer International
444   Publishing, Cham, pp. 21-38
445   14.   Hung L-L, Trappe JM (1983) Growth Variation Between and Within Species of
446   Ectomycorrhizal Fungi in Response to pH in Vitro. Mycologia 75: 234-241. doi:
447   10.1080/00275514.1983.12021660
448   15.   Wheeler KA, Hurdman BF, Pitt JI (1991) Influence of pH on the growth of some
449   toxigenic species of *Aspergillus*, *Penicillium* and *Fusarium*. Int J Food Microbiol 12:
450   141-149. doi: https://doi.org/10.1016/0168-1605(91)90063-U
451   16.   Nevarez L, Vasseur V, Le Madec A, Le Bras MA, Coroller L, Leguérinel I, Barbier G
452   (2009) Physiological traits of *Penicillium glabrum* strain LCP 08.5568, a filamentous
453   fungus isolated from bottled aromatised mineral water. Int J Food Microbiol 130: 166-
454   171. doi: https://doi.org/10.1016/j.ijfoodmicro.2009.01.013
455   17.   Redman RS, Litvintseva A, Sheehan KB, Henson JM, Rodriguez RJ (1999) Fungi from
456   Geothermal Soils in Yellowstone National Park. Appl Environ Microbiol 65: 5193. doi:
457   10.1128/AEM.65.12.5193-5197.1999
458   18.   Appoloni S, Lekberg Y, Tercek MT, Zabinski CA, Redecker D (2008) Molecular
459   Community Analysis of Arbuscular Mycorrhizal Fungi in Roots of Geothermal Soils in
460   Yellowstone National Park (USA). Microb Ecol 56: 649-659.
461   19.   Lekberg Y, Meadow J, Rohr JR, Redecker D, Zabinski CA (2011) Importance of
462   dispersal and thermal environment for mycorrhizal communities: lessons from
463   Yellowstone National Park. Ecology 92: 1292-1302. doi: https://doi.org/10.1890/10-
464   1516.1
465   20.   Meadow JF, Zabinski CA (2012) Spatial heterogeneity of eukaryotic microbial
466   communities in an unstudied geothermal diatomaceous biological soil crust: Yellowstone
467   National Park, WY, USA. FEMS Microbiol Ecol 82: 182-191. doi: 10.1111/j.1574-
468   6941.2012.01416.x %J FEMS Microbiology Ecology
469   21.   Das S, Roy G, Najar IN, Sherpa MT, Thakur N (2021) Diversity and composition of the
470   North Sikkim hot spring mycobiome using a culture-independent method. Folia
471   Microbiol 66: 457-468. doi: 10.1007/s12223-021-00859-z
472   22.   Stout RG, Al☐Niemi TS (2002) Heat☐tolerant Flowering Plants of Active Geothermal
473   Areas in Yellowstone National Park. Ann Bot 90: 259-267. doi: 10.1093/aob/mcf174 %J
474   Annals of Botany
475   23.   Bazzicalupo AL, Bálint M, Schmitt I (2013) Comparison of ITS1 and ITS2 rDNA in 454
476   sequencing of hyperdiverse fungal communities. Fungal Ecology 6: 102-109.
477   24.   Gardes M, Bruns TD (1993) ITS primers with enhanced specificity for
478   basidiomycetes☐application to the identification of mycorrhizae and rusts. Mol Ecol 2:
479   113-118. doi: 10.1111/j.1365-294X.1993.tb00005.x
480   25.   White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal
481   ribosomal RNA genes for phylogenetics. In: Innis, M, Gelfand, D, Sninsky, J, White, TJ
482   (eds.) PCR protocols: a guide to methods and applications. Academic Press Inc., New
483   York, pp. 315-322
484   26.   Nguyen NH, Smith D, Peay K, Kennedy P (2015) Parsing ecological signal from noise in
485   next generation amplicon sequencing. New Phytol 205: 1389-1393. doi:
486   https://doi.org/10.1111/nph.12923

27. Palmer JM, Jusino MA, Banik MT, Lindner DL (2018) Non-biological synthetic spike-in controls and the AMPtk software pipeline improve mycobiome data. PeerJ 6: e4925. doi: 10.7717/peerj.4925

28. Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, Alexander H, Alm EJ, Arumugam M, Asnicar F, Bai Y, Bisanz JE, Bittinger K, Brejnrod A, Brislawn CJ, Brown CT, Callahan BJ, Caraballo-Rodríguez AM, Chase J, Cope EK, Da Silva R, Diener C, Dorrestein PC, Douglas GM, Durall DM, Duvallet C, Edwardson CF, Ernst M, Estaki M, Fouquier J, Gauglitz JM, Gibbons SM, Gibson DL, Gonzalez A, Gorlick K, Guo J, Hillmann B, Holmes S, Holste H, Huttenhower C, Huttley GA, Janssen S, Jarmusch AK, Jiang L, Kaehler BD, Kang KB, Keefe CR, Keim P, Kelley ST, Knights D, Koester I, Kosciolek T, Kreps J, Langille MGI, Lee J, Ley R, Liu Y-X, Loftfield E, Lozupone C, Maher M, Marotz C, Martin BD, McDonald D, McIver LJ, Melnik AV, Metcalf JL, Morgan SC, Morton JT, Naimey AT, Navas-Molina JA, Nothias LF, Orchanian SB, Pearson T, Peoples SL, Petras D, Preuss ML, Pruesse E, Rasmussen LB, Rivers A, Robeson MS, Rosenthal P, Segata N, Shaffer M, Shiffer A, Sinha R, Song SJ, Spear JR, Swafford AD, Thompson LR, Torres PJ, Trinh P, Tripathi A, Turnbaugh PJ, Ul-Hasan S, van der Hooft JJJ, Vargas F, Vázquez-Baeza Y, Vogtmann E, von Hippel M, Walters W, Wan Y, Wang M, Warren J, Weber KC, Williamson CHD, Willis AD, Xu ZZ, Zaneveld JR, Zhang Y, Zhu Q, Knight R, Caporaso JG (2019) Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. Nat Biotechnol 37: 852-857. doi: 10.1038/s41587-019-0209-9

29. Nguyen N (2020) Amplicon analysis pipeline with QIIME2: https://github.com/nnguyenlab/amplicon-pipeline.

30. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP (2016) DADA2: High-resolution sample inference from Illumina amplicon data. Nat Methods 13: 581-583. doi: 10.1038/nmeth.3869

31. Abarenkov K, Nilsson RH, Larsson KH, Alexander IJ, Eberhardt U, Erland S, Høiland K, Kjøller R, Larsson E, Pennanen T (2010) The UNITE database for molecular identification of fungi–recent updates and future perspectives. New Phytol 186: 281-285.

32. Kõljalg U (2020) UNITE general FASTA release for Fungi. In: community, U (ed.), 10.15156/BIO/786368.

33. De Caceres M, Jansen F, De Caceres MJRp, Version (2016) Indicspecies: relationship between species and groups of sites1: 2014.

34. Dixon P (2003) VEGAN, a package of R functions for community ecology. Journal of Vegetation Science 14: 927-930. doi: https://doi.org/10.1111/j.1654-1103.2003.tb02228.x

35. Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin P, O'hara R, Simpson G, Solymos P, Stevens M, Wagner H (2017) Vegan: community ecology package. R package. Version.

36. van den Boogaart KG, Tolosana-Delgado R (2008) "compositions": A unified R package to analyze compositional data. Computers & Geosciences 34: 320-338. doi: https://doi.org/10.1016/j.cageo.2006.11.017

37. Nguyen NH, Song Z, Bates ST, Branco S, Tedersoo L, Menke J, Schilling JS, Kennedy PG (2016) FUNGuild: An open annotation tool for parsing fungal community datasets by ecological guild. Fungal Ecology 20: 241-248. doi: https://doi.org/10.1016/j.funeco.2015.06.006

38. Marx DH (1970) The influence of ectotrophic mycorrhizal fungi on the resistance of Pine
roots to pathogenic infections. V. Resistance of mycorrhizae to infection by vegetative
mycelium of Phytophthora cinnamomi. Phytopathology 60: 1472-1473. doi:
10.1094/Phyto-60-1472
39. Abràmoff MD, Magalhães PJ, Ram SJ (2004) Image processing with ImageJ.
Biophotonics international 11: 36-42.
40. Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of
image analysis. Nat Methods 9: 671-675.
41. Mouchacca J (2007) Heat tolerant fungi and applied research: Addition to the previously
treated group of strictly thermotolerant species. World Journal of Microbiology and
Biotechnology 23: 1755. doi: 10.1007/s11274-007-9426-3
42. Zalar P, Novak M, de Hoog GS, Gunde-Cimerman N (2011) Dishwashers – A man-made
ecological niche accommodating human opportunistic fungal pathogens. Fungal Biology
115: 997-1007. doi: https://doi.org/10.1016/j.funbio.2011.04.007
43. Gümral R, Özhak-baysan B, Tümgör A, Saraçli MA, Yildiran ST, Ilkit M, Zupancic J,
Novak-babic M, Gunde-cimerman N, Zalar P, de Hoog GS (2016) Dishwashers provide a
selective extreme environment for human-opportunistic yeast-like fungi. Fungal
Diversity 76: 1-9. doi: http://dx.doi.org/10.1007/s13225-015-0327-8
44. de la Cerda KA, Douhan GW, Wong FP (2007) Discovery and Characterization of
Waitea circinata var. *circinata* Affecting Annual Bluegrass from the Western United
States. Plant Dis 91: 791-797. doi: 10.1094/PDIS-91-7-0791
45. Wang Y-N, Liu X-Y, Zheng R-Y (2014) Umbelopsis changbaiensis sp. nov. from China
and the typification of *Mortierella vinacea*. Mycological Progress 13: 657-669. doi:
10.1007/s11557-013-0948-9
46. Henderson L, Ly M-A, Robinson K, Gleason FH, Lilje O (2018) Maximum temperature
for growth and reproduction is similar in two soil isolates of *Gaertneriomyces*
semiglobifer (Spizellomycetales, Chytridiomycetes) from different soil environments in
north eastern New South Wales, Australia. Nova Hedwigia 106: 485-497. doi:
10.1127/nova_hedwigia/2017/0448
47. Houbraken J, Spierenburg H, Frisvad JC (2012) *Rasamsonia*, a new genus comprising
thermotolerant and thermophilic *Talaromyces* and *Geosmithia* species. Antonie Van
Leeuwenhoek 101: 403-421. doi: 10.1007/s10482-011-9647-1
48. Selbmann L, Zucconi L, Isola D, Onofri S (2015) Rock black fungi: excellence in the
extremes, from the Antarctic to space. Curr Genet 61: 335-345. doi: 10.1007/s00294-014-
0457-7
49. Hirose D, Hobara S, Matsuoka S, Kato K, Tanabe Y, Uchida M, Kudoh S, Osono T
(2016) Diversity and community assembly of moss-associated fungi in ice-free coastal
outcrops of continental Antarctica. Fungal Ecology 24: 94-101. doi:
https://doi.org/10.1016/j.funeco.2016.09.005
50. Řezanka T, Kolouchová I, Sigler K (2016) Lipidomic analysis of psychrophilic yeasts
cultivated at different temperatures. Biochim Biophys Acta 1861: 1634-1642. doi:
https://doi.org/10.1016/j.bbalip.2016.07.005
51. Thomas-Hall SR, Turchetti B, Buzzini P, Branda E, Boekhout T, Theelen B, Watson K
(2010) Cold-adapted yeasts from Antarctica and the Italian Alps—description of three
novel species: *Mrakia robertii* sp. nov., *Mrakia blollopis* sp. nov. and *Mrakiella*
niccombsii sp. nov. Extremophiles 14: 47-59. doi: 10.1007/s00792-009-0286-7

52.   Vishniac HS (1999) Psychrophilic Yeasts. In: Seckbach, J (ed.) Enigmatic
Microorganisms and Life in Extreme Environments. Springer Netherlands, Dordrecht,
pp. 315-321

53.   Białkowska AM, Szulczewska KM, Krysiak J, Florczak T, Gromek E, Kassassir H, Kur
J, Turkiewicz M (2017) Genetic and biochemical characterization of yeasts isolated from
Antarctic soil samples. Polar Biol 40: 1787-1803. doi: 10.1007/s00300-017-2102-7

54.   Talbot JM, Bruns TD, Taylor JW, Smith DP, Branco S, Glassman SI, Erlandson S,
Vilgalys R, Liao H-L, Smith ME, Peay KG (2014) Endemism and functional
convergence across the North American soil mycobiome. Proceedings of the National
Academy of Sciences 111: 6341. doi: 10.1073/pnas.1402584111

55.   Cavello I, Albanesi A, Fratebianchi D, Garmedia G, Vero S, Cavalitto S (2017)
Pectinolytic yeasts from cold environments: novel findings of *Guehomyces pullulans*,
Cystofilobasidium infirmominiatum and *Cryptococcus adeliensis* producing pectinases.
Extremophiles 21: 319-329. doi: http://dx.doi.org/10.1007/s00792-016-0904-0

56.   Vu D, Groenewald M, Szöke S, Cardinali G, Eberhardt U, Stielow B, de Vries M,
Verkleij GJM, Crous PW, Boekhout T, Robert V (2016) DNA barcoding analysis of
more than 9 000 yeast isolates contributes to quantitative thresholds for yeast species
and genera delimitation. Studies in Mycology 85: 91-105. doi:
https://doi.org/10.1016/j.simyco.2016.11.007

57.   Steidinger BS, Bhatnagar JM, Vilgalys R, Taylor JW, Qin C, Zhu K, Bruns TD, Peay KG
(2020) Ectomycorrhizal fungal diversity predicted to substantially decline due to climate
changes in North American Pinaceae forests. J Biogeogr 47: 772-782. doi:
https://doi.org/10.1111/jbi.13802

58.   Tsuji M, Kudoh S (2020) Soil Yeasts in the Vicinity of Syowa Station, East Antarctica:
Their Diversity and Extracellular Enzymes, Cold Adaptation Strategies, and Secondary
Metabolites. Sustainability 12. doi: 10.3390/su12114518

59.   Sharp CE, Brady AL, Sharp GH, Grasby SE, Stott MB, Dunfield PF (2014) Humboldt's
spa: microbial diversity is controlled by temperature in geothermal environments. The
ISME Journal 8: 1166-1174. doi: 10.1038/ismej.2013.237

60.   de Oliveira TB, Rodrigues A (2019) Ecology of Thermophilic Fungi. In: Tiquia-Arashiro,
SM, Grube, M (eds.) Fungi in Extreme Environments: Ecological Role and
Biotechnological Significance. Springer International Publishing, Cham, pp. 39-57

61.   Zhou W-N, White JF, Soares MA, Torres MS, Zhou Z-P, Li H-Y (2015) Diversity of
fungi associated with plants growing in geothermal ecosystems and evaluation of their
capacities to enhance thermotolerance of host plants. Journal of Plant Interactions 10:
305-314. doi: 10.1080/17429145.2015.1101495

62.   Inskeep W, Jay Z, Tringe S, Herrgard M, Rusch D (2013) The YNP Metagenome Project:
Environmental Parameters Responsible for Microbial Distribution in the Yellowstone
Geothermal Ecosystem4. doi: 10.3389/fmicb.2013.00067

63.   Yamazaki A, Toyama K, Nakagiri A (2010) A new acidophilic fungus *Teratosphaeria*
acidotherma (Capnodiales, Ascomycota) from a hot spring. Mycoscience 51: 443-455.
doi: 10.1007/s10267-010-0059-2

64.   Orwa P, Mugambi G, Wekesa V, Mwirichia R (2020) Isolation of haloalkaliphilic fungi
from Lake Magadi in Kenya. Heliyon 6: e02823. doi:
https://doi.org/10.1016/j.heliyon.2019.e02823

65.   Rousk J, Bååth E, Brookes PC, Lauber CL, Lozupone C, Caporaso JG, Knight R, Fierer
624   N (2010) Soil bacterial and fungal communities across a pH gradient in an arable soil.
625   The ISME Journal 4: 1340-1351. doi: 10.1038/ismej.2010.58
626   66.   Ishida TA, Nara K, Ma S, Takano T, Liu S (2009) Ectomycorrhizal fungal community in
627   alkaline-saline soil in northeastern China. Mycorrhiza 19: 329-335. doi:
628   http://dx.doi.org/10.1007/s00572-008-0219-9
629   67.   Grum-Grzhimaylo AA, Georgieva ML, Debets AJM, Bilanenko EN (2013) Are
630   alkalitolerant fungi of the Emericellopsis lineage (Bionectriaceae) of marine origin? IMA
631   Fungus 4: 213-228. doi: 10.5598/imafungus.2013.04.02.07
632   68.   Sato H, Tsujino R, Kurita K, Yokoyama K, Agata K (2012) Modelling the global
633   distribution of fungal species: new insights into microbial cosmopolitanism. Mol Ecol 21:
634   5599-5612.
635   69.   Bazzicalupo AL, Whitton J, Berbee ML (2019) Over the hills, but how far away?
636   Estimates of mushroom geographic range extents. J Biogeogr 46: 1547-1557. doi:
637   10.1111/jbi.13617
638