Fungal communities from geothermal soils in Yellowstone National Park

Anna Bazzicalupo*[1], Sonya Erlandson[2], Margaret Branine[3], Megan Ratz[2], Lauren Ruffing[2], Nhu H. Nguyen[4], Sara Branco[5]

[1]Department of Zoology, University of British Columbia, Vancouver, BC
[2] Department of Microbiology and Immunology, Montana State University, Bozeman, MT
[3] Department of Microbiology, Cornell University, Ithaca, NY
[4] Department of Tropical Plant and Soil Sciences, University of Hawaii, Honolulu, HI
[5] Department of Integrative Biology, University of Colorado Denver, Denver, CO
*Corresponding author: bazzicalupo@zoology.ubc.ca

Abstract

Geothermal soils offer unique insight into the way extreme environmental factors shape communities of organisms. However, little is known about the fungi growing in these environments and in particular how localized steep abiotic gradients affect fungal diversity. We used metabarcoding to characterize soil fungi surrounding a hot spring-fed thermal creek with water up to ~85 ℃ and pH ~10 in Yellowstone National Park. No soil variable we measured determined fungal community composition. However, soils with pH >8 had lower fungal richness and different fungal assemblages when compared to less extreme soils. Saprotrophic fungi community profile followed more closely overall community patterns while ectomycorrhizal fungi did not, highlighting potential differences in the factors that structure these different fungal trophic guilds. In addition, *in vitro* growth experiments in four target fungal species revealed a wide range of tolerances to pH levels but not to heat. Overall, our results documenting fungal communities within a few hundred meters suggest stronger statistical power and wider sampling are needed to untangle so many co-varying environmental factors affecting such diverse species communities.

Keywords
Soil, pH, hot spring, geothermal activity, mycorrhizal, saprobe, *Pisolithus*, *Agaricus*, *Fusarium*

Introduction

Soil abiotic factors are known to strongly influence global fungal diversity [1], with soil

moisture and chemistry as key drivers of large-scale fungal richness and composition [2].

However, little is known about whether these same drivers function to structure fungal

communities at a scale of <1,000 m, especially in environments with dramatic small-scale soil

abiotic gradients. Geothermal areas are ideal systems to investigate such effects. Thermal sites

are geologic units characterized by hot geothermal water, steam, and gases such as carbon

dioxide, sulfur dioxide, and hydrogen chloride. They are also notable for increased ambient soil

temperature and landscape level thermal features such as hot springs and fumaroles. Thermal

areas are by definition hot but can vary greatly in temperature ranging from mildly hot (~30˚C)

to boiling point. In addition, geothermal water can cover the range of the pH scale [3]. Thermal

soils can therefore display strong edaphic gradients within localized areas, with high variation in

soil chemistry, moisture, and temperature.

Thermal areas and hot springs in particular are known for hosting diverse and specialized

bacterial and archaeal communities, composed of thermophilic lineages, including novel ones,

that evolved to tolerate these extreme environmental conditions [4-7]. However, much less is

known about eukaryotic communities from thermal areas [8]. For the most part, members of

Eukarya lack thermally stable membranes and are much less resilient to high temperatures

compared to Archaea and Bacteria [9]. While the vast majority of Fungi fit this pattern, a very

small number of thermophilic fungi have been documented. Thermophily evolved independently

multiple times across the fungal tree of life [10] and a few species are able to withstand

temperatures up to ~60 ℃ [11]. Thermophilic fungi have been found in a wide variety of habitats

responding to a variety of different environmental pressures and can also occur outside of

thermal areas [12]. Additionally, fungi are well known to withstand a range of other extreme

abiotic factors including broad pH ranges [13]. Several fungal species withstand 5-9 pH unit

differences even when originating from non-extreme habitats [14-16], thus indicating that

perhaps at the local scale, pH does not contribute strongly to structuring fungal communities.

The rarity of thermophily in fungi, as opposed to tolerance of other environmental factors, make

it logical to hypothesize that in thermal areas, soil temperature is likely the main factor driving

fungal richness and composition, while other parameters such as pH are expected to be less

relevant for structuring fungal diversity.

Little is known about the soil fungi from Yellowstone National Park (YNP)'s thermal

habitats which are famous for extremophile research. Available studies have been either based on

fungal culturing from soil (known to detect only a small portion of fungal diversity) or used low-

resolution molecular approaches that do not allow to fully document fungal species diversity [17-

20]. Here we report on the effects of steep, localized soil abiotic gradients on fungi from an

alkaline thermal area in YNP. We expected geothermal water to form strong abiotic gradients

forming in surrounding soils, directly impacting soil fungal diversity. Specifically, we

hypothesized that more extreme soil conditions (with higher temperature and pH) will host

differentiated and depauperate fungal communities relative to less extreme conditions. Given the

low fungal tolerance to high temperatures, we predicted temperature to be the main factor

affecting fungal diversity in this site. We used amplicon sequencing of the fungal ITS rRNA

gene to characterize the fungal communities in soils across a gradient surrounding a hot spring-

fed thermal creek with water up to ~85 ℃ and pH ~10. We found soil temperature was much

lower than that of nearby boiling thermal water and covaried with moisture and pH. Our data

indicated that no single variable was a determining factor driving fungal community

composition. However, we found the highest pH (>8) hosted lower fungal richness when

compared to less extreme soils. We also found saprotrophic fungi matched the pattern in the

overall fungal community much more closely than mycorrhizal (tree-associated) fungi. In

addition, we conducted *in vitro* growth assays testing for temperature and pH tolerance in four

target fungal species (three saprobes and one mycorrhizal species) from Rabbit Creek. Although

none of our target fungi were found to be thermophilic, they strongly differed in the ability to

tolerate high or low soil pH.

Methods

Sampling site and sample processing

We sampled soils around Rabbit Creek, located in the Yellowstone National Park Lower Geyser

Basin which is a thermal area characterized by numerous thermal features. Rabbit Creek flows

directly out of a set of hot springs (Table S1, Figure S1). Water temperatures in Rabbit Creek

start at approximately 84 °C at the main hot spring source and cool to 30 °C downstream. The

water stays consistently alkaline at approximately pH 10. Site vegetation consisted of *Pinus*

contorta forest with herbaceous plants including the heat tolerant hot springs panic grass,

Dichanthelium lanuginosum (see Stout and Al□Niemi [21] for a complete geothermal plant

survey of Yellowstone National Park).

In September 2018, we collected a total of 70 soil cores (2.5 × 10 cm) along 14 transects in the

Rabbit Creek area. For each 20 m transect we sampled a total of five cores with one core every 5

m. We sampled eight transects perpendicular to Rabbit Creek (T2-T8) which were 150 m apart

along the creek. We also collected six transects perpendicular to three nearby hot springs (two

transects per hot spring, T1, T9-T13) and an additional transect (T14) away from surface water

and in the pine forest (Table S1).

We measured the soil temperature 10 cm deep next to each soil core and homogenized the soil in

each core before flash freezing and storing approximately 2 g of soil. We saved the remaining

soil for chemical analyses. All cores were processed within 24 h of collection. Soil chemistry and

moisture analyses were conducted at the Environmental Analytical Laboratory (Land Resources

and Environmental Sciences, Montana State University). See Tables S1 and S2 for complete list

of measurements.

DNA extraction and library preparation for sequencing

We extracted genomic DNA from soil with the QIAGEN DNA Power Soil Pro kit using 0.5 g of

soil instead of the recommended 0.25 g. For library preparation, we amplified the ITS1 region

using indexed primers ITS1F [22] and ITS2 [23] in the first round of PCR , followed by a second

PCR to attach the 8 bp barcodes and Illumina adapters, using Hi-Fidelity Phusion DNA

Polymerase for both reactions. Between each amplification step the product was cleaned from

reagents and primers using the NGS SPRI Bead clean-up kit (ABM). Negative PCR controls

were sequenced along with the experimental samples [24]. We included a synthetic mock

community for ITS1 as a positive control to recover tag-switching among samples [25].

Amplicons of equimolar concentrations were pooled from each of 71 samples. The libraries were

sent for paired 250-nucleotide reads on the Illumina MiSeq sequencing platform at the UC Davis

DNA Sequencing Facility (Davis, California).

Sequence processing

Raw sequence reads were demultiplexed at the UC Davis DNA Sequencing Facility (Davis,

California). We used the QIIME2 software package [26], and a modified QIIME2 pipeline for

fungal ITS by Nguyen [27] with some additional modifications. We quality filtered the

demultiplexed raw sequences (q > 30) setting the truncation of sequences 240 bp for forward and

200 bp for reverse sequences, and denoised the raw reads with DADA2 [28]. After quality

filtering, we *de novo* clustered sequences into operational taxonomic unit (OTU) with

VSEARCH at a 97% sequence similarity followed by a re-clustering step with UCLUST at 97%

to ensure the best recovery of our OTUs according to our input mock community [24]. We then

excluded all sequences that were ≤ 85% coverage in the BLAST search since this primer set

amplifies sequences that only partially match to a small part of the ITS gene. We used the

UNITE database v8, [29, 30] for chimera checking, OTU clustering, and assigning taxonomy.

And finally, we rarefied each sample to a sampling depth of 2831 sequences.

Initial ordination analyses of the OTU community matrix led us to exclude 30 samples as they

clustered due to partial, very short sequences spanning only ~20 nucleotides at the beginning of

the gene making them unidentifiable. In short, the non-metric dimensional scaling (NMDS)

ordination including all remaining samples clustered in three groups (Figure S2) separated by pH

but with high environmental variable overlap (Figure S3). A species indicator analysis showed

the group with pH>8 clustered due to shared unidentified sequences (Table S3) and in fact, the

ITS1 sequences for the OTUs in this group could not be identified by performing additional

BLAST searches. We therefore excluded cores S5, S6, S7, S8, S9, S60, S63, S64, S67, and S69

from subsequent analyses. These cores were sampled near the three pools and the forest transect

away from surface water (Table S1 and S2). After filtering, we used 40 cores to perform all

further analyses. All scripts and data tables are available online at

https://github.com/abazzical/YellowstoneFungi.

Soil Chemistry and Fungal Community Analyses

We performed Principal Coordinates Analysis (PCoA) to assess chemistry and community

similarities across soil cores using the vegan package [31, 32] and Non-Metric Dimensional

Scaling (NMDS) based on Bray-Curtis distances implemented in R [33]. Because our data had

no multivariate normality and unequal variances, we tested if the clusters identified were

significantly different with the *adonis* function in the *vegan* package in R. We performed an

indicator species analysis to determine if specific OTUs reflected soil abiotic parameters using

the *indicspecies* package (ver. 1.7.8) [34] in R with 9999 permutations. We produced a species

accumulation curve using the *specaccum* function estimating the mean curve and its standard

deviation from 100 random permutations of the species accumulation data.

We categorized fungi into different trophic guilds using FUNGuild [35]. We further used the

ectomycorrhizal (EM) OTUs and saprotrophic OTUs for additional NMDS and MANOVA

analyses to test if specific guilds drive the patterns found in the total community. We excluded

core S42 from EM fungi NMDS as it contained only one taxon, the orchid associated

Ceratobasidium that was unique to that core.

To test for significant differences in the number of OTUs per core between the clusters identified

in the NMDS analysis of the communities, we performed three Welch's T-tests (with a

Bonferroni correction p=0.01667) for the whole data set, EM OTUs and for saprotrophic OTUs.

Fungal culture isolation and growth assays

We obtained pure cultures for four species growing at the Rabbit Creek site. We cultured the

saprotophic *Agaricus campestris* and ectomycorrhizal *Pisolithus tinctorius* from fruitbodies on

Potato Dextrose Agar (PDA) and Modified Melin-Norkans (MMN, [36]) media respectively.

We also isolated cultures of potential plant pathogens *Fusarium oxysporum* and *F. avenae* from

soil by plating serial soil dilutions in PDA. For all four species, we measured mycelial radial

expansion at 30, 35, and 40°C and pH 4, 7, and 9 separately. Each culture was grown in triplicate

and measured at 2-4 day intervals. *Fusarium* and *Agaricus* cultures were grown on PDA and

Pisolithus was grown on MMN media. The growth in pH was calculated as the average of four

radius measurements per colony for each time point. The temperature measurements were made

by calculating the area of the fungal colony on the agar plate by importing the drawn outline of

the fungus into ImageJ [37, 38].

We obtained culture identifications by sequencing the ITS rRNA gene. In short, we extracted

genomic DNA using the Extract-N-Amp Sigma kit and amplified the ITS region with primers

ITS1F [22] and ITS4 [23]. The PCR product was Sanger sequenced at GenScript and the

sequences were checked and trimmed for low quality bases in the software FinchTV

(http://www.geospiza.com/finchtv/). To identify the fungi, we used the BLAST tool in the NCBI

database. Sequences are deposited in GenBank (MW471687 - MW472279).

Results

Soil abiotic variation

Soils in the Rabbit Creek area showed wide variation in temperature, moisture content, and

chemistry (Fig. 1). Despite the creek's very high temperature water sources (84ºC), soil

temperature was relatively low and varied between 10 ºC and 31ºC. Soil pH and moisture content

varied broadly, with pH ranging from 4 to 10 and moisture from 10% to 100%. Temperature, pH,

and moisture were correlated as soil cores collected closest to the creek were wetter, hotter, and

more alkaline. The remaining soil chemical parameters also varied considerably (Table S1 and

S2). Manganese (Mn), cobalt (Co), and cadmium (Cd) were highly correlated and inversely

proportional to pH, temperature, and moisture. Total Nitrogen (N), total Carbon (C), and zinc

(Zn) were orthogonal to these other parameters.

Figure 1 – Rabbit Creek soil chemistry variation. Principal components analysis on abiotic
parameters measured across soil samples. Colors reflect the pH level measured for each soil core.

Overall soil fungal community composition

We found a total of 593 fungal OTUs at Rabbit Creek and were able to obtain taxonomic

identifications other than "fungi" for close to 65% of them. This thermal site was heavily

dominated by members of Ascomycota (38.4%) and Basidiomycota (18.7%) but also included

Chytridiomycota, Rozellomycota, Mucoromycota, Glomeromycota, Kickxellomycota,

Mortierellomycota, and Calcarisporiellomycota. We detected a few fungi known for thriving at

high temperatures (>30 ℃), but not resilient enough to be considered heat tolerant fungi (with

the ability to grow at 40 ˚C [39]). These included *Exophiala opportunistica*, a black yeast that

also withstands high moisture and alkalinity and is commonly found in dishwashers [40, 41],

Waitea circinata, a grass pathogen with optimal growth at 25-30 ℃ [42], *Umbelopsis vinacea*, a

soil fungus that remains active up to 37 ℃ [43], *Gaertneriomyces semiglobifer*, a chytrid that

also growth up to 37 ℃ [44], and *Fusarium kerasplasticarum*, an opportunistic animal pathogen

that can also grow up to 37 ℃. In addition, we found one unidentified species of *Talaromyces*, a

genus that includes thermophilic and thermotolerant species [45]. Notably, we recovered a

number of cold-adapted fungi in our survey, including *Extremus aquaticus*, *Elasticomyces*

elasticus, *Endophoma alongata*, *Alternaria chlamydosporigena*, *Mrakiella aquatica*, *Mrakia*

frigida, *Tausonia pamirica*, *T. pulullans*, *Naganishia adeliensis*, and *Udeniomyces*

megalosporus. Many of these species occur in Polar regions and have documented optimal

growth temperatures <20 ℃ [46-51].

Our species accumulation curve does not plateau (Figure S4), suggesting that community

diversity is not saturated with our sampling.

Community clustering and indicator species analysis

We found fungal community differences across soil cores were not significantly different.

Although the soil pH values mapped to the community NMDS analysis showed a pattern of

samples clustering by pH, the two clusters of pH > and < 8 were not statistically significant

(Figure 2A, S3A). A set of 42 OTUs was present only at pH >8 (Figure 2B).

Figure 2 – Fungal community composition across soil cores collected in Rabbit Creek. (A) Non-Metric Dimensional Scaling across cores with respective pH measurements >8 (yellow to light green) and <8 (green – dark blue) (stress = 0.1577); (B) Venn diagram depicting fungal OTU overlap between the two clusters matching cores with pH measurements >8 (yellow) and <8 (dark blue).

The indicator species analysis on the two NMDS clusters detected 33 species associated with pH

>8 and seven species associated with pH <8 (Table 1 and Table S3). 51% of taxa from the group

pH >8 and all of the taxa from the group pH <8 belonged to the phylum Ascomycota. Three

OTUs not identified to species were part of Glomeromycotina and six in Basidiomycota (Table

S3).

Table 1 Indicator species identified to species for clusters of cores with pH >8 and <8. The
complete list including unidentified taxa can be found in Table S3.

Cluster	Phylum	Class	Order	Species
pH >8	Ascomycota	Dothideomycetes	Pleosporales	*Septoriella phragmitis*
	Ascomycota	Sordariomycetes	Hypocreales	*Acremonium exuviarum*
	Ascomycota	Sordariomycetes	Hypocreales	*Acremonium tubakii*
	Ascomycota	Sordariomycetes	Hypocreales	*Fusarium keratoplasticum*
	Ascomycota	Sordariomycetes	Hypocreales	*Metarhizium pemphigi*

Ascomycota	Sordariomycetes	Hypocreales	*Purpureocillium lavendulum*
Ascomycota	Sordariomycetes	Hypocreales	*Sarocladium bactrocephalum*
Ascomycota	Sordariomycetes	Sordariales	*Schizothecium miniglutinans*
Basidiomycota	Agaricomycetes	Agaricales	*Agaricus gastronevadensis*
Basidiomycota	Agaricomycetes	Agaricales	*Coprinus cordisporus*
Basidiomycota	Agaricomycetes	Agaricales	*Simocybe centunculus*
Basidiomycota	Agaricostilbomycetes	Agaricostilbales	*Kondoa miscanthi*
Basidiomycota	Tremellomycetes	Cystofilobasidiales	*Udeniomyces megalosporus*
Basidiomycota	Ustilaginomycetes	Ustilaginales	*Ustilago nunavutica*
pH <8 Ascomycota	Dothideomycetes	Capnodiales	*Elasticomyces elasticus*
Ascomycota	Eurotiomycetes	Eurotiales	*Penicillium nodositatum*
Ascomycota	Leotiomycetes	Helotiales	*Infundichalara minuta*
Ascomycota	Leotiomycetes	Helotiales	*Phialocephala fortinii*

Indicator species for cores with pH >8 included *Acremonium exuviarum* and *Acremonium tubakii*

and closely related *Sarocladium bactrocephalum* [52], which belong to the Emericellopsis clade

characterized by alkalitolerant fungi [53]. We also identified the plant-associated and

psychrophilic fungus *Udeniomyces megalosporus* as an indicatory species for cores with pH >8

[54]. For cores with pH <8, we found indicator species *Elasticomyces elasticus* reported as being

able to grow at 0 as well as at 25 °C [55], and *Infundichalara minuta* also recorded from Pb-Zn

mining sites with pH 7.3, although no other information is available for pH tolerance of these

species [56].

Saprotrophic taxa follow overall fungal community clustering and pH gradient

Although not significant, we found that saprotrophic fungi but not ectomycorrhizal (EM) fungi

closely followed the overall community pattern regarding pH (Figure 2). EM fungi (Figure 3A &

B) showed much less separation between cores of higher and lower pH compared to saprotrophic

taxa (Figure 3C & D). We found an almost complete EM OTU overlap between cores with high

and low pH and only a single OTU unique to pH >8 cores while six out of 26 saprotrophic fungi

were unique to the pH >8 cores (Figure 3). A total of 234 (out of 593) OTUs were classified into

trophic guilds, of which 22.2% (52) OTUs were mycorrhizal (including 16.7% (39)

ectomycorrhizal (EM) fungi), 34.6% (81) saprotrophic, 9.8% (23) plant pathogens, 15% (35)

animal pathogens, and 6.8% (16) dung-associated fungi (Table S4).

269
Figure 3 – EM and saprotrophic fungal community composition and OTU overlap. (A) EM
fungi Non-Metric Dimensional Scaling across soil samples (stress = 0.205) with color scale of
pH >8 (yellow to light green) and pH <8 (green – dark blue). (B) Venn diagram depicting EM
fungal OTU overlap between soil cores with pH measurements >8 (yellow) and <8 (dark blue).
(C) Saprotrophic fungi Non-Metric Dimensional Scaling across soil samples (stress = 0.196)
with color scale of pH >8 (yellow to light green) and pH <8 (green – dark blue). (D) Venn
diagram depicting saprotrophic fungal OTU overlap between soil cores with pH measurements
>8 (yellow) and <8 (dark blue).

Saprotrophic OTU numbers decrease in more alkaline soils but EM OTU numbers do not

We found the environmental change from pH <8 to pH >8 reduced the number of total soil fungi and saprotrophic fungi, but not EM fungi. Soils with pH > 8 to host lower OTU richness (31 OTUs) compared to soils with pH <8 (40 OTUs, Figure 4, Welch's T-test: t = -3.0974, df = 30.356, p-value = 0.00418) . Saprotrophic OTUs followed a similar pattern (Welch's T-test: t = -3.1753, df = 17.662, p-value = 0.005335), with a mean of 6 OTUs in soils with pH >8 and a mean of 10 OTUs in soils with pH <8. EM fungi did not show a significant difference in numbers of OTUs across soils with pH > and < 8 (Welch's T-test: t = 1.5064, df = 12.7, p-value = 0.1564).

Figure 4 – Number of fungal OTUs detected in soil samples with pH <8 (dark blue) and >8 (yellow). (A) Number fungal OTUs; (B) Number of EM fungal OTUs; (C) Number of

293 saprotrophic fungal OTUs. Note the scale bar differences. Asterisks (*) indicate significance
294 between two groups.
295

Effect of temperature and pH on four fungi from Yellowstone National Park soils

The four fungal species tested for temperature and pH tolerance lacked thermal tolerance but had

the ability to tolerate a wide range of pH (Figures 5 and 6). Notably, all four species performed

well at pH 7. Both *Fusarium oxysporum* and *F. avenae* grew equally well at pH 9 and pH 7

showing less growth at pH 4. Both *Agaricus campestris* and *Pisolithus tinctorius* grew best at pH

7, however, *P. tinctorius* was the only fungus with no growth at pH 9 (Figure 5). All four species

grew well at 30 ºC but very poorly at 40 ºC (Figure 6). *P. tinctorius* and *F. oxysporum* grew

equally well at 30 ˚C and 35 ˚C, while *A. campestris* and *F. avenae* grew better at 30 ˚C than 35

˚C.

Figure 5 – Mycelial growth (average of 4 radius measurement per fungal thallus) at pH 4 (dark blue), pH 7 (green), and pH 9 (yellow) in *Agaricus campestris* (A), *Pisolithus tinctorius* (B), *Fusarium oxysporum* (C), and *Fusarium avenae* (D).

307
308
309
310

Figure 6 – Mycelial growth (fungus growth area mm^3) at 30ºC (peach), 35ºC (pink), and 40ºC (burgundy) in *Agaricus campestris* (A), *Pisolithus tinctorius* (B), *Fusarium oxysporum* (C), and *Fusarium avenae* (D).

Discussion

We investigated soil fungal communities from a geothermal area in Yellowstone National Park

and found that although no single soil variable explained or structured fungal community

composition, higher soil pH hosted lower fungal OUT richness. Saprotrophic fungi followed

more closely the observed community trends rather than ectomycorrhizal fungi. Notably, we did

not detect any thermophilic fungi in Rabbit Creek. In contrast, we detected several cold-adapted

species previously recorded in the region. Specifically, we found the psychrophilic

basidiomycete yeast *Guehomyces pulullans* [57] and a few cold adapted Ascomycete yeasts

identified in Vu, Groenewald, Szöke, Cardinali, Eberhardt, Stielow, de Vries, Verkleij, Crous,

Boekhout and Robert [58], including *Naganishia adeliensis* and *Udenomyces megalosporus* (=*U.*

pyricola in Vu et al 2016). We also found the psychrophiles *Elasticomyces elasticus* and *Mrakia*

sp., which had been detected in previous soil surveys in both West Yellowstone and Hyalite near

Cooke City, Montana [59]. Additionally, our *Tausonia pamirica* OTU matched 100% to the

sequence of the type specimen (Genbank # NR_154490), a known psychrotolerant yeast from

Antarctica [60]. The occurrence of these species is most likely unrelated to the thermal nature of

Rabbit Creek and rather may reflect the regional climate characterized by severe cold winters.

We were surprised by the absence of thermophilic fungi and the presence of cold-adapted

lineages in a thermal area and that none of the fungi we isolated in culture were able to grow at

temperatures higher than 35°C. Soil temperature is thought to be the most significant factor

influencing bacterial and archaeal species composition in geothermally heated environments [21,

61, 62]. Thermophilic fungi are mostly known from studies on plant-associated fungi from

geothermal environments [63] and surveys such as the YNP metagenome project [64]. There is

also experimental evidence documenting previously unknown thermophilic and acidophilic

fungal species from hot spring habitats (Yamazaki, Toyama and Nakagiri [65]. However, the soil

temperature of our sampled cores was relatively low, ranging from 10 ºC to 31ºC, which is well

within the range of known temperature tolerance of most fungi including the detected

psychrophilic species [57, 58] and the fungi tested in this study (Fig. 6). It is important to note

that our findings do not imply the absence of heat-adapted fungi in Yellowstone, as other thermal

areas characterized by warmer soils may well host thermophilic fungi of unknown taxonomic

identity.

Conditions in highly acidic or alkaline environments often co-vary with other environmentally

extreme variables such as elevated salt concentrations or temperatures (Orwa, Mugambi,

Wekesa, & Mwirichia, 2020), and in our case, soil moisture content. It is known that edaphic

parameters structure the communities of soil fungi and variation in chemistry (including pH)

affects fungal diversity on global, continental and local scales [1, 2]. Individual fungi appear to

have a wider pH tolerance range compared to bacteria [66], however there are several examples

of fungi (including *P. tinctorus* which was studied here) being able to withstand both low and

high pH levels [14]. Studies on acid- or alkaline-loving fungi have recorded new species and

have expanded our knowledge of the possible habitat types for commonly found fungal species

[13]. In addition, research on fungi associated with hot spring sediment across the globe has

documented the phylum Ascomycota as dominant in these habitats. These included newly

discovered species [65] and common members of widespread genera, such as *Aspergillus*,

Penicillium, and *Cladosporium*. Mycorrhizal species belonging to Ascomycota have also been

detected soda lake (pH range) habitats [67]. We found the Rabbit Creek soils with higher pH are

inhabited by many members of the Ascomycota lineage, which includes several alkalitolerant

fungi such as *Acremonium* from the Emericellopsis clade [53].

In conclusion, we found thermophilic fungi to be absent from the soils surrounding thermal

water in Rabbit Creek. We also found no statistical support suggesting soil variables determined

the structure of fungal community composition, but both overall and saprotrophic fungal richness

decreased with higher pH values. Further identifying the abiotic variables driving the fungal

community assemblage of these soils will require more statistical power to untangle the weight

of several co-varying factors. Further research on thermal areas characterized by hot soils is

however needed, as it will shed light on the effects of high soil temperature on the ecology and

evolution of fungi.

Acknowledgements

We thank Maddie Trent, Rio Wofford, Colin Kennedy, Seamus Hoolahan, and Kathryn Gannon

for assistance in the field and laboratory, and the Quandt lab for suggestions on the manuscript.

Cathy Zabinski helped obtaining the collecting permit, Dan Colman assisted with sampling

design, and Kabir Peay provided raw data from another study that allowed fungal community

comparisons. We also thank Yellowstone National Park for allowing the collection of samples in

Rabbit Creek (YELL-2018-SCI8062).

Cited Literature

1. Tedersoo L, Bahram M, Põlme S, Kõljalg U, Yorou NS, Wijesundera R, Ruiz LV,
Vasco-Palacios AM, Thu PQ, Suija A (2014) Global diversity and geography of soil
fungi. Science 346: 1256688.
2. Peay KG, Kennedy PG, Talbot JM (2016) Dimensions of biodiversity in the Earth
mycobiome. Nature Reviews Microbiology 14: 434-447. doi: 10.1038/nrmicro.2016.59
3. Amenábar M, Urschel M, Boyd E (2015) Metabolic and taxonomic diversification in
continental magmatic hydrothermal systems
4. Brock TD, Freeze H (1969) *Thermus aquaticus* gen. n. and sp. n., a Nonsporulating
Extreme Thermophile. J Bacteriol 98: 289.
5. Meyer-Dombard DR, Shock EL, Amend JP (2005) Archaeal and bacterial communities
in geochemically diverse hot springs of Yellowstone National Park, USA. Geobiology 3:
211-227. doi: https://doi.org/10.1111/j.1472-4669.2005.00052.x
6. Colman DR, Jay ZJ, Inskeep WP, Jennings Rd, Maas KR, Rusch DB, Takacs-Vesbach
CD (2016) Novel, Deep-Branching Heterotrophic Bacterial Populations Recovered from
Thermal Spring Metagenomes7. doi: 10.3389/fmicb.2016.00304

7. Colman DR, Lindsay MR, Boyd ES (2019) Mixing of meteoric and geothermal fluids
supports hyperdiverse chemosynthetic hydrothermal communities. Nature
Communications 10: 681. doi: 10.1038/s41467-019-08499-1
8. Brown SR, Fritz SC (2019) Eukaryotic organisms of continental hydrothermal systems.
Extremophiles 23: 367-376. doi: 10.1007/s00792-019-01101-y
9. Tansey MR, Brock TD (1972) The Upper Temperature Limit for Eukaryotic Organisms.
Proceedings of the National Academy of Sciences 69: 2426. doi: 10.1073/pnas.69.9.2426
10. Morgenstern I, Powlowski J, Ishmael N, Darmond C, Marqueteau S, Moisan M-C,
Quenneville G, Tsang A (2012) A molecular phylogeny of thermophilic fungi. Fungal
Biology 116: 489-502. doi: https://doi.org/10.1016/j.funbio.2012.01.010
11. Hutchinson MI, Powell AJ, Herrera J, Natvig DO (2019) New Perspectives on the
Distribution and Roles of Thermophilic Fungi. In: Tiquia-Arashiro, SM, Grube, M (eds.)
Fungi in Extreme Environments: Ecological Role and Biotechnological Significance.
Springer International Publishing, Cham, pp. 59-80
12. Powell AJ, Parchert KJ, Bustamante JM, Ricken JB, Hutchinson MI, Natvig DO (2012)
Thermophilic fungi in an aridland ecosystem. Mycologia 104: 813-825. doi: 10.3852/11-
298
13. Aguilera A, González-Toril E (2019) Eukaryotic Life in Extreme Environments:
Acidophilic Fungi. In: Tiquia-Arashiro, SM, Grube, M (eds.) Fungi in Extreme
Environments: Ecological Role and Biotechnological Significance. Springer International
Publishing, Cham, pp. 21-38
14. Hung L-L, Trappe JM (1983) Growth Variation Between and Within Species of
Ectomycorrhizal Fungi in Response to pH in Vitro. Mycologia 75: 234-241. doi:
10.1080/00275514.1983.12021660
15. Wheeler KA, Hurdman BF, Pitt JI (1991) Influence of pH on the growth of some
toxigenic species of *Aspergillus*, *Penicillium* and *Fusarium*. Int J Food Microbiol 12:
141-149. doi: https://doi.org/10.1016/0168-1605(91)90063-U
16. Nevarez L, Vasseur V, Le Madec A, Le Bras MA, Coroller L, Leguérinel I, Barbier G
(2009) Physiological traits of *Penicillium glabrum* strain LCP 08.5568, a filamentous
fungus isolated from bottled aromatised mineral water. Int J Food Microbiol 130: 166-
171. doi: https://doi.org/10.1016/j.ijfoodmicro.2009.01.013
17. Redman RS, Litvintseva A, Sheehan KB, Henson JM, Rodriguez RJ (1999) Fungi from
Geothermal Soils in Yellowstone National Park. Appl Environ Microbiol 65: 5193. doi:
10.1128/AEM.65.12.5193-5197.1999
18. Appoloni S, Lekberg Y, Tercek MT, Zabinski CA, Redecker D (2008) Molecular
Community Analysis of Arbuscular Mycorrhizal Fungi in Roots of Geothermal Soils in
Yellowstone National Park (USA). Microb Ecol 56: 649-659.
19. Lekberg Y, Meadow J, Rohr JR, Redecker D, Zabinski CA (2011) Importance of
dispersal and thermal environment for mycorrhizal communities: lessons from
Yellowstone National Park. Ecology 92: 1292-1302. doi: https://doi.org/10.1890/10-
1516.1
20. Meadow JF, Zabinski CA (2012) Spatial heterogeneity of eukaryotic microbial
communities in an unstudied geothermal diatomaceous biological soil crust: Yellowstone
National Park, WY, USA. FEMS Microbiol Ecol 82: 182-191. doi: 10.1111/j.1574-
6941.2012.01416.x %J FEMS Microbiology Ecology

21.   Stout RG, Al☐Niemi TS (2002) Heat☐tolerant Flowering Plants of Active Geothermal
445   Areas in Yellowstone National Park. Ann Bot 90: 259-267. doi: 10.1093/aob/mcf174 %J
446   Annals of Botany
447   22.   Gardes M, Bruns TD (1993) ITS primers with enhanced specificity for
448   basidiomycetes☐application to the identification of mycorrhizae and rusts. Mol Ecol 2:
449   113-118. doi: 10.1111/j.1365-294X.1993.tb00005.x
450   23.   White TJ, Bruns T, Lee S, Taylor J (1990) Amplification and direct sequencing of fungal
451   ribosomal RNA genes for phylogenetics. In: Innis, M, Gelfand, D, Sninsky, J, White, TJ
452   (eds.) PCR protocols: a guide to methods and applications. Academic Press Inc., New
453   York, pp. 315-322
454   24.   Nguyen NH, Smith D, Peay K, Kennedy P (2015) Parsing ecological signal from noise in
455   next generation amplicon sequencing. New Phytol 205: 1389-1393. doi:
456   https://doi.org/10.1111/nph.12923
457   25.   Palmer JM, Jusino MA, Banik MT, Lindner DL (2018) Non-biological synthetic spike-in
458   controls and the AMPtk software pipeline improve mycobiome data. PeerJ 6: e4925. doi:
459   10.7717/peerj.4925
460   26.   Bolyen E, Rideout JR, Dillon MR, Bokulich NA, Abnet CC, Al-Ghalith GA, Alexander
461   H, Alm EJ, Arumugam M, Asnicar F, Bai Y, Bisanz JE, Bittinger K, Brejnrod A,
462   Brislawn CJ, Brown CT, Callahan BJ, Caraballo-Rodríguez AM, Chase J, Cope EK, Da
463   Silva R, Diener C, Dorrestein PC, Douglas GM, Durall DM, Duvallet C, Edwardson CF,
464   Ernst M, Estaki M, Fouquier J, Gauglitz JM, Gibbons SM, Gibson DL, Gonzalez A,
465   Gorlick K, Guo J, Hillmann B, Holmes S, Holste H, Huttenhower C, Huttley GA, Janssen
466   S, Jarmusch AK, Jiang L, Kaehler BD, Kang KB, Keefe CR, Keim P, Kelley ST, Knights
467   D, Koester I, Kosciolek T, Kreps J, Langille MGI, Lee J, Ley R, Liu Y-X, Loftfield E,
468   Lozupone C, Maher M, Marotz C, Martin BD, McDonald D, McIver LJ, Melnik AV,
469   Metcalf JL, Morgan SC, Morton JT, Naimey AT, Navas-Molina JA, Nothias LF,
470   Orchanian SB, Pearson T, Peoples SL, Petras D, Preuss ML, Pruesse E, Rasmussen LB,
471   Rivers A, Robeson MS, Rosenthal P, Segata N, Shaffer M, Shiffer A, Sinha R, Song SJ,
472   Spear JR, Swafford AD, Thompson LR, Torres PJ, Trinh P, Tripathi A, Turnbaugh PJ,
473   Ul-Hasan S, van der Hooft JJJ, Vargas F, Vázquez-Baeza Y, Vogtmann E, von Hippel M,
474   Walters W, Wan Y, Wang M, Warren J, Weber KC, Williamson CHD, Willis AD, Xu
475   ZZ, Zaneveld JR, Zhang Y, Zhu Q, Knight R, Caporaso JG (2019) Reproducible,
476   interactive, scalable and extensible microbiome data science using QIIME 2. Nat
477   Biotechnol 37: 852-857. doi: 10.1038/s41587-019-0209-9
478   27.   Nguyen N (2020) Amplicon analysis pipeline with QIIME2:
479   https://github.com/nnguyenlab/amplicon-pipeline.
480   28.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP (2016)
481   DADA2: High-resolution sample inference from Illumina amplicon data. Nat Methods
482   13: 581-583. doi: 10.1038/nmeth.3869
483   29.   Abarenkov K, Nilsson RH, Larsson KH, Alexander IJ, Eberhardt U, Erland S, Høiland K,
484   Kjøller R, Larsson E, Pennanen T (2010) The UNITE database for molecular
485   identification of fungi–recent updates and future perspectives. New Phytol 186: 281-285.
486   30.   Kõljalg U (2020) UNITE general FASTA release for Fungi. In: community, U (ed.),
487   10.15156/BIO/786368.
488   31.   Dixon P (2003) VEGAN, a package of R functions for community ecology. Journal of
489   Vegetation Science 14: 927-930. doi: https://doi.org/10.1111/j.1654-1103.2003.tb02228.x

32. Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin P, O'hara R, Simpson G,
Solymos P, Stevens M, Wagner H (2017) Vegan: community ecology package. R
package. Version.

33. R Core Team (2014) R: A language and environment for statistical computing. R
Foundation for Statistical Computing, Vienna, Austria, 2012.

34. De Caceres M, Jansen F, De Caceres MJRp, Version (2016) Indicspecies: relationship
between species and groups of sites1: 2014.

35. Nguyen NH, Song Z, Bates ST, Branco S, Tedersoo L, Menke J, Schilling JS, Kennedy
PG (2016) FUNGuild: An open annotation tool for parsing fungal community datasets by
ecological guild. Fungal Ecology 20: 241-248. doi:
https://doi.org/10.1016/j.funeco.2015.06.006

36. Marx DH (1970) The influence of ectotrophic mycorrhizal fungi on the resistance of Pine
roots to pathogenic infections. V. Resistance of mycorrhizae to infection by vegetative
mycelium of Phytophthora cinnamomi. Phytopathology 60: 1472-1473. doi:
10.1094/Phyto-60-1472

37. Abràmoff MD, Magalhães PJ, Ram SJ (2004) Image processing with ImageJ.
Biophotonics international 11: 36-42.

38. Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of
image analysis. Nat Methods 9: 671-675.

39. Mouchacca J (2007) Heat tolerant fungi and applied research: Addition to the previously
treated group of strictly thermotolerant species. World Journal of Microbiology and
Biotechnology 23: 1755. doi: 10.1007/s11274-007-9426-3

40. Zalar P, Novak M, de Hoog GS, Gunde-Cimerman N (2011) Dishwashers – A man-made
ecological niche accommodating human opportunistic fungal pathogens. Fungal Biology
115: 997-1007. doi: https://doi.org/10.1016/j.funbio.2011.04.007

41. Gümral R, Özhak-baysan B, Tümgör A, Saraçli MA, Yildiran ST, Ilkit M, Zupancic J,
Novak-babic M, Gunde-cimerman N, Zalar P, de Hoog GS (2016) Dishwashers provide a
selective extreme environment for human-opportunistic yeast-like fungi. Fungal
Diversity 76: 1-9. doi: http://dx.doi.org/10.1007/s13225-015-0327-8

42. de la Cerda KA, Douhan GW, Wong FP (2007) Discovery and Characterization of
Waitea circinata var. *circinata* Affecting Annual Bluegrass from the Western United
States. Plant Dis 91: 791-797. doi: 10.1094/PDIS-91-7-0791

43. Wang Y-N, Liu X-Y, Zheng R-Y (2014) Umbelopsis changbaiensis sp. nov. from China
and the typification of *Mortierella vinacea*. Mycological Progress 13: 657-669. doi:
10.1007/s11557-013-0948-9

44. Henderson L, Ly M-A, Robinson K, Gleason FH, Lilje O (2018) Maximum temperature
for growth and reproduction is similar in two soil isolates of *Gaertneriomyces*
semiglobifer (Spizellomycetales, Chytridiomycetes) from different soil environments in
north eastern New South Wales, Australia. Nova Hedwigia 106: 485-497. doi:
10.1127/nova_hedwigia/2017/0448

45. Houbraken J, Spierenburg H, Frisvad JC (2012) *Rasamsonia*, a new genus comprising
thermotolerant and thermophilic *Talaromyces* and *Geosmithia* species. Antonie Van
Leeuwenhoek 101: 403-421. doi: 10.1007/s10482-011-9647-1

46. Selbmann L, Zucconi L, Isola D, Onofri S (2015) Rock black fungi: excellence in the
extremes, from the Antarctic to space. Curr Genet 61: 335-345. doi: 10.1007/s00294-014-
0457-7

47. Hirose D, Hobara S, Matsuoka S, Kato K, Tanabe Y, Uchida M, Kudoh S, Osono T
(2016) Diversity and community assembly of moss-associated fungi in ice-free coastal
outcrops of continental Antarctica. Fungal Ecology 24: 94-101. doi:
https://doi.org/10.1016/j.funeco.2016.09.005
48. Řezanka T, Kolouchová I, Sigler K (2016) Lipidomic analysis of psychrophilic yeasts
cultivated at different temperatures. Biochim Biophys Acta 1861: 1634-1642. doi:
https://doi.org/10.1016/j.bbalip.2016.07.005
49. Thomas-Hall SR, Turchetti B, Buzzini P, Branda E, Boekhout T, Theelen B, Watson K
(2010) Cold-adapted yeasts from Antarctica and the Italian Alps—description of three
novel species: *Mrakia robertii* sp. nov., *Mrakia blollopis* sp. nov. and *Mrakiella*
niccombsii sp. nov. Extremophiles 14: 47-59. doi: 10.1007/s00792-009-0286-7
50. Vishniac HS (1999) Psychrophilic Yeasts. In: Seckbach, J (ed.) Enigmatic
Microorganisms and Life in Extreme Environments. Springer Netherlands, Dordrecht,
pp. 315-321
51. Białkowska AM, Szulczewska KM, Krysiak J, Florczak T, Gromek E, Kassassir H, Kur
J, Turkiewicz M (2017) Genetic and biochemical characterization of yeasts isolated from
Antarctic soil samples. Polar Biol 40: 1787-1803. doi: 10.1007/s00300-017-2102-7
52. Giraldo A, Gené J, Sutton DA, Madrid H, de Hoog GS, Cano J, Decock C, Crous PW,
Guarro J (2015) Phylogeny of Sarocladium (Hypocreales). Persoonia 34: 10-24. doi:
10.3767/003158515X685364
53. Grum-Grzhimaylo AA, Georgieva ML, Debets AJM, Bilanenko EN (2013) Are
alkalitolerant fungi of the Emericellopsis lineage (Bionectriaceae) of marine origin? IMA
Fungus 4: 213-228. doi: 10.5598/imafungus.2013.04.02.07
54. Han P-J, Qiu J-Z, Wang Q-M, Bai F-Y (2012) *Udeniomyces kanasensis* sp. nov., a
ballistoconidium-forming yeast species in the Cystofilobasidiales. Antonie Van
Leeuwenhoek 102: 45-51. doi: 10.1007/s10482-012-9711-5
55. Selbmann L, de Hoog GS, Zucconi L, Isola D, Ruisi S, van den Ende AHGG, Ruibal C,
De Leo F, Urzì C, Onofri S (2008) Drought meets acid: three new genera in a
dothidealean clade of extremotolerant fungi. Studies in Mycology 61: 1-20. doi:
https://doi.org/10.3114/sim.2008.61.01
56. Böhmer M, Ozdín D, Račko M, Lichvár M, Budiš J, Szemes T (2020) Identification of
Bacterial and Fungal Communities in the Roots of Orchids and Surrounding Soil in
Heavy Metal Contaminated Area of Mining Heaps. Applied Sciences 10: 7367.
57. Cavello I, Albanesi A, Fratebianchi D, Garmedia G, Vero S, Cavalitto S (2017)
Pectinolytic yeasts from cold environments: novel findings of *Guehomyces pullulans*,
Cystofilobasidium infirmominiatum and *Cryptococcus adeliensis* producing pectinases.
Extremophiles 21: 319-329. doi: http://dx.doi.org/10.1007/s00792-016-0904-0
58. Vu D, Groenewald M, Szöke S, Cardinali G, Eberhardt U, Stielow B, de Vries M,
Verkleij GJM, Crous PW, Boekhout T, Robert V (2016) DNA barcoding analysis of
more than 9 000 yeast isolates contributes to quantitative thresholds for yeast species
and genera delimitation. Studies in Mycology 85: 91-105. doi:
https://doi.org/10.1016/j.simyco.2016.11.007
59. Steidinger BS, Bhatnagar JM, Vilgalys R, Taylor JW, Qin C, Zhu K, Bruns TD, Peay KG
(2020) Ectomycorrhizal fungal diversity predicted to substantially decline due to climate
changes in North American Pinaceae forests. J Biogeogr 47: 772-782. doi:
https://doi.org/10.1111/jbi.13802

60. Tsuji M, Kudoh S (2020) Soil Yeasts in the Vicinity of Syowa Station, East Antarctica:
Their Diversity and Extracellular Enzymes, Cold Adaptation Strategies, and Secondary
Metabolites. Sustainability 12. doi: 10.3390/su12114518
61. Sharp CE, Brady AL, Sharp GH, Grasby SE, Stott MB, Dunfield PF (2014) Humboldt's
spa: microbial diversity is controlled by temperature in geothermal environments. The
ISME Journal 8: 1166-1174. doi: 10.1038/ismej.2013.237
62. de Oliveira TB, Rodrigues A (2019) Ecology of Thermophilic Fungi. In: Tiquia-Arashiro,
SM, Grube, M (eds.) Fungi in Extreme Environments: Ecological Role and
Biotechnological Significance. Springer International Publishing, Cham, pp. 39-57
63. Zhou W-N, White JF, Soares MA, Torres MS, Zhou Z-P, Li H-Y (2015) Diversity of
fungi associated with plants growing in geothermal ecosystems and evaluation of their
capacities to enhance thermotolerance of host plants. Journal of Plant Interactions 10:
305-314. doi: 10.1080/17429145.2015.1101495
64. Inskeep W, Jay Z, Tringe S, Herrgard M, Rusch D (2013) The YNP Metagenome Project:
Environmental Parameters Responsible for Microbial Distribution in the Yellowstone
Geothermal Ecosystem4. doi: 10.3389/fmicb.2013.00067
65. Yamazaki A, Toyama K, Nakagiri A (2010) A new acidophilic fungus *Teratosphaeria
acidotherma* (Capnodiales, Ascomycota) from a hot spring. Mycoscience 51: 443-455.
doi: 10.1007/s10267-010-0059-2
66. Rousk J, Bååth E, Brookes PC, Lauber CL, Lozupone C, Caporaso JG, Knight R, Fierer
N (2010) Soil bacterial and fungal communities across a pH gradient in an arable soil.
The ISME Journal 4: 1340-1351. doi: 10.1038/ismej.2010.58
67. Ishida TA, Nara K, Ma S, Takano T, Liu S (2009) Ectomycorrhizal fungal community in
alkaline-saline soil in northeastern China. Mycorrhiza 19: 329-335. doi:
http://dx.doi.org/10.1007/s00572-008-0219-9
607