1 **Unique protein features of SARS-CoV-2 relative to other *Sarbecoviruses***
2
3

4 Matthew Cotten[1,2], David L. Robertson[2], My V.T. Phan[1]
5

Affiliations

1. MRC/UVRI & LSHTM Uganda Research Unit, Entebbe, Uganda

2. MRC-University of Glasgow Centre for Virus Research, Glasgow, UK

Correspondence: Matthew Cotten (Matthew.Cotten@lshtm.ac.uk)

Keywords: SARS-CoV-2, proteome changes, *Sarbecovirus* evolution, spike protein changes

Abstract

Defining the unique protein features of SARS-CoV-2, the viral agent causing Coronavirus
Disease 2019, may guide efforts to control this pathogen. We examined proteins encoded by the
Sarbecoviruses closest to SARS-CoV-2 using profile Hidden Markov Model similarities to identify
features unique to SARS-CoV-2. Consistent with previous reports, a small set of bat and pangolin-
derived *Sarbecovirus*es show the greatest similarity to SARS-CoV-2. The analysis provided a measure
of total proteome similarity and showed that a small subset of bat *Sarbecoviruses* are closely related
but unlikely to be the direct source of SARS-CoV-2. Spike analysis reveals that the current SARS-CoV-
2 variants of concern have sampled only 36% of the possible spikes changes which have occurred
historically in *Sarbecovirus* evolution. It is likely that new SARS-CoV-2 variants with changes in these
regions are compatible with virus replication and are to be expected in the coming months, unless global
viral replication is severely reduced.

Introduction

Since the first report of Coronavirus Disease 2019 (COVID-19) caused by SARS-CoV-2 in
December 2019 in Wuhan city, China(1)(2) and the World Health Organisation declaring COVID-19 a
global pandemic in March 2020, the disease has continued to affect every part of the world. The SARS-
CoV-2 virus belongs to the *Coronaviridae* family of enveloped positive-sense single-stranded RNA
viruses, *Betacoronavirus* genus, *Sarbecovirus* subgenus. Other *Sarbecoviruses* include SARS-CoV
(the coronavirus causing the SARS outbreak in 2002-2004) and a large number of SARS-like bat
viruses that have been identified. The genomes of *Sarbecoviruses* are 30kb in length, encoding >14
open reading frames (ORFs). Among the structural proteins, the spike protein plays a crucial role in the
virus cell tropism, host range, cell entrance and infectivity and is considered the main protein target for
the host immune response. Other ORFs encode for accessory proteins, many of which modulate host
responses to infection. Investigation of the evolutionary history of SARS-CoV-2 show a clear link to
Sarbecoviruses from bats although no direct animal precursor for SARS-CoV-2 has been identified (3)
(4) (5) (6) (7). We sought to identify unique peptide regions of SARS-CoV-2 compared to all available
Sarbecoviruses to determine the features that might have allowed SARS-CoV-2 to replicate and

transmit efficiently in humans. Comparative analyses of viral proteins would aid in better understanding

of virus biology and pathology, providing insights into the origin of the virus and the conditions that led

to its zoonosis to humans, efficient spread without the need for adaptation, as well as providing leads

for drug and immune targets for effective treatments.

Results and Discussion

Protein domains and profile hidden Markov models. We have explored the genomes across

the *Sarbecovirus* subgenus using profile hidden Markov models (pHMMs) based on viral proteins

which provide a statistical description of the properties of amino acid sequences of viral proteins. The

differences in pHMMs between virus genomes can reveal differences among viral proteins (8) (9).

Efficient tools for preparing and comparing pHMMs are available with HMMER-3 (10) and the method

is useful for comparing large or difficult to align genomes and for identifying changes in functional

regions of the genomes. We have recently used these methods to identify and classify diverse

coronaviruses in the *Coronaviridae* family (11) and to explore large and unwieldy genomes such as

those from the African Swine Fever Virus (12). These methods were employed here to explore the

relationship between SARS-CoV-2 and the other known *Sarbecoviruses* to gain understanding of their

evolutionary history.

a. Prepare Agnostic Domains from a small set of early SARS-CoV-2 genomes

57
Figure 1. Analysis scheme. (A) Profile Hidden Markov Model (pHMM) domains were generated from
a set of 35 early lineage B SARS-CoV-2 genome sequences. All open reading frames were translated
and then sliced into either 44 amino acid peptides with a step size of 22 amino acids or 15 amino acid
peptides with a step size of 8 amino acid. The peptides were clustered using Uclust (13), aligned with
MAFFT (14) and then each alignment was built into a pHMM using HMMER-3 (10). **(B)** The set of
pHMMs were used to query *Sarbecovirus* genome sequences, bit scores were collected as a
measure of similarity between each pHMM and the query sequence. **(C)** Bit-scores were gathered an
analyzed to detect regions that differ between early SARS-CoV-2 genomes and query genomes.
66

Genome scans using custom pHMM domains. We generated overlapping 44 or 15 amino

acid (aa) alignments of all SARS-CoV-2 encoded proteins derived from a set of 35 lineage B SARS-

CoV-2 genomes from early in the pandemic and then prepared pHMMs from each alignment (see Figure
1). The resulting libraries of pHMMs were then used to survey variations across all *Sarbecoviruses*
relative to the initial SARS-CoV-2 found in human cases in late 2019/early 2020.

Using this method, ten early SARS-CoV-2 genomes were compared to a representative subset
of 29 *Sarbecovirus* genomes obtained from humans, bats, civet cats and pangolins (Figure 2) selected
after applying the same analysis to all available *Betacoronavirus* genomes to ensure that we were not
missing any unexpectedly close viral genome regions (results not shown). For each query genome, the
bit-score from the corresponding pHMM from early B lineage SARS-CoV-2 were accumulated and
hierarchical clustering based on the normalized domain bits-scores was performed. The bit-score
describes the similarity between the 44aa (or 15aa) query sequence and the related sequence from
early SARS-CoV-2. The central region of the *Sarbecovirus* genome is conserved across the genome
set with all domains marked as dark or light grey in the Figure 2A clustermap indicating normalized bit-
scores close to 1. This is not unexpected as this central genomic region spans nsp5 to nsp16 proteins
encoding the viral polymerase, other enzymes and non-surface exposed structural proteins of the virus,
which are presumably more constrained and less likely to change than other regions of the virus. In
contrast, the domains displayed in yellow, orange and red in Figure 2A indicate more increasingly
divergent regions between SARS-CoV-2 and the query *Sarbecovirus* genomes (much lower normalized
bit scores). For each domain across the proteome, a measure of the difference between the query and
SARS-CoV-2 domain (1-mean bit-score) was plotted to identify regions that differed extensively across
the *Sarbecoviruses* genome set (Figure 2B). Further details on the relationship between amino acid
changes across a genome or gene set and the metric 1-mean bit-score are found in the Supplemental
Figure 1. The higher 1-mean bit-score values (≥0.3) were found in nsp2, nsp3, nsp4, spike and orf8
and orf9N (Figure 2B), indicating that these proteins differ in the set of SARS-CoV-2 and closely related
Sarbecoviruses. These regions may be dispensable, flexible and can tolerate change and may be
products of neutral evolution they may differ because selective pressure has resulted in new functional
roles or immune escape.
95
96

a.

b.

Figure 2. Proteome differences in SARS-CoV-2 vs close Bat, Human and Civet cat
Sarbecoviruses. All forward open reading frames from the 35 early lineage B SARS-CoV-2 genomes
were translated, and processed into 44 aa peptides (with 22 aa overlap), clustered at 0.65 identity using
Uclust (11), aligned with MAAFT (12) and converted into pHMMs using HMMER-3 (10). The presence
of these domains was sought in a set of *Sarbecovirus* genomes plus the SARS-CoV-2 genomes and
genomes were then clustered using hierarchical clustering based on the normalized domain bit-scores
(e.g. the similarity of the identified query domain to the reference lineage B SARS-CoV-2 domain). Each
row represents a genome, each column represents a domain. Domains are displayed in their order
across the SARS-CoV-2 genome, Red = low normalized domain bit-score (lower similarity to lineage B
SARS-CoV-2) = distant from SARS-CoV-2, Darkest grey = normalized domain bit-score = 1 = highly
similar to lineage B SARS-CoV-2. Groups of coronaviruses were indicated to the right of the figure. **(A)**
Domain differences across the *Sarbecovirus* subgenus. **(B)** For each domain the mean bit-score was
calculated across the entire set of *Sarbecovirus genomes* and the value 1-mean bit-score was plotted
for each domain. Domains are coloured by the proteins from which they were derived with the colour
code indicated below the figure.

The role of pangolins as an amplifying intermediate host of SARS-CoV-2 is important to
document securely, to guide efforts to prevent or prepare for future zoonotic events. A small number of
Sarbecoviruses have been identified in samples from trafficked pangolins in China (15) (7) (16), yet
there is no direct evidence that pangolins host the virus in their natural environment. It remains possible
that the positive pangolins identified in China were infected by viruses encountered after transport to
China. Five CoV sequences from pangolins were included in this analysis (Figure 2), including four
(EPI_ISL_410538,-39,-40,-42) generated by Lam *et al.* (7) after sequencing the original samples
described by Liu *et al.*(15); a 5th genome (Guandong_1_2019_EPI_ISL_410721_2019) deposited by
Xiao *et al.* (16) was included despite unclear progeny for this sequence (17). Four additional CoV
sequences from pangolins from GISAID were excluded due to excessive gaps in the sequence which
precluded the domain analysis. Overall, the pangolin CoVs show many genomic regions of very close

domain homology although the overall similarity across the entire genome is not close enough to
support of direct pangolin-human jump as the source of SARS-CoV-2 (see below).

A number of closely related *Sarbecoviruses* have been reported in bats (18) (19). We focused
on the four bat *Sarbecoviruses* with closest similarity to SARS-CoV-2 (close bats: RaTG13, RmYn02,
MG772933,34). The clustermap and variance analysis (Figure 2A) showed higher similarity across most
of the genome (dark grey sectors) with three proteins (nsp3, spike and orf9) displaying high reduced
bit-scores from the SARS-CoV-2 sequences (Figure 2A, close bats, yellow/red domains). These limited
regions that differ between the 4 closely related bat *Sarbecoviruses* and SARS-CoV-2 indicate virus
elements that needed to adjust to human transmission. The spike differences have been explored in
detail (refs and see below) however it may be important to consider nsp3 and orf9 changes in the
complete analysis.

Spike changes with 15 amino acid domains. To provide more detailed resolution, 15 amino
acid pHMMs across the early lineage B spike protein were prepared and used to examine changes
across the set of close *Sarbecoviruses* and SARS-CoV-2 (Figure 3). The regions of change observed
in the spike protein show an expanded version of the total proteome pattern observed in Figure 2 plus
additional details. Adding the current knowledge of SARS-CoV-2 spike evolution reveals some
important patterns. The S1 domain of spike (the amino-terminal half of the protein) tolerates at large
amount of change with most of the low score domains (red) concentrated here (Figure 3A), consistent
with the surface exposure of the S1 region, and driven by pressure to avoid immune responses which
in turn require the virus to maintain a malleable structure in the exposed S1. Except for the
Guangdong_1 pangolin and RatG13 spikes, the central ACE2 receptor binding region is very different
between the close *Sarbecoviruses* and SARS-CoV-2. The furin cleavage site at the junction between
the S1 and S2 domains is also a region showing a lot of malleability in the *Sarbecovirus* spikes (Figure
3A) and is completely unique to SARS-CoV-2. This has been discussed in detail (20) and is also a site
of frequent change in the current SARS-CoV-2 Variant of Concern (VOC) spike sequence with Q677,
P681 and T717 flanking the furin site showing changes (Figure 3B).

Thirdly, the amino acid changes that have accumulated in the VOC spike protein are marked
in color (Figure 3B) and cluster in regions with high variation (1-mean bit-score > 0.3) suggesting that
Sarbecoviruses have made changes in these regions in previous evolutionary periods and are
continuing to change in SARS-CoV-2 evolution. If all domains with 1-mean bit-score of 0.3 and above
tolerate change in the current SARS-CoV-2 spike protein, this suggests that SARS-CoV-2 has many
additional changes available for future immune evasion. Important regions that show high levels of
historical change are RBD, the furin cleavage site and flanking regions and the NTD.

Figure 3. Spike differences in SARS-CoV-2 vs close Bat, Human and Civet cat *Sarbecoviruses*. All forward spike open reading frames from the 35 early lineage B SARS-CoV-2 genomes were translated, and processed into 15 aa peptides (with 8 aa overlap) and processed into an pHMM library as described in Figure 2. **(A)** shows a hierarchical clustering of 15 amino acid domain bit-scores. **(B)** shows the 1-mean of each domain bit-scores across the genome set, domain values, individual domains that span known amino acid changes in the 5 VOC are colored (see key below panel B). **(C)** The locations of important spike protein features are indicated. NTD: N-terminal domain, RBD: receptor-binding domain, S1: spike 1, S1: Spike 2, TM: transmembrane domain, HR1: helical repeat 1, HR2: helical repeat 2, NTD super: N-terminal domain supersite.

Global proteome similarities. A measure of the total protein distance between the SARS-CoV-2 and any query *Sarbecovirus* can be obtained by summing the normalized bit-scores (SNBS) across the entire query proteome. The encoded proteomes from more distant viruses will show lower SNBS values. We examined SNBSs grouped by virus host for both the 44 amino acid total genome analysis as well as the 15 amino acid spike gene analysis. Overall, the patterns showed that the similarity to SARS-CoV-2 follows the order Pangolin > Bat > human=civet cat (Figure 4) although the small set of close bat *Sarbecoviruses* show some exceptions. Of special interest, the bat coronavirus genome RaTG13 (GenBank MN996532.1) was identified as closely related to the SARS-CoV-2 lineage {Citation} and supports a bat coronavirus being the zoonotic source of the epidemic, although the genetic distance is too far for RaTG13 itself to be a direct source of the pandemic SARS-CoV-2 virus (21)(3). Another close bat coronavirus RmYN02, shows some regions of close identity to SARS-CoV-2 (6), yet overall it is more distant from SARS-CoV-2 than the strain RaTG13 (Figure 4A) due its possible recombinant nature. A single pangolin derived SARS-CoV-2 (Guangdong_1) showed an SNBS value that was also elevated but not as high as the RaTG13 (Figure 4a), the 15 aa spike analysis showed

185   similar patterns except that only the RaTG13 spike displayed the high similarity to SARS-CoV-2 (Figure

186   4b).

188
189
Figure 4. Total domain distances between virus groups. Normalized bit-score sums (NBSS) grouped
into SARS-CoV-2 and *Sarbecoviruses* from pangolin, bat, human and civet cat NBSS for all domains
for each genome were summed. The boxplot shows individual values marked in orange, median values
indicated by horizontal black lines, 1st interquartile ranges marked with a box. The identities of several
high scoring bat and pangolin genomes are indicated. **(A)** NBSS for 44 aa domains across the entire
coronavirus genome. **(B)** NBSS for 15 aa domains across the spike protein.
196

Conclusions

What is special about SARS-CoV-2? Spike changes in SARS-CoV-2 compared to a large set

of known *Sarbecovirus* indicate that the immediate zoonotic source of SARS-CoV-2 is yet to be

identified and reinforces the unique nature of the SARS-CoV-2 genome. The more global analysis of

spike regions in SARS-CoV-2 genomes (Figure 3) revealed the changes that have occurred across the

Sarbecovirues. Combined with the current VOC spike changes, the patterns suggest that SARS-CoV-

2 has a great deal of evolutionary possibilities to avoid immune pressure. This suggests that genomic

variant surveillance should continue and vaccine producers should be prepared to accommodate such

spike changes in the next generation of vaccine updates. In addition to the spike protein, additional

regions of high variance were observed in the nsp3 across all *Sarbecoviruses* (Figure 2) in close bat

and pangolins (Figure 3). The high variance regions flanked and partially overlapped the Macro domain,

which is frequently associated with ADP-deribosylase activity (22)ˌ(23). Variance observed in the ORF8

changes across the set was due to frequent deletion of this ORF, suggesting that the encoded protein
may be dispensable for human infection. Similar loss of ORF8 was observed with the original SARS-
CoV (24) (25) and has been observed in several SARS-CoV-2 lineages as the virus adapted to humans
(26) (27) (28). The ORF9 (N protein) variance observed across *Sarbecoviruses* and the changes in this
protein in VOC strains suggest an additional region that may be adapting to human replication. The
regions of variance identified here may indicate either functional changes in SARS-CoV-2 proteins or
amino acid positions that can be changed without impairing the necessary functions of the protein. The
relatively high mutation rate of SARS-CoV-2 combined with the unprecedented number of SARS-CoV-
2 infections in the world is resulting in massive viral adaptation. Additional experiments are required to
distinguish true functional changes from neutral evolution.

Finally, the detailed spike analysis of Figure 3 reveals 82 domains of 15-aa that have shown
high variation (1-mean bit-score>= 0.3) in the *Sarbecoviruses* while 29 of these domains show changes
in VOC relative to early lineage B SARS-CoV-2. In broad terms, the SARS-CoV-2 evolution observed
in the current VOC has sampled only 36% (29/82) of the possible spikes changes which have occurred
historically in *Sarbecovirus* evolution. It is highly likely that a large number of new SARS-CoV-2 variants
with changes in these regions are possible, compatible with virus replication and expected in the coming
months, unless global viral replication is severely reduced.

Acknowledgements

We thank all global SARS-CoV-2 sequencing groups for their open and rapid sharing of sequence
data and GISAID for providing an effective platform for making these data available. The study is funded
by the Wellcome, DFID - Wellcome Epidemic Preparedness – Coronavirus (AFRICO19, grant
agreement number 220977/Z/20/Z) awarded to MC. DLR and MC receive funding from the MRC
(MC_UU_1201412).

References

1. Li,Q., Guan,X., Wu,P., Wang,X., Zhou,L., Tong,Y., Ren,R., Leung,K.S.M., Lau,E.H.Y., Wong,J.Y., *et al.* (2020)
Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus–Infected Pneumonia. *N. Engl. J.*
Med., 10.1056/NEJMoa2001316.

2. Yang,X., Yu,Y., Xu,J., Shu,H., Xia,J., Liu,H., Wu,Y., Zhang,L., Yu,Z., Fang,M., *et al.* (2020) Clinical course and
outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered,
retrospective, observational study. *Lancet Respir. Med.*, 10.1016/S2213-2600(20)30079-5.

3. Andersen,K.G., Rambaut,A., Lipkin,W.I., Holmes,E.C. and Garry,R.F. (2020) The proximal origin of SARS-CoV-
2. *Nat. Med.*, 10.1038/s41591-020-0820-9.

4. Boni,M.F., Lemey,P., Jiang,X., Lam,T.T.-Y., Perry,B.W., Castoe,T.A., Rambaut,A. and Robertson,D.L. (2020)
Evolutionary origins of the SARS-CoV-2 sarbecovirus lineage responsible for the COVID-19 pandemic.
Nat. Microbiol., **5**, 1408–1417.

5. Zhang,T., Wu,Q. and Zhang,Z. (2020) Probable Pangolin Origin of SARS-CoV-2 Associated with the COVID-19
Outbreak. *Curr. Biol.*, 10.1016/j.cub.2020.03.022.

6. Zhou,H., Chen,X., Hu,T., Li,J., Song,H., Liu,Y., Wang,P., Liu,D., Yang,J., Holmes,E.C., *et al.* (2020) A novel bat
coronavirus reveals natural insertions at the S1/S2 cleavage site of the Spike protein and a possible
recombinant origin of HCoV-19 Microbiology.

7. Lam,T.T.-Y., Jia,N., Zhang,Y.-W., Shum,M.H.-H., Jiang,J.-F., Zhu,H.-C., Tong,Y.-G., Shi,Y.-X., Ni,X.-B., Liao,Y.-S.,
et al. (2020) Identifying SARS-CoV-2-related coronaviruses in Malayan pangolins. *Nature*, **583**, 282–
285.

8. Eddy,S.R. (1998) Profile hidden Markov models. *Bioinformatics*, **14**, 755–763.

9. Eddy,S.R. (1996) Hidden Markov models. *Curr. Opin. Struct. Biol.*, **6**, 361–365.

10. Eddy,S.R. (2011) Accelerated Profile HMM Searches. *PLOS Comput. Biol.*, **7**, e1002195.

11. Phan,M.V.T., Ngo Tri,T., Hong Anh,P., Baker,S., Kellam,P. and Cotten,M. (2018) Identification and
characterization of Coronaviridae genomes from Vietnamese bats and rats based on conserved
protein domains. *Virus Evol.*, **4**.

12. Masembe,C., Phan,M.V.T., Robertson,D.L. and Cotten,M. (2020) Increased resolution of African Swine
Fever Virus genome patterns based on profile HMM protein domains Genomics.

13. Edgar,R.C. (2010) Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*, **26**, 2460–
2461.

14. Katoh,K. and Standley,D.M. (2013) MAFFT Multiple Sequence Alignment Software Version 7:
Improvements in Performance and Usability. *Mol. Biol. Evol.*, **30**, 772–780.

15. Liu,P., Chen,W. and Chen,J.-P. (2019) Viral Metagenomics Revealed Sendai Virus and Coronavirus Infection
of Malayan Pangolins (Manis javanica). *Viruses*, **11**, 979.

16. Xiao,K., Zhai,J., Feng,Y., Zhou,N., Zhang,X., Zou,J.-J., Li,N., Guo,Y., Li,X., Shen,X., *et al.* (2020) Isolation of
SARS-CoV-2-related coronavirus from Malayan pangolins. *Nature*, **583**, 286–289.

17. Chan,Y.A. and Zhan,S.H. (2020) Single source of pangolin CoVs with a near identical Spike RBD to SARS-CoV-
2 Genomics.

18. Tang,X.C., Zhang,J.X., Zhang,S.Y., Wang,P., Fan,X.H., Li,L.F., Li,G., Dong,B.Q., Liu,W., Cheung,C.L., *et al.*
(2006) Prevalence and Genetic Diversity of Coronaviruses in Bats from China. *J. Virol.*, **80**, 7481–7490.

19. Li,W. (2005) Bats Are Natural Reservoirs of SARS-Like Coronaviruses. *Science*, **310**, 676–679.

20. Hoffmann,M., Kleine-Weber,H. and Pöhlmann,S. (2020) A Multibasic Cleavage Site in the Spike Protein of
SARS-CoV-2 Is Essential for Infection of Human Lung Cells. *Mol. Cell*, **78**, 779-784.e5.

21. Andrew Rambaut (2020) Divergence of nCoV-2019 to closest non-human relative. *Virological.org*.

22. Frick,D.N., Virdi,R.S., Vuksanovic,N., Dahal,N. and Silvaggi,N.R. (2020) Molecular Basis for ADP-Ribose
Binding to the Mac1 Domain of SARS-CoV-2 nsp3. *Biochemistry*, **59**, 2608–2615.

23. Lei,J., Kusov,Y. and Hilgenfeld,R. (2018) Nsp3 of coronaviruses: Structures and functions of a large multi-
domain protein. *Antiviral Res.*, **149**, 58–74.

24. Chiu,R.W.K., Chim,S.S.C., Tong,Y., Fung,K.S.C., Chan,P.K.S., Zhao,G. and Lo,Y.M.D. (2005) Tracing SARS-
Coronavirus Variant with Large Genomic Deletion. *Emerg. Infect. Dis.*, **11**, 168–170.

25. Tang,J.W., Cheung,J.L.K., Chu,I.M.T., Sung,J.J.Y., Peiris,M. and Chan,P.K.S. (2006) The Large 386-nt Deletion
in SARS-Associated Coronavirus: Evidence for Quasispecies? *J. Infect. Dis.*, **194**, 808–813.

26. Su,Y.C.F., Anderson,D.E., Young,B.E., Linster,M., Zhu,F., Jayakumar,J., Zhuang,Y., Kalimuddin,S., Low,J.G.H.,
Tan,C.W., *et al.* (2020) Discovery and Genomic Characterization of a 382-Nucleotide Deletion in
ORF7b and ORF8 during the Early Evolution of SARS-CoV-2. *mBio*, **11**, e01610-20,
/mbio/11/4/mBio.01610-20.atom.

27. Gong,Y.-N., Tsao,K.-C., Hsiao,M.-J., Huang,C.-G., Huang,P.-N., Huang,P.-W., Lee,K.-M., Liu,Y.-C., Yang,S.-L.,
Kuo,R.-L., *et al.* (2020) SARS-CoV-2 genomic surveillance in Taiwan revealed novel ORF8-deletion
mutant and clade possibly associated with infections in Middle East. *Emerg. Microbes Infect.*, **9**, 1457–
1466.

28. Young,B.E., Fong,S.-W., Chan,Y.-H., Mak,T.-M., Ang,L.W., Anderson,D.E., Lee,C.Y.-P., Amrun,S.N., Lee,B.,
Goh,Y.S., *et al.* (2020) Effects of a major deletion in the SARS-CoV-2 genome on the severity of
infection and the inflammatory response: an observational cohort study. *Lancet Lond. Engl.*, **396**,
603–611.

298
299

Supplementary Figure 1 demonstrates the relationship between the 1-mean bit-score and amino acid
changes across a sequence alignment. A set of twelve, 15 aa pHMM domains across the central region
of the spike protein were used to demonstrate the relationship between the total amino acid changes
and the resulting pHMM 1-mean bit-scores. An alignment of 38 spike proteins from 10 early SARS-
CoV-2 genomes, 4 close bat coronavirus *Sarbecoviruses* plus 24 additional close *Sarbecoviruses* was
prepared and trimmed to an 87 amino acid region spanning conserved and variable sequences across
the furin cleavage site. The bit-scores(10) for the twelve, 15 aa pHMMs across each query sequence
were gathered, and the mean value for each domain across the set was calculated and converted to a
1-mean bit-score value for ease of visualization (higher value = greater difference from SARS-CoV-2).
The amino acid changes across the alignment are shown in Supplementary Figure 1 panel B.
with each colored bar indicating an amino acid change (or gap, indicated in grey) from the spike of
reference genome NC-045512, The 3' end of the spike S1, the furin cleavage site and the 5' start of the
spike S2 region are indicated in Panel B. Peaks of variability are seen flanking and within the furin site
and in the S2 region (Supplementary Figure 1 panel A). The 1-mean bit-score values for the 12 domains
are plotted in Panel C. and show a related but more softened pattern of variation across the region. The
advantage the pHMM method and the 1-mean bit-score metrics is that the score is weighted to reflect
the type of amino acid change or insertion/deletions and can be determined quickly in large genome
sets using the HMMER tools(10). For our comparative analysis, the cutoff of 0.3 for the 1-mean bit-
score was set to reflect about 25 amino changes in the total aa positions covered by each domain query
(15 aa domain across 38 sequences = 570 total position differences possible).

Supplemental Figure 1. Demonstrating relationship between amino acid changes and 1-mean bit score. (B) The spike proteins encoded by the set of *Sarbecovirus* genomes examined in Figure 2 and 3 were aligned and the amino acid changes from the early SARS-CoV-2 genomes (= NC-045512 sequence) were depicted as colored lines across each sequence. **(A)** The total number of proteins showing a change from the early SARS-CoV-2 genomes (NC-045512) were plotted for each position of the alignment. **(C)** The set of 12, 15 aa pHMMs spanning the central region of the spike (a subset of the pHMMs used for Figure 3) were used to examine the *Sarbecovirus* spike sequences. For each pHMM, the mean bit-score for the entire sequence sequencer set was calculate and the value 1-mean bit-score is plotted for each of the 12 pHMMs. Domains containing known amino acid changes in VOC (Q677, P681, T716) and the domains spanning the furin cleavage site are indicated by color (with the legend found below panel C).