1 **Unique protein features of SARS-CoV-2 relative to other *Sarbecoviruses***
2
3

4 Matthew Cotten[1,2], David L. Robertson[2], My V.T. Phan[1]
5

Affiliations

1. MRC/UVRI & LSHTM Uganda Research Unit, Entebbe, Uganda

2. MRC-University of Glasgow Centre for Virus Research, Glasgow, UK

Correspondence: Matthew Cotten (Matthew.Cotten@lshtm.ac.uk)

Keywords: SARS-CoV-2, proteome changes, *Sarbecovirus* evolution, spike protein changes

Abstract

Defining the unique properties of SARS-CoV-2 protein sequences, has potential to explain the
range of Coronavirus Disease 2019 (COVID-19) severity. To achieve this we compared proteins
encoded by all *Sarbecoviruses* using profile Hidden Markov Model similarities to identify protein
features unique to SARS-CoV-2. Consistent with previous reports, a small set of bat and pangolin-
derived *Sarbecovirus*s show the greatest similarity to SARS-CoV-2 but unlikely to be the direct source
of SARS-CoV-2. Three proteins (nsp3, spike and orf9) showed differing regions between the bat
Sarbecoviruses and SARS-CoV-2 and indicate virus protein features that might have evolved to support
human infection and/or transmission. Spike analysis identified all regions of the protein that have
tolerated change and revealed that the current SARS-CoV-2 variants of concern (VOCs) have sampled
only a fraction (~31%) of the possible spike domain changes which have occurred historically in
Sarbecovirus evolution. This result emphasises the evolvability of these coronaviruses and potential for
further change in virus replication and transmission properties over the coming years.

Introduction

Since the first report of Coronavirus Disease 2019 (COVID-19) caused by SARS-CoV-2 in
December 2019 in Wuhan city, China (Li et al. 2020)(Yang et al. 2020) and the World Health
Organisation declaring COVID-19 a global pandemic in March 2020, the disease has proceeded to
affect every part of the world. The SARS-CoV-2 virus belongs to the *Coronaviridae* family of enveloped
positive-sense single-stranded RNA viruses, *Betacoronavirus* genus, *Sarbecovirus* subgenus. Other
Sarbecoviruses include SARS-CoV (the coronavirus causing the SARS outbreak in 2002-2004) and a
large number of SARS-like bat viruses. The genomes of *Sarbecoviruses* are 30kb in length, encoding
>14 open reading frames (ORFs). Among the structural proteins, the spike protein plays a crucial role
in virus host-cell tropism, host range, cell entry and infectivity, and is considered the main protein target
for protective immune responses. Other virus ORFs encode structural and accessory proteins, many of
which modulate important host responses to infection.
Investigation of the evolutionary history of SARS-CoV-2 shows a clear link to *Sarbecoviruses*
circulating in horseshoe bats although no direct animal precursor for SARS-CoV-2 has been identified
(Andersen et al. 2020) (Boni et al. 2020) (Zhang et al. 2020) (H. Zhou et al. 2020) (Lam et al. 2020).

We sought to identify unique peptide regions of SARS-CoV-2 compared to all available *Sarbecoviruses*
to determine viral features that might be unique to SARS-CoV-2 and that might have allow the virus to
infect, replicate and transmit efficiently in humans. Such a comparative analysis of viral proteins might
provide insights into the origin of the virus and identify the conditions that led to the zoonosis to humans,
efficient spread without the need for much, if any, adaptation (MacLean et al. 2021), as well as providing
leads for drug and immune targets for effective treatments.
47
Results and Discussion
Protein domains and profile hidden Markov models. We have explored the genomes across
the *Sarbecovirus* subgenus using profile hidden Markov models (pHMMs). pHMMs can provide a
detailed statistical description of an amino acid sequence and can be used to detect related domains
found and to document their differences from a reference domain (Eddy 1998) (Eddy 1996). Efficient
tools for preparing and comparing pHMMs are available in the HMMER-3 package (Eddy 2011). This
approach is particularly useful for comparing large or evolutionary divergent genomes. We have
recently used these methods to identify and classify diverse coronaviruses in the *Coronaviridae* family
(Phan et al. 2018) and to explore large and unwieldy genomes such as those from the African Swine
Fever Virus (Masembe et al. 2020). Here pHMMs were used to explore the relationship between SARS-
CoV-2 and the other known *Sarbecoviruses* to gain understanding of their evolutionary history and to
identify regions of encoded viral proteins that are with static to change or are altered across the
Sarbecoviruses.

a. Prepare Agnostic Domains from a small set of early SARS-CoV-2 genomes

b. Use **library** to query related Sarbecovirus genomes

c. Identify protein domains (15 or 44 amino acids) that show differences from early SARS-CoV-2

61
Figure 1. Analysis scheme. (a) Profile Hidden Markov Model (pHMM) domains were generated from
a set of 35 35 early (Pango) lineage B SARS-CoV-2 genome sequences. All open reading frames
were translated and then sliced into either 44 amino acid peptides with a step size of 22 amino acids
or 15 amino acid peptides with a step size of 8 amino acid. The peptides were clustered using Uclust
(Edgar 2010), aligned with MAFFT (Katoh and Standley 2013) and then each alignment was built into
a pHMM using HMMER-3 (Eddy 2011). **(b)** The set of pHMMs were used to query *Sarbecovirus*
genome sequences, bit scores were collected as a measure of similarity between each pHMM and
the query sequence. **(b)** Bit-scores were gathered and analyzed to detect regions that differ between
early SARS-CoV-2 genomes and query genomes.
71

Genome scans using custom pHMM domains. First, some background on the strategy used here. We sought to define the distance of any query virus genome from the early SARS-CoV-2 genome that first began to infect humans in December 2019. To give two levels of resolution, we generated overlapping 44 or 15 amino acid (aa) peptides from all early lineage B SARS-CoV-2 encoded proteins then prepared pHMMs using HMMER-3 (see Figure 1a). The resulting libraries of pHMMs were then used to survey domain diversity across query coronaviruses relative to the initial 2019 SARS-CoV-2. For each pHMM match to a related sequence, a bit-score is generated which provides a metric for how close the query sequence is to the pHMM (Figure 1b). These bit-scores, when collected across an entire viral genome, can provide a sensitive description of similarities and difference between a query genome and the reference genome (Figure 1c). For additional background on the method, Supplementary Figure 1 demonstrates the the sensitivity of pHMMs to detect and distinguish single amino acid substitutions and Supplementary Figure 2 demonstrates the use of pHMMs to identify single amino acid substitution in a crucial region of the SARS-CoV-2 spike protein.

An initial triage was performed using all available Betacoronavirus genomes from GenBank. All full genomes with the taxon id 694002 were retrieved, genomes with gaps were removed to yield a set of 1480 Betacoronavirus genomes. SARS-CoV-2 genomes were initially excluded from the retrieval and then a set of 27 early lineage B genomes were added as a reference. An additional 5 recently reported bat CoV genomes not yet in GenBank (see Supplemental Table 1) were also added. The 44 aa pHMM library was used to query the Betacoronavirus set. For each genome, the bit-score each of 384 pHMMs from early lineage B SARS-CoV-2 sequences was collected and hierarchical clustering based on the normalized domain bits-scores was performed (Figure 2a). Scores colored with dark to light grey indicating domains identical or close to the corresponding domain from early lineage B SARS-CoV-2 and yellow to orange to red indicating increasing distance. Within the Betacoronavirus set the genomes clustered by their taxonomic group and clusters of OC43, MERS-CoV and SARS-CoV and SARS-CoV-2 were observed. The central region of the *Sarbecovirus* genome is conserved across the genome set with all domains marked as dark or light grey in the Figure 2a This is not unexpected as this central region encodes the viral polymerase, other enzymes and non-surface exposed structural proteins of the virus, which are functionally constrained and less likely to allow change than other regions of the virus. In contrast, the domains displayed in yellow, orange and red in Figure 2a indicated more increasingly divergent regions between early SARS-CoV-2 and the query *Sarbecovirus* genomes (much lower normalized bit scores).

The sum of the entire set of bit-scores for a genome was then used to calculate a distance from the early SARS-CoV-2 genome. A histogram of these bit-scores sums show several peaks (Figure 2b) with majority of the Betacoronavirus genomes (mostly OC43 and MERS-CoV) clustering around 100 units and a subset of virus genomes with bit-scores >200 units. This >200 set included the SARS-CoV-2, SARS-like CoVs from bats and all the SARS-CoV genomes (Figure 2c). A second triage retained a set of close genomes all with bit-score sums >320 (Figure 2e) that was used more detailed analysis. For simplicity, the 27 early B SARS-CoV-2 genomes in the set (which were nearly identical) were reduced to 5 resulting in a 19 genomes in the close set: 14 bat/pangolin CoV and 5 SARS-CoV-2 (Figure 2e).

Figure 2 Triage of Betacoronaviruses. All Betacoronavirus genomes (excluding SARS-CoV-2) were
retrieved from GenBank using the query ((txid694002[Organism] AND 24000[SLEN]:40000[SLEN] NOT
patent)) NOT txid2697049[Organism]) generating a set of 1581 genomes that were screened to remove
genomes with gaps to yield a set of 1480 genomes. A small set of 27 early lineage B SARS-CoV-2
genomes from December 2019/January 2020 were added as markers. A library of 44 amino acid
pHMMs prepared from early B SARS-CoV-2 genomes was used and the bit-scores for each pHMM
were gathered and used to cluster the genomes **(a)** with each domain score indicated by color (dark
grey = 1= very similar to SARS-CoV-2 to red = low = distant from SARS-CoV-2). The total bit-scores
sum for each genome was calculated (see histograms of all total bitscore sums **(b))**. A total bitscore
sum of 200 was used to select for the CoV genomes most similar to SARS-CoV-2. The clustering of
this subset of CoV genomes **(c)** included the SARS-CoV-2 genomes, a large number of SARS-CoV
genomes and a smaller number of SARS-like CoVs. A cut off of 320 for total bit-scores sum **(d)** was
used to identify the closest CoV genomes which were then used for the subsequent analyses reported
in Figures 3, 4 and 5.

We next focused on the bat *Sarbecoviruses* with closest similarity to SARS-CoV-2 in at least

part of their genomes due to recombinant histories (see Supplementary Table 1 for genome details and

references). The clustermap and variance analysis (Figure 3a) showed higher similarity across most of

the genome (dark grey sectors) with three proteins (nsp3, spike and orf9) displaying reduced bit-scores

compared to SARS-CoV-2 (Figure 3a, yellow, red domains). These differing regions between the bat

Sarbecoviruses and SARS-CoV-2 indicate virus protein features that might have evolved to support

human infection and/or transmission. The spike differences are explored in detail below however it may

be important to consider nsp3, ORF9 (and perhaps nsp4 and ORF8) in future analyses.

Figure 3. Proteome differences in SARS-CoV-2 versus close bat *Sarbecoviruses*. All forward open reading frames from the 35 early lineage B SARS-CoV-2 genomes were translated, and processed into 44 aa peptides (with 22 aa overlap), clustered at 0.65 identity using Uclust (11), aligned with MAAFT (12) and converted into pHMMs using HMMER-3 (Eddy 2011). The presence of these domains was sought in a set of *Sarbecovirus* genomes plus the SARS-CoV-2 genomes. These were then clustered using hierarchical clustering based on the normalized domain bit-scores (e.g. the similarity of the identified query domain to the reference lineage B SARS-CoV-2 domain). Each row represents a genome, each column represents a domain. Domains are displayed in their order across the SARS-CoV-2 genome, Red = low normalized domain bit-score (lower similarity to lineage B SARS-CoV-2), i.e., higher distance from lineage B SARS-CoV-2, darkest grey = normalized domain bit-score = 1, i.e., highly similar to lineage B SARS-CoV-2. Groups of coronaviruses are indicated to the right of the figure. **(a)** Domain differences across the *Sarbecovirus* subgenus. **(b)** For each domain the mean bit-score was calculated across the entire set of *Sarbecovirus genomes* and the value 1-mean bit-score was plotted for each domain. Domains are coloured by the proteins from which they were derived with the colour code indicated below the figure. **(c)** Schematic of open reading frames or protein products of SARS-CoV-2.

Spike changes with 15 amino acid domains. Using the same strategy described in Figure 3, we performed a triage of the Betacoronaviruses with 15 amino acid pHMMs prepared from early lineage B spike protein (Figure 4) and selected for CoV genomes encoding close Spike proteins. The high bit-score spike set largely overlapped with the high bit-score full genome set suggesting that spike is a good surrogate for full genome homology.

Key features of the spike protein are outlined in Figure 4c. The analysis revealed regions of spike that historically have tolerated change. In general the S1 subunit of spike (the amino-terminal half of the protein) displayed a large amount of diversity with most of the low score domains (more distant from SARS-CoV-2, marked in red) concentrated here (Figure 4a). This is consistent with the surface exposure of S1 on the virion and with protein changes driven by pressure to avoid immune responses. The central ACE2 receptor binding region (Figure 4c) was very different between the close *Sarbecoviruses* and SARS-CoV-2. The furin cleavage site at the junction between the S1 and S2 domains (Figure 4c) is also a region showing a lot of diversity across the *Sarbecovirus* spikes (Figure 4a) and appears completely unique to SARS-CoV-2. This has been discussed in detail (Hoffmann et al. 2020) and is also a site of frequent change in the current SARS-CoV-2 Variant of Concern (VOC) spike sequences with Q677, P681 and T717 flanking the furin site showing changes (Figure 4c).

The domains that where amino acid changes have appeared in VOC spike proteins are marked

in color (Figure 4b) and largely appear in domains with high variation (Figure 4b, 1-mean bit-score >

0.3) suggesting that *Sarbecoviruses* have made changes in these regions in previous evolutionary

periods and are continuing to change in SARS-CoV-2 evolution. Of the 88 spike domains showing high

variation in *Sarbecoviruses (*1 - mean bit-scores >= 0.3 units), only 27 of the domains (31%) have

accumulated substitutions or deletions. This indicates a very large potential in the SARS-CoV-2 spike

protein for tolerating future change. Important regions that have shown high levels of historical change

are the NTD, the RBD and the furin cleavage site and flanking regions.

180 Figure 4. Spike differences in SARS-CoV-2 versus close bat *Sarbecoviruses*. All forward spike
open reading frames from the 35 early lineage B SARS-CoV-2 genomes were translated, and
processed into 15 aa peptides (with 8 aa overlap) and processed into an pHMM library as described in
Figure 2. **(a)** Shows a hierarchical clustering of 15 amino acid domain bit-scores. **(b)** Shows the 1-mean
of each domain bit-scores across the genome set, domain values, individual domains that span known
amino acid changes in the 6 VOC and VOIs (B.1.1.7, B.1.351, B.1.525, P.1, B.1.617.2 and A.23.1) are
colored (see key below panel B). **(c)** The locations of important spike protein features are indicated.
NTD: N-terminal domain, RBD: receptor-binding domain, S1: spike 1, S1: Spike 2, TM: transmembrane
domain, HR1: helical repeat 1, HR2: helical repeat 2, NTD super: N-terminal domain supersite.
189

Global proteome similarities. As described in Figure 2, a measure of the total protein distance

between the SARS-CoV-2 and any query *Sarbecovirus* can be obtained by summing the normalized

bit-scores (SNBS) across the entire query proteome. We examined SNBSs grouped by virus host for

the 44 amino acid total genome analysis and the 15 amino acid spike gene analysis. The potential role

of pangolins as an amplifying intermediate host of SARS-CoV-2 is important to document securely, to

guide efforts to prevent or prepare for future zoonotic events. A small number of *Sarbecoviruses* have

been identified in samples from trafficked pangolins in China (Liu et al. 2019) (Lam et al. 2020) (Xiao et

al. 2020), yet there is no direct evidence that pangolins host the virus in their natural environment. It is

thus likely these pangolins identified in China were infected by viruses encountered after transport to
China, consistent with reports of disease in these animals. Five CoV sequences from pangolins were
included in this analysis (Supplementary Table 1), including four generated by Lam *et al.* (Lam et al.
2020) after sequencing the original samples described by Liu *et al.*(Liu et al. 2019); a 5th genome
(MP789) was deposited by Liu et al..
The bat coronavirus genome RaTG13 (GenBank MN996532.1) was identified as closely related
to the SARS-CoV-2 lineage (P. Zhou et al. 2020) and supports a bat coronavirus being the zoonotic
source of the epidemic, although despite the close genetic distance it is too far in time (decades) for
RaTG13 itself to be a direct source of the pandemic SARS-CoV-2 virus (Boni et al. 2020). The next
closest bat coronavirus RsYN06, shows some regions of even close identity to SARS-CoV-2 than
RaTG13 (H. Zhou et al. 2020) (Figure 4a) due its possible recombinant nature. A single pangolin derived
SARS-CoV-2 (MP789) showed an SNBS value that was also elevated but not as high as the RaTG13
(Figure 5a), the 15 aa spike analysis showed similar patterns except that only the RaTG13 spike
displayed the high similarity to SARS-CoV-2 (Figure 5b).
212

213
214
Figure 5. Total domain distances between virus groups. Normalized bit-score sums (NBSS) grouped
into SARS-CoV-2 and *Sarbecoviruses* from pangolin, bat, for all domains for each genome were
summed. The boxplot shows individual values marked in orange, median values indicated by horizontal
black lines, 1st interquartile ranges marked with a box. The identities of several high scoring bat and
pangolin genomes are indicated. **(A)** NBSS for 44 aa domains across the entire coronavirus genome.
(B) NBSS for 15 aa domains across the spike protein.
221
222

Conclusions

What is special about SARS-CoV-2? Spike changes in SARS-CoV-2 compared to the close set
of *Sarbecovirus* genomes indicate that the immediate zoonotic source of SARS-CoV-2 is yet to be
identified due to the unique nature of the SARS-CoV-2 genome. The more detailed analysis of spike
regions in SARS-CoV-2 genomes (Figure 3) revealed the extent of the changes that have occurred
across the *Sarbecovirues*. Combined with the current VOC spike changes (from lineages B.1.1.7,
B.1.351, B.1.525, P.1, B.1.617.2 and A.23.1), the patterns suggest that SARS-CoV-2 has a great deal

of possibilities for further evolution, presumably enabling persistence and avoid immune responses.
This emphasises the importance of genomic variant surveillance for monitoring for further changes in
virus biology that may have implications for spread and disease severity. Vaccine producers should be
prepared to accommodate such spike changes in the next generation of vaccine updates. In addition
to the spike protein, additional regions of high variance were observed in the nsp3 across all
Sarbecoviruses (Figure 2) in close bat and pangolins (Figure 3).

The high variance regions flanked and partially overlapped the Macro domain, which is
frequently associated with ADP-deribosylase activity (Frick et al. 2020),(Lei et al. 2018). Variance
observed in the ORF8 changes across the set was due to frequent deletion of this ORF, suggesting
that the encoded protein may be dispensable for human infection. Similar loss of ORF8 was observed
with the original SARS-CoV (Chiu et al. 2005) (Tang et al. 2006) and has been observed in several
SARS-CoV-2 lineages as the virus adapted to humans (Su et al. 2020) (Gong et al. 2020) (Young et al.
2020). The ORF9 (N protein) variance observed across *Sarbecoviruses* and the changes in this protein
in VOC strains suggest an additional region that may be adapting to human replication. The regions of
variance identified here may indicate either functional changes in SARS-CoV-2 proteins or amino acid
positions that can be changed without impairing the necessary functions of the protein. The relatively
high mutation rate of SARS-CoV-2 combined with the unprecedented number of SARS-CoV-2
infections in the world is resulting in massive viral adaptation. Additional experiments are required to
distinguish true functional changes from neutral evolution.

Finally, the detailed spike analysis of Figure 4 revealed 88 15aa spike domains showing high
variation while only 27 (31%) have accumulated substitutions or deletions in the current epidemic in
VOCs and VOIs indicating a large potential for tolerating future change. It is highly likely that a large
number of new SARS-CoV-2 variants with changes in these regions will evolve, compatible with similar
levels of virus replication but tolerating significant antigenic change in the coming years, unless global
SARS-CoV-2 spread is severely curtailed.

Acknowledgements
We thank all global SARS-CoV-2 sequencing groups for their open and rapid sharing of sequence
data and GISAID for providing an effective platform for making these data available. This work was
supported by the Wellcome, DFID - Wellcome Epidemic Preparedness – Coronavirus (AFRICO19,
grant agreement number 220977/Z/20/Z) awarded to MC. DLR and MC receive funding from the MRC
(MC_UU_1201412), by the UK Medical Research Council (MRC/UKRI) and the UK Department for
International Development (DFID) under the MRC/DFID Concordat agreement (grant agreement
number NC_PC_19060).

References
Andersen KG, Rambaut A, Lipkin WI, Holmes EC, Garry RF. 2020. The proximal origin of SARS-CoV-2. *Nat. Med.*
[Internet]. Available from: http://www.nature.com/articles/s41591-020-0820-9

Boni MF, Lemey P, Jiang X, Lam TT-Y, Perry BW, Castoe TA, Rambaut A, Robertson DL. 2020. Evolutionary
origins of the SARS-CoV-2 sarbecovirus lineage responsible for the COVID-19 pandemic. *Nat.*
Microbiol. 5:1408–1417.

Chan YA, Zhan SH. 2020. Single source of pangolin CoVs with a near identical Spike RBD to SARS-CoV-2.
Genomics Available from: http://biorxiv.org/lookup/doi/10.1101/2020.07.07.184374

Chiu RWK, Chim SSC, Tong Y, Fung KSC, Chan PKS, Zhao G, Lo YMD. 2005. Tracing SARS-Coronavirus Variant
with Large Genomic Deletion. *Emerg. Infect. Dis.* 11:168–170.

Eddy SR. 1996. Hidden Markov models. *Curr. Opin. Struct. Biol.* 6:361–365.

Eddy SR. 1998. Profile hidden Markov models. *Bioinformatics* 14:755–763.

Eddy SR. 2011. Accelerated Profile HMM Searches. *PLOS Comput. Biol.* 7:e1002195.

Edgar RC. 2010. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* 26:2460–2461.

Frick DN, Virdi RS, Vuksanovic N, Dahal N, Silvaggi NR. 2020. Molecular Basis for ADP-Ribose Binding to the
Mac1 Domain of SARS-CoV-2 nsp3. *Biochemistry* 59:2608–2615.

Gong Y-N, Tsao K-C, Hsiao M-J, Huang C-G, Huang P-N, Huang P-W, Lee K-M, Liu Yi-Chun, Yang S-L, Kuo R-L, et
al. 2020. SARS-CoV-2 genomic surveillance in Taiwan revealed novel ORF8-deletion mutant and clade
possibly associated with infections in Middle East. *Emerg. Microbes Infect.* 9:1457–1466.

Henikoff S, Henikoff JG. 1992. Amino acid substitution matrices from protein blocks. *Proc. Natl. Acad. Sci. U. S.
A.* 89:10915–10919.

Hoffmann M, Kleine-Weber H, Pöhlmann S. 2020. A Multibasic Cleavage Site in the Spike Protein of SARS-CoV-
2 Is Essential for Infection of Human Lung Cells. *Mol. Cell* 78:779-784.e5.

Katoh K, Standley DM. 2013. MAFFT Multiple Sequence Alignment Software Version 7: Improvements in
Performance and Usability. *Mol. Biol. Evol.* 30:772–780.

Lam TT-Y, Jia N, Zhang Y-W, Shum MH-H, Jiang J-F, Zhu H-C, Tong Y-G, Shi Y-X, Ni X-B, Liao Y-S, et al. 2020.
Identifying SARS-CoV-2-related coronaviruses in Malayan pangolins. *Nature* 583:282–285.

Lei J, Kusov Y, Hilgenfeld R. 2018. Nsp3 of coronaviruses: Structures and functions of a large multi-domain
protein. *Antiviral Res.* 149:58–74.

Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, Ren R, Leung KSM, Lau EHY, Wong JY, et al. 2020. Early
Transmission Dynamics in Wuhan, China, of Novel Coronavirus–Infected Pneumonia. *N. Engl. J.
Med.*:NEJMoa2001316.

Liu P, Chen W, Chen J-P. 2019. Viral Metagenomics Revealed Sendai Virus and Coronavirus Infection of
Malayan Pangolins (Manis javanica). *Viruses* 11:979.

MacLean OA, Lytras S, Weaver S, Singer JB, Boni MF, Lemey P, Kosakovsky Pond SL, Robertson DL. 2021.
Natural selection in the evolution of SARS-CoV-2 in bats created a generalist virus and highly capable
human pathogen.Tully D, editor. *PLOS Biol.* 19:e3001115.

Masembe C, Phan MVT, Robertson DL, Cotten M. 2020. Increased resolution of African Swine Fever Virus
genome patterns based on profile HMM protein domains. Genomics Available from:
http://biorxiv.org/lookup/doi/10.1101/2020.01.12.903104

Phan MVT, Ngo Tri T, Hong Anh P, Baker S, Kellam P, Cotten M. 2018. Identification and characterization of
Coronaviridae genomes from Vietnamese bats and rats based on conserved protein domains. *Virus
Evol.* [Internet] 4. Available from:
https://academic.oup.com/ve/article/doi/10.1093/ve/vey035/5250438

Su YCF, Anderson DE, Young BE, Linster M, Zhu F, Jayakumar J, Zhuang Y, Kalimuddin S, Low JGH, Tan CW, et al.
2020. Discovery and Genomic Characterization of a 382-Nucleotide Deletion in ORF7b and ORF8

during the Early Evolution of SARS-CoV-2.Schultz-Cherry S, editor. *mBio* 11:e01610-20,
/mbio/11/4/mBio.01610-20.atom.

Tang JW, Cheung JLK, Chu IMT, Sung JJY, Peiris M, Chan PKS. 2006. The Large 386-nt Deletion in SARS-
Associated Coronavirus: Evidence for Quasispecies? *J. Infect. Dis.* 194:808–813.

Xiao K, Zhai J, Feng Y, Zhou N, Zhang X, Zou J-J, Li N, Guo Y, Li X, Shen X, et al. 2020. Isolation of SARS-CoV-2-
related coronavirus from Malayan pangolins. *Nature* 583:286–289.

Yang X, Yu Y, Xu J, Shu H, Xia J, Liu H, Wu Y, Zhang L, Yu Z, Fang M, et al. 2020. Clinical course and outcomes of
critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective,
observational study. *Lancet Respir. Med.*:S2213260020300795.

Young BE, Fong S-W, Chan Y-H, Mak T-M, Ang LW, Anderson DE, Lee CY-P, Amrun SN, Lee B, Goh YS, et al.
2020. Effects of a major deletion in the SARS-CoV-2 genome on the severity of infection and the
inflammatory response: an observational cohort study. *Lancet Lond. Engl.* 396:603–611.

Zhang T, Wu Q, Zhang Z. 2020. Probable Pangolin Origin of SARS-CoV-2 Associated with the COVID-19
Outbreak. *Curr. Biol.*:S0960982220303602.

Zhou H, Chen X, Hu T, Li J, Song H, Liu Y, Wang P, Liu D, Yang J, Holmes EC, et al. 2020. A novel bat coronavirus
reveals natural insertions at the S1/S2 cleavage site of the Spike protein and a possible recombinant
origin of HCoV-19. Microbiology Available from:
http://biorxiv.org/lookup/doi/10.1101/2020.03.02.974139

Zhou P, Yang X-L, Wang X-G, Hu B, Zhang L, Zhang W, Si H-R, Zhu Y, Li B, Huang C-L, et al. 2020. A pneumonia
outbreak associated with a new coronavirus of probable bat origin. *Nature* 579:270–273.

Supplementary Material.

Supplementary Table 1. Close bat coronaviruses.

Genome name	GenBank or GISAID	Year	Reference
RpYN06	EPI_ISL_1699446	2020	unpublished
RmYN02	EPI_ISL_412977	2019	Zhou et al. Curr Biol. 2020 Jun 8;30(11):2196-2203.e3. PMID: 32416074
PrC31	EPI_ISL_1098866	2018	unpublished
RaTG13	MN996532	2013	Zhou et al. Nature. 2020 Mar;579(7798):270-273 PMID: 32015507
RshSTT182	EPI_ISL_852604	2010	http://biorxiv.org/lookup/doi/10.1101/2021.01.26.428212
RshSTT200	EPI_ISL_852605	2010	http://biorxiv.org/lookup/doi/10.1101/2021.01.26.428212
CoVZ45	MG772933	2017	Hu et al. Emerg Microbes Infect 7 (1), 154 (2018) PMID: 30209269
CoVZXC21	MG772934	2015	Hu et al. Emerg Microbes Infect 7 (1), 154 (2018) PMID: 30209269
RaCS203	MW251308	2020	Wacharapluesadee et al. Nat Commun. 2021 12(1):972 PMID: 33563978
Rc-0319	LC556375	2013	Murakami et al. EID 2020 Dec;26(12):3025-3029 PMID: 33219796
GX-P4L	MT040333	2017	Lam et al. 2020. *Nature* 583:282–285. PMID: 32218527
GX-P1E	MT040334	2017	Lam et al. 2020. *Nature* 583:282–285. PMID: 32218527
GX-P5L	MT040335	2017	Lam et al. 2020. *Nature* 583:282–285. PMID: 32218527
GX-P5E	MT040336	2017	Lam et al. 2020. *Nature* 583:282–285. PMID: 32218527
MP789	MT121216	2019	Liu et al. PLoS Pathog. 16 (5), e1008421 (2020) PMID: 32407364

Supplementary Figure 1 to illustrate pHMM detection of amino acid changes.

We sought to illustrate the ability of pHMMs to detect amino acid differences between a

reference and a query sequence. A reference peptide containing the twenty amino acids was used to

prepare a pHMM. A test set of mutant sequences was prepared by sequentially changing each amino

acid to each of the other 20 amino acids. This set of 400 sequences was then queried with the wildtype

20aa profile HMM, the bit-scores describing each match were collection. The distribution of bit-scores

from the 400 pHMM matches (Supplementary Figure 1a) was broad, consistent with the methods ability

to report not only an amino acid changes but the type of amino acid change. The pattern of all amino

acid changes across all 20 AA peptide is displayed in clustermap (Supplementary Figure 1b) with each

column corresponding to a single amino acid and each row showing the score if that amino acid were

changed to another amino acid. An amino acid that is frequent (e.g. alanine (A)) shows higher bit-scores

across the set of changes than rarer amino acids such as cysteine (C), histidine (H), tryptophan (W) or

proline (P). This spectrum closely reflects the BLOSUM62 substitution matrix (Henikoff and Henikoff

1992) and demonstrates the capacity of a pHMM match both to detect changes in proteins as they

evolve and to distinguish different types of changes.

Supplementary Figure 1. pHMM bit-score values as a measure of the type of amino acid change. A
sequence encoding all 20 amino acids (ACDEFGHIKLMNPQRSTVWY) was used to prepare a pHMM.
A test set of mutant sequences was prepared by changing each amino acid to each of the other 20
amino acids. This set of 400 sequences was then queried with the wildtype 20aa profile HMM, bit-
scores for each match were collected in a matrix. **(a)** a histogram of all observed normalized bit-scores,
the peak at 1.00 is due to changes to self (e.g. A to A change). **(b)** heatmap of normalized bit-scores,
each columns represents a position in the 20 AA wt peptide, each row represents a change at that
position to the indicated amino acid. The normalized bit-scores were color coded with no change from
wildtype amino acid (dark grey) to the largest change from the wildtype amino acid (dark red). **(c)**
Variance of normalized bit-scores from panel b were calculated for each position.

In a second analysis we examined a peptide sequence spanning the important furin cleavage

site in the SARS-CoV-2 spike protein. Mutations in this region have appeared in several VOCs (A.23.1:

P681R, B.1.1.7: P681H, B.1.525: Q677H) and we wanted to document the sensitivity of pHMM

matching to detect single amino acid changes. Similar to Supplementary Figure 1, we prepared a pHMM

from the wildtype 14aa sequence spanning the furin site. For a test set we systematically change

position to each of the 20 amino acids and then gathered the bit-scores for the wildtype pHMM matching

each test peptide. Similar to Supplementary Figure 1, the range of normalized bit-scores scores

included a peak at 1.00 (self sequence matched to self) plus a range of lower values demonstrating

the breadth of possible pHMM bits-scores for any possible single amino acid changes in the 14 amino

acid peptide (Supplementary Figure 2a). The heatmap of the resulting normalized bit-scores

(Supplementary Figure 2b) reveals some patterns. Most changes of the proline adjacent to the

cleavage site resulted in a large reduction in bit-scores, whereas other changes resulted in detectable,

distinct, but less dramatic bit-scores.

Supplementary Figure 2. Amino acid changes across the P681 region of the spike protein. A 14
amino acid sequence spanning the SARS-CoV-2 spike position 681 and the adjacent furin cleavage
site   (YQTQTNSPRRARSV) was used to prepare a pHMM. A test set of mutant sequences was
prepared by changing each amino acid to each of the other 20 amino acids. This set of 280 sequences
was then queried with the wildtype 14aa pHMM, bit-scores were collected. Panel a, a histogram of all
observed normalized bit-scores, the peak at 1.00 due to changes to self (e.g. A to A change). Panel b,
heatmap of normalized bit-scores, each columns represents a position in the 14 AA wt peptide, each
row represents a change at that position to the indicated amino acid. The normalized bit-scores were
color coded with no change from wildtype amino acid (dark grey) to the largest change from the wildtype
amino acid (dark red). Panel C. Variance of normalized bit-scores from Panel b were calculated for
each amino acid position across the peptide.