The determinants of the subjective perception of sleep: insights from a high-density EEG study with serial awakenings

Aurélie Stephan[1,] Sandro Lecci[1], Jacinthe Cataldi[1], Francesca Siclari[1,2*]

[1] Center for Investigation and Research on Sleep and [2] Department of Clinical Neurosciences
Lausanne University Hospital (CHUV), 1010 Switzerland.
*Correspondence: francesca.siclari@chuv.ch

Abstract. What determines the feeling of being asleep? Standard sleep recordings only incompletely reflect subjective aspects of sleep and some individuals with so-called sleep misperception frequently feel awake although sleep recordings indicate clear-cut sleep. Here we performed 787 awakenings in 20 good sleepers and 10 individuals with severe sleep misperception to interview them about their subjective sleep depth while they underwent high-density EEG sleep recordings (256-channels). Surprisingly, in good sleepers, sleep was subjectively lightest in the first two hours of Non-rapid eye movement (NREM) sleep, generally considered the 'deepest' sleep, and deepest in rapid eye movement (REM) sleep. Compared to good sleepers, sleep misperceptors felt more frequently awake during sleep, reported overall lighter REM sleep and had more thought-like conscious experiences. In both groups, subjective sleep depth positively correlated with dream-like features of conscious experiences. At the EEG level, spatially widespread high-frequency power was inversely related to subjective sleep depth in NREM sleep in both groups and in REM sleep in misperceptors. Taken together, these findings challenge the widely held notion that 'deep' (slow wave) sleep best accounts for feeling soundly asleep. Instead, they suggest that subjective sleep depth is inversely related to a neurophysiological process that predominates in NREM sleep early in the night, becomes quiescent in REM sleep and is reflected in high-frequency EEG-activity. In sleep misperceptors, this neurophysiological process is more active and spatially widespread, and abnormally persists into REM sleep. Thus, it is not the presence of 'sleep rhythms' but rather the absence of 'wake-like' EEG activity that predicts the feeling of being deeply asleep. These findings will help identify the neuromodulatory systems involved in subjective sleep depth and are therefore relevant for future studies aiming to improve subjective sleep quality.

What accounts for the subjective perception of being asleep? Why do we sometimes feel deeply asleep while at other times we do not? These questions may well appear trivial, as most individuals intuitively seem to know whether they were awake or asleep and how deep their sleep was. Yet the relationship between objective and subjective sleep parameters is not as straightforward as one may think. A recent study involving 1483 participants showed for instance that conventional polysomnographic parameters only explain a small part of the variance in subjective sleep quality [1]. Several studies have shown that most individuals tend to consistently overestimate sleep latencies [2–5] and conversely, appear to perceive a large part of intranight awakenings as sleep [4,6,7]. An extreme mismatch between subjective estimations and objective measures of sleep is seen in patients with so-called 'paradoxical insomnia' or 'sleep state misperception', who feel awake during large parts of the night although sleep recordings indicate clear-cut sleep. Despite the high prevalence of insomnia in the general population, it still remains unknown what accounts for sleep misperception in this population [2,8]. Determining what accounts for feeling deeply asleep is crucial if one wants to develop targeted approaches to improve sleep quality, both in the genral population and in insomnia sufferers. Despite this obvious interest, few research studies have systematically evaluated the determinants of the subjective perception of sleep.

Sleep can be broadly divided into rapid eye movement (REM) sleep and NREM sleep. REM sleep is characterized by rapid eye movements and fast, low-amplitude electroencephalographic (EEG) activity, whereas NREM sleep features high-amplitude, low-frequency EEG slow waves and graphoelements called spindles. According to the current classification of sleep stages, NREM sleep can be further divided into three stages, in which the EEG becomes progressively slower (stages NREM 1, 2 and 3). Stage NREM 3, in which high-amplitude slow waves occur nearly continuously, is referred to as 'slow wave sleep'. This stage is also called 'deep' sleep, presumably because early studies showed that arousal thresholds were high in this stage and varied as a function of slow wave activity [9]. In addition, subjects awoken from deep sleep often display impaired cognitive performances (sleep inertia) [10] and frequently report having been unconscious before awakening [11,12]. But is objectively defined 'deep' sleep necessarily perceived as subjectively deep?

Early work in which awakenings were performed serially in the falling asleep period revealed that approximately 50% of subjects felt asleep 4 min after the first sleep spindle and 90% after 16 min. However, subsequent studies performing awakenings in stable NREM 2 sleep at different times of the night revealed that subjects felt awake during polysomnographically defined sleep in a high proportion of cases [3,5,13–16], reaching 40-45% in some studies [5,17,18], suggesting a dissociation between standard sleep parameters and the subjective estimation of sleep. Studies sampling sleep

perception in REM sleep reported variable, although overall lower objective-subjective discrepancies in this stage compared to NREM sleep stage 2 [5,14,16,19].

Other studies have evaluated the correlates of sleep misperception more indirectly, by comparing EEG spectral power between insomnia patients and good sleepers. These studies have documented relative increases in high-frequency spectral power, comprising the alpha (8-12 Hz), sigma (12-16 Hz)) or beta (16-32 Hz) frequency bands [20–23], and/or relative decreases in low-frequency power including the delta and theta band [23–26] in insomnia patients compared to healthy controls. Some studies have extended these findings to insomnia patients with sleep misperception [4,24,25]. It remains however open whether the EEG differences between the populations are due to sleep misperception *per se* or secondary to other factors that differ between the groups, including cognitive or psychological changes [27], or insomnia itself.

Studies based on retrospective estimations of sleep duration given upon awakening in the morning reported positive correlations between the degree of sleep underestimation and some EEG features, including alpha [28], sigma [28] and/or beta power [28,29], the delta/beta power ratio [30], delta-alpha sleep patterns [31] and signatures of 'light' sleep within 'deep' sleep [32]. One study combining serial awakenings with EEG spectral power measures found that feeling awake instances in light NREM 1 and 2 sleep (n=14) were associated with higher relative alpha, beta and gamma power compared to instances of feeling asleep (n=27)[33].

To determine what accounts for the moment-to-moment perception of sleep across different sleep stages, we sampled subjective experiences extensively by awakening subjects 787 times while they were sleeping in the sleep laboratory. We recorded brain activity with high-density EEG, which combines the excellent temporal resolution of EEG with an increased spatial resolution upon source modeling. This technique has been used successfully in the past to identify local correlates of subjective experiences during sleep, including dream contents [34]. Recent studies using techniques with high spatial resolution, including positron emission tomography [35], functional magnetic resonance [33] and high-density EEG [22] have indeed reported localized changes in brain activity in insomnia patients, opening the possibility that they may play a role in sleep perception. To understand whether sleep perception shares a similar substrate in healthy subjects and insomnia patients, we performed the same paradigm in 20 good sleepers as well as in 10 insomnia sufferers with sleep misperception (sleep misperceptors), who severely underestimated their sleep duration with respect to the general population [4].

Results

Participants underwent two overnight high-density EEG (hd-EEG) sleep recordings during which they were awakened multiple times with a computerized alarm sound [12]. After each awakening, they were asked to estimate whether they had been feeling awake (FAW) or asleep (FAS) prior to the alarm sound. When they had felt asleep, they also had to estimate the degree of sleep depth on a scale from 1 to 5. In addition, they had to describe what was going through their mind immediately before the alarm sound and to rate their experiences on previously validated sclaes [12,34].

Distribution of awakenings

Overall, we performed 787 awakenings. Good sleepers and sleep misperceptors did not differ in the number of awakenings per subject [26.7 +/- 7.1 vs. 25.3 +/- 5.7, t-test p=0.56], the proportion of NREM (N2 and N3) and REM awakenings [good sleepers: 76.8 +/- 4.6% vs 23.2 +/- 4.6%; sleep misperceptors: 75.7+/- 2.8% vs 24.3 +/- 2.9%; t-test on NREM p=0.42] or the time spent asleep before each awakening [NREM: 13.7 +/- 1.2 min vs 14.9 +/- 1.8 min, t-test p=0.07; REM: 21.6 +/- 7.9 min vs 20.1 +/- 3.8 min, t-test p=0.49].

Effect of group, sleep stage and time of the night on perceived state (FAW/FAS) and perceived sleep depth

To evaluate whether the proportion of FAS and FAW differed between sleep misperceptors and good sleepers, REM and NREM (stages N2 & N3) sleep, and according to the time of the night, we computed a generalized linear mixed model, with perceived state (FAW vs. FAS) as the categorical dependent variable, group (good sleepers vs. sleep misperceptors), stage (NREM vs. REM) and/or time (hours after lights off) as fixed factors, and subject and time spent asleep as random factors (#1, #2).

As expected, sleep misperceptors felt more often awake than good sleepers *[main effect of group, $\chi^2(1) = 11.0$, p=0.001, **Figure 1A**, estimates and post-hoc results in **Table S.1B-C**]*. Both groups felt more often awake in NREM than REM sleep *[main effect of stage: $\chi^2(1) = 7.7$, p=0.005, **Figure 1A, Table S.1B-C**, no group-stage interaction, NREM: 12.4% and 41.5% in good sleepers and sleep misperceptors respectively, REM: 0.5% and 9.0%; $\chi^2(1) = 0.28$, p=0.59, **Table S.1A**]*. In both groups, within NREM sleep, the proportion of FAW changed significantly in the course of the night *[main effect of time of night: $\chi^2(7) = 48.1$, p<0.0001, **Figure 1B**, no group-time interaction effect, **Table S.2A**]*, with most FAW instances occurring in the first two, followed by the last hour of the night (**Table 2B., Table S.2B-C**).

We then evaluated, for all FAS instances, how perceived sleep depth differed between groups, sleep stage and time of night. Here, we performed a linear mixed model, with perceived sleep depth (from 1 to 5, 5=deepest) as the ordinal dependent variable, and the same fixed and random factors as in the model described above (#3). Good sleepers felt more deeply asleep in REM than NREM sleep, while sleep misperceptors reported equivalent subjective sleep depths between REM and NREM sleep *[significant group - sleep stage interaction effect: $\chi^2(1)$ 4.2, p=0.041, **Table S.3B-C**]*. Similarly to the proportion of FAW, subjective sleep depth within NREM sleep significantly changed in the course of the night – with subjects feeling less deeply asleep in the first two hours, followed by the last hour of the night [*main effect of time, $\chi^2(7) = 59.1$, p <0.0001, interaction effect between time and stage $\chi^2(6)=13.2$, p=0.04, **Figure 1D**, **Table S.3B-C**].

The same analyses including stages N2 and N3 separately showed that perceived state and sleep depth were comparable between N2 and N3, but differed from REM sleep in both cases *[main effect of group: perceived state: $\chi^2(1)=16.82$, p<0.0001 and of stage: perceived state: $\chi^2(1)=12.37$, p=0.002; perceived sleep depth: $\chi^2(1)=9.32$, p=0.01, **Table S.4B-C, Figure S.1**].* For further analyses, N2 and N3 were therefore analyzed together (as NREM).

Figure 1. Determinants of sleep perception. Perceived state (feeling awake vs feeling asleep; A, B) and degree of perceived sleep depth (scale from 1 to 5; C, D) as a function of group (good sleepers/misperceptors), stage (REM/NREM) and time of the night. **A.** Proportion of feeling awake instances in good sleepers (n=20, grey) and misperceptors (n=10, black) in NREM and REM sleep. **B.** Average proportion of feeling awake (red) and feeling asleep instances (blue) +/- standard error across trials in good sleepers (upper row) and misperceptors (bottom row) in NREM sleep in the course of the night. **C.** Degree of perceived sleep depth in good sleepers (grey) and misperceptors (black) in NREM and REM sleep. **D.** Average degree of perceived sleep depth +/- standard error across trials in good sleepers (green) and misperceptors (orange) in NREM (solid line) and REM (dashed line) in the course of the night. In boxplot figures (A, C), each point represents an individual, the box displays the 25th to 75th percentile of data and the vertical bars shows the confidential interval. Post-hoc of significant main and interaction effects evaluated with generalized linear mixed models are indicated with lines and asterisks indicating the level of significance: *$p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, including probability of FAW/FAS as a function of group and stage (A); group and time of night (B). Degree of perceived sleep depth as a function of group, stage and time of night (C, D). All models included subject identity and time spent asleep as random

Perceived sleep depth and conscious experiences

Next, we investigated how conscious experiences related to subjective sleep perception. Reports of conscious experiences (CE) were considered separately from instances in which participants reported conscious experiences but could not report the content (herein 'conscious experiences without report of content'), and those of no experience (NE) [12]. As expected [36] both groups were more likely to report a CE in REM than NREM sleep (#4) [*main effect of stage, $\chi^2(1) = 12.6$; p=0.0004,* **Figure 2.C, Table S.13B-C***, no effect of group; no interaction effect between group and stage*]. In both groups, experiences reported upon awakening from REM sleep were more perceptual, richer, longer and associated with lower degrees of lucidity compared to NREM experiences (#5) [*significant main effect of stage on perception $\chi^2(1) = 32.01$; p<0.0001,* **Table S.15B-C***; richness & complexity $\chi^2(1) = 10.58$, p=0.001,* **Table S.16B-C***; duration $\chi^2(1) = 28.23$, p<0.0001,* **Table S.17B-C***; lucidity $\chi^2(1) =5.33$, p=0.02, ,* **Table S.19B-C; Figure 2.D**]. Good sleepers and sleep misperceptors differed only in the thinking dimension, with sleep misperceptors reporting significantly more thought-like experiences compared to good sleepers [*main effect of group, $\chi^2(1) = 9.01$, p=0.003,* **Table S.14B-C**].

We then evaluated whether perceived state was influenced by the presence (CE) or absence (NE) of conscious experiences (C.Exp, #6). As FAW instances were very rare in REM sleep, this analysis was performed only for NREM sleep. Suprisingly, in both groups, FAW instances were more likely to be associated with NE than CE [*main effect of C.Exp, $\chi^2(1)=5.54$, p=0.018,* **Figure 2.A, Table S.6B-C***, no group – C.Exp interaction effect*]. We then evaluated how the characteristics of reported conscious experiences differed depending on whether participants felt awake or asleep in NREM sleep (#7). In both good sleepers and sleep misperceptors, FAW experiences were more thought-like and associated with a higher degree of control over the experience compared to FAS experiences [*main effect of perceived state on thinking $\chi^2(1) = 8.71$, p=0.003,* **Table S.7B-C** *and voluntary control $\chi^2(1) =7.68$, p=0.006,* **Table S.11B-C, Figure 2.B**]. Sleep misperceptors displayed more thought-like experiences than good sleepers irrespective of perceived state [*main effect of group, $\chi^2(1) = 11.75$, p=0.0006,* **Figure 2.B***, no group - perceived state interaction effect,* **Table S.7B-C**)]. When feeling awake, sleep misperceptors reported richer and more complex experiences compared to when feeling asleep, but no such difference was seen for good sleepers [*group - state interaction effect, $\chi^2(1) = 4.59$, p=0.03,* **Table S.9D**]. Examples of experiences associated with FAS and FAW reports can be found in Annex S.1.

No relation was found between the degree of subjective sleep depth and the presence or absence of conscious experiences (#8) *[main effect of C.Exp, $\chi^2(1) = 2.66$, p>0.05; no interaction effect between C.Exp and group, $\chi^2(2) = 2.06$, p>0.05 ; no interaction C.Exp and stage effect]*. The more subjects felt asleep, the richer, more complex and perceptual the experience *[main effect of perception, $\chi^2(4)=19.8$, p=0.0006, **Table S.22B-C**; and richness & complexity, $\chi^2(4)=17.86$, p=0.001, **Table S.23B-C**; no interaction effects]*.

Figure 2. Conscious experiences in sleep and sleep perception. A. Probability of presence (CE) and absence (NE) of conscious experience. **B.** Average scores for different scales evaluating qualitative aspects of experiences, shown for good sleepers (grey) and misperceptors (black), in NREM and REM sleep. **C.** Proportion of feeling awake instances in presence and absence of conscious experiences **D.** Average scores for different scales evaluating qualitative aspects of experiences when FAW and FAS. In all figures, each point represents an individual, the box displays the interquartile range and the vertical bars shows the confidential interval. Post-hoc of significant main and interaction effects evaluated with generalized linear mixed models are indicated with lines and asterisks indicating the level of significance: * *p < 0.05,* ** *p < 0.01,* *** *p < 0.001.* (A) probability of CE/NE and (B) CE content score as a function of group and stage; (C) probability of FAW as a function of group and C.Exp. and (D) CE content score as a function of group and FAW/FAS. All models included subject identity and time spent asleep as random factors.

EEG activity and perceived sleep depth in NREM sleep

To determine how absolute EEG spectral power in different frequency bands related to perceived sleep depth in NREM sleep, we computed a linear mixed model per frequency band and electrode, including subject identity and time of night as random factors (#10). We found that sigma power (12 16 Hz) in good sleepers and beta power (18-30 Hz) in both groups significantly predicted lower subjective sleep depth (**Figure 3A, first two rows**). No interaction effect between power and groups was found (**Figure 3A, last row**). To map the topography of the effects in the sigma and beta power range on sleep depth, we performed the same analysis on source reconstructed data (**Figure 3B**). Here we found a significant effect for sigma in good sleepers and beta in sleep misperceptors on subjective sleep depth. In both cases, the effects of sigma and beta power were spatially widespread, peaking in central brain regions, and sparing only small portions of the frontopolar, inferior and medial temporo-occipital cortices. For beta power, sleep misperceptors displayed more widespread effects compared to good sleepers, extending to fronto-polar and occipital areas, although this difference did not resist our stringent correction for multiple comparisons.

As our findings indicated that within NREM, subjective sleep depth decreased after the first two hours, then increased in the very last hour in both groups, we investigated how spectral power changed across the night (***Figure S2***). All frequency bands showed significant changes across the night in NREM sleep in both good sleepers and sleep misperceptors. Despite the serial awakenings, delta power decreased across the night in both groups, as would be expected for an undisturbed night of sleep. Beta power across the night inversely mirrored subjective sleep depth in both groups, decreasing early in the night in both groups and increasing towards the end of the night, consistent with our findings relating beta power to lower subjective sleep depth.

Since sigma power is known to reflect sleep spindle activity and predicted lower sleep depth in both populations, we performed a spindle detection analysis and examined how different spindle parameters related to sleep depth (#11). We found that the amplitude, but not the density of spindles predicted a lower perceived sleep depth (**Figure 7**). In good sleepers, fast spindle density also predicted lower sleep depth. Similar to sigma power findings, these effects were spatially widespread.

NREM

Figure 3. Absolute power spectral density and perceived sleep depth in NREM sleep. A. Results of linear mixed models explaining perceived sleep depth by power spectral density in good sleepers (n=20, top row) and misperceptors (n=10; middle row). Group interaction effect is displayed in bottom row. Absolute power spectral density was averaged over the 120s preceding the awakening. The effect of each frequency band was evaluated in separate models. All models included subject identity and time of night as random factors. Wald statistics values (squared ratio of the fixed factor estimate over its standard error) are shown at the scalp level for each electrode; electrodes with a significant effect after a cluster-and probability-based correction for multiple comparisons (p<0.05) are marked white. **B.** Cortical distribution of Wald statistics at the source level for selected frequency bands (based on results shown in A). The same procedure described in A was used. Voxels with non-significant results are colored in grey. Contrasts that did not survive the correction for multiple comparisons are marked with a hash sign. *LL = left lateral, RL = right lateral, LM = left medial, RM = right medial.*

Sleep spindles (NREM)

Figure 4. Spindles and perceived sleep depth in NREM sleep. Results of linear mixed models explaining perceived sleep depth by spindle parameters in good sleepers (n=20, top row), misperceptors (middle row), and group interaction effect (bottom row). Spindle parameters were extracted and averaged over the 120s preceding the awakening. The effect of each parameter was evaluated in separate models. All models included subject identity and time of night as random factors. Wald statistics values (squared ratio of the fixed factor estimate over its standard error) are shown at the scalp level for each electrode; electrodes with a significant effect after a cluster-and probability-based correction for multiple comparisons ($p<0.05$) are marked white.

EEG activity and perceived sleep depth in REM sleep

In REM sleep, beta (18-29.5 Hz) and gamma power (30-45 Hz) significantly predicted lower subjective sleep depth in sleep misperceptors (**Figure 5A**). A non-significant negative correlation was also seen for all other frequency bands in sleep misperceptors. In good sleepers, no significant effects were found, although a weak and non-significant tendency for alpha, sigma and beta power to predict lower sleep depth was seen. Models explaining perceived sleep depth with source reconstructed EEG activity showed again a widespread effect of beta and gamma on perceived sleep depth in sleep misperceptors, sparing only small islands of the lateral temporal and fronto-polar cortices (**Figure 5B**). As expected, corrected group interaction effects suggested that the predictive power of beta and gamma activity was significantly higher in sleep misperceptors than good sleepers (**Figure 5B**, last column).

Since subjective sleep depth in good sleepers was lower in NREM sleep compared to REM sleep, a sleep state in which the EEG is traditionally described as 'fast' and 'wake-like', we directly compared spectral power between REM and NREM sleep (***Figure S3***). As expected, low-

frequency power (1-16 Hz) was lower in REM sleep compared to NREM sleep. However, high-frequency power in the beta (18-29.5 Hz) and gamma range (30-45 Hz) was not significantly different between REM and NREM sleep, and in some frontal electrodes was even higher in REM sleep. These findings indicate that REM sleep is 'wake-like' in terms of low low-frequency power but does not necessarily display higher high-frequency power values.

REM

Figure 5. Power spectral density and perceived sleep depth in REM sleep. A. Results of linear mixed models explaining perceived sleep depth by power spectral density in good sleepers (n=20, top row) and misperceptors (middle row). Group interaction effect is displayed in bottom row. Power spectral density was averaged over the 120s preceding the awakening. The effect of each frequency band was evaluated in separate models. All models included subject identity and time of night as random factors. Wald statistics values (squared ratio of the fixed factor estimate over its standard error) are shown at the scalp level for each electrode and electrodes with a significant effect after a cluster-and probability-based correction for multiple comparisons (p<0.05) are marked white. **B.** Cortical distribution of Wald statistics at the source level for selected frequency bands (based on results shown in A). The same procedure described in A was used. Voxels with non-significant results are colored in grey. Contrasts that did not survive the correction for multiple comparisons are marked with a hash sign. *LL = left lateral, RL = right lateral, LM = left medial, RM = right medial.*

Discussion

Sleep perception in good sleepers

Both the feeling of being asleep as well as subjective sleep depth (1-5 scale) showed considerable variability within the same night of sleep. Good sleepers reported feeling awake despite full-fledged sleep in ~ 10% of probes, most frequently so during the first two hours of NREM sleep, and again towards the end of the night, during the last hour of sleep. Perceived sleep depth mirrored this course. This observation suggests that the feeling of being asleep sets in progressively, sometime after objective sleep onset, and fades out gradually, before the re-establishment of full wakefulness. The observation that early NREM sleep was associated with the highest proportion of feeling awake instances and among the lowest sleep depth ratings is surprising, since early NREM sleep is generally considered the "deepest" sleep, with the highest amount of slow wave activity. Even when accounting for the time of the night, we did not find a relation between subjective sleep depth and slow wave activity (SWA). This cannot be ascribed to the serial awakening paradigm we used, since in our dataset, SWA decreased progressively across the night (**Figure S.3**), similar to an undisturbed night of sleep, and displayed the highest values during the first two hours of sleep. These results therefore suggest that SWA does not mediate subjective sleep depth. The fact that subjective sleep depth was highest in the middle of the night, near the circadian nadir, could indicate that circadian factors, instead of homeostatic sleep factors (reflected in SWA) may play a role in subjective sleep depth [37,38]. Another argument suggesting that the presence of high SWA may not be necessary for feeling (deeply) asleep comes from the observation that REM sleep, in which SWA is reduced and highly localized [39–41], featured the highest sleep depth ratings and only 1 out of 126 FAW instances. Since we controlled for the time spent asleep before the awakening, the high subjective sleep depth in this stage cannot be merely a consequence of having spent more time asleep in this stage than in NREM sleep before being awakened.

While we did not find a relation between subjective sleep depth and SWA, we documented a negative relation between high-frequency power in the sigma and beta range and sleep depth in NREM sleep in good sleepers. Consistent with these findings, the course of beta power across the night followed a U-shaped course, inversely mirroring subjective sleep depth (**Figure S.3**). Just as it is surprising to note that subjective sleep depth is lowest in early NREM sleep, it is puzzling to notice that NREM sleep displays equal or even higher high-frequency (beta and gamma) power in some frontal regions compared to REM sleep (**Figure S.4**). The EEG in REM sleep is typically

described as fast and wake-like, but systematic comparisons in spectral power across behavioral states and within subjects are hardly ever reported.

Since sigma power reflects spindle activity, we performed a spindle analysis, which revealed that spindle amplitude and fast spindle density predicted lower sleep depth in good sleepers. These findings are in line with a previous study, showing faster sleep spindles in insomnia patients compared to controls [23]. Intraspindle frequency has been suggested to reflect levels of thalamic polarization within a relatively narrow range [42]. Spindle frequency typically decreases in the transition to sleep and fast spindles tend to occur during slow wave up-states during which thalamocortical neurons are depolarized [43,44]. It is thus possible that relative thalamo-cortical depolarization, closer to the 'wake state', which is reflected in fast spindle density, may account for subjectively lighter sleep. It is however unlikely that our spectral power findings are explained by changes in spindle frequency and amplitude alone. Although only sigma and beta levels were significantly related to lower sleep depth (Fig 3.A), similar non-significant results were obtained for a broad range of frequencies from alpha upwards. In addition, centro-parietal electrodes, in which fast spindle density is usually highest [43] displayed the weakest relationship with sleep depth compared to other electrodes. Overall, these results suggest that high-frequency EEG activity, including but not limited to the spindle range, predicts lower sleep depth. Abrupt increases in EEG activity in the alpha, beta and gamma band are the defining features of micro-arousals of sleep, which can occur either spontaneously or in response to sensory stimuli, and can be associated with slow waves. It is therefore possible that the high-frequency increases that predict lower sleep depth reflect the (possibly subthreshold) activation of arousal systems. The spatially diffuse distribution of sigma and beta power changes are consistent with the widespread projections of arousal systems [45]. In good sleepers, EEG power did not significantly predict subjective sleep depth in REM sleep, possibly because of the overall high ratings of subjective sleep depth in this stage and group.

In line with previous studies, we found that experiences reported upon awakening from REM sleep were more perceptual, richer, longer and associated with lower degrees of lucidity compared to NREM experiences [12,46,47]. Feeling awake in NREM sleep was associated with a lower proportion of conscious experiences compared to FAS, suggesting that although subjects felt awake, they did at times, not experience anything. This is somehow surprising, as one could expect that feeling awake would necessarily be associated with conscious experiences, similar to the waking state [48]. Yet recent studies probing the contents of mental experiences during wakefulness found an unusually high proportion of mind blanking (no experience) in this state [49],

in relation to sleep-like slow waves. The incidence of such mind blanking periods can be expected to occur with even greater frequency while one is asleep, even if one feels awake. In addition, mental experiences in wakefulness are typically more discontinuous than full-fledged 'dreams', possibly because during wakefulness, attention is continuously redirected towards environmental events, which is not the case during sleep, when one is disconnected from the environment. It is possible that subjects who feel awake during sleep experience such interruptions, which leads them to answer that they did not experience anything before the alarm sound. Alternatively, such instances could represent a « dreamless sleep experience, or the persistence of temporal experience devoid of further intentional content or any specific objects of awareness during sleep » [50].

When experiences were reported in association with FAW instances in NREM sleep, they were associated with more thought-like content, over which participants had greater control compared to FAS instances. The richer, the more complex and more perceptual the experience, the more participants felt asleep. These reports are perfectly in line with previous studies showing that thinking and reflective consciousness (comprising lucidity and voluntary control) are typical features of wake mentation, while sleep experiences, especially REM sleep dreams, are highly perceptual [12,13,15]. In addition, the disappearance of control over thoughts has been shown to relate to the perception of having fallen asleep [51]. Thus, ratings of conscious experiences during FAW instances are in line with mental activity during wakefulness, even though participants were asleep.

We previously showed that perceptual aspects of dreams correlate with high-frequency power in posterior brain regions, while thought-like experiences correlate with high-frequency power in anterior brain regions [34]. Here we showed that high-frequency EEG activity predicted lower sleep depth in almost all cortical regions except for parts of the occipital lobe, involved in the perceptual aspects of dreams. It is thus possible that relative frontal "deactivation" [52], brought about by quiescence of arousal systems, allows for highly perceptual experiences to emerge and thereby accounts for the feeling of being deeply asleep.

Sleep perception in sleep misperceptors

As expected [19], sleep misperceptors felt more frequently awake during sleep than good sleepers (roughly three times as often) and perceived their sleep as shallower. The distribution of FAW instances across sleep stages was similar between the two groups, and subjective sleep depth in NREM followed a comparable course (no interaction effects). These findings suggest that feeling

awake could be mediated by the same neurophysiological mechanisms in NREM sleep, which appear enhanced in sleep misperceptors, possibly as a consequence of hyperarousal (suggested by metabolic studies in insomnia patients [53,54]). Unlike good sleepers however, who felt most asleep in REM sleep, sleep misperceptors did not display differences in subjective sleep depth between REM and NREM sleep, suggesting that the neurophysiological mechanisms that mediate the feeling of being awake abnormally persist into REM sleep only in sleep misperceptors. These findings are consistent with a previous serial awakening study showing that patients with insomnia felt more frequently awake than control subjects in REM sleep, but not in N2 sleep [16].

Some studies have shown that REM sleep fragmentation by arousals is a prominent feature of insomnia that contributes to the overestimation of wake time during the night [55] and interferes with the overnight resolution of emotional distress [56]. In addition, REM sleep fragmentation in insomnia sufferers has been related to thought-like nocturnal mentation [56]. In our dataset, insomnia sufferers displayed more thought-like nocturnal mentation compared to controls, irrespective of sleep stage. Interestingly, unlike good sleepers, insomnia sufferers described their mental content as richer and more complex when they felt awake as opposed to when they felt asleep, which may reflect the ruminations (repetitive thoughts) that these patients experience when trying to fall asleep. In sleep misperceptors high-frequency EEG changes predicting sleep depth were more spatially widespread compared to good sleepers, in, as they involved additional fronto-polar and posterior brain areas, suggesting a more extensive recrutement of arousal systems in this group.

We previously showed that this group of sleep misperceptors had a lower proportion of slow wave sleep in an undisturbed baseline night compared to control subjects [4], a finding that is in line with several previous studies [24,25,57]. Although sleep misperceptors had overall lower delta power also in this dataset compared to controls, delta power did not predict higher sleep depth. It is thus conceivable that increased activity of arousal systems, which mediate the feeling of being awake, may prevent subjects to enter or spend a long time in N3 sleep, without necessarily correlating with reduced SWA.

Overall interpretation

Our results suggest that subjective sleep depth is inversely related to a neurophysiological process that predominates in NREM sleep early in the night, becomes silent in REM sleep, is reflected in high-frequency EEG activity and associated with highly thought-like conscious experiences. Our

results also suggest that this neurophysiological process is amplified in sleep misperceptors: it comes into play more frequently and more strongly (more FAW instances, lower sleep depth), recruits more brain regions (spatially more diffuse changes) and abnormally persists into REM sleep in this group (no REM-NREM difference in sleep perception). A positive relationship between subjective sleep depth and traditional hallmarks of sleep was not found (sleep spindles, slow wave activity). Thus, it is not the presence of sleep rhythms but rather the absence of 'wake-like' activity (high-frequency power) that appears to account for feeling asleep.

Our data does not provide direct information regarding the origin of the underlying neurophysiological processes, but several of our findings suggest that the noradrenergic arousal system could be a potential mediator of feeling awake during sleep and of low subjective sleep depth. When averaged within behavioral states, noradrenaline levels and locus coeruleus (LC) activity are highest in wakefulness, intermediate in NREM sleep, and lowest during REM sleep, paralleling the incidence of feeling awake instances [58,59]. Although there are many other neuromodulator systems that show similar profiles across behavioral states (histaminergic, orexinergic, serotoninergic, etc.), several observations make noradrenaline a better candidate. First, despite overall low LC activity levels in sleep, several studies have shown that the LC discharges intermittently during NREM sleep, time-locked to slow wave up-states and spindles [60,61]. Second, while the LC may promote EEG activation and arousal, some studies have documented transient increases in LC activity (to wake levels) during full-fledged slow wave sleep at the beginning of the night, when FAW preferentially occurred and subjective sleep depth was lowest in our study [62–64]. These findings suggest that LC activity can be high even during full-fledged slow wave sleep and that these dissociations may account for feeling awake in deep sleep. Another dissociation between SWA and noradrenaline has been documented in fur seals, in which cortical and subcortical noradrenaline levels, unlike EEG activity, do not show lateralization during uni-hemispheric slow wave sleep [64]. Finally, our source analysis results revealed that high-frequency power changes related to sleep depth were spatially widespread, consistent with the diffuse cortical projections of the noradrenergic system. They tended to spare parts of the temporal and occipital cortices, especially in good sleepers, which is in line with studies showing that the occipital cortex is one of the regions with the lowest levels of cortical noradrenaline [65,66]. While noradrenergic activity has been linked to the memory benefits of NREM sleep in good sleepers [67], its excessive activity has been suggested to disrupt REM sleep in insomnia patients and to lead to dysfunctional emotion regulation [68]. Future studies should try to determine whether the

suppression of LC activity, for instance with alpha-2 agonists, leads to an increase in subjective sleep depth in insomnia patients [68].

Conclusions and perspectives

Our results challenge the commonly held notion that slow wave sleep, generally referred to as deep sleep, best accounts for feeling deeply asleep. Instead, our findings suggest that the subjective perception of sleep is more closely related to the absence of wake-like high-frequency EEG activity: the lower it is, the more one feels asleep. Our results also indicate that the feeling of being deeply asleep is incompatible with highly thought-like mental activity and instead associated with perceptual, dream-like experiences, which most often occur in REM sleep.

In light of these and other recent findings, one should reconsider the use of the term 'deep' sleep to designate slow wave sleep. Indeed, NREM sleep is more heterogeneous than previously thought. It comprises alternating phases of sleep stability and fragility and is associated with intermittent noradrenergic surges [60,61,69–71]. Once considered largely dreamless, it is now known to be associated with mental activity in up to 43% of cases [36]. Conversely, sleep inertia and high arousal thresholds, traditionally regarded as a hallmark of slow wave sleep, can be seen to a similar extent in REM sleep [72,73] and according to some studies[76], REM sleep appears to better account for subjective sleep quality than slow wave sleep. Not surprisingly, some early sleep electrophysiologists observing cats in different behavioral states chose the term 'deep sleep' to refer to REM sleep instead of slow wave sleep [74,75]. Continuing to use the term 'deep sleep' for slow wave sleep could also be misleading for future studies, which may focus exclusively on slow wave enhancement to improve subjective aspects of sleep, and neglect REM sleep. Finally, the existence of an objective EEG correlate of subjective sleep depth, which is present in both good sleepers and insomnia patients to a different degree, strongly argues against purely psychological factors that have been suggested to underlie the sleep underestimation in insomnia. Instead, our findings suggest that so-called 'misperceptors' are not truly 'misperceptors', but instead may accurately perceive subtle shifts towards 'wake-like' activity during sleep, which are not not visible in standard sleep recordings.

Methods

Selection of participants

Good sleepers (n=20, age 38,3 ± 7,4 yrs (mean±SD), range: 25 – 51, 15 females) were recruited through advertisement and word by mouth. They had to have regular bed and rise times and a good subjective sleep quality (Pittsburgh Sleep Quality Index[77] <5). Subjects with extreme chronotypes (Horne and Ostberg morningness-eveningness questionnaire[78] scores >70 or <30), excessive daytime sleepiness (Epworth Sleepiness Scale[79] >10), suffering from neurological, psychiatric or medical disorders affecting sleep, taking regular medication besides birth control and pregnant females were not included in the study.

Sleep misperceptors of similar age and gender distribution (n=10, age 40,8 ± 6,5 yrs, range: 29 – 49, 8 females, t-test p=0.4) were recruited among outpatients of the Center for Investigation and Research in Sleep at the Lausanne University Hospital. They had to have a diagnosis of chronic insomnia fulfilling the criteria of the international classification of sleep disorders [80]. Only patients who had had a polysomnography for clinical reasons were included, which had to fulfil the following criteria: 1) normal sleep efficiency (>85%), 2) sleep latency <60 min. As a routine clinical procedure, immediately after the clinical polysomnography, patients had to estimate how much time they had spent asleep during the night. Only subjects who presented a mismatch between subjective and objective total sleep time (TST) > than 60% were included. The mismatch was evaluated by means of the Sleep Perception Index (SPI), which expresses the ratio between subjective (sTST) and objective TST (oTST) in percent ([sTST/oTST] *100). The cutoff of <60% was chosen based on a previous study, in which a sample of the general population (n=2092) underwent polysomnography and estimated TST the next day. In this population, an SPI lower than 58.82% defined the lowest 2.5% of the population distribution. The sample of the general population was comparable, in terms of subjective complaints, PSG and spectral EEG parameters, to the population of insomnia sufferers of this study (for a comparison, see [4]).

Exclusion criteria for patients comprised major psychiatric and neurological comorbidities, medication other than birth control, pregnancy, apnea-hypopnea index >15/h and periodic leg movement index in sleep >15/h. Written informed consent was obtained by all the participants and the study was approved by the local ethical committee (commission cantonale éthique de la recherche sur l'être humain du canton de Vaud).

Experimental procedure

Both sleep misperceptors and good sleepers underwent an undisturbed baseline hd-EEG sleep recording with 256 electrodes in the sleep laboratory, starting at approximately 11:30 pm and ending at 6:30 am the next morning. Sleep architecture and EEG findings of this baseline night have been published separately [4]. In brief, sleep parameters in insomnia sufferers did not differ significantly from controls, except for a significantly lower proportion of stage N3 sleep [4]. Participants subsequently underwent two experimental hd-sleep EEG recordings, in which a a previously validated serial awakening paradigm was employed [12]. The two experimental nights were scheduled at least one week apart to limit the effects of the experimental sleep fragmentation on dream recall. One healthy participant completed only one experimental night because of organizational constraints. Subjects were asked to maintain a regular sleep-wake schedule during the week before the recordings, and compliance was verified with wrist-worn actigraphy. Subjects did not have access to watches or phones during the recordings.

Awakenings were carried out with a 1.5s computerized alarm sound that was played through intercom by the examiner who continuously monitored the subjects via EEG and video/audio from a control room [12]. NREM awakenings were performed approximately 10 min after the first sleep spindle, while rapid eye movement (REM) sleep awakenings were carried out 10 min after the beginning of REM sleep.

After each awakening, participants were asked to estimate whether they had been feeling awake or asleep before the alarm sound. Next, they had to estimate, on a 1-5 scale, how deeply they had felt asleep before before the alarm sound (depending on the answer to the previous question). They also had to describe what they had in mind immediately before the alarm sound ("What was going through your mind immediately before the alarm sound?"). When an experience was reported, participants had to estimate the following dimensions on a 1 to 5 point scale: 1) the degree of lucidity ("How much did you realize that the experience was not happening in reality?"), 2) degree of voluntary control over the experience ("How much did you have control over the content of the experience?"), 3) the thought-like character of the experience ("How much did it resemble a thought or reflection ?"), 4) the perceptual character of the experience ("How much was it perceptual?"), 5) the length of the experience ("How long was the experience ?") as well as 6) the richness and complexity of the experience ("How rich and complex was the experience ?"). Once the interview was finished, subjects were allowed to fall back to sleep.

Sleep scoring and EEG preprocessing

A 256-channel system (Electrical Geodesics, Inc., Eugene, Oregon) was used for EEG acquisition, with a sampling rate of 500 Hz. Four of the 256 electrodes near the eyes were used to monitor eye movements, electrodes overlying the masseter muscles and close to the chin were used to monitor muscle tone [34]. The EEG signal was off-line band-pass filtered between 0.5 and 45 Hz, and sleep scoring was performed over 30s epochs according to standard criteria [81].

The two minutes of data before each awakening were extracted for analysis. Channels containing artefactual activity were visually identified and replaced by interpolation from nearby channels using spherical splines (NetStation, Electrical Geodesic Inc.). Ocular, muscular, and cardiac artefacts were eliminated with Independent Component Analysis (ICA) using EEGLAB routines [82]. ICA components with specific activity patterns and component maps characteristic of artefactual activity were removed [83].

To determine the time spent asleep before each awakening, all the recordings were visually inspected. Wake periods were scored as at least 15s of clear-cut EEG wakefulness [81]. The time between the end of the last period scored as wake and the alarm sound was extracted for each awakening.

EEG analysis

Power spectral densities (PSD) were calculated using the Fast Fourier transform method on artifact-free consecutive, non-overlapping 2s epochs and used to compute signal power in typical frequency bands, including delta (1–4 Hz), theta (5–7.5 Hz), alpha (8–11.5 Hz), sigma (12–16 Hz), beta (18–29.5 Hz) and gamma (30 Hz – 45 Hz). In order to avoid overlap between frequency bands, we excluded the upper 0.5 Hz range of neighbouring bands.

Source localization was performed using a previously described procedure [34]. In brief, the 120s sleep period prior to each awakening was extracted from the already cleaned and filtered. Source modeling was performed using the GeoSource software (Electrical Geodesics, Inc., Eugene, Oregon). Individualized geocoordinates of the electrodes were used to construct the forward model. The source space was restricted to 2,447 dipoles that were distributed over 7×7×7-mm cortical voxels. The inverse matrix was computed using the standardized low resolution brain electromagnetic tomography (sLORETA) constraint [84]. A Tikhonov regularization (λ=10-2) procedure was applied to account for the variability in the signal-to noise ratio [84].

Spindle detection was based on an automatic algorithm, adapted from a previous study [85]. The EEG signal was average-referenced, downsampled to 128 Hz and bandpass-filtered between 11 and 16 Hz (~3 dB at 10 and 17 Hz). The following spindle parameters were extracted: density, density of low and high frequency, maximal amplitude, duration and frequency. Based on previous studies [42,43,86], we categorized spindles as fast or slow based on an individualized threshold, which was defined as the intermediate value between the average spindle frequency in one centroparietal channel (Pz) and one frontal channel (Fz) [87].

Statistical analysis

Statistical analyses were performed in R-Studio (version 1.1.463). In the following section and in the result section, reference numbers preceded by hash signs refer to the statistical models listed in Table 1.

Generalized mixed models were computed to evaluate the predictors of perceived state (FAW vs FAS, categorical dependent variable), including group and stage (#1, Fig.1A), group and time in NREM (#2, Fig.1B), and group and conscious experience presence and recall (#6, Fig.2A). Linear mixed models were computed to evaluate the predictors of perceived sleep depth (Degree, ordinal dependent variable), including group, stage and time (#3, Fig.1C and Fig.1D), group, stage and C.Exp (#8) and group, stage and dream content characteristics (#9). Linear mixed models were computed to evaluate the predictors of dream content characteristics (listed in the experimental procedure section), including group and sleep stage in FAS (#5, Fig.2B) and group and perceiveed state in NREM (#7, Fig.2D). A generalized mixed model was computed to evaluate the effect of group and stage of night on presence or absence of conscious experience (CE_presence, categorical dependent variable, #4, Fig.2C). Subject identity and time spent asleep before each awakening was included as a random factor in all analyses. Post-hoc contrasts were computed with estimated marginal means (emmeans package) and a Tukey's HSD (honestly significant difference) test was applied to adjust for the risk of false discoveries (type I error) consequent to multiple comparisons. In the results section, all main statistics and associated p-values are reported in the text, post-hoc statistics are displayed in the figures and estimates and post-hoc values are reported in supplementary tables referenced in the text.

For EEG data analyses, linear mixed models were computed for each electrode (at the scalp level) or voxel (at the source level) to evaluate the effect on perceived sleep depth of absolute power spectral density in different frequency bands (#10, Fig.3 and 4) and of different spindle parameters

(#11, Fig.6). For models #10, one model per frequency band was performed. For models #11, one model per parameter was performed. Subject identity and time of night was included as a random factor in all the analyses, and frequency bands as fixed factors. As between-subject variability is accounted for by including subject identity as random factor, all EEG power spectral density models were performed on absolute (non-noramlized) power values. For each model, we obtained the Wald statistics, which is the squared ratio of the fixed factor estimate over its standard error. To account for multiple comparisons, a cluster and probability-based correction was used. In order to create a dummy population, labels corresponding to the dependent variables were shuffled within subjects 1000 times. For each permutation, the model was applied and neighboring electrodes with p-values inferior to 0.05 were identified as a cluster. A cluster statistics was produced by summing Wald statistics inside each significant cluster and by dividing by the number of electrodes in the cluster, resulting in one value for each identified cluster. For each permutation, only the absolute maximal value of all cluster statistics was retained. A threshold for significance was set at the 95th percentile of the dummy cluster statistics distribution. The cluster statistics obtained in the real dataset were considered significant when above this threshold. This procedure was applied for each electrode/voxel.

	Formula	Details	Figure
#1	Perceived State ~ Group*Stage + (1|sub) + (1|time_asleep)	Stage = NREM vs REM	Fig.1A
#2	Perceived State ~ Group*Time + (1|sub) + (1|time_asleep)	In NREM only as FAW are quasi inexistent in REM	Fig.1B
#3	Degree ~ Group*Stage*Time + (1|sub) + (1|time_asleep)		Fig.1C-D
#4	CE_presence ~ Group*Stage	CE_presence = CE vs NE, FAS only	Fig.2A
#5	DC ~ Group*Stage + (1|sub) + (1|time_asleep)	One model per DC, FAS only	Fig.2B
#6	Perceived State ~ Group *CE + (1|sub) + (1|time_asleep)	NREM only	Fig.2C
#7	DC ~ Group*Perceived State + (1|sub) + (1|time_asleep)	One mode per DC, NREM only	Fig.2D
#8	Degree ~ Group*Stage*CE + (1|sub) + (1|time_asleep)	One mode per DC, FAS only	
#9	Degree ~ Group*Stage*DC + (1|sub) + (1|time_asleep)	One mode per DC, FAS only	
#10	Degree ~ PSD + (1|sub) + (1|time)	One model per frequency band in NREM and REM separately	Fig.3 Fig.5
#11	Degree ~ SP + (1|sub) + (1|time)	One model per spindle parameter in NREM	Fig.4

Table 1. List of generalized and linear mixed models. All models referred to in the result section are summarized in this table, with specifications and reference to the figure in which their result are displayed. DC = Dream Content = Thought, Perception, Rich, Long, Control, Consciousness ; Event = FAW vs FAS ; time = time since lights off; time_asleep = time since end of occipital alpha.

References

1.   Kaplan, K. A., Hirshman, J., Hernandez, B., Stefanick, M. L., Hoffman, A. R., Redline, S., Ancoli-Israel, S., Stone, K., Friedman, L. & Zeitzer, J. M. When a gold standard isn't so golden: Lack of prediction of subjective sleep quality from sleep polysomnography. *Biol. Psychol.* **123**, 37–46 (2017).

2.   Castelnovo, A., Ferri, R., Punjabi, N. M., Castronovo, V., Garbazza, C., Zucconi, M., Ferini-Strambi, L. & Manconi, M. The paradox of paradoxical insomnia: A theoretical review towards a unifying evidence-based definition. *Sleep Med. Rev.* **44**, 70–82 (2019).

3.   Bonnet, M. H. & Moore, S. E. The threshold of sleep: Perception of sleep as a function of time asleep and auditory threshold. *Sleep* **5**, 267–276 (1982).

4.   Lecci, S., Cataldi, J., Betta, M., Bernardi, G., Heinzer, R. & Siclari, F. Electroencephalographic changes associated with subjective under- and overestimation of sleep duration. *Sleep* (2020) doi:10.1093/sleep/zsaa094.

5.   Sewitch, D. E. The Perceptual Uncertainty of Having Slept: The Inability to Discriminate Electroencephalographic Sleep From Wakefulness. *Psychophysiology* **21**, 243–259 (1984).

6.   Scott, S. C. & Wilse, B. W. The Perception of Wakefulness Within Sleep. *Sleep* **4**, 177–183 (1981).

7.   Knab, B. & Engel, R. R. Perception of waking and sleeping: Possible implications for the evaluation of insomnia. *Sleep* **11**, 265–272 (1988).

8.   Rezaie, L., Fobian, A. D., McCall, W. V. & Khazaie, H. Paradoxical insomnia and subjective–objective sleep discrepancy: A review. *Sleep Med. Rev.* **40**, 196–202 (2018).

9.   Blake, H. & Gerard, R. W. Potentials During. *Am. J. Physiol.* **119**, 692–703 (1937).

10.   Dinges, D. F., Orne, M. T. & Orne, E. C. Assessing performance upon abrupt awakening from naps during quasi-continuous operations. *Behav. Res. Methods, Instruments, Comput.* **17**, 37–45 (1985).

11.   Noreika, V., Valli, K., Lahtela, H. & Revonsuo, A. Early-night serial awakenings as a new paradigm for studies on NREM dreaming. *Int. J. Psychophysiol.* **74**, 14–18 (2009).

12.   Siclari, F., LaRocque, J. J., Postle, B. R. & Tononi, G. Assessing sleep consciousness within subjects using a serial awakening paradigm. *Front. Psychol.* **4**, 1–9 (2013).

13.   Foulkes, D. & Vogel, G. Mental activity at sleep onset. *J. Abnorm. Psychol.* **70**, 231–243 (1965).

14.   Antrobus, J. S & Saul, H. N. Sleep onset: Subjective, behavioral and

electroencephalographic comparisons. *Waking Sleeping* (1980).

15. Gibson, E., Perry, F., Redington, D. & Kamiya, J. Discrimination of sleep onset stages: behavioral responses and verbal reports. *Percept. Mot. Skills* (1982) doi:10.2466/pms.1982.55.3f.1023.

16. Feige, B., Nanovska, S., Baglioni, C., Bier, B., Cabrera, L., Diemers, S., Quellmalz, M., Siegel, M., Xeni, I., Szentkiralyi, A., Doerr, J. P. & Riemann, D. Insomnia - Perchance a dream? Results from a NREM/REM sleep awakening study in good sleepers and patients with insomnia. *Sleep* **41**, 1–11 (2018).

17. Borkovec, T. D., Lane, T. W. & VanOot, P. H. Phenomenology of sleep among insomniacs and good sleepers: Wakefulness experience when cortically asleep. *J. Abnorm. Psychol.* **90**, 607–609 (1981).

18. Coates, T. J., Killen, J. D., Silverman, S., George, J., Marchini, E., Hamilton, S. & Thoresenz, C. E. Cognitive Activity, Sleep Disturbance, and Stage Specific Differences Between Recorded and Reported Sleep. *Psychophysiology* vol. 20 243–250 (1983).

19. Mercer, J. D., Bootzin, R. R. & Lack, L. C. Insomniacs' Perception of Wake Instead of Sleep. *Sleep* **25**, 559–566 (2002).

20. Buysse, D. J., Germain, A., Hall, M. L., Moul, D. E., Nofzinger, E. A., Begley, A., Ehlers, C. L., Thompson, W. & Kupfer, D. J. EEG spectral analysis in primary insomnia: NREM period effects and sex differences. *Sleep* **31**, 1673–1682 (2008).

21. Spiegelhalder, K., Regen, W., Feige, B., Holz, J., Piosczyk, H., Baglioni, C., Riemann, D. & Nissen, C. Increased EEG sigma and beta power during NREM sleep in primary insomnia. *Biol. Psychol.* **91**, 329–333 (2012).

22. Riedner, B. A., Goldstein, M. R., Plante, D. T., Rumble, M. E., Ferrarelli, F., Tononi, G. & Benca, R. M. Regional Patterns of Elevated Alpha and High-Frequency Electroencephalographic Activity during Nonrapid Eye Movement Sleep in Chronic Insomnia: A Pilot Study. *Sleep* **39**, 801–812 (2016).

23. Andrillon, T., Solelhac, G., Bouchequet, P., Romano, F., Le Brun, M. P., Brigham, M., Chennaoui, M. & Léger, D. Revisiting the value of polysomnographic data in insomnia: more than meets the eye. *Sleep Med.* (2020) doi:10.1016/j.sleep.2019.12.002.

24. St-Jean, G., Turcotte, I., Pérusse, A. D. & Bastien, C. H. REM and NREM power spectral analysis on two consecutive nights in psychophysiological and paradoxical insomnia sufferers. *Int. J. Psychophysiol.* **89**, 181–194 (2013).

25. Krystal, A. D., Edinger, J. D., Wohlgemuth, W. K. & Marsh, G. R. NREM Sleep EEG Frequency Spectral Correlates of Sleep Complaints in Primary Insomnia Subtypes. *Sleep*

25, 630–640 (2002).

26. Colombo, M. A., Ramautar, J. R., Wei, Y., Gomez-Herrero, G., Stoffers, D., Wassing, R., Benjamins, J. S., Tagliazucchi, E., van der Werf, Y. D., Cajochen, C. & Van Someren, E. J. W. Wake High-Density Electroencephalographic Spatiospectral Signatures of Insomnia. *Sleep* **39**, 1015–1027 (2016).

27. Fernandez-Mendoza, J., Calhoun, S. L., Bixler, E. O., Karataraki, M., Liao, D., Vela-Bueno, A., Jose Ramos-Platon, M., Sauder, K. A., Basta, M. & Vgontzas, A. N. Sleep Misperception and Chronic Insomnia in the General Population: Role of Objective Sleep Duration and Psychological Profiles. *Psychosom. Med.* **73**, 88–97 (2011).

28. Kang, S. G., Mariani, S., Marvin, S. A., Ko, K. P., Redline, S. & Winkelman, J. W. Sleep EEG spectral power is correlated with subjective-objective discrepancy of sleep onset latency in major depressive disorder. *Prog. Neuro-Psychopharmacology Biol. Psychiatry* **85**, 122–127 (2018).

29. Perlis, M. L., Smith, M. T., Andrews, P. J., Orff, H. & Giles, D. E. Beta/Gamma EEG activity in patients with primary and secondary insomnia and good sleeper controls. *Sleep* **24**, 110–117 (2001).

30. Maes, J., Verbraecken, J., Willemen, M., De Volder, I., van Gastel, A., Michiels, N., Verbeek, I., Vandekerckhove, M., Wuyts, J., Haex, B., Willemen, T., Exadaktylos, V., Bulckaert, A. & Cluydts, R. Sleep misperception, EEG characteristics and Autonomic Nervous System activity in primary insomnia: A retrospective study on polysomnographic data. *Int. J. Psychophysiol.* **91**, 163–171 (2014).

31. Martinez, D., Breitenbach, T. C. & Lenz, M. do C. S. Light sleep and sleep time misperception - Relationship to alpha-delta sleep. *Clin. Neurophysiol.* **121**, 704–711 (2010).

32. Christensen, J. A. E., Wassing, R., Wei, Y., Ramautar, J. R., Lakbila-Kamal, O., Jennum, P. J. & Van Someren, E. J. W. Data-driven analysis of EEG reveals concomitant superficial sleep during deep sleep in insomnia disorder. *Front. Neurosci.* (2019) doi:10.3389/fnins.2019.00598.

33. Hsiao, F. C., Tsai, P. J., Wu, C. W., Yang, C. M., Lane, T. J., Lee, H. C., Chen, L. C., Lee, W. K., Lu, L. H. & Wu, Y. Z. The neurophysiological basis of the discrepancy between objective and subjective sleep during the sleep onset period: An EEG-fMRI study. *Sleep* **41**, 1–10 (2018).

34. Siclari, F., Baird, B., Perogamvros, L., Bernardi, G., LaRocque, J. J., Riedner, B., Boly, M., Postle, B. R. & Tononi, G. The neural correlates of dreaming. *Nat. Neurosci* **20**,

872–878 (2017).

35.   Kay, D. B., Karim, H. T., Soehner, A. M., Hasler, B. P., James, J. A., Germain, A., Hall, M. H., Franzen, P. L., Price, J. C., Nofzinger, E. A. & Buysse, D. J. Subjective-Objective Sleep Discrepancy Is Associated With Alterations in Regional Glucose Metabolism in Patients With Insomnia and Good Sleeper Controls. *Sleep* **40**, (2017).

36.   Nielsen, T. A. A review of mentation in REM and NREM sleep: "Covert" REM sleep as a possible reconciliation of two opposing models. *Behav. Brain Sci.* **23**, 851–866 (2000).

37.   O'Donnel, D., Silva, E. J., Münch, M., Ronda, J. M., Wang, W. & Duffy, J. F. Comparison of subjective and objective assessments of sleep in healthy older subjects without sleep complaints. *J. Sleep Res.* **18**, 254–263 (2009).

38.   Akersted, T., Hum, K., Minors, D. & Waterhouse, J. Good sleep - its timing and physiological sleep characteristics. *J. Sleep Res.* **6**, 221–229 (1997).

39.   Bernardi, G., Betta, M., Ricciardi, E., Pietrini, P., Tononi, G. & Siclari, F. Regional delta waves in human rapid eye movement sleep. *J. Neurosci.* **39**, 2686–2697 (2019).

40.   Baird, B., Castelnovo, A., Riedner, B. A., Lutz, A., Ferrarelli, F., Boly, M., Davidson, R. J. & Tononi, G. Human Rapid Eye Movement Sleep Shows Local Increases in Low-Frequency Oscillations and Global Decreases in High-Frequency Oscillations Compared to Resting Wakefulness. *eNeuro* **5**, ENEURO.0293-18.2018 (2018).

41.   Funk, C. M., Honjoh, S., Rodriguez, A. V., Cirelli, C. & Tononi, G. Local Slow Waves in Superficial Layers of Primary Cortical Areas during REM Sleep. *Curr. Biol.* **26**, 396–403 (2016).

42.   Nir, Y., Staba, R. J., Andrillon, T., Vyazovskiy, V. V., Cirelli, C., Fried, I. & Tononi, G. Regional slow waves and spindles in human sleep. *Neuron* **70**, 153–69 (2011).

43.   Mölle, M., Bergmann, T. O., Marshall, L. & Born, J. Fast and Slow Spindles during the Sleep Slow Oscillation: Disparate Coalescence and Engagement in Memory Processing. *Sleep* (2011) doi:10.5665/sleep.1290.

44.   Siclari, F., Bernardi, G., Riedner, B. A., LaRocque, J. J., Benca, R. M. & Tononi, G. Two Distinct Synchronization Processes in the Transition to Sleep: A High-Density Electroencephalographic Study. *Sleep* **37**, 1621–1637 (2014).

45.   Jones, B. E. Arousal systems. *Frontiers in Bioscience* vol. 8 (2003).

46.   Fosse, R., Stickgold, R. & Hobson, J. A. Thinking and hallucinating: Reciprocal changes in sleep. *Psychophysiology* **41**, 298–305 (2004).

47.   Hobson, J. A., Stickgold, R. & Pace-Schott, E. F. The neuropsychology of REM sleep dreaming. *Salud Ment.* **22**, 9–22 (1999).

48. Windt, J. M., Nielsen, T. & Thompson, E. Does Consciousness Disappear in Dreamless Sleep? *Trends in Cognitive Sciences* vol. 20 871–882 (2016).

49. Andrillon, T., Burns, A., MacKay, T., Windt, J. & Tsuchiya, N. Wandering minds, sleepy brains: Lapses of attention and local sleep in wakefulness. *bioRxiv* (2020) doi:10.1101/2020.06.23.166991.

50. Windt, J. M. Just in Time—Dreamless Sleep Experience as Pure Subjective Temporality. in *Open MIND* (eds. Metzinger, T. K. & Windt, J. M.) (MIND Group, 2015). doi:10.15502/9783958571174.

51. Yang, C. M., Han, H. Y., Yang, M. H., Su, W. C. & Lane, T. What subjective experiences determine the perception of falling asleep during sleep onset period? *Conscious. Cogn.* **19**, 1084–1092 (2010).

52. Maquet, P., Peters, J. M., Aerts, J., Delfiore, G., Degueldre, C., Luxen, A. & Franck, G. Functional neuroanatomy of human rapid-eye-movement sleep and dreaming. *Nature* **383**, 163–166 (1996).

53. Nofzinger, E. A., Boysse, D. J., Germain, A., Price, J. C., Miewald, J. M. & Kopfer, D. J. Functional neuroimaging evidence for hyperarousal in insomnia. *Am. J. Psychiatry* **161**, 2126–2129 (2004).

54. Bonnet, M. H., Burton, G. G. & Arand, D. L. Physiological and medical findings in insomnia: Implications for diagnosis andcare. *Sleep Medicine Reviews* vol. 18 111–122 (2014).

55. Feige, B., Al-Shajlawi, A., Nissen, C., Voderholzer, U., Hornyak, M., Spiegelhalder, K., Kloepfer, C., Perlis, M. & Riemann, D. Does REM sleep contribute to subjective wake time in primary insomnia? A comparison of polysomnographic and subjective sleep in 100 patients. *J. Sleep Res.* **17**, 180–190 (2008).

56. Wassing, R., Benjamins, J. S., Dekker, K., Moens, S., Spiegelhalder, K., Feige, B., Riemann, D., Van Der Sluis, S., Van Der Werf, Y. D., Talamini, L. M., Walker, M. P., Schalkwijk, F. & Van Someren, E. J. W. Slow dissolving of emotional distress contributes to hyperarousal. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 2538–2543 (2016).

57. Merica, H., Blois, R. & Gaillard, J.-M. Spectral characteristics of sleep EEG in chronic insomnia. *Eur. J. Neurosci.* **10**, 1826–1834 (2001).

58. Aston-Jones, G. & Bloom, F. E. Activity of norepinephrine-containing locus coeruleus neurons in behaving rats anticipates fluctuations in the sleep-waking cycle. *J. Neurosci.* **1**, 876–886 (1981).

59. Léna, I., Parrot, S., Deschaux, O., Muffat-Joly, S., Sauvinet, V., Renaud, B., Suaud-

Chagny, M.-F. & Gottesmann, C. Variations in extracellular levels of dopamine, noradrenaline, glutamate, and aspartate across the sleep-wake cycle in the medial prefrontal cortex and nucleus accumbens of freely moving rats. *J. Neurosci. Res.* **81**, 891–899 (2005).

60. Aston-Jones, G., Chiang, C. & Alexinsky, T. Discharge of noradrenergic locus coeruleus neurons in behaving rats and monkeys suggests a role in vigilance. in *Progress in Brain Research* 501–520 (1991). doi:10.1016/S0079-6123(08)63830-3.

61. Eschenko, O., Magri, C., Panzeri, S. & Sara, S. J. Noradrenergic neurons of the locus coeruleus are phase locked to cortical up-down states during sleep. *Cereb. Cortex* **22**, 426–35 (2012).

62. Eschenko, O. & Sara, S. J. Learning-dependent, transient increase of activity in noradrenergic neurons of locus coeruleus during slow wave sleep in the rat: Brain stem-cortex interplay for memory consolidation? *Cereb. Cortex* **18**, 2596–2603 (2008).

63. Bellesi, M., Tononi, G., Cirelli, C. & Serra, P. A. Region-Specific Dissociation between Cortical Noradrenaline Levels and the Sleep/Wake Cycle. *Sleep* **39**, 143–154 (2016).

64. Lyamin, O. I., Lapierre, J. L., Kosenko, P. O., Kodama, T., Bhagwandin, A., Korneva, S. M., Peever, J. H., Mukhametov, L. M. & Siegel, J. M. Monoamine Release during Unihemispheric Sleep and Unihemispheric Waking in the Fur Seal. *Sleep* **39**, 625–636 (2016).

65. Javoy-Agid, F., Scatton, B., Ruberg, M., L'heureux, R., Cervera, P., Raisman, R., Maloteaux, J.-M., Beck, H. & Agid, Y. Distribution of monoaminergic, cholinergic, and GABAergic markers in the human cerebral cortex. *Neuroscience* **29**, 251–259 (1989).

66. Fuxe, K., Hamberger, B. & Hökfelt, T. Distribution of noradrenaline nerve terminals in cortical areas of the rat. *Brain Res.* **8**, 125–131 (1968).

67. Gais, S., Rasch, B., Dahmen, J. C., Sara, S. & Born, J. The memory function of noradrenergic activity in NON-REM sleep. *J. Cogn. Neurosci.* **23**, 2582–2592 (2011).

68. Van Someren, E. J. W. Brain mechanisms of insomnia: new perspectives on causes and consequences. *Physiol. Rev.* (2020) doi:10.1152/physrev.00046.2019.

69. Vanhatalo, S., Palva, J. M., Holmes, M. D., Miller, J. W., Voipio, J. & Kaila, K. Infraslow oscillations modulate excitability and interictal epileptic activity in the human cortex during sleep. *Proc. Natl. Acad. Sci. U. S. A.* **101**, 5053–5057 (2004).

70. Parrino, L., Ferri, R., Bruni, O. & Terzano, M. G. Cyclic alternating pattern (CAP): The marker of sleep instability. *Sleep Med. Rev.* **16**, 27–45 (2012).

71. Lecci, S., Fernandez, L. M. J., Weber, F. D., Cardis, R., Chatton, J.-Y., Born, J. & Lüthi,

A. Coordinated infraslow neural and cardiac oscillations mark fragility and offline periods in mammalian sleep. *Sci. Adv.* **3**, e1602026 (2017).

72. Ermis, U., Krakow, K. & Voss, U. Arousal thresholds during human tonic and phasic REM sleep. *J. Sleep Res.* **19**, 400–406 (2010).

73. Hilditch, C. J. & McHill, A. W. Sleep inertia: Current insights. *Nature and Science of Sleep* (2019) doi:10.2147/NSS.S188911.

74. Dagnino, N., Favale, E., Loeb, C., Manfredi, M. & Seitun, A. Accelerated synaptic transmission in nucleus ventralis-postero-lateralis during deep sleep. *Experientia* **22**, 329–30 (1966).

75. Favale, E., Loeb, C., Manfredi, M. & Sacco, G. Somatic afferent transmission and cortical responsiveness during natural sleep and arousal in the cat. *Electroencephalogr. Clin. Neurophysiol.* (1965) doi:10.1016/0013-4694(65)90055-6.

76. Della Monica, C., Johnsen, S., Atzori, G., Groeger, J. A. & Dijk, D. J. Rapid eye movement sleep, sleep continuity and slow wave sleep as predictors of cognition, mood, and subjective sleep quality in healthy men and women, aged 20-84 years. *Front. Psychiatry* **9**, 1–17 (2018).

77. Buysse, D. J., Reynolds, C. F., Monk, T. H., Berman, S. R. & Kupfer, D. J. The Pittsburgh sleep quality index: A new instrument for psychiatric practice and research. *Psychiatry Res.* **28**, 193–213 (1989).

78. Horne, J. A. & Ostberg, O. A self-assessment questionnaire to determine morningness-eveningness in human circadian rhythms. *Int. J. Chronobiol.* **4**, 97–110 (1976).

79. Johns, M. W. A new method for measuring daytime sleepiness: The Epworth sleepiness scale. *Sleep* **14**, 540–5 (1991).

80. American Academy of Sleep Medicine. *International Classification of Sleep Disorders*. (American Academy of Sleep Medicine, 2014).

81. Berry, R. B., Budhiraja, R., Gottlieb, D. J., Gozal, D., Iber, C., Kapur, V. K., Marcus, C. L., Mehra, R., Parthasarathy, S., Quan, S. F., Redline, S., Strohl, K. P., Ward, S. L. D. & Tangredi, M. M. Rules for scoring respiratory events in sleep: Update of the 2007 AASM manual for the scoring of sleep and associated events. *J. Clin. Sleep Med.* **15**, 597–619 (2012).

82. Delorme, A. & Makeig, S. EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics. *J. Neurosci. Methods* **134**, 9–21 (2004).

83. Jung, T. P., Makeig, S., Humphries, C., Lee, T. W., Mckeown, M. J., Iragui, V. & Sejnowski, T. J. Removing electroencephalographic artifacts by blind source separation.

Psychophysiology **37**, 163–78 (2000).

84. Pascual-Marqui, R. D. Standardized low-resolution brain electromagnetic tomography (sLORETA): technical details. *Methods Find. Exp. Clin. Pharmacol.* **24 Suppl D**, 5–12 (2002).

85. Ferrarelli, F., Huber, R., Peterson, M. J., Massimini, M., Murphy, M., Riedner, B. A., Watson, A., Bria, P. & Tononi, G. Reduced sleep spindle activity in schizophrenia patients. *Am. J. Psychiatry* (2007) doi:10.1176/ajp.2007.164.3.483.

86. Anderer, P., Klösch, G., Gruber, G., Trenker, E., Pascual-Marqui, R. D., Zeitlhofer, J., Barbanoj, M. J., Rappelsberger, P. & Saletu, B. Low-resolution brain electromagnetic tomography revealed simultaneously active frontal and parietal sleep spindle sources in the human cortex. *Neuroscience* (2001) doi:10.1016/S0306-4522(01)00028-8.

87. Siclari, F., Bernardi, G., Cataldi, J. & Tononi, G. Dreaming in NREM sleep: A high-density EEG study of slow waves and spindles. *J. Neurosci.* **38**, 9175–9185 (2018).