Morphology of migration: Associations between wing shape, bill morphology, and migration in kingbirds (*Tyrannus*)

Short Title: Ecomorphology of migration in kingbirds (*Tyrannus*)

Authors & Affiliations
Maggie P. MacPherson [1,2,*], Alex E. Jahn [3,4], Nicholas A. Mason [2]
[1] Department of Ecology and Evolutionary Biology, Tulane University, New Orleans, LA 70118
[2] Louisiana State University Museum of Natural Sciences, Louisiana State University, Baton Rouge, LA 70803
[3] Departamento de Zoologia, Universidade Estadual Paulista, Av. 24a No. 1515, Rio Claro, Brazil
[4] Environmental Resilience Institute, Indiana University, 717 E 8th St., Bloomington, IN 47408
[*] Corresponding author: Maggie P. MacPherson, E-mail: maggie.macpherson@gmail.com

Abstract

Morphology is closely linked to locomotion and diet in animals. In animals that undertake long-distance migrations, limb-morphology is under selection to maximize mobility and minimize energy expenditure. Migratory behaviors also interact with diet, such that migratory animals tend to be dietary generalists, while sedentary taxa tend to be dietary specialists. Despite a hypothesized link between migration status and morphology, phylogenetic comparative studies have yielded conflicting findings. In this study, we tested for evolutionary associations between migratory status and limb and bill morphology across kingbirds, a pan-American genus of birds with migratory, partially migratory, and sedentary taxa. We found that migratory kingbirds had longer and more pointed wings, in agreement with expectations if selection favors improved aerodynamics for long-distance migration. We also found an association between migratory status and bill shape, such that more migratory taxa had wider and deeper bills and sedentary taxa had longer bills. However, there was no difference in levels of intraspecific morphological variation among migrants, partial migrants, and residents, suggesting that dietary specialization has evolved independently of migration strategy. Evolutionary links between migration, diet, and morphology in kingbirds further strengthen ecomorphological associations that underlie long-distance seasonal movements in animals.

Keywords: ecomorphology, flycatcher, migration, movement ecology, natural selection, partial migration, Tyrannidae

Introduction

Animal movement is linked to morphology at various taxonomic scales. At a macroevolutionary scale, more streamlined or aerodynamic body shapes have been associated with the evolution

of long-distance migration in fish (Common Roach (*Rutilus rutilus*); Chapman et al. 2015),

insects (dragonflies; Johansson et al. 2009), and birds (Fiedler, 2005). Within species, migratory

distance has also been associated with streamlined body shapes in fish (e.g., sockeye salmon

(*Oncorhynchus nerka*); Crossin *et al.*, 2004) and aerodynamic shapes in birds (Voelker, 2001;

Minias *et al.*, 2015; Vágási *et al.*, 2016). While certain taxa with long-distance movements

exhibit strong selection for energy-efficient body shapes, this is not universal (Mulvihill &

Chandler, 1990; Mönkkönen, 1995; Wang & Clarke, 2015). Thus, there is a persistent need to

expand the taxonomic breadth of studies linking migration and morphology to better understand

which lineages exhibit migration-morphology associations, and why these associations vary

among taxa.

Morphology is also shaped by foraging strategies. For example, dietary niche is

associated with head and body shape in fish (Arctic char (*Salvelinus alpinus*); Knudsen *et al.*,

2011, brown trout (*Salmo trutta*); Závorka *et al.*, 2020), birds (Felice *et al.*, 2019), and mammals

(Swanson, Oliveros, & Esselstyn, 2019). Additionally, phenotypic plasticity is expected in dietary

generalists, as shown in comparative common-garden experiments in stickleback minnows

(*Gasterosteus aculeatus*; Svanbäck & Schluter, 2012). In populations recently released from

interspecific competition, such as island colonizers (Wilson, 1961; Clegg & Owens, 2002),

phenotypic plasticity is thought to support increased morphological variation to limit intraspecific

competition (i.e., "niche variation hypothesis", Van Valen, 1965). Migratory lineages of dietary

generalists may have strong preferences for food resources that are easiest to access (Sherry,

1984; Levey & Stiles, 1992; Bell, 2011) or that are superabundant (Moreau, 1952; Morse, 1971;

Willis, 1974). In contrast, sedentary individuals may mitigate intra- and interspecific competition

by seeking more temporally-stable food resources in microhabitats that are more buffered from

environmental fluctuation (e.g., temperature fluctuations; Bell, 2011). Comparisons across avian

species (Levey & Stiles, 1992) and families (Chesser & Levey, 1998) have shown that migratory

taxa that forage on seasonally variable resources tend to exhibit more morphological variation,

presumably associated with opportunistic foraging across a wider dietary breadth among or

within individuals (Bell, 2011).

Testing how migratory status shapes morphological variation among taxa requires a

phylogenetic comparative framework with comprehensive inter- and intraspecific sampling.

Studies comparing migratory to sedentary bird taxa support that long-distance migration favors

longer (Rayner, 1988; Wiedenfeld, 1991; Pérez-Tris & Tellería, 2001; Förschler & Bairlein,

2011; Tellería *et al.*, 2013), and more pointed (Carvalho Provinciato, Araújo, & Jahn, 2018;

Gómez-Bahamón *et al.*, 2020a) wings for increased aerodynamics. Hypotheses that link bill

morphology and migratory status among birds are less well-studied than those involving limb

morphology, but hinge on differences in diet that covary with migratory status (Bell, 2011; but

see Cox, 1968; Herrera, 1978; Leisler, 1990). Sedentary insectivorous taxa often have longer

bills, presumably to improve closing speed for capturing highly mobile prey (Leisler, 1990). In

contrast, migratory taxa may have shorter bills for capturing slow-moving prey such as

caterpillars to feed young during the breeding season and small, abundant arthropods during

the winter (Thiollay, 1988; Leisler, 1990). Furthermore, studies on bill size in relation to foraging

strategies disagree, such that both narrow and wide bills have been associated with generalist

diets (Tellería & Carbonell, 1999; Tellería *et al.*, 2013), and bill size has been shown to differ

between isolated specialist populations of the same species (Alonso *et al.*, 2020).

Early assessments of morphological variation among migratory and sedentary avian

species did not account for their shared evolutionary history (Cox, 1968, 1985; Leisler, 1990).

Although Boyle and Conway (2007) advanced this approach by performing phylogenetically

independent contrasts to address selective pressures for the evolution of migration between

species (see also Gómez *et al.*, 2016; Vágási *et al.*, 2016), the literature on the role of migratory

status in shaping morphological phenotypes in birds draws predominantly from comparisons

among unrelated migratory and sedentary taxa (e.g., Rappole's (1995) critique of Herrera

(1978)). Thus, detailed comparisons among taxa that differ in migratory status within a modern

phylogenetic comparative framework are necessary for a more comprehensive test of whether

and how migration is associated with morphology at different taxonomic scales (e.g., Bolnick *et*

al., 2007).

In this study, we used phylogenetic comparative methods to test for ecomorphological

associations in wing and bill morphology among migratory, partially migratory, and sedentary

kingbirds (*Tyrannus*; Fitzpatrick *et al.*, 2004). Kingbirds are tyrant flycatchers (family

Tyrannidae) with considerable variation in migration status and morphology within and among

species, as well as a rich body of literature linking ecology and morphology (Fitzpatrick &

Schauensee, 1980; Fitzpatrick, 1981, 1985; Sherry, 1984; Cintra, 1997; Fitzpatrick *et al.*, 2004;

Gabriel & Pizo, 2005; Carvalho Provinciato *et al.*, 2018; Gómez-Bahamón *et al.*, 2020a). As

aerodynamic theory predicts that longer, more pointed wings and shorter tails are more efficient

for long-distance migratory flights (Norberg, 1995; Pennycuick, 2008), we expected migratory

taxa to have longer and more pointed wings compared to sedentary taxa (e.g., Kipp, 1942,

1958; Winkler & Leisler, 1992; Mönkkönen, 1995, and references therein; Lockwood, Swaddle,

& Rayner, 1998). Tails of long-distance migrants should also be shorter than more sedentary

individuals to reduce drag during long-distance flights (Rayner, 1988; Norberg, 1990; Winkler &

Leisler, 1992; Förschler & Bairlein, 2011). However, some species' tails are also targets of

sexual selection (Winquist & Lemon, 1994; Mobley, 2002), and tail lengths are often

unassociated with migration strategy (e.g., Voelker, 2001; Neto *et al.*, 2013). Within-population

variation in flight capability could be associated with migration (Fernández & Lank, 2007) or

foraging ecology (Hromada & Tryjanowski, 2003), so we also tested whether more migratory

taxa have more variable wing morphology compared to less migratory taxa.

Bill morphology is closely tied to foraging niche (Snow, 1953, 1954; Lack, 1971) and bill

size and shape is typically related to broad dietary categories in birds (e.g., >50% insectivorous,

granivorous, frugivorous etc.; e.g., Reaney *et al*., 2020). We therefore predicted that there would

be no relationship between migratory status and bill size because all kingbirds are primarily

insectivorous. Additionally, if migration ecology imparts an 'ecological release' that is associated

with more variable, opportunistic diet (Bell, 2011), we expect that more migratory taxa will have

higher coefficients of variation in bill morphology compared to less migratory taxa.

Methods

Phylogeny construction

We extracted the clade of 15 kingbird taxa from the Harvey et al. (2020) phylogeny of the

suboscines to construct a kingbird phylogeny that includes 28 operational taxonomic units

(OTUs). Harvey et al. (2020) included all species (13/13) but only 15% of the subspecific

diversity (3/20). We therefore added *T. savana* taxa following phylogenetic relationships and

branch lengths estimated by Gómez-Bahamón (2020a), while the remaining subspecies were

added as polytomies assuming a most recent common ancestor of 0.5 million years from

members of their species group (Figure 1A)**.** To test the sensitivity of analyses to this

assumption, we tried a range of different dates for the most recent common ancestor for the

added subspecies and this did not change the results (see Output Files at

https://github.com/mmacphe/Tyrannus_morphology).

Morphological measurements

We measured bill length, width, and depth at the distal end of the nares. We also measured

unflattened wing chord length, Kipp's Index (a measure of wing pointedness: the distance

between the tip of the first secondary feather to the tip of the longest primary feather Kipp, 1942,

1958); tail length, and tarsus length on 2108 kingbird study skins from across the ranges of

each species and subspecies (28 operational taxonomic units (OTUs); Figure 1A; Table S1.1).

MM measured 2008 specimens, and identified all individuals to the lowest level of taxonomic

identification (using Clements *et al.*, 2019) and classified each individual as migratory, partially

migratory, or sedentary (following Fitzpatrick *et al.*, 2004). Partially migratory taxa are those in

which individuals vary in migratory tendency (Boyle, 2008). JIGA (see Acknowledgements)

measured 100 specimens to improve sampling of some taxa. A complete list of the specimens

in this study and their associated metadata will be made available at Dryad data repository (doi

to be provided upon acceptance). Bill and tail measurements were taken twice and were

confirmed to be within one mm of one another. Both right and left wings and tarsi were

measured and averaged. Measurements by JIGA were only taken once. We measured tail

length as the longest rectrix to the nearest 0.1 cm (Pyle *et al.*, 1997). For most measurements,

we used a Mitutoyo brand IP 67 digital calipers (part number 573-271) with a range of up to

15.24 cm, with 0.00127 cm resolution. For tails longer than 15.24 cm, we used a 30.48 cm

stainless steel ruler placed between the two middle rectrices, and when tails were longer than

30.48 cm photos were taken of the tails above a 0.64 x 0.64 cm square grid with the calipers

measuring to their extent and ImageJ was used to calculate the full tail length (ImageJ protocol

available in Supporting Information 2).

We used linear model comparisons to test whether accounting for age (juvenile versus

adult) and sex (female versus male) classes improved model performance. We found that AIC

values were lower in models that did not include age (except in tarsus length), and although AIC

values were lower in some models that included sex (bill length, bill width, bill depth, and tarsus

length), we present the results of all adult individuals in the main text as our results did not differ

when analyzing sexes independently (Supporting Information 3). To test associations between

migratory strategy and morphology, we calculated the coefficient of variation (mean/standard

deviation) for each taxon.

Phylogenetic Principal Components Analysis

As selection is thought to act independently on bill morphology compared to flight feather

morphology, we conducted separate principal component analyses for bills and flight feathers to

estimate their overall relative size and shape using the phyl.pca function from phytools package

(Revell, 2009; Table 1). Because two species have extremely long tail feathers (*T. savana* and

T. forficatus), we explored whether the PPCA would be substantially improved with their

omission, but the amount of variance explained by PPC1 was similar regardless of whether

these taxa were included or excluded (Table S1.2). The PPCA scores were included in the

following phylogenetic analysis of variance (described in the next section). PPCA was also

conducted for females and males separately and results can be found in Supplementary

Material (Supporting Information 3).

Figure 1. Ancestral state reconstruction of migration ecology category and Kipp's Index, and ordination of 28 kingbird OTUs for bill (length, width, depth) and feather (wing chord length and Kipp's Index) morphometrics. A) Kingbird phylogeny showing ancestral state reconstruction of migration ecology strategy (pie charts at each node), and Kipp's Index (branch greyscale). B-D) PPCA plots showing individuals color-coded to species (identified at the tip labels of the phylogeny), shape-coded to subspecies, and their status as migratory, partially migratory, or sedentary is distinguished via the shape outline. Pie charts for each node in the phylogeny show the ancestral state reconstruction of migration type.

171 **Table 1**. Results of phylogenetic principal component analysis (PPCA) of bill and feather

172 morphometrics. Percent variance explained by each eigenvector is in brackets for each principal

173 component.

Loadings	Bill PPC1 (94.68%)	Bill PPC2 (4.30%)	lambda
Bill length	0.99	0.14	0.96
Bill width	0.93	-0.33	
Bill depth	0.94	-0.25	
Feather PPC1 (93.18%)		**Feather PPC2 (6.24%)**	
Kipp's Index	-0.33	-0.64	0.96
Wing chord	-0.072	-0.99	
Tail length	-0.99	0.003	
CV Bill PPC1 (80.17%)			
CV Bill length	0.92		0.03
CV Bill width	0.91		
CV Bill depth	0.82		
CV Feather PPC1 (82.52%)			
CV Kipp's Index	-0.38		0.50
CV Wing chord	-0.42		
CV Tail length	-0.99		

Phylogenetic Analysis of Variance

We compared morphology between migratory, partially migratory, and sedentary taxa via a

phylogenetic analysis of variance (Garland *et al.*, 1993) with the phylANOVA function in the

phytools package (Revell, 2012). As differences in body size can account for much of the

variation among species (Albrecht, Gelvin, & Hartman, 1993; McCoy *et al.*, 2006; Revell, 2009;

Berner, 2011), we extracted phylogenetic residuals (Revell, 2009) for each dependent variable

(bill length, bill width, bill depth, wing chord, Kipp's Index, and tail length) with tarsus length as

an approximation of body size and the independent variable. We opted to use tarsus instead of

mass to adjust for body size (Rising & Somers, 1989; Senar & Pascual, 1997) because mass

can change intra-annually (particularly in migratory birds; Lindström & Piersma, 1993), mass

can also vary by sex in some kingbirds (e.g., *T. melancholicus*, Jahn *et al.*, 2010; *T. savana*,

Carvalho Provinciato *et al.*, 2018; *T. tyrannus*, Murphy, 2007), and mass data were not available

for many of the museum specimens we measured. We then conducted a phylogenetic ANOVA

using the residual values, presenting the overall average value of P-values of 1000 replicates.

Post-hoc comparisons identified differences in mean values between migration categories (a

default setting in the phylANOVA function), which were subsequently displayed with boxplots

(Figure 2 and 3). We performed an ancestral state reconstruction of Kipp's Index after finding it

was associated with migration status via the phylANOVA, and we also performed an ancestral

state reconstruction of migratory status. Specifically, we used the contMap function in the

phytools package for continuous variables (Revell, 2012) and the fit.mle function in the

diversitree package (FitzJohn, 2012; Figure 1A, Figure S1.1, Figure S1.2).

All statistical analyses were performed using program R version 4.0.3 (R Core Team,

2020). R code is provided for all analytical procedures undertaken in this manuscript at

https://github.com/mmacphe/Tyrannus_morphology.

Results

The average lengths for each morphometric +/- standard deviation for each OTU are reported in Table S1.1. We found that migratory taxa had longer wings than sedentary taxa (Figure 2F, Table 2), and pointier wings than both partially migratory and sedentary taxa (Figure 2D, Table 2). However, we found no association between migratory status and coefficients of variation in bill morphology (Figure 2G-I, Table 2). In agreement with our prediction, we found no relationship between migratory status and bill length, width, depth, or bill PPC1 (Figure 2A-C, Figure 3A, Table 2). However, migratory and partially migratory taxa had shorter, wider, and deeper bills (bill PPC2) compared to sedentary taxa, which had longer, shallower, and more narrow bills (Figure 3B, Table 2).

Figure 2. Phylogenetic ANOVA results comparing residuals of bill and feather morphometrics

(A-F), and coefficient of variation in bill and feather morphometrics (G-L) across migratory,

partially migratory, and sedentary kingbird OTUs. Significant differences are shown by different

letters: A versus B.

216 **Figure 3**. Phylogenetic ANOVA results comparing bill and feather phylogenetic principal

217 component (PPC) scores across migratory, partially migratory, and sedentary kingbird OTUs.

218 Letters above each barplot show groupings resulting from post-hoc pairwise comparisons.

220 **Table 2**. T and P values from phylogenetic ANOVA analysis. Significant results are in bold.

221 Model F and Pr(>F) values can be found in Table S1.3.

Morphometric	migratory v. partially migratory		migratory v. sedentary		partially migratory v. sedentary	
	T	P	T	P	T	P
Bill length	0.31	0.72	0.09	0.96	0.26	0.86
CV Bill length	1.12	0.26	1.36	0.38	2.40	0.07
Bill width	0.19	0.84	1.62	0.31	1.58	0.27
CV Bill width	0.03	0.98	2.12	0.18	1.82	0.20
Bill depth	0.46	0.63	2.10	0.17	1.27	0.38
CV Bill depth	0.15	0.89	1.97	0.22	1.83	0.19
Bill PC1 (size)	0.08	0.92	0.66	0.69	0.46	0.74
CV Bill PC1 (size)	0.06	0.95	0.48	0.73	0.34	0.83

Bill PC2 (shape)	0.64	0.52	3.94	**<0.05**	2.63	**0.05**
CV Bill PC2 (shape)	0.59	0.55	2.89	0.09	1.79	0.20
Kipp's Index	1.99	**0.05**	3.57	**0.02**	0.83	0.54
CV Kipp's Index	1.08	0.29	1.24	0.42	0.15	0.91
Wing chord length	0.57	0.56	3.58	**0.02**	2.41	0.07
CV Wing chord length	0.18	0.86	0.39	0.82	0.14	0.92
Tail length	0.88	0.38	0.48	0.74	1.38	0.31
CV Tail length	1.18	0.23	0.48	0.81	0.90	0.57
Feather PC1 (size)	0.81	0.36	0.49	0.74	1.31	0.33
CV Feather PC1 (size)	0.85	0.40	0.40	0.80	1.28	0.37
Feather PC2 (shape)	0.92	0.31	2.45	0.12	1.06	0.44
CV Feather PC2 (shape)	0.16	0.87	1.54	0.32	1.48	0.28
CV Tarsus length	0.25	0.79	2.56	0.10	2.45	0.07

Discussion

We found evidence of an association between migratory status and wing morphology in the

kingbirds. This corroborates the idea that migratory birds have evolved morphological features

that help in sustained, long-distance flight, such as longer wings (Winkler & Leisler, 1992; Milá,

Wayne, & Smith, 2008; Neto *et al.*, 2013; but not Carvalho Provinciato *et al.*, 2018). There is a

strong genetic component determining wing length (Böhning-Gaese & Oberrath, 1999; Tarka *et*

al., 2010), and among migratory taxa, wing length is positively associated with migratory

distance (Förschler & Bairlein, 2011; Rönn, Shafer, & Wolf, 2016; Vágási *et al.*, 2016; but see

Carvalho Provinciato *et al.*, 2018). Partial migration, wherein only some individuals migrate, is

thought to be an evolutionary stepping-stone between sedentarism and obligate migration

(Berthold, 1999; Bell, 2000; Chapman *et al.*, 2011). Interestingly, we found that migratory taxa

had significantly pointier wings than partially migratory taxa and although not significant, partially

migratory taxa appear to have intermediate wing chord lengths between sedentary and

migratory taxa. Longer wings may allow for more flexibility in altering migration speeds

according to conditions experienced during migration (Hahn *et al.*, 2016), but there may be

ecological limits and trade-offs imposed on wing length by other selective pressures (e.g.,

shorter wings for agility in flight for arboreal species, or as an adaptation for evading predators,

Fernández & Lank, 2007; or a consequence of year-round territoriality, Sheard *et al.*, 2020), or

by foraging strategy (e.g., shorter wings for more efficient hovering flight; Marchetti, Price, &

Richman, 1995).

Our finding of differences in bill shape between migratory and sedentary taxa suggest

bill shape is a target of selection pressures related to migration ecology. However, our

comparisons of coefficients of variation in the current study did not support that intraspecific

competition has driven diversifying selection in bill morphology amongst generalist migratory

kingbirds. A simple ratio of bill length/bill width has been shown to hold potential significance in

foraging behaviors in other tyrannids (Botero-Delgadillo & Bayly, 2012), but our findings

disagree with other research across the Tyrannidae describing generalists as having

intermediate (neither large or small) bill morphologies (Fitzpatrick, 1985). We found that

sedentary taxa had longer, narrower, and shallower bills compared to migratory and partially

migratory taxa (Figure 1B; Table 1). Previous work has shown that directional selection

increasing bill length in dietary specialists may be driven by interspecific competition, including

from closely related taxa (Freed, Medeiros, & Cann, 2016). Thus, other selective pressures may

be acting upon the bill shape and size among migratory, partially migratory, and sedentary taxa.

For example, bill morphology has been linked with thermoregulation in dietary generalists, and

bill width and depth are adaptive for dissipating heat during migration for improved

thermoregulation (Danner & Greenberg, 2015; Friedman *et al.*, 2017; Danner *et al.*, 2017). If

convergence on bill shape is driven mainly by diet, future research integrating multivariate or

nonlinear bill morphometrics (e.g., hooked bills, lateral and longitudinal bill curvature, bill surface

area; Shao *et al.*, 2016) with degrees of intraguild dietary specialization and foraging behaviors

would broaden our understanding of selection mechanisms shaping bill phenotypes.

Aspects of kingbird ecology may partially explain the associations between migratory

status and morphology that may or may not be present in other clades. Kingbirds are open-

habitat foragers and thus may be predisposed to evolve migration under the 'evolutionary

precursor hypothesis' (Fitzpatrick, 1985; Levey & Stiles, 1992). Furthermore, gains and losses

of migration are common in this group (Gómez-Bahamón *et al.*, 2020a). As such, plasticity in

migratory life history strategies of our study taxa could work in tandem with flexibility in wing and

bill morphology. The difference in bill shape that we found seemed to be driven by the very

short, thick, and deep bills present in two migratory *T. crassirostris* subspecies, and partially

migratory *T. couchii* compared to very long, narrow, and shallow bills in sedentary *T.*

caudifasciatus subspecies (six OTUs with island distributions distinct from one another, Mobley

& de Juana, 2020; Figure S1.2). Thus, this trend could be the result of ecological release

commonly experienced by island taxa (Wilson, 1961; Clegg & Owens, 2002; Cooney *et al.*,

2017). Tail length, and variation in wing chord length, and Kipp's Index could also be under

sexual selection. This could particularly be the case in the kingbird species where tail length is

suspected to be an honest indicator of mate quality (e.g., *T. forficatus,* Regosin & Pruett-Jones,

2001). Additionally, other forms of natural selection could be acting on flight morphology in

species that use aerial courtship displays or engage in territorial encounters with conspecifics

(e.g., *T. savana*; Jahn & Tuero, 2020; Gómez-Bahamón *et al.*, 2020b), or in which sexual

dimorphism may occur (e.g., *T. dominicensis*; Haberman, MacKenzie, & Rising, 1991).

Additional studies spanning multiple genera will aid in further elucidating how morphology is

shaped by migratory status.

Phenotypes shaped by migratory status may be more complex than has been historically

assumed with linear measurements of morphology alone. Divergence in bill shape between

migratory and sedentary taxa may define an important morphological feature that is a part of the

suite of co-adaptations for migratory life histories (Bell, 2000). Morphology may be driven by

selection related to migration itself such as increased needs of heat dissipation or conservation

found in other studies (see above). Alternatively, morphology may be shaped by a more

complex competitive landscape that affects migratory and sedentary taxa differently. In taxa

occupying tropical regions, like many kingbird taxa, the dry season is when both intra- and

interspecific competition increase due to 'winter food limitation' and competition with migrants

returning to wintering grounds, in addition to competition with juvenile conspecifics

(Hespenheide, 1975; Emlen, 1977; Orejuela, Raitt, & Álvarez, 1980; Stiles, 1980; Waide, 1980;

Rappole, Ramos, & Winker, 1989; Rappole, 1995; Sherry, 2005; Danner *et al.*, 2013). Selection

may favor longer bills among sedentary taxa to resolve both intra- and interspecific conflict

(Table 1, Figure 3B), allowing access to more temporally stable food resources in microhabitats

buffered from spatial and temporal environmental fluctuations (Bell, 2011).

Migratory and sedentary taxa may not differ in linear bill metrics if a combination of

characteristics (e.g., Table 1, Figure 3B) support a more generalist foraging strategy. For

example, specific bill shapes could accommodate spatiotemporal changes in the competitive

landscape (Navalón *et al.*, 2019) or provide increased stability for prey capture in perch-

gleaning or hover-gleaning foraging techniques (Fitzpatrick, 1985; Fitzpatrick *et al.*, 2004;

Botero-Delgadillo, 2011; Botero-Delgadillo & Bayly, 2012). The diversity of habitats that

migratory taxa encounter during their annual cycle may shape phenotypes in more complex

ways than previously thought, and techniques that incorporate more advanced characterizations

of morphological variation may expose novel insight into how movement life history strategies

shape phenotypes (e.g., Pol *et al.*, 2009; Mallarino *et al.*, 2011; Navalón *et al.*, 2019; Alonso *et*

al., 2020; Medina *et al.*, 2020).

Here we assessed the impact of variation in migration strategies on morphology in the

kingbirds (genus *Tyrannus*). Aspects of both wing and bill morphology appear to evolve in

association with migratory status in this group consisting of migratory, partially migratory, and

sedentary taxa. Our results suggest that migratory status shapes wing morphology in a

widespread avian genus, and that multivariate bill shape metrics (i.e., PPC2) may distinguish

sedentary from migratory and partially migratory lineages. Thus, adaptive phenotypes may be

related to migratory status in more complex ways than previously understood. Assessments of

the various mechanisms driving patterns in bill shape (e.g., heat dissipation, foraging strategy,

and competitive landscape) across a broader range of taxonomic groups that differ in migration

strategies would complement our refinement of the morphology of migration in kingbirds.

Data Availability

324 R code for all statistical analyses and results shared in tables and figures is available from our

325 github repository (https://github.com/mmacphe/Tyrannus_morphology). The measurements of

326 all individuals used in this research will be available at an online data repository upon

327 acceptance (for now, this can be found here).

Acknowledgements

We thank José Ignacio Giraldo Arango ("Nacho") for measuring specimens, and Brenna and Michael Wells, and Shelbey Hearn for assistance measuring specimens used in this study. Ben Marks, Shannon Hacket, and John Bates from the Field Museum of Natural History, Bradley Millen, and Allan Baker from the Royal Ontario Museum, Gary Stiles from the Universidad Nacional Instituto de Ciencias Naturales, Jacob Saucier, Helen James, and Gary Graves from the Smithsonian National Museum of Natural History, and Steven Cardiff, and Van Remsen from the Louisiana State University Museum of Natural Science provided assistance in accessing the specimens used in this study. Thanks to Robb Brumfield and Michael Harvey for providing an early-access version of the suboscine phylogeny for our study. We also thank Dr. Calandra Stanley, Dr. Caleb McMahan, the Toronto MacPhersons and the Bogotá Acevedos for their special help aiding the lead author to access collections. Samantha Rutledge, David Vander Pluym, and Subir Shakya provided helpful feedback on an earlier version of this manuscript. We wish to acknowledge www.gbif.org for providing the ROM database (accessed September 30, 2020) and Field Museum database (accessed October 20, 2020) online that was used to validate locality data previously recorded by hand. Amethys E'etessam, Clare Lister, and Trey Hendrix also helped with georeferencing from locality data that was a part of this project.

Literature Cited

Albrecht GH, Gelvin BR & Hartman SE. **1993**. Ratios as a size adjustment in morphometrics. *American Journal of Physical Anthropology* **91**: 441–468.

Alonso D, Fernández-Eslava B, Edelaar P & Arizaga J. **2020**. Morphological divergence among Spanish Common Crossbill populations and adaptations to different pine species. *Ibis* **162**: 1279–1291.

Bell C. **2000**. Process in the evolution of bird migration and pattern in avian ecogeography. *Journal of Avian Biology* **31**: 258–265.

Bell C. **2011**. Resource buffering and the evolution of bird migration. *Evolutionary Ecology* **25**:
91–106.

Berner D. **2011**. Size correction in biology: how reliable are approaches based on (common)
principal component analysis? *Oecologia* **166**: 961–971.

Berthold P. **1999**. A comprehensive theory for the evolution, control and adaptability of avian
migration. *Ostrich* **70**: 1–11.

Böhning-Gaese K & Oberrath R. **1999**. Phylogenetic effects on morphological, life-history,
behavioural and ecological traits of birds. : 18.

Bolnick DI, Svanbäck R, Araújo MS & Persson L. **2007**. Comparative support for the niche
variation hypothesis that more generalized populations also are more heterogeneous.
Proceedings of the National Academy of Sciences **104**: 10075–10079.

Botero-Delgadillo E. **2011**. Cuantificando el comportamiento: Estrategias de búsqueda y
ecología de forrajeo de 12 especies sintópicas de Atrapamoscas (Tyrannidae) en la Sierra
Nevada de Santa Marta, Colombia. *Revista Brasileira de Ornitologia* **19**: 343–357.

Botero-Delgadillo E & Bayly NJ. **2012**. Does morphology predict behavior? Correspondence
between behavioral and morphometric data in a Tyrant-flycatcher (Tyrannidae) assemblage in
the Santa Marta Mountains, Colombia. *Journal of Field Ornithology* **83**: 329–342.

Boyle WA. **2008**. Partial migration in birds: tests of three hypotheses in a tropical lekking
frugivore. *Journal of Animal Ecology* **77**: 1122–1128.

Boyle WA & Conway CJ. **2007**. Why Migrate? A Test of the Evolutionary Precursor
Hypothesis. *The American Naturalist* **169**: 344–359.

Carvalho Provinciato IC, Araújo MS & Jahn AE. **2018**. Drivers of wing shape in a widespread
Neotropical bird: a dual role of sex-specific and migration-related functions. *Evolutionary*
Ecology **32**: 379–393.

Chapman BB, Brönmark C, Nilsson JÅ & Hansson LA. **2011**. The ecology and evolution of
partial migration. *Oikos* **120**: 1764–1775.

Chapman BB, Hulthén K, Brönmark C, Nilsson PA, Skov C, Hansson LA & Brodersen J.
2015. Shape up or ship out: migratory behaviour predicts morphology across spatial scale in a
freshwater fish. *Journal of Animal Ecology* **84**: 1187–1193.

Chesser RT & Levey DJ. **1998**. Austral Migrants and the Evolution of Migration in New World
Birds: Diet, Habitat, and Migration Revisited. *The American Naturalist* **152**: 311–319.

Cintra R. **1997**. Spatial Distribution and Foraging Tactics of Tyrant Flycatchers in Two Habitats
in the Brazilian Amazon. *Studies on Neotropical Fauna and Environment* **32**: 17–27.

Clegg SM & Owens PF. **2002**. The 'island rule' in birds: medium body size and its ecological
explanation. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **269**:
1359–1365.

Clements JF, Schulenberg TS, Iliff MJ, Billerman SM, Fredericks TA, Sullivan BL & Wood CL. **2019**. The eBird/Clements Checklist of Birds of the World: v2019.

Cooney CR, Bright JA, Capp EJR, Chira AM, Hughes EC, Moody CJA, Nouri LO, Varley ZK & Thomas GH. **2017**. Mega-evolutionary dynamics of the adaptive radiation of birds. *Nature* **542**: 344–347.

Cox GW. **1968**. The Role of Competition in the Evolution of Migration. *Evolution* **22**: 180–192.

Cox GW. **1985**. The Evolution of Avian Migration Systems between Temperate and Tropical Regions of the New World. *The American Naturalist* **126**: 451–474.

Crossin GT, Hinch SG, Farrell AP, Higgs DA, Lotto AG, Oakes JD & Healey MC. **2004**. Energetics and morphology of sockeye salmon: effects of upriver migratory distance and elevation. *Journal of Fish Biology* **65**: 788–810.

Danner RM, Greenberg RS, Danner JE, Kirkpatrick LT & Walters JR. **2013**. Experimental support for food limitation of a short-distance migratory bird wintering in the temperate zone. *Ecology* **94**: 2803–2816.

Danner RM, Gulson-Castillo ER, James HF, Dzielski SA, Frank DC III, Sibbald ET & Winkler DW. **2017**. Habitat-specific divergence of air conditioning structures in bird bills. *The Auk* **134**: 65–75.

Danner RM & Greenberg R. **2015**. A critical season approach to Allen's rule: Bill size declines with winter temperature in a cold temperate environment. *Journal of Biogeography* **42**: 114–120.

Emlen JT. **1977**. Land Bird Communities of Grand Bahama Island: The Structure and Dynamics of an Avifauna. *Ornithological Monographs*: iii–129.

Felice RN, Tobias JA, Pigot AL & Goswami A. **2019**. Dietary niche and the evolution of cranial morphology in birds. *Proceedings of the Royal Society B: Biological Sciences* **286**: 20182677.

Fernández G & Lank DB. **2007**. Variation in the Wing Morphology of Western Sandpipers (Calidris Mauri) in Relation to Sex, Age Class, and Annual Cycle. *The Auk* **124**: 1037–1046.

Fiedler W. **2005**. Ecomorphology of the External Flight Apparatus of Blackcaps (Sylvia atricapilla) with Different Migration Behavior. *Annals of the New York Academy of Sciences* **1046**: 253–63.

FitzJohn RG. **2012**. Diversitree: comparative phylogenetic analyses of diversification in R. *Methods in Ecology and Evolution* **3**: 1084–1092.

Fitzpatrick JW. **1981**. Search strategies of tyrant flycatchers. *Animal Behaviour* **29**: 810–821.

Fitzpatrick JW. **1985**. Form, foraging behavior, and adaptive radiation in the Tyrannidae BT - Neo-Tropical Ornithology Ornithological Monographs No. 26. In: *Neo-Tropical Ornithology Ornithological Monographs No. 26.*, 447–470.

Fitzpatrick JW, Bates JM, Bostwick KS, Caballero IC, Clock BM, Farnsworth A, Hosner
PA, Joseph L, Langham GM & Lebbin DJ. **2004**. Family Tyrannidae (tyrant-flycatchers).
Handbook of the birds of the world **9**: 170–462.

Fitzpatrick W & Schauensee D. **1980**. Foraging behavior of Tyrant flycatchers. : 43–57.

Förschler MI & Bairlein F. **2011**. Morphological Shifts of the External Flight Apparatus across
the Range of a Passerine (Northern Wheatear) with Diverging Migratory Behaviour. *PLOS ONE*
6: e18732.

Freed LA, Medeiros MCI & Cann RL. **2016**. Multiple Reversals of Bill Length over 1.7 Million
Years in a Hawaiian Bird Lineage. *The American Naturalist* **187**: 363–371.

Friedman NR, Harmáčková L, Economo EP & Remeš V. **2017**. Smaller beaks for colder
winters: Thermoregulation drives beak size evolution in Australasian songbirds. *Evolution* **71**:
2120–2129.

Gabriel V de A & Pizo MA. **2005**. Foraging behavior of tyrant flycatchers (Aves, Tyrannidae) in
Brazil. *Revista Brasileira de Zoologia* **22**: 1072–1077.

Garland T, Dickerman AW, Janis CM & Jones JA. **1993**. phylogenetic analysis of covariance
by computer simulation. *Systematic Biology* **42**: 265–292.

Gómez C, Tenorio EA, Montoya P & Cadena CD. **2016**. Niche-tracking migrants and niche-
switching residents: evolution of climatic niches in New World warblers (Parulidae). *Proceedings*
of the Royal Society B: Biological Sciences **283**: 20152458.

Gómez-Bahamón V, Márquez R, Jahn AE, Miyaki CY, Tuero DT, Laverde-R O, Restrepo S
& Cadena CD. **2020a**. Speciation Associated with Shifts in Migratory Behavior in an Avian
Radiation. *Current Biology* **30**: 1312-1321.e6.

Gómez-Bahamón V, Tuero DT, Castaño MI, Jahn AE, Bates JM & Clark CJ. **2020b**.
Sonations in Migratory and Non-migratory Fork-tailed Flycatchers (*Tyrannus savana*).
Integrative and Comparative Biology **60**: 1147–1159.

Haberman K, MacKenzie DI & Rising JD. **1991**. Geographic Variation in the Gray Kingbird
(Variación geográfica en Tyrannus dominicensis). *Journal of Field Ornithology* **62**: 117–131.

Hahn S, Korner-Nievergelt F, Emmenegger T, Amrhein V, Csörgő T, Gursoy A, Ilieva M,
Kverek P, Pérez-Tris J, Pirrello S, Zehtindjiev P & Salewski V. **2016**. Longer wings for faster
springs – wing length relates to spring phenology in a long-distance migrant across its range.
Ecology and Evolution **6**: 68–77.

Harvey M, Bravo G, Claramunt S, Cuervo A, Derryberry G, Battilana J, Seeholzer G,
McKay JS, O'Meara BC, Faircloth BC, Edwards S, Pérez-Emán J, Moyle RG, Sheldon FH,
Aleixo A, Smith BT, Chesser RT, Silveira LF, Cracraft J, Brumfield RT & Derryberry EP.
2020. The evolution of a tropical biodiversity hotspot. *Science* **370**: 1343–1348.

Herrera CM. **1978**. Ecological Correlates of Residence and Non-Residence in a Mediterranean
Passerine Bird Community. *Journal of Animal Ecology* **47**: 871–890.

Hespenheide HA. **1975**. Selective Predation by Two Swifts and a Swallow in Central America.
Ibis **117**: 82–99.

Hromada M & Tryjanowski P. **2003**. Animals of different phenotype differentially utilise dietary
niche - the case of the Great Grey Shrike Lanius excubitor. **80**: 8.

Jahn AE, Levey DJ, Mamani AM, Saldias M, Alcoba A, Ledezma MJ, Flores B, Vidoz JQ &
Hilarion F. **2010**. Seasonal differences in rainfall, food availability, and the foraging behavior of
Tropical Kingbirds in the southern Amazon Basin. *Journal of Field Ornithology* **81**: 340–348.

Jahn AE & Tuero DT. **2020**. Fork-tailed Flycatcher (Tyrannus savana). *Birds of the World*.

Johansson F, Söderquist M & Bokma F. **2009**. Insect wing shape evolution: independent
effects of migratory and mate guarding flight on dragonfly wings. *Biological Journal of the*
Linnean Society **97**: 362–372.

Kipp F. **1942**. Über Flügelbau und Wanderzug der Vögel. *Biologisches Zentralblatt* **62**: 289–
299.

Kipp FA. **1958**. Zur geschichte des Vogelzuges auf der grundlage der Flügelanpassungen.
Vogelwarte **19**: 233–242.

Knudsen R, Siwertsson A, Adams CE, Garduño-Paz M, Newton J & Amundsen PA. **2011**.
Temporal stability of niche use exposes sympatric Arctic charr to alternative selection
pressures. *Evolutionary Ecology* **25**: 589–604.

Lack D. **1971**. *Ecological isolation in birds*.

Leisler B. **1990**. Selection and Use of Habitat of Wintering Migrants. In: Gwinner E, ed. *Bird*
Migration. Berlin, Heidelberg: Springer Berlin Heidelberg, 156–174.

Levey DJ & Stiles FG. **1992**. Evolutionary Precursors of Long-Distance Migration: Resource
Availability and Movement Patterns in Neotropical Landbirds. *The American Naturalist* **140**:
447–476.

Lindström Å & Piersma T. **1993**. Mass changes in migrating birds: the evidence for fat and
protein storage re-examined. *Ibis* **135**: 70–78.

Lockwood R, Swaddle JP & Rayner JMV. **1998**. Avian Wingtip Shape Reconsidered: Wingtip
Shape Indices and Morphological Adaptations to Migration. *Journal of Avian Biology* **29**: 273–
292.

Mallarino R, Grant PR, Grant BR, Herrel A, Kuo WP & Abzhanov A. **2011**. Two
developmental modules establish 3D beak-shape variation in Darwin's finches. *Proceedings of*
the National Academy of Sciences **108**: 4057–4062.

Marchetti K, Price T & Richman A. **1995**. Correlates of Wing Morphology with Foraging
Behaviour and Migration Distance in the Correlates of wing morphology with foraging behaviour
and migration distance in the genus Phylloscopus. *Journal of Avian Biology* **26**: 177–181.

McCoy MW, Bolker BM, Osenberg CW, Miner BG & Vonesh JR. **2006**. Size correction:
comparing morphological traits among populations and environments. *Oecologia* **148**: 547–554.

Medina JJ, Maley JM, Sannapareddy S, Medina NN, Gilman CM & McCormack JE. **2020**. A
rapid and cost-effective pipeline for digitization of museum specimens with 3D photogrammetry.
PLOS ONE **15**: e0236417.

Milá B, Wayne RK & Smith TB. **2008**. Ecomorphology of Migratory and Sedentary Populations
of the Yellow-Rumped Warbler (Dendroica Coronata). *The Condor* **110**: 335–344.

**Minias P, Meissner W, Włodarczyk R, Ożarowska A, Piasecka A, Kaczmarek K &
Janiszewski T**. **2015**. Wing shape and migration in shorebirds: a comparative study. *Ibis* **157**:
528–535.

Mobley JA. **2002**. *Molecular phylogenetics and the evolution of nest building in kingbirds and
their allies (Aves: Tyrannidae)*. University of California, Berkeley.

Mobley JA & de Juana E. **2020**. Loggerhead Kingbird (Tyrannus caudifasciatus). *Birds of the
World*.

Mönkkönen M. **1995**. Do migrant birds have more pointed wings?: A comparative study.
Evolutionary Ecology **9**: 520–528.

Moreau RE. **1952**. The Place of Africa in the Palaearctic Migration System. *Journal of Animal
Ecology* **21**: 250–271.

Morse DH. **1971**. The Insectivorous Bird as an Adaptive Strategy. *Annual Review of Ecology
and Systematics* **2**: 177–200.

Mulvihill RS & Chandler CR. **1990**. The Relationship between Wing Shape and Differential
Migration in the Dark-Eyed Junco. *The Auk* **107**: 490–499.

Murphy MT. **2007**. A Cautionary Tale: Cryptic Sexual Size Dimorphism in a Socially
Monogamous Passerine. *The Auk* **124**: 515–525.

Navalón G, Bright JA, Marugán-Lobón J & Rayfield EJ. **2019**. The evolutionary relationship
among beak shape, mechanical advantage, and feeding ecology in modern birds*. *Evolution* **73**:
422–435.

Neto JM, Gordinho L, Belda EJ, Marín M, Monrós JS, Fearon P & Crates R. **2013**.
Phenotypic Divergence among West European Populations of Reed Bunting Emberiza
schoeniclus: The Effects of Migratory and Foraging Behaviours. *PLOS ONE* **8**: e63248.

Norberg UM. **1990**. *Vertebrate flight: mechanics, physiology, morphology, ecology and
evolution*. Berlin: Springer.

Norberg UM. **1995**. WING DESIGN AND MIGRATORY FLIGHT. *Israel Journal of Ecology and
Evolution* **41**: 297–305.

Orejuela JE, Raitt RJ & Álvarez H. **1980**. *Differential use by North American migrants of three
types of Colombian forests: 253-264 (en) KEAST, A. & MORTON, ES (eds.). Migrant Birds in*

the Neotropics: Ecology, Behavior, Distribution and Conservation. Smithsonian Institution Press.
Washington DC, USA.

Pennycuick C. **2008**. The membrane wings of bats and pterosaurs Modelling the Flying Bird ed
CJ Pennycuick.

Pérez-Tris J & Tellería JL. **2001**. Age-related variation in wing shape of migratory and
sedentary Blackcaps Sylvia atricapilla. *Journal of Avian Biology* **32**: 207–213.

Pol MVD, Ens BJ, Oosterbeek K, Brouwer L, Verhulst S, Tinbergen JM, Rutten AL & Jong
MD. **2009**. Oystercatchers' Bill Shapes as a Proxy for Diet Specialization: More Differentiation
than Meets the Eye. *Ardea* **97**: 335–347.

Pyle P, Howell SNG, Institute for Bird Populations, & Point Reyes Bird Observatory. **1997**.
Identification guide to North American birds: a compendium of information on identifying,
ageing, and sexing 'near-passerines' and passerines in the hand. Part I, Part I,. Bolinas, Calif.:
Slate Creek Press.

R Core Team. **2020**. *R: A Language and Environment for Statistical Computing (4.0.*
3)[Computer software]. R Foundation for Statistical Computing. Retrieved from http://www. R-
proje ….

Rappole JH. **1995**. *The ecology of migrant birds: a neotropical perspective*. Washington;
London: Smithsonian institution Press.

Rappole JH, Ramos MA & Winker K. **1989**. Wintering Wood Thrush Movements and Mortality
in Southern Veracruz. *The Auk* **106**: 402–410.

Rayner JMV. **1988**. The evolution of vertebrate flight. *Biological Journal of the Linnean Society*
34: 269–287.

Reaney AM, Bouchenak-Khelladi Y, Tobias JA & Abzhanov A. **2020**. Ecological and
morphological determinants of evolutionary diversification in Darwin's finches and their relatives.
Ecology and Evolution **10**: 14020–14032.

Regosin JV & Pruett-Jones S. **2001**. Sexual Selection and Tail-Length Dimorphism in Scissor-
Tailed Flycatchers. *The Auk* **118**: 167–175.

Revell LJ. **2009**. Size-Correction and Principal Components for Interspecific Comparative
Studies. *Evolution* **63**: 3258–3268.

Revell LJ. **2012**. phytools: an R package for phylogenetic comparative biology (and other
things). *Methods in Ecology and Evolution* **3**: 217–223.

Rising JD & Somers KM. **1989**. The Measurement of Overall Body Size in Birds. *The Auk* **106**:
666–674.

Rönn JAC von, Shafer ABA & Wolf JBW. **2016**. Disruptive selection without genome-wide
evolution across a migratory divide. *Molecular Ecology* **25**: 2529–2541.

Senar JC & Pascual J. **1997**. Keel and tarsus length may provide a good predictor of avian body size. *Ardea* **85**.

Shao S, Quan Q, Cai T, Song G, Qu Y & Lei F. **2016**. Evolution of body morphology and beak shape revealed by a morphometric analysis of 14 Paridae species. *Frontiers in Zoology* **13**: 30.

Sheard C, Neate-Clegg MHC, Alioravainen N, Jones SEI, Vincent C, MacGregor HEA, Bregman TP, Claramunt S & Tobias JA. **2020**. Ecological drivers of global gradients in avian dispersal inferred from wing morphology. *Nature Communications* **11**: 2463.

Sherry TW. **1984**. Comparative Dietary Ecology of Sympatric, Insectivorous Neotropical Flycatchers (Tyrannidae). *Ecological Monographs* **54**: 313–338.

SHERRY TW. **2005**. Does winter food limit populations of migratory birds? *Birds of two worlds: the ecology and evolution of migration*: 414–425.

Snow DW. **1953**. Systematics and comparative ecology of the genus Parus in the Palaearctic region.

Snow DW. **1954**. The habitats of Eurasian tits (Parus spp.). *Ibis* **96**: 565–585.

Stiles FG. **1980**. Evolutionary implications of habitat relations between permanent and winter resident landbirds in Costa Rica. *Migrant Birds in the Neotropics (A. Keast and ES Morton, Eds.) Smithsonian Institution Press, Washington DC*: 421–435.

Svanbäck R & Schluter D. **2012**. Niche Specialization Influences Adaptive Phenotypic Plasticity in the Threespine Stickleback. *The American Naturalist* **180**: 50–59.

Swanson MT, Oliveros CH & Esselstyn JA. **2019**. A phylogenomic rodent tree reveals the repeated evolution of masseter architectures. *Proceedings of the Royal Society B: Biological Sciences* **286**: 20190672.

Tarka M, Åkesson M, Beraldi D, Hernández-Sánchez J, Hasselquist D, Bensch S & Hansson B. **2010**. A strong quantitative trait locus for wing length on chromosome 2 in a wild population of great reed warblers. *Proceedings of the Royal Society B: Biological Sciences* **277**: 2361–2369.

Tellería LJ, Blázquez M, De La Hera I & Pérez-Tris J. **2013**. Migratory and resident Blackcaps Sylvia atricapilla wintering in southern Spain show no resource partitioning. *Ibis* **155**: 750–761.

Tellería JL & Carbonell R. **1999**. Morphometric Variation of Five Iberian Blackcap Sylvia atricapilla Populations. *Journal of Avian Biology* **30**: 63–71.

Thiollay JM. **1988**. Comparative Foraging Success of Insectivorous Birds in Tropical and Temperate Forests: Ecological Implications. *Oikos* **53**: 17–30.

Vágási CI, Pap PL, Vincze O, Osváth G, Erritzøe J & Møller AP. **2016**. Morphological Adaptations to Migration in Birds. *Evolutionary Biology* **43**: 48–59.

Van Valen L. **1965**. Morphological Variation and Width of Ecological Niche. *The Amercian Naturalist* **99**: 377–390.

Voelker G. **2001**. Morphological correlates of migratory distance and flight display in the avian
genus Anthus. *Biological Journal of the Linnean Society* **73**: 425–435.

Waide RB. **1980**. Resource partitioning between migrant and resident birds: the use of irregular
resources. *Migrant birds in the Neotropics: ecology, behavior, distribution, and conservation*:
337–352.

Wang X & Clarke JA. **2015**. The evolution of avian wing shape and previously unrecognized
trends in covert feathering. *Proceedings of the Royal Society B: Biological Sciences* **282**:
20151935.

Wiedenfeld DA. **1991**. Geographical Morphology of Male Yellow Warblers. *The Condor* **93**:
712–723.

Willis EO. **1974**. Populations and Local Extinctions of Birds on Barro Colorado Island, Panama.
Ecological Monographs **44**: 153–169.

Wilson EO. **1961**. The Nature of the Taxon Cycle in the Melanesian Ant Fauna. *The American
Naturalist* **95**: 169–193.

Winkler H & Leisler B. **1992**. On the ecomorphology of migrants. *Ibis* **134**: 21–28.

Winquist T & Lemon RE. **1994**. Sexual Selection and Exaggerated Male Tail Length in Birds.
The American Naturalist **143**: 95–116.

**Závorka L, Larranaga N, Lovén Wallerius M, Näslund J, Koeck B, Wengström N,
Cucherousset J & Johnsson JI**. **2020**. Within-stream phenotypic divergence in head shape of
brown trout associated with invasive brook trout. *Biological Journal of the Linnean Society* **129**:
347–355.

Supporting Information

628 Supporting Information 1 – Supplementary Results

629 Supporting Information 2 – ImageJ protocol for measuring long *Tyrannus* tail lengths

630 Supporting Information 3 – Sex-specific results: averages for each morphometric, phylogenetic

631 principal components analysis (PPCA), and phylogenetic ANOVA