Morphology of migration: Associations between wing, and bill morphology and migration in kingbirds (*Tyrannus*)

Short Title: Ecomorphology of migration in kingbirds (*Tyrannus*)

Authors & Affiliations
Maggie P. MacPherson [1,2*], Alex E. Jahn [3,4], Nicholas A. Mason [2]
[1] Department of Ecology and Evolutionary Biology, Tulane University, New Orleans, LA 70118
[2] Louisiana State University Museum of Natural Sciences, Louisiana State University, Baton Rouge, LA 70803
[3] Departamento de Zoologia, Universidade Estadual Paulista, Av. 24a No. 1515, Rio Claro, Brazil
[4] Environmental Resilience Institute, Indiana University, 717 E 8th St., Bloomington, IN 47408
[*]Corresponding author: Maggie P. MacPherson, E-mail: maggie.macpherson@gmail.com

Abstract

Morphology is closely linked to locomotion and diet in animals. In animals that undertake long-distance migrations, limb-morphology is under selection to maximize mobility and minimize energy expenditure. Migratory behaviors also interact with diet, such that migratory animals tend to be dietary generalists, while sedentary taxa tend to be dietary specialists. Despite a hypothesized link between migration status and morphology, phylogenetic comparative studies have yielded conflicting findings. We tested for evolutionary associations between migratory status and limb and bill morphology across kingbirds, a pan-American genus of birds with migratory, partially migratory, and sedentary taxa. Migratory kingbirds had longer wings, in agreement with expectations if selection favors improved aerodynamics for long-distance migration. We also found an association between migratory status and bill shape, such that more migratory taxa had wider, deeper, and shorter bills compared to sedentary taxa. However, there was no difference in intraspecific morphological variation among migrants, partial migrants, and residents, suggesting that dietary specialization has evolved independently of migration strategy. The evolutionary links between migration, diet, and morphology in kingbirds uncovered here further strengthen ecomorphological associations that underlie long-distance seasonal movements in animals.

Keywords: ecomorphology, flycatcher, migration, movement ecology, natural selection, partial migration, Tyrannidae

Introduction

Animal movement is linked to morphology at various taxonomic scales. At a macroevolutionary scale, streamlined or aerodynamic body shapes have been associated with the evolution of

long-distance migration in fish (Chapman *et al.*, 2015), insects (Johansson, Söderquist, &

Bokma, 2009), and birds (Fiedler, 2005). Within species, migratory distance has also been

associated with streamlined body shapes in fish (e.g., Crossin *et al.*, 2004) and aerodynamic

shapes in birds (Voelker, 2001; Minias *et al.*, 2015; Vágási *et al.*, 2016). While certain taxa with

long-distance movements exhibit strong selection for energy-efficient body shapes, this is not

universal (Mulvihill & Chandler, 1990; Mönkkönen, 1995; Wang & Clarke, 2015). Thus, there is

a persistent need to expand the taxonomic breadth of studies linking migration and morphology

to better understand which lineages exhibit migration-morphology associations, and why these

associations vary among taxa.

Morphology is also shaped by foraging strategies. For example, dietary niche is

associated with head and body shape in fish (Knudsen *et al.*, 2011; Závorka *et al.*, 2020), birds

(Felice *et al.*, 2019), and mammals (Swanson, Oliveros, & Esselstyn, 2019). Additionally,

phenotypic plasticity is expected in dietary generalists, as shown in comparative common-

garden experiments in stickleback minnows (Svanbäck & Schluter, 2012). In populations

recently released from interspecific competition, such as island colonizers (Wilson, 1961; Clegg

& Owens, 2002), phenotypic plasticity is thought to support increased morphological variation to

limit intraspecific competition (i.e., 'niche variation hypothesis', Van Valen, 1965). Migratory

lineages of dietary generalists may have strong preferences for food resources that are easiest

to access (Sherry, 1984; Levey & Stiles, 1992; Bell, 2011) or that are superabundant (Moreau,

1952; Morse, 1971; Willis, 1974). In contrast, sedentary individuals may mitigate intra- and

interspecific competition by seeking temporally stable food resources in microhabitats that are

buffered from environmental fluctuation (e.g., temperature fluctuations; Bell, 2011).

Comparisons across avian species (Levey & Stiles, 1992) and families (Chesser & Levey, 1998)

have shown migratory taxa that forage on seasonally variable resources tend to exhibit more

morphological variation, presumably associated with opportunistic foraging across a wider

dietary breadth among or within individuals (Bell, 2011).

Testing how migration shapes morphological variation among taxa requires a

phylogenetic comparative framework with comprehensive inter- and intraspecific sampling.

Studies comparing migratory to sedentary birds support that long-distance migration favors

longer (Rayner, 1988; Wiedenfeld, 1991; Pérez-Tris & Tellería, 2001; Förschler & Bairlein,

2011; Tellería *et al.*, 2013), and more pointed (Carvalho Provinciato, Araújo, & Jahn, 2018;

Gómez-Bahamón *et al.*, 2020b) wings for increased aerodynamics. Hypotheses that link bill

morphology and migratory status are less well-studied, but hinge on differences in diet that

covary with migratory status (Bell, 2011; but see Cox, 1968; Herrera, 1978; Leisler, 1990).

Sedentary insectivorous taxa often have longer bills, presumably to improve closing speed for

capturing highly mobile prey (Leisler, 1990). In contrast, migratory taxa may have shorter bills

for capturing slow-moving prey like caterpillars to feed young during the breeding season and

small, abundant arthropods during winter (Thiollay, 1988; Leisler, 1990). Furthermore, studies

on bill size in relation to foraging strategies disagree, such that both narrow and wide bills are

associated with generalist diets (Tellería & Carbonell, 1999; Tellería *et al.*, 2013), and bill size

differs between isolated specialist populations of the same species (Alonso *et al.*, 2020).

Early assessments of morphological variation among migratory and sedentary avian

species did not account for their shared evolutionary history (Cox, 1968, 1985; Leisler, 1990).

Boyle and Conway (2007) advanced this approach by performing phylogenetically independent

contrasts to address selective pressures for the evolution of migration between species (see

also Gómez *et al.*, 2016; Vágási *et al.*, 2016). However, the literature on the role of migratory

status in shaping morphological phenotypes in birds draws predominantly from comparisons

among distantly related taxa (e.g., Rappole's 1995; critique of Herrera 1978). Thus, detailed

comparisons among taxa that differ in migratory status within a modern phylogenetic

comparative framework are necessary for a more comprehensive test of how migration is

associated with morphology at different taxonomic scales (e.g., Bolnick *et al.*, 2007).

We used phylogenetic comparative methods to test for ecomorphological associations in

wing and bill morphology among migratory, partially migratory, and sedentary kingbirds

(*Tyrannus*; Fitzpatrick *et al.*, 2004). Kingbirds are flycatchers (Tyrannidae) with considerable

variation in migration status and morphology within and among species, as well as a rich body

of literature linking ecology and morphology (Fitzpatrick & Schauensee, 1980; Fitzpatrick, 1981;

Sherry, 1984; Fitzpatrick, 1985; Cintra, 1997; Fitzpatrick *et al.*, 2004; Gabriel & Pizo, 2005;

Carvalho Provinciato *et al.*, 2018; Gómez-Bahamón *et al.*, 2020b). As aerodynamic theory

predicts that longer, more pointed wings and shorter tails are more efficient for long-distance

migratory flights (Norberg, 1995; Pennycuick, 2008), we expected migratory taxa to have longer

and more pointed wings compared to sedentary taxa (Kipp, 1942, 1958; Winkler & Leisler,

1992; Mönkkönen, 1995 and references therein; Lockwood, Swaddle, & Rayner, 1998). Tails of

long-distance migrants should also be shorter than more sedentary individuals to reduce drag

during long-distance flights (Rayner, 1988; Norberg, 1990; Winkler & Leisler, 1992; Förschler &

Bairlein, 2011). However, tails are targets of sexual selection (Winquist & Lemon, 1994; Mobley,

2002), and tail lengths are often unassociated with migration strategy (e.g., Voelker, 2001; Neto

et al., 2013). Within-population variation in flight capability could be associated with migration

(Fernández & Lank, 2007) or foraging ecology (Hromada & Tryjanowski, 2003), so we also

tested whether more migratory taxa have more variable wing morphology compared to less

migratory taxa.

Bill morphology is closely tied to foraging niche (Snow, 1953, 1954; Lack, 1971) and bill

size and shape is typically related to broad dietary categories in birds (e.g., >50% insectivorous,

granivorous, frugivorous etc.; e.g., Reaney *et al.*, 2020). We therefore predicted that there would

be no relationship between migratory status and bill size because all kingbirds are primarily

insectivorous. Additionally, migration may impart an 'ecological release' associated with the

evolution of a more variable, opportunistic diet (Bell, 2011). We thus predicted that more

migratory taxa will have higher coefficients of variation in bill morphology compared to less

migratory taxa.

Methods

Phylogeny construction

We extracted the clade of 15 kingbird taxa from the Harvey et al. (2020) suboscine phylogeny to

construct a kingbird phylogeny that includes 28 operational taxonomic units (OTUs). Harvey et

al. (2020) included all species (13/13) but only 15% of the subspecific diversity (3/20). We

therefore added *T. savana* taxa following phylogenetic relationships and branch lengths

estimated by Gómez-Bahamón (2020a). The remaining subspecies were added as polytomies

assuming a most recent common ancestor (MRCA) of 0.5 million years from members of their

species group (Figure 1A). To test the sensitivity of analyses to this assumption, we tested a

range of different dates for the MRCA for the added subspecies, which did not change the

results (Supplementary Information 1).

Figure 1. Ancestral state reconstruction of migration ecology category and wing chord length, as well as ordination of 28 kingbird OTUs for bill (length, width, depth) and feather (wing chord length, tail length, and Kipp's distance). A) Kingbird phylogeny showing ancestral state reconstruction of migration ecology strategy (pie charts at each node), and wing chord length (branch greyscale). B-D) PPCA plots showing individuals color-coded to species (identified at the tip labels of the phylogeny), shape-coded to subspecies, and their status as migratory, partially migratory, or sedentary is distinguished via the shape outline. Pie charts for each node in the phylogeny show the ancestral state reconstruction of migration type. E-H) photos of representative kingbird taxa with corresponding color- and shape-coded points in the top right of

each photo. Taxa are as follows: E) *Tyrannus savana savana* (photo credit: Rodrigo Conte), F)
Tyrannus caudifasciatus caudifasciatus (photo credit: Yeray Seminario / Whitehawk), G)
Tyrannus crassirostris pompalis (photo credit: Martin Molina), and H) *Tyrannus cubensis* (photo
credit: Dubi Shapiro).

Morphological measurements

We measured bill length, width, and depth at the distal end of the nares. We also

measured unflattened wing chord length, Kipp's distance (a measure of wing pointedness: the

distance between the tip of the first secondary feather to the tip of the longest primary feather;

Kipp, 1942, 1958; Baldwin *et al.*, 2010), tail length, and tarsus length on 2108 study skins from

across the ranges of each species and subspecies (28 operational taxonomic units (OTUs);

Figure 1A; Table S1.1). MM measured 2008 specimens, and identified all individuals to the

lowest level of taxonomic identification (using Clements *et al.*, 2019) and classified each

individual as migratory, partially migratory, or sedentary (*sensu* Fitzpatrick *et al.*, 2004). Partially

migratory taxa are those in which individuals vary in migratory tendency (Boyle, 2008). JIGA

(see Acknowledgements) measured 100 specimens to improve sampling of some taxa. Bill and

tail measurements were taken twice and were confirmed to be within one mm of one another.

Both right and left wing chords and tarsus lengths were measured and averaged.

Measurements by JIGA were only taken once. We measured tail length as the longest rectrix to

the nearest 0.1 cm (Pyle *et al.*, 1997). For most measurements, we used a Mitutoyo brand IP 67

digital calipers (part number 573-271) with a range of up to 15.24 cm, with 0.00127 cm

resolution. For tails longer than 15.24 cm, we used a 30.48 cm stainless steel ruler placed

between the two middle rectrices. When tails were longer than 30.48 cm photos were taken of

the tails above a 0.64 x 0.64 cm square grid with the calipers measuring to their extent and

ImageJ was used to calculate the full tail length (Supporting Information 2). We then averaged

measurements across individuals within each OTU for downstream analyses. To test

associations between migratory strategy and intraspecific morphological variation, we also

calculated the coefficient of variation (mean/standard deviation) for each character and each

taxon. This provides a scaled measure of variance for each character and allows testing

whether migratory strategies differ in the amount of intraspecific variation.

We tested whether accounting for age (juvenile versus adult) and sex (female versus

male) classes improved linear models explaining morphological measurements among taxa. We

did this using the mulTree function of the mulTree package (Guillerme & Healy, 2020) in R,

which assesses intraspecific variation while accounting for phylogenetic relatedness (e.g.,

Nations *et al.*, 2019). We found that including age class improved model fit for all characters

except tarsus length and including sex class improved model fit for all characters except bill

length and tarsus length (Supporting Information 1). We subsequently omitted juveniles from our

analyses, but presented the results of both sexes combined in the main text because our results

did not differ when analyzing sexes independently (Supporting Information 3).

Phylogenetic Principal Components Analysis

We conducted a phylogenetic principal component analysis (PPCA) for bill measurements using

the phyl.pca function from the phytools package (Revell, 2009) because this gives information

on bill volume and shape that is not reflected in individual bill measurements (Table 1). Bill

PPCA scores were included in the following phylogenetic analysis of variance (see next

section). PPCA was also conducted for sexes separately (Supporting Information 3).

Table 1. Results of phylogenetic principal component analysis (PPCA) of bill measurements.

Percent variance explained by each eigenvector is in brackets for each principal component.

Loadings	Bill PPC1 (94.52%)	Bill PPC2 (4.47%)	lambda
Bill length	0.99	0.14	0.95
Bill width	0.93	-0.33	
Bill depth	0.94	-0.27	

Phylogenetic Analysis of Variance

We compared morphology between migratory, partially migratory, and sedentary taxa via a

phylogenetic analysis of variance (Garland *et al.*, 1993) with the phylANOVA function in the

phytools package (Revell, 2012). As differences in body size can account for much of the

variation among species (Albrecht, Gelvin, & Hartman, 1993; McCoy *et al.*, 2006; Revell, 2009;

Berner, 2011), we extracted phylogenetic residuals (Revell, 2009) for each dependent variable

(bill length, bill width, bill depth, wing chord, Kipp's distance, and tail length) with tarsus length

as an approximation of body size and the independent variable. We opted to use tarsus instead

of mass to adjust for body size (Rising & Somers, 1989; Senar & Pascual, 1997) because mass

can change seasonally (particularly in migratory birds; Lindström & Piersma, 1993) and also

varies by sex in some kingbirds (e.g., *T. melancholicus*, Jahn *et al.*, 2010; *T. savana*, Carvalho

Provinciato *et al.*, 2018; *T. tyrannus*, Murphy, 2007). To additionally support this decision, we

used the mulTree function of the mulTree package to test whether tarsus length had the highest

correlation among measurements while accounting for phylogeny, finding that tarsus length the

highest correlation with body mass among all variables for the individuals that had mass data

(Supporting Information 1). We then conducted a phylogenetic ANOVA for each character using

the residual values (with 1000 replicates) and reported mean P-values for pairwise, post-hoc

comparisons of mean values and coefficients of variation between migration categories. As

such, we conducted 51 tests of statistical significance, which raises the issue of multiple

hypothesis testing (Shaffer, 1995). Opinions differ on whether or not to adjust P-values when

testing multiple hypotheses (Curran-Everett, 2000) and the 0.05 conventional threshold is

ultimately arbitrary (Wasserstein, Schirm, & Lazar, 2019). Nonetheless, we report both

uncorrected P-values and Bonferroni-corrected P-values (n = 3 tests for each set of pairwise

comparisons of means or coefficients of variation). We also performed ancestral state

reconstructions of wing chord length with the contMap function in the phytools package (Revell,

2012) and of migratory status with the fit.mle function in the diversitree package (FitzJohn,

2012). All statistical analyses were performed using program R version 4.0.3 (R Core Team,

2020).

203   Results

The mean values for each morphometric character +/- standard deviation for all adults in each

OTU are reported in Table S1.1. We found that migratory taxa had longer wing chords (Figure

2F, Table 2) and pointier wings than sedentary taxa (Figure 2D, Table 2). However, we found no

association between migratory status and coefficients of variation in bill morphology (Figure 2G-

I, Table 2). In agreement with our prediction, we found no relationship between migratory status

and bill length, width, depth, or bill PPC1 (Figure 2A-C, Figure 3A, Table 2). However, migratory

taxa had shorter, wider, and deeper bills (bill PPC2) compared to sedentary taxa, which had

longer, shallower, and more narrow bills (Figure 3B, Table 2).

Figure 2. Phylogenetic ANOVA results comparing residuals of bill and feather morphometrics
(A-F), and coefficient of variation in bill and feather morphometrics (G-L) across migratory,
partially migratory, and sedentary kingbird OTUs. Note: although Kipp's distance was
significantly different between migratory and sedentary taxa for all adults, this was not
significantly different after Bonferroni correction when males were assessed separately
(Supporting Information Table 3.6). Letters above each barplot show groupings resulting from
post-hoc pairwise comparisons.

Table 2. T and P values from phylogenetic ANOVA analysis of adult *Tyrannus* flycatchers.

Values shown in brackets are P values with Bonferroni adjustment (n = 3) to account for

multiple hypothesis testing. Significant results are in bold. Model F and Pr(>F) values can be

found in Table S1.4.

Morphometric	**migratory v. partially migratory**	**migratory v. sedentary**	**partially migratory v. sedentary**

	T	P (adj. p)	T	P (adj. p)	T	P (adj. p)
Bill length	0.32	0.73 (1.00)	0.06	0.96 (1.00)	0.30	0.82 (1.00)
CV Bill length	0.39	0.72 (1.00)	1.90	0.23 (0.66)	2.04	0.15 (0.39)
Bill width	0.20	0.82 (1.00)	1.68	0.30 (0.86)	1.65	0.24 (0.74)
CV Bill width	1.01	0.29 (0.99)	2.49	0.10 (0.33)	1.00	0.49 (1.00)
Bill depth	0.50	0.60 (1.00)	2.17	0.17 (0.49)	1.29	0.40 (1.00)
CV Bill depth	0.74	0.44 (1.00)	0.34	0.85 (1.00)	1.10	0.45 (1.00)
Bill PC1 (size)	0.07	0.94 (1.00)	0.62	0.70 (1.00)	0.45	0.74 (1.00)
CV Bill PC1 (size)	0.03	0.99 (1.00)	0.073	0.96 (1.00)	0.096	0.95 (1.00)
Bill PC2 (shape)	0.64	0.53 (1.00)	3.91	**0.01 (0.03)**	2.60	0.06 (0.18)
CV Bill PC2 (shape)	0.19	0.85 (1.00)	1.32	0.40 (1.00)	0.91	0.50 (1.00)
Kipp's distance	2.11	**0.05** (0.11)	3.73	**0.02 (0.04)**	0.83	0.57 (1.00)
CV Kipp's distance	0.71	0.43 (1.00)	0.79	0.62 (1.00)	0.12	0.94 (1.00)
Wing chord length	0.71	0.48 (1.00)	3.78	**0.02 (0.03)**	2.42	0.09 (0.20)
CV Wing chord length	0.56	0.54 (1.00)	0.28	0.87 (1.00)	0.39	0.78 (1.00)
Tail length	0.93	0.32 (1.00)	0.34	0.84 (1.00)	1.32	0.35 (1.00)
CV Tail length	1.15	0.23 (0.76)	0.35	0.83 (1.00)	0.98	0.49 (1.00)
CV Tarsus length	0.29	0.78 (1.00)	2.81	0.06 (0.16)	2.71	**0.04** (0.16)

Figure 3. Phylogenetic ANOVA results comparing bill phylogenetic principal component (PPC) scores across For Peer Review migratory, partially migratory, and sedentary kingbird OTUs. Letters above each barplot show groupings resulting from post-hoc pairwise comparisons

Discussion

We found evidence of an association between migratory status and wing morphology in the kingbirds. This corroborates the idea that migratory birds have evolved morphological features that help in sustained, long-distance flight, such as longer wings (Winkler & Leisler, 1992; Milá, Wayne, & Smith, 2008; Neto *et al.*, 2013; Carvalho Provinciato *et al.*, 2018). There is a strong genetic component determining wing length (Böhning-Gaese & Oberrath, 1999; Tarka *et al.*, 2010), and among migratory taxa, wing length is positively associated with migratory distance (Förschler & Bairlein, 2011; Rönn, Shafer, & Wolf, 2016; Vágási *et al.*, 2016; Carvalho Provinciato *et al.*, 2018). Partial migration, wherein only some individuals migrate, is thought to be an evolutionary stepping-stone between sedentarism and obligate migration (Berthold, 1999; Bell, 2000; Chapman *et al.*, 2011). However, partial migration can also be sex and age-class dependent, as is the case for *T. melancholicus melancholicus* (Jahn *et al.*, 2010). Longer wings may allow for more flexibility in altering migration speeds according to conditions experienced

during migration (Hahn *et al.*, 2016), but there may be ecological limits and trade-offs imposed

on wing length by other selective pressures, like shorter wings for aerial agility among arboreal

species (Sheard *et al.*, 2020), for evading predators (Fernández & Lank, 2007), or for hovering

flight (Marchetti, Price, & Richman, 1995).

Our finding of differences in bill shape between migratory and sedentary taxa suggests

that bill shape is a potential target of selection pressures related to migration ecology. However,

our comparisons of coefficients of variation did not support intraspecific competition as a driver

of diversifying selection in bill morphology among generalist migratory kingbirds. The ratio of bill

length/bill width holds potential significance in foraging behaviors in other tyrannids (Botero-

Delgadillo & Bayly, 2012), but our findings disagree with other research across the Tyrannidae

describing generalists as having intermediate bill morphologies (Fitzpatrick, 1985). We found

that sedentary taxa had longer, narrower, and shallower bills compared to migratory taxa

(Figure 1B; Table 1). Previous work has shown that directional selection increasing bill length in

dietary specialists may be driven by interspecific competition, including from closely related taxa

(Freed, Medeiros, & Cann, 2016). Thus, other selective pressures may be acting upon the bill

shape and size. For example, bill morphology has been linked with thermoregulation in dietary

generalists, and bill width and depth are adaptive for dissipating heat during migration for

improved thermoregulation (Danner & Greenberg, 2015; Friedman *et al.*, 2017; Danner *et al.*,

2017). If convergence on bill shape is driven mainly by diet, future research integrating

multivariate or nonlinear bill morphometrics (e.g., hooked bills, lateral and longitudinal bill

curvature, bill surface area; Shao *et al.*, 2016) with degrees of intraguild dietary specialization

and foraging behaviors would broaden our understanding of selection mechanisms shaping bill

phenotypes.

The range of phenotypes shaped by migratory status is likely to be more varied than has

been historically detectable using linear measurements of morphology alone. Divergence in bill

shape between migratory and sedentary taxa may define an important component of the suite of

co-adaptations for migratory life histories (Bell, 2000). Morphology may be driven by selection

related to migration itself, such as an increased need for heat dissipation or conservation in

flight(see above). Alternatively, morphology may be shaped by a more complex competitive

landscape that affects migratory and sedentary taxa differently. In taxa occupying tropical

regions, like many kingbirds, the dry season is when both intra- and interspecific competition

increase due to 'winter food limitation' and competition with migrants returning to wintering

grounds, and competition with juvenile conspecifics (Hespenheide, 1975; Emlen, 1977;

Orejuela, Raitt, & Álvarez, 1980; Stiles, 1980; Waide, 1980; Rappole, Ramos, & Winker, 1989;

Rappole, 1995; Sherry, 2005; Danner *et al.*, 2013). Selection may favor longer bills among

sedentary taxa to resolve both intra- and interspecific conflict (Table 1, Figure 3B), allowing

access to more temporally stable food resources in microhabitats buffered from spatial and

temporal environmental fluctuations (Bell, 2011).

Migratory and sedentary taxa may not differ in linear, individual bill metrics if a

combination of characteristics (e.g., Table 1, Figure 3B) support a more generalist foraging

strategy. For example, specific bill shapes could accommodate spatiotemporal changes in the

competitive landscape (Navalón *et al.*, 2019) or provide increased stability for prey capture in

perch-gleaning or hover-gleaning foraging techniques (Fitzpatrick, 1985; Fitzpatrick *et al.*, 2004;

Botero-Delgadillo, 2011; Botero-Delgadillo & Bayly, 2012). The diversity of habitats that

migratory taxa encounter during their annual cycle may shape phenotypes in more complex

ways than previously thought, and techniques that incorporate more advanced characterizations

of morphological variation may expose novel insight into how movement life history strategies

shape phenotypes (e.g., Pol *et al.*, 2009; Mallarino *et al.*, 2011; Navalón *et al.*, 2019; Alonso *et*

al., 2020; Medina *et al.*, 2020).

Aspects of both wing and bill morphology appear to have evolved in association with

migratory status in kingbirds that have migratory, partially migratory, and sedentary taxa. Our

results suggest that migratory status has shaped wing morphology in a widespread avian

genus, and that multivariate bill shape metrics may differ between sedentary and migratory

lineages. Thus, adaptive phenotypes may be related to migratory status in more complex ways

than previously understood. Assessments of the various mechanisms driving patterns in bill

shape (e.g., heat dissipation, foraging strategy, and competitive landscape) across a broader

range of taxonomic groups that differ in migration strategies would complement our refinement

of the morphology of migration in kingbirds.

304   **Data Availability**

305   The data underlying the work, and the R code for all statistical analyses and results shared in

306   tables and figures is available from the following public Github repository

307   (https://github.com/mmacphe/Tyrannus_morphology). The voucher table and measurements of

308   all individuals will also be made available at Dryad upon publication.

310   **Acknowledgements**

This manuscript has been Accepted at the Biological Journal of the Linnean Society (August 02,

2021). We thank José Ignacio ("Nacho") Giraldo Arango for measuring specimens, and Brenna

and Michael Wells, and Shelbey Hearn for their assistance measuring specimens used in this

study. Ben Marks, Shannon Hacket, and John Bates from the Field Museum of Natural History,

Bradley Millen, and Allan Baker from the Royal Ontario Museum, Gary Stiles from the Instituto

de Ciencias Naturales, Universidad Nacional de Colombia, Jacob Saucier, Helen James, and

Gary Graves from the Smithsonian National Museum of Natural History, and Steven Cardiff, and

Van Remsen from the Louisiana State University Museum of Natural Science provided

assistance in accessing the specimens used in this study. Thanks to Robb Brumfield and

Michael Harvey for providing an early-access version of the suboscine phylogeny for our study.

We also thank Calandra Stanley, Caleb McMahan, the Toronto MacPhersons and the Bogotá

Acevedos for their special help aiding the lead author to visit collections. Travel to measure

specimens was supported in part from a scholar award from the James S. McDonnell

Foundation. Samantha Rutledge, David Vander Pluym, and Subir Shakya provided helpful

feedback on an earlier version of this manuscript, and we thank an anonymous reviewer and

Prof John A. Allen for their insightful peer review. We wish to acknowledge www.gbif.org for

providing the ROM database (accessed September 30, 2020) and the Field Museum database

(accessed October 20, 2020) online that was used to validate locality data previously recorded

by hand. Amethys E'etessam, Clare Lister, and Trey Hendrix also helped with georeferencing

from locality data that was a part of this project.

Literature Cited

Albrecht GH, Gelvin BR & Hartman SE. **1993**. Ratios as a size adjustment in morphometrics.
American Journal of Physical Anthropology **91**: 441–468.
Alonso D, Fernández-Eslava B, Edelaar P & Arizaga J. **2020**. Morphological divergence
among Spanish Common Crossbill populations and adaptations to different pine species. *Ibis*
162: 1279–1291.
Bartoń K. **2020**. *MuMIn: multi-model inference*.
Bell C. **2000**. Process in the evolution of bird migration and pattern in avian ecogeography.
Journal of Avian Biology **31**: 258–265.
Bell C. **2011**. Resource buffering and the evolution of bird migration. *Evolutionary Ecology* **25**:
91–106.
Berner D. **2011**. Size correction in biology: how reliable are approaches based on (common)
principal component analysis? *Oecologia* **166**: 961–971.
Berthold P. **1999**. A comprehensive theory for the evolution, control and adaptability of avian
migration. *Ostrich* **70**: 1–11.
Böhning-Gaese K & Oberrath R. **1999**. Phylogenetic effects on morphological, life-history,
behavioural and ecological traits of birds. : 18.

Bolnick DI, Svanbäck R, Araújo MS & Persson L. **2007**. Comparative support for the niche
variation hypothesis that more generalized populations also are more heterogeneous.
Proceedings of the National Academy of Sciences **104**: 10075–10079.
Botero-Delgadillo E. **2011**. Cuantificando el comportamiento: Estrategias de búsqueda y
ecología de forrajeo de 12 especies sintópicas de Atrapamoscas (Tyrannidae) en la Sierra
Nevada de Santa Marta, Colombia. *Revista Brasileira de Ornitologia* **19**: 343–357.
Botero-Delgadillo E & Bayly NJ. **2012**. Does morphology predict behavior? Correspondence
between behavioral and morphometric data in a Tyrant-flycatcher (Tyrannidae) assemblage in
the Santa Marta Mountains, Colombia. *Journal of Field Ornithology* **83**: 329–342.
Boyle WA. **2008**. Partial migration in birds: tests of three hypotheses in a tropical lekking
frugivore. *Journal of Animal Ecology* **77**: 1122–1128.
Boyle WA & Conway CJ. **2007**. Why Migrate? A Test of the Evolutionary Precursor
Hypothesis. *The American Naturalist* **169**: 344–359.
Carvalho Provinciato IC, Araújo MS & Jahn AE. **2018**. Drivers of wing shape in a widespread
Neotropical bird: a dual role of sex-specific and migration-related functions. *Evolutionary*
Ecology **32**: 379–393.
Chapman BB, Brönmark C, Nilsson JÅ & Hansson LA. **2011**. The ecology and evolution of
partial migration. *Oikos* **120**: 1764–1775.
Chapman BB, Hulthén K, Brönmark C, Nilsson PA, Skov C, Hansson LA & Brodersen J.
2015. Shape up or ship out: migratory behaviour predicts morphology across spatial scale in a
freshwater fish. *Journal of Animal Ecology* **84**: 1187–1193.
Chesser RT & Levey DJ. **1998**. Austral Migrants and the Evolution of Migration in New World
Birds: Diet, Habitat, and Migration Revisited. *The American Naturalist* **152**: 311–319.
Cintra R. **1997**. Spatial Distribution and Foraging Tactics of Tyrant Flycatchers in Two Habitats
in the Brazilian Amazon. *Studies on Neotropical Fauna and Environment* **32**: 17–27.
Clegg SM & Owens PF. **2002**. The 'island rule' in birds: medium body size and its ecological
explanation. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **269**:
1359–1365.
Clements JF, Schulenberg TS, Iliff MJ, Billerman SM, Fredericks TA, Sullivan BL & Wood
CL. **2019**. The eBird/Clements Checklist of Birds of the World: v2019.
Cooney CR, Bright JA, Capp EJR, Chira AM, Hughes EC, Moody CJA, Nouri LO, Varley
ZK & Thomas GH. **2017**. Mega-evolutionary dynamics of the adaptive radiation of birds. *Nature*
542: 344–347.
Cox GW. **1968**. The Role of Competition in the Evolution of Migration. *Evolution* **22**: 180–192.
Cox GW. **1985**. The Evolution of Avian Migration Systems between Temperate and Tropical
Regions of the New World. *The American Naturalist* **126**: 451–474.
Crossin GT, Hinch SG, Farrell AP, Higgs DA, Lotto AG, Oakes JD & Healey MC. **2004**.
Energetics and morphology of sockeye salmon: effects of upriver migratory distance and
elevation. *Journal of Fish Biology* **65**: 788–810.
Danner RM, Greenberg RS, Danner JE, Kirkpatrick LT & Walters JR. **2013**. Experimental
support for food limitation of a short-distance migratory bird wintering in the temperate zone.
Ecology **94**: 2803–2816.
Danner RM, Gulson-Castillo ER, James HF, Dzielski SA, Frank DC III, Sibbald ET &
Winkler DW. **2017**. Habitat-specific divergence of air conditioning structures in bird bills. *The*
Auk **134**: 65–75.
Danner RM & Greenberg R. **2015**. A critical season approach to Allen's rule: Bill size declines
with winter temperature in a cold temperate environment. *Journal of Biogeography* **42**: 114–120.
Emlen JT. **1977**. Land Bird Communities of Grand Bahama Island: The Structure and Dynamics
of an Avifauna. *Ornithological Monographs*: iii–129.
Felice RN, Tobias JA, Pigot AL & Goswami A. **2019**. Dietary niche and the evolution of
cranial morphology in birds. *Proceedings of the Royal Society B: Biological Sciences* **286**:

20182677.
Fernández G & Lank DB. **2007**. Variation in the Wing Morphology of Western Sandpipers
(Calidris Mauri) in Relation to Sex, Age Class, and Annual Cycle. *The Auk* **124**: 1037–1046.
Fiedler W. **2005**. Ecomorphology of the External Flight Apparatus of Blackcaps (Sylvia
atricapilla) with Different Migration Behavior. *Annals of the New York Academy of Sciences*
1046: 253–63.
FitzJohn RG. **2012**. Diversitree: comparative phylogenetic analyses of diversification in R.
Methods in Ecology and Evolution **3**: 1084–1092.
Fitzpatrick JW. **1981**. Search strategies of tyrant flycatchers. *Animal Behaviour* **29**: 810–821.
Fitzpatrick JW. **1985**. Form, foraging behavior, and adaptive radiation in the Tyrannidae BT -
Neo-Tropical Ornithology Ornithological Monographs No. 26. In: *Neo-Tropical Ornithology*
Ornithological Monographs No. 26., 447–470.
Fitzpatrick JW, Bates JM, Bostwick KS, Caballero IC, Clock BM, Farnsworth A, Hosner
PA, Joseph L, Langham GM & Lebbin DJ. **2004**. Family Tyrannidae (tyrant-flycatchers).
Handbook of the birds of the world **9**: 170–462.
Fitzpatrick W & Schauensee D. **1980**. Foraging behavior of Tyrant flycatchers. : 43–57.
Förschler MI & Bairlein F. **2011**. Morphological Shifts of the External Flight Apparatus across
the Range of a Passerine (Northern Wheatear) with Diverging Migratory Behaviour. *PLOS ONE*
6: e18732.
Freed LA, Medeiros MCI & Cann RL. **2016**. Multiple Reversals of Bill Length over 1.7 Million
Years in a Hawaiian Bird Lineage. *The American Naturalist* **187**: 363–371.
Friedman NR, Harmáčková L, Economo EP & Remeš V. **2017**. Smaller beaks for colder
winters: Thermoregulation drives beak size evolution in Australasian songbirds. *Evolution* **71**:
2120–2129.
Gabriel V de A & Pizo MA. **2005**. Foraging behavior of tyrant flycatchers (Aves, Tyrannidae) in
Brazil. *Revista Brasileira de Zoologia* **22**: 1072–1077.
Garland T, Dickerman AW, Janis CM & Jones JA. **1993**. phylogenetic analysis of covariance
by computer simulation. *Systematic Biology* **42**: 265–292.
Gómez C, Tenorio EA, Montoya P & Cadena CD. **2016**. Niche-tracking migrants and niche-
switching residents: evolution of climatic niches in New World warblers (Parulidae). *Proceedings*
of the Royal Society B: Biological Sciences **283**: 20152458.
Gómez-Bahamón V, Márquez R, Jahn AE, Miyaki CY, Tuero DT, Laverde-R O, Restrepo S
& Cadena CD. **2020a**. Speciation Associated with Shifts in Migratory Behavior in an Avian
Radiation. *Current Biology* **30**: 1312-1321.e6.
Gómez-Bahamón V, Tuero DT, Castaño MI, Jahn AE, Bates JM & Clark CJ. **2020b**.
Sonations in Migratory and Non-migratory Fork-tailed Flycatchers (*Tyrannus savana*).
Integrative and Comparative Biology **60**: 1147–1159.
Guillerme T & Healy K. **2020**. *mulTree: performs MCMCglmm on Multiple Phylogenetic Trees*.
Haberman K, MacKenzie DI & Rising JD. **1991**. Geographic Variation in the Gray Kingbird
(Variación geográfica en Tyrannus dominicensis). *Journal of Field Ornithology* **62**: 117–131.
Hahn S, Korner-Nievergelt F, Emmenegger T, Amrhein V, Csörgő T, Gursoy A, Ilieva M,
Kverek P, Pérez-Tris J, Pirrello S, Zehtindjiev P & Salewski V. **2016**. Longer wings for faster
springs – wing length relates to spring phenology in a long-distance migrant across its range.
Ecology and Evolution **6**: 68–77.
Harvey M, Bravo G, Claramunt S, Cuervo A, Derryberry G, Battilana J, Seeholzer G,
McKay JS, O'Meara BC, Faircloth BC, Edwards S, Pérez-Emán J, Moyle RG, Sheldon FH,
Aleixo A, Smith BT, Chesser RT, Silveira LF, Cracraft J, Brumfield RT & Derryberry EP.
2020. The evolution of a tropical biodiversity hotspot. *Science* **370**: 1343–1348.
Herrera CM. **1978**. Ecological Correlates of Residence and Non-Residence in a Mediterranean
Passerine Bird Community. *Journal of Animal Ecology* **47**: 871–890.
Hespenheide HA. **1975**. Selective Predation by Two Swifts and a Swallow in Central America.

Ibis **117**: 82–99.

Hromada M & Tryjanowski P. **2003**. Animals of different phenotype differentially utilise dietary niche - the case of the Great Grey Shrike Lanius excubitor. **80**: 8.

Jahn AE, Levey DJ, Mamani AM, Saldias M, Alcoba A, Ledezma MJ, Flores B, Vidoz JQ & Hilarion F. **2010a**. Seasonal differences in rainfall, food availability, and the foraging behavior of Tropical Kingbirds in the southern Amazon Basin. *Journal of Field Ornithology* **81**: 340–348.

Jahn AE, Levey DJ, Hostetler JA & Mamani AM. **2010b**. Determinants of partial bird migration in the Amazon Basin. *Journal of Animal Ecology* **79**: 983–992.

Jahn AE & Tuero DT. **2020**. Fork-tailed Flycatcher (Tyrannus savana). *Birds of the World*.

Johansson F, Söderquist M & Bokma F. **2009**. Insect wing shape evolution: independent effects of migratory and mate guarding flight on dragonfly wings. *Biological Journal of the Linnean Society* **97**: 362–372.

Kipp F. **1942**. Über Flügelbau und Wanderzug der Vögel. *Biologisches Zentralblatt* **62**: 289–299.

Kipp FA. **1958**. Zur geschichte des Vogelzuges auf der grundlage der Flügelanpassungen. *Vogelwarte* **19**: 233–242.

Knudsen R, Siwertsson A, Adams CE, Garduño-Paz M, Newton J & Amundsen PA. **2011**. Temporal stability of niche use exposes sympatric Arctic charr to alternative selection pressures. *Evolutionary Ecology* **25**: 589–604.

Lack D. **1971**. *Ecological isolation in birds*.

Leisler B. **1990**. Selection and Use of Habitat of Wintering Migrants. In: Gwinner E, ed. *Bird Migration*. Berlin, Heidelberg: Springer Berlin Heidelberg, 156–174.

Levey DJ & Stiles FG. **1992**. Evolutionary Precursors of Long-Distance Migration: Resource Availability and Movement Patterns in Neotropical Landbirds. *The American Naturalist* **140**: 447–476.

Lindström Å & Piersma T. **1993**. Mass changes in migrating birds: the evidence for fat and protein storage re-examined. *Ibis* **135**: 70–78.

Lockwood R, Swaddle JP & Rayner JMV. **1998**. Avian Wingtip Shape Reconsidered: Wingtip Shape Indices and Morphological Adaptations to Migration. *Journal of Avian Biology* **29**: 273–292.

Mallarino R, Grant PR, Grant BR, Herrel A, Kuo WP & Abzhanov A. **2011**. Two developmental modules establish 3D beak-shape variation in Darwin's finches. *Proceedings of the National Academy of Sciences* **108**: 4057–4062.

Marchetti K, Price T & Richman A. **1995**. Correlates of Wing Morphology with Foraging Behaviour and Migration Distance in the Correlates of wing morphology with foraging behaviour and migration distance in the genus Phylloscopus. *Journal of Avian Biology* **26**: 177–181.

McCoy MW, Bolker BM, Osenberg CW, Miner BG & Vonesh JR. **2006**. Size correction: comparing morphological traits among populations and environments. *Oecologia* **148**: 547–554.

Medina JJ, Maley JM, Sannapareddy S, Medina NN, Gilman CM & McCormack JE. **2020**. A rapid and cost-effective pipeline for digitization of museum specimens with 3D photogrammetry. *PLOS ONE* **15**: e0236417.

Milá B, Wayne RK & Smith TB. **2008**. Ecomorphology of Migratory and Sedentary Populations of the Yellow-Rumped Warbler (Dendroica Coronata). *The Condor* **110**: 335–344.

Minias P, Meissner W, Włodarczyk R, Ożarowska A, Piasecka A, Kaczmarek K & Janiszewski T. **2015**. Wing shape and migration in shorebirds: a comparative study. *Ibis* **157**: 528–535.

Mobley JA. **2002**. *Molecular phylogenetics and the evolution of nest building in kingbirds and their allies (Aves: Tyrannidae)*. University of California, Berkeley.

Mobley JA & de Juana E. **2020**. Loggerhead Kingbird (Tyrannus caudifasciatus). *Birds of the World*.

Mönkkönen M. **1995**. Do migrant birds have more pointed wings?: A comparative study.

Evolutionary Ecology **9**: 520–528.

Moreau RE. **1952**. The Place of Africa in the Palaearctic Migration System. *Journal of Animal Ecology* **21**: 250–271.

Morse DH. **1971**. The Insectivorous Bird as an Adaptive Strategy. *Annual Review of Ecology and Systematics* **2**: 177–200.

Mulvihill RS & Chandler CR. **1990**. The Relationship between Wing Shape and Differential Migration in the Dark-Eyed Junco. *The Auk* **107**: 490–499.

Murphy MT. **2007**. A Cautionary Tale: Cryptic Sexual Size Dimorphism in a Socially Monogamous Passerine. *The Auk* **124**: 515–525.

Nations JA, Heaney LR, Demos TC, Achmadi AS, Rowe KC & Esselstyn JA. **2019**. A simple skeletal measurement effectively predicts climbing behaviour in a diverse clade of small mammals. *Biological Journal of the Linnean Society* **128**: 323–336.

Navalón G, Bright JA, Marugán-Lobón J & Rayfield EJ. **2019**. The evolutionary relationship among beak shape, mechanical advantage, and feeding ecology in modern birds*. *Evolution* **73**: 422–435.

Neto JM, Gordinho L, Belda EJ, Marín M, Monrós JS, Fearon P & Crates R. **2013**. Phenotypic Divergence among West European Populations of Reed Bunting Emberiza schoeniclus: The Effects of Migratory and Foraging Behaviours. *PLOS ONE* **8**: e63248.

Norberg UM. **1990**. *Vertebrate flight: mechanics, physiology, morphology, ecology and evolution*. Berlin: Springer.

Norberg UM. **1995**. WING DESIGN AND MIGRATORY FLIGHT. *Israel Journal of Ecology and Evolution* **41**: 297–305.

Orejuela JE, Raitt RJ & Álvarez H. **1980**. *Differential use by North American migrants of three types of Colombian forests: 253-264 (en) KEAST, A. & MORTON, ES (eds.). Migrant Birds in the Neotropics: Ecology, Behavior, Distribution and Conservation*. Smithsonian Institution Press. Washington DC, USA.

Pennycuick C. **2008**. The membrane wings of bats and pterosaurs Modelling the Flying Bird ed CJ Pennycuick.

Pérez-Tris J & Tellería JL. **2001**. Age-related variation in wing shape of migratory and sedentary Blackcaps Sylvia atricapilla. *Journal of Avian Biology* **32**: 207–213.

Pol MVD, Ens BJ, Oosterbeek K, Brouwer L, Verhulst S, Tinbergen JM, Rutten AL & Jong MD. **2009**. Oystercatchers' Bill Shapes as a Proxy for Diet Specialization: More Differentiation than Meets the Eye. *Ardea* **97**: 335–347.

Pyle P, Howell SNG, Institute for Bird Populations, & Point Reyes Bird Observatory. **1997**. *Identification guide to North American birds: a compendium of information on identifying, ageing, and sexing 'near-passerines' and passerines in the hand. Part I, Part I,*. Bolinas, Calif.: Slate Creek Press.

R Core Team. **2020**. *R: A Language and Environment for Statistical Computing (4.0. 3)[Computer software]*. R Foundation for Statistical Computing. Retrieved from http://www. R-proje ….

Rappole JH. **1995**. *The ecology of migrant birds: a neotropical perspective*. Washington; London: Smithsonian institution Press.

Rappole JH, Ramos MA & Winker K. **1989**. Wintering Wood Thrush Movements and Mortality in Southern Veracruz. *The Auk* **106**: 402–410.

Rayner JMV. **1988**. The evolution of vertebrate flight. *Biological Journal of the Linnean Society* **34**: 269–287.

Reaney AM, Bouchenak-Khelladi Y, Tobias JA & Abzhanov A. **2020**. Ecological and morphological determinants of evolutionary diversification in Darwin's finches and their relatives. *Ecology and Evolution* **10**: 14020–14032.

Regosin JV & Pruett-Jones S. **2001**. Sexual Selection and Tail-Length Dimorphism in Scissor-Tailed Flycatchers. *The Auk* **118**: 167–175.

Revell LJ. **2009**. Size-Correction and Principal Components for Interspecific Comparative
Studies. *Evolution* **63**: 3258–3268.
Revell LJ. **2012**. phytools: an R package for phylogenetic comparative biology (and other
things). *Methods in Ecology and Evolution* **3**: 217–223.
Rising JD & Somers KM. **1989**. The Measurement of Overall Body Size in Birds. *The Auk* **106**:
666–674.
Rönn JAC von, Shafer ABA & Wolf JBW. **2016**. Disruptive selection without genome-wide
evolution across a migratory divide. *Molecular Ecology* **25**: 2529–2541.
Senar JC & Pascual J. **1997**. Keel and tarsus length may provide a good predictor of avian
body size. *Ardea* **85**.
Shao S, Quan Q, Cai T, Song G, Qu Y & Lei F. **2016**. Evolution of body morphology and beak
shape revealed by a morphometric analysis of 14 Paridae species. *Frontiers in Zoology* **13**: 30.
Sheard C, Neate-Clegg MHC, Alioravainen N, Jones SEI, Vincent C, MacGregor HEA,
Bregman TP, Claramunt S & Tobias JA. **2020**. Ecological drivers of global gradients in avian
dispersal inferred from wing morphology. *Nature Communications* **11**: 2463.
Sherry TW. **1984**. Comparative Dietary Ecology of Sympatric, Insectivorous Neotropical
Flycatchers (Tyrannidae). *Ecological Monographs* **54**: 313–338.
SHERRY TW. **2005**. Does winter food limit populations of migratory birds? *Birds of two worlds:*
the ecology and evolution of migration: 414–425.
Snow DW. **1953**. Systematics and comparative ecology of the genus Parus in the Palaearctic
region.
Snow DW. **1954**. The habitats of Eurasian tits (Parus spp.). *Ibis* **96**: 565–585.
Stiles FG. **1980**. Evolutionary implications of habitat relations between permanent and winter
resident landbirds in Costa Rica. *Migrant Birds in the Neotropics (A. Keast and ES Morton,*
Eds.) Smithsonian Institution Press, Washington DC: 421–435.
Svanbäck R & Schluter D. **2012**. Niche Specialization Influences Adaptive Phenotypic
Plasticity in the Threespine Stickleback. *The American Naturalist* **180**: 50–59.
Swanson MT, Oliveros CH & Esselstyn JA. **2019**. A phylogenomic rodent tree reveals the
repeated evolution of masseter architectures. *Proceedings of the Royal Society B: Biological*
Sciences **286**: 20190672.
Tarka M, Åkesson M, Beraldi D, Hernández-Sánchez J, Hasselquist D, Bensch S &
Hansson B. **2010**. A strong quantitative trait locus for wing length on chromosome 2 in a wild
population of great reed warblers. *Proceedings of the Royal Society B: Biological Sciences* **277**:
2361–2369.
Tellería LJ, Blázquez M, De La Hera I & Pérez-Tris J. **2013**. Migratory and resident Blackcaps
Sylvia atricapilla wintering in southern Spain show no resource partitioning. *Ibis* **155**: 750–761.
Tellería JL & Carbonell R. **1999**. Morphometric Variation of Five Iberian Blackcap Sylvia
atricapilla Populations. *Journal of Avian Biology* **30**: 63–71.
Thiollay JM. **1988**. Comparative Foraging Success of Insectivorous Birds in Tropical and
Temperate Forests: Ecological Implications. *Oikos* **53**: 17–30.
Vágási CI, Pap PL, Vincze O, Osváth G, Erritzøe J & Møller AP. **2016**. Morphological
Adaptations to Migration in Birds. *Evolutionary Biology* **43**: 48–59.
Van Dijk A, Nakamura G, Rodrigues AV, Maestri R & Duarte L. **2021**. Imprints of tropical
niche conservatism and historical dispersal in the radiation of Tyrannidae (Aves:
Passeriformes). *Biological Journal of the Linnean Society*.
Van Valen L. **1965**. Morphological Variation and Width of Ecological Niche. *The American*
Naturalist **99**: 377–390.
Voelker G. **2001**. Morphological correlates of migratory distance and flight display in the avian
genus Anthus. *Biological Journal of the Linnean Society* **73**: 425–435.
Waide RB. **1980**. Resource partitioning between migrant and resident birds: the use of irregular
resources. *Migrant birds in the Neotropics: ecology, behavior, distribution, and conservation*:

337–352.
Wang X & Clarke JA. **2015**. The evolution of avian wing shape and previously unrecognized
trends in covert feathering. *Proceedings of the Royal Society B: Biological Sciences* **282**:
20151935.
Wiedenfeld DA. **1991**. Geographical Morphology of Male Yellow Warblers. *The Condor* **93**:
712–723.
Willis EO. **1974**. Populations and Local Extinctions of Birds on Barro Colorado Island, Panama.
Ecological Monographs **44**: 153–169.
Wilson EO. **1961**. The Nature of the Taxon Cycle in the Melanesian Ant Fauna. *The American*
Naturalist **95**: 169–193.
Winkler H & Leisler B. **1992**. On the ecomorphology of migrants. *Ibis* **134**: 21–28.
Winquist T & Lemon RE. **1994**. Sexual Selection and Exaggerated Male Tail Length in Birds.
The American Naturalist **143**: 95–116.
Závorka L, Larranaga N, Lovén Wallerius M, Näslund J, Koeck B, Wengström N,
Cucherousset J & Johnsson JI. **2020**. Within-stream phenotypic divergence in head shape of
brown trout associated with invasive brook trout. *Biological Journal of the Linnean Society* **129**:
347–355.

620 Supporting Information

621 Supporting Information 1 – Supplementary Results

622 Supporting Information 2 – ImageJ protocol for measuring long *Tyrannus* tail lengths

623 Supporting Information 3 – Sex-specific results: means for each morphometric, phylogenetic

624 principal components analysis (PPCA), and phylogenetic ANOVA