Chained structure of dimeric F$_1$-like ATPase in *Mycoplasma mobile* gliding machinery

Takuma Toyonaga[a], Takayuki Kato[b], Akihiro Kawamoto[b], Noriyuki Kodera[c], Tasuku Hamaguchi[a,d], Yuhei O Tahara[a,d], Toshio Ando[c], Keiichi Namba[e,f,g], Makoto Miyata[a,d,#]

[a]Graduate School of Science, Osaka City University, 3-3-138 Sugimoto, Sumiyoshi-ku, Osaka 558-8585, Japan.

[b]Institute for Protein Research, Osaka University, 3-2 Yamadaoka, Suita, Osaka, 565-0871, Japan.

[c]Nano Life Science Institute (WPI-NanoLSI), Kanazawa University, Kakuma-chou, Kanazawa, Ishikawa 920-1192, Japan.

[d]The OCU Advanced Research Institute for Natural Science and Technology (OCARINA), Osaka City University, 3-3-138 Sugimoto, Sumiyoshi-ku, Osaka 558-8585, Japan.

[e]Graduate School of Frontier Biosciences, Osaka University, 1-3 Yamadaoka, Suita, Osaka, 565-0871, Japan.

[f]RIKEN Center for Biosystems Dynamics Research and SPring-8 Center, 1-3 Yamadaoka, Suita, Osaka, 565-0871, Japan.

[g]JEOL YOKOGUSHI Research Alliance Laboratories, Osaka University, 1-3

Yamadaoka, Suita, Osaka, 565-0871, Japan.

[#]Address correspondence to Makoto Miyata, miyata@osaka-cu.ac.jp

Present address: Tasuku Hamagichi: Biostructural Mechanism Laboratory, RIKEN

SPring-8 Center, 1-1-1 Kouto, Sayo, Hyogo, 679-5148, Japan.

Running title: Gliding machinery of *Mycoplasma mobile*

Keywords: F_1-ATPase, rotary motor, parasitic bacteria, electron microscopy, atomic

force microscopy, bacterial motility

ABSTRACT

Mycoplasma mobile, a fish pathogen, exhibits gliding motility using ATP hydrolysis on solid surfaces, including animal cells. The gliding machinery can be divided into surface and internal structures. The internal structure of the motor is composed of 28 so-called 'chains' that are each comprised of 17 repeating protein units called 'particles'. These proteins include homologs of the catalytic α- and β-subunits of F_1-ATPase. In this study, we isolated the particles and determined their structures using negative-staining electron microscopy and high-speed atomic force microscopy. The isolated particles were composed of five proteins, MMOBs 1660 (α-subunit homolog), 1670 (β-subunit homolog), 1630, 1620, and 4530, and showed ATP hydrolyzing activity. The 2D structure, with dimensions of 35 and 26 nm, showed a hexameric ring dimer about 12 nm in diameter, resembling F_1-ATPase catalytic $(αβ)_3$. We isolated the F_1-like ATPase unit, which is composed of MMOBs 1660, 1670, and 1630. Furthermore, we isolated the chain and analyzed the 3D structure, showing that dimers of mushroom-like structures resembling F_1-ATPase were connected and aligned along the dimer axis at 31 nm intervals. An atomic model of F_1-ATPase catalytic $(αβ)_3$ from *Bacillus* PS3 was successfully fitted to each hexameric ring of the mushroom-like structure. These results suggest that the motor for *M. mobile* gliding shares an evolutionary origin with F_1-ATPase. Based on the obtained structure, we propose possible force transmission processes in the gliding mechanism.

IMPORTANCE

F_1F_o- ATPase, a rotary ATPase, is widespread in the membranes of mitochondria, chloroplasts, and bacteria, and converts ATP energy with a proton motive force across the membrane by its physical rotation. Homologous protein complexes play roles in ion and protein transport. *Mycoplasma mobile*, a pathogenic bacterium, was recently suggested to have a special motility system evolutionarily derived from F_1-ATPase. The present study isolated the protein complex from *Mycoplasma* cells and supported this conclusion by clarifying the detailed structures containing common and novel features as F_1-ATPase relatives.

INTRODUCTION

Mycoplasmas are parasitic bacteria characterized by small cell size, a short genome and lack of a peptidoglycan layer (1-3). Many *Mycoplasma* species exhibit a unique gliding motility, which is necessary for their infection (4-6). *Mycoplasma mobile* glides on solid surfaces at 2.0 to 4.5 µm/s in the direction of a protrusion on one side of the cell (Fig. S1) (5). The gliding machinery is localized to the cell protrusion and can be divided into surface and internal structures (Fig. 1A, upper). The surface structure has approximately 450 repeats of a complex of three large proteins, Gli123, Gli521, and Gli349, inserted into the cell membrane (Fig. 1A, lower) (7-11). Fifty-nm-long leg structures corresponding to Gli349 molecules can be seen jutting out from cell protrusion by electron microscopy (EM) (12). The tip of Gli349 is characterized by a "foot" with an oval structure that can bind to sialylated oligosaccharides (SOs) (13-20). Gli521 and Gli123 serve as the "crank" that transfers force to Gli349 and the "mount" that localizes the other two surface proteins to the gliding machinery, respectively. A working model of the gliding mechanism has been proposed in which the cells are propelled by Gli349 molecules that repeatedly catch, pull, and release SOs on solid surfaces (5, 21-23).

The internal structure consists of a lumpy structure, called "bell" at the tip of the cell protrusion and 28 "chains" lining the inner membrane surface (Fig. 1A) (23-26). Each chain is characterized by 17 repeating particle structures, resulting in a total of 476 particles in one cell. The chains tend to form sheets when they are isolated from cells, suggesting lateral interaction with the adjacent chains (23, 24). The

internal structure consists of least ten proteins (Fig. 1B). Six of these proteins,

MMOBs 1620, 1630, 1640, 1650, 1660 and 1670 were coded tandemly in a locus.

Interestingly, MMOBs 1660 and 1670, which have Walker A and B motifs that are

involved in ATP binding and hydrolysis (27), show high amino acid sequence

identity with the catalytic subunits α and β of F_1-ATPase, respectively. MMOB1670

has an extra N-terminal region (amino acids 1–299), which is not present in the β

subunit. The chains most likely contain the motor for gliding, because the gliding

motility is coupled to ATP hydrolysis (23, 28, 29).

F_1F_o-ATPases, found in most organisms, are rotary motors that perform

biological energy conversion (30, 31). Their role is to both synthesize ATP using a

proton motive force and, conversely, to hydrolyze ATP to drive protons to maintain

the membrane potential. Their structure is composed of a soluble catalytic

F_1-domain for ATP catalysis and a membrane-embedded F_o-domain for the proton

pathway. In the F_1-domain, the catalytic subunits α and β alternate to form a

hexameric ring $(αβ)_3$ that rotates the central stalk penetrating the ring using ATP

hydrolysis. Phylogenetic studies have shown that *Mycoplasma* have three F_1-like

ATPase clusters, which are referred to as Type 1–3 ATPases (26, 32). Type 1, found

in all mycoplasmas, is a typical operon encoding F_1F_o-ATPase and is likely to

function as a proton pump to maintain membrane potential. Type 3 is found in

mycoplasmas that have an MIB-MIP system to cleave host immunoglobulins (33).

Type 2 is only found in four *Mycoplasma* species, including *M. mobile*. Interestingly,

the Type 2 ATPase of *M. mobile*, which encodes MMOB1620–70, is involved in the

internal structure of the gliding machinery.

Recently, the chains of the internal structure were shown to have structural

changes linked to ATP hydrolysis. However, it is still unclear how the chain

generates and transmits the force to the outside, because its detailed structure has

not been clarified. In this study, we isolated the chains and elucidated their

structures. The structure had a common architecture with F_1-ATPase, suggesting

that the chain shares a common evolutionary origin with F_1-ATPase. Based on our

findings, we suggest two possible force transmission models for the gliding

machinery.

Results

Isolation and biochemical analyses of stable unit complex. In this study, we

isolated and analyzed three fractions including "Dimer", "Monomer", and "Chain"

structures which are schematically shown in Fig. 1A (Fig. S2, S3).

First, to isolate a stable unit complex from the chain, we lysed cultured *M.*

mobile cells with 1% Triton X-100 and recovered the insoluble fraction by

centrifugation (24). This fraction named as "Pellet-1" was used for further

preparations. We suspended Pellet-1 in suspension buffer, which contains 137 mM

NaCl, and solubilized the putative unit complex for 8 h. Here, we used 137 mM NaCl

rather than 400 mM to reduce the contamination of other proteins. The soluble

fraction was then subjected to Superdex 200 gel filtration chromatography. The

obtained peak fraction in the void contained MMOBs 1620, 1630, 1660, and 1670,

which are coded on the mycoplasma Type 2 ATPase operon, and MMOB4530

annotated as phosphoglycerate kinase (PGK) (Fig. 2A). These proteins are known

to be components of the internal structure (Fig. 1B) (23-26).

To examine the assembly of these proteins, we applied the isolated fraction to

gel filtration chromatography using a Sephacryl S-400 HR column, which can

fractionate up to 8000 kDa globular proteins (Fig. 2B). The proteins eluted as a

single peak at a non-void position and were larger than 669 kDa, suggesting that

they form a large complex. The molar ratios of the components were estimated to

be 3.2:2.9:3.0:1.0:2.3 for MMOBs 1670, 4530, 1660, 1630, and 1620, respectively,

from the relative intensity of the SDS-PAGE bands. We then analyzed the isolated

fraction by blue-native (BN) PAGE (Fig. 2C, left). A single band was detected, which

is consistent with the result of gel filtration chromatography showing a single peak.

Next, we applied the band to an In-gel ATPase activity assay, which detects the

activity as a white precipitation of lead caused by released of inorganic phosphate

(Fig. 2C, right). The band with the complex showed precipitation, whereas the band

with thyroglobulin, the negative control, did not. This result indicates that the

isolated complex has ATPase activity.

In addition, we assayed the isolated fraction for phosphate release from solution.

The complex hydrolyzed ATP at a maximum turnover rate of 0.18 molecules/s per

MMOB1670 subunit, β-subunit paralog with a K_m of 74 µM at 25°C (Fig. 2D). The

ATPase activity was inhibited by addition of 15.4 mM sodium azide, an inhibitor to

ATPases with Walker A motifs (35), with a K_m of 108 µM and a maximum turnover

rate of 0.055 molecules/s. In a previous study, the Triton-insoluble fraction, which

included the internal structure, showed ATPase activity with a K_m of 66 µM and a

maximum turnover rate of 0.09 molecules/s and was suppressed by 15.4 mM

sodium azide, showing a K_m of 84 µM and a maximum rate of 0.063 molecules/s

(23). The values obtained here are comparable to these previous data. The above

results suggest that the isolated complex is the motor in the internal structure of the

gliding machinery. We used this unit complex for further analyses.

Hexamers resembling F_1-ATPase catalytic $(\alpha\beta)_3$ form a dimer. We observed

the unit complex by EM using the negative-staining method. A field image showed

uniform particles with dimensions of 40 and 20 nm (Fig. 3A and B). As the particle

frequency depended on the protein concentration, we concluded that the observed

particles were a part of the protein complex. We picked 2148 particle images

automatically using RELION software (36) for 2D-classification. From the

2D-classification, we obtained four clear particle images (Fig. S4). We adopted

mirror images according to the structure observed in high-speed atomic force

microscopy (HS-AFM) (see below). Structural handedness cannot be judged from

EM images because they are projections of electrons transmitted through the

sample. We focused on an image showing a complex structure with dimensions of

35 and 26 nm featuring nearly two-fold symmetry (Fig. 3C and D). Interestingly, the

characteristic hexamer of about 12 nm in diameter formed a dimer and was

reminiscent of F_1-ATPase catalytic $(\alpha\beta)_3$. Considering that the amino acid

sequences of MMOBs 1660 and 1670 have high identity to the α- and β-subunits of

F_1-ATPase, respectively, the dimeric complex is likely an evolutionary related

F_1-ATPase. The distance between the centers of the two hexamers was 11.0 nm.

The complex had ten filamentous structures around the two hexamers, four of which

appeared to form bridges across the two hexamers. The filamentous structures are

unlikely artifact, because they showed common features in four independent

averaged images (Fig. S4). Hereafter, we refer to this structural unit as "Dimer."

The hexamer featured a ring and a peak in HS-AFM of Dimer. Next, we

visualized Dimer using HS-AFM to clarify the structure under liquid conditions,

because the molecules are in dry condition under negative-staining EM. HS-AFM is

a powerful method that can visualize the structure and dynamics of single

molecules in liquid conditions at a video rate (37, 38). In this method, a specimen is

placed on the stage surface and with a probe is scanned in buffer at high speed. In

this study, we placed Dimer on a mica surface and scanned it in an area of 70 × 70

nm^2 at 56 × 56 pixels with a scanning rate of 100 ms per frame. HS-AFM images

showed a complex with dimensions of approximately 30 and 20 nm composed of

two globules and attached by 2–4 lateral protrusions shorter than 15 nm (Fig. 3E;

Movie S1 and S2). The Dimer images were categorized into two patterns as either a

ring (pattern I) or a peak (pattern II), based on the central part. Then, we observed

them at a higher resolution (area, 40 × 40 nm^2 with 50 × 50 pixels; scanning rate,

100 ms per frame) (Fig. 3F). In pattern I, the slice image near the top end of the

particle between 9.8 and 11.3 nm above the substrate surface showed two

hexameric rings (Fig. 3G). The position and direction of the two rings are consistent

with those of the hexamers in the negative-staining EM image. In addition, the

distance between the centers of the two hexameric rings was 10.4 nm (Fig. 3H),

similar with the distance between the centers of the hexamers in the

negative-staining EM image (Fig. 3C). These observations suggest that the shape

of the Dimer structure in liquid is preserved in negative-staining EM conditions and

that the hexamers form rings like F_1-ATPase catalytic $(\alpha\beta)_3$. In pattern II, the two

central peaks were positioned 11.2 nm apart (Fig. 3H, lower), similar to the distance

between the centers of the hexameric rings in pattern I (Fig. 3H, upper), suggesting

that patterns I and II are two sides of the same coin (Fig. 3E, left). The distances

between the hexameric rings are slightly different for patterns I and II, as 10.4 and

11.2 nm, respectively. This difference suggests that the central axes of two

hexamers are not parallel. Considering that HS-AFM detects surface structures

while EM shows projection images, the corresponding distance, 11.0 mm in EM is

consistent with the numbers obtained from HS-AFM. Interestingly, in most of the

particles, the two peaks at 6 and 5 nm became invisible in 20 s, between frames 1

and 3 (Fig. 3I and J; Movie S3). We concluded that these subunits dropped out from

the complex, because they did not reappear until the complex was disrupted. Next,

we focused on the lateral protrusions of these particles, which may be related to the

sheet formation of chains (23). To visualize them more clearly, we scanned Dimer

by HS-AFM with an area of 120 × 120 nm^2, 120 × 120 pixels, and scanning rate 500

ms per frame. Dimer showed seven lateral protrusions around the two globules (Fig.

3K). These protrusions swayed without being fixed (Fig. 3L; Movie S4).

Isolation and structure of Monomer. To clarify the components and structure

of the hexamer, we focused on the monomeric hexamer, called Monomer (Fig. S2).

We treated Dimer fraction with 1.5% sodium cholate, an anionic detergent.

BN-PAGE and In-gel ATPase activity assays showed a single band with ATPase

activity at a position lower than the original one, corresponding to 720–1048 kDa,

indicating that Dimer dissociated into smaller units with ATPase activity (Fig. 2C).

Therefore, to isolate Monomer, we applied Pellet-1 sequentially to the treatments

with 250 mM NaCl for 8 h, 1.5% sodium cholate for 7 h, and to Sephacryl S-400HR

gel filtration chromatography (Fig. S2). The elution pattern showed a rather isolated

small peak and following overlapping peaks (Fig. S3C). An SDS-PAGE gel showed

that MMOBs 1670, 1660, and 1630 eluted in the same fractions, while MMOBs

4530 and 1620 eluted at later fractions (Fig. S3CD), indicating that MMOBs 4530

and 1620 were dissociated from Dimer by sodium cholate treatment. Then, we

focused on the small peak fraction, which mainly contained MMOBs 1670, 1660,

and 1630 (Fig. 2A, middle, S3D). This complex presumably corresponds to the

BN-PAGE band showing ATPase activity (Fig. 2C), because only MMOBs 1660 and

1670 have the Walker A and Walker B motifs in the Dimer components. EM

observation using the negative-staining method showed uniform globular particles

10–15 nm in diameter (Fig. 4A and B). As the particle frequency depended on the

protein concentration, we concluded that the observed particles were a part of

Monomer. We picked 11687 particle images automatically using RELION software

for 2D-classification. By 2D-classifing the images in 50 classes, we obtained 15

clear particle images, which were averaged (Fig. S5). Fig. 4C shows a 12 nm

diameter globule characterized by a single hexameric ring, corresponding to a part

of the Dimer image in Fig. 3C. Three of the subunits were larger than the others with

hook structures on either side of the edge. Three averaged images (II–IV) showed a

mushroom-like structure resembling F_1-ATPase, which is characterized by a 12

nm-diameter umbrella and a 3 nm-long stalk (Fig. 4C). Now, we can suggest that

MMOBs 1670, 1660, and 1630 form Monomer, i.e., F_1-like ATPase unit. MMOB4530

was probably not included in this unit because it probably binds to the complex and

could not be distinguished in the image due to the low proportion of bound entities

(Fig. 2A, middle). Thus, the hexameric ring is likely formed by the α-subunit

homolog MMOB1660 and the β-subunit homolog MMOB1670, and the stalk is

formed by MMOB1630.

Isolation and structure of Chain. In gliding machinery, Dimer link to form

chains. To characterize these chains, we isolated "Chain fraction" with milder

mechanical treatment than those for other fractions. Pellet-1 was treated by 387 mM

NaCl and soluble fraction was isolated. Chain fraction contained more than 30

proteins, including the Dimer component proteins MMOBs 1670, 4530, 1660, 1630,

and 1620 as major components (Fig. 2A, right). EM observation using the

negative-staining method showed chain structures with lengths longer than 70 nm

and particles of various sizes (Fig. 5A and B). This time we manually picked 2127

particles from the chain images, overlapping approximately 50% of the 71 × 71 nm^2

box area. From 2D-classifcation, we obtained seven clear particle images (Fig. 5C).

The particle images show the various orientations required for 3D reconstruction.

We then created a 3D map by combining a total of 1709 particle images of good

quality (Fig. 5D and S6). The 3D map with dimensions of 70, 20, and 15 nm at a

density threshold (contour level = 0.026) was composed of two dimers of

mushroom-like structures resembling F_1-ATPase, aligned along the dimer axis (Fig.

5D). Dimers were connected by a bulge structure with a diameter of 5 nm. The

chain interval was 31 nm, consistent with the corresponding dimension in a 2D

image from electron cryotomography (ECT) (23), suggesting that the 3D model

obtained reflects the original structure from a cell. The mushroom-like structure with

a diameter of 15 nm, consisting of a hexameric ring and a central stalk, was

connected to the dimer by two bridge structures with a diameter of 3–6 nm. An

atomic model of F_1-ATPase catalytic $(\alpha\beta)_3$ from *Bacillus* PS3 (PDB ID 6N2Y) (39)

was fitted into each hexameric ring of the mushroom-like structure (Fig. 5E). The

distance between the centers of the fitted $(\alpha\beta)_3$ in Dimer was 12.5 nm, which is

consistent with that of Dimer observed by negative-staining EM and HS-AFM (Fig.

3C and H). The fitted model showed that each hexameric ring had two protrusions

of 3–6 nm pointing laterally (Fig. 5E). The cross-sections of each mushroom-like

structure showed the central stalk length of 5 nm (Fig. 5F). A cavity was observed at

the center of the hexameric ring. However, it may be an artifact of the low-resolution

map of negative-staining EM, because metal coating tends to emphasize the

peripheral part of large particles (40). Next, we compared a reprojection image of

the 3D chain map with the 2D averaged image of Dimer from negative-staining EM

(Fig. 5G). Two short filaments marked by asterisks in Dimer (Fig. 5G, left) are

positioned facing each other in the connecting bulge (Fig. 5G). Previously, ECT of a

permeabilized *M. mobile* cell showed a chain structure characterized by repeats of

two globules and two types of projections to the cell membrane (Fig. 5H, left) (23).

The hexameric ring and the central stalk in the 3D map here correspond to the

globule and one type of projection to the cell membrane in the ECT image,

respectively, suggesting that the chain is oriented with the central stalk facing the

membrane, which is common in F_1-ATPases. At the interface between Dimers, the

3D map here did not include a structure composed of another type of projection and

a globule as observed in the ECT image. The subunits corresponding to these

structures probably had structural variations or dissociation during the isolation

process.

Discussion

Outline of internal structure of gliding machinery. Previously, sequence

analysis suggested that the chain of *M. mobile* gliding machinery evolved from

F_1-ATPase (5, 23-26). The present study supports this conclusion by structural data

showing that the chain has hexameric rings similar to the F_1-ATPase catalytic $(\alpha\beta)_3$.

Integrating available information, we can now describe the outline of the internal

structure of *the M. mobile* gliding machinery (Fig. 1A). *M. mobile* cells have 28

individual 530 nm long chains, each of which contains 17 Dimer units composed of

two F_1-like ATPases and filamentous structures (23). The central stalk of the F_1-like

ATPase and another protrusion from the connecting bulge project to the cell

membrane.

Unique role of F_1-ATPase related complex. To date, several complexes are

known to be evolutionarily related to F_1-ATPase, all of which are responsible for

transporting substances across the membrane (41). However, the motor we

identified here most likely plays a role in motility. This case may be reminiscent of

dynein, a motor in eukaryotes, which evolved from a widely conserved AAA

(ATPases associated with diverse cellular activities)+ protein, in which multiple

subunits of ATPases perform functional rotation (42, 43). Sequence analyses have

shown that mycoplasma Type 3 ATPase is also related to F_1-ATPase, and its role

has been suggested to promote substrate turnover in the MIB-MIP system (33). If

Type 3 ATPase provides the force to change the conformation of a hydrolytic

enzyme, its role in force generation is common with Type 2, the gliding motor.

F_1F_o-ATPases are known to be dimerized through interactions between

F_o-domains and are usually arranged in rows along the short axis in the tightly

curved cristae ridges of mitochondria (31, 44, 45). The dimer structure found in the

present study is not related to this, because the F_1-like domain is dimerized through

the filament structure and is linked in the long axis direction. However, some roles

may be common in part if the dimerization and chain formation observed in the

gliding motor identified in this study stabilizes the membrane structure, as seen in

the F_1F_o-ATPase dimer (46). Moreover, dimerization may result in cooperativity in

motor functions.

Protein assignment. The α-subunit homolog MMOB1660 (58.7 kDa) and the

β-subunit homolog MMOB1670 (88.4 kDa) likely correspond to the smaller and

larger subunits, respectively, of the hexameric ring of an F_1-like ATPase unit, as

suggested by the estimated 1:1 molar ratio in Dimer (Fig. 2A, 4C). This means that

the hook structure of the larger subunit may be formed by the extra N-terminal

region (34.8 kDa) of MMOB1670. Previously, 3D structure modeling based on

secondary structure suggested that MMOB1630 is structurally similar to the γ

subunit, the principal component of the central stalk of F_1-ATPase (5). In general,

the γ subunit of F_1-ATPase is composed of a coiled-coil and a globular domain and

penetrates the hexameric ring (47). In the F_1-like ATPase unit and chain 3D model,

a stalk structure, suggesting the globular domain of the γ subunit, was found in the

center of the hexameric ring (Fig. 4C and 5F), implying that MMOB1630 penetrates

the hexameric ring like the γ subunit.

Using HS-AFM observations, the peak at approximately 5 nm at the center of the

hexameric ring dropped out with time (Fig. 3I). The peak height agrees with the

length of the estimated globular domain of MMOB1630 in Chain 3D model (Fig. 3J

and 5F), suggesting that the peak is composed of MMOB1630 and was pulled out

from the hexameric ring by the scanning cantilever during HS-AFM observation.

The pull-out event is thought to be common to that of the F_1-ATPase, in which the γ

subunit is removed from the hexameric ring by optical tweezers (48).The

filamentous structures around the hexameric ring probably correspond to lateral

protrusions in the HS-AFM images and are formed by the remaining proteins,

MMOB1620 and MMOB4530 (PGK). These proteins probably play roles in ATPase

dimerization, chain formation, and lateral chain interaction (23). MMOB1620 is an

unannotated protein specific to the Type 2 ATPase cluster (26, 32). MMOB4530 is

annotated as an enzyme that transfers phosphate groups from

1,3-bisphosphoglycerate to ADP in glycolysis to yield ATP and 3-phosphoglycerate

(49). In *M. mobile*, ATP is probably provided by glycolysis (50). MMOB4530 may

supply ATP efficiently to the gliding motor by its close proximity. Yeast V-ATPase,

which belongs to the rotary ATPase family like F_1F_o-ATPase, is also attached by two

glycolytic enzymes, 6-phosphofructo-1-kinase and aldolase (51-53). These

glycolytic enzymes are involved in the regulation of V-ATPase assembly and

activity.

Ten proteins have been identified as the components of the internal structure (Fig.

1) (23, 24, 26). In the present study, we identified five proteins as the Dimer

components. Other five proteins are likely involved in other parts, for example "bell"

at the front part of internal structure (Fig. 1A) and parts of the intact chain structure

which were lost in the fractionation (Fig. 5H).

Possible force transmission mechanisms for gliding. The involvement of an

internal ATPase in the gliding mechanism is based on the following five

observations from the analysis of the "gliding head" of *M. mobile* protrusions and of

the isolated gliding machinery: (a) The affinity for ATP estimated by the saturation

extent is comparable in the ATPase activity of the internal structure and the speed

of the gliding head (23). (b) Substrate binding and gliding speed of the gliding head

are inhibited by azide, as well as the ATPase activity of the internal structure (23).

(c) The chain in the internal structure undergoes conformational changes based on

ATP hydrolysis (23, 34). (d) Among the 21 proteins identified from the gliding head,

only MMOBs 1660 and 1670 could be suggested for ATPase from the amino acid

sequences alone (23, 24). (e) Fluorescent protein tagging of components of the

internal structure significantly affects the substrate binding of cell and the gliding

speed (26).

The structure elucidated in the present study allows us to discuss the gliding

mechanism in more detail. In F_1-ATPase, the three catalytic sites in the hexameric

ring cooperatively hydrolyze ATP, and each catalytic β-subunit undergoes a

conformational change that drives the rotation of the central stalk (47). Previously,

structural changes linked to ATP hydrolysis were reported: (I) EM studies showed 2

nm contraction of Dimer intervals in the isolated chains (23), and (II) HS-AFM

studies showed movements of individual Dimers in the cell 9 nm perpendicular to

the chain long axis and 2 nm into the cell (34). Considering these observations, we

propose two different working models for the force transmission mechanism in

gliding (Fig. 6). In the "contraction model" (Fig. 6 i), the force generated by the

hexameric ring shortens the chain. The resulting displacement of the projections

from Dimer to the cell membrane drives the hook structure of Gli521 like a "lever."

Then, the leg moves with the catch, pull, and release of the SOs. In the "rotation

model" (Fig. 6 ii), the force generated by the hexameric ring rotates MMOB1630 in

the same way as F_1-ATPase. This rotation is transmitted across the cell membrane

to the Gli521. The hook structure converts rotational motion into linear motion of the

leg, similar to a crank. Previous studies have reported that *M. mobile* exhibits

unitary steps of approximately 70 nm in size at no load (29, 54). In our models, both

the rotation and contraction displacements are expected to be a few nanometers.

These displacements may be amplified by the large surface structure complex

formed by the 100 nm long Gli349 and 120 nm long Gli521, which show dimensions

comparable to the step size, acting as a large gear (22). This conjecture could

explain how a single leg exerts a force of 1.5 pN, smaller than that of conventional

motor proteins such as myosin, dynein, and kinesin.

Evolution of *M. mobile* gliding. A previous study suggested that Gli349 evolved

from a static binding receptor to parasitize the host (16). Considering this, the

evolutionary origin of *M. mobile* gliding can now be discussed. F_1F_o-ATPase, which

is abundant on the cell membrane, could have been accidentally associated with

the binding receptor and turned into a primitive motility system, which may have

provided random cell spreading. The system was then refined under survival

pressure, because motility might be beneficial to infection and evading the host's

immune system. For dimerization and chain formation, PGK was then incorporated

into the gliding machinery, because PGK was working in close proximity to

F_1F_o-ATPase.

Materials and Methods

Strains and culture conditions. We used P476R *gli521*, a mutant strain of *M. mobile* that can glide normally but binds SOs more tightly than wild-type strains (10, 22, 28). *M. mobile* cells were cultured as described previously (55, 56).

Optical microscopy. The cultured cells were inserted into a tunnel chamber assembled with two coverslips and double-sided tapes and observed by phase-contrast microscopy using an inverted microscope (IX71; Olympus, Tokyo, Japan) (17, 19). Movement was recorded using a complementary metal-oxide semiconductor (CMOS) camera (DMK33UX174; The Imaging Source, Bremen, Germany). Video was analyzed using the ImageJ software, version 1.53a (http://rsb.info.nih.gov/ij/).

Solubility test. All procedures for fractionation were performed at 4°C unless otherwise noted and focused protein bands were identified by peptide mass fingerprinting (PMF), as previously reported (24, 57). To investigate the solubility of the chain components, *M. mobile* cells from 60 mL of culture medium were collected by centrifugation at 14000 × *g* for 20 min and washed twice with PBS consisting of 8.1 mM Na_2HPO_4, 1.5 mM KH_2PO_4, pH 7.3, 2.7 mM KCl and 137 mM NaCl. Cells were resuspended in PBS to a 12-fold higher concentration than the culture and sonicated for 1 min at 24–27°C to be dispersed in microtubes using an ultrasonic generator (2510 J-MT; BRANSON, Kanagawa, Japan). The cells were then treated with Triton solution (1% Triton X-100, 0.1 mg/mL DNase, 5 mM $MgCl_2$, and 1 mM phenylmethylsulfonyl fluoride in PBS) in a total volume of 10 mL. After gentle

shaking for 1 h, the suspensions were centrifuged at 20000 × g for 20 min, and

pellets were collected and washed once with suspension buffer (PBS including 5

mM MgCl$_2$). This fraction is Pellet-1. Pellet-1 was then resuspended separately in

suspension buffer adjusted to contain different concentrations of NaCl (0, 50, 137,

200, and 400 mM) by pipetting several times. After treatment for 8 h, suspensions

were centrifuged at 20000 × g for 20 min, and supernatants and pellets collected for

SDS-PAGE analysis.

Preparation of fractions. For isolation of Dimer, Pellet-1 from 1.2-liter culture

was resuspended in 5 mL suspension buffer by pipetting up and down and allowed

to dissolve for 8 h. The soluble fraction was collected by centrifugation at 20000 × g

for 20 min and loaded onto a HiLoad 16/600 Superdex 200 column (Cytiva, Tokyo,

Japan) equilibrated with 1 mM MgCl$_2$ in PBS at a flow rate of 0.8 mL/min. The

fractions were analyzed by SDS-PAGE and CBB staining.

For isolation of Monomer, Pellet-1 from 1.2-liter culture was suspended by 5 mL

Tris buffer consisting of 20 mM Tris-HCl (pH 7.5), 250 mM NaCl, 1 mM

phenylmethylsulfonyl fluoride, and 1 mM MgCl$_2$ with pipetting and allowed to

dissolve for 8 h. The soluble fraction was collected by centrifugation at 20000 × g for

20 min and mixed with 1.5% (w/v) sodium cholate. After 7 h of incubation, the

complexes were loaded onto a Sephacryl S-400 HR column (Cytiva) equilibrated

with 0.7% sodium cholate, 20 mM Tris-HCl (pH 7.5), 250 mM NaCl, and 1 mM

MgCl$_2$ at a flow rate of 0.5 mL/min. The fractions were analyzed by SDS-PAGE and

CBB- and reverse-staining (58, 59). The fraction of the complex composed of

MMOBs 1670, 1660, and 1630 was collected. Samples were concentrated using an

Amicon Ultra 100 K spin filter (Merck KGaA, Darmstadt, Germany), if necessary.

For isolation of Chain, Pellet-1 from 15 mL culture was resuspended in 60 µL

suspension buffer. The suspension was then gently mixed with an equal volume in a

suspension buffer adjusted to contain 637 mM NaCl. Chain was recovered as the

supernatant by centrifugation at 5000 × g for 5 min.

Analytical gel filtration. Dimer solution was loaded onto a Sephacryl S-400

HR column equilibrated with gel filtration buffer containing 20 mM Tris-HCl (pH 7.5),

200 mM NaCl and 1 mM $MgCl_2$ at a flow rate of 0.5 mL/min at 4°C. Thyroglobulin

(669 kDa; Gel Filtration Calibration Kits; Cytiva) was dissolved in gel filtration buffer

and loaded onto the column as a size standard at a flow rate of 0.5 mL/min. The

stoichiometry of protein complexes was estimated by densitometry of SDS-PAGE

gels stained with CBB, using a scanner (GT-9800F; Epson, Nagano, Japan) and

ImageJ (9).

BN-PAGE and in-gel ATPase activity assays BN-PAGE was performed

according to the user manual of the Native PAGE Novex Bis-Tris Gel System

(Thermo Fisher Scientific, Waltham, MA). For BN-PAGE of sodium cholate treated

Dimer, SC treated, Dimer fraction was mixed with sodium cholate (1.5%) and

incubated at 4°C for 9 h. When this sample was mixed with a sample buffer,

NativePAGE™ 5% (w/v) G-250 sample additive was also added at 0.5% (w/v) to

prevent protein aggregation. Thyroglobulin was dissolved in water and used as a

negative control for the In-gel ATPase activity assay. For the In-gel ATPase activity

assay (60, 61), native gels were incubated with gentle shaking for 8h at 24–27°C in

activity buffer containing 270 mM glycine, 35 mM Tris (pH 8.4), 4 mM ATP, 14 mM

$MgSO_4$, and 0.2% (w/v) $Pb(NO_3)_2$. The gels were rinsed once with water and

images were taken using ImageQuant LAS 4000 mini (Cytiva). White precipitates

were then dissolved by gentle shaking for 8 h at 24–27°C with 50% (v/v) methanol

and 10% (v/v) acetic acid in water. The gels were restained with 0.025% (w/v) CBB

G-250 and 10% acetic acid in water for 80 min at 24–27°C with gentle shaking and

destained with 10% (v/v) ethanol and 10% acetic acid in water for 180 min at

24–27°C with gentle shaking. The gels were rinsed once with water and images

were taken using ImageQuant LAS 4000 mini.

Phosphate-release assay Dimer solution was dialyzed for 8 h using 20 mM

Tris-HCl (pH 7.5), 150 mM NaCl, and 2 mM $MgCl_2$. ATPase activity was assayed by

a continuous spectrophotometric method using a

2-amino-6-mercapto-7-methylpurine ribonucleoside–purine nucleoside

phosphorylase reaction to detect released inorganic phosphate (EnzChek kit;

Thermo Fisher Scientific) (62). The reaction mixture was as follows: 15.7 nM Dimer,

20 mM Tris-HCl (pH 7.5), 150 mM NaCl, 2 mM $MgCl_2$ and 0.01–1 mM ATP in a

total volume of 0.2 mL at 25°C. Sodium azide was added to 15.4 mM final

concentration when the reaction was started. The protein amount of the

MMOB1670 comprising F_1-ATPase β-subunit paralogs was estimated using

densitometric of SDS-PAGE.

Negative-staining EM and image processing. Dimer solution was placed on

a glow-discharged (PIB-10; VACUUM DEVICE, Ibaraki, Japan) carbon-coated grid

(F-400; Nisshin EM Co., Tokyo, Japan) and incubated for 1 min at 24–27°C. The

solution was then removed, and the grid was stained with 2% uranyl acetate (w/v)

for 30 s. The stain was then removed, and the grid was air-dried. To observe

Monomer and Chain, the grids were washed with water after 1 min of incubation

and then treated as described for Dimer solution. Samples were observed using a

transmission EM (JEM1010, JEOL) at 80 kV, equipped with a FastScan-F214 (T)

charge-coupled-device (CCD) camera (TVIPS, Gauting, Germany), and images

were captured at 2.58 Å/pix.

The contrast transfer function parameters for electron micrographs were

estimated using Gctf (63). Further image processing was performed using RELION

3.0 (36). A total of 2148 particles for Dimer and 11687 particles for Monomer were

automatically selected with box sizes of 180 × 180 and 100 × 100 pixels using

reference-based auto-picking, respectively. These particle images were binned to

5.16 Å/pix. For Dimer, the particle images were 2D-classified into 100 classes. For

Monomer, particle images were 2D-classified in four rounds, and the selected 7381

particles were re-extracted with the pixel size returned to the unbinned image and

then 2D-classified into 50 classes.

For reconstruction of the 3D chain structure, 2127 particles were manually

selected for chains with a box size of 276 × 276 pixels with ~50% overlap. These

particle images were binned to 5.16 Å/pix. Particle images were 2D-classified in two

rounds, and the selected 1709 particles were used to reconstruct the initial model

with a final resolution limit of 50 Å. The initial model and selected particles were

used to perform 3D refinement. Reprojection images were produced from the 3D

map using the relion_project command in RELION. The 3D map was visualized

using UCSF Chimera 1.14 (64). Atomic models of F_1-ATPase catalytic $(\alpha\beta)_3$ from *B.*

subtilis (PDB ID 6N2Y) (39) were fitted into the 3D map using command of "Fit in

map" in UCSF Chimera.

HS-AFM. Imaging was performed with a laboratory-built tapping mode HS-AFM

(65, 66), using small cantilevers (BLAC10DS-A2, Olympus; resonant frequency,

~0.5 MHz in water; quality factor, ~1.5 in water; spring constant, ~0.1 N/m). The

cantilever's free-oscillation peak-to-peak amplitude (A_0) and set-point amplitude

were set at ~2.5 nm and ~0.8 × A_0, respectively. The probe tip was grown on the

original tip end of a cantilever through electron beam deposition and further

sharpened using a radio frequency plasma etcher (PE-2000, South Bay Technology,

Redondo Beach, CA) under an argon gas atmosphere (typically at 180 mTorr and

15 W for 3 min). The sample was deposited on a freshly cleaved mica disc glued to

a glass stage beforehand. After 3.5 min, the stage surface was immersed in a liquid

cell containing an observation buffer [20 mM Tris-HCl (pH 7.5), 50 mM KCl, 2 mM

$MgCl_2$]. Imaging was performed at 24–27°C. AFM images were processed using a

low-pass filter to remove spike noise and make the *xy*-plane flat and analyzed using

Kodec software (version 4.4.7.39) (67). Surface profiles and smoothing were

performed using ImageJ software.

Acknowledgments

We thank Toshiaki Arata, Ikuko Fujiwara, Kohei Kobayashi, and Hiroki Sato at the

Graduate School of Science, Osaka City University, and Takayuki Uchihashi at the

Department of Physics and Structural Biology Research Center, Nagoya University,

for helpful discussions. We also thank Aya Takamori at the Graduate School of

Science, Osaka City University, for performing MALDI-TOF MASS spectrometry. TT

learned RELION software in an instruction course on 20180927–28 provided by the

Cyclic Innovation for Clinical Empowerment (CiCLE) from AMED.

This work was supported by a Grant-in-Aid for Scientific Research on the Innovative

Area "Harmonized Supramolecular Motility Machinery and Its Diversity" (MEXT

KAKENHI Grant Number JP24117002), by a Grants-in-Aid for Scientific Research

(B) and (A) (MEXT KAKENHI Grant Numbers JP24390107, JP17H01544), by JST

CREST Grant Number JPMJCR19S5, Japan, by the Osaka City University (OCU)

Strategic Research Grant 2018 for top priority researches and by a Grant-in-aid of

the FUGAKU TRUST FOR MEDICINAL RESEARCH to MM, and JSPS KAKENHI

(Grant Number JP25000013), the Platform Project for Supporting Drug Discovery

and Life Science Research (BINDS) from AMED (Grant Number JP19am0101117

and support number 1282), CiCLE (Grant Number JP17pc0101020), and JEOL

YOKOGUSHI Research Alliance Laboratories of Osaka University to KN.

Figure Legends

FIG 1. Gliding machinery of *Mycoplasma mobile*. (A) Schematic illustration of

gliding machinery based on the present study. In the whole cell shown in upper

illustration, the internal structure and legs are colored gray and red, respectively.

The actual cell has about 28 chains, each consisting of 17 particles, although a

more limited number is illustrated here. In this paper we refer to particle and F_1-like

ATPase unit as the "Dimer" and the "Monomer", respectively. A single unit of the

surface structure and a chain of the internal structure are magnified in the lower

illustration. (B) ORFs for the internal structure. The components of the internal

structure are colored gray. Type 2 ATPase operon is at the top. Dimer components

revealed in the present study are marked by colored boxes, corresponding to the

colored components of the lower illustration in panel A.

FIG 2. Protein profile and characterization of three fractions. (A) Protein profile

of each fraction. Dimer, Monomer, and Chain fractions were subjected to

SDS-12.5% PAGE gel and stained with CBB. The bands marked by black triangles

were identified by peptide mass fingerprinting (PMF). The intensity of band marked

by an asterisk is reduced in Monomer fraction. Molecular masses are shown on the

left.

(B) Gel filtration assay of unit complex fraction using Sephacryl S-400 HR column.

The peak positions of unit complex (Dimer) and thyroglobulin (669 kDa) are marked

by orange and purple triangles. (C) BN-PAGE (left) and In-gel ATPase activity assay

(right). Dimer, Dimer treated with 1.5% sodium cholate (SC treated), and

thyroglobulin, which has no ATPase activity, were subjected to 3 to 12% gradient

BN-PAGE. The bands of Dimer and Monomer after sodium cholate treatment are

marked by closed and open triangles. White lead phosphate bands, indicating

ATPase activity appeared in the right panel. Molecular masses are shown on the left.

(D) Phosphate release assay of Dimer under various ATP concentrations with and

without sodium azide. The ATPase activities under 0- and 15.4-mM sodium azide

are marked by red and green filled circles, respectively ($n = 3$). These data were

fitted by the Michaelis-Menten equation as solid lines.

FIG 3. Negative-staining EM and HS-AFM of Dimer. (A) Electron micrograph of

negatively stained particles in unit complex fraction. (B) Images of individual

particles. (C) Representative 2D averaged image. A mirror image is shown to match

the orientation of the hexameric ring observed by HS-AFM. (D) Illustration based on

the averaged image in panel C. Filamentous structures are marked by asterisks at

an end. The double-headed arrow shows the distance between the centers of the

hexamers. (E) Two patterns of HS-AFM images. A Dimer was scanned at 56 × 56

pixels in an area of 70 × 70 nm^2 with a scanning rate of 100 ms per frame.

Illustrations for patterns I and II (left side) were depicted based on 3D chain model

shown in Fig. 5. Observation directions are indicated by arrows. Protrusions are

marked by blue triangles. Images of patterns I and II are shown in green and purple

frames, respectively. (F) Averaged images for patterns I (green frame) and II (purple

frame). Dimer was scanned at 50 × 50 pixels in an area of 40 × 40 nm^2 with a

scanning rate of 100 ms per frame. The images were produced by averaging three

successive video frames. (G) HS-AFM slice image showing two hexameric rings

(upper) and averaged EM image (lower). Upper: The red broken boxed area in

panel F was sliced for the height 9.8–11.3 nm from the substrate surface, processed

for smoothing, and magnified. The angle alignments of two hexamers are

schematically shown in the left upper. Lower: The central part of panel C was

excised and aligned to compare with the upper panel. Subunits of the hexamer are

marked by red circles. (H) Surface profiles along the lines in pattern I (green) and II

(purple). The upper and lower images in panel F were each profiled at the green

and purple lines passing the globule centers. The dimples and the peaks are

marked by open and black triangles, respectively. The slice height in panel G is

shown by broken lines. (I) Shedding process of the peaks of pattern II particle

shown in panel E. The peaks are marked by red triangles. (J) Surface profile

showing the disappeared peaks. The images in panel I were each profiled at the

blue, pink, and light blue lines passing the globule centers. The peaks are marked

by black triangles. The double-headed arrows show the peak heights. (K) HS-AFM

image of Dimer with seven lateral protrusions. Dimer was scanned at 120 × 120

pixels in an area of 120 × 120 nm^2 with a scanning rate of 500 ms per frame. Lateral

protrusions are indicated by blue triangles. (L) Fluctuations of the protrusions of the

particle shown in panel K. The images were sliced for the height 0–7.0 nm from the

substrate surface. The moving directions are indicated by arrows. In all HS-AFM

images, the color bar on the right shows the range of image heights.

FIG 4. Negative staining EM of Monomer fraction. (A) Electron micrograph of negatively stained ATPase in Monomer fraction. (B) Images of individual particles. (C) Representative 2D averaged images (upper) and depictions of their structures (lower). Upper: Hook structures in the hexameric ring and the stalks are marked by green and orange triangles, respectively. Lower: The hexameric part, hook structures and the stalk are colored rose, green and orange, respectively.

FIG 5. Chain structure. (A) Electron micrograph of negatively stained Chain. The chain structures are marked by blue lines. (B) Magnified chain images. (C) Representative 2D averaged images. (D) Three-dimensional reconstruction of Chain containing two F_1-like ATPase dimers. The 3D map is visualized at a density threshold (contour level = 0.026). The central stalks are marked by open triangles. (E) Superposition of the atomic model of *Bacillus* F_1-ATPase catalytic $(\alpha\beta)_3$ (PDB ID 6N2Y) (39) onto the 3D chain structure. The α and β subunits are colored salmon and turquoise, respectively. The central stalk and protrusions from hexameric rings are marked by open and black triangles, respectively. (F) Cross section of mushroom-like structures. Central stalks are marked by broken circles. The double-headed arrow shows the length of the protrusion. Corresponding mushroom-like structures are marked (a)–(d) in panel E. (G) Comparison between Dimer image from Fig. 3C (left) and the Chain reprojection (middle). The reprojected image is viewed from the angle used for the right image of panel D. Short filaments

corresponding to the position of the connecting bulge are marked by asterisks. A

depiction of Chain model based on the comparison (right). (H) Comparison between

the averaged chain image from ECT (left) and the chain reprojection (middle). Left:

The averaged chain image was modified from the whole cell ECT image in the

previous study (23). Middle: Chain was reprojected from an angle close to the

middle image in panel D. Inner sides and membrane relative to the chain are

marked by i and m, respectively. The protrusion from the globule corresponding to

the central stalk from the hexameric ring, one from the connecting bulge and the

globule attached to the connecting bulge are marked by orange, light blue and

green triangles, respectively. The areas of image densities that were visualized only

in the ECT image are marked by broken lines. Right: An illustration depicts a chain

model based on the comparison.

FIG 6. Working models for force transmission mechanism. The gliding direction

is indicated by a red arrow. The regions marked in gray were visualized only in the

ECT image. The crank protein Gli521 and the leg protein Gli349 are colored green

and red, respectively. (i) Contraction model: The force generated by the hexameric

ring displaces Dimer along the gliding direction. The displacements are transmitted

through the membrane to Gli521. (ii) Rotation model: The force generated by the

hexameric ring rotates the central stalk in a mechanism similar to that of F_1-ATPase.

The rotational motion is transmitted across the membrane to Gli521.The generation

and transmission of forces are presented by black arrows for both models.

FIG S1 Gliding of *M. mobile* cells. Optical microscopy of cells (left) and trajectories of gliding cells (right). All cells are gliding in the direction of tapered end as indicated by a black arrow. For trajectories, video frames of every 0.2 s were colored differently from red to blue, and stacked for 4 s.

FIG S2 Isolation procedure for three fractions. Each fraction was obtained from Pellet-1 fraction. The adjusted concentrations of NaCl in the solution are indicated in parentheses. Supernatant is abbreviated as "Sup". See Materials and Methods for details.

FIG S3 Isolation profile of Dimer (unit complex) and Monomer fractions. (A) Solubility of the chain components. Pellet-1 fraction was treated with buffers containing the specified concentrations of NaCl and centrifuged. Pellet-1 fraction, the supernatants (S) and the pellets (P) were analyzed by SDS-12.5% PAGE. See Materials and Methods for details. (B) Isolation of Dimer (unit complex). The fractions of each step were subjected to SDS-12.5% PAGE gel and stained with CBB. Lane 1, lysate of *M. mobile* cells; lane 2, Triton-soluble fraction; lane 3, Triton-insoluble (Pellet-1) fraction; lane 4, supernatant after incubation in suspension buffer containing 137 mM NaCl; lane 5, peak fraction of Superdex 200 gel filtration chromatography. The bands of Dimer components are marked by black triangles. (C) Gel filtration profile in Monomer isolation using a Sephacryl S-400 HR

column. The small peak is marked by a green triangle. (D) The F1–F13 fractions in

the gel filtration indicated by the green line in panel C were subjected to SDS-12.5%

PAGE and stained with CBB. F6, which corresponds to the small peak in the gel

filtration, was used for further analyses as Monomer fraction. Bands of Dimer

components are marked by black triangles. For panels (A, B, D), Focused proteins

were identified by PMF, and molecular masses are shown on the left.

**FIG S4 Two-dimensional averaged images of Dimer obtained by
negative-staining EM.** Four classes of clear particle images from 20 classes are

represented. These images have common features as represented long and short

protrusions marked by light blue and orange triangles, respectively. As mentioned in

Fig. 3C, the images were mirrored to uniform handedness.

**FIG S5 Two-dimensional averaged images of globular complex in Monomer
fraction.** Fifteen classes of clear particle images from 50 classes are represented.

FIG S6 Reprojection images of Chain. Two-dimensional averaged images (upper)

and the corresponding reprojection images (lower) calculated from the 3D map of

Chain are compared.

Movie S1 HS-AFM movie showing pattern I particles The particles were scanned

at 10 fps. The scanning field was 70 × 70 nm^2 with 56 × 56 pixels. The video was

played at 10 fps.

Movie S2 HS-AFM movie showing pattern II particles The particles were scanned at 10 fps. The scanning field was 70 × 70 nm^2 with 56 × 56 pixels. The video was played at 10 fps.

Movie S3 HS-AFM movie showing the shedding process of the peaks. The particles were scanned at 10 fps. The scanning field was 70 × 70 nm^2 with 56 × 56 pixels. The video was played at 10 fps. The peaks are indicated by the red triangles.

Movie S4 HS-AFM movie showing fluctuations in protrusions. The particles were scanned at 2 fps. The scanning field was 120 × 120 nm^2 with 120 × 120 pixels. The video was played at 1 fps. The protrusions are indicated by the red triangles.

References

1.   Razin S, Hayflick L. 2010. Highlights of mycoplasma research--an historical perspective. Biologicals 38:183-90.

2.   Grosjean H, Breton M, Sirand-Pugnet P, Tardy F, Thiaucourt F, Citti C, Barre A, Yoshizawa S, Fourmy D, de Crecy-Lagard V, Blanchard A. 2014. Predicting the minimal translation apparatus: lessons from the reductive evolution of mollicutes. PLoS Genet 10:e1004363.

3.   Miyata M, Robinson RC, Uyeda TQP, Fukumori Y, Fukushima SI, Haruta S,

Homma M, Inaba K, Ito M, Kaito C, Kato K, Kenri T, Kinosita Y, Kojima S,

Minamino T, Mori H, Nakamura S, Nakane D, Nakayama K, Nishiyama M,

Shibata S, Shimabukuro K, Tamakoshi M, Taoka A, Tashiro Y, Tulum I, Wada

H, Wakabayashi KI. 2020. Tree of motility - A proposed history of motility

systems in the tree of life. Genes Cells 25:6-21.

4.   Miyata M. 2010. Unique centipede mechanism of *Mycoplasma* gliding. Annu

Rev Microbiol 64:519-37.

5.   Miyata M, Hamaguchi T. 2016. Prospects for the gliding mechanism of

Mycoplasma mobile. Curr Opin Microbiol 29:15-21.

6.   Miyata M, Hamaguchi T. 2016. Integrated information and prospects for

gliding mechanism of the pathogenic bacterium *Mycoplasma pneumoniae*.

Front Microbiol 7:960.

7.   Uenoyama A, Kusumoto A, Miyata M. 2004. Identification of a 349-kilodalton

protein (Gli349) responsible for cytadherence and glass binding during

gliding of *Mycoplasma mobile*. J Bacteriol 186:1537-45.

8.   Seto S, Uenoyama A, Miyata M. 2005. Identification of a 521-kilodalton

protein (Gli521) involved in force generation or force transmission for

Mycoplasma mobile gliding. J Bacteriol 187:3502-10.

9.   Uenoyama A, Miyata M. 2005. Identification of a 123-kilodalton protein

(Gli123) involved in machinery for gliding motility of *Mycoplasma mobile*. J

Bacteriol 187:5578-84.

10. Uenoyama A, Seto S, Nakane D, Miyata M. 2009. Regions on Gli349 and

764 Gli521 protein molecules directly involved in movements of *Mycoplasma*

765 *mobile* gliding machinery, suggested by use of inhibitory antibodies and

766 mutants. J Bacteriol 191:1982-5.

767 11. Kusumoto A, Seto S, Jaffe JD, Miyata M. 2004. Cell surface differentiation of

768 *Mycoplasma mobile* visualized by surface protein localization. Microbiology

769 150:4001-8.

770 12. Miyata M, Petersen JD. 2004. Spike structure at the interface between

771 gliding *Mycoplasma mobile* cells and glass surfaces visualized by

772 rapid-freeze-and-fracture electron microscopy. J Bacteriol 186:4382-6.

773 13. Adan-Kubo J, Uenoyama A, Arata T, Miyata M. 2006. Morphology of isolated

774 Gli349, a leg protein responsible for *Mycoplasma mobile* gliding via glass

775 binding, revealed by rotary shadowing electron microscopy. J Bacteriol

776 188:2821-8.

777 14. Lesoil C, Nonaka T, Sekiguchi H, Osada T, Miyata M, Afrin R, Ikai A. 2010.

778 Molecular shape and binding force of *Mycoplasma mobile*'s leg protein

779 Gli349 revealed by an AFM study. Biochem Biophys Res Commun

780 391:1312-7.

781 15. Metsugi S, Uenoyama A, Adan-Kubo J, Miyata M, Yura K, Kono H, Go N.

782 2005. Sequence analysis of the gliding protein Gli349 in *Mycoplasma mobile*.

783 Biophysics (Nagoya-shi) 1:33-43.

784 16. Hamaguchi T, Kawakami M, Furukawa H, Miyata M. 2019. Identification of

785 novel protein domain for sialyloligosaccharide binding essential to

Mycoplasma mobile gliding. FEMS Microbiol Lett 366:fnz016.

17.   Morio H, Kasai T, Miyata M. 2016. Gliding direction of *Mycoplasma mobile*. J

Bacteriol 198:283-90.

18.   Kasai T, Hamaguchi T, Miyata M. 2015. Gliding motility of *Mycoplasma*

mobile on uniform oligosaccharides. J Bacteriol 197:2952-7.

19.   Kasai T, Nakane D, Ishida H, Ando H, Kiso M, Miyata M. 2013. Role of

binding in *Mycoplasma mobile* and *Mycoplasma pneumoniae* gliding

analyzed through inhibition by synthesized sialylated compounds. J Bacteriol

195:429-35.

20.   Nagai R, Miyata M. 2006. Gliding motility of *Mycoplasma mobile* can occur by

repeated binding to *N*-acetylneuraminyllactose (sialyllactose) fixed on solid

surfaces. J Bacteriol 188:6469-75.

21.   Chen J, Neu J, Miyata M, Oster G. 2009. Motor-substrate interactions in

Mycoplasma motility explains non-Arrhenius temperature dependence.

Biophys J 97:2930-8.

22.   Mizutani M, Tulum I, Kinosita Y, Nishizaka T, Miyata M. 2018. Detailed

analyses of stall force generation in *Mycoplasma mobile* gliding. Biophys J

114:1411-1419.

23.   Nishikawa M, Nakane D, Toyonaga T, Kawamoto A, Kato T, Namba K, Miyata

M. 2019. Refined mechanism of *Mycoplasma mobile* gliding based on

structure, ATPase activity, and sialic acid binding of machinery. mBio

10:e02846-19.

24.   Nakane D, Miyata M. 2007. Cytoskeletal "jellyfish" structure of *Mycoplasma*

809   *mobile*. Proc Natl Acad Sci U S A 104:19518-23.

810   25.   Tulum I, Yabe M, Uenoyama A, Miyata M. 2014. Localization of P42 and

811   F_1-ATPase alpha-subunit homolog of the gliding machinery in *Mycoplasma*

812   *mobile* revealed by newly developed gene manipulation and fluorescent

813   protein tagging. J Bacteriol 196:1815-24.

814   26.   Tulum I, Kimura K, Miyata M. 2020. Identification and sequence analyses of

815   the gliding machinery proteins from *Mycoplasma mobile*. Sci Rep 10:3792.

816   27.   Walker JE, Saraste M, Runswick MJ, Gay NJ. 1982. Distantly related

817   sequences in the alpha- and beta-subunits of ATP synthase, myosin, kinases

818   and other ATP-requiring enzymes and a common nucleotide binding fold.

819   Embo j 1:945-51.

820   28.   Uenoyama A, Miyata M. 2005. Gliding ghosts of *Mycoplasma mobile*. Proc

821   Natl Acad Sci USA 102:12754-8.

822   29.   Kinosita Y, Nakane D, Sugawa M, Masaike T, Mizutani K, Miyata M,

823   Nishizaka T. 2014. Unitary step of gliding machinery in Mycoplasma mobile.

824   Proc Natl Acad Sci U S A 111:8601-6.

825   30.   Abrahams JP, Leslie AG, Lutter R, Walker JE. 1994. Structure at 2.8 A

826   resolution of F1-ATPase from bovine heart mitochondria. Nature 370:621-8.

827   31.   Kühlbrandt W. 2019. Structure and Mechanisms of F-Type ATP Synthases.

828   Annu Rev Biochem 88:515-549.

32.   Beven L, Charenton C, Dautant A, Bouyssou G, Labroussaa F, Skollermo A,

Persson A, Blanchard A, Sirand-Pugnet P. 2012. Specific evolution of F_1-like

ATPases in mycoplasmas. PLoS One 7:e38793.

33. Nottelet P, Bataille L, Gourgues G, Anger R, Lartigue C, Sirand-Pugnet P,

Marza E, Fronzes R, Arfi Y. 2021. The mycoplasma surface proteins MIB and

MIP promote the dissociation of the antibody-antigen interaction. Sci Adv 7.

34. Kobayashi K, Kodera N, Kasai T, Tahara YO, Toyonaga T, Mizutani M,

Fujiwara I, Ando T, Miyata M. 2021. Movements of *Mycoplasma mobile*

gliding machinery detected by high-speed atomic force microscopy. mBio in

press:mBio00040-21.

35. Bowler MW, Montgomery MG, Leslie AG, Walker JE. 2006. How azide

inhibits ATP hydrolysis by the F-ATPases. Proc Natl Acad Sci U S A

103:8646-9.

36. Zivanov J, Nakane T, Forsberg BO, Kimanius D, Hagen WJ, Lindahl E,

Scheres SH. 2018. New tools for automated high-resolution cryo-EM

structure determination in RELION-3. eLife 7:e42166.

37. Ando T. 2018. High-speed atomic force microscopy and its future prospects.

Biophys Rev 10:285-292.

38. Uchihashi T, Iino R, Ando T, Noji H. 2011. High-speed atomic force

microscopy reveals rotary catalysis of rotorless F_1-ATPase. Science

333:755-8.

39. Guo H, Suzuki T, Rubinstein JL. 2019. Structure of a bacterial ATP synthase.

eLife 8:e43128.

40.   Cheng Y, Wolf E, Larvie M, Zak O, Aisen P, Grigorieff N, Harrison SC, Walz T.

853   2006. Single particle reconstructions of the transferrin-transferrin receptor

854   complex obtained with different specimen preparation techniques. J Mol Biol

855   355:1048-65.

856   41.   Dibrova DV, Konovalov KA, Perekhvatov VV, Skulachev KV, Mulkidjanian AY.

857   2017. COGcollator: a web server for analysis of distant relationships

858   between homologous protein families. Biol Direct 12:29.

859   42.   Neuwald AF, Aravind L, Spouge JL, Koonin EV. 1999. AAA+: A class of

860   chaperone-like ATPases associated with the assembly, operation, and

861   disassembly of protein complexes. Genome Res 9:27-43.

862   43.   Snider J, Thibault G, Houry WA. 2008. The AAA+ superfamily of functionally

863   diverse proteins. Genome Biol 9:216.

864   44.   Minauro-Sanmiguel F, Wilkens S, García JJ. 2005. Structure of dimeric

865   mitochondrial ATP synthase: novel F0 bridging features and the structural

866   basis of mitochondrial cristae biogenesis. Proc Natl Acad Sci U S A

867   102:12356-8.

868   45.   Allegretti M, Klusch N, Mills DJ, Vonck J, Kühlbrandt W, Davies KM. 2015.

869   Horizontal membrane-intrinsic α-helices in the stator a-subunit of an F-type

870   ATP synthase. Nature 521:237-40.

871   46.   Blum TB, Hahn A, Meier T, Davies KM, Kühlbrandt W. 2019. Dimers of

872   mitochondrial ATP synthase induce membrane curvature and self-assemble

873   into rows. Proc Natl Acad Sci U S A 116:4250-4255.

874   47.   Noji H, Ueno H, McMillan DGG. 2017. Catalytic robustness and torque

875   generation of the F_1-ATPase. Biophys Rev 9:103-118.

876   48.   Naito TM, Masaike T, Nakane D, Sugawa M, Okada KA, Nishizaka T. 2019.

877   Single-molecule pull-out manipulation of the shaft of the rotary motor

878   F_1-ATPase. Sci Rep 9:7451.

879   49.   Watson HC, Walker NP, Shaw PJ, Bryant TN, Wendell PL, Fothergill LA,

880   Perkins RE, Conroy SC, Dobson MJ, Tuite MF. 1982. Sequence and

881   structure of yeast phosphoglycerate kinase. Embo j 1:1635-40.

882   50.   Jaffe JD, Stange-Thomann N, Smith C, DeCaprio D, Fisher S, Butler J, Calvo

883   S, Elkins T, FitzGerald MG, Hafez N, Kodira CD, Major J, Wang S, Wilkinson

884   J, Nicol R, Nusbaum C, Birren B, Berg HC, Church GM. 2004. The complete

885   genome and proteome of *Mycoplasma mobile*. Genome Res 14:1447-61.

886   51.   Lu M, Holliday LS, Zhang L, Dunn WA, Jr., Gluck SL. 2001. Interaction

887   between aldolase and vacuolar H^+-ATPase: evidence for direct coupling of

888   glycolysis to the ATP-hydrolyzing proton pump. J Biol Chem 276:30407-13.

889   52.   Lu M, Ammar D, Ives H, Albrecht F, Gluck SL. 2007. Physical interaction

890   between aldolase and vacuolar H^+-ATPase is essential for the assembly and

891   activity of the proton pump. J Biol Chem 282:24495-503.

892   53.   Chan CY, Dominguez D, Parra KJ. 2016. Regulation of Vacuolar H^+-ATPase

893   (V-ATPase) Reassembly by Glycolysis Flow in 6-Phosphofructo-1-kinase

894   (PFK-1)-deficient Yeast Cells. J Biol Chem 291:15820-9.

895   54.   Kinosita Y, Miyata M, Nishizaka T. 2018. Linear motor driven-rotary motion of

896   a membrane-permeabilized ghost in *Mycoplasma mobile*. Sci Rep 8:11513.

897   55.   Miyata M, Yamamoto H, Shimizu T, Uenoyama A, Citti C, Rosengarten R.

898   2000. Gliding mutants of *Mycoplasma mobile*: relationships between motility

899   and cell morphology, cell adhesion and microcolony formation. Microbiology

900   146:1311-20.

901   56.   Aluotto BB, Wittler RG, Williams CO, Faber JE. 1970. Standardized

902   bacteriologic techniques for the characterization of *Mycoplasma* species. Int

903   J Syst Bacteriol 20:35-58.

904   57.   Kawakita Y, Kinoshita M, Furukawa Y, Tulum I, Tahara YO, Katayama E,

905   Namba K, Miyata M. 2016. Structural study of MPN387, an essential protein

906   for gliding motility of a human-pathogenic bacterium, *Mycoplasma*

907   *pneumoniae*. J Bacteriol 198:2352-9.

908   58.   Lee C, Levin A, Branton D. 1987. Copper staining: a five-minute protein stain

909   for sodium dodecyl sulfate-polyacrylamide gels. Anal Biochem 166:308-12.

910   59.   Dzandu JK, Johnson JF, Wise GE. 1988. Sodium dodecyl sulfate-gel

911   electrophoresis: staining of polypeptides using heavy metal salts. Anal

912   Biochem 174:157-67.

913   60.   Zerbetto E, Vergani L, Dabbeni-Sala F. 1997. Quantification of muscle

914   mitochondrial oxidative phosphorylation enzymes via histochemical staining

915   of blue native polyacrylamide gels. Electrophoresis 18:2059-64.

916   61.   Wittig I, Schägger H. 2005. Advantages and limitations of clear-native PAGE.

917   Proteomics 5:4338-46.

918   62.   Webb MR. 1992. A continuous spectrophotometric assay for inorganic

919   phosphate and for measuring phosphate release kinetics in biological

920   systems. Proc Natl Acad Sci U S A 89:4884-7.

921   63.   Zhang K. 2016. Gctf: Real-time CTF determination and correction. J Struct

922   Biol 193:1-12.

923   64.   Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC,

924   Ferrin TE. 2004. UCSF Chimera--a visualization system for exploratory

925   research and analysis. J Comput Chem 25:1605-12.

926   65.   Ando T, Kodera N, Takai E, Maruyama D, Saito K, Toda A. 2001. A

927   high-speed atomic force microscope for studying biological macromolecules.

928   Proc Natl Acad Sci U S A 98:12468-72.

929   66.   Uchihashi T, Kodera N, Ando T. 2012. Guide to video recording of structure

930   dynamics and dynamic processes of proteins by high-speed atomic force

931   microscopy. Nat Protoc 7:1193-206.

932   67.   Ngo KX, Kodera N, Katayama E, Ando T, Uyeda TQ. 2015. Cofilin-induced

933   unidirectional cooperative conformational changes in actin filaments revealed

934   by high-speed atomic force microscopy. eLife 4:e04806.

Figure 1

A

Chain

Chain
Dimer (Particle)
Monomer (F_1-like ATPase unit)

Membrane
Surface structure

Gli123
Gli521
Gli349
Foot

Sialylated oligosaccharides

B

MMOB
1610 1620 1630 1640 1650 1660 1670

0120 0130 0140 0150 4530

4830 4840 4850 4860 5430 5440

⇨ Component ▮ Transmembrane segment ▮ Walker A motif (P-loop)
▮ Walker B motif ▨ Extra N-terminal region

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Cell membrane

SOs

(i) Contraction model

(ii) Rotation model

FIG S1 Gliding of *M. mobile* cells. Optical microscopy of cells (left) and trajectories of gliding cells (right). All cells are gliding in the direction of tapered end as indicated by a black arrow. For trajectories, video frames of every 0.2 s were colored differently from red to blue, and stacked for 4 s.

FIG S2 Isolation procedure for three fractions. Each fraction was obtained from Pellet-1 fraction. The adjusted concentrations of NaCl in the solution are indicated in parentheses. Supernatant is abbreviated as "Sup". See Materials and Methods for details.

FIG S3 Isolation profile of Dimer (unit complex) and Monomer fractions. (A) Solubility of the chain components. Pellet-1 fraction was treated with buffers containing the specified concentrations of NaCl and centrifuged. Pellet-1 fraction, the supernatants (S) and the pellets (P) were analyzed by SDS-12.5% PAGE. See Materials and Methods for details. (B) Isolation of Dimer (unit complex). The fractions of each step were subjected to SDS-12.5% PAGE gel and stained with CBB. Lane 1, lysate of *M. mobile* cells; lane 2, Triton-soluble fraction; lane 3, Triton-insoluble (Pellet-1) fraction; lane 4, supernatant after incubation in suspension buffer containing 137 mM NaCl; lane 5, peak fraction of Superdex 200 gel filtration chromatography. The bands of Dimer components are marked by black triangles. (C) Gel filtration profile in Monomer isolation using a Sephacryl S-400 HR column. The small peak is marked by a green triangle. (D) The F1– F13 fractions in the gel filtration indicated by the green line in panel C were subjected to SDS-12.5% PAGE and stained with CBB. F6, which corresponds to the small peak in the gel filtration, was used for further analyses as Monomer fraction. Bands of Dimer components are marked by black triangles. For panels (A, B, D), Focused proteins were identified by PMF, and molecular masses are shown on the left.

10 nm

FIG S4 Two-dimensional averaged images of Dimer obtained by negative-staining EM. Four classes of clear particle images from 20 classes are represented. These images have common features as represented long and short protrusions marked by light blue and orange triangles, respectively. As mentioned in Fig. 3C, the images were mirrored to uniform handedness.

10 nm

FIG S5 Two-dimensional averaged images of globular complex in Monomer fraction. Fifteen classes of clear particle images from 50 classes are represented.

2D averages

Reprojections

20 nm

FIG S6 Reprojection images of Chain. Two-dimensional averaged images (upper) and the corresponding reprojection images (lower) calculated from the 3D map of Chain are compared.