

Rab10 regulates the sorting of internalised TrkB to retrograde axonal transport

Authors: Oscar M. Lazo^{1,2} and Giampietro Schiavo^{1,2} *

Affiliations:

¹ Department of Neuromuscular Diseases, UCL Queen Square Institute of Neurology, University College London, London WC1N 3BG, UK

² UK Dementia Research Institute at UCL, London WC1E 6BT, UK

* To whom correspondence should be addressed: Giampietro Schiavo (giampietro.schiavo@ucl.ac.uk)

Word count main text: 4015

Figures: 6

References: 64

Supplementary tables: 2

Key words: Brain-derived neurotrophic factor, signalling endosome, small GTPases

Abstract

The extreme, complex morphology of neurons provides an unrivalled model to study the coordination between local signalling and long-range cell responses. A cogent example is provided by the binding of brain-derived neurotrophic factor (BDNF) to its receptor TrkB, which triggers signalling cascades at axon terminals that result in responses at the level of the cell body, including modulation of gene expression. Retrograde propagation of these critical signals relies on the sorting of activated TrkB receptors to retrograde axonal transport organelles termed signalling endosomes. In this work, we show that the small GTPase Rab10 is critical for the sorting of activated TrkB receptors to axonal retrograde carriers and the propagation of neurotrophin signalling from the axon terminal to the soma. Moreover, our data indicate that Rab10 defines a novel class of axonal organelles that are mobilised towards the axon terminal upon BDNF stimulation, thus enabling the axon to dynamically adjust the retrograde signalling flow to changes in BDNF availability at the synapse.

Introduction

Communication between cells depends on their ability to respond as integrated units to spatial and temporal signalling patterns. The complex morphology of neurons provides an unrivalled model to study how sorting and trafficking of signalling complexes coordinate local signalling at the axon terminal and the propagation of messages to the cell body. A clear example is provided by neurotrophic factors secreted by target tissues and sensed by axon terminals, where, among other functions, they regulate local cytoskeletal dynamics to promote or cease axon elongation, induce branching, as well as maturation and plasticity of synapses¹⁻³. In parallel, a population of activated receptors are internalised and targeted to the retrograde axonal transport pathway within signalling endosomes^{4,5}, which propagate neurotrophic signalling towards the cell body, regulating gene expression, dendritic branching, and the equilibrium between survival and apoptosis⁶⁻⁹. How local and central responses are coordinated across the massive distance from axon terminals to the soma is a crucial and still unanswered question for neuronal cell biology.

Brain-derived neurotrophic factor (BDNF), a member of the neurotrophin family, is widely expressed in the central nervous system, together with its receptor tropomyosin-related kinase B (TrkB). By activating diverse signalling cascades, comprising the phosphoinositide 3-kinase (PI3K)-Akt pathway, mitogen-activated protein kinases (MAPKs) and phospholipase C-gamma (PLCγ), BDNF and TrkB play a critical role in the formation, maintenance and plasticity of neuronal circuits^{10,11}. Binding of BDNF at synaptic sites leads to endocytosis of TrkB and accumulation of the activated ligand-receptor complex in early endosomes at axon terminals¹². Whilst part of the internalised TrkB receptors recycle back to the plasma membrane, a pool of receptors is sorted to signalling endosomes and engage with the cytoplasmic dynein motor, which mediates the transport of these organelles to the soma^{1,9,13}. Several lines of evidence indicate that this compartment is generated from early endosomes that then mature into a more specialised organelle that escape acidification and lysosomal degradation⁵. Sorting of TrkB receptors from early to signalling endosomes constitutes the critical regulatory node controlling the intensity of the retrogradely-propagated signal; however, to date no clear mechanism coordinating this sorting process has been proposed.

The Rab family of monomeric GTPases plays a central role regulating post-endocytic trafficking of TrkB. Whilst early endosome formation is regulated by Rab5, maturation and processive transport of signalling endosomes in different neuronal models are controlled by Rab7^{12,14-16}. However, it is currently unclear whether or not Rab7 is the only member of the Rab family controlling the latter process. Because axonal retrograde signalling endosomes appear to be a diverse group of organelles⁵, we hypothesised that other members of the Rab family also contribute to the segregation of TrkB and sorting to retrograde axonal carriers.

In this work, we specifically focused on Rab10, since it has been shown that Rab10-positive organelles are transported both anterogradely and retrogradely along the axon in hippocampal neurons¹⁷. Previous work from our laboratory using an affinity purification approach to isolate neurotrophin signalling endosomes from mouse embryonic stem cell-derived motor neurons found that Rab10 was significantly enriched in this axonal compartment, together with Rab5 and Rab7¹⁸. By manipulating Rab10 expression and activity in hippocampal neurons, as well as analysing the axonal dynamics of Rab10 organelles, we have explored its ability to regulate the sorting of TrkB to the retrograde axonal transport pathway and respond to increasing concentrations of BDNF, adjusting retrograde signalling on demand.

Results

Rab10 is required for retrograde TrkB trafficking and signalling

To study axonal TrkB dynamics, we cultured hippocampal neurons in microfluidic chambers, which allow cellular and fluidic compartmentalisation of axon terminals. After 7 days in vitro, neurons plated in one of the compartments (designated as 'somatic') displayed axons reaching the axonal compartment (Figure 1a). Maintaining a higher volume of media in the somatic compartment allows a micro-flow along the grooves, keeping any substance added to the axonal compartment confined to this area. Using this experimental set up, we incubated axon terminals with an antibody against the extracellular domain of TrkB and induced its endocytosis by adding 20 ng/mL BDNF. After 2.5 hours, we were able to detect the axonal TrkB in the cell body of neurons stimulated with BDNF, but not in neurons depleted of BDNF by addition of an anti-BDNF blocking antibody (Figure 1b), confirming the reliability of our retrograde accumulation assay.

The same experimental design was applied to neurons transduced with a doxycycline-inducible lentivirus encoding shRNA targeting Rab10. Neurons treated with the shRNA for 18-22 hours showed lower levels of Rab10 (Figure 1c, grey) and significantly reduced retrograde accumulation of TrkB when compared with neurons expressing control lentivirus (Figure 1c-d). Since expression of Rab10 is variable among neurons, we tested the correlation between the abundance of endogenous Rab10 with the levels of retrograde accumulation of TrkB (Figure 1e). We found that, even though there was a large overlap, control and knocked-down neurons clustered as expected. Moreover, when we take both populations together, we found a significant correlation between Rab10 expression and TrkB accumulation, strongly suggesting that Rab10 plays a role in retrograde axonal transport of internalised TrkB.

To confirm the functional consequences of this decrease on the retrograde transport of TrkB, we treated neurons in the axonal compartment with BDNF and analysed the levels of phosphorylated cAMP response element binding protein (pCREB) in the nucleus. pCREB is a well-established proxy for neurotrophic signalling in neurons and has been shown to be critical for global neuronal responses to neurotrophins, such as BDNF-induced dendritic branching¹⁹. Neurons treated with a shRNA directed against Rab10 showed a significant decrease in nuclear pCREB (Figure 1f-g), which was completely rescued by treatment with a lentivirus encoding shRNA-resistant Rab10, further confirming the specificity of this effect.

Rab10 associates transiently to TrkB-containing retrograde carriers

Given its critical role regulating retrograde accumulation of TrkB, we investigated whether Rab10 was present in retrograde signalling endosomes. As stated before, signalling endosomes are likely to be a heterogeneous collection of functionally-related organelles with diverse molecular composition⁵. Most of these axonal carriers are positive for Rab7¹², which enables the recruitment of the dynein motor complex and their processive retrograde transport along microtubules^{9,13}. Endogenous Rab10 (orange) and Rab7 (green) were detected with specific antibodies and imaged by confocal microscopy (Figure 2a). Co-localisation was evaluated using Manders index and significance determined using confined-displacement algorithm (CDA)²⁰. Rab10 and Rab7 were found to exhibit very low co-localisation levels (Figure 2a, yellow arrowheads), both in neurons stimulated with BDNF (20 ng/mL for 30 min), as well as in neurons depleted of BDNF by treatment a blocking antibody (Figure 2b). Super-resolution radial fluctuations (SRRF) microscopy was used to examine the distribution of Rab10, Rab7 and Rab5 along the axon of neurons stimulated with BDNF as above. Consistently, we found that Rab10 and Rab7 showed a very low degree of overlap (see yellow regions in Figure 2c and superposition of intensity peaks in 2d). On the other hand, super resolution microscopy showed few organelles where Rab10 and Rab5 partially co-localise (pink regions, Figure 2c-d), suggesting that Rab5-positive retrograde carriers²¹ or stationary early endosomes could contain Rab10.

To identify retrograde signalling endosomes containing TrkB, we took advantage of the property of the non-toxic carboxy-terminal domain of the heavy chain of tetanus neurotoxin (HcT) of being transported almost exclusively in retrograde axonal organelles in neurons¹². We co-internalised HcT-AlexaFluor647 and mouse anti-Flag antibodies in neurons transfected with a TrkB-Flag construct, and stimulated them with BDNF. Double-positive TrkB/HcT puncta in the axon were considered retrograde signalling endosomes, and we analysed the proportion of them co-localising with Rab10. Figure 2e represents the relative area of TrkB/HcT and the triple positive area of TrkB/HcT/Rab10 across three different time points (30, 60 and 90 minutes) post-endocytosis. Whereas the amount of TrkB in retrograde carriers increases with time (cyan) as expected, the proportion of TrkB present in Rab10 organelles remains low and does not increase any further after 60 minutes of incubation (white). These results suggest that TrkB localises transiently on Rab10-positive organelles *en route* to its delivery to axonal retrograde carriers. This observation is supported by super-resolution images (Figure 2f), showing an increase of TrkB/HcT double-positive puncta (cyan arrowheads), but very few TrkB/HcT/Rab10 triple-positive puncta (white empty triangles) at 60 and 90 minutes.

Overexpressed Rab10 is co-transported with retrograde TrkB

The data presented so far support a model in which Rab10 is critical for retrograde transport of TrkB, but does not specifically define a stable population of retrograde carriers. Rather, the evidence shown in Figure 2e-f suggests that TrkB is transiently associated to Rab10-positive organelles, opening the possibility that Rab10 participates in the sorting of internalised receptors to the retrograde axonal transport compartment. Therefore, we decided to overexpress Rab10-EGFP, since it is well documented that overexpression of Rab GTPases promotes their activity due to increased concentration in the membrane²². We expected that an increase in the abundance and/or activity of Rab10 stabilises its presence in TrkB retrograde carriers for enough time to give us the chance to visualise this potential interaction in live neurons.

To test this hypothesis, neurons were co-transfected with Rab10-EGFP and TrkB-Flag and, after 1 hour of starvation in non-supplemented Neurobasal, we internalised AlexaFluor647-labelled anti-Flag antibodies in the presence of BDNF for 45 min (Figure 3a). Live-cell confocal imaging of axon segments was performed at least 200 µm from the cell body. Representative frames of the time-lapse movie are shown in Figure 3b, where

retrograde co-transport of TrkB and Rab10-EGFP is indicated with yellow arrowheads. Kymographs (Figure 3c) were generated for Rab10-EGFP (orange) and TrkB-Flag (green). Tracks are shown in the bottom panel, with examples of co-transport as yellow lines. Quantification of five independent experiments (Figure 3d) confirmed that approximately 60% of retrograde TrkB carriers are positive for Rab10. Interestingly, no anterograde TrkB/Rab10 double-positives compartments were observed, suggesting that TrkB is present in organelles with a strong retrograde bias.

To extend this analysis to the other neurotrophin receptor responding to BDNF, a similar experiment was done by monitoring the internalisation of endogenous p75^{NTR} in neurons transfected with Rab10-EGFP. After depletion of trophic factors, p75^{NTR} uptake was visualised by incubating neurons for 45 min with an AlexaFluor647-labelled antibody against the extracellular domain of p75^{NTR} in the presence of BDNF (Figure 3e). Live-cell imaging of axon segments was done under conditions identical to those for TrkB. Representative frames and kymographs of Rab10-EGFP (orange) and internalised p75^{NTR} (green) are shown in the Figure 3f-g. Kymograph traces show that p75^{NTR} receptor can be found in both retrograde and anterograde Rab10 organelles. On average, 29.6% of anterograde and 15.5% of retrograde p75^{NTR} carriers were found positive for Rab10 across five independent experiments (Figure 3h).

Increasing BDNF favours anterograde trafficking of Rab10 organelles

In agreement with the bidirectional transport observed for p75^{NTR} organelles (Figure 3h), previous work suggested that Rab10-positive compartments are transported along the axon in both directions¹⁷. In light of these results, we hypothesised that the dynamics of Rab10 organelles may respond to BDNF signalling to regulate the sorting of TrkB to retrograde carriers. We therefore examined the axonal transport of Rab10-positive organelles in hippocampal neurons under two opposite conditions: depletion of BDNF using an anti-BDNF blocking antibody, followed by stimulation with 50 ng/mL BDNF.

Figure 4a shows representative frames of a Rab10-EGFP organelle moving in the retrograde direction along the axon (white arrows) in the absence of BDNF. Five-minutes segments of video microscopy have been integrated in each kymograph (Figure 4b) at different time points of BDNF stimulation: before BDNF (top), immediately after BDNF addition (centre) and after 10 min of BDNF incubation (bottom). Traces have been colour-coded as retrograde (cyan), anterograde (pink) or stationary/bidirectional (yellow) to reveal changes in the direction bias of the Rab10 organelles in the same axon, before and after the addition of BDNF. Quantification of five independent experiments shows that BDNF-depleted axons exhibit a bias towards retrograde Rab10 transport, which significantly switches to anterograde after 10 min of stimulation with BDNF.

These surprising results reveals a novel mechanism ensuring a tight balance between retrograde and anterograde transport of Rab10 organelles, which is fine-tuned by extracellular BDNF levels. As a consequence, any local increase in BDNF release from post-synaptic compartments will escalate the abundance of Rab10 organelles in the immediate vicinity, either by mobilising or retaining them at the distal axonal sites.

Rab10 regulates the sorting of TrkB in early endosomes, with no effect on recycling

After endocytosis, TrkB accumulates in early endosomes, from which it is sorted either to the recycling route or to endosomal organelles with signalling capabilities^{9,12}. Rab10, on the other hand, has been shown to regulate trafficking across early endosomes, the formation of specialised tubular endosomes, recycling of cargoes back to the plasma membrane, as well as targeting of plasmalemmal precursor vesicles (PPVs), among other functions^{17,23-26}. To understand the mechanism linking Rab10-positive compartments with the

retrograde axonal transport of TrkB in signalling endosomes, we designed experiments to discern between a potential role of Rab10 on recycling of internalised TrkB back to the plasma membrane, and sorting of TrkB out to the early endosome into retrograde transport carriers (Figure 5a).

To assess the contribution of Rab10 to the recycling of TrkB in the axon, hippocampal neurons were transfected with EGFP or a dominant negative mutant of Rab10 (Rab10^{T23N}; referred as Rab10DN) and TrkB-Flag. Endocytosis of anti-Flag M1 antibodies was allowed for 30 minutes in the presence of BDNF, and then the remaining antibody still bound to the neuronal surface was removed using EDTA, which dissociates this antibody from the bound Flag peptide²⁷. Recycling receptors were then chased using AlexaFluor647-conjugated secondary antibodies. After fixation, internalised TrkB-Flag was revealed using AlexaFluor555-conjugated secondary antibodies (Figure 5b). Comparison between recycling ratio (recycled/internalised) of EGFP- and Rab10DN-expressing neurons shows no significant differences (Figure 5c).

In contrast, if the sorting of TrkB to signalling endosomes was regulated by Rab10, expression of Rab10DN would result in the trapping of TrkB in early endosomes, driving an increase of co-localisation between internalised TrkB and Rab5. Therefore, we incubated neurons transfected with TrkB-Flag and either EGFP or Rab10DN, with anti-Flag antibodies for 30 min in the presence of BDNF, and then analysed TrkB/Rab5 co-localisation using Manders index and CDA (Figure 5d-e). Rab10DN caused a statistically significant increase of internalised TrkB in Rab5-positive domains from $13.2\% \pm 1.7$ to $22.2\% \pm 2.8$, and an increase in the reliability of the co-localisation measurements (CDA $p < 0.05$) (Figure 5e). These results indicate that the probability of finding axonal TrkB in a Rab5-positive early endosome increases in neurons expressing Rab10DN, thus indicating that Rab10 modulates the sorting of activated TrkB receptors from axonal early endosomes to retrograde transport organelles.

Discussion

Our results unravel a novel role for Rab10 in regulating the sorting of internalised TrkB receptors to the retrograde axonal transport pathway. This function appears to be necessary not only for efficient trafficking of TrkB from axons to soma, but also for the propagation of neurotrophic signalling to the nucleus, as shown by the decrease of BDNF-induced CREB activation upon Rab10 knock down (Figure 1). CREB is a transcription factor well known for supporting survival, differentiation and morphology of neurons, which is activated by phosphorylation downstream to the Akt and MAPK branches of neurotrophic signalling²⁸. Some of the better characterised early response genes to neurotrophic factors (e.g., Egr1, Egr2, Arc and cFos) are transcriptional targets of CREB, which is also required for BDNF-induced dendritic branching^{19,29,30}. Therefore, CREB phosphorylation is one of the best proxies for global cell responses to neurotrophic signalling. Long-distance activation of CREB has been reported from distal axons and dendrites, and trafficking across endosomal compartments of pCREB has been shown to play a crucial role in propagating neurotrophin signalling from the periphery to the nucleus^{8,31,32}. Interestingly, in our study we observed that expression of functional Rab10 was critical for maintenance and survival of differentiated neurons, since whenever we forced the downregulation of Rab10 or expressed Rab10DN for 48 hours or more, changes in dendritic and axonal arborisation and a decrease in cell viability were observed (not shown). Similar effects have been reported upon the overexpression of dominant-negative mutants of Rab5, the GTPase mediating formation of early endosomes, but not for Rab11, which controls the slow recycling of receptors to the plasma membrane^{33,34}, suggesting that the distinct arms of the endosomal network differentially impact on neuronal homeostasis.

In general, small GTPases of the Rab family have been defined by the specificity of the trafficking process they facilitate. Acting as a highly specialised network, they constitute one of the master regulators of membrane trafficking in eukaryotic cells³². Rab10 is one of the few exceptions to this rule and during the last 30 years, it

has been associated to multiple trafficking pathways, including polarised exocytosis from early endosomes, specific exocytic mechanisms in adipocytes and neurons, endoplasmic reticulum dynamics and the formation of tubular endosomes, to cite but a few³⁵. This multiplicity of functions is reflected by the diversity of its interactors and effectors, emphasising the importance of the specific context in which Rab10 operates. Here we have confirmed previous findings showing that Rab10-positive organelles are present in the axon of hippocampal neurons and display bi-directional transport¹⁷. Importantly, in this work, we have demonstrated that the balance between anterograde and retrograde transport is regulated by BDNF stimulation, indicating the ability of Rab10 to specifically respond to extracellular cues in a signalling context. A similar function of Rab10 balancing anterograde and retrograde transport has been shown to be required for correct dendritic patterning in *Drosophila*³⁶.

Although the mechanism by which Rab10-positive organelles switch between retrograde and anterograde transport has not been the focus of this work, two key observations suggest that directionality does not depend on Rab10 active status. First, both anterograde and retrograde organelles have membrane-bound Rab10 (Figure 4), which is generally accepted to be its GTP-bound active form. Accordingly, the Rab10-T23N dominant-negative mutant, which resembles GDP-bound Rab10, displays a diffuse cytoplasmic distribution. Second, we observed that the Rab10-Q68L constitutively-active mutant is transported predominantly in retrograde direction in neurons depleted of BDNF (data not shown). Altogether these observations suggest that the GTP-bound conformation of Rab10 is required to interact with the membrane of axonal organelles, but is not sufficient to determine the direction of transport.

Phosphorylation of Rab10 by the Leucine-rich repeat kinase 2 (LRRK2) has been shown to regulate interaction with JIP3 and JIP4³⁷, which are adaptors for the plus-end-directed microtubule-dependent molecular motor kinesin-1^{38,39}. Interestingly, JIP3 has been shown to mediate TrkB anterograde transport in neurons and, by that mechanism, to enhance BDNF signalling^{40,41}. Although the mechanism by which TrkB signalling promotes recruitment and activation of LRRK2 in Rab10-positive organelles is currently only a matter of speculation, one of the most interesting candidates to mediate this process is Vps35, a component of the retromer complex, which has been shown to modulate LRRK2⁴². Consistent with this view, TrkB binds SorLA, a key factor for retromer assembly⁴³. It is therefore possible that the first wave of TrkB-containing signalling endosomes induces the phosphorylation of Rab10 through the Vps35-LRRK2 recruitment, leading to the mobilisation (or retention) of Rab10-positive organelles at the cell periphery. This proposed mechanism could be particularly attractive in the context of certain models of neurodegeneration, given the genetic links between LRRK2 and Parkinson's disease (PD)^{44,45} and the neuroprotective role shown for BDNF in neurons of the substantia nigra in PD models^{46,47}.

In addition, activity and phosphorylation of Rab10 could be controlled by canonical TrkB signalling pathways, including PI3K-Akt. In adipocytes and muscle cells, Rab10 is known to regulate the plasma membrane delivery of the glucose transporter GLUT4 in response to insulin, where activation of Akt leads to phosphorylation of the Rab GAP Akt substrate of 160 kDa (AS160). Rab10 associated to GLUT4-containing endosomes is kept inactive by AS160, until Akt signalling releases the brake promoting fusion with the plasma membrane⁴⁸. Interestingly, it has been recently shown that LRRK2 is required for efficient insulin-stimulated translocation of GLUT4, and the effects of LRRK2 depletion can be compensated by increasing Akt signalling⁴⁹. Whether TrkB-PI3K-Akt is able to positively regulate phosphorylation of Rab10 is an interesting possibility we aim to explore in the near future.

Once internalised, axonal TrkB reaches early endosomes, from where it can either recycle back to the plasma membrane, thus fine-tuning the response of the nerve terminal to BDNF (reviewed in⁵⁰), or undergo sorting to the retrograde axonal transport route, propagating the neurotrophic signal to the soma^{9,12,13}. Given that

Rab10 downregulation decreased retrograde transport of TrkB and stimulation with BDNF promoted anterograde transport of Rab10, we reasoned that Rab10 delivery to the axon terminal facilitated sorting of TrkB to retrogradely transported compartments. We demonstrated that the expression of Rab10DN increases the amount of internalised TrkB accumulated in early endosomes without significantly affecting recycling (Figure 5), a result that indicates functional Rab10 aids the sorting of TrkB away from early endosomes and to a distinct axonal compartment.

Internalised Rab10 is found in retrograde signalling endosomes positive for TrkB or p75^{NTR} (Figure 3). However, when we studied triple co-localisation of internalised HcT and TrkB with endogenous Rab10, we found the absolute amount of retrograde TrkB in Rab10 organelles to remain constantly low, even when the total amount of retrogradely-transported TrkB increases three times from 30 to 90 min (Figure 2e), suggesting that this interaction is transient. This is in line with previous evidence showing that in MDCK cells, Rab10 mediates transport from early endosomes to a polarised trafficking route²³. Moreover, in *C. elegans*, Rab10 is recruited to early endosomes where it downregulates Rab5, helping to select cargoes for delivery to other endosomal compartments, in this case recycling endosomes⁵¹. Among other functions, Rab10 has been shown to participate in the biogenesis of tubular endosomes in mammalian cells²⁵, which together with evidence indicating that Rab10 regulates endosomal phosphatidylinositol-4,5-bisphosphate in *C. elegans*, suggests that Rab10 modulates the membrane recruitment of factors altering membrane curvature and vesicle budding⁵².

Our data suggests that Rab10 regulates the amount of internalised TrkB that is sorted to retrograde signalling endosomes, depending on the concentration of BDNF at the axon terminal. Since BDNF is known to be released from the post-synaptic membrane depending on neuronal activity⁵³, retrograde neurotrophic signalling from the axon terminal is a substantial feedback mechanism regulating growth and survival, and therefore, ensuring that the active circuit is preserved. To keep this feedback signal meaningful, any change in the availability of BDNF at the synapse has to translate into proportional changes in the intensity of the signal perceived at the soma. We propose that Rab10 organelles deliver crucial components of the sorting machinery on demand. At steady state, anterograde and retrograde transport of Rab10 organelles are in equilibrium (Figure 6; *low BDNF*). Further decrease in BDNF concentration makes retrograde transport of Rab10 compartments predominant, as observed in live-cell imaging experiments performed in neurons treated with anti-BDNF blocking antibodies (Figure 4, no BDNF). In contrast, adding BDNF reverts the directional bias of Rab10 organelles to anterograde (Figure 6; *high BDNF*), which allows the delivery of appropriate levels of the sorting machinery to the nerve terminal, thus increasing the efficiency of TrkB retrograde transport.

Little is known about the specific sorting machinery required for the biogenesis of signalling endosomes. Endophilins A1 and A3 have been shown to regulate the trafficking of TrkB across early endosomes and mediate survival signalling^{14,54}. Interestingly, LRRK2 substrates include endophilin A1 as well as Rab10, opening the possibility of both being found on the same organelle⁵⁵. Further identification of components of the sorting machinery delivered to the axonal terminal by Rab10-positive compartments will be crucial to understand not only how this mechanism allows coordination between local signalling and global neuronal responses, but also how this process may fail in neurodegeneration. In this light, promoting the delivery of Rab10 organelles to nerve terminals may be explored as an innovative therapeutic strategy for diseases in which the endo-lysosomal system is overloaded or dysfunctional, such as Alzheimer's disease^{56,57}, or to increase the ability of axons to respond to trophic factors during regeneration.

Methods

A summary of all the antibodies, plasmids and lentiviral vectors used in this work can be found in the [Supplementary Table 1](#).

Neuronal cultures

Embryonic hippocampal neurons from C57BL/6 mice of either sex and embryonic age of 16-17 days were dissected adapting previously described protocols⁵⁸. Dissection was performed in cold Hanks' balanced salt solution (HBSS) and the tissue was collected in cold Hibernate E medium (ThermoFisher, #A1247601). After incubating for 10 min in 300 μ L of Accumax (Innovative Cell Technologies, #AM105) diluted in HBSS (1:1), tissue was washed in HBSS, resuspended in warm plating medium (Minimum Essential Medium supplemented with 10% horse serum, 6% glucose and 2 mM glutamine) and mechanically dissociated by pipetting. 10,000-12,000 neurons per cm^2 were then seeded on glass coverslips or microfluidic chambers, pre-coated with 1 mg/mL poly-L-lysine. Before coating, glass coverslips were treated overnight with NoChromix (Godax Laboratories), washed three times and sterilised in 70% ethanol. Microfluidic chambers were produced in-house as previously described^{59,60}. Polydimethylsiloxane inserts were fabricated from resin moulds, which are replicas of the master template produced by soft lithography, and then irreversibly bound to glass-bottom dishes (WillCo Wells, #HBSB-3512) by plasma treatment. Neurons were left in plating medium for 1.5 h and then shifted to maintenance medium (Neurobasal supplemented with B27, 2 mM glutamine, 6% glucose and antibiotics). Half of the culture medium was replaced by fresh medium every 3-4 days.

Immunofluorescence.

Neurons were washed in phosphate buffer saline (PBS) and fixed for 15 min in 3% PFA and 4% sucrose dissolved in PBS. Next, cells were incubated in 0.15 M glycine dissolved in PBS for 10 minutes and then blocked and permeabilised simultaneously by incubation for 1 h in 5% BSA and 0.1% saponin in PBS. Samples were incubated at 4°C overnight with primary antibodies diluted in 5% BSA, 0.05% saponin, 0.1 mM CaCl_2 and 0.1 mM MgCl_2 dissolved in PBS at the following concentrations: 1:50 goat anti-Rab10 (Santa Cruz Biotechnologies, cat no. sc-6564, RRID:AB_2237844); 1:200 mouse anti-Rab10 (Abcam, cat no. ab104859, RRID:AB_10711207); 1:200 rabbit anti-Rab10 (Cell Signalling Technology, cat no. 8127, RRID:AB_10828219); 1:300 chicken anti- β III tubulin (Synaptic Systems, cat no. 302 306, AB_2620048); 1:250 rabbit anti-phosphorylated CREB S133 (Abcam, cat no. ab32096, RRID:AB_731734); 1:200 mouse anti-Rab7 (Abcam, cat no. ab50533, RRID:AB_882241); and 1:200 rabbit anti-Rab5 (Abcam, cat no. ab13253, RRID:AB_299796). Then, neurons were washed three times with PBS and incubated for 90 min with AlexaFluor-conjugated secondary antibodies 1:400 (ThermoFisher) in 5% BSA, 0.05% saponin, 0.1 mM CaCl_2 and 0.1 mM MgCl_2 dissolved in PBS. Nuclear staining 4',6-diamidino-2-phenylindole (DAPI) was added with the secondary antibodies when appropriate. Finally, coverslips were washed in PBS and mounted with Mowiol.

Transfection and plasmids

Hippocampal neurons were transfected at 7 days in vitro (DIV) using Lipofectamine 2000 (ThermoFisher). Experiments were carried out after 20-24 h. The pEGFP-C1 plasmid is from Clontech (Addgene plasmid # 13031, RRID:Addgene_13031), the plasmids for Rab10 WT EGFP (RRID:Addgene_49472) and Rab10 T23N EGFP (RRID:Addgene_49545) were a gift from Marci Scidmore⁶¹, TrkB-FLAG plasmid was a gift from Francis Lee²⁷.

Retrograde accumulation and signalling assays

After 7 DIV, microfluidic chambers with overt axon crossing were selected. Rab10 was knocked down by transducing neurons with an inducible shRNA Rab10 lentivirus (pTightPuro-shRNA Rab10) and its doxycycline-dependent regulator TET-ON Advance. The day before the experiment, doxycycline 1 μ g/mL was added to the cell bodies and 18-22 h later, the media was replaced with Neurobasal in somatic and axonal compartments to deplete cells from endogenous growth factors for 1 h. For analysing the retrograde accumulation of TrkB, we added polyclonal antibodies against the extracellular domain of TrkB (1:50 rabbit anti-TrkB, Millipore, cat no. AB9872, RRID:AB_2236301) together with 20 ng/mL BDNF for 2.5 h. After PFA fixation, the transport of internalized antibodies was revealed by incubating the somatic compartment with fluorescently-labelled

secondary antibodies. The same protocol was used to study retrograde propagation of neurotrophic signalling; after 1 h of growth factor-depletion, axons were stimulated with 20 ng/mL BDNF for 2.5 h and after fixation, phosphorylation of CREB in the nucleus was analysed by immunofluorescence.

Co-localisation studies

To analyse the presence of two markers in the same organelle, we used confocal z-stack images (voxel size: 0.197 x 0.197 x 0.5 μm) and the confined displacement algorithm to measure Manders' correlation index within axons and determine its statistical significance compared to random images of identical total intensity and shape²⁰. To compute random scenarios, seven random radial displacements were taken at a maximum radial distance of 12 pixels (a total of 353 samples), and histograms binning = 16. CDA was implemented by using the plugin from the GDSC University of Sussex (<http://www.sussex.ac.uk/gdsc/intranet/microscopy/UserSupport/AnalysisProtocol/imagej/colocalisation>). All the data points were plotted and the mean and standard error is indicated for each group and compared using Student t-test. Statistical significance of the individual data point is colour coded (see figure legends).

Super-resolution radial fluctuations

High-fidelity super-resolution information was extracted from time series of 1,000 confocal images per channel by using super-resolution radial fluctuations (SRRF) algorithm⁶². Super-resolution images were then quality-controlled by using Super-Resolution Quantitative Image Rating and Reporting of Error Locations (SQUIRREL) algorithm⁶³. Implementation of the algorithms was done in FIJI by using the open-source plugin NanoJ-core (<https://henriqueslab.github.io/resources/NanoJ>).

Immunoendocytosis

Hippocampal neurons transfected with TrkB-Flag were kept in Neurobasal media for 1 h and then incubated on ice with 1:50 mouse anti-Flag antibody (Sigma-Aldrich Cat no. F3040, RRID:AB_439712). In selected samples, 20 nM HcT was also added. Internalisation of receptors was then induced by incubation with 50 ng/mL BDNF for 30 min at 37°C. Antibodies bound to receptors still at the cell surface were dissociated by washing twice for 1 min in PBS supplemented with 1 mM EDTA. In experiments to measure recycling of internalised receptors, neurons were further stimulated with BDNF for other 60 min in the presence of an AlexaFluor647-conjugated anti-mouse secondary antibody, fixed, permeabilised and incubated with an AlexaFluor555-conjugated anti-mouse secondary antibody to detect total internalised receptor.

Axonal transport

Axonal transport of overexpressed fluorescent proteins and internalised fluorescent antibodies was analysed from confocal time series of 1 frame/s and a pixel size of $\sim 0.1 \times 0.1 \mu\text{m}^2$, captured during 5 min intervals at different time points by using a Zeiss LSM 780 NLO multiphoton confocal microscope with an oil immersion 63 \times objective and equipped with an environmental chamber (Zeiss XL multi S1 DARK LS set at 37°C and environmental CO₂). For these experiments, neurons were cultured on 25 mm coverslips kept in Neurobasal for 1 h prior to cell imaging and mounted inside Attofluor chambers (Thermo-Fisher Scientific, Cat no. A7816) with BrightCell NEUMO photostable media (Sigma-Aldrich, Cat no. SCM145). Speed, pausing and direction of labelled organelles were analysed from kymographs by using Kymoanalyzer set of macros from Encalada lab (<https://www.encalada.scripps.edu/kymoanalyzer>)⁶⁴.

Statistical Analysis

Samples collected in independent experiments were tested for normality and homoscedasticity to apply the appropriate corrections to the statistical tests. Each specific test, its degrees of freedom and level of significance are indicated in the respective figure legend. All the details are summarised in the statistical annex (Supplementary Table 2). Plots show mean \pm standard error and exact p values are indicated when relevant.

Software

Images were handled, edited and analysed using ImageJ/FIJI (version 2.1.0, 1.53c). Figures were checked with Coblis (<https://www.color-blindness.com/coblis-color-blindness-simulator/>), using The Colour Blind Simulator algorithms from Matthew Wickline and the Human-Computer Interaction Resource Network. Palettes were adjusted to maximise visibility. The Orange/Green/Purple balanced look-up table was obtained from Christophe Leterrier's GitHub repository (<https://github.com/cleterrier/ChrisLUTs>). Database files were imported, analysed and sorted as R files using RStudio (version 1.0.44). GraphPad Prism for Mac (version 6.00, GraphPad Software) was used for running statistical analysis and generate the plots included in the figures. Illustrations were created with BioRender (<http://www.biorender.com>). Updated versions of ImageJ macros and R scripts used in this article, as well as the specific implementation of Kymoanalyzer used to analyse our datasets, can be found on our GitHub repository (<https://github.com/omlazo>).

Acknowledgements

We thank Dr James N Sleight (UCL Queen Square Institute of Neurology) for critical reading of the manuscript. This work was supported by the MRC Project grant MR/T001976/1 [OL], the Wellcome Trust Senior Investigator Award (107116/Z/15/Z) [GS] and the UK Dementia Research Institute Foundation award UKDRI-1005 [GS].

Author's contributions

OL performed the research and drafted the manuscript, which was revised by GS. OL and GS interpreted the data and approved submission of this work.

Conflict of Interest

The authors have no competing interest to declare.

References

- 1 Andres-Alonso, M. *et al.* SIPA1L2 controls trafficking and local signaling of TrkB-containing amphisomes at presynaptic terminals. *Nat Commun* **10**, 5448, doi:10.1038/s41467-019-13224-z (2019).
- 2 Szobota, S. *et al.* BDNF, NT-3 and Trk receptor agonist monoclonal antibodies promote neuron survival, neurite extension, and synapse restoration in rat cochlea ex vivo models relevant for hidden hearing loss. *PLoS One* **14**, e0224022, doi:10.1371/journal.pone.0224022 (2019).
- 3 Woo, D. *et al.* Locally Activating TrkB Receptor Generates Actin Waves and Specifies Axonal Fate. *Cell Chem Biol* **26**, 1652-1663 e1654, doi:10.1016/j.chembiol.2019.10.006 (2019).
- 4 Barford, K., Deppmann, C. & Winckler, B. The neurotrophin receptor signaling endosome: Where trafficking meets signaling. *Dev Neurobiol* **77**, 405-418, doi:10.1002/dneu.22427 (2017).
- 5 Villarroel-Campos, D., Schiavo, G. & Lazo, O. M. The many disguises of the signalling endosome. *FEBS Lett* **592**, 3615-3632, doi:10.1002/1873-3468.13235 (2018).
- 6 Du, J. L. & Poo, M. M. Rapid BDNF-induced retrograde synaptic modification in a developing retinotectal system. *Nature* **429**, 878-883, doi:10.1038/nature02618 (2004).
- 7 Pazyra-Murphy, M. F. *et al.* A retrograde neuronal survival response: target-derived neurotrophins regulate MEF2D and bcl-w. *J Neurosci* **29**, 6700-6709, doi:10.1523/JNEUROSCI.0233-09.2009 (2009).
- 8 Watson, F. L. *et al.* Rapid nuclear responses to target-derived neurotrophins require retrograde transport of ligand-receptor complex. *J Neurosci* **19**, 7889-7900 (1999).
- 9 Zhou, B., Cai, Q., Xie, Y. & Sheng, Z. H. Snapin recruits dynein to BDNF-TrkB signaling endosomes for retrograde axonal transport and is essential for dendrite growth of cortical neurons. *Cell Rep* **2**, 42-51, doi:10.1016/j.celrep.2012.06.010 (2012).
- 10 Huang, E. J. & Reichardt, L. F. Trk receptors: roles in neuronal signal transduction. *Annu Rev Biochem* **72**, 609-642, doi:10.1146/annurev.biochem.72.121801.161629 (2003).
- 11 Minichiello, L. TrkB signalling pathways in LTP and learning. *Nat Rev Neurosci* **10**, 850-860, doi:10.1038/nrn2738 (2009).
- 12 Deinhardt, K. *et al.* Rab5 and Rab7 control endocytic sorting along the axonal retrograde transport pathway. *Neuron* **52**, 293-305, doi:10.1016/j.neuron.2006.08.018 (2006).
- 13 Ha, J. *et al.* A neuron-specific cytoplasmic dynein isoform preferentially transports TrkB signaling endosomes. *J Cell Biol* **181**, 1027-1039, doi:10.1083/jcb.200803150 (2008).
- 14 Burk, K. *et al.* EndophilinAs regulate endosomal sorting of BDNF-TrkB to mediate survival signaling in hippocampal neurons. *Sci Rep* **7**, 2149, doi:10.1038/s41598-017-02202-4 (2017).
- 15 Kucharava, K. *et al.* Sodium-hydrogen exchanger 6 (NHE6) deficiency leads to hearing loss, via reduced endosomal signalling through the BDNF/Trk pathway. *Sci Rep* **10**, 3609, doi:10.1038/s41598-020-60262-5 (2020).
- 16 Bucci, C., Alifano, P. & Cogli, L. The role of Rab proteins in neuronal cells and in the trafficking of neurotrophin receptors. *Membranes (Basel)* **4**, 642-677, doi:10.3390/membranes4040642 (2014).
- 17 Deng, C. Y. *et al.* JIP1 mediates anterograde transport of Rab10 cargos during neuronal polarization. *J Neurosci* **34**, 1710-1723, doi:10.1523/JNEUROSCI.4496-13.2014 (2014).
- 18 Debaisieux, S., Encheva, V., Chakravarty, P., Snijders, A. P. & Schiavo, G. Analysis of Signaling Endosome Composition and Dynamics Using SILAC in Embryonic Stem Cell-Derived Neurons. *Mol Cell Proteomics* **15**, 542-557, doi:10.1074/mcp.M115.051649 (2016).
- 19 Gonzalez-Gutierrez, A., Lazo, O. M. & Bronfman, F. C. The Rab5-Rab11 Endosomal Pathway is Required for BDNF-Induced CREB Transcriptional Regulation in Hippocampal Neurons. *J Neurosci* **40**, 8042-8054, doi:10.1523/JNEUROSCI.2063-19.2020 (2020).

- 20 Ramirez, O., Garcia, A., Rojas, R., Couve, A. & Hartel, S. Confined displacement algorithm determines true and random colocalization in fluorescence microscopy. *J Microsc* **239**, 173-183, doi:10.1111/j.1365-2818.2010.03369.x (2010).
- 21 Goto-Silva, L. *et al.* Retrograde transport of Akt by a neuronal Rab5-APPL1 endosome. *Sci Rep* **9**, 2433, doi:10.1038/s41598-019-38637-0 (2019).
- 22 Zhen, Y. & Stenmark, H. Cellular functions of Rab GTPases at a glance. *J Cell Sci* **128**, 3171-3176, doi:10.1242/jcs.166074 (2015).
- 23 Babbey, C. M. *et al.* Rab10 regulates membrane transport through early endosomes of polarized Madin-Darby canine kidney cells. *Mol Biol Cell* **17**, 3156-3175, doi:10.1091/mbc.e05-08-0799 (2006).
- 24 Brewer, P. D., Habtemichael, E. N., Romenskaia, I., Mastick, C. C. & Coster, A. C. Glut4 Is Sorted from a Rab10 GTPase-independent Constitutive Recycling Pathway into a Highly Insulin-responsive Rab10 GTPase-dependent Sequestration Pathway after Adipocyte Differentiation. *J Biol Chem* **291**, 773-789, doi:10.1074/jbc.M115.694919 (2016).
- 25 Etoh, K. & Fukuda, M. Rab10 regulates tubular endosome formation through KIF13A and KIF13B motors. *J Cell Sci* **132**, doi:10.1242/jcs.226977 (2019).
- 26 Xu, X. H. *et al.* MARCKS regulates membrane targeting of Rab10 vesicles to promote axon development. *Cell Res* **24**, 576-594, doi:10.1038/cr.2014.33 (2014).
- 27 Chen, Z. Y., Ieraci, A., Tanowitz, M. & Lee, F. S. A novel endocytic recycling signal distinguishes biological responses of Trk neurotrophin receptors. *Mol Biol Cell* **16**, 5761-5772, doi:10.1091/mbc.e05-07-0651 (2005).
- 28 Wang, H., Xu, J., Lazarovici, P., Quirion, R. & Zheng, W. cAMP Response Element-Binding Protein (CREB): A Possible Signaling Molecule Link in the Pathophysiology of Schizophrenia. *Front Mol Neurosci* **11**, 255, doi:10.3389/fnmol.2018.00255 (2018).
- 29 Esvald, E. E. *et al.* CREB Family Transcription Factors Are Major Mediators of BDNF Transcriptional Autoregulation in Cortical Neurons. *J Neurosci* **40**, 1405-1426, doi:10.1523/JNEUROSCI.0367-19.2019 (2020).
- 30 Kwon, M., Fernandez, J. R., Zegarek, G. F., Lo, S. B. & Firestein, B. L. BDNF-promoted increases in proximal dendrites occur via CREB-dependent transcriptional regulation of cypin. *J Neurosci* **31**, 9735-9745, doi:10.1523/JNEUROSCI.6785-10.2011 (2011).
- 31 Cohen, M. S., Bas Orth, C., Kim, H. J., Jeon, N. L. & Jaffrey, S. R. Neurotrophin-mediated dendrite-to-nucleus signaling revealed by microfluidic compartmentalization of dendrites. *Proc Natl Acad Sci U S A* **108**, 11246-11251, doi:10.1073/pnas.1012401108 (2011).
- 32 Riccio, A., Pierchala, B. A., Ciarallo, C. L. & Ginty, D. D. An NGF-TrkA-mediated retrograde signal to transcription factor CREB in sympathetic neurons. *Science* **277**, 1097-1100, doi:10.1126/science.277.5329.1097 (1997).
- 33 Lazo, O. M. *et al.* BDNF regulates Rab11-mediated recycling endosome dynamics to induce dendritic branching. *J Neurosci* **33**, 6112-6122, doi:10.1523/JNEUROSCI.4630-12.2013 (2013).
- 34 Moya-Alvarado, G., Gonzalez, A., Stuardo, N. & Bronfman, F. C. Brain-Derived Neurotrophic Factor (BDNF) Regulates Rab5-Positive Early Endosomes in Hippocampal Neurons to Induce Dendritic Branching. *Front Cell Neurosci* **12**, 493, doi:10.3389/fncel.2018.00493 (2018).
- 35 Chua, C. E. L. & Tang, B. L. Rab 10-a traffic controller in multiple cellular pathways and locations. *J Cell Physiol* **233**, 6483-6494, doi:10.1002/jcp.26503 (2018).
- 36 Taylor, C. A., Yan, J., Howell, A. S., Dong, X. & Shen, K. RAB-10 Regulates Dendritic Branching by Balancing Dendritic Transport. *PLoS Genet* **11**, e1005695, doi:10.1371/journal.pgen.1005695 (2015).
- 37 Waschbusch, D. *et al.* Structural Basis for Rab8a Recruitment of RILPL2 via LRRK2 Phosphorylation of Switch 2. *Structure* **28**, 406-417 e406, doi:10.1016/j.str.2020.01.005 (2020).
- 38 Cockburn, J. J. B. *et al.* Insights into Kinesin-1 Activation from the Crystal Structure of KLC2 Bound to JIP3. *Structure* **26**, 1486-1498 e1486, doi:10.1016/j.str.2018.07.011 (2018).

- 39 Isabet, T. *et al.* The structural basis of Arf effector specificity: the crystal structure of ARF6 in a complex with JIP4. *EMBO J* **28**, 2835-2845, doi:10.1038/emboj.2009.209 (2009).
- 40 Huang, S. H. *et al.* JIP3 mediates TrkB axonal anterograde transport and enhances BDNF signaling by directly bridging TrkB with kinesin-1. *J Neurosci* **31**, 10602-10614, doi:10.1523/JNEUROSCI.0436-11.2011 (2011).
- 41 Sun, T. *et al.* JIP1 and JIP3 cooperate to mediate TrkB anterograde axonal transport by activating kinesin-1. *Cell Mol Life Sci* **74**, 4027-4044, doi:10.1007/s00018-017-2568-z (2017).
- 42 Mir, R. *et al.* The Parkinson's disease VPS35[D620N] mutation enhances LRRK2-mediated Rab protein phosphorylation in mouse and human. *Biochem J* **475**, 1861-1883, doi:10.1042/BCJ20180248 (2018).
- 43 Rohe, M., Hartl, D., Fjorback, A. N., Klose, J. & Willnow, T. E. SORLA-mediated trafficking of TrkB enhances the response of neurons to BDNF. *PLoS One* **8**, e72164, doi:10.1371/journal.pone.0072164 (2013).
- 44 Kluss, J. H., Mamais, A. & Cookson, M. R. LRRK2 links genetic and sporadic Parkinson's disease. *Biochem Soc Trans* **47**, 651-661, doi:10.1042/BST20180462 (2019).
- 45 Tolosa, E., Vila, M., Klein, C. & Rascol, O. LRRK2 in Parkinson disease: challenges of clinical trials. *Nat Rev Neurol* **16**, 97-107, doi:10.1038/s41582-019-0301-2 (2020).
- 46 Qin, Q. *et al.* Effects of LRRK2 Inhibitors on Nigrostriatal Dopaminergic Neurotransmission. *CNS Neurosci Ther* **23**, 162-173, doi:10.1111/cns.12660 (2017).
- 47 Rocha, E. M. *et al.* LRRK2 inhibition prevents endolysosomal deficits seen in human Parkinson's disease. *Neurobiol Dis* **134**, 104626, doi:10.1016/j.nbd.2019.104626 (2020).
- 48 Sano, H. *et al.* Rab10, a target of the AS160 Rab GAP, is required for insulin-stimulated translocation of GLUT4 to the adipocyte plasma membrane. *Cell Metab* **5**, 293-303, doi:10.1016/j.cmet.2007.03.001 (2007).
- 49 Funk, N. *et al.* The Parkinson's disease-linked Leucine-rich repeat kinase 2 (LRRK2) is required for insulin-stimulated translocation of GLUT4. *Sci Rep* **9**, 4515, doi:10.1038/s41598-019-40808-y (2019).
- 50 Andreska, T., Luningschror, P. & Sendtner, M. Regulation of TrkB cell surface expression-a mechanism for modulation of neuronal responsiveness to brain-derived neurotrophic factor. *Cell Tissue Res* **382**, 5-14, doi:10.1007/s00441-020-03224-7 (2020).
- 51 Liu, O. & Grant, B. D. Basolateral Endocytic Recycling Requires RAB-10 and AMPH-1 Mediated Recruitment of RAB-5 GAP TBC-2 to Endosomes. *PLoS Genet* **11**, e1005514, doi:10.1371/journal.pgen.1005514 (2015).
- 52 Shi, A. & Grant, B. D. Interactions between Rab and Arf GTPases regulate endosomal phosphatidylinositol-4,5-bisphosphate during endocytic recycling. *Small GTPases* **4**, 106-109, doi:10.4161/sgtp.23477 (2013).
- 53 Matsuda, N. *et al.* Differential activity-dependent secretion of brain-derived neurotrophic factor from axon and dendrite. *J Neurosci* **29**, 14185-14198, doi:10.1523/JNEUROSCI.1863-09.2009 (2009).
- 54 Fu, X. *et al.* Retrolinkin cooperates with endophilin A1 to mediate BDNF-TrkB early endocytic trafficking and signaling from early endosomes. *Mol Biol Cell* **22**, 3684-3698, doi:10.1091/mbc.E11-04-0308 (2011).
- 55 Matta, S. *et al.* LRRK2 controls an EndoA phosphorylation cycle in synaptic endocytosis. *Neuron* **75**, 1008-1021, doi:10.1016/j.neuron.2012.08.022 (2012).
- 56 Van Acker, Z. P., Bretou, M. & Annaert, W. Endo-lysosomal dysregulations and late-onset Alzheimer's disease: impact of genetic risk factors. *Mol Neurodegener* **14**, 20, doi:10.1186/s13024-019-0323-7 (2019).
- 57 Xu, W., Fang, F., Ding, J. & Wu, C. Dysregulation of Rab5-mediated endocytic pathways in Alzheimer's disease. *Traffic* **19**, 253-262, doi:10.1111/tra.12547 (2018).
- 58 Kaech, S. & Banker, G. Culturing hippocampal neurons. *Nat Protoc* **1**, 2406-2415, doi:10.1038/nprot.2006.356 (2006).

- 59 Sannerud, R. *et al.* ADP ribosylation factor 6 (ARF6) controls amyloid precursor protein (APP) processing by mediating the endosomal sorting of BACE1. *Proc Natl Acad Sci U S A* **108**, E559-568, doi:10.1073/pnas.1100745108 (2011).
- 60 Restani, L. *et al.* Botulinum neurotoxins A and E undergo retrograde axonal transport in primary motor neurons. *PLoS Pathog* **8**, e1003087, doi:10.1371/journal.ppat.1003087 (2012).
- 61 Huang, B. *et al.* The Anaplasma phagocytophilum-occupied vacuole selectively recruits Rab-GTPases that are predominantly associated with recycling endosomes. *Cell Microbiol* **12**, 1292-1307, doi:10.1111/j.1462-5822.2010.01468.x (2010).
- 62 Culley, S., Tosheva, K. L., Matos Pereira, P. & Henriques, R. SRRF: Universal live-cell super-resolution microscopy. *Int J Biochem Cell Biol* **101**, 74-79, doi:10.1016/j.biocel.2018.05.014 (2018).
- 63 Culley, S. *et al.* Quantitative mapping and minimization of super-resolution optical imaging artifacts. *Nat Methods* **15**, 263-266, doi:10.1038/nmeth.4605 (2018).
- 64 Neumann, S., Chassefeyre, R., Campbell, G. E. & Encalada, S. E. KymoAnalyzer: a software tool for the quantitative analysis of intracellular transport in neurons. *Traffic* **18**, 71-88, doi:10.1111/tra.12456 (2017).

Figures

Figure 1. Rab10 is required for retrograde TrkB trafficking and signalling.

(a) Schematic of two-compartment microfluidic chambers highlighting compartmentalisation of somata (left) and axon terminals (right). Micro-flow from somatic to axonal compartments provides fluidic isolation of the axonal compartment. **(b)** Representative images from the cell bodies of neurons incubated for 2.5 hours with anti-TrkB in the axonal compartment, with (+) or without (-) BDNF. Pink arrowheads indicate examples of retrogradely transported TrkB-positive organelles. Scale bar: 10 μ m. **(c)** Neurons treated with an inducible shRNA targeting Rab10 were compared to control neurons. Immunofluorescence revealed similar neuronal density (see β III-tubulin in orange and nuclear staining in green, top panel), but a decrease in both, expression of Rab10 (gray, middle panel) and retrograde accumulation of TrkB after 2.5 hours (colour intensity scale, bottom panel). Scale bar: 50 μ m. **(d)** Quantification of retrograde TrkB accumulation in three independent experiments show statistically significant differences (unpaired t-Student test, $t(140)$, $p < 0.0001$). **(e)** Analysis of the correlation between expression level of Rab10 and retrograde TrkB accumulation in the whole population of control and Rab10-knockdown neurons show a significant linear correlation (Goodness of fit $R^2 = 0.61$; Pearson r , XY pairs=131, $p < 0.0001$). **(f)** Axonal stimulation with BDNF for 2.5 hours leads to robust appearance of phosphorylated CREB in the nucleus of control neurons (left panel). This was restricted in neurons depleted of Rab10 (middle panel), which was rescued by the co-expression of a shRNA-resistant mutant Rab10 (right panel). Immunofluorescence for Rab10 is shown in gray, with the nuclei indicated with a pink mask, and nuclear phosphorylated CREB is shown in a colour intensity scale. Scale bar: 50 μ m. **(g)** Quantification from three independent experiments showing the statistically significant effect of manipulating Rab10 expression on the levels of phosphorylated CREB in the nucleus (one-way ANOVA, $F(2,280)$, $p < 0.0001$; p values for the Bonferroni multiple comparison tests, $t(280)$, are indicated in the plot).






Figure 2. Internalised TrkB transiently co-localises with Rab10 in axons.




(a) The distribution of endogenous Rab10 (orange) and Rab7 (green) was monitored using confocal microscopy in axons of neurons with or without BDNF treatment for 30 min. Points where both markers overlap are indicated with yellow arrowheads. Scale bar = 5 μ m **(b)** Co-localisation was analysed using confined-displacement algorithm (CDA), and Manders coefficients were compared for control versus BDNF-treated axons (Welch's corrected Unpaired t-Student; M1: $t(72.32)$, p value = 0.0621; M2: $t(62.33)$, p value = 0.1043). Data points showing significant co-localisation compared to random scenarios are indicated in coloured dots. No significant differences were found between starved and BDNF-stimulated neurons. **(c)** Representative super-resolution radial fluctuations (SRRF) microscopy of an axon stained for endogenous Rab5 (purple), Rab7 (green) and Rab10 (orange). The inset at 6x higher magnification shows examples of partial co-localisation of Rab10 with Rab5 and Rab7. Scale bar = 5 μ m. **(d)** Intensity along the same axonal segment shown in (c) was plotted to observe the correlation between the three markers. The region of the inset is indicated. Several examples of partial coincidence between Rab10 and Rab5 are observed. Overlap between Rab10 and Rab7 peaks is less frequent, confirming the data in (a) and (b). **(e)** Labelled HcT and anti-TrkB were co-internalised in the presence of BDNF for 30, 60 and 90 min, and then their level of overlap with endogenous Rab10 in axons was evaluated using confocal microscopy. Relative area positive for HcT and TrkB is shown in cyan (normalised to 30 min) and the fraction of the normalised area that was triple positive for HcT, TrkB and Rab10 is plotted in white. While the double TrkB/HcT-positive area significantly increased by 90 min (one-way ANOVA, $F(2,72)$, p value < 0.0001; Bonferroni multiple comparison test p value is shown in the plot), the triple TrkB/HcT/Rab10 surface remained low and fairly constant at all the time points (one-way ANOVA, $F(2,72)$, p value = 0.2730; Bonferroni multiple comparisons test for 60 versus 90 min, $t(72)$, p value > 0.9999). **(f)** Representative SRRF microscopy from the same three time points. Double positive puncta for TrkB (green) and HcT (purple) is indicated with cyan arrowheads. Triple positive puncta of TrkB/HcT (cyan) and Rab10 (orange) are indicated with white empty triangles. Scale bar = 5 μ m.

Figure 3. Over-expressed Rab10 is co-transported with TrkB and p75 receptors in the axon.




(a) Hippocampal neurons were co-transfected with Rab10-EGFP and TrkB-Flag plasmids. Fluorescently-labelled anti-Flag antibodies were internalised in the presence of BDNF and their axonal dynamics monitored by live-cell microscopy. **(b)** Representative images of the axon of a double transfected neuron, where a double-positive organelle for Rab10-EGFP (orange) and anti-Flag (green) is indicated with yellow arrowheads. Scale bar = 5 μ m. **(c)** Representative kymograph of the axon of a double transfected neuron during 5 minutes of imaging showing Rab10 and TrkB-Flag channels, and traced tracks with double-positive tracks in yellow showing transport predominantly in the retrograde direction (right to left). Scale bar = 10 μ m. **(d)** Quantification of five experiments showing the proportion of TrkB mobile organelles that were positive for Rab10. No anterograde TrkB organelles were found; therefore, presence of Rab10 could not be determined (N.D.). **(e)** Same experiment was performed by transfecting Rab10 and visualising it together with fluorescently-labelled anti-p75^{NTR} antibodies. **(f)** Representative sequence of images from a live axon shows a double-positive organelle for Rab10 (orange) and p75^{NTR} (green) moving anterogradely. Scale bar = 5 μ m. **(g)** Representative kymograph showing Rab10 and p75^{NTR} channels, and traced tracks in yellow showing transport in both anterograde (left to right) and retrograde directions. Scale bar = 10 μ m. **(h)** Quantification of five experiments showing the proportion of p75^{NTR} mobile organelles that were positive for Rab10 in the anterograde and retrograde direction.

Figure 4. BDNF regulates the directionality of Rab10 organelles.



Hippocampal neurons were transfected with Rab10-EGFP and depleted of BDNF for 60 mins. **(a)** Representative axon of a live neuron showing retrograde (right to left) transport of a Rab10-positive organelle (white arrowheads). Scale bar = 5 μ m. **(b)** In the panels on the left, representative kymographs (colour-coded as in a) are presented from the same axon after BDNF depletion (top), immediately after the addition of 50 ng/mL of BDNF (middle) and 10 minutes after (bottom). In the panels on the right side, the tracks have been traced and categorised as retrograde (cyan), anterograde (pink) or stationary/bidirectional (yellow). Scale bar = 10 μ m. **(c)** The frequencies of tracks from each of the three categories have been quantified and plotted comparing no BDNF and 10 min post-addition of 50 ng/mL BDNF. N= 14 axonal segments from 10 independent experiments. Unpaired t-Student, t(14), showed a significant increase in anterograde carriers (p value = 0.0150, *) at the expense of retrograde carriers (p value = 0.003, **). Stationary and bidirectional carriers did not show any significant change (p value = 0.4278).


Figure 5. Rab10 regulates sorting of TrkB out of early endosomes.



(a) Main hypotheses about the role of Rab10 regulating the sorting of TrkB to signalling endosomes include recycling back to the plasma membrane or sorting of TrkB receptors out of early endosomes to retrograde carriers. **(b)** Top, diagram of the experiment showing TrkB-flag receptors on the axonal surface bound to anti-Flag antibodies (green). After internalisation, the remaining anti-Flag is removed from the surface and the labelled receptor that recycled to the plasma membrane is chased with a secondary antibody (orange). Bottom, representative examples of internalised TrkB (green) and recycled TrkB (orange) in axons from neurons transfected with EGFP or a Rab10 DN mutant. **(c)** Normalised recycling shows no difference between EGFP and Rab10 DN transfected neurons. Unpaired t-Student, $t(60.40)$, p value = 0.3914. **(d)** Top, diagram of the internalisation of TrkB-flag labelled with anti-Flag antibodies (orange) to Rab5-positive early endosomes (green). Bottom, representative thresholded microscopy images from the axon of neurons transfected with EGFP or Rab10 DN mutant. While the amount of orange puncta are similar in both conditions, yellow areas showing co-localisation of internalised TrkB and Rab5 are increased upon Rab10 DN expression. **(e)** Quantification of co-localisation between internalised TrkB-Flag and endogenous Rab5 is significantly higher in neurons expressing Rab10 DN compared to EGFP. Unpaired t-Student, $t(38.22)$, p value = 0.008. Significant co-localisation according to CDA (p value < 0.05 compared to randomised signal) is shown with black circles, while inconclusive co-localisation (p value > 0.05) is shown in grey. Scale bars = 5 μ m.

Figure 6. Model: Role of Rab10 in the sorting of TrkB to retrograde axonal transport.

At steady state, low concentrations of BDNF (grey side of the terminal) induce basal levels of TrkB internalisation. Rab10 supply (blue arrows) is in equilibrium and it mediates baseline level of TrkB retrograde transport (pink arrows). Upon increase of BDNF concentration (green side of the axon terminal), TrkB endocytosis as well as the proportion of Rab10 organelles moving towards the axon terminal are increased. Increased amounts of Rab10 result in further facilitation of the sorting of TrkB out of the early endosome and an augmented flux of retrograde signalling carriers.



Supplementary tables

Table S1 — Resources

Primary antibodies					
antigen	host species	manufacturer	cat number	RRID	concentration
Rab10	goat	Santa Cruz Biotechnologies	sc-6564	AB_2237844	IF: 1:50
Rab10	mouse	Abcam	ab104859	AB_10711207	IF: 1:200
Rab10	rabbit	Cell Signalling	8127	AB_10828219	IF: 1:200
TrkB	rabbit	Merck (Millipore)	AB9872	AB_11214317	IE: 1:50
Tubulin β III	chicken	Synaptic Systems	302 306	AB_2620048	IF: 1:300
phospho CREB	rabbit	Abcam	ab32096	AB_731734	IF: 1:250
Rab7	mouse	Abcam	ab50533	AB_882241	IF: 1:200
Rab5	rabbit	Abcam	ab13253	AB_299796	IF: 1:200
Flag (M1)	mouse	Sigma	F3040	AB_439712	IE: 1:100
p75NTR	rabbit	Cancer Research UK	CRD5410	AB_2864325	IE: 1:500
Lentiviral Vectors					
name		manufacturer	cat number		
TET ON Advance		Takara Bio (Clontech)	630930		
pLVX shRab10		Schiavo Lab	-		
pLVX Rab10 myc		Schiavo Lab	-		
Plasmids					
product	manufacturer		Addgene	RRID	reference
pEGFP-C1	Clontech			(discontinued)	
Rab10 WT EGFP	Marci Scidmore lab		49472	Addgene_49472	Huang 2010 <i>Cell Microbial</i>
Rab10 T23N EGFP	Marci Scidmore lab		49545	Addgene_49545	Huang 2010 <i>Cell Microbial</i>
TrkB-FLAG	Francis Lee lab		-	-	Chen 2005 <i>Mol Biol Cell</i>

Table S1: Primary antibodies, lentiviral vectors and plasmids used in this study. RRID: Research Resource Identifier. IF: immunofluorescence. IE: immuno-endocytosis.

Table S2 — Statistical summary

Figure		Variable	Test	Groups	Degrees of freedom	p value
1	D	TrkB accumulation	Unpaired t-Student	control : shRNA Rab10	t(140)	< 0.0001
	E	TrkB accumulation and Rab10 expression	Pearson r	control : shRNA Rab10	XY pairs= 131	< 0.0001
	G	pCREB abundance	one-way ANOVA	control, shRNA Rab10 and rescue	F(2,280)	< 0.0001
			Multiple comparisons	control : shRNA Rab10	t(280)	< 0.0001
			Multiple comparisons	control : rescue	t(280)	0.0336
2	B	Co-localisation Rab10 and Rab7	Unpaired t-Student	M1 control : BDNF	t(72.32)	0.0621
			Unpaired t-Student	M2 control : BDNF	t(62.33)	0.1043
	E	Area of overlay HcT and TrkB (retrograde TrkB)	one-way ANOVA	30, 60 and 90 mins	F(2,72)	< 0.0001
			Multiple comparisons	30 : 60 mins	t(72)	> 0.9999
			Multiple comparisons	60 : 90 mins	t(72)	0.0002
		Area of overlay retrograde TrkB and Rab10	one-way ANOVA	30, 60 and 90 mins	F(2,72)	0.2730
			Multiple comparisons	30 : 60 mins	t(72)	0.5717
			Multiple comparisons	60 : 90 mins	t(72)	> 0.9999
4	C	Direction of Rab10 organelles	Unpaired t-Student	Anterograde pre : post BDNF	t(14)	0.0150
			Unpaired t-Student	Retrograde pre : post BDNF	t(14)	0.0030
			Unpaired t-Student	Non-mobile pre : post BDNF	t(14)	0.4278
5	C	Recycling of TrkB	Unpaired t-Student	EGFP : Rab10 DN	t(60.40)	0.3914
	E	Co-localisation TrkB and Rab5	Unpaired t-Student	EGFP : Rab10 DN	t(38.22)	0.0080

Table S2: Summary of the statistical tests and their results for the data in the figures.