Maternal age affects equine Day 8 embryo gene expression both in

trophoblast and inner cell mass

E. Derisoud[1,2], L. Jouneau[1,2], C. Dubois[3], C. Archilla[1,2], Y. Jaszczyszyn[4], R. Legendre[5], N. Daniel[1,2], N.

Peynot[1,2], M. Dahirel[1,2], J. Auclair-Ronzaud[3], L. Wimel[3], V. Duranthon[1,2], P. Chavatte-Palmer[1,2]

[1] Université Paris-Saclay, UVSQ, INRAE, BREED, 78350, Jouy-en-Josas, France

[2] Ecole Nationale Vétérinaire d'Alfort, BREED, 94700, Maisons-Alfort, France

[3] IFCE, Plateau technique de la Valade, Chamberet, France

[4] Institute for Integrative Biology of the Cell (I2BC), UMR 9198 CNRS, CEA, Paris-Sud University F-

91198, Gif-sur-Yvette, France

[5] Institut Pasteur—Bioinformatics and Biostatistics Hub—Department of Computational Biology, USR

3756 IP CNRS, Paris, France

Corresponding authors:

Emilie Derisoud

emilie.derisoud@gmail.com

Pascale Chavatte-Palmer

pascale.chavatte-palmer@inrae.fr

Declarations

Ethics approval and consent to participate

The experiment was performed at the experimental farm of IFCE (research agreement C1903602 valid until March 22, 2023). The protocol was approved by the local animal care and use committee ("Comité des Utilisateurs de la Station Expérimentale de Chamberet") and by the regional ethical committee ("Comité Régional d'Ethique pour l'Expérimentation Animale du Limousin", approved under N° C2EA - 33 in the National Registry of French Ethical Committees for animal experimentation) under protocol number APAFIS#14963-2018050316037888 v2. All experiments were performed in accordance with the European Union Directive 2010/63EU.

Consent for publication

Not applicable

Availability of data and materials

The RNA sequençing data supporting the conclusions of this article are available in the GEO repository, [accession: GSE162893; https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE162893].

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the "Institut Français du Cheval et de l'Equitation" (grant numbers CS_2018_23, 2018). The National Research Institute for Agriculture, Food and Environment (INRAE) department Animal Physiology and Breeding Systems also supported this research.

Author contribution statement

PCP obtained the funding. PCP and VD conceived the project. LW, PCP, and VD supervised the study. ED, CD, CA, ND, NP, LW, VD and PCP adapted the methodology for the project. ED, CD, CA, YJ, JAR, and LW performed the experiments. CD, CA, ND, NP, MD and LW provided the resources. ED, LJ, YJ and RL performed data curation. ED and LJ analysed the data. ED wrote the original draft. All authors read, revised, and approved the submitted manuscript.

Acknowledgments

The authors are grateful to the staff of the Institut Français du Cheval et de l'Equitation (IFCE) experimental farm (Domaine de la Valade, Chamberet, France) for care and management of animals. We acknowledge the High-throughput sequencing facility of I2BC for its sequencing and bioinformatics expertise. The bioinformatics analyses were performed thanks to Core Cluster of the Institut Français de Bioinformatique (IFB) (ANR-11-INBS-0013). Many thanks to Matthias Zytnicki and Christophe Klopp for their advice on RNA-seq de novo analysis. Many thanks to Pablo Ross who kindly provided the coordinates for the XIST gene. The authors thank Shavahn Loux for her information on PLAC8A. The authors are grateful to Alice Jouneau and Sophie Calderari who kindly revised this article.

Abstract

Background: Increased embryo loss as mares become older is a major consideration for breeders as older animals are currently used for reproduction in the equine industry. Lower embryo quality has been pointed out as partly responsible for this reduced fertility. Here, the effect of mare's age on blastocysts' gene expression was explored. Day 8 post ovulation embryos were collected by uterine flushing from multiparous young (YM, 6-year-old, N = 5) and older (OM, > 10-year-old, N = 6) non-

nursing Saddlebred mares, inseminated with the semen of one stallion. Pure (TE_part) or inner cell

mass enriched (ICMandTE) trophoblast were obtained by embryo bisection and paired end, non-

oriented RNA sequencing (Illumina, NextSeq500) was performed on each hemi-embryo. To

discriminate gene expression in the ICM from that in the TE, deconvolution (DeMixT R package) was

used on the ICMandTE dataset. Differential expression was analyzed (DESeq2) with embryo sex and

diameter as cofactors using a false discovery rate < 0.05 cutoff. Functional annotation and classification

of differentially expressed genes and gene set enrichment analysis were also performed.

Results: Maternal aging did not affect embryo recovery rate, embryo diameter nor total RNA quantity

but modified gene expression in equine D8 embryos. The expression of genes in the ICM seemed to be

more altered by maternal aging than in the TE. In both compartments, the expression of genes involved

in mitochondrial function, translation and transcription due to chromatin modification were disturbed

by maternal age. Mitosis, signaling and adhesion pathways and embryo development were particularly

decreased in the ICM hemi-embryos from old mares. Finally, in TE, ion movement related genes were

affected.

Conclusions: This is the first study showing an effect of maternal age on gene expression in the equine

blastocyst at Day 8 post ovulation. Maternal age, even for mares as young as 10 years old, disturbs

mitosis, translation and transcription, cell signaling and adhesion as well as mitochondrial function and

cell commitment of horse embryos. These perturbations may affect further embryo development and

contribute to decreased fertility due to aging.

Keywords

horse; aging; blastocyst; methylation; mitochondria; mitosis; cell commitment

Background

Unlike other farm animals, the main purpose for the use of horses is athletic performance and leisure. In many cases, mares produce foals regularly to remain profitable after their sport career so that their reproductive career may last until advanced age, even though mares' fertility declines with age [1]. Indeed, in Thoroughbreds, pregnancy rates per estrous at Day 42 post-ovulation were shown to be reduced from 63.0% in young mares to 58.5% in mares older than 9 , and to reach 43.8% in mares older than 18 [2]. One study on 535,746 fertility records in mixed breeds in France showed that if a 10% of fertility decline is chosen as cutoff value for culling breeding mares, then mares older than 8 should not be bred [3].

Embryo collections and transfers at various developmental stages have been performed to determine at what stage embryos are affected and whether embryo or uterine quality is responsible for the fertility decline. Uterine blood flow during the luteal phase (Day 12-20 post ovulation) was reduced in mares >15 years old compared to mares <6 years old [4]. Moreover, at 2 days post ovulation, using oviductal flushes, the fertilization rate was similar between young and fertile mares *vs* old and subfertile mares (respectively, 10/14 *vs* 11/14 embryos recovered per group on Day 2). In contrast, in the same study, using the same mares on another estrus period, a significantly higher pregnancy rate at 14 days post ovulation was observed in young and fertile mares (respectively, 12/15 *vs* 3/15 pregnant mares on Day 14 by ultrasonography) [5]. Moreover, in practice, embryo loss is generally higher between the first (14 - 22 days) and the second (42 - 45 days) pregnancy check in old compared to young mares [2,6,7]. Using a ROC analysis based on 11 years of data from nursing Thoroughbred mares in Brazil, the cutoff value for embryo mortality to reach above average values was calculated to be 10 years of age [7]. In addition, Day 4 embryos collected from old and subfertile mares and directly transferred in synchronized young and fertile mares, were shown to result in lower pregnancy rates at 14 days post ovulation compared to embryos from young and fertile mares transferred in the same recipient group. This was true even when only good quality embryos were considered [8]. Thus,

embryo loss seems to be the principal cause of the fertility decline in old mares, even if uterine

degeneration is also observed in old mares.

In terms of embryo morphology, the effect of maternal age is not so clear, as studies demonstrated a

reduction of embryo size in old mares while others did not detect any effect of maternal aging on

embryo size. In these studies, however, the developmental stage was not uniform, which might lead

to these disparities [5,8–15]. For stages of development that take place in the oviduct, *i.e.*, from

fertilization to the late morula stage, one study on embryos collected *in vivo* on Day 1.5 post ovulation

showed that embryos have the same number of cells but a poorer morphology grade (uniformity of

cell size and texture, imperfections and abnormalities) when mares are older than 15 [11]. In addition,

when mares were older than 20 years, Day 3 post ovulation embryos were shown to combine a poorer

morphology grade with reduced cell number [11]. In contrast, in other studies, no difference according

to maternal age was observed in terms of diameter or number, size, texture of blastomeres of embryos

collected in oviducts on Days 2 and 4 post ovulation [5,8]. For later stages, the size of embryos collected

on Days 7 to 10 post-ovulation, in commercial embryo transfer programs in Italy, was reduced in mares

older than 16 years of age, without any morphological grade alteration [10]. In contrast, smaller

embryos, with a higher incidence of morphological abnormalities, were collected when donors were

subfertile and/or 18 years old mares compared to 4 years old nulliparous mares [12]. In embryos

produced by intracytoplasmic sperm injection (ICSI), maternal age did not affect blastomere size, nor

the presence of ooplasm fragments, nor membrane extrudates through zona pellucida at 24h post-

ICSI [9]. Embryo cell number at the time of transfer (between 6 and 9 days post-ICSI), however, was

higher in embryos collected from mares older than 20 [8].

In terms of functional potential, a reduced quantity of mitochondrial DNA, but not genomic DNA, has

been reported in Day 7 embryos from mares older than 16 [16]. These data suggest that old mares'

blastocysts have the same number of cells than those of younger mares, but with less mitochondria

[16], which could be a cause or an effect of compromised embryo development in old mares.

Moreover, embryos from mares older than 15 seem to be less mobile before implantation than those

from mares aged 5-6 [14]. This could indicate impaired embryo development, alteration of uterine

contractions or both.

Altogether, these data indicate that the quality of embryos from older mares is reduced. Subsequently,

dam age could also affect placental and fetal intrauterine and offspring post-natal growth and long-

term health, in the context of the Developmental Origins of Health and Diseases (DOHaD) [17]. To our

knowledge, the role of the embryo *per se*, however, has not yet been explored. Moreover, age and

parity are generally positively correlated and their respective impact have not been determined yet.

The aim of this study was to determine the effect of maternal age in the mare on embryo gene

expression at the blastocyst stage. Young (<10 years) and older (>10 years) multiparous mares were

inseminated with the same stallion. Blastocysts were collected and bisected to separate the pure

trophoblast (TE_part) from the inner cell mass enriched hemi-embryo (ICMandTE). Gene expression

was analyzed by RNA-seq in each compartment. Deconvolution was used to isolate gene expression of

inner cell mass from trophoblast cells in ICMandTE part. Results shows that maternal age alters gene

expression of both equine Day 8 trophoblast and inner cell mass.

Materials and methods

Ethics

The experiment was performed at the experimental farm of IFCE (research agreement C1903602 valid

until March 22, 2023). The protocol was approved by the local animal care and use committee ("Comité

des Utilisateurs de la Station Expérimentale de Chamberet") and by the regional ethical committee

("Comité Régional d'Ethique pour l'Expérimentation Animale du Limousin", approved under N° C2EA -

33 in the National Registry of French Ethical Committees for animal experimentation) under protocol

number APAFIS#14963-2018050316037888 v2. All experiments were performed in accordance with

the European Union Directive 2010/63EU.

Embryo collection

Thirty-one multiparous Saddlebred mares (French Anglo-Arabian and Selle Français breeds) were

included in this study. Multiparous mares were defined as dams that had already foaled at least once.

Mares were bred and raised in the "Institut Français du Cheval et de l'Equitation" experimental farm

(Chamberet, France, 45° 34'55.17"N, 1°43'16.29"E, 442m). During the experimental protocol, mares

were managed in herds in natural pastures 24h/day with free access to water. The experiments took

place from April 1st to May 23rd, 2019. All mares remained healthy during this period.

Mares were allocated to 2 groups according to their age: young mares (YM, < 10 years, n = 11) and

older mares (OM, ≥ 10 years, n = 20). The age threshold was defined in accordance with the literature

[3,7]. Before experimentation, mare's withers' height was measured. During the experimentation,

mares were weighted and mares' body condition score (BCS, scale 1 to 5) was determined by one

trained operator [18]. Mares' characteristics are detailed in Table 1.

Table 1: Mares' characteristics at embryo collection time.

Characteristics	Young mares (YM) (n=11)	Old mares (OM) (n=20)
Breed	AA n = 1; SF n =9; SB n=1	AA n = 13; SF n = 7
Age *(in years)*	6.00 ± 0.00	11.95 ± 2.11
Parity *(number of foalings)*	1.00 ± 0.00	3.20 ± 1.15
Weight *(in kg)*	536.62 ± 44.12	589.35 ± 57.34
BCS *(scale 1-5)*	2.16 ± 0.28	2.61 ± 0.32
Withers' height *(in cm)*	157.95 ± 3.74	161.21 ± 4.23

All mares were multiparous.

AA: Anglo Arab or Anglo-Arabian type; SF: Selle Français; SB: Saddlebred; BCS: Body Condition Score.

Age, parity, weight, BCS and height are presented as mean ±SD

The mares' estrous period was monitored routinely by ultrasound with a 5MHz trans-rectal transducer.

During estrus, ovulation was induced with a single injection of human chorionic gonadotropin (i.v.; 750

- 1500IU; Chorulon® 5000; MSD Santé animale, France) as soon as one ovarian follicle > 35mm in

diameter was observed, together with marked uterine edema. Ovulation usually takes place within

48h, with > 80% occurring 25 to 48h after injection [19,20]. At the same time, mares were inseminated

once with fresh or refrigerated semen containing at least 1 billion motile spermatozoa from a single

fertile stallion. Ovulation was confirmed within the next 48 hours by ultrasonography.

Embryos were collected by non-surgical uterine lavage using prewarmed (37°C) lactated Ringer's

solution (B.Braun, France) and EZ-Way Filter (IMV Technologies, France) 10 days after insemination,

i.e. approximately 8 days post ovulation. Just after embryo collection, mares were treated with

luprotiol an analogue of prostaglandin F2α (i.m; 7.5 mg; Prosolvin, Virbac, France).

Embryo bisection and RNA extraction

Collected embryos were immediately observed using a binocular magnifying glass and photographed

with a size standard. Embryo diameter was subsequently determined on photographs using ImageJ®

software (version 1.52a; National Institutes of Health, Bethesda, MD, USA). Embryos were washed 4

times in commercially available Embryo holding medium (IMV Technologies, France) at 34°C and then

bisected with a microscalpel under binocular magnifying glass to separate trophoblast part, at this

stage composed of trophectoderm and endoderm (TE_part) from inner cell mass (composed of

epiblast cells) enriched hemi-embryos (ICMandTE) (Figure 1). Each hemi-embryo was transferred to an

Eppendorf DNA LoBind tube (Eppendorf, Montesson, France) with extraction buffer (PicoPure RNA

isolation kit, Applied Biosystems, France) and incubated for 30 min at 42°C prior to storage at - 80°C.

RNA was extracted from each hemi-embryo following the manufacturer's instructions using PicoPure

RNA isolation kit (PicoPure RNA isolation kit, Applied Biosystems, France), which included a DNAse

treatment. RNA quality and quantity were assessed with the 2100 Bioanalyzer system using RNA 6000

Pico or nano kit (Agilent Technologies, France) according to the manufacturer's instructions (Additional

file 1).

RNA sequencing

Five nanograms of total RNA were mixed (Additional file 1) with ERCC spike-in mix (Thermofisher

Scientific, France) according to manufacturer recommendations. Messenger RNAs were reverse

transcribed and amplified using the SMART-Seq V4 ultra low input RNA kit (Clontech, France) according

to the manufacturer recommendations. Nine PCR cycles were performed for each hemi-embryo. cDNA

quality was assessed on an Agilent Bioanalyzer 2100, using an Agilent High Sensitivity DNA Kit (Agilent

Technologies, France). Libraries were prepared from 0.15 ng cDNA using the Nextera XT Illumina library

preparation kit (Illumina, France). Libraries were pooled in equimolar proportions and sequenced

(Paired end 50-34 pb) on NextSeq500 instrument, using a NextSeq 500 High Output 75 cycles kit

(Illumina, France) (Additional file 1). Demultiplexing was performed with bcl2fastq2 version 2.2.18.12

(Illumina, France) and adapters were trimmed with Cutadapt version 1.15 [21]. Only reads longer than

10pb were kept.

Embryo sexing

Embryo sex was determined for each embryo in order to take in consideration embryo sex in the

statistical analysis as sexual dimorphism was observed in blastocysts in other species [22].

The expression of *X Inactive Specific Transcript* (*XIST*) was analyzed. *XIST* is a long noncoding RNA from

chromosome X responsible for X inactivation when two X chromosomes are present, meaning that

only female embryos express *XIST*. *XIST* was shown to be expressed in equine female blastocysts on

Day 8 post ovulation [23,24]. *XIST* is not annotated in the available equine Ensembl 3.0.99 or NCBI

annotation. Thus, a *de novo* transcript assembly was made using StringTie version 2.1.2 [25] following

manufacturer instructions to identify *XIST* in equine embryo samples (Additional file 1). The genome

annotation file output was aligned on Ensembl 99 EquCab3.0 assembly using IGV version 2.7.2 [26]. As

XIST position and length on X chromosome has been previously determined (Pablo Ross, personal

communication, [23]) and as *XIST* is annotated in humans, the most similar contig was selected. The

obtained *XIST* sequence was 22,600 bp long and composed of 8 exons and 7 introns. *XIST* sequence

was added to FASTA files from Ensembl 99 EquCab3.0 assembly and to the genome annotation file

from Ensembl 99 EquCab3.0 assembly as a unique fictitious chromosome with one gene composed of

one transcript.

XIST expression differed between individuals but was relatively close in quantification between related

hemi-embryos. Therefore, seven embryos were determined as females (4 in the YM group and 3 in the

OM group) while 4 were considered as males (1 in the YM group, and 3 in the OM group).

RNA mapping and counting

Alignment was performed using STAR version 2.6 [27] (2-pass mapping with the genome annotation

file using the following parameters: --alignSoftClipAtReferenceEnds No --alignEndsType EndToEnd --

alignIntronMax 86545 --alignMatesGapMax 86545) on previously modified Ensembl 99 EquCab3.0

assembly and annotation. Genes were then counted with featureCounts [28] from Subreads package

version 1.6.1 (parameters: -p -B -C -M) (Additional file 1).

Data analysis

All statistical analyses were performed by comparing OM to YM (set as reference group) using R version

4.0.2 [29] on Rstudio software version 1.3.1056 [30].

Embryo recovery rate, embryo diameter and total RNA content analysis

Embryo recovery rates (ERR) were calculated as number of attempts with at least one embryo

collected/total number of attempts. They were analyzed using the Exact Fisher test to determine if age

influences embryo recovery rate.

For total RNA content analysis, as embryos were bisected without strict equality for each hemi-

embryo, a separate analysis of ICMandTE and TE_part RNA quantities would not be meaningful. Thus,

ICMandTE and TE_part RNA quantities were initially summed up and analyzed using a mixed linear

model from nlme package version 3.1-148 [31], followed by 1000 permutations using PermTest

function from pgirmess package version 1.6.9 [32]. Maternal age and embryo sex were considered as

random effects.

Embryo diameter was analyzed with a fixed effects linear model of nlme package version 3.1-148 [31]

including maternal age and embryo sex, followed by 1000 permutations using PermTest function from

pgirmess package version 1.6.9 [32]. Variables were kept in models when statistically significant

differences were observed. Differences were considered as significant for $p < 0.05$.

Moreover, the relation between embryo size and total RNA quantity/embryo was studied and

represented using the nonlinear regression analysis (fitness to exponential growth equation) in

GraphPad Prism software 8.0.1 for Windows (Graphpad Software, San Diego, California USA,

www.graphpad.com).

Deconvolution of gene expression in ICMandTE using DeMixT

A filter on counts obtained with featureCounts was applied: all genes with more than 3 non null count

values in at least one group (OM or YM) per hemi-embryo (ICMandTE or TE_part) were conserved. The

ICMandTE hemi-embryo was composed of both trophoblast and inner cell mass in unknown relative

proportions. In order to estimate the relative gene expression of both cell types, the DeMixT R package

version 1.4.0 was used [33,34]. Starting from a dataset obtained from one cell type, DeMixT estimates

the proportion of cells in the mixed samples and performs a deconvolution algorithm on gene

expression. TE_part (reference cells) and ICMandTE (mixed cells type) datasets were thus analyzed

using the following options: niter=10; nbin=50; if.filter=TRUE; ngene.selected.for.pi=1500; ngene.

Profile.selected=1500; nspikein=0. Since the deconvolution algorithm cannot be performed in the

presence of null values, the value "1" was added to all count values in ICMandTE and TE_part data

tables. Moreover, variance in reference cell dataset (TE_part) must not be null and gene with null

variance had been removed prior to the deconvolution. Output datasets were DeMixT_ICM_cells and

DeMixT_TE_cells, corresponding to the deconvoluted gene expression in ICM cells and TE cells of

ICMandTE, respectively.

Quality checks for deconvolution

At the end of deconvolution, a quality check is automatically performed by the analysis: for every gene,

if the difference of average of deconvoluted expression of reference cells in mixed samples (here, gene

expression of TE cells in ICMandTE dataset, *i.e.,* DeMixT_TE_cells) and average of expression of

reference cells (here, gene expression of TE cells in TE_part) is > 4, then, the deconvolution for

concerned genes is not reliable. Therefore, genes in this case are filtered out.

Moreover, outputs of DeMixT_ICM_cells *vs* DeMixT_TE_cells, DeMixT_ICM_cells *vs* TE_part and

ICMandTE *vs* TE_part were compared with Deseq2 version 1.28.1 [35] to confirm that the

deconvolution was effective at separating gene expression. Moreover, several genes proper to ICM

and TE cells in equine embryos were identified using literature search [23]. Their raw and deconvoluted

expressions were analyzed to check the reliability of deconvolution. Results of these analyses were

represented through manually drawn Venn diagrams as well as principal component analysis graphics

of individuals, using ggplot2 version 3.3.3 [36] and factoextra version 1.0.7 [37].

Maternal age comparison for gene expression

For the comparison of maternal age, ICMandTE, DeMixT_ICM_cells and TE_part datasets were used

and a filter was applied on datasets: all genes with an average expression equal or above 10 counts in

at least one group (OM or YM) per hemi-embryo (ICM or TE) were conserved. Differential analyses

were performed with Deseq2 version 1.28.1 [35] with YM group as reference, without independent

filtering and taking into account embryo diameter and sex in the model. Genes were considered

differentially expressed (DEG) for FDR < 0.05 after Benjamini-Hochberg correction (also known as false

discovery rate, FDR).

Equine Ensembl IDs were converted into Human Ensembl IDs and Entrez Gene names using gorth

function in gprofiler2 package version 0.1.9 [38]. Genes without Entrez Gene names using gprofiler2

were manually converted when Entrez Gene names were available, using Ensembl web search function

[39].

Moreover, DEG were analyzed for classification, over- and under- representation (FDR < 0.05) using

Fisher's exact test corrected with FDR and using the database Gene Ontology (GO) Biological Process

complete on PANTHER version 16.0 web-based software [40]. Only the most specific subclass

pathways, provided by PANTHER's automated hierarchy sorting, were conserved and represented.

Gene set enrichment analyses (GSEA)

The count values were log transformed using RLOG function of DESeq2 version 1.28.1. Gene set

enrichment analyses (GSEA) were performed on expressed genes using GSEA software version 4.0.3

(Broad Institute, Inc., Massachusetts Institute of Technology, and Regents of the University of

California) [41,42] to identify biological gene sets disturbed by age. GSEA was performed using the

following parameters: 1000 gene set permutations, weighted enrichment analysis, gene set size

between 15 and 500, signal-to-noise metrics. Molecular Signatures Database [43] version 7.1 (C2:

KEGG: Kyoto Encyclopedia of Genes and Genomes, C5: BP: GO biological process) were used to identify

most perturbed pathways. Pathways were considered significantly enriched for FDR < 0.05. When

normalized enrichment score (NES) was positive, the gene set was enriched in the OM group while

when NES was negative, the gene set was enriched in the YM group.

Enriched terms from GO BP and KEGG database were represented using SUMER analysis from SUMER

R package version 1.1.5 and using FDR q-values [44]. First, SUMER analysis reduces redundancy using

weighted set cover algorithm and then clusters the results using affinity propagation algorithm. The

weighted set cover algorithm performs a selection of the fewest number of gene sets that include all

genes associated with the enriched sets, giving priority for the most significant terms. Then, the affinity

propagation algorithm groups similar gene sets and identifies one representative gene set. This

algorithm will repeat this clustering to obtain the most representative gene sets. Results were

represented with graph modified using Cytoscape version 3.8.2 [45]. Gene sets were represented by

nodes and the gene set size was represented by the size of the node. Node shape represented the

gene set database (GO BP or KEGG). Blue nodes represented gene sets enriched in YM (NES < 0) while

red nodes represented gene sets enriched in OM (NES > 0). Edge width represents the level of

connection between representative gene sets (thinner edges represent the first clustering while

thicker edges represent the second clustering of the affinity propagation algorithm)

Results

Embryo rates, diameter, total RNA content and quality

Altogether, 36 uterine flushings were performed (12 in YM and 24 in OM, 5 mares being flushed twice).

No twin embryos were obtained. Embryo recovery rate was 42% and 25% in YM and OM, respectively

and did not differ between groups (p = 0.45). All embryos were expanded blastocysts grade I or II

according to the embryo classification of McKinnon and Squires [46].

Embryo diameter ranged from 409 µm to 2643 µm, with no effect of group on embryo diameter (p =

0.97). RNA yield per embryo ranged from 16 ng to 2915.5 ng and was not related to mares' age (p =

0.92). There is a clear exponential relation between RNA yield/embryo and embryo size (Figure 2).

The median RNA Integrity Number (RIN) was 9.8 (8.7 - 10 range). Between 29.4 and 69.5 million reads

per sample were obtained after trimming. On average, 83.61% of reads were mapped on the modified

EquCab 3.0 using STAR and 66.66% of reads were assigned to genes by featureCounts.

Deconvolution of gene expression to discriminate ICM and TE gene expression in ICMandTE

hemi-embryos

After keeping genes with less than 3 non null count values in at least one group (OM or YM) per hemi-

embryo (ICMandTE or TE_part), 16,741 genes were conserved for deconvolution. In addition, twenty-

seven genes were removed because their variance was null in the TE_part. However, for these genes,

the mean counting in ICMandTE samples was above 10 counts, excepted for one which had a mean

expression of 34 counts. On the other hand, this gene was only expressed in 5 ICMandTE over the 10

counts threshold.

During the analysis, 3 additional genes were excluded because the deconvolution quality for these

genes was not sufficient. Therefore, at the end of deconvolution algorithm, 16,711 genes were

available for differential analysis.

Before deconvolution, 341 genes were differentially expressed (FDR < 0.05) between the ICMandTE

and the TE_part (Figure 3A). After deconvolution, comparison between DeMixT_ICM_cells and

DeMixT_TE_cells yielded 7,565 differentially expressed genes while the comparison DeMixT_ICM_cells

vs TE_part yielded 6,085 differentially expressed genes, with 5,925 in common (72%). Moreover,

except one of the initially 341 differentially expressed genes before deconvolution, were also identified

as differentially expressed in both post-deconvolution analyses. On the PCA graphic of individuals,

ICMandTE and TE_part were not really separated before deconvolution (Figure 3B). DeMixT_TE_cells

and TE_part were partly superposed, suggesting that datasets before and after deconvolution have a

similar global gene expression; whereas the DeMixT_ICM_cells group is clearly separated from both

on Axis 2 (14.3% of variance), indicating that the deconvolution effectively enabled the separation of

gene expression in the two cell types.

Six of the 12 genes previously identified as more expressed in ICM [23] were also more expressed in

ICMandTE *vs* TE_part comparison (Table 2). After deconvolution (comparison DeMixT_ICM_cells *vs*

TE_part), 3 more genes previously identified were also differentially expressed. However, for two of

them (*POU Class 5 Homeobox 1*, *POU5F1* and *Undifferentiated Embryonic Cell Transcription Factor 1*,

UTF1), their expression was increased in TE_part after deconvolution. In TE, any genes previously

identified were differentially expressed in the comparison ICMandTE *vs* TE_part, *i.e.*, before

deconvolution. After deconvolution, the expression of 2 genes was increased in TE_part compared to

DeMixT_ICM_cells.

Table 2: Comparison of selected genes expression before and after deconvolution

	Gene name	Ensembl ID	ICMandTE *vs* TE_part		DeMixT_ICM_cells *vs* TE_part	
			log2FC from DeSeq2	padj	log2FC from DeSeq2	padj
ICM	SOX2	ENSECAG00000010653	4.49	0.02	6.49	1.7E-06
	NANOG	ENSECAG00000012614	4.24	1.0E-07	5.98	2.4E-15
	SPP1	ENSECAG00000017191	4.02	8.4E-09	5.74	5.1E-16
	LIN28B	ENSECAG00000020994	3.23	1.3E-10	5.00	4.7E-22
	SMARCA2	ENSECAG00000024187	0.62	0.97	1.25	0.07
	POU5F1 (OCT4)	ENSECAG00000008967	0.60	0.009	-2.04	4.6E-07
	ID2	ENSECAG00000008738	0.44	0.29	1.24	9.0E-05
	DNMT3B	ENSECAG00000012102	0.35	0.24	0.8	0.06
	DPPA4	ENSECAG00000013271	0.35	0.04	1.14	1.8E-05
	SALL4	ENSECAG00000018533	0.10	1	0.60	0.01
	KLF4	ENSECAG00000010613	-0.04	1	-0.81	0.39
	UTF1	ENSECAG00000039888	-0.13	1	-2.21	7.3E-05

TE	**TFAP2A**	ENSECAG00000017468	-0.17	0.46	-0.30	0.004
	CDX2	ENSECAG00000027754	-0.14	0.77	-0.23	0.07
	ELF3	ENSECAG00000014608	-0.13	1	-0.19	0.27
	GATA2	ENSECAG00000016768	-0.13	1	-0.29	0.048
	GATA3	ENSECAG00000024574	-0.10	0.87	-0.05	0.64
	TEAD4	ENSECAG00000011303	-0.10	0.96	0.00	1
	FREM2	ENSECAG00000020410	0.07	1	0.19	0.44

Gene expressions were obtained from RNA of 11 equine embryos bissected in two hemi-embryos: one

part is composed only of trophoblast (TE), TE_part, while the other part is composed of TE and inner

cell mass (ICM), ICMandTE. As it is impossible to estimate the proportion of each cells in ICMandTE,

deconvolution algorithm (package DeMixT) was used to estimate gene expression of these different

kind of cells. DeMixT_ICM_cells dataset is the deconvoluted gene expression of ICM cells from

ICMandTE. Log2 fold change (log2FC) and padj (adjusted p-value with Benjamini-Hochberg correction)

were obtained with Deseq2 package. TE_part is the reference group in both analyses: when log2 fold

changes (log2FC) > 0, gene is more expressed in ICMandTE or DeMixT_ICM_cells while when log2FC <

0, gene is more expressed in TE_part. Green is used to represent gene differentially expressed in the

present study. Orange is used to represent gens that have been previously identified as predominant

in the ICM [23] but which are identified here as predominant in the TE.

For further analysis, DeMixT_ICM_cells and TE_part gene expression datasets are presented.

Nevertheless, the comparison of maternal age on ICMandTE dataset (graphics, differential gene

expression analysis and GSEA) was also performed and this analysis is available in Additional file 2, 3

and 4.

Differential gene expression in deconvoluted ICM cells' genes expression according to maternal

age

After filtering out of genes with an average expression < 10 counts/maternal age group/hemi-embryo,

13,735 genes were considered as expressed in the ICM cells from OM or YM embryos including 255

differentially expressed genes (227 downregulated and 28 upregulated in OM) (Figure 4A and

Additional file 5). Only 218 and 12 genes out of the down- and upregulated genes, respectively, were

associated to a known protein in human. These downregulated genes in ICM of embryos from OM

mainly belonged to cellular process, metabolic process and biological regulation in GO biological

process gene sets (Figure 4A). From downregulated genes in OM ICM, 84 pathways were statistically

overrepresented in GO Biological process leading to 16 most specific subclass pathways according to

hierarchy sorting by PANTHER (Figure 4B). Overrepresented pathways in downregulated genes in OM

ICM were linked to cell division, embryo morphogenesis, histone methylation, protein modification

processes and cytoskeleton organization.

Upregulated DEGs in OM ICM were mainly part of biological regulation, response to stimulus, cellular

process, signaling and metabolic process in GO biological process gene sets (Figure 4A). No pathway

was statistically overrepresented in upregulated genes in ICM of embryos from old mares using

PANTHER overrepresentation testing.

Differential gene expression in pure TE part gene expression

In the TE, 13,178 genes were considered as expressed in OM or YM. Fourteen were differentially

expressed (Additional file 6) with 11 genes being downregulated and 3 being up regulated in OM

(Figure 4A). Only 12 genes were associated to a known protein in human. Downregulated genes in OM

mainly belonged to both cellular and metabolic process in GO biological process gene sets.

Gene set enrichment analysis in deconvoluted ICM cells' genes expression according to

maternal age

After Entrez Gene ID conversion, 12,344 genes were considered expressed in ICM. In the GO Biological

Process database, 106 identified gene sets differed according to maternal age in the ICM (11 enriched

in OM and 95 enriched in YM) (Additional file 7). In the KEGG database, 14 gene sets were perturbed

(Additional file 7).

Using SUMER analysis (Figure 5), gene sets enriched in OM ICM were mainly grouped under terms

"Respiratory electron transport chain" and "Translational elongation". The first term represents

mitochondrial function while the second represents translation function. Moreover, terms relative to

"Systemic lupus erythematosus" are conserved by SUMER analysis and enriched in OM ICM. Genes

involved in these enrichments were also present in dysregulated metabolic pathways.

Gene sets enriched in YM ICM were involved in different biological processes such as "Mitotic nuclear

division" which is relative to mitosis. Under this term, "Spindle organization" and "Negative regulation

of supramolecular fiber organization" terms represent the assembly of spindle during mitosis.

"Chromosome segregation" is also found in this grouping. Furthermore, terms under "Golgi vesicle

transport" seem to be involved in cytokinesis and vesicle trafficking. The catabolism of protein and

mRNA is represented by "Ubiquitin mediated proteolysis". Translation function is represented by

several terms grouped under "covalent chromatin modification" in the mitosis part.

"Leukocyte transendothelial migration" term is also enriched in YM ICM and seems to represent cell

signaling and adhesion. Indeed, this term regroups several pathways such as "Focal adhesion",

"Adherens junction" and "WNT signaling pathway".

Finally, further terms related to embryo development could be found in different groups such as

"Dendrite development", "Cell differentiation in spinal cord" or "endothelial cell development".

Gene set enrichment analysis in TE

After Entrez Gene ID conversion, 11,874 genes were considered expressed in TE from OM or YM

embryos. In GO BP database, 38 gene sets were perturbed by mares' age in the TE of embryos (11

enriched in YM and 27 enriched in OM) (Additional file 8). In the KEGG database, 5 pathways were

enriched in the TE of OM embryos (Additional file 8).

As in ICM part, using SUMER analysis, translation (represented by both "Translational termination" and

"Ribosome" terms) and mitochondrial function (represented by "Oxidative phosphorylation"

associated terms) were enriched in OM embryos. "Proteasome" term is enriched in OM embryos

representing protein catabolism.

Moreover, transcription (represented by "Peptidyl lysine trimethylation" term) and embryo

development (represented by "spinal cord development") were, as in ICM, also, enriched in YM

embryos. Finally, in TE, "Regulation of action potential" term and associated terms were enriched in

YM. These terms could be related to ion movement.

Discussion

Maternal aging did not affect embryo recovery rate, embryo size nor total RNA quantity. Moreover,

total RNA content in equine embryo was exponentially linked to embryo diameter and was not affected

by maternal age.

As in tumor, today, the only way to analyze separate expression of ICM and TE cells in mammalian

embryos is to use micromanipulation but micromanipulation is expensive in time and requires adapted

skills and material. In this study, micromanipulation to properly separate ICM and TE was not possible.

Therefore, deconvolution appears to be a good option to analyze the effects of maternal ICM gene

expression in mixed ICM and TE samples. Deconvolution seemed to have been effective on embryo to

separate ICM and TE gene expression. Indeed, the 341 genes differentially expressed in ICMandTE *vs*

TE_part were also identified by comparing TE with post-deconvolution ICM cells gene expression

dataset. Moreover, after deconvolution, more genes were differentially expressed between TE and

ICM datasets and among them, more genes identified in literature were differentially expressed.

Finally, PCA showed that deconvolution separate DeMixT_ICM_cells from other datasets while before

deconvolution ICMandTE and TE_part were partly superposed. Only POU5F1 and UTF1 differed in

terms of allocation between previously published data [23] and the present study. In data before

deconvolution, expression of POU5F1 and UTF1 were extremely variable according to embryo

diameter in both compartments. Deconvolution reduced the variability of both gene expression in ICM

but not in TE. Studies showed that POU5F1 is expressed in both compartments until Day 10 post

ovulation in equine in vivo produced embryos [47] but its expression decreases progressively from Day

7 in TE cells. Thus, the statistically increased expression of POU5F1 in TE on Day 8 is compatible with

previously published data.

Therefore, it was chosen to only present the analysis of the effects of age in DeMixT_ICM_cells dataset.

In all cases, deconvolution did not change the principal results of maternal aging on embryo gene

expression because the comparison of maternal age in ICMandTE and DeMixT_ICM_cells gave

comparable results.

In both ICM and TE, the expression of genes involved in mitochondrial function and translation with

ribosome biogenesis were enriched in OM embryos while transcription with chromatin modification

was enriched in YM embryos (Figure 6). Mitosis, signaling and adhesion pathways were additionally

altered and the expression of genes involved in embryo development was particularly reduced in the

ICM hemi-embryos from old mares. Finally, in the TE, ion movement related genes were affected.

Several studies on mature oocytes in women and mice, using microarrays and RNAseq, had shown a

differential expression of genes involved in mitochondrial, cell cycle, cytoskeleton functions and

related pathways according to maternal age [48–50], as observed in the present study in equine

blastocysts, especially in the ICM. In equine follicles and oocytes, gene expression, as studied by RT-

qPCR, quantitatively and temporally differed with maternal age, suggesting a desynchronization of

essential genes for maturation [51]. Moreover, a study on the expression of 48 genes in cumulus-

oocytes complexes from young (< 12 years of age) and old (> 18 years old) mares showed differential

expression that may be related with the reduced developmental competence observed in old mares'

oocytes [52]. Finally, genes related to mitochondrial replication and function were down-regulated in

oocytes of old mares [16]. Even effects of maternal aging on gene expression in equine oocytes, using

high throughput techniques, are not yet available, results on equine Day 8 blastocysts suggest that

mitochondrial perturbations present in the oocyte persist until the blastocyst stage. In humans, single-

embryo RNA-seq analysis showed that maternal aging alters the transcriptome of blastocysts derived

from ICSI. Genes altered by maternal age were related to segregation of chromosomes, embryonic

growth and cell signaling [53,54], as observed in our study on equine blastocysts.

The equine embryo remains spherical, grows very fast and is surrounded by its glycoprotein capsule

up to 22 days post ovulation [55]. At 8 days post ovulation, equine embryos are therefore growing and

the endodermal layer entirely delineates the future yolk sac (Figure 6) [56]. Moreover, whereas in Day

7 embryos, the epiblast is lined by a continuous layer of trophoblast named the polar trophoblast. As

the blastocyst grows, epiblast cells phagocyte polar trophoblast cells that become discontinuous.

During this time, epiblast and polar trophoblast cells divided and are bound by tight junctions [57]. The

turnover in focal adhesion is essential for cell movement and phagocytosis [58].

Cell signaling and adhesion function were altered in OM embryos. Indeed, KEGG pathways "focal

adhesion" and "adherens junction" regrouped in "Leukocyte transendothelial migration" on Figure 4

are disturbed in old mares' embryos. Focal adhesions are subcellular structures observed in adherent

cells, that are clusters of different structural and signaling molecules and integrins that form a physical

link between the actin cytoskeleton and the extracellular matrix [59,60]. Integrins are the major

receptors for cell adherence to extracellular matrix proteins in focal adhesions. Indeed, they play an

important role in transmembrane connections to the cytoskeleton and activation of several signaling

pathways, as growth factor receptors [61]. *Integrin Subunit Alpha V* (*ITGAV*) expression was reduced

in the ICM (log2FC = -1.66; FDR < 0.05) of embryos collected from OM. In different human breast cancer

cell lines, cell proliferation, invasion and renewal are restricted when *ITGAV* is inhibited [62].

Therefore, the reduction of *ITGAV* could lead to an impaired cell signalization in OM embryos.

Moreover, small GTPases within the Rho family are required for the turnover of actin filaments and

cell adhesions necessary for cell migration, that are essential processes for the proper development of

the embryo. Activity and localization of each Rho GTPase must be precisely regulated as the

communication between them is indispensable to achieve this process [63]. The *Rho GTPase Activating*

Protein 9 and *35* (*ARHGAP9* and *ARHGAP35*) were downregulated (respectively, log2FC = -1.83, FDR <

0.05 and log2FC = -0.74, FDR < 0.05) in OM ICM. *ARHGAP35* codes for P190 Rho GTPase activating

protein that mediates signalization through integrin-dependent adhesion in cells. Aberrant neural

morphogenesis, related to an accumulation of polymerized actin in embryoneural tube cells, is

reported in mice lacking P190 Rho GTPase activating proteins [64]. Furthermore, P21 activated kinases

(PAK) are effector kinases for other small Rho GTPases such as Rac and regulate the polymerization of

actin and can affect the microtubule organization and therefore the regulation of cytoskeleton [65].

Once activated by this signaling pathway, PAKs promote the turnover of focal adhesions. PAKs will

then, trigger p38 complex activation that regulates transcription [66]. *PAK3* was downregulated in the

OM ICM (log2FC = -3.76, FDR < 0.001). However, in TE, *SLIT-ROBO Rho GTPase Activating Protein 3,*

SRGAP3 was one of the upregulated genes in OM (log2FC = 2.52, FDR < 0.01). This gene regulates

cytoskeletal reorganization and plays an important role in the formation of cell-cell adhesion [67].

Therefore, maternal aging seems to alter focal adhesion in both ICM and TE of embryos but, while in

ICM, it seems to reduce cohesion and signalization, in TE it seems to reinforce them.

Several signaling pathways required for development were altered and numerous genes related to

focal adhesions and signaling pathways were down-regulated in OM ICM (*ARHGAP9* and *ARHGAP35*;

B-Raf Proto-Oncogene, Serine/Threonine Kinase, *BRAF*; *Filamin C*, *FLNC*; *FYN Proto-Oncogene, Src*

Family Tyrosine Kinase, *FYN*; *PAK3*; *Protein Phosphatase 1 Regulatory Subunit 12A*, *PPP1R12A*; *Talin 2,*

TLN2; *X-Linked Inhibitor Of Apoptosis*, *XIAP*). Among all signaling pathways, transforming growth factor

beta (TGF-β), Wg et Int (Wnt), mitogen-activated protein kinases (MAPK) and apoptosis signaling were

altered by maternal age in equine embryos. Indeed, *transforming growth factor beta receptor 1*

(*TGFBR1*) was downregulated (log2FC = -4.34; FDR < 0.05) in the ICM of embryos from old mares.

TGFBR1 is one of the two dimers for the receptor for transforming growth factors beta. This factor

family is essential in the generation of axes and cell fate commitment during mammal embryogenesis

[68]. Therefore, the reduced production of TGF-β receptor in ICM could impair the establishment of

cell lines in old mares' embryos. Moreover, *Casein Kinase 1 Alpha 1*, (*CSNK1A1*) and *Protein*

Phosphatase 2 Regulatory Subunit B'Epsilon (*PPP2R5E*), involved in Wnt signaling pathway, were

downregulated in OM embryos. Wnt signaling pathway is also involved in nervous development

system and may act on cell determination in mice embryos [69]. In horses, gastrulation begins by Day

12 post ovulation and the primitive streak develops quickly after mesoderm formation [70]. It is

therefore possible that the alteration of the Wnt signaling pathway could have a negative impact on

the brain development of the future fetus. This hypothesis could be confirmed by the fact that *DMRT3*

(*Doublesex And Mab-3 Related Transcription Factor 3*) was downregulated in the OM ICM (log2FC = -

3.07, FDR < 0.05). DMRT3 is required for the proper development of the brain in prenatal chicken and

mice [71]. Moreover, mutations of this gene are responsible for different gaits in horses (specific to

aptitude to trot instead of cantering) [72]. Therefore, altogether, the data indicate that cell fate

commitment in ICM seems to be altered by maternal aging.

Cell cycle related pathways and genes are also disturbed by maternal aging. Indeed, the expression of

Pleckstrin Homology domain-Interacting Protein (*PHIP*), also known as replication initiation

determinant, was reduced (log2FC = -0.86, FDR < 0.01) in the OM ICM. This gene is required for the

initiation of DNA replication. Fibroblasts originated from mice embryos with depleted *PHIP* expression

presented less replication-initiation as well as abnormal replication fork progression events [73].

Moreover, several cyclins (CCN) were less expressed in the ICM of embryos from old mares (*CCNB3*,

CCNI and *CCNT2*). Cyclins are key cell cycle regulators and, for instance cyclin B3 degradation allows

the passage from metaphase (chromatids aligned on metaphase plate) to anaphase (segregation of

sister chromatids) [74]. During metaphase and anaphase, spindle assembly and chromosome

segregation could also be compromised. Indeed, genes involved in the formation of microtubules

(*Tubulin Gamma 2*, *TUBG2*) [75], the generation of the mitotic spindle (*Never in Mitosis A related kinase*

9, *NEK9*) [76], the production of cohesion and condensing complexes (*Structural Maintenance Of*

Chromosomes 4, *SMC4*) [77], the control of centriole length, important for centrosomes and

microtubules organization (*Centriolar Coiled-Coil Protein 110*, *CCP110*) [78], the amplification of intra-

spindle microtubules allowing the formation of a sufficient quantity of kinetochore (*HAUS Augmin Like*

Complex Subunit 6, *HAUS6*) [79], the stabilization and protection of the cohesin complex association

with chromatin (*Cell Division Cycle Associated 5*, *CDCA5*; *shugosin 2*, *SGO2*) [80,81] and the production

of microtubule-based motor proteins (*Kinesin Family Member 14* and *23*, *KIF14* and *KIF23*) [82] were

downregulated in OM ICM. Furthermore, *Centromere protein F* (*CENPF*, log2FC = -1.11, FDR < 0.0001),

an important regulator of chromosome segregation during mitosis, was reduced in OM ICM. CENPF is

indispensable for murine embryo development from the first stages of development until at least the

blastocyst stage [83]. Impaired CENPF expression could induce chromosome mis-segregation in the

ICM of old mares' embryos.

Cytokinesis seems to also be altered by maternal age. *CD2-associated protein* (*CD2AP*) and *Vacuolar*

Protein Sorting 4 Homolog A (*VPS4A*) were downregulated in OM ICM (respectively, log2FC = -0.90,

FDR < 0.05 and log2FC = -0.77, FDR < 0.05). Both are involved in the late phase of cytokinesis which

correspond to the abscission of the cell, producing two daughter cells [84,85]. Globally, results suggest

that cell division is impaired or indicate that there is less division in the ICM of old mares' embryos.

Transcription is regulated by epigenetic marks. Several pathways linked to methylation and acetylation

were enriched in YM ICM and TE in relation to numerous upregulated genes in YM ICM (i.e., *Histone*

Deacetylase 1 and *2*, *HDAC1* and *HDAC2*; *Helicase, Lymphoid Specific*, *HELLS*; *DNA Methyltransferase 3*

Alpha, *DNMT3A*; *Lysine Acetyltransferase 6B*, *KAT6B*; *Lysine Demethylase 4A*, *KDM4A*; *Lysine*

Methyltransferase 2A, *KMT2A*; *Nuclear Receptor Binding SET Domain Protein 3*, *NSD3*; *SET Domain*

Containing 5, *SETD5*; *Tet Methylcytosine Dioxygenase 1*, *TET1*). At blastocyst stage, *de novo*

methylation is more important in the inner cell mass than in trophoblast cells and is essential for the

proper development of mammalian embryos [86]. *DNMT3A* is essential for *de novo* methylation and

required for embryo viability [87]. *DNMT3A* expression was reduced in OM ICM (log2FC = -1.01, FDR <

0.05). As it has been shown that maternal aging increases the acetylation of lysine 12 of histone H4 in

mice oocytes, affecting fertilization and embryo development [88], it is likely that the methylation and

acetylation dysregulation observed in the embryos of old mares could be inherited from the oocyte

rather than directly due to embryo environment.

Many of the perturbed gene sets were involved in protein metabolism. In OM ICM, translation was

enriched but protein catabolism is not, whilst in TE, both are enriched. The biogenesis of ribosomes

and protein metabolism regulation are crucial for embryo development. Still in the ICM, proteasome

and mRNA catabolism pathways were also affected. In embryonic stem cells, the proteasome plays a

role in the restriction of permissive transcriptional activity by promoting a dynamic turnover of

transcription factors [89].

Both in OM ICM and TE, several pathways related to mitochondrial function were enriched.

Mitochondria are long known to be the "powerhouse" of cells, producing most of the necessary ATP

for cellular function [90]. ATP production is possible through oxidative phosphorylation enabled by the

electron transport chain in the inner membrane of mitochondria. This process begins with the

oxidation of NADH or FADH that generates a proton gradient across the mitochondrial membrane,

driving ATP synthesis through the phosphorylation of ADP *via* ATPase [91]. Enriched pathways in OM

embryos demonstrate that the NADH dehydrogenase (complex I), the oxidoreductase acting on a

heme group of donors (complex III) with the global electron carrier, and the proton-transporting ATP

synthase (1^{rst} part of the complex V) activities were particularly enriched while the second part of

complex V is depleted. Since all complexes are required to produce ATP, the impaired complex

generation may lead to mitochondrial stress and uncoupled respiratory chain, resulting in poor ATP

production. In human IVF embryos, reduced mitochondrial respiration efficiency, with a negative effect

on embryo development, is observed when oocytes are collected from old women [92]. In mammalian

expanded blastocysts, energy requirement is high, probably because of the need for ion transport

enabling quick blastocoel cavity expansion. Defects in metabolic requirements during blastocyst

development are associated with impaired implantation [93]. Equine embryos are not an exception

[94].

Mitochondria being maternally inherited, uncoupling of the mitochondrial respiratory chain may

originate from mitochondrial defects already present in oocytes [95]. Studies have shown that, in

human, poor oocyte quality is responsible for the decline in fertility associated with maternal aging.

The reduced oocyte developmental competence in older women is related to mitochondrial dysfunction [96]. Regardless of the cell type, it is well known that mitochondrial DNA is more sensitive to mutation than nuclear DNA and that aging is an important risk factor. In mammalian oocytes, mitochondrial DNA mutations increase with age and are inversely correlated with oocyte quality [97–99]. In mice, oocyte mitochondrial DNA content increases during *in vivo* oogenesis without difference according to mice age [100]. In horses' oocytes, before *in vitro* maturation, no difference between young and old mares is observed while after, reduced mitochondrial DNA copy numbers, increased swelling and reduced mitochondrial cristae are reported in oocytes collected from mares >12 years. This suggests that oocyte mitochondria from old mares present deficiencies that become problematic when energy needs are high, as during *in vitro* maturation or subsequent embryo development [101]. Because mitochondrial replication in equine embryos does not begin until the blastocyst stage [102], oocyte mitochondrial numerical or functional deficiencies may compromise early embryogenesis. Alternatively, perturbations in mitochondrial gene expression as observed in old mares' embryos may also result from alterations of the maternal environment. Indeed, mitochondrial function can be improved in mouse embryos by adding antioxidants to the *in vitro* culture medium [103]. To our knowledge, there is no study on oviductal or uterine fluid composition according to maternal age at least in the horse, at the present time.

Finally, ATP production defects could affect embryo blastocoel expansion. Indeed, in TE, the GO biological process "Regulation of action potential" and "Transmission nerve impulse", related to ion movement, were enriched in YM embryos. These processes are associated with genes involved in the action potential dependent ion channels. These ion channels such as NA^+, K^+, ATPase are essential for the accumulation of blastocoel and subsequently yolk sac fluid in equine embryos [104]. Their inhibition is detrimental for embryo development [105]. Thus, both reduced ATP production and disturbed ion channels' function may impact embryo growth. Studies on blastocyst growth and size according to mares' age, however, are controversial [9–15], probably due to the fact that the size of the embryo at the same stage is very variable, as it is the case in this study. To this time, no study on

the effect of mares' age on equine NA^+, K^+, ATPase abundance and function in equine embryos is

available.

Altogether these data indicate that several processes related to chromatin segregation, cytokinesis,

apposition of epigenetic marks and mitochondrial function are affected in older mares' embryos and

that the ICM appears to be more affected than the TE. These perturbations most certainly contribute

to the increased subfertility and early embryo loss observed in aged mares [2,6,7]

Conclusion

This is the first study showing an effect of maternal age on gene expression in the equine blastocyst at

Day 8 post ovulation. Maternal age, even for mares as young as 10 years old, disturbs mitosis,

translation and transcription, cell signaling and adhesion as well as mitochondrial function and cell

commitment in horse embryos. These perturbations on Day 8 post ovulation may affect the further

development of the embryo and as a result, may contribute to decreased fertility due to aging. Even if

the oocyte seems to be in great part responsible for these alterations, maternal environment could

also be responsible. So far, maternal age does not appear to have an impact on foal morphology at

birth but, as for placental function, more subtle effects may exist. Here, as embryos were collected

long before implantation, it is impossible to know if these embryos would have implanted and if they

would have produced a live foal or not. Studies on long-term growth and health of foals born to the

same mares are currently being performed.

List of abbreviations

ARHGAP35: Rho GTPase Activating Protein 35

ARHGAP9: Rho GTPase Activating Protein 9

BRAF: B-Raf Proto-Oncogene, Serine/Threonine Kinase

CCN: cyclin

CCP110: Centriolar Coiled-Coil Protein 110

CD2AP: CD2-associated protein

CD69: CD69 Molecule

CDCA5: Cell Division Cycle Associated 5

CENPF: Centromere protein F

CSNK1A1: Casein Kinase 1 Alpha 1

DEG: differential expressed genes

DeMixT_ICM_cells: deconvoluted gene expression in ICM cells

DeMixT_TE_cells: deconvoluted gene expression in TE cells

DMRT3: Doublesex And Mab-3 Related Transcription Factor 3

DNMT3A: DNA Methyltransferase 3 Alpha

ERR: embryo collection rate

FDR: false discovery rate

FLNC: Filamin C

FYN: FYN Proto-Oncogene, Src Family Tyrosine Kinase

GO BP: Gene Ontology biological process

GO: Gene Ontology

GSEA: gene set enrichment analyses

HAUS6: HAUS Augmin Like Complex Subunit 6

HDAC: Histone Deacetylase

HELLS: Helicase, Lymphoid Specific

ICM: inner cell mass

ICMandTE: inner cell mass enriched hemi-embryo

ICSI: intracytoplasmic sperm injection

ITGAV: Integrin Subunit Alpha V

KAT6B: Lysine Acetyltransferase 6B

KDM4A: Lysine Demethylase 4A

KEGG: Kyoto

KEGG: Kyoto Encyclopedia of Genes and Genomes

KIF: Kinesin Family Member

KMT2A: Lysine Methyltransferase 2A

Log2FC: log2 fold change

MAPK: mitogen-activated protein kinases

NEK9: Never in Mitosis A related kinase 9

NES: normalized enrichment score

NSD3: Nuclear Receptor Binding SET Domain Protein 3

OM: old mares

PAK: P21 activated kinases

PHIP: Pleckstrin Homology domain-Interacting Protein

PLAC8A: placenta Associated 8 A

POU5F1: POU Class 5 Homeobox 1

PPP1R12A: Protein Phosphatase 1 Regulatory Subunit 12A

PPP2R5E: Protein Phosphatase 2 Regulatory Subunit B'Epsilon

ROCK: Rho-associated coiled-coil kinases

RPS29: Ribosomal Protein S29

SETD5: SET Domain Containing 5

SGO2: shugosin 2

SMC4: Structural Maintenance Of Chromosomes 4

SRGAP3: SLIT-ROBO Rho GTPase Activating Protein 3

SRY: sex determining region Y

TE: trophoblast

TE_part: pure trophoblast hemi-embryo

TET1: Tet Methylcytosine Dioxygenase 1

TGF- β: transforming growth factor beta

TGFBR1: transforming growth factor beta receptor 1

TLN2: Talin 2

TUBG2: Tubulin Gamma 2

UTF1: Undifferentiated Embryonic Cell Transcription Factor 1

VPS4A: Vacuolar Protein Sorting 4 Homolog A

Wnt: Wg et Int

XIAP: X-Linked Inhibitor Of Apoptosis

XIST: X inactive Specific Transcript

YM: young mares

Bibliography

765   [1]   Scoggin CF. Not just a number: effect of age on fertility, pregnancy and offspring vigour in

766   thoroughbred brood-mares. Reprod Fertil Dev 2015;27:872. https://doi.org/10.1071/RD14390.

767   [2]   Allen WR, Brown L, Wright M, Wilsher S. Reproductive efficiency of Flatrace and National Hunt

768   Thoroughbred mares and stallions in England. Equine Veterinary Journal 2007;39:438–45.

769   https://doi.org/10.2746/042516407X1737581.

770   [3]   Langlois B, Blouin C. Statistical analysis of some factors affecting the number of horse births in

771   France. Reprod Nutr Dev 2004;44:583–95. https://doi.org/10.1051/rnd:2004055.

772   [4]   Ferreira JC, Canesin HS, Ignácio FS, Rocha NS, Pinto CR, Meira C. Effect of age and endometrial

773   degenerative changes on uterine blood flow during early gestation in mares. Theriogenology

774   2015;84:1123–30. https://doi.org/10.1016/j.theriogenology.2015.06.013.

775   [5]   Ball BA, Little TV, Hillman RB, Woods GL. Pregnancy rates at Days 2 and 14 and estimated

776   embryonic loss rates prior to day 14 in normal and subfertile mares. Theriogenology

777   1986;26:611–9. https://doi.org/10.1016/0093-691X(86)90168-8.

778   [6]   Chevalier-Clément F. Pregnancy loss in the mare. Animal Reproduction Science 1989;20:231–

779   44. https://doi.org/10.1016/0378-4320(89)90088-2.

780   [7]   Souza JRM de, Gonçalves PBD, Bertolin K, Ferreira R, Ribeiro ASS, Ribeiro DB, et al. Age-

781   Dependent Effect of Foal Heat Breeding on Pregnancy and Embryo Mortality Rates in

Thoroughbred Mares. Journal of Equine Veterinary Science 2020;90:102982.

https://doi.org/10.1016/j.jevs.2020.102982.

[8]   Ball BA, Little TV, Weber JA, Woods GL. Survival of Day-4 embryos from young, normal mares

and aged, subfertile mares after transfer to normal recipient mares. Reproduction

1989;85:187–94. https://doi.org/10.1530/jrf.0.0850187.

[9]   Frank BL, Doddman CD, Stokes JE, Carnevale EM. Association of equine oocyte and cleavage

stage embryo morphology with maternal age and pregnancy after intracytoplasmic sperm

injection. Reprod Fertil Dev 2019;31:1812. https://doi.org/10.1071/RD19250.

[10] Panzani D, Rota A, Marmorini P, Vannozzi I, Camillo F. Retrospective study of factors affecting

multiple ovulations, embryo recovery, quality, and diameter in a commercial equine embryo

transfer program. Theriogenology 2014;82:807–14.

https://doi.org/10.1016/j.theriogenology.2014.06.020.

[11] Carnevale EM, Bergfelt DR, Ginther OJ. Aging effects on follicular activity and concentrations of

FSH, LH, and progesterone in mares. Animal Reproduction Science 1993;31:287–99.

https://doi.org/10.1016/0378-4320(93)90013-H.

[12] Woods GL, Baker CB, Hillman RB, Schlafer DH. Recent studies relating to early embryonic death

in the mare. Equine Veterinary Journal 1985;17:104–7. https://doi.org/10.1111/j.2042-

3306.1985.tb04609.x.

[13] Carnevale EM, Ginther OJ. Defective Oocytes as a Cause of Subfertility in Old Mares. Biology of

Reproduction 1995;52:209–14.

https://doi.org/10.1093/biolreprod/52.monograph_series1.209.

[14] Camargo Ferreira J, Linhares Boakari Y, Sousa Rocha N, Saules Ignácio F, Barbosa da Costa G, de

Meira C. Luteal vascularity and embryo dynamics in mares during early gestation: Effect of age

and endometrial degeneration. Reprod Dom Anim 2019;54:571–9.

https://doi.org/10.1111/rda.13396.

[15] Morel MCGD, Newcombe JR, Swindlehurst JC. The effect of age on multiple ovulation rates,

multiple pregnancy rates and embryonic vesicle diameter in the mare. Theriogenology

2005;63:2482–93. https://doi.org/10.1016/j.theriogenology.2004.09.058.

[16] Hendriks WK, Colleoni S, Galli C, Paris DBBP, Colenbrander B, Roelen BAJ, et al. Maternal age

and in vitro culture affect mitochondrial number and function in equine oocytes and embryos.

Reprod Fertil Dev 2015;27:957. https://doi.org/10.1071/RD14450.

[17] Chavatte-Palmer P, Velazquez MA, Jammes H, Duranthon V. Review: Epigenetics,

developmental programming and nutrition in herbivores. Animal 2018;12:s363–71.

https://doi.org/10.1017/S1751731118001337.

[18] Arnaud MG, Dubroeucq MH, Rivot MD. Notation de l'état corporel des chevaux de selle et de

sport: guide pratique. 1ère. Paris: Institut de l'élevage; 1997.

[19] Duchamp G, Bour B, Combarnous Y, Palmer E. Alternative solutions to hCG induction of

ovulation in the mare. J Reprod Fertil Suppl 1987;35:221–8.

[20] Bucca S, Carli A. Efficacy of human chorionic gonadotropin to induce ovulation in the mare,

when associated with a single dose of dexamethasone administered at breeding time: Efficacy

of human chorionic gonadotropin to induce ovulation when associated with dexamethasone.

Equine Veterinary Journal 2011;43:32–4. https://doi.org/10.1111/j.2042-3306.2011.00488.x.

[21] Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.

EMBnet Journal 2011;17:10–2.

[22] Gutiérrez-Adán A, Perez-Crespo M, Fernandez-Gonzalez R, Ramirez M, Moreira P, Pintado B, et

al. Developmental Consequences of Sexual Dimorphism During Pre-implantation Embryonic

Development. Reprod Domest Anim 2006;41:54–62. https://doi.org/10.1111/j.1439-

0531.2006.00769.x.

[23] Iqbal K, Chitwood JL, Meyers-Brown GA, Roser JF, Ross PJ. RNA-Seq Transcriptome Profiling of

Equine Inner Cell Mass and Trophectoderm. Biology of Reproduction 2014;90.

https://doi.org/10.1095/biolreprod.113.113928.

[24] Beckelmann J, Budik S, Bartel C, Aurich C. Evaluation of Xist expression in preattachment equine

embryos. Theriogenology 2012;78:1429–36.

https://doi.org/10.1016/j.theriogenology.2012.05.026.

[25] Pertea M, Pertea GM, Antonescu CM, Chang T-C, Mendell JT, Salzberg SL. StringTie enables

improved reconstruction of a transcriptome from RNA-seq reads. Nat Biotechnol 2015;33:290–

5. https://doi.org/10.1038/nbt.3122.

[26] Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, et al. Integrative

genomics viewer. Nat Biotechnol 2011;29:24–6. https://doi.org/10.1038/nbt.1754.

[27] Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal

RNA-seq aligner. Bioinformatics 2013;29:15–21.

https://doi.org/10.1093/bioinformatics/bts635.

[28] Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning

sequence reads to genomic features. Bioinformatics 2014;30:923–30.

https://doi.org/10.1093/bioinformatics/btt656.

[29] R Core Team. A Language and Environment for Statistical Computing. Vienna, Austria: R

foundation for Statistical Computing; 2020.

[30] Rstudio Team. RStudio: Integrated Development for R. RStudio. Boston, USA: PBC; 2020.

[31] Pinheiro J, Bates D, Debroy S, Sarkar D, R Core Team. nlme: Linear and Nonlinear Mixed Effects

Models. 2020.

[32] Giraudoux P. pgirmess: Spatial Analysis and Data Mining for Field Ecologists. 2018.

[33] Cao S, Wang JR, Ji S, Yang P, Chen J, Shen JP, et al. Differing total mRNA expression shapes the

molecular and clinical phenotype of cancer. BioRxiv 2020:57.

https://doi.org/10.1101/2020.09.30.306795.

[34] Wang Z, Cao S, Morris JS, Ahn J, Liu R, Tyekucheva S, et al. Transcriptome Deconvolution of

Heterogeneous Tumor Samples with Immune Infiltration. IScience 2018;9:451–60.

https://doi.org/10.1016/j.isci.2018.10.028.

[35]   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq

860   data with DESeq2. Genome Biol 2014;15:550. https://doi.org/10.1186/s13059-014-0550-8.

861   [36]   Gómez-Rubio V. **ggplot2** - Elegant Graphics for Data Analysis (2nd Edition). J Stat Soft 2017;77.

862   https://doi.org/10.18637/jss.v077.b02.

863   [37]   Kassambara A, Mundt F. factoextra: Extract and Visualize the Results of Multivariate Data

864   Analyses. 2020.

865   [38]   Kolberg L, Raudvere U. gprofiler2: Interface to the "g:Profiler" Toolset. 2020.

866   [39]   Yates AD, Achuthan P, Akanni W, Allen J, Allen J, Alvarez-Jarreta J, et al. Ensembl 2020. Nucleic

867   Acids Research 2020;48:D682–8. https://doi.org/10.1093/nar/gkz966.

868   [40]   Mi H, Ebert D, Muruganujan A, Mills C, Albou L-P, Mushayamaha T, et al. PANTHER version 16: a

869   revised family classification, tree-based classification tool, enhancer regions and extensive API.

870   Nucleic Acids Research 2021;49:D394–403. https://doi.org/10.1093/nar/gkaa1106.

871   [41]   Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set

872   enrichment analysis: A knowledge-based approach for interpreting genome-wide expression

873   profiles. Proceedings of the National Academy of Sciences 2005;102:15545–50.

874   https://doi.org/10.1073/pnas.0506580102.

875   [42]   Mootha VK, Lindgren CM, Eriksson K-F, Subramanian A, Sihag S, Lehar J, et al. PGC-1α-

876   responsive genes involved in oxidative phosphorylation are coordinately downregulated in

877   human diabetes. Nat Genet 2003;34:267–73. https://doi.org/10.1038/ng1180.

878   [43]   Liberzon A, Birger C, Thorvaldsdóttir H, Ghandi M, Mesirov JP, Tamayo P. The Molecular

879   Signatures Database Hallmark Gene Set Collection. Cell Systems 2015;1:417–25.

880   https://doi.org/10.1016/j.cels.2015.12.004.

881   [44]   Savage SR, Shi Z, Liao Y, Zhang B. Graph Algorithms for Condensing and Consolidating Gene Set

882   Analysis Results. Molecular & Cellular Proteomics 2019;18:S141–52.

883   https://doi.org/10.1074/mcp.TIR118.001263.

[45]   Shannon P. Cytoscape: A Software Environment for Integrated Models of Biomolecular

Interaction Networks. Genome Research 2003;13:2498–504.

https://doi.org/10.1101/gr.1239303.

[46]   McKinnon AO, Squires EL. Morphologic assessment of the equine embryo. Journal of the

American Veterinary Medical Association 1988;192:401–6.

[47]   Hisey E, Ross PJ, Meyers SA. A Review of OCT4 Functions and Applications to Equine Embryos.

Journal of Equine Veterinary Science 2021;98:103364.

https://doi.org/10.1016/j.jevs.2020.103364.

[48]   Grøndahl ML, Andersen CY, Bogstad J, Nielsen FC, Meinertz H, Borup R. Gene expression

profiles of single human mature oocytes in relation to age. Human Reproduction 2010;25:957–

68. https://doi.org/doi:10.1093/humrep/deq014.

[49]   Reyes JM, Silva E, Chitwood JL, Schoolcraft WB, Krisher RL, Ross PJ. Differing molecular

response of young and advanced maternal age human oocytes to IVM. Human Reproduction

2017;32:2199–208. https://doi.org/10.1093/humrep/dex284.

[50]   Hamatani T, Falco G, Carter MG, Akutsu H, Stagg CA, Sharov AA, et al. Age-associated alteration

of gene expression patterns in mouse oocytes. Human Molecular Genetics 2004;13:2263–78.

https://doi.org/10.1093/hmg/ddh241.

[51]   Campos-Chillon F, Farmerie TA, Bouma GJ, Clay CM, Carnevale EM. Effects of aging on gene

expression and mitochondrial DNA in the equine oocyte and follicle cells. Reprod Fertil Dev

2015;27:925. https://doi.org/10.1071/RD14472.

[52]   Cox L, Vanderwall DK, Parkinson KC, Sweat A, Isom SC. Expression profiles of select genes in

cumulus–oocyte complexes from young and aged mares. Reprod Fertil Dev 2015;27:914.

https://doi.org/10.1071/RD14446.

[53]   Kawai K, Harada T, Ishikawa T, Sugiyama R, Kawamura T, Yoshida A, et al. Parental age and gene

expression profiles in individual human blastocysts. Sci Rep 2018;8:2380.

https://doi.org/10.1038/s41598-018-20614-8.

[54] McCallie BR, Parks JC, Trahan GD, Jones KL, Coate BD, Griffin DK, et al. Compromised global

embryonic transcriptome associated with advanced maternal age. J Assist Reprod Genet

2019;36:915–24. https://doi.org/10.1007/s10815-019-01438-5.

[55] Betteridge KJ, Eaglesome MD, MITCHELLt D, Flood PF, Beriault R. Development of horse

embryos up to twenty two days after ovulation: observations on fresh specimens. Journal of

Anatomy 1982;135:191–209.

[56] Enders AC, Schlafke S, Lantz KC, Liu IKM. Endoderm cells of the equine yolk sac from Day 7 until

formation of the definitive yolk sac placenta. Equine Veterinary Journal 1993;25:3–9.

https://doi.org/10.1111/j.2042-3306.1993.tb04814.x.

[57] Enders AC, Lantz KC, Liu IKM, Schlafke S. Loss of polar trophoblast during differentiation of the

blastocyst of the horse. Journal of Reproduction and Fertility 1988;83:447–60.

https://doi.org/10.1530/jrf.0.0830447.

[58] Hauck CR, Hsia DA, Schlaepfer DD. The Focal Adhesion Kinase--A Regulator of Cell Migration

and Invasion. IUBMB Life 2002;53:115–9. https://doi.org/10.1080/15216540211470.

[59] Provenzano PP, Keely PJ. Mechanical signaling through the cytoskeleton regulates cell

proliferation by coordinated focal adhesion and Rho GTPase signaling. Journal of Cell Science

2011;124:1195–205. https://doi.org/10.1242/jcs.067009.

[60] Zhao X, Guan J-L. Focal adhesion kinase and its signaling pathways in cell migration and

angiogenesis. Advanced Drug Delivery Reviews 2011;63:610–5.

https://doi.org/10.1016/j.addr.2010.11.001.

[61] Hynes, RO. Integrins: bidirectional, allosteric signaling machines. Cells 2002;110:673–87.

https://doi.org/10.1016/S0092-8674(02)00971-6.

[62] Cheuk IW-Y, Siu MT, Ho JC-W, Chen J, Shin VY. ITGAV targeting as a therapeutic approach for

treatment of metastatic breast cancer. American Journal of Cancer Research 2020;10:211–23.

[63] Riento K, Ridley AJ. ROCKs: multifunctional kinases in cell behaviour. Nat Rev Mol Cell Biol

2003;4:446–56. https://doi.org/10.1038/nrm1128.

[64] Brouns MR. p190 RhoGAP is required for neural development. Development 2000;127:4891–

903.

[65] Rane CK, Minden A. P21 activated kinases: Structure, regulation, and functions. Small GTPases

2014;5:e28003. https://doi.org/10.4161/sgtp.28003.

[66] Chan PM, Lim L, Manser E. PAK Is Regulated by PI3K, PIX, CDC42, and PP2Cα and Mediates

Focal Adhesion Turnover in the Hyperosmotic Stress-induced p38 Pathway. J Biol Chem

2008;283:24949–61. https://doi.org/10.1074/jbc.M801728200.

[67] Bacon C, Endris V, Rappold GA. The cellular function of srGAP3 and its role in neuronal

morphogenesis. Mechanisms of Development 2013;130:391–5.

https://doi.org/10.1016/j.mod.2012.10.005.

[68] Wu MY, Hill CS. TGF-β Superfamily Signaling in Embryonic Development and Homeostasis.

Developmental Cell 2009;16:329–43. https://doi.org/10.1016/j.devcel.2009.02.012.

[69] Kemp C, Willems E, Abdo S, Lambiv L, Leyns L. Expression of all Wnt genes and their secreted

antagonists during mouse blastocyst and postimplantation development. Dev Dyn

2005;233:1064–75. https://doi.org/10.1002/dvdy.20408.

[70] Gaivão MMF, Rambags BPB, Stout TAE. Gastrulation and the establishment of the three germ

layers in the early horse conceptus. Theriogenology 2014;82:354–65.

https://doi.org/10.1016/j.theriogenology.2014.04.018.

[71] Smith CA, Hurley TM, McClive PJ, Sinclair AH. Restricted expression of DMRT3 in chicken and

mouse embryos. Mechanisms of Development 2002;119:S73–6.

https://doi.org/10.1016/S0925-4773(03)00094-7.

[72] Andersson LS, Larhammar M, Memic F, Wootz H, Schwochow D, Rubin C-J, et al. Mutations in

DMRT3 affect locomotion in horses and spinal circuit function in mice. Nature 2012;488:642–6.

https://doi.org/10.1038/nature11399.

[73] Zhang Y, Huang L, Fu H, Smith OK, Lin CM, Utani K, et al. A replicator-specific binding protein

essential for site-specific initiation of DNA replication in mammalian cells. Nat Commun

2016;7:11748. https://doi.org/10.1038/ncomms11748.

[74] Yuan K, O'Farrell PH. Cyclin B3 Is a Mitotic Cyclin that Promotes the Metaphase-Anaphase

Transition. Current Biology 2015;25:811–6. https://doi.org/10.1016/j.cub.2015.01.053.

[75] Vinopal S, Černohorská M, Sulimenko V, Sulimenko T, Vosecká V, Flemr M, et al. γ-Tubulin 2

Nucleates Microtubules and Is Downregulated in Mouse Early Embryogenesis. PLoS ONE

2012;7:e29919. https://doi.org/10.1371/journal.pone.0029919.

[76] Fry AM, O'Regan L, Sabir SR, Bayliss R. Cell cycle regulation by the NEK family of protein kinases.

Journal of Cell Science 2012;125:4423–33. https://doi.org/10.1242/jcs.111195.

[77] Losada A, Hirano T. Shaping the metaphase chromosome: coordination of cohesion and

condensation. Bioessays 2001;23:924–35. https://doi.org/10.1002/bies.1133.

[78] Schmidt TI, Kleylein-Sohn J, Westendorf J, Le Clech M, Lavoie SB, Stierhof Y-D, et al. Control of

Centriole Length by CPAP and CP110. Current Biology 2009;19:1005–11.

https://doi.org/10.1016/j.cub.2009.05.016.

[79] Watanabe S, Shioi G, Furuta Y, Goshima G. Intra-spindle Microtubule Assembly Regulates

Clustering of Microtubule-Organizing Centers during Early Mouse Development. Cell Reports

2016;15:54–60. https://doi.org/10.1016/j.celrep.2016.02.087.

[80] Ladurner R, Kreidl E, Ivanov MP, Ekker H, Idarraga-Amado MH, Busslinger GA, et al. Sororin

actively maintains sister chromatid cohesion. EMBO J 2016;35:635–53.

https://doi.org/10.15252/embj.201592532.

[81] Rivera T, Losada A. Shugoshin and PP2A, shared duties at the centromere. Bioessays

2006;28:775–9. https://doi.org/10.1002/bies.20448.

[82] Welburn JPI. The molecular basis for kinesin functional specificity during mitosis. Cytoskeleton

2013;70:476–93. https://doi.org/10.1002/cm.21135.

[83] Zhou C-J, Wang X-Y, Han Z, Wang D-H, Ma Y-Z, Liang C-G. Loss of CENPF leads to developmental

failure in mouse embryos. Cell Cycle 2019;18:2784–99.

https://doi.org/10.1080/15384101.2019.1661173.

[84] Monzo P, Gauthier NC, Keslair F, Loubat A, Field CM, Marchand-Brustel YL, et al. Clues to CD2-

associated Protein Involvement in Cytokinesis□D □V. Molecular Biology of the Cell 2005;16:12.

[85] Babst M, Davies BA. Regulation of Vps4 During MVB Sorting and Cytokinesis. Traffic

2011;12:1298–305.

[86] Hales BF, Grenier L, Lalancette C, Robaire B. Epigenetic programming: From gametes to

blastocyst. Birth Defects Research Part A: Clinical and Molecular Teratology 2011;91:652–65.

https://doi.org/10.1002/bdra.20781.

[87] Marcho C, Cui W, Mager J. Epigenetic dynamics during preimplantation development.

REPRODUCTION 2015;150:R109–20. https://doi.org/10.1530/REP-15-0180.

[88] Suo L, Meng Q-G, Pei Y, Yan C-L, Fu X-W, Bunch TD, et al. Changes in acetylation on lysine 12 of

histone H4 (acH4K12) of murine oocytes during maternal aging may affect fertilization and

subsequent embryo development. Fertility and Sterility 2010;93:945–51.

https://doi.org/10.1016/j.fertnstert.2008.12.128.

[89] Szutorisz H, Georgiou A, Tora L, Dillon N. The Proteasome Restricts Permissive Transcription at

Tissue-Specific Gene Loci in Embryonic Stem Cells. Cell 2006;127:1375–88.

https://doi.org/10.1016/j.cell.2006.10.045.

[90] Dumollard R, Duchen M, Carroll J. The Role of Mitochondrial Function in the Oocyte and

Embryo. The mitochondrion in the germline and early development, vol. 77, Elsevier; 2007, p.

21–49. https://doi.org/10.1016/S0070-2153(06)77002-8.

[91] Balaban RS, Nemoto S, Finkel T. Mitochondria, Oxidants, and Aging. Cell 2005;120:483–95.

https://doi.org/10.1016/j.cell.2005.02.001.

[92]   Wilding M, Dale B, Marino M, di Matteo L, Alviggi C, Pisaturo ML, et al. Mitochondrial

aggregation patterns and activity in human oocytes and preimplantation embryos. Human

Reproduction 2001;16:909–17. https://doi.org/10.1093/humrep/16.5.909.

[93]   Gardner DK, Harvey AJ. Blastocyst metabolism. Reprod Fertil Dev 2015;27:638.

https://doi.org/10.1071/RD14421.

[94]   Lane M, O'Donovan MK, Squires EL, Seidel GE, Gardner DK. Assessment of metabolism of

equine morulae and blastocysts. Mol Reprod Dev 2001;59:33–7.

https://doi.org/10.1002/mrd.1004.

[95]   Hutchison CA, Newbold JE, Potter SS, Edgell MH. Maternal inheritance of mammalian

mitochondrial DNA. Nature 1974;251:536–8. https://doi.org/10.1038/251536a0.

[96]   Krisher RL. Maternal age affects oocyte developmental potential at both ends of the age

spectrum. Reprod Fertil Dev 2019;31:1. https://doi.org/10.1071/RD18340.

[97]   Yang L, Lin X, Tang H, Fan Y, Zeng S, Jia L, et al. Mitochondrial DNA mutation exacerbates female

reproductive aging via impairment of the $NADH/NAD^+$ redox. Aging Cell 2020;19.

https://doi.org/10.1111/acel.13206.

[98]   May-Panloup P, Boucret L, Desquiret-Dumas V, Ferré-L V, Morinière C, Descamps P, et al.

Ovarian ageing: the role of mitochondria in oocytes and follicles. Human Reproduction Update

2016;22:725–43. https://doi.org/10.1093/humupd/dmw028.

[99]   Duran HE, Simsek-Duran F, Oehninger SC, Jones HW, Castora FJ. The association of reproductive

senescence with mitochondrial quantity, function, and DNA integrity in human oocytes at

different stages of maturation. Fertility and Sterility 2011;96:384–8.

https://doi.org/10.1016/j.fertnstert.2011.05.056.

[100] May-Panloup P, Brochard V, Hamel JF, Desquiret-Dumas V, Chupin S, Reynier P, et al. Maternal

ageing impairs mitochondrial DNA kinetics during early embryogenesis in mice. Human

Reproduction 2019;34:1313–24. https://doi.org/10.1093/humrep/dez054.

[101] Rambags BPB, van Boxtel DCJ, Tharasanit T, Lenstra JA, Colenbrander B, Stout TAE. Advancing

maternal age predisposes to mitochondrial damage and loss during maturation of equine

oocytes in vitro. Theriogenology 2014;81:959–65.

https://doi.org/10.1016/j.theriogenology.2014.01.020.

[102] Hendriks WK, Colleoni S, Galli C, Paris DBBP, Colenbrander B, Stout TAE. Mitochondrial DNA

replication is initiated at blastocyst formation in equine embryos. Reprod Fertil Dev

2019;31:570. https://doi.org/10.1071/RD17387.

[103] Silva E, Greene AF, Strauss K, Herrick JR, Schoolcraft WB, Krisher RL. Antioxidant

supplementation during in vitro culture improves mitochondrial function and development of

embryos from aged female mice. Reprod Fertil Dev 2015;27:975.

https://doi.org/10.1071/RD14474.

[104] Waelchli RO, MacPhee DJ, Kidder GM, Betteridge KJ. Evidence for the Presence of Sodium- and

Potassium-Dependent Adenosine Triphosphatase alpha 1 and beta 1 Subunit Isoforms and

Their Probable Role in Blastocyst Expansion in the Preattachment Horse Conceptus. Biology of

Reproduction 1997;57:630–40.

[105] Nascimento AD do, Marques JCC, Cezar ARR, Batista AM, Kastelic JP, Câmara DR. Inhibition of

Na+, K+ -ATPase with ouabain is detrimental to equine blastocysts. Anim Reprod

2020;17:e20190079. https://doi.org/10.21451/1984-3143-AR2019-0079.

1052

Figures

Figure 1: Bisection of equine embryos at 8 days post-ovulation into an ICMandTE and a TE_part.

A) Step 1: Identification of the ICM and the TE; B) Step 2: Definition of the cutting plane to isolate the

ICM in one of the parts; C) Step 3: Cutting of the embryo and separation of the two parts. ICMandTE:

inner cell mass + trophoblast; TE part: pure trophoblast.

NB: Here the trophoblast represents trophectoderm + endoderm and ICM is composed of epiblast

cells.

1060

1061   Figure 2: Total RNA yield per embryo in relation to embryo diameter

1062   Green circles represent embryos from young mares (YM) and yellow squares represent embryos from

1063   old mares (OM). The used relation is exponential.

1064

Figure 3: Gene expression in ICM and TE before and after deconvolution using DeMixT

A) Venn diagrams of the differential gene expression in ICMandTE *vs* TE part (before deconvolution),

DeMixT_ICM_cells *vs* DeMixT_TE_cells (after deconvolution) and DeMixT_ICM_cells *vs* TE_part (gene

expression of ICM after deconvolution *vs* gene expression in TE_part without deconvolution); B)

Principal Component Analysis of gene expression of DeMixT_ICM_cells, DeMixT_TE_cells, ICMandTE

and TE part datasets.

Deconvolution was used to isolate gene expression of ICM and TE cells in ICMandTE hemi-embryos.

ICMandTE: inner cell mass + trophoblast; TE part: pure trophoblast. Here trophoblast represents

trophectoderm + endoderm.

1074

Figure 4: Analysis of differentially expressed genes (DEG) in embryos according to maternal age

A) representation of down (blue) and upregulated (red) DEG in ICM (from DeMixT_ICM_cells data

obtained after deconvolution of ICMandTE using DeMixT R package [33,34]) and TE (from TE_part

dataset) of embryos from OM *vs* YM. Functional classifications (for down regulated genes in ICM and

TE and for up-regulated genes in ICM) using GO Biological Process obtained with PANTHER online

software are presented as pie charts. For upregulated DEG in TE, no functional classification was made.

B) Bar chart presenting p-adjusted of most specific subclass pathways provided by PANTHER online

software after statistical overrepresentation test (Fisher's Exact test and Benjamini-Hochberg

correction) with the Human GO Biological Process annotation on down-regulated DEGs in ICM.

DEG: Differentially Expressed Genes (FDR < 0.05); TE: Trophoblast; ICM: Inner Cell Mass; OM: Old

multiparous mares; YM: young multiparous mares

1086

Figure 5: GSEA clustering of the most perturbed terms in ICM and TE according to mares' age

Nodes represent altered gene sets in ICM and TE (FDR < 0.05). Node size represents the gene set size.

Node shape represents the gene set database: GO BP (circle) or KEGG (diamond). Blue nodes represent

enriched gene sets in YM (NES < 0) while red nodes represent enriched gene sets in OM (NES > 0).

Edges represent the level of connection between representative gene sets. This graph was performed

using SUMER R package [44] and modified using cytoscape 3.8.2 [45]

ICM: Inner Cell Mass; TE: trophoblast; FDR: False Discovery Rate; GO BP: Gene Ontology Biological

Process; Kyoto Encyclopedia of Genes and Genomes; NES: Normalized Enrichment Score; YM: Young

mares; OM: Old mares

1096

1097   Figure 6: Schematic representation of the observed effects of maternal aging on ICM and TE in equine

1098   blastocysts

1099   Normal blastocyst development from day 7 to day 9 is represented inside black square using literature

1100   [56,57]. Light blue boxes represent biological processes that are enriched in YM embryos and light red

1101   boxes represent biological processes that are enriched in OM embryos.

1102

Additional file 1:

Additional file 1.png

Pipeline of biostatistical analysis

Once extracted, total RNA qualification and quantification were obtained using a Bioanalyzer. For

sequencing, 5 ng/sample of total RNA were used for paired end RNA sequencing (I2BC platform) using

Illumina NextSeq 500 high technology. Sequences were trimmed using Cutadapt. To determine XIST

sequence, a *de novo* assembly was made using a mapping with STAR and StringTie. The GTF observed

was aligned in IGV and XIST sequence was found at the known position (according to Pablo Ross

personal communication). XIST sequence was included in EquCab 3.0 and a new mapping was

performed using STAR. Counting was performed using featureCounts. The differential analysis was

performed including embryo sex and diameter using Deseq2. A functional enrichment analysis of DEGs

was performed using PANTHER web software. In another time, counts of all genes were normalized

using Deseq2 and a gene set analysis (GSEA) was performed using the Gene Ontology (GO) biological

process (BP) and the Kyoto Encyclopedia of Genes and Genomes (KEGG) databases with GSEA software

form the Broad Institute. To reduce redundancy between terms, SUMER analysis was performed on

enriched gene sets.

1119

Additional file 2:

Additional file 2.pdf

Analysis of ICM enriched and TE part before gene expression deconvolution using DeMixT

On first page, the analysis of differential expressed genes is represented. The second page represent

the results of the gene set enrichment analysis of ICMandTE gene expression. The third page presents

the clustering of gene sets altered by maternal age from SUMER analysis of the GO BP and KEGG terms

of mixed ICMandTE and TE_part.

1127

Additional file 3:

Additional file 3.csv

Differential gene analysis using DeSeq2 in ICMandTE of equine embryo at Day 8 post-ovulation

Equine ensemble ID, orthologue human Ensembl ID, Orthologue human Entrez Gene ID, gene

description, normalized counts for each sample of ICM and parameters obtained after Deseq2 analysis

(log2FoldChange, pvalue and padj (after FDR correction)) of genes expressed in ICM enriched part

(before gene expression deconvolution using DeMixT) of OM and YM embryos

ICM: Inner cell mass; OM: old mares; YM: young mares

1136

Additional file 4:

Additional file 4.csv

Gene set enrichment analysis results on gene expression comparing ICMandTE of embryos from old

mares to young mares

Gene Set Enrichment Analysis results (pathway name, GO accession when possible and size,

Normalized Enrichment Score, p-value and FDR corrected q-value) for GO biological process and KEGG

databases on ICMandTE gene expression table.

1144

1145

Additional file 5:

Additional file 5.csv

Differential gene analysis using DeSeq2 in DeMixT_ICM_cells of equine embryo at Day 8 post-ovulation

Equine ensemble ID, orthologue human Ensembl ID, Orthologue human Entrez Gene ID, gene

description, normalized counts for each sample of ICM and parameters obtained after Deseq2 analysis

(log2FoldChange, pvalue and padj (after FDR correction)) of genes expressed in ICM (after gene

expression deconvolution of ICMandTE using DeMixT) of OM and YM embryos

ICM: Inner cell mass; OM: old mares; YM: young mares

1154

Additional file 6:

Additional file 6.csv

Differential gene analysis using DeSeq2 in TE_part of equine embryo at Day 8 post-ovulation

Equine ensemble ID, orthologue human Ensembl ID, Orthologue human Entrez Gene ID, gene

description, normalized counts for each sample of ICM and parameters obtained after Deseq2 analysis

(log2FoldChange, pvalue and padj (after FDR correction)) of genes expressed in TE of OM and YM

embryos

TE: Trophoblast; OM: old mares; YM: young mares

1163

Additional file 7:

Additional file 7.csv

Gene set enrichment analysis results on gene expression of DeMixT_ICM_cells of embryos from young

and old mares

Gene Set Enrichment Analysis results (pathway name, GO accession when possible and size,

Normalized Enrichment Score, p-value and FDR corrected q-value) for GO biological process and KEGG

databases on DeMixT_ICM_cells gene expression table (after gene expression deconvolution on

ICMandTE using DeMixT).

1172

Additional file 8:

Additional file 8.csv

Gene set enrichment analysis results on gene expression of TE_part of embryos from young and old

mares

Gene Set Enrichment Analysis results (pathway name, GO accession when possible and size,

Normalized Enrichment Score, p-value and FDR corrected q-value) for GO biological process and KEGG

databases on TE_part gene expression table.

Inner Cell Mass

Trophoblast

MATERNAL AGING

Cell cycle / Mitotic division
→ Assembly of spindle
→ Chromosome segregation
→ Cytokinesis/vesicle trafficking
→ Catabolism of protein and mRNA

Transcription
→ Chromatin modification

Cell signaling & adhesion
→ Wnt signaling pathway
→ Focal adhesion
→ Adherens junction

Embryo development
→ Cell fate commitment

Translation
→ Ribosome biogenesis

Mitochondrial function

Intermembrane space

Mitochondrial matrix

NADH NAD⁺ FADH₂ FAD 2H⁺ + ½O₂ H₂0 ADP + Pᵢ ATP

ATP synthase

↑ Up-regulated in OM   ↓ Down-regulated in OM

ALTERED ICM DEVELOPMENT

Normal **ICM** development

Capsule
Polar TE
Epiblast (ICM)

The polar TE becomes discontinuous → isolated cells are still healthy and dividing and adhere to the epiblast
The epiblast grows → cells are transcriptionally active and ingest polar TE cells by phagocytosis

Day 7   Day 8   Day 9

The mural TE divides following blastocyst growth
The endoderm develops to be continuous and fully surround the blastocoel, with some areas too thin to be seen

Normal **TE** development

Endoderm
Mural TE

ALTERED TE DEVELOPMENT

Ion movement
→ Transmission of action potential

Transcription
→ Chromatin modification

Embryo development

Protein catabolism
→ Proteasome

Mitochondrial function

Translation
→ Ribosome biogenesis

MATERNAL AGING