The quantitative genetics of the prevalence of infectious diseases: hidden genetic variation due to Indirect Genetic Effects dominates heritable variation and response to selection

Piter Bijma[1], Andries D. Hulst[1,2], Mart C. M. de Jong[2]

[1]Animal Breeding and Genomics, Wageningen University and Research, Wageningen, the Netherlands

[2]Quantitative Veterinary Epidemiology, Wageningen University and Research, Wageningen, the Netherlands

Abstract

Infectious diseases have profound effects on life, both in nature and agriculture. Despite the availability of well-established epidemiological theory, however, a quantitative genetic theory of the host population for the endemic prevalence of infectious diseases is almost entirely lacking. While several studies have demonstrated the relevance of the transmission dynamics of infectious diseases for heritable variation and response to selection of the host population, our current theoretical framework of quantitative genetics excludes these dynamics. As a consequence, we do not know which genetic effects of the host population determine the prevalence of an infection, and have no concepts of breeding value and heritable variation for endemic prevalence.

Here we integrate quantitative genetics and epidemiology, and propose a quantitative genetic theory for R_0 and for the endemic prevalence of an infectious disease. We first identify the genetic factors that determine the prevalence of an infection, using an approach founded in epidemiological theory. Subsequently we investigate the population level consequences of individual genetic variation, both for R_0 and for the endemic prevalence. Next, we present expressions for the breeding value and heritable variation, for both prevalence and individual binary disease status, and show that these parameters depend strongly on the level of the prevalence. Results show that heritable variation for endemic prevalence is substantially greater than currently believed, and increases when prevalence approaches zero, while heritability of individual disease status goes to zero. As a consequence, response of prevalence to selection accelerates considerably when prevalence goes down, in contrast to predictions from classical theory. Finally, we show that most of the heritable variation for the endemic prevalence of an infection is hidden due to indirect genetic effects, suggesting a key role for kin-group selection both in the evolutionary history of current populations and for genetic improvement strategies in animals and plants.

Introduction

Pathogens have profound effects on life on earth, both in nature and agriculture, and also directly on the human population (Schrag and Wiener, 1995; Russel, 2013). In nature, infectious pathogens are a major force shaping evolution of populations by natural selection, both in animals and plants (reviewed in Karlsson *et al.* 2014 and Ebert and Fields 2020). In livestock, the annual cost of fighting and controlling epidemic and endemic infectious diseases is substantial, and much greater than the annual value of genetic improvement (Rushton, 2009; Knap and Doeschl-Wilson, 2020). Moreover, while antimicrobials have revolutionized medicine, the rapid appearance of resistant strains has resulted in a global health problem, both in the human population and in livestock (EFSA 2012; Thanner *et al.* 2016). Thus there is an urgent need for additional methods and tools to combat infectious diseases. For livestock and plant production, artificial genetic selection of (host) populations for infectious disease traits may provide such a tool. To quantify and optimize the potential benefits of such selection, however, we need to understand the quantitative genetics of infectious disease traits.

Despite the availability of well-established epidemiological theory (*e.g.*, Diekmann *et al.* 2012), a quantitative genetic theory of the host population for the prevalence of infectious diseases is almost entirely lacking. Current approaches for genetic selection against infectious diseases in livestock and crops are entirely based on the individual host response, ignoring transmission of the infection in the population. While several studies have demonstrated the relevance of the transmission dynamics of infectious diseases for heritable variation and response to selection in the host population (Lipschutz-Powell *et al.*, 2012; Anche *et al.*, 2014; Tsairidou *et al.*, 2019; Hulst *et al.*, 2021), mostly using stochastic simulations, the current theoretical framework of quantitative genetics does not include these dynamics. Moreover, we lack general expressions for the genetic variance in key epidemiological parameters, in particular the basic reproduction number R_0, even though such parameters may have a genetic basis.

Infections for which recovery does not confer any long-lasting immunity typically show endemic behaviour, where the infection remains present in the population. For such infections, the endemic prevalence is defined as the expected fraction of the population that is infected. Because we lack a theoretical quantitative genetic framework for infectious diseases, we do not know which genetic effects of the host population determine the prevalence of an infectious disease, and have no concepts of breeding value and heritable variation for endemic prevalence. Hence, we do not understand response to genetic selection in the endemic prevalence of infectious diseases at present. The main parameter determining the prevalence of endemic infections is the basic reproduction number R_0, defined as the average number of individuals that gets infected by a typical infected individual in an otherwise non-infected population. In this manuscript we will propose a quantitative genetic framework for heritable variation and response to selection for R_0 and for the endemic prevalence of infectious diseases.

Individual phenotypes for infectious diseases are often recorded as the binary infection status of an individual, zero indicating non-infected and one indicating infected. The prevalence of an infection is then defined as the fraction of individuals that is infected, which is the fraction of individuals that has infection status $y = 1$. Because the average value of individual binary infection status is equal to the fraction of individuals infected, response to genetic selection in binary infection status is identical to response in prevalence, and *vice versa*. Binary infection status (0/1) typically shows low heritability, which suggests that response to selection is limited, also for prevalence (Bishop and Woolliams 2010; Bishop *et al.*, 2012; Martin *et al.*, 2018).

Geneticists have long realized that the categorical distribution of binary traits does not agree well with quantitative genetic models for polygenic traits, such as the infinitesimal model (Fisher 1918). For this reason, models have been developed that link an underlying normally-distributed trait to the observed binary phenotype, such as the threshold model (Dempster & Lerner, 1950; Gianola, 1982) and the equivalent generalized linear mixed model with a probit link function (*e.g.,* de Villemereuil *et al.* 2016). In such models, the underlying scale is interpreted as causal, and genetic parameters are assumed to represent "biological constants" on this scale. The genetic parameters on the observed scale, in contrast, depend on the mean of the trait, and thus change with the mean even when the change in allele frequencies at causal loci is infinitesimally small. In a landmark paper, Robertson (1950) showed that the observed-scale heritability of binary traits reaches a maximum at a prevalence of 0.5, and approaches zero when the prevalence is close to 0 or 1. Hence, observed-scale heritability vanishes when artificial selection moves prevalence close to zero, hampering further genetic change.

Infectious disease status, however, differs fundamentally from binary phenotypes for non-communicable traits, such as, say, heart failure. Because pathogens can be transmitted between host individuals, either directly or via the environment, the infection status of an individual depends on the status of other individuals in the population. This suggests that Indirect Genetic Effects (IGE) may play a role, which would fundamentally alter heritable variation and response to selection (Griffing 1967; Moore *et al.*, 1997; Wolf *et al.*, 1998; Bijma and Wade 2008; Bijma 2011). Results of simulation studies indeed suggest that selection response in the prevalence of infectious diseases may differ qualitatively from response in non-communicable traits (Nieuwhof *et al.*, 2009; Doeschl-Wilson *et al.*, 2011; Anche *et al.*, 2014; Hulst *et al.* 2021), and this has also been observed in an actual population (Heringstad *et al.* 2007). Results of Hulst *et al.* (2021), for example, show that genetic selection may result in the eradication of an infection via the mechanism of herd immunity, just like with vaccination (FINE 1993). This result contradicts predictions based on the observed-scale heritability for non-communicable binary traits, where heritability vanishes when prevalence approaches zero (Robertson, 1950).

While quantitative geneticists and breeders typically focus on individual disease status and (implicitly) interpret prevalence as an average of individual trait values, epidemiologists interpret the endemic prevalence of an infectious disease as the result of a population level process of transmission of the infection (Kermack and McKendrick, 1927; Keeling and Rohani, 2011; Diekmann *et al.* 2012). In the latter perspective, both R_0 and the prevalence are emergent properties of a population, similar to the size of a termite colony or the number of prey caught by a hunting pack, rather than an average of individual trait values. Because such emergent traits do not belong to single individuals, we cannot apply the common partitioning of individual phenotypic values into individual additive genetic values (breeding values) and non-heritable residuals ("environment"). Nevertheless, the genetic effects that determine the response to selection in an emergent trait and the heritable variation for an emergent trait can be defined based on the so-called total heritable variation (Bijma 2011). The total heritable variation in a trait is based on the individual genetic effects on the level of the emergent trait, rather than on a decomposition of individual trait values into genetic and residual effects. This suggests we can develop a quantitative genetic theory for the endemic prevalence of infectious diseases by combining epidemiological theory with the theory of total heritable variation.

Here we propose a quantitative genetic theory for the basic reproduction number R_0 and for the endemic prevalence of infectious diseases. We first identify the genetic factors that determine the prevalence of an infectious disease. Similar to the threshold model, we will assume an underlying additive infinitesimal model for those genetic factors. However, the link between the underlying additive

scale and the observed endemic prevalence will be founded in epidemiological theory, with a key role
for R_0. Subsequently we investigate the population level consequences of genetic variation in individual
disease traits for R_0 and for the endemic prevalence. Next, we move to the individual level, and derive
expressions for the breeding value and heritable variation, for R_0, endemic prevalence and individual
binary infection status, and show how these parameters depend on the level of the endemic prevalence.
Results will show that heritable variation for endemic prevalence increases when prevalence approaches
zero, while heritability of individual infection status goes to zero. Then we investigate response to
selection, and show that response of prevalence to selection accelerates considerably when prevalence
goes down. Finally, we partition the breeding value for prevalence into direct and indirect genetic
effects, and show that most of the heritable variation in the endemic prevalence of the infection is
indirect, and thus hidden to classical genetic analysis and selection. We focus solely on the development
of quantitative genetic theory, and do not consider the statistical estimation of the genetic effects
underlying prevalence. Such methods have been developed elsewhere (Anacleto *et al.* 2015; Biemans
et al. 2017; Pooley *et al.* 2020).

Theory and Results

1. The genetic factors that determine R_0 and the endemic prevalence

We consider an endemic infectious disease, where individuals can either be susceptible (*i.e.*, in the non-
infected state), denoted by S, or in the infected state, denoted by I. We use corresponding symbols in
italics to denote the number of individuals with that status. Thus, with a total of *N* individuals in the
population in which the endemic takes place, *S* denotes the number of susceptible individuals, *I* the
number of infected individuals, and $S + I = N$ (see Table 1 for a notation key). We will assume that
infected individuals are also infectious, and can thus infect others. When individuals recover they
become susceptible again. This model is known as the SIS compartmental model (Hethcote, 1989), and
was first discussed by Weiss and Dishon (1971; In the Discussion we will consider the validity of our
results for other compartmental models).

The prevalence (*P*) of an endemic infection is defined as the fraction of the population infected
(Diekmann *et al.* 2012),

$$P = \frac{I}{N} \tag{1}$$

151 When individual infection status is coded in a binary fashion, using $y = 0$ for non-infected individuals
152 and $y = 1$ for infected individuals, the prevalence is also equal to the average individual infection status
153 in the population,

$$P = \bar{y}. \tag{2}$$

The prevalence of an infectious disease is determined by R_0. The R_0 is defined as the average
number of individuals that get infected by a typical (*i.e.*, average) infected individual in an otherwise
non-infected population, and is a property of the population (Kermack and McKendrick, 1927; Anderson
and May, 1979; Diekmann *et al.* 1990). When $R_0 > 1$, an average infected individual on average infects
more than one new individual in an infection free population, and the infection can persist in the
population.

The prevalence of an endemic infection reaches an equilibrium value, known as the endemic
prevalence, when a single typical infected individual on average infects one other individual ($R = 1$; the
endemic steady state). In a population where all individuals are the same, *i.e.,* in the absence of genetic
heterogeneity among host individuals, this occurs when the product of R_0 and the fraction of contact
individuals that is susceptible is equal to one; $R_0 (1 - P) = 1$. For example, when $R_0 = 3$, an infected
individual could in principle infect three other individuals. However, when only one third of its contact
individuals is susceptible (*i.e.*, not infected), meaning $1–P = 1/3$, then the effective reproduction number
equals $3 \times 1/3 = 1$. Hence, when $1–P = 1/3$, an infected individual is on average replaced by a single
newly infected individual, so that an equilibrium occurs at $P = 1 - 1/3 = 2/3$. In the absence of
heterogeneity, therefore, the endemic prevalence is given by (Weiss and Dishon 1971).

$$P = 1 - 1/R_0. \tag{3}$$

Throughout, we will use the symbol P to denote the endemic prevalence. The actual prevalence tends
to fluctuate around the equilibrium value because of random perturbations and transient effects, for
example when new animals replace some of the resident animals. Equation 3 is an approximation when
there is variation among individuals, which is commonly referred to as "heterogeneity" in the
epidemiological literature, and which will be addressed in section 4 below.

Figure 1 illustrates the relationship between the endemic prevalence and R_0. When R_0 is smaller
than one the endemic prevalence is zero (the infection is not present in the long run), and Equation 3
does not apply. For large R_0 the endemic prevalence asymptotes to 1. This threshold phenomenon, *i.e.*,
$P = 0$ when $R_0 < 1$ and $P > 0$ when $R_0 \geq 1$, is exact also with heterogeneity (Diekmann *et al*., 1990).
Note that the curve is steeper the closer R_0 is to 1. This pattern will have considerable consequences for
the relationship between the heritable variation in the endemic prevalence and the level of the endemic
prevalence, as will be shown in section 6 of this manuscript.

Figure 1 – The relationship between the endemic prevalence (P) and the basic reproduction number (R_0) for a
homogeneous population. From Equation 3.

Because the endemic prevalence is determined by R_0 (Equation 3), the response of prevalence
to selection, *i.e.,* the genetic change in the endemic prevalence from one (host) generation to the next,
follows from the genetic change in R_0. Thus, to measure the value of an individual with respect to
response to selection, we should base this measure on the genetic impact of the individual on R_0. In other
words, the definition of an individual breeding value for endemic prevalence should be based on R_0.
The next step, therefore, is to find the individual genetic factors underlying R_0.

In the absence of variation among individuals (heterogeneity), R_0 is the product of the
transmission rate parameter (β) and the mean duration of the infectious period ($1/\alpha$; Kermack and
McKendrick, 1927; Diekmann *et al*., 1990),

$$R_0 = \beta/\alpha. \tag{4}$$

where α is the recovery rate parameter. The β is the average number of individuals infected per unit of
time by a single infected individual when all its contact individuals are susceptible, and α is the
probability per unit of time for an infected individual to recover. With heterogeneity, Equation 4 is an
approximation (Diekmann *et al*., 1990); the effect of heterogeneity on R_0 will be addressed in section 3
of this manuscript.

With heterogeneity, the transmission rate parameter may vary between pairings (contacts) of
individuals. The transmission rate parameter between infectious individual j and susceptible individual
i may be modelled as the product of an overall effective contact rate (c), the susceptibility (γ) of recipient
individual i and the infectivity (φ) of donor individual j (*e.g.,* De Jong *et al.*, 1996; Lipschutz-Powell *et*
al., 2014; Anacleto *et al.* 2015; Biemans *et al.* 2017),

$$\beta_{ij} = c\gamma_i\varphi_j \tag{5}$$

Hence, β_{ij} refers to transmission from individual j to i, and may differ from β_{ji}. Note that i and j are
distinct individuals in Equation 5, so that γ_i and φ_j are independent when i and j are genetically
unrelated.

Equations 3 through 5 show that the factors underlying the endemic prevalence of an infection are
the contact rate c, the susceptibility, γ, the infectivity, φ, and the recovery rate α. We define c as a fixed
parameter for the population (or, for example, for a sex, herd or age class combination), whereas γ, φ
and α may show random variation among individuals. Note, while the actual contact rate may vary
among individuals, it is convenient to include such variation in the individual susceptibility and
infectivity traits. Thus we also assume that all the individuals are mixing randomly within the (local)
population, such as a herd. In principle, β_{ij} might also depend on the specific combination of i and j, so
that we cannot separate β_{ij} into a product of components due to i an j. However, in a quantitative genetic
perspective, such an interaction represents between-individual epistasis, which does not contribute to
the heritable variation, and which we will therefore ignore (In epidemiological terminology, we assume
separable mixing; Diekmann *et al*. 1990). Hence, conceptually we define c as the average effective
contact rate for the population, while variation in contact rate among individuals is included in γ and φ.
Moreover, to define the scale of Equations 4 and 5, it is convenient to include the scale in c, and to
express γ, φ and α relative to a value of 1. Hence, with this parameterisation, the c is on the scale of R_0,

and R_0 and c are identical in the absence of heterogeneity. With heterogeneity, however, R_0 may deviate
from c (See section 3 below).

2. Genetic models for susceptibility, infectivity, recovery and R_0

Genetic variation is potentially present in susceptibility, infectivity and the recovery rate. In this section
we propose a genetic model for these traits, which subsequently leads to a genetic model for R_0.

We assume that susceptibility, infectivity and the recovery rate are affected by a large number
of loci, each of small effect, so that genetic effects approximately follow a Normal distribution.
However, as γ, φ, and α represent rates, *i.e.,* probabilities per unit of time, their values are strictly
positive. Moreover, in the expressions for the epidemiological parameters of the previous section
(Equations 4 and 5), all these parameters appear in products. For these reasons, following Anacleto *et*
al. (2015), we define normally distributed additive genetic effects for the logarithm of these rates, so
that effects are multiplicative on the actual scale, and the rates themselves follow a log-normal
distribution,

$$\gamma_i = e^{A_{l\gamma,i}} \tag{6a}$$

$$\varphi_i = e^{A_{l\varphi,i}} \tag{6b}$$

$$\alpha_i = e^{A_{l\alpha,i}} \tag{6c}$$

243 where $A_{l..,i}$ denotes the Normally distributed additive genetic value (breeding value) for the logarithm
244 of the corresponding rate for individual i, and has a mean of zero,

$$\begin{bmatrix} A_{l\gamma} \\ A_{l\varphi} \\ A_{l\alpha} \end{bmatrix} \sim N \left[\begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} \sigma^2_{A_{l\gamma}} & \sigma_{A_{l\gamma}A_{l\varphi}} & \sigma_{A_{l\gamma}A_{l\alpha}} \\ \sigma_{A_{l\gamma}A_{l\varphi}} & \sigma^2_{A_{l\varphi}} & \sigma_{A_{l\varphi}A_{l\alpha}} \\ \sigma_{A_{l\gamma}A_{l\alpha}} & \sigma_{A_{l\varphi}A_{l\alpha}} & \sigma^2_{A_{l\alpha}} \end{pmatrix} \right]. \tag{7}$$

Throughout, we use subscript l to denote the natural logarithm. Thus, the breeding values for $\log(\gamma)$,
$\log(\varphi)$ and $\log(\alpha)$ follow a multivariate normal distribution, as common in quantitative genetics.
Moreover, for the average individual the $A_{l..} = 0$, so that its rates are equal to one ($\gamma = \varphi = \alpha = 1$).
Hence, those rates should be interpreted relative to a value of 1. An individual with $\gamma = 2$, for example,
is twice as susceptible as the average individual.

The breeding values on the log-scale can approximately be interpreted as a relative change of
the corresponding rate. For example, since $e^{0.1} \approx 1.1$, an $A_{l\gamma}$ of 0.1 corresponds approximately to a
10% greater than average susceptibility ($\gamma \approx 1.1$). Similarly, an $A_{l\gamma}$ of -0.1 corresponds approximately
to a 10% smaller than average susceptibility ($\gamma \approx 0.9$). Realistic values for the genetic variances on the
log-scale are probably smaller than $\sim 0.5^2$ (Hulst *et al.*, 2021). For example, with $\sigma^2_{A_{l\gamma}} = 0.5^2$, the 10%
least susceptible individuals have $\bar{\gamma} = e^{-0.88} = 0.42$, while the 10% most susceptible individuals have
$\bar{y} = e^{0.88} = 2.40$. Thus the average susceptibilities of these top and bottom 10% of individuals differ
by a factor of 5.7, which is substantial. Therefore, we will consider additive genetic variances on the
log-scale no greater than 0.5^2. With a prevalence of 0.3, this value corresponds to an observed-scale
heritability of individual binary infection status of about 0.05 (Hulst *et al.*, 2021).

Genotypic value and breeding value for R_0: Based on Equations 4 and 5, we may define an individual
genotypic value for R_0 (see also Anche *et al.* 2014 and Biemans *et al.* 2019),

$$G_{R_0,i} = c\gamma_i\varphi_i/\alpha_i \tag{8}$$

In contrast to the pair-wise transmission rate parameter β_{ij} in Equation 5, an individual's genotypic
value for R_0 is entirely a function of its own rates, as can be seen from the index i on all elements of
Equation 8. This is because $G_{R_0,i}$ refers to the genetic effects that originate from the individual, rather
than to those that affect its trait value. As a consequence these rates may be correlated, as defined in
Equation 7 above. Hence, $G_{R_0,i}$ represents a total genotypic value, including both direct and indirect
genetic effects (Bijma *et al.*, 2007; Bijma 2011). We focus on the total genotypic value, because our
ultimate interest is in response to selection (Section 7). In section 3 of this manuscript, we will show
that R_0 is indeed the simple population average of G_{R_0}.

From Equations 6 and 8,

$$G_{R_0,i} = c\, e^{A_{l\gamma,i}}\, e^{A_{l\varphi,i}}/e^{A_{l\alpha,i}}$$

$$= e^{\ln(c) + A_{l\gamma,i} + A_{l\varphi,i} - A_{l\alpha,i}}$$

$$= e^{\ln(c) + A_{lR_0,i}}, \tag{9}$$

276 where $A_{lR_0,i}$ is a Normally distributed additive genetic effect (breeding value) for the logarithm of R_0,

$$A_{lR_0,i} = A_{l\gamma,i} + A_{l\varphi,i} - A_{l\alpha,i} \tag{10a}$$

$$A_{lR_0} \sim N(0, \sigma^2_{A_{lR_0}}) \tag{10b}$$

$$\sigma^2_{A_{lR_0}} = \sigma^2_{A_{l\gamma}} + 2\sigma_{A_{l\gamma}A_{l\varphi}} - 2\sigma_{A_{l\gamma}A_{l\alpha}} + \sigma^2_{A_{l\varphi}} - 2\sigma_{A_{l\varphi}A_{l\alpha}} + \sigma^2_{A_{l\alpha}} \tag{10c}$$

280 Hence, our model of the genotypic value for R_0 is additive with Normally distributed effects on the log-
281 scale. Thus the genotypic value for R_0, as defined in Equations 8 and 9, follows a log-normal
282 distribution,

$$G_{R_0} \sim logN\left(\mu = \ln(c), \sigma^2 = \sigma^2_{A_{lR_0}}\right). \tag{11}$$

The genotypic value for R_0 for the average individual, which has $A_{lR_0} = 0$, is equal to the contact rate,
c. Hence, the genotypic value is defined here including its average, it is not expressed as a deviation
from the mean. Moreover, we refer to G_{R_0} as a genotypic value, rather than a breeding value, because
the $e^{A_{lR_0}}$ in Equation 9 is a non-linear function, so that G_{R_0} will show some non-additive genetic
variance, even though A_{lR_0} is additive.

The log-normal distribution of G_{R_0} agrees with the infinitesimal model and the strictly positive
values for R_0 (Anacleto *et al.*, 2015), and is also convenient because the mean and variance of G_{R_0}
follow from the known properties of the log-normal distribution,

$$\mathrm{E}\left(G_{R_0}\right) = c\, e^{\frac{1}{2}\sigma^2_{A_{lR_0}}} \tag{12}$$

$$\text{var}(G_{R_0}) = c^2 (e^{2\sigma^2_{A_{lR_0}}} - e^{\sigma^2_{A_{lR_0}}}) \tag{13}$$

Equations 12 and 10c show that genetic (co)variation in susceptibility, infectivity and/or recovery, and thus in the breeding value for the logarithm of R_0, leads to an increase in the mean genotypic value for R_0. For example, for $\sigma^2_{A_{lR_0}} = 0.5^2$, $\text{E}(G_{R_0}) \approx 1.13c$. While this 13% increase in G_{R_0} may suggest limited impact of heterogeneity, a 13% increase in R_0 has a considerable impact on the endemic prevalence when R_0 is close to one (Figure 1).

Equations 12 and 13 show that a log-normal distribution for susceptibility, infectivity and recovery results in a positive mean-variance relationship for G_{R_0}. Figure 2 illustrates this relationship, for $\sigma^2_{A_{lR_0}} = 0.3^2$ and genetic variation in susceptibility only. The *x*-axis shows the contact rate, which is equal to the genotypic value for R_0 of the average individual. Hence, the *x*-axis reflects the level of R_0. The small circle represents a population with a prevalence of ~0.33, for which observed-scale heritability of binary infection status is ~0.02 (Hulst *et al.*, 2021). For that population, R_0 is ~1.5, and the genetic standard deviation in R_0 is ~0.48. Hence, despite the small observed-scale heritability, R_0 has considerable genetic variation and some individuals will have a genotypic value smaller than 1, which agrees with the findings of Hulst *et al.* (2021). In the context of artificial selection against infectious diseases, the positive mean-variance relationship resulting from our model may be interpreted as conservative, because it implies a reduction of the genetic variance in R_0 with continued selection for lower prevalence.

Figure 2 - Genetic standard deviation in R_0 as a function of the level of R_0 (as approximated by the contact rate c here). From Equation 13, for three values of $\sigma^2_{A_{lR_0}} = \sigma^2_{A_{l\gamma}} = 0.1^2$, 0.3^2 and 0.5^2, and no variation in φ and α. The circle represents a population with a prevalence of ~0.33, for which observed-scale heritability of binary infection status is ~0.02 (Hulst *et al.*, 2021).

In summary, this section has presented a genetic model for susceptibility, infectivity and recovery, leading to expressions for the genotypic value and genetic variance in R_0 (Equations 8, 9 and 11). Note however, that we have not yet provided formal proof that the individual genotypic value for R_0 indeed predicts the actual R_0 (of the population so to say). In fact, the definition of G_{R_0} in Equation 8 is an educated guess based on the expression for R_0 in a homogeneous population (Equation 4). In epidemiology, however, R_0 is an emerging property of a population level process of the transmission of an infection, rather than an average of individual (genotypic) values. Thus it remains to be proven that the G_{R_0} defined in Equation 8 indeed predicts the R_0 of a genetically heterogeneous population. In the next two sections, therefore, we will focus on the population-level consequences of genetic heterogeneity, and investigate the impact of genetic variation on the level of R_0 and on the endemic prevalence.

3. The impact of genetic heterogeneity on R_0

R_0 is a key parameter for infectious diseases, because infections can persist in a population if and only if R_0 is greater than one (Kermack & McKendrik,1927; Diekmann *et al.* 1990). In other words, an endemic equilibrium can exist only when R_0 is greater than one. Conversely, eradication of an infectious disease, either by vaccination or other measures such as genetic selection of the host population, requires that R_0 is reduced to a value smaller than one. Here we address the consequences of genetic (co)variation in susceptibility, infectivity and recovery for the value of R_0, and provide a proof that R_0 is indeed the simple population average of the individual genotypic values for R_0, as defined in Equations 8, 9 and 11. Note that R_0 is strictly defined for the infection free state of the population (*i.e.*, where the infected fraction is infinitesimally small). Hence, in this section we consider the infection free state, while the endemic equilibrium will be addressed in section 4.

Genetic (co)variation in susceptibility, infectivity and recovery has two consequences for R_0. First, it increases the mean genotypic value for R_0. This effect is trivial; it follows directly from Equations 12 and 10c and is not the main focus of this section. Second, as stated above, R_0 is the average number of individuals that gets infected by a *typical* infected individual in an otherwise non-infected (large) population (Kermack and McKendrick, 1927; Diekmann *et al.,* 1990). The expression for R_0 given in Equation 4 ignores the "typical" term in the definition of R_0, and is therefore an approximation in case of heterogeneity (Diekmann *et al*., 1990). The focus of this section is on the consequences of heterogeneity for the properties of the typical infected individual, and thus for R_0.

The properties of the "typical infected individual" will depend on the magnitude and nature of the heterogeneity among the individuals in the population, because the susceptibility and recovery determine which animals are infected, while the infectivity of those individuals may differ from the population average. In contrast to the conclusion of Springbett *et al.* (2003), therefore, genetic heterogeneity can affect R_0 (Diekmann *et al.* 1990, 2012). Suppose, for example, that individuals differ in both susceptibility and infectivity, and that susceptibility is positively correlated to infectivity. Because individuals with greater susceptibility are more likely to become infected, the typical infected individual will have an above-average susceptibility. Moreover, because of the positive correlation with infectivity, this will also translate into an above average infectivity of the typical infected individual, leading to higher R_0. Hence, variation among individuals together with a positive (negative) correlation

between susceptibility and infectivity results in an increase (decrease) in R_0 (Diekmann *et al.* 1990). A similar argument holds for genetic covariation between recovery and infectivity. For this reason, R_0 in general deviates from the right-hand side of Equation 4 obtained using the averages of α, γ and φ.

In Appendix 1, we derive the relationship between R_0 and the genetic parameters for susceptibility, infectivity and recovery. The first step is the derivation of the lifetime infectivity of the typical infected individual. Lifetime infectivity is the total infectivity of an individual, aggregated over its average infectious period, and is the product of its infectivity per unit of time (φ, Equations 5 & 6) and the mean duration of its infectious period, $1/\alpha_i$,

$$\phi_i = \varphi_i/\alpha_i. \tag{14}$$

By introducing life time infectivity, we summarize the infectivity (per unit of time, φ) and the recovery of an individual into a single variable (ϕ), which simplifies the analysis. Appendix 1 shows that the average lifetime infectivity of the typical (*i.e.*, average) infected individual equals

$$\bar{\phi}_{typ} = \bar{\phi} e^{\sigma_{A_{l\gamma}A_{l\phi}}} \tag{15}$$

where $\bar{\phi}$ is the simple average of lifetime infectivity in the entire population, and $\sigma_{A_{l\gamma}A_{l\phi}}$ the covariance between the breeding values for the logarithms of susceptibility and lifetime infectivity. Thus Equation 15 is the average lifetime infectivity of infected individuals in the population at the infection free state, and can be interpreted as an average weighted for differences in susceptibility. It shows that the typical infected individual has an above (below) average life-time infectivity when the covariance between susceptibility and life-time infectivity is positive (negative), as argued verbally in the previous paragraph.

From the definition of R_0 and Equations 4, 5 and 15, it follows that

$$R_0 = c\,\bar{\gamma}\,\bar{\phi}\,e^{\sigma_{A_{l\gamma}A_{l\phi}}} \tag{16}$$

where $\bar{\gamma}$ is the simple population average value of susceptibility. The last term of this expression shows that a positive covariance between susceptibility and life-time infectivity indeed increases R_0. Note that this R_0 is the reproduction number in the large infection free population, as it is normally defined in epidemiology. Hence the distribution of susceptibility in the susceptible individuals remains equal to the population distribution, in contrast to the distribution of infectivity in the infected individuals (See Appendix 1).

Equation 16 can be simplified by substituting the expression for $\bar{\gamma}$ and $\bar{\phi}$, which follow from the log-normal distribution,

$$\bar{\gamma} = e^{\frac{1}{2}\sigma^2_{A_{l\gamma}}} \tag{17a}$$

$$\bar{\phi} = e^{\frac{1}{2}\sigma^2_{A_{l\phi}}} \tag{17b}$$

Substituting Equations 17a and b into Equation 16, and expressing the genetic variance of life-time infectivity in terms of infectivity per unit of time and recovery, reveals that R_0 is equal to the simple population average of the individual genotypic value for R_0 (see Appendix 1 and Equation 12),

$$R_0 = c\, e^{\frac{1}{2}\sigma^2_{A_{lR_0}}} = \mathrm{E}(G_{R_0}) \tag{18}$$

Thus, R_0 depends on the variance in the breeding value for the logarithm of R_0, but is still equal to the mean genotypic value for R_0. In other words, a positive covariance between susceptibility and life-time infectivity indeed increases R_0, but this effect is fully captured by the effect of the variance in the breeding values for the logarithm of R_0 on the mean genotypic value for R_0 (the $e^{\frac{1}{2}\sigma^2_{A_{lR_0}}}$ term in Equation 18). This result, therefore, provides formal proof that the genotypic value for R_0, as defined in Equations $8 - 11$, indeed represents the individual genetic value for R_0.

Note that, while R_0 is equal to the simple average genotypic value for R_0, it still differs from the product of the simple averages of the rates when susceptibility, infectivity and/or recovery are correlated; $R_0 \neq c\,\bar{\gamma}\,\bar{\varphi}\,/\bar{\alpha}$. Moreover, R_0 may also differ from the simple $\bar{\beta}/\bar{\alpha}$ with heterogeneity. A numerical investigation of the $e^{\sigma_{A_{l\gamma}A_{l\phi}}}$ term in Equation 16 shows that a correlation between susceptibility and life time infectivity may change R_0 by a maximum of about 25% for realistic levels of heterogeneity and log-normally distributed genetic effects. For example, for $\sigma^2_{A_{l\gamma}} = \sigma^2_{A_{l\phi}} = 0.5^2$, and a correlation $r_{A_\gamma A_\phi} = 0.8$, R_0 is 22% greater than $c\,\bar{\gamma}\,\bar{\varphi}\,/\bar{\alpha}$. For values of R_0 close to 1, this 22% may be the difference between absence of an infection *vs.* a significant endemic prevalence. Thus a correlation between susceptibility, infectivity and/or recovery may have a meaningful impact on R_0.

In summary, this section has shown that heterogeneity and a positive correlation between susceptibly and life-time infectivity lead to an increase of R_0, and thus increase the probability that an infectious disease persists in the population. However, when genotypic values for R_0 follow a log-normal distribution, R_0 is still equal to the simple average of those genotypic values.

4. The impact of genetic variation on the endemic prevalence

In this section we present an expression for the endemic prevalence in a population with genetic variation in susceptibility, infectivity and recovery, and also briefly investigate the quantitative effect of such variation for the endemic prevalence. Figure 1 and Equation 3 show the relationship between R_0 and the endemic prevalence for a homogeneous population. With variation among individuals, however, more susceptible individuals are more likely to be in the infected state in the endemic equilibrium. For this reason, the mean susceptibility of the remaining non-infected individuals will be lower than the population average susceptibility. This in turn translates into an endemic prevalence lower than expected based on R_0 (Equation 3; Diekmann *et al.* 2012; Springbett *et al.*, 2003; Note, however, that the threshold value of $R_0 = 1$ remains, so that endemic prevalence is zero if and only if $R_0 \leq 1$, and in that sense the R_0 given in Equation 18 is exact). Similar arguments can be used to show that prevalence depends on the variation in the recovery rate, and on the covariation of infectivity, susceptibility and recovery. Thus Equation 3 is exact only in the absence of heterogeneity in these parameters.

The endemic prevalence in a heterogeneous population can be found by realizing that the prevalence must have reached an equilibrium value for each type of individual (Biemans *et al.*, 2017; Aznar *et al.*, 2018). Suppose, for example, that susceptibility, infectivity, and recovery would be governed by the same single bi-allelic locus in a diploid organism. Then, for the entire population to be

in equilibrium, each of the three genotypic classes should be in equilibrium as well. In other words, the
prevalence should have reached an equilibrium value within each genotypic class, but this value may
differ among the three classes. Here we adapt this approach to continuous variation in polygenic traits.

In the endemic equilibrium, the number of susceptible individuals of each type, say i, should
not change over time (apart from random fluctuation). Thus, for each type i, the number of newly
infected susceptibles should be equal to the number of recovering infecteds,

$$c\gamma_i\bar{\varphi}_{\text{inf}}\,S_i(t)\frac{I}{N} = \alpha_i I_i(t) \tag{19}$$

where S_i is the number of susceptible individuals of type i, t denotes time, c the contact rate, γ_i the
susceptibility of type i, $\bar{\varphi}_{\text{inf}}$ the mean infectivity of the infected individuals in the endemic equilibrium,
I the total number of infected individuals in the endemic equilibrium, N total population size, α_i the
recovery rate for type i, and I_i the number of infected individuals of type i. The left hand side in Equation
19 represents the decrease in the number of susceptibles due to transmission (infection), while the right
hand side represents the increase of the number of susceptibles due to recovery of infected individuals.
Our interest is in the solution of Equation 19 for I_i (or equivalently, for $S_i = N_i - I_i$). Above, we used i to
index individuals. Here we use i also to index types, since each individual will be genetically unique for
polygenic traits, so that a type corresponds to an individual. Moreover, we treat S_i and I_i as non-integer
because our interest is in their expectation. Note that the mean infectivity of the infected individuals in
the endemic equilibrium ($\bar{\varphi}_{\text{inf}}$) will differ from the simple population average of infectivity ($\bar{\varphi}$) when
infectivity is correlated to susceptibility and/or recovery.

Equation 19 can be solved for the endemic prevalence in type i, $P_i = I_i/N_i$, N_i denoting the total
number of individuals of type i in the population,

$$P_i = \frac{\mathcal{R}_{0,i}P}{\mathcal{R}_{0,i}P+1} \tag{20a}$$

453 where P denotes the overall endemic prevalence in the population (Equation 1), and

$$\mathcal{R}_{0,i} = \frac{c\gamma_i\bar{\varphi}_{\text{inf}}}{\alpha_i} \tag{20b}$$

Equations 20a&b make no assumptions on the distribution of γ, φ and α, and are thus not restricted to
log-normal distributions. Although Equation 20b is similar to Equation 8, note that $\mathcal{R}_{0,i}$ differs from the
genotypic value for R_0 ($G_{R_0,i}$; We use a symbol slightly different from R to highlight this difference).
The $\mathcal{R}_{0,i}$ is a function of the mean infectivity of the infected individuals in the endemic equilibrium
($\bar{\varphi}_{\text{inf}}$), while $G_{R_0,i}$ is a function of the infectivity of the individual itself (φ_i). Our interest here is in the
prevalence for an individual with susceptibility γ_i and recovery rate α_i in the endemic equilibrium,
where i is exposed to the mean infectivity of the infected individuals. Hence the $\bar{\varphi}_{\text{inf}}$ term in $\mathcal{R}_{0,i}$. The
$G_{R_0,i}$, in contrast, defines the contribution of an individual's genes to R_0 (Equation 18), which is relevant
for response to selection (section 7 below). Note that $\bar{\varphi}_{\text{inf}}$ depends on the multivariate distribution of γ,
φ and α.

To find the endemic prevalence, we need to solve Equations 20a and b for P. While we found
an approximate analytical solution for the case without (correlated) genetic variation in infectivity, the
resulting expression is very complex (not shown). We therefore used a numerical solution, which is

easily obtained (see Appendix 2 for methods, and Supplementary Material 1 for an R-code). We validated the numerically obtained solution using full stochastic simulation of actual endemics, following standard methods in epidemiology. Results of these simulations confirmed the numerically obtained solutions (Appendix 3).

The solutions of Equations 20a and b show that variation in susceptibility and/or recovery reduces the endemic prevalence, compared to the simple prediction based on R_0 (Equation 3). Hence, with variation in susceptibility and/or recovery, prevalence is always lower than predicted by Equation 3 (Figure 3; as expected with heterogeneity; Greenhalgh *et al.* 2000). Note that genetic variation in infectivity has no effect on the prevalence (beyond its trivial effect on the mean of G_{R_0}, Equation 18), as long as infectivity is not correlated to susceptibility and/or recovery.

Moreover, it follows from Equation 20 that the effects of genetic variation on the endemic prevalence are identical for susceptibility and recovery, since the $\mathcal{R}_{0,i}$ of an individual depends on the difference between its breeding values for log-susceptibility and log-recovery, $A_{l\gamma,i} - A_{l\alpha,i}$,

$$\mathcal{R}_{0,i} = c\bar{\varphi}_{\text{inf}}\, e^{A_{l\gamma_i} - A_{l\alpha_i}} \tag{21}$$

Hence, with a log-normal distribution of susceptibility and recovery, and in the absence of correlated variation in infectivity, the equilibrium prevalence depends only on (R_0 and) the variance of this difference,

$$var\big(A_{l\gamma} - A_{l\alpha}\big) = \sigma^2_{A_{l\gamma}} - 2\sigma_{A_{l\gamma}A_{l\alpha}} + \sigma^2_{A_{l\alpha}}. \tag{22}$$

Figure 3 illustrates the impact of heterogeneity on the endemic prevalence for a limited number of scenarios with genetic variation in susceptibility only. For $\sigma_{A_{l\gamma}} = 0.1$, the effect of heterogeneity is imperceptible. For $\sigma_{A_{l\gamma}} = 0.3$, the true prevalence is up to 2 percent point lower than the value from Equation 3. For $\sigma_{A_{l\gamma}} = 0.5$, true prevalence is up to 6 percent point lower. This maximum difference occurs at a contact rate of two. Moreover, when $c = 2$ and there is no variation in infectivity, prevalence is always equal to $1 - 1/c = 0.5$, irrespective of the genetic variation in susceptibility and recovery. (This is not visible in Figure 3, because the x-axis shows R_0 rather than c). This occurs because the two opposing effects mentioned at the beginning of this paragraph exactly cancel each other. More detailed results can be found in Supplementary Material 2.

Figure 3 - The impact of heterogeneity on the endemic prevalence. The solid line shows the prevalence predicted from Equation 3. The three other lines show the true prevalence (From numerically solving Equations 20a & b), for three levels of the additive genetic standard deviation in log susceptibility ($\sigma_{A_{l\gamma}}$), and no genetic variation in infectivity or recovery. The line for $\sigma_{A_{l\gamma}} = 0.1$ is almost identical to the solid line. Note that identical results would have been obtained with the same amount of heterogeneity in the recovery rate, or more generally in $A_{l\gamma,i} - A_{l\alpha,i}$, instead of in susceptibility.

5. Genotypic value for individual binary infection status

In the previous two sections, we have considered the population-level effects of genetic heterogeneity. In the next two sections, we move to the individual level. This section focusses on the effects of an individual's genes on its own infection status, while section 6 focusses on the effect of an individual's genes on the prevalence in the population.

By definition, the genotypic value for binary disease status is the expected infection status of an individual given its genotype,

$$G_{y,i} = E(y_i | A_{l\gamma,i}, A_{l\alpha,i}) \tag{23}$$

Thus, the G_y represents the direct genetic effect (DGE) on the own phenotypic value (including the mean, $\bar{y}$, here; Note the distinction between subscript *y*, indicating individual binary infection status,

and γ, indicating susceptibility). The genotypic value of an individual is not a function of its breeding

value for log-infectivity, since an individual's infectivity does not affect its own infection status. Hence,

Equation 23 does not condition on $A_{l\varphi,i}$.

In the previous section, we used Equations 20a and b to investigate the effect of heterogeneity

on the endemic prevalence in the population. Equation 20a shows the expected prevalence of an

individual of type i. However, since prevalence is simply the mean of binary infection status, Equation

20a may also be interpreted as the expected phenotypic value for infection status ($y = 0,1$) of an

individual, given its genotype (specifically, the γ_i and α_i components of $\mathcal{R}_{0,i}$). Hence, Equation 20a also

represents the genotypic value for binary infection status,

522
$$G_{y,i} = \frac{\mathcal{R}_{0,i}P}{\mathcal{R}_{0,i}P+1} \tag{24}$$

where $\mathcal{R}_{0,i}$ follows from Equation 20b. The same result was found by Bijma (2020), but based on a

different approach. Thus the G_y refers to the expected binary infection status of individual i, conditional

on its genotype, in a population with prevalence P. Equations 21 and 24 imply that susceptibility and

recovery are equally important for the infection status of an individual. For example, an individual with

$A_{l\gamma} = -0.1$ has the same expected infection status as an individual with $A_{l\alpha} = +0.1$.

Calculation of G_y from Equation 24 requires knowledge of the endemic prevalence P. In the

previous section we used a numerical approach to find P, because our interest was in the effects of

heterogeneity on P. In applied breeding, however, breeders may often have a reasonable idea of realistic

values for the endemic prevalence, and a numerical solution may not be needed to find G_y, or have little

added value.

Validation: We used stochastic simulation of endemics, following standard methods in epidemiology

(Appendix 3), to validate Equation 24. Figures 4a-c show the mean observed infection status of

individuals as a function of their genotypic value G_y. For all three panels in Figure 4, regression

coefficients were very close to 1, showing that G_y is an unbiased linear predictor of individual infection

status.

Figure 4. Validation of the genotypic value for individual binary infection status. Panels show a scatter plot of the

mean observed infection status of individuals (y-axis) as a function of their genotypic value for infection status

(G_y, x-axis, Equation 24). For genetic variation in susceptibility only, with $\sigma_{A_\gamma}^2 = 0.3^2$, $N = 2000$ individuals, and

a total of 300,000 events (sum of recoveries and infections). Panel A: P = 0.2; $\hat{b}_{y,G_\gamma} = 0.998$. Panel B: P = 0.5,

$\hat{b}_{y,G_\gamma} = 1.006$. Panel C: P = 0.8, $\hat{b}_{y,G_\gamma} = 1.005$.

We numerically investigated the relative amount of non-additive genetic variance in G_y. Results
(not shown) revealed only little non-additive genetic variance. For example, for $P = 0.2$ and $\sigma^2_{A_{G_y}} = 0.5^2$,
more than 96% of the genotypic variance in y was additive. Thus the breeding value for own infection
status is very similar to the genotypic value,

548
$$A_y \approx G_y - P, \tag{25}$$

where the "$-P$" term simply reflects subtraction of the average, $\overline{G_y} = P$, so that the mean breeding value
is zero by definition. We defer further investigation of the breeding value and the additive genetic
variance for individual infection status to the next section, to facilitate comparison with the
corresponding measures for prevalence.

6.   Breeding value and heritable variation for the endemic prevalence

The previous section focussed on the genetic effects of individuals on their own infection status. In this
section we will consider the genetic effects of individuals on the endemic prevalence in the population.
In other words, the previous section focussed on the contribution of genetic effects to the variation in
infection status among individuals, while this section considers the genetic effects that are relevant for
response to selection. We will present expressions for the genotypic value, breeding value and additive
genetic variance for the endemic prevalence. The genotypic value will reflect the full genetic effect of
an individual on the endemic prevalence in the population, while the breeding value reflects the additive
component thereof. The last part of this section contains a comparison of the breeding value for endemic
prevalence and that for individual infection status.

The relationship between R_0 and the endemic prevalence (Equation 3) suggests we can translate
the individual genotypic value for R_0 (Equations 8 and 9) to the scale of prevalence, by defining an
individual genotypic value for prevalence as

567
$$G_{P,i} = 1 - \frac{1}{G_{R_0,i}} \tag{26a}$$

568   Because this definition is based on Equation 3, it ignores the of effect heterogeneity on the relationship
569   between P and R_0 (Figure 3). We will investigate the relevance of this approximation numerically in
570   section 7, which focusses on response to selection. Substituting Equation 9 into 26a yields an expression
571   for the genotypic value of an individual for the endemic prevalence, in terms of its breeding value for
572   the logarithm of R_0,

573
$$G_{P,i} = 1 - e^{-\ln(c) - A_{lR_0,i}} \tag{26b}$$

574   Because the term in the exponent is Normally distributed, $1 - G_P$ follows a log-normal distribution,

575
$$(1 - G_P) \sim logN\left[\mu = -\ln(c), \sigma^2 = \sigma^2_{A_{lR_0}}\right] \tag{27}$$

576   The mean and variance of the genotypic values for prevalence, therefore, follow from the properties of
577   the log-normal distribution,

578
$$\mathrm{E}(G_P) = 1 - c^{-1} e^{\frac{1}{2}\sigma^2_{A_{lR_0}}} \tag{28}$$

$$\text{var}(G_P) = c^{-2}(e^{2\sigma^2_{A_{lR_0}}} - e^{\sigma^2_{A_{lR_0}}}) \tag{29}$$

To enhance the interpretation of Equation 29, we can express it as a function of R_0 or of the endemic prevalence. Substituting Equation 18 into Equation 29 yields an expression for the genetic variance in prevalence as a function of R_0,

$$\text{var}(G_P) = \frac{1}{R_0^2}(e^{3\sigma^2_{A_{lR_0}}} - e^{2\sigma^2_{A_{lR_0}}}) \tag{30a}$$

Given the definition of G_P in Equation 26a, Equation 30a is exact. Next, substituting Equation 3 into Equation 30a yields an expression for $\text{var}(G_P)$ as a function of the endemic prevalence,

$$\text{var}(G_P) \approx (1 - P)^2(e^{3\sigma^2_{A_{lR_0}}} - e^{2\sigma^2_{A_{lR_0}}}) \tag{30b}$$

Equation 30b is approximate, because the relationship between P and R_0 given in Equation 3 is approximate with heterogeneity. Equations 30a and b show how the genotypic variance for endemic prevalence depends on the level of R_0 or equivalently, on the level of the endemic prevalence. Hence, in contrast to ordinary additive genetic traits, the genetic variance for endemic prevalence is a function of the level of the endemic prevalence (Equation 30b).

Figures 5a and b illustrate that the standard deviation in genetic values for endemic prevalence is considerably larger at lower R_0, or equivalently, at lower prevalence. Hence, even though the genetic variance in R_0 decreases with the level of R_0 (Figure 2), the genetic variance for prevalence increases strongly when R_0 decreases. This result originates from the increasing slope of the relationship between prevalence and R_0 when R_0 decreases (Figure 1). In other words, an equal change in R_0 has much greater impact on the endemic prevalence at low R_0 than at high R_0, which is well-known in epidemiology (*e.g.*, Metz, 1978; Bolker and Grenfell, 1996). Hence, for a constant variance in the breeding value for the logarithm of R_0, the genetic variance for endemic prevalence is much greater at lower prevalence. Moreover, genetic selection for lower prevalence will lead to an increase in the genetic variance for prevalence.

Figure 5 – Genetic standard deviation for endemic prevalence as a function of R_0 (panel A), and as a function of the level of the endemic prevalence (panel B). From Equations 30a and b. For $\sigma^2_{A_{lR_0}} = 0.3^2$. In panel A, *x*-axis values below $R_0 = 1$ are omitted, because equilibrium prevalence is zero (the infection is not present) and Equation 30a does not apply.

Figure 6 shows some examples of the distribution of the genotypic value for endemic prevalence, for different values of R_0 and the corresponding endemic prevalence. For the scenarios in Figure 6, the observed-scale heritability of individual infection status does not exceed 0.022 (see Figure 7 below). The panels illustrate that the genotypic standard deviation for endemic prevalence is relatively large, particularly when prevalence is small. For example, for $R_0 = 1.67$ ($P \approx 0.4$; Panel B), the standard deviation in genotypic values for prevalence is around 0.19 (See also Figure 5), and values between ~0 and ~0.7 are quite probable. Hence, despite the low observed-scale heritability of individual infection status, the probable values of G_P span as much as 70% of the full 0-1 range of endemic prevalence.

Figure 6 – Distribution of individual genotypic values for prevalence (G_P), for different values of R_0, or equivalently, different (approximate) values of the endemic prevalence. The distribution is given by $f(G_P) = \frac{1}{(1-G_P)\sigma_{A_{lR_0}}\sqrt{2\pi}} exp\left(-\frac{(\log(1-G_P)+\log(c))^2}{2\sigma_{A_{lR_0}}^2}\right)$, with domain $G_P = (-\infty, 1)$. For $\sigma_{A_{lR_0}}^2 = 0.3^2$. The dashed vertical line shows the mean of G_P. Note that G_P can take negative values while prevalence cannot. This is because G_P reflects the genetic effect of an individual on the prevalence of the population, not the expected value of its own infection status. Thus, negative values for G_P are possible, as long as P is positive. Note that P is very close to the average of the distributions shown.

Breeding value and additive genetic variance for prevalence: The genotypic value for prevalence is not identical to the additive genetic value (*i.e.*, breeding value) for prevalence, because the exponential function in Equation 26b is non-linear, so that G_P contains a non-additive component. Appendix 4 shows that the linear regression coefficient of G_P on A_{lR_0} is equal to $c^{-1} \exp\left(\frac{1}{2}\sigma^2_{A_{lR_0}}\right)$. Substituting $c^{-1} = e^{\frac{1}{2}\sigma^2_{A_{lR_0}}}/R_0$ (from Equation 18), shows that the breeding value for prevalence is given by

$$A_{P,i} = \frac{1}{R_0} e^{\sigma^2_{A_{lR_0}}} A_{lR_0,i} \tag{31a}$$

Given the definition of G_P in Equation 26a, this result is exact. With limited heterogeneity, this result is approximately equal to

$$A_{P,i} \approx \frac{1}{R_0} A_{lR_0,i} \tag{31b}$$

Thus the breeding value for prevalence is proportional to the reciprocal of R_0. The additive genetic variance for prevalence equals

$$\sigma^2_{A_P} = \frac{1}{R_0^2} e^{2\sigma^2_{A_{lR_0}}} \sigma^2_{A_{lR_0}} \tag{32a}$$

and, with limited heterogeneity,

$$\sigma^2_{A_P} \approx \frac{1}{R_0^2} \sigma^2_{A_{lR_0}} \tag{32b}$$

or, expressed as a function of endemic prevalence,

$$\sigma^2_{A_P} \approx (1-P)^2 \sigma^2_{A_{lR_0}} \tag{32c}$$

Equation 32c shows that the additive genetic variance in endemic prevalence increases strongly when prevalence decreases, similar to the relationship between the genotypic variance and endemic prevalence (Figure 5). This result suggests that response of endemic prevalence to selection will be greater at lower levels of the prevalence, which we will further investigate in Section 7 below.

The relative amount of non-additive genetic variance in the endemic prevalence is determined by the magnitude of $\sigma^2_{A_{lR_0}}$ (Appendix 4). For realistic values of $\sigma^2_{A_{lR_0}}$, the vast majority of the genotypic variance in prevalence is additive. For example, for $\sigma^2_{A_{lR_0}} = 0.5^2$, 88% of the variance in G_P is additive. Hence, the distinction between the breeding value for prevalence (A_P) and the genotypic value for prevalence (G_P) seems of minor importance, and results in Figures 5 and 6 will closely resemble those for the additive genetic effects.

Breeding value and heritability for infection status *vs.* prevalence: Appendix 5 shows that, in the absence of genetic variation in infectivity, the breeding value for endemic prevalence is approximately a factor $1/P$ greater than the breeding value for individual infection status,

$$A_{P,i} \approx \frac{1}{P} A_{y,i} \tag{33}$$

Note, in contrast to genotypic values, breeding values are expressed as a deviation from their mean here.
The A_y is the ordinary observed-scale breeding value for binary infection status that breeders are familiar
with.

Equation 33 implies that the impact of an individual's genes on the response of the endemic
prevalence to selection is considerably larger than their impact on the infection status of the individual
itself, particularly when the endemic prevalence is small. Consider, for example, an individual with
$A_{y,i} = -0.02$ in a population with an endemic prevalence of $P = 20\%$. The expected infection status of
this individual in the current population equals $0.20 - 0.02 = 0.18$. Hence, on average, this individual
will be infected 18% of the time. However, its breeding value for prevalence equals $A_{P,i} = -0.02/0.2 =$
-0.10. Hence, if we select individuals with $A_{y,i} = -0.02$ as parents of the next generation, then the
endemic prevalence will go down to $0.20 - 0.10 = 0.10$. In other words, the response to selection will
be fivefold greater than suggested by the ordinary breeding value for individual infection status (since
$1/P = 1/0.2 = 5$). We will numerically validate this theoretical result in Section 7.

The relationship between the breeding value for prevalence and the breeding value for own
infection status shown in Equation 33 suggests a relatively simple expressions for A_y. Such a simple
expression would be convenient, because the alternative is to calculate A_y from Equation 25, which
requires solving Equations 20a and b numerically. On combining Equations 31a and 33, and assuming
limited heterogeneity, so that $e^{\sigma^2_{A_{lR_0}}} \approx 1$ and $1/R_0 \approx (1-P)$, the breeding value for individual
infection status becomes

$$A_{y,i} \approx P(1-P)A_{lR_0,i} \tag{34}$$

We used stochastic simulation to validate this expression and investigate its precision. Results
show that Equation 34 closely matches the regression of individual binary infection status on the
breeding value for the logarithm of R_0 for realistic levels of heterogeneity ($\sigma^2_{A_{lR_0}} \leq 0.5^2$; Supplementary
Material 3; the good fit results from compensating errors due to the approximations). Hence, Equation
34 is sufficiently precise for practical purposes. Note that, since infectivity does not affect the infection
status of an individual itself, a potential component due to infectivity has to be left out of the $A_{lR_0,i}$ term
when calculating Equation 34. In other words, in Equation 34 the $A_{lR_0,i}$ should include only the breeding
values for the logarithm of susceptibility and recovery (see Equation 10a).

It follows from Equation 34 that the additive genetic variance in individual binary infection status
equals

$$\sigma^2_{A_y} \approx P^2(1-P)^2\sigma^2_{A_{lR_0}} \tag{35}$$

688 Next, the observed-scale heritability of binary infection status follows from dividing Equation 35 by the
689 phenotypic variance of binary infection status, $\sigma^2_y = P(1-P)$, giving

$$h^2_y \approx P(1-P)\,\sigma^2_{A_{lR_0}}. \tag{36}$$

Hence, the observed-scale heritability for binary infection status has a maximum at a prevalence of 0.5,
and goes to zero at a prevalence of zero or one, just like the heritability of binary phenotypes for non-

communicable polygenic traits (Robertson, 1950; Figure 7A; assuming the infinitesimal model at the
level of the logarithm of R_0, so that $\sigma^2_{A_{lR_0}}$ is constant).

The ratio of additive genetic variance for prevalence over phenotypic variance in binary infection
status is given by,

$$T^2_P = \frac{\sigma^2_{A_P}}{P(1-P)} \approx \frac{1-P}{P}\sigma^2_{A_{lR_0}} \tag{37}$$

with $\sigma^2_{A_P}$ taken from Equation 32a or c. The T^2_P is an analogy of heritability, but the numerator represents
the additive genetic variance relevant for response to selection in endemic prevalence, rather than for
individual binary infection status. The T^2_P, therefore, reflects the genetic variance that can be used for
response to selection, whereas h^2_y reflects the contribution of additive genetic effects to phenotypic
variance (Bijma *et al.*, 2007; Bijma 2011).

Figures 7 A & B show a comparison of h^2_y and T^2_P for a population without genetic variation in
infectivity, with genetic variances in the logarithm of R_0 ranging from 0.1^2 through 0.5^2. In Figure 7A,
the maximum value of h^2_y equals 0.0625, for $P = 0.5$ and $\sigma^2_{A_{lR_0}} = 0.5^2$. Given that genetic variances
greater than $\sigma^2_{A_{lR_0}} = 0.5^2$ are very large (as argued above), observed-scale heritabilities of binary
infection status greater than ~0.06 are unlikely for endemic infectious diseases. The heritabilities in
Figure 7A agree with the findings of Hulst *et al.* (2021), who used stochastic simulation of actual
endemics and analysis of the resulting binary infection status data with a linear animal model. Figure
7B shows that T^2_P increases strongly when prevalence goes down. Figure 7 illustrates that T^2_P and h^2_y
differ by a factor of approximately P^2, so that the additive genetic variance in prevalence is (much)
greater than the additive genetic variance in individual infection status, and may even exceed the
phenotypic variance at low values of the endemic prevalence (*i.e.*, $T^2_P > 1$).

In conclusion, in this section we have presented expressions for the breeding value for prevalence
(Equation 31) and for individual infection status (Equation 34), and for the corresponding genetic
variances. With realistic levels of heterogeneity, the breeding value for prevalence is a factor $1/P$ greater
than the breeding value for individual infection status. This result suggests that response to selection
should be considerably greater than expected based on ordinary heritability of individual infection status.
We will test this hypothesis in the next section.

Figure 7. Panel A: Observed-scale heritability (h_y^2) of individual binary infection status ($y = 0,1$) as a function of the endemic prevalence, for different additive genetic standard deviations in the logarithm of R_0 (SDAlR0). From Equation 36. Panel B: Ratio of additive genetic variance for prevalence and phenotypic variance in infection status (T_P^2), as a function of the endemic prevalence. From Equations 37 and 32a. In both panels, there is no genetic variation in infectivity.

7. Response to selection

The higher genetic variance for prevalence at lower values of the prevalence (Equations 30 & 32, Figure 5b & 6) suggests that response to selection should increase when the prevalence decreases. To validate and illustrate this hypothesis, we stochastically simulated an endemic infectious disease in a large population undergoing mass selection for individual infection status. Simulations were based on standard methods in epidemiology, not making use of the above theory (Appendix 6).

Figure 8A shows the observed prevalence (*i.e.*, the mean binary infection status in each generation), the mean breeding value for prevalence and the mean breeding value for binary infection status, for ~70 generations of selection. Response in prevalence increases strongly when prevalence decreases, and the infection disappears in the final generation. There is excellent agreement between the observed prevalence and the breeding value for prevalence, showing that the change in $\bar{A}_P$ indeed predicts the change in prevalence. In contrast, the response in prevalence deviates substantially from the response in the breeding value for individual infection status ($\bar{A}_y$), particularly at lower values of the prevalence. Hence, while the breeding value for infection status correctly predicts the average individual infection status within a generation (Figure 4), the change in $\bar{A}_y$ considerably underestimates the response to selection. Furthermore, given the weak selection and the low value of the observe-scale heritability of binary infection status, which did not exceed 0.022 in Figure 8A, response to selection in prevalence is remarkably large, unless prevalence is high. This result agrees with findings of Hulst *et al.* (2021).

Figure 8 – Response to selection in prevalence for 70 generations of mass selection of the host population. For two populations, one starting at a prevalence of ~90% ($c = 10$), the other starting at a prevalence of ~50% ($c = 2$). Each generation, the 50% individuals with the lowest average infection status were selected as parents of the next generation. With genetic variation in susceptibility only, and $\sigma^2_{A_{l\gamma}} = 0.3^2$. For a population of $N = 4{,}000$ individuals, a total of 15,000 events (sum of infections and recoveries) per generation, consisting of a burn-in of 10,000 events and 5,000 recorded events. Hence, selection is based on 1.25 events per individual on average, indicating a limited amount of phenotypic data. Observed-scale heritability for binary infection status in any generation can be read from Figure 7A using an *x*-axis value corresponding to the prevalence in that generation. *Panel A:* Observed prevalence (circles), breeding value for prevalence ($\bar{A}_P$, solid blue line) and breeding value for individual infection status ($\bar{A}_y$, dashed red line). Results for breeding values are the cumulative change in breeding value in each generation plus the initial prevalence. Breeding values were taken from Equation 31a for A_P and from Equation 34 for A_y. *Panel B*: Predicted (lines) versus observed (circles) prevalence. Prevalence was predicted from Equation 38 (blue solid line) or Equation 39 (red dashed line).

We also briefly investigated the prediction of response to mass selection with a very simple expression assuming linearity and with genetic variation in susceptibility only. We based our predictions on breeding values for binary infection status, because these are typically estimated by breeders. Because the breeding value for prevalence and the breeding value for individual infection status differ by a factor $1/P$ when there is no genetic variation in infectivity (Equation 33), we simply upscaled the response to selection predicted for individual infection status from the breeder's equation (Walsh and Lynch, 2018) by this factor, giving

$$R_P = \iota \rho_{A_y,\bar{y}} \sigma_{A_y} \frac{1}{P} \tag{38}$$

where ι is the intensity of selection, defined as the standardized selection differential in mean individual infection status, $\iota = (\bar{y}_{\text{selected}} - \bar{\bar{y}})/\sigma_{\bar{y}}$, $\rho_{A_y,\bar{y}}$ is the accuracy of selection for individual infection status, which is the correlation between the selection criterion ($\bar{y}_i$) and the true breeding value for individual infection status, and P is the prevalence in the generation of the selection candidates. Hence, the numerator of Equation 38 represents the predicted response for individual binary infection status, which is multiplied by a factor $1/P$ to find response in prevalence. To implement Equation 38, we calculated the $\rho_{A_y,\bar{y}}$ as the observed correlation between the true breeding values for binary infection status (A_y;

Eqn. 34) and the selection criterion ($\bar{y}$) in the candidates for selection. Hence, we did not attempt to predict the accuracy of selection.

Figure 8B shows a comparison of observed and predicted prevalence. Above a prevalence of ~0.5, response predicted from Equation 38 is somewhat larger than observed response, while the reverse is true below a prevalence of ~0.5 (Note, response to selection in a generation is reflected by the slope of the figure). Nevertheless, agreement between observed and predicted response is remarkably good given the very unrealistic assumption of linearity in Equation 38 (*i.e.*, bivariate normality of A_y and $\bar{y}$). Because selection was based on mean individual infection status recorded over a period lasting on average only 1.25 events per individual (see legend Figure 8), many values were either 0 or 1, implying strong deviations from normality.

When the prevalence was smaller than 0.5, response to selection was quite large. Hence, there was a meaningful difference in prevalence between parent and offspring generations. Because the P in Equation 38 refers to the prevalence in the parent generation, while response is realized in the offspring generation, Equation 38 resulted in underprediction of response to selection when response was large. This underprediction disappeared when using prevalence in the offspring generation in the $1/P$ term in Equation 38. However, because prevalence in the offspring generation is initially unknown, as it depends on the response to selection, this prediction required solving the expression $R = \iota\rho_{A_y,\bar{y}}\sigma_{A_y}/(P + R)$, yielding

$$R_P = \frac{1}{2}\left(-P + \sqrt{P^2 + 4\iota\rho_{A_y,\bar{y}}\sigma_{A_y}}\right) \tag{39}$$

For a prevalence smaller than ~0.5, predictions from Equation 39 were very close to the observed response in prevalence (Figure 8B; for $P > 0.5$, results of Equations 38 and 39 are almost identical).

In conclusion, results in this section show that response to selection in the prevalence of endemic infectious diseases is a factor $1/P$ greater than suggested by the ordinary breeding values for individual binary infection status. Thus breeders can predict response to selection by upscaling the selection differential in the usual estimated breeding values for binary infection status by a factor $1/P$.

8.   Direct and indirect genetic variance for endemic prevalence

In this section, we partition the total additive genetic variance for endemic prevalence into direct and indirect genetic components. This partitioning is relevant, because IGE respond fundamentally different to selection than DGE (Griffing 1967, 1977; Wright 1985; Moore *et al.*, 1997; Muir 2005; Bijma 2010, 2011; see Discussion). We can partition the total breeding value for prevalence into a direct and an indirect component,

$$A_P = A_{P_D} + A_{P_I} \tag{40}$$

Analogously, we can partition the full additive genetic variance in prevalence into components due to direct genetic variance, indirect genetic variance and a covariance,

$$\sigma_{A_P}^2 = \sigma_{A_{P_D}}^2 + 2\sigma_{A_{P_D}A_{P_I}} + \sigma_{A_{P_I}}^2 \tag{41}$$

In the absence of genetic variation in infectivity, the breeding value for own infection status is
a fraction P of the breeding value for prevalence (Equation 33). Hence, a fraction P of the additive
genetic effects of susceptibility and recovery on prevalence affects the infection status of the individual
itself and thus represents a direct effect, while the remaining fraction $(1 - P)$ represents an indirect
effect. For infectivity, the entire genetic effect is indirect, because an individual's infectivity does not
affect its own infection status. It follows from Equations 31a that

$$A_{P_D} = \frac{e^{\sigma^2_{A_{lR_0}}}}{R_0} \, P \left(A_{l\gamma} - A_{l\alpha} \right) \tag{42a}$$

$$A_{P_I} = \frac{e^{\sigma^2_{A_{lR_0}}}}{R_0} \left((1 - P)A_{l\gamma} + A_{l\varphi} - (1-P)A_{l\alpha} \right) \tag{42b}$$

820   Note that Equation 42a represents the breeding value for individual infection status (A_y), but the
821   current expression emphasizes the partitioning of A_P into direct and indirect effects. The direct and
822   indirect genetic (co)variances are given by

$$\sigma^2_{A_{P_D}} = \frac{e^{2\sigma^2_{A_{lR_0}}}}{R_0^2} P^2 \left(\sigma^2_{A_{l\gamma}} - 2\sigma_{A_{l\gamma}A_{l\alpha}} + \sigma^2_{A_{l\alpha}} \right) \tag{43a}$$

$$\sigma_{A_{P_D}A_{P_I}} = \frac{e^{2\sigma^2_{A_{lR_0}}}}{R_0^2} P \left\{ (1 - P) \left(\sigma^2_{A_{l\gamma}} - 2\sigma_{A_{l\gamma}A_{l\alpha}} + \sigma^2_{A_{l\alpha}} \right) + \sigma_{A_{l\gamma}A_{l\varphi}} - \sigma_{A_{l\gamma}A_{l\alpha}} \right\} \tag{43b}$$

$$\sigma^2_{A_{P_I}} = \frac{e^{2\sigma^2_{A_{lR_0}}}}{R_0^2} \left\{ (1 - P)^2 \left(\sigma^2_{A_{l\gamma}} - 2\sigma_{A_{l\gamma}A_{l\alpha}} + \sigma^2_{A_{l\alpha}} \right) + 2(1 - P)\left(\sigma_{A_{l\gamma}A_{l\varphi}} - \sigma_{A_{l\varphi}A_{l\alpha}} \right) + \sigma^2_{A_{l\varphi}} \right\}$$
$$\tag{43c}$$

Figure 9 shows the total additive genetic variance for the endemic prevalence and the fractions
due to DGE, IGE and their covariance, for a scenario with equal genetic variances in susceptibility,
infectivity and recovery and covariances equal to zero. For an endemic prevalence smaller than 0.5, IGE
contribute the majority of the genetic variance. For example, for an endemic prevalence of 0.3, the total
additive genetic variance consists of 6% direct genetic variance, 66% indirect genetic variance and 28%
direct-indirect genetic covariance. These results imply that IGE dominate the heritable variation and
response to selection for the endemic prevalence of infectious diseases, unless prevalence is high.

Figure 9 – Total additive genetic variance in endemic prevalence (V_{total}; secondary y-axis) and the relative contributions of DGE, IGE and their covariance (f_{direct}, f_{indirect}, and $f_{\text{covariance}}$; primary y-axis). For constant values of $\sigma^2_{A_\gamma} = \sigma^2_{A_\varphi} = \sigma^2_{A_\alpha} = 0.3^2$ and covariances equal to zero. Results are obtained from Equations 32b, 3, and 43a-c.

Discussion

We have presented a quantitative genetic theory for endemic infectious diseases, with a focus on the genetic factors that determine the endemic prevalence. We defined an additive model for the logarithm of individual susceptibility, infectivity and rate of recovery, which results in normally distributed breeding values for the logarithm of R_0. Next, we investigated the impact of genetic heterogeneity on the population level, for both R_0 and the endemic prevalence. Results show that, despite heterogeneity, R_0 remains equal to the mean individual genotypic value for R_0. Subsequently, we considered genetic effects of individuals on their own infection status and on the endemic prevalence in the population. Building on the breeding value for the logarithm of R_0, we showed that genotypic values and genetic parameters for the prevalence follow from the known properties of the log-normal distribution. In the absence of genetic variation in infectivity, genetic effects for the endemic prevalence are a factor 1/prevalence greater than the ordinary genetic values for individual binary infection status. Hence, even though prevalence is the simple average of individual binary infection status, breeding values for prevalence show much more variation than those for individual infection status. These results imply that the genetic variance that determines the potential response of the endemic prevalence to selection is largely due to IGE, and thus hidden to classical genetic analysis and selection. For susceptibility and recovery, a fraction 1-P of the full genetic effect on endemic prevalence is due to IGE, whereas the effect of infectivity is entirely due to IGE. Hence, the genetic variance that determines the potential response of the endemic prevalence to selection must be much greater than expected based on classical

quantitative genetic theory, particularly at low levels of the prevalence (Figure 7). We evaluated this implication using stochastic simulation of endemics following standard methods in epidemiology, where parents of the next generation were selected based on their own infection status (mass selection). The results of these simulations show that response to selection in the observed prevalence and in the breeding value for prevalence increases strongly when prevalence decreases, and closely matches our predictions, which supports the theoretical findings presented here.

Model assumptions: Following Anacleto *et al.* (2015, 2019), Biemans *et al.* (2019) and Pooley *et al.* (2020), we assumed a linear additive model with normally distributed effects for the logarithm of susceptibility, infectivity and recovery, leading to a normal distribution of the additive genetic values for the logarithm of R_0 (Equation 10). For complex traits, it is common to assume normally distributed genetic effects, based on the central limit theorem (Fisher 1918). Because susceptibility, infectivity and recovery act multiplicatively in the expression for R_0, and because R_0 is non-negative, we specified a normal distribution for its logarithm. This resulted in an additive model on the log scale, which fits with the infinitesimal model, and also translates the $[0, \infty)$ domain of R_0 to the $(-\infty, +\infty)$ domain of the normal distribution. Hence, we assumed constant genetic parameters for the logarithm of R_0. The same approach has been used to model genetic variation in the residual variance, which is also restricted to non-negative values (SanCristobal-Gaudy *et al.* 1998; Hill and Mulder 2010). The log-normal distribution of genotypic values for R_0 results in a decrease of the genetic standard deviation in R_0 with decreasing R_0 (Figure 2), which seems reasonable given the presence of a lower bound for R_0. Moreover, the log-normal distribution for R_0 is convenient, because it results in simple expressions for the breeding value and the genetic variance for prevalence.

The assumption of a normal distribution for the logarithm of genotypic values for R_0 also agrees with the standard implementation of generalized linear (mixed) models (GLMM; Nelder and Wedderburn, 1972). R_0 refers to an expected number of infected individuals; In other words, R_0 is the expected value of count data. In GLMM, the default link function for count data is the log-link (McCullagh and Nelder, 2019). Hence, our linear model for the logarithm of R_0 also agrees with common statistical practise.

The strong increase of the genetic variance in prevalence with decreasing R_0 (Figure 5A) is not due to the assumption of lognormality of R_0. On the contrary, the log-normal distribution results in a decrease of the genetic standard deviation in R_0 with decreasing R_0 (Figure 2). The strong increase in the genetic variance in prevalence results from the relationship between R_0 and the prevalence in the endemic steady state (Figure 1; Equation 3), which becomes steeper when R_0 is closer to one. This relationship is very well established in epidemiology since Weiss and Dishon (1971; *e.g.*, Keeling & Rohani, 2011).

While we defined an additive genetic model for the logarithm of R_0, we can also find the additive genetic effect (breeding value) for R_0 itself. Using results of Appendix 4,

$$A_{R_0,i} = c e^{\frac{1}{2}\sigma^2_{A_{lR_0}}} A_{lR_0,i}$$

Hence, the A_{R_0} represents the additive component of the genotypic value for R_0. However, because our model is additive on the log-scale, while the genotypic value for R_0 includes non-additive genetic effects, we decided to build our theory on the breeding value for the logarithm of R_0.

Other compartmental models: In this work, we focused on endemic infectious diseases following a SIS-model, where individuals can be either susceptible (S, *i.e.*, non-infected) or infected (I). Hence, we assumed the infection does not confer any long-lasting immunity, and we ignored the potential existence of infected classes ("compartments") other than S and I, such as recovered infected individuals that are not yet susceptible again. Moreover, we ignored the influx of new individuals into the population due to births, and the removal of individuals due to deaths.

A key condition for validity of our results is that the pathogen can replicate only in the host individual, meaning that a reduction in infected individuals fully translates into reduced exposure of the host population to the pathogen. The mere survival of the pathogen in the environment does not violate our assumptions; see Hulst *et al*., 2021 for a discussion. Our conclusions are not limited to SIS models if this condition is met, but apply to all models with no longer lasting immunity. For models with temporary immunity (*e.g.,* SIRS) or lifelong immunity (*e.g.,* SIR) the conclusions with respect to infectivity and susceptibility will be true, but the genetic variation in recovery may play no role (*e.g.,* SIR) or a different more restricted role (*e.g.,* SIRS when the R compartment lasts long).

Also infections that do confer long-lasting immunity may show endemic behavior when a population is large enough. Measles in the human population before the introduction of vaccination are is a well-known example. For such infections, the same mechanisms as discussed above will play a role and the endemic prevalence for a homogeneous population still follows from Equation 3. However, the introduction of new susceptibles by birth can no longer be ignored, and recovery of infected individuals does not result in new susceptible individuals. Thus the role of recovery will change, and the genetic make-up of the newborn individuals becomes relevant, particularly in populations undergoing selection.

Positive feedback: The increasing difference between the breeding value for prevalence and the breeding value for individual infection status at lower prevalence (Equation 33) is a result of the increasing slope of the relationship between R_0 and the endemic prevalence (Equation 3, Figure 1). Equation 3 follows directly from a simple equilibrium condition (see text above Equation 3). However, the focus on the equilibrium partly obscures the underlying mechanism. Figures 10A and B illustrate that the difference between A_P and A_y originates from positive feedback effects in the transmission dynamics. (Figure 10 shows results for selection against susceptibility, selection for faster recovery would yield identical results). With lower susceptibility fewer individuals will become infected, which subsequently translates into a reduced transmission rate, followed by a further reduction in the number of infected individuals, *etc*, resulting in a positive feedback loop (Figure 10A). The initial change in prevalence before feedback effects manifest is equal to the selection differential in breeding value for individual infection status ($\Delta \bar{A}_y$; horizontal lines in Figure 9). This change represents the direct response due to reduced susceptibility, and does not include any change in exposure of susceptible individuals to infected herd mates. Next, prevalence decreases further because the initial decrease in prevalence reduces the exposure of susceptible individuals to infected herd mates. This additional decrease represents the indirect response to selection via the "social" environment. Without genetic variation in infectivity, the direct response makes up a fraction P of the total response in prevalence, and the indirect response a fraction $1 - P$.

Figure 10 – Positive feedback after selection for lower susceptibility. Panel A: Diagram of the SIS compartmental model illustrating the feedback, with the number of susceptible (*S*) and infectious (*I*) individuals and the transmission and recovery rates (ignoring heterogeneity for simplicity). A reduction in the transmission rate parameter β reduces *I*, which in turn reduces the transmission rate, leading to a further reduction in *I*, *etc*. Panel B: Convergence of the prevalence to the new equilibrium after selection. For two populations; one starting at $P = 0.25$ (red triangles; $c = 1.333$), the other at $P = 0.125$ (blue circles; $c = 1.143$). The *x*-axis represents cycles of the transmission loop. The horizontal dotted lines show the prevalence predicted by the breeding value for binary infection status, and represent the direct effect. The Asterix shows the equilibrium prevalence after convergence, which occurs a little later than $t = 50$ for the lower line. The genetic selection differential for binary infection status equals $\Delta \bar{A}_y = -0.01$ for each of the two populations. The initial response to selection (the *y*-axis difference between $t = 0$ and $t = 1$) is equal to the $\Delta \bar{A}_y$ of -0.01 for both scenarios. Total response is -0.04 for the scenario with $P = 0.25$, and -0.08 for the scenario with $P = 0.125$, corresponding to -0.01/0.25 and -0.01/0.125. Results in panel B follow from iterating on Equation 20a, using a single value for $\mathcal{R}_{0,i}$, with $\alpha_i = \bar{\varphi}_{\text{inf}} = 1$ and choosing γ so that the selection differential $\Delta \bar{A}_y = -0.01$ (using $\gamma = \frac{G_y}{P(1 - G_y)}/c$ from Equation 20a). In each iteration, the *P* in the righthand side of Equation 20a is replaced by the P_i calculated from Equation 20a in the previous iteration. This iteration converges to the prevalence given by Equation 3 (assuming negligible heterogeneity).

Herd immunity: In Figure 8A, the infection ultimately goes extinct due to mass selection for individual infection status. This happens due to a phenomenon known as herd immunity (Fine, 1993). In the final generation, the infection disappears because R_0 falls below a value of one; not because all the individuals have become fully resistant to infection. This result is similar to the eradication of an infection by means of vaccination, which also does not require full immunity of all individuals and can also be achieved when only part of a population is vaccinated (Anderson and May, 1985). As can be seen in Figure 10 and in simulation results of Hulst *et al*. (2021), herd immunity develops over cycles of the transmission-recovery loop. Thus the full benefits of genetic selection or vaccination do not manifest immediately, as it takes some time for a population to converge to the new endemic steady state.

The relevance of herd immunity for response to genetic selection can be illustrated using the data underlying Figure 8A. For the population starting at a prevalence of 0.5, the contact rate is equal to two, and the mean breeding value for log-susceptibility is equal to zero in the initial generation ($c = 2$, $\bar{A}_{l\gamma} = 0$, so that $R_0 \approx ce^{\bar{A}_{l\gamma}} = 2$). In the final generation, the mean breeding value for log-susceptibility has dropped to -0.73, so that $R_0 \approx 2e^{-0.73} = 0.96$. Hence, $R_0 < 1$, explaining extinction. However, if the average individual of the final generation would have been exposed to the infection pressure of the first generation, then the expected prevalence for this individual would have been 0.32 (From Equation 20a, with $R_{0,i} = 0.96$ and $P = 0.5$). Hence, the individual would have been infected 32% of the time. Nevertheless, in a population consisting entirely of this type of individual, as is the case in the final generation, the infection will no longer be present in the long term. This example illustrates the relevance of indirect effects for herd immunity and for response to selection of infectious diseases.

Utilization of hidden genetic variation for genetic improvement: In this work, we have shown that a fraction $1 - P$ of the full individual genetic effect on the endemic prevalence represents an IGE, because only a fraction P of the full effect surfaces in the infection status of the individual itself (excluding genetic variation in infectivity; Equation 33 and Appendix 5). In other words, a fraction $1 - P$ of the individual genetic effects of susceptibility and recovery on the prevalence are hidden to direct selection and classical genetic analysis. Nevertheless, results in Figure 8 show that prevalence responds rapidly to selection, particularly when prevalence is small. Hence, prevalence responds faster to selection when a greater proportion of its heritable variation is hidden, and when heritability is low (Figure 7A), which seems a paradox.

However, the IGEs due to susceptibility and recovery are a special kind, because they are fully correlated to the corresponding DGE. For each of the two traits, there is only a single genetic effect ($A_{l\gamma}$ and $A_{l\alpha}$, respectively), which has both a direct effect and an indirect effect on the prevalence. Hence, when selection changes the mean DGE, the mean IGE changes correspondingly. This can be seen from Equation 38, where the term σ_{A_y}/P represents the full additive genetic standard deviation in prevalence (as is clear from Equation 33), while the accuracy ($\rho_{A_y,\bar{y}}$) refers to selection for the direct effect only. Hence, without genetic variation in infectivity, the total response to selection based on individual infection status can be interpreted as the sum of a direct response in DGE and a correlated response in IGE,

$$R_{P,direct} = \iota \rho_{A_y,\bar{y}} \sigma_{A_y}$$

$$R_{P,correlated} = \iota \rho_{A_y,\bar{y}} \sigma_{A_y} \frac{1 - P}{P}$$

and the sum of $R_{P,direct}$ and $R_{P,correlated}$ is equal to Equation 38. The $R_{P,direct}$ is the expected response to selection based on ordinary genetic analysis of individual infection status. The $R_{P,correlated}$ represents the additional response due to IGE. The direct response occurs immediately in the first cycle of the transmission loop (Figure 10B), while the indirect response manifests gradually over several cycles of the transmission-recovery loop, particularly when prevalence is small (see also result in Hulst *et al.* 2021).

The response due to the IGE of susceptibility and recovery arises naturally when selecting for lower individual infection status (*i.e.*, for the direct effect); it does not require any specific measures of

the breeder. Thus, on the one hand, our results imply that response to genetic selection against infectious diseases should be considerably greater than currently believed, even when no changes are made to the selection strategy. While empirical studies are scarce, the available results support this expectation (discussed in Hulst *et al.* 2021).

On the other hand, however, classical selection for direct effects is not the optimal way to reduce prevalence, for the following two reasons. First, classical selection does not target genetic effects on infectivity, because an individual's infectivity does not affect its own infection status (Lipschutz-Powell et al., 2012). Hence, infectivity changes merely due to a potential genetic correlation with susceptibility and/or recovery. When this correlation is unfavourable, infectivity will increase and response in prevalence will be smaller than expected based on the genetic selection differentials for susceptibility and recovery. (And thus smaller than the result of Equation 38). In theory, this could even lead to a negative net response (Griffing 1967). This is similar to the case with social behaviour-related IGEs on survival in laying hens and Japanese quail, where selection for individual survival has sometimes increased mortality (Craig and Muir, 1996; Muir 2005). This scenario seems unlikely for infectious diseases, but at present we lack knowledge of the multivariate genetic parameters of susceptibility, infectivity and recovery to make well-founded statements.

Second, even in the absence of genetic variation in infectivity, individual selection for susceptibility and recovery is non-optimal because the accuracy of selection is limited due to limited heritability, particularly at low prevalence (Figure 7A). The response to selection in traits affected by IGE can be increased by using kin selection and/or group selection (Griffing 1976; Muir 1996; Bijma 2011), or by including IGE in the genetic analysis (Muir 2005, Bijma *et al.* 2007b; Muir *et al.* 2013; Biemans *et al.* 2019, Anacleto *et al.* 2015, Pooley *et al.* 2020). Kin selection occurs when transmission takes place between related individuals, for example within groups of relatives (Anche *et al.* 2014). Group selection refers to the selection of parents for the next generation based on the prevalence in the group in which transmission takes place, rather than on individual infection status (Griffing 1976). Both theoretical and empirical work shows that kin and group selection lead to utilization of the full genetic variation, including both DGE and IGE (Griffing 1976, Muir 1996, 2005; Bijma and Wade, 2008; Bijma 2010, 2011). For infectious diseases, the work of Anche *et al.* (2014) illustrates the effect of kin selection, where favourable alleles for susceptibility increase much faster in frequency when disease transmission is between related individuals. Simulation studies on IGE in pig populations suggest that the benefits of kin selection should also extend to breeding schemes based on genomic prediction (Chu *et al.* 2021).

Do pathogens create kin selection? Exposure to infectious pathogens is a major driver of the evolution of host populations by natural selection, both in animals and plants (reviewed in Karlsson *et al.* 2014 and Ebert and Fields 2020). In the human species, for example, a study of genetic variation in 50 worldwide populations reveals that selection on infectious pathogens is the primary driver of local adaptation and the strongest selective force that shapes the human genome (Barreiro and Quintana-Murci 2010; Fumagalli *et al.* 2011). The key role of infectious pathogens in natural selection, together with the large contribution of IGE to the genetic variation in prevalence in the host population, indicates that IGE must have been an important fitness component in the evolutionary history of populations. This, in turn, suggests that associating with kin may have evolved as an adaptive behaviour. In other

words, the key role of infectious diseases in natural selection might lead to social structures where individuals associate preferably with kin, because such behaviour has indirect fitness benefits. This is because interactions among kin lead to utilisation of the full heritable variation in fitness, including both DGE and IGE (Bijma, 2010), and thus accelerate response of fitness to selection. At low to moderate levels of the endemic prevalence the indirect genetic variance in prevalence might be sufficiently large for such behaviour to evolve, even in the absence of direct fitness benefits such as preferential behaviour towards kin. While this is a complex issue requiring careful quantitative modelling, including migration and the emergence of selfish mutants, the key role of pathogens in natural selection together with the large IGE demonstrated here strongly suggest the importance of kin selection in the history of life.

In agriculture, the implementation of kin selection may be feasible when animals can be kept in kin groups or plants can be grown in plots of a single genotype or a family in the breeding population. In many cases, however, this will not be feasible, and other methods are required to optimally capture the IGE underlying the prevalence of infectious diseases. Current developments in sensing technology and artificial intelligence enable the development of tools for large scale automated collection of longitudinal data on individual infection status, and also on the contact structure between individuals (relevant mainly in animals). These advances, together with genomic prediction and recently developed statistical methods for the estimation of the direct and indirect genetic effects underlying the transmission of the infection (Biemans *et al*., 2019; Pooley *et al*. 2020), could represent a much-needed breakthrough in artificial selection against infectious diseases in agriculture. Our results on genetic variation and response to selection suggest that such selection is way more promising than currently believed.

References

Anacleto, O., Garcia-Cortés, L. A., Lipschutz-Powell, D., Woolliams, J. A., & Doeschl-Wilson, A. B. (2015). A novel statistical model to estimate host genetic effects affecting disease transmission. *Genetics*, *201*(3), 871-884.

Anacleto, O., Cabaleiro, S., Villanueva, B., Saura, M., Houston, R. D., Woolliams, J. A., & Doeschl-Wilson, A. B. (2019). Genetic differences in host infectivity affect disease spread and survival in epidemics. Scientific reports, 9(1), 1-12.

Anche, M. T., De Jong, M. C. M., & Bijma, P. (2014). On the definition and utilization of heritable variation among hosts in reproduction ratio R 0 for infectious diseases. *Heredity*, *113*(4), 364-374.

Anderson, R., May, R. Population biology of infectious diseases: Part I. *Nature* **280,** 361–367 (1979). https://doi.org/10.1038/280361a0

Anderson, R. M., & May, R. M. (1985). Vaccination and herd immunity to infectious diseases. Nature, 318(6044), 323-329.

Aznar, I., Frankena, K., More, S. J., O'Keeffe, J., McGrath, G., & De Jong, M. C. M. (2018). Quantification of Mycobacterium bovis transmission in a badger vaccine field trial. *Preventive veterinary medicine*, *149*, 29-37.

Barreiro LB, Quintana-Murci L (2010) From evolutionary genetics to human immunology: How selection shapes host defence genes. Nat Rev Genet 11(1): 17–30.

Biemans, F., de Jong, M. C., & Bijma, P. (2017). A model to estimate effects of SNPs on host susceptibility and infectivity for an endemic infectious disease. *Genetics Selection Evolution*, *49*(1), 1-13.

Biemans, F., De Jong, M. C., & Bijma, P. (2019). Genetic parameters and genomic breeding values for digital dermatitis in Holstein Friesian dairy cattle: host susceptibility, infectivity and the basic reproduction ratio. *Genetics Selection Evolution*, *51*(1), 1-13.

Bijma, P., Muir, W. M., & Van Arendonk, J. A. (2007). Multilevel selection 1: quantitative genetics of inheritance and response to selection. Genetics, 175(1), 277-288.

Bijma, P. (2010). Fisher's fundamental theorem of inclusive fitness and the change in fitness due to natural selection when conspecifics interact. Journal of evolutionary biology, 23(1), 194-206.

Bijma, P. (2011). A general definition of the heritable variation that determines the potential of a population to respond to selection. *Genetics*, *189*(4), 1347-1359.

Bijma, P., & Wade, M. J. (2008). The joint effects of kin, multilevel selection and indirect genetic effects on response to genetic selection. Journal of evolutionary biology, 21(5), 1175-1188.

Bishop SC, Woolliams JA (2010) On the Genetic Interpretation of Disease Data. PLoS ONE 5(1): e8940. https://doi.org/10.1371/journal.pone.0008940.

Bishop, S., Doeschl-Wilson, A. B., & Woolliams, J. A. (2012). Uses and implications of field disease data for livestock genomic and genetics studies. Frontiers in genetics, 3, 114

Bolker, B. M., & Grenfell, B. T. (1996). Impact of vaccination on the spatial correlation and persistence of measles dynamics. *Proceedings of the National Academy of Sciences*, 93(22), 12648-12653.

Chu, T. T., Henryon, M., Jensen, J., Ask, B., & Christensen, O. F. (2021). Statistical model and testing designs to increase response to selection with constrained inbreeding in genomic breeding programs for pigs affected by social genetic effects. Genetics Selection Evolution, 53(1), 1-16.

Craig, J. V., & Muir, W. M. (1996). Group selection for adaptation to multiple-hen cages: beak-related mortality, feathering, and body weight responses. *Poultry Science*, *75*(3), 294-302.

De Villemereuil, P., Schielzeth, H., Nakagawa, S., & Morrissey, M. (2016). General methods for evolutionary quantitative genetic inference from generalized mixed models. *Genetics*, *204*(3), 1281-1294.

DeJong, M. C. M., vanderPoel, W. H. M., Kramps, J. A., Brand, A., & vanOirschot, J. T. (1996). Quantitative investigation of population persistence and recurrent outbreaks of bovine respiratory syncytial virus on dairy farms. American Journal of Veterinary Research, 57(5), 628-633.

Dempster, E. R., & Lerner, I. M. (1950). Heritability of threshold characters. Genetics, 35(2), 212.

O Diekmann; J.A.P. Heesterbeek; J.A.J. Metz (1990). "On the definition and the computation of the basic reproduction ratio   in models for infectious diseases in heterogeneous populations". Journal of Mathematical Biology. 28 (4): 356–382. doi:10.1007/BF00178324.

Diekmann, O., Heesterbeek, H., & Britton, T. (2012). Mathematical tools for understanding infectious disease dynamics (Vol. 7). Princeton University Press.

Ebert, D., & Fields, P. D. (2020). Host–parasite co-evolution and its genomic signature. Nature Reviews Genetics, 21(12), 754-768.

EFSA Panel on Animal Health and Welfare (AHAW). (2012). Scientific Opinion on Review of the European Union Summary Report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2009 and 2010 specifically for the data related to bovine tuberculosis, Echinococcus, Q fever, brucellosis and non-food borne diseases. EFSA Journal, 10(6), 2765.

Fine, P. E. (1993). Herd immunity: history, theory, practice. Epidemiologic reviews, 15(2), 265-302.

Fisher, R. A. (1918). XV.—The correlation between relatives on the supposition of Mendelian inheritance. Earth and Environmental Science Transactions of the Royal Society of Edinburgh, 52(2), 399-433.

Fumagalli, M., Sironi, M., Pozzoli, U., Ferrer-Admettla, A., Pattini, L., & Nielsen, R. (2011). Signatures of environmental genetic adaptation pinpoint pathogens as the main selective pressure through human evolution. PLoS Genet, 7(11), e1002355.

Gianola, D. (1982). Theory and analysis of threshold characters. *Journal of animal Science*, *54*(5), 1079-1096.

Greenhalgh, D., Diekmann, O., & de Jong, M. C. M. (2000). Subcritical endemic steady states in mathematical models for animal infections with incomplete immunity. Mathematical Biosciences, 165(1), 1-25. doi:10.1016/S0025-5564(00)00012-2.

Griffing, B. (1967). Selection in reference to biological groups I. Individual and group selection applied to populations of unordered groups. Australian Journal of Biological Sciences, 20(1), 127-140.

Griffing, B. (1976). Selection in reference to biological groups. V. Analysis of full-sib groups. Genetics, 82(4), 703-722.

Griffing, B. (1977) Selection for populations of interacting genotypes, in Proceedings of the International Congress on Quantitative Genetics, August 16-21, 1976 (Pollack, E., Kempthorne, O. and Bailey, T.B., eds), pp. 413–434, Iowa State University Press

Hethcote, H. W., 1989 Three basic epidemiological models, pp. 119-144 in *Applied mathematical ecology*. Springer.

Heringstad, B., G. Klemetsdal and T. Steine, 2007 Selection responses for disease resistance in two selection experiments with Norwegian red cows. Journal of dairy science 90**:** 2419-2426.

Hill, W. G., & Mulder, H. A. (2010). Genetic analysis of environmental variation. Genetics Research, 92(5-6), 381-395.

Hulst, A. D., de Jong, M. C., & Bijma, P. (2021). Why genetic selection to reduce the prevalence of infectious diseases is way more promising than currently believed. Genetics, 217(4), iyab024.

Karlsson, E. K., Kwiatkowski, D. P., & Sabeti, P. C. (2014). Natural selection and infectious disease in human
populations. Nature Reviews Genetics, 15(6), 379-393. : Host genetics influences susceptibility to infectious
disease.

Keeling, M. J., & Rohani, P. (2011). *Modeling infectious diseases in humans and animals*. Princeton university
press.

Kermack, W. O. and McKendrick, A. G. 1927. A contribution to the mathematical theory of epidemics. *Proc. R.*
Soc. Lond. Ser. A, 115: 700–721.

Knap, P. W., & Doeschl-Wilson, A. (2020). Why breed disease-resilient livestock, and how?. *Genetics Selection*
Evolution, *52*(1), 1-18.

Lipschutz-Powell, D., Woolliams, J. A., Bijma, P., & Doeschl-Wilson, A. B. (2012). Indirect genetic effects and
the spread of infectious disease: are we capturing the full heritable variation underlying disease
prevalence?. *PloS one*, *7*(6), e39551.

Lipschutz-Powell, D., Woolliams, J. A., & Doeschl-Wilson, A. B. (2014). A unifying theory for genetic
epidemiological analysis of binary disease data. Genetics Selection Evolution, 46(1), 1-12.

Martin, P., H. Barkema, L. Brito, S. Narayana and F. Miglior, 2018 Symposium review: Novel strategies to
genetically improve mastitis resistance in dairy cattle. Journal of dairy science 101**:** 2724-2736.

McCullagh, P., & Nelder, J. A. (2019). Generalized linear models. Routledge.

Metz, J. A. J. (1978). The epidemic in a closed population with all susceptibles equally vulnerable; some results
for large susceptible populations and small initial infections. *Acta biotheoretica*, *27*(1), 75-123.

Moore, A. J., Brodie III, E. D., & Wolf, J. B. (1997). Interacting phenotypes and the evolutionary process: I. Direct
and indirect genetic effects of social interactions. Evolution, 51(5), 1352-1362.

Muir, W. M. (1996). Group selection for adaptation to multiple-hen cages: selection program and direct
responses. *Poultry Science*, *75*(4), 447-458.

Muir, W. M. (2005). Incorporation of competitive effects in forest tree or animal breeding
programs. *Genetics*, *170*(3), 1247-1259.

Muir, W. M., Bijma, P., & Schinckel, A. (2013). Multilevel selection with kin and non-kin groups, experimental
results with Japanese quail (Coturnix japonica). *Evolution*, *67*(6), 1598-1606.

Nelder, J. A., & Wedderburn, R. W. (1972). Generalized linear models. Journal of the Royal Statistical Society:
Series A (General), 135(3), 370-384.

Nieuwhof, G. J., J. Conington and S. C. Bishop, 2009 A genetic epidemiological model to describe resistance to
an endemic bacterial disease in livestock: application to footrot in sheep. Genetics Selection Evolution 41**:** 19.

Pooley, C. M., Marion, G., Bishop, S. C., Bailey, R. I., & Doeschl-Wilson, A. B. (2020). Estimating individuals'
genetic and non-genetic effects underlying infectious disease transmission from temporal epidemic data. *PLOS*
Computational Biology, *16*(12), e1008447.

Robertson, A. (1950). Proof that the additive heritability on the p scale is given by the expression z2hx2/pq.
Genetics, 35, 234-36.

Rushton J. The economics of animal health and production. Wallingford: CABI; 2009.

Russell, G. E. (2013). Plant breeding for pest and disease resistance: studies in the agricultural and food sciences.

SanCristobal-Gaudy, M., Elsen, J. M., Bodin, L., & Chevalet, C. (1998). Prediction of the response to a selection
for canalisation of a continuous trait in animal breeding. Genetics Selection Evolution, 30(5), 423-451.

Schrag, S. J., & Wiener, P. (1995). Emerging infectious disease: what are the relative roles of ecology and evolution?. *Trends in ecology & evolution*, *10*(8), 319-324.

A J Springbett, K MacKenzie, J A Woolliams, S C Bishop, The Contribution of Genetic Diversity to the Spread of Infectious Diseases in Livestock Populations, *Genetics*, Volume 165, Issue 3, 1 November 2003, Pages 1465–1474, https://doi.org/10.1093/genetics/165.3.1465

Thanner, S., Drissner, D., & Walsh, F. (2016). Antimicrobial resistance in agriculture. *MBio*, *7*(2).

Tsairidou, S., Anacleto, O., Woolliams, J. A., & Doeschl-Wilson, A. (2019). Enhancing genetic disease control by selecting for lower host infectivity and susceptibility. *Heredity*, *122*(6), 742-758.

Walsh, B., & Lynch, M. (2018). Evolution and selection of quantitative traits. Oxford University Press.

Weiss, G. H., & Dishon, M. (1971). On the asymptotic behavior of the stochastic and deterministic models of an epidemic. *Mathematical Biosciences*, 11(3-4), 261-265.

Wolf, J. B., Brodie III, E. D., Cheverud, J. M., Moore, A. J., & Wade, M. J. (1998). Evolutionary consequences of indirect genetic effects. *Trends in ecology & evolution*, *13*(2), 64-69.

Wright, A. J. (1985). Selection for improved yield in inter-specific mixtures or intercrops. *Theoretical and applied genetics*, 69(4), 399-407.

Table 1 – Notation key

N	Total number of individuals in the population in which the endemic takes place; $N = I + S$
I	Number of infected individuals in the population
S	Number of susceptible (*i.e.,* non-infected) individuals in the population
P	Prevalence in the endemic equilibrium; $P = I/N$; $P = \bar{y}$
R_0	Basic reproduction number
R_P	Response of prevalence to selection, *i.e.,* change in prevalence per generation
$\mathcal{R}_{0,i}$	R_0-like quantity that determines the prevalence for type i (see Equation 20)
A	Breeding value
$A_{l\gamma}, A_{l\varphi}, A_{l\alpha}$	Breeding value for the logarithm of susceptibility, infectivity and recovery rate
A_{lR_0}, A_{R_0}	Breeding value for the logarithm of R_0; breeding value for R_0
A_y, A_P	Breeding value for individual binary disease status; breeding value for prevalence
G_{R_0}	Genotypic value for R_0
G_y, G_P	Genotypic value for individual disease status; genotypic value for prevalence.
T_P^2	Ratio of variance in breeding value for prevalence over phenotypic variance in y
c	Effective contact rate. Without heterogeneity $R_0 = c$.
h_y^2	Heritability of individual binary disease status
y	Individual binary infection status; infected: $y = 1$; non-infected: $y = 0$.
i,j	Subscript denoting an individual.
α	Recovery rate (relative to a value of 1)
β_{ij}	Transmission rate parameter from individual j to individual i.
γ	Susceptibility (relative to a value of 1)
φ	Infectivity (per unit of time, relative to a value of 1)
$\bar{\varphi}_{\text{inf}}$	Mean infectivity of the infected individuals in the endemic equilibrium
ϕ	Life time infectivity (relative to a value of 1);
$\bar{\phi}_{\text{typ}}$	$\bar{\phi}$ for the typical infected individual in an otherwise non-infected population
ι	Intensity of selection; selection differential expressed in SD units
$\rho_{A_y,\bar{y}}$	Accuracy of mass selection, correlation of A_y and $\bar{y}$ in the selection candidates
$\sigma^2_{A_{l\gamma}}$	Variance of $A_{l\gamma}$ among individuals; analogous for $\sigma^2_{A_{l\varphi}}$ and $\sigma^2_{A_{l\alpha}}$
$\sigma_{A_{l\gamma}A_{l\varphi}}$	Covariance of $A_{l\gamma}$ and $A_{l\varphi}$; analogous for $\sigma_{A_{l\gamma}A_{l\alpha}}$ and $\sigma_{A_{l\varphi}A_{l\alpha}}$
$\sigma^2_{A_{lR_0}}$	Variance of the breeding values for the logarithm of R_0
$\sigma^2_{A_P}$	Additive genetic variance for endemic prevalence
$\sigma^2_{A_y}$	Additive genetic variance in individual binary infection status12/07/2021
$\sigma^2_{A_{P_D}}, \sigma^2_{A_{P_I}}$	Direct and indirect additive genetic variance for endemic prevalence, respectively
$\sigma_{A_{P_D}A_{P_I}}$	Direct-indirect additive genetic covariance for endemic prevalence

1256

1257 **List of appendices**

1266

Appendix 1

R_0 with heterogeneity and log-normally distributed susceptibility, infectivity en recovery

We assume that the transmission rate from infected individual j to susceptible individual i is proportional to the product of the infectivity of j and the susceptibility of i (Equation 5),

$$\beta_{ij} = c\gamma_i\varphi_j.$$

So there is no interaction between i and j. (This property is known as separable mixing in the epidemiological literature; Diekmann *et al*. 1990; 2013). Moreover, we assume that susceptibility, infectivity and recovery follow a log-normal distribution (Equations 6 and 7). We also assume that the population is not very small, so that in the early phase of an endemic where only few individuals are infected, the composition of the remaining susceptible individuals is not affected.

Because R_0 refers to the "total number of individuals that become infected by a typical infected individual over its entire infectious lifetime", we define an individual lifetime infectivity, which is the product of an individual's infectivity per unit of time and the average duration of its infectious lifetime,

$$\phi_i = \varphi_i/\alpha_i,$$

which follows a log-normal distribution with parameters following from those of φ and α. Hence, we have condensed our three genetic effects into two.

We can find R_0 from

$$R_0 = c\,\bar{\gamma}\,\bar{\phi}_{typ}$$

where $\bar{\phi}_{typ}$ is the lifetime infectivity of the typical infected individual, and $\bar{\gamma}$ is the simple average susceptibility in the population,

$$\bar{\gamma} = \int_0^\infty \gamma\, g(\gamma)\, d\gamma.$$

where $g(\gamma)$ is the pdf of γ. We can use the simple average of susceptibility in this expression because we assume the population is large.

With separable mixing, the typical infected individual is created immediately in the first generation of disease transmission. This is the case because there is no interaction between γ and φ, so that the properties of the typical infected individual are determined entirely by susceptibility. Hence, the pdf of γ for the typical infected individual follows from weighing $g(\gamma)$ by γ,

$$g_{typ}(\gamma) = \frac{1}{\bar{\gamma}}\gamma\, g(\gamma)$$

Since the properties of the typical infected individual depend on susceptibility only, we can find $\bar{\phi}_{typ}$ by averaging ϕ over its distribution conditional on γ, and subsequently averaging over the distribution of γ,

$$1299 \qquad \bar{\phi}_{typ} = \int_0^\infty \left(\int_0^\infty \phi\, f(\phi|\gamma)\, d\phi \right) g_{typ}(\gamma)\, d\gamma$$

1300   Hence, we now have the elements of R_0, but still need to solve the integral expression.

1301   Because conditional Normal distributions are also Normal and the logarithm is a bijective
1302   function, $\phi|\gamma$ follows a log-normal distribution with parameters being the conditional mean and variance
1303   of the Normal distribution,

$$1304 \qquad \phi|\gamma \sim Lnorm\left(\mu = b_{\phi,\gamma} A_{l\gamma}\, ;\, \sigma^2 = \left(1 - \rho_{\gamma,\phi}^2\right)\sigma_{A_{l\phi}}^2 \right)$$

1305   with $b_{\phi,\gamma} = cov(A_{l\gamma}, A_{l\phi})/var(A_{l\gamma})$ denoting the regression coefficient of $A_{l\phi}$ on $A_{l\gamma}$, and $\rho_{\gamma,\phi}^2 =$
1306 $cov^2(A_{l\gamma}, A_{l\phi})/[var(A_{l\gamma})var(A_{l\phi})]$ the squared correlation, where $A_{l\phi}$ denotes the breeding value
1307   for logarithm of lifetime infectivity.

1308   Hence, the inner integral is the mean of a log-normal variate, which is of the form $\exp(\mu +$
1309 $\frac{\sigma^2}{2})$,

$$1310 \qquad \int_0^\infty \phi\, f(\phi|\gamma)\, d\phi = E[\phi|\gamma] = \exp\left(b_{\phi,\gamma} A_{l\gamma} + \frac{1}{2}\left(1 - \rho_{\gamma,\phi}^2\right)\sigma_{A_{l\phi}}^2 \right)$$

$$1311 \qquad = e^{\frac{1}{2}\left(1 - \rho_{\gamma,\phi}^2\right)\sigma_{A_{l\phi}}^2}\, e^{b_{\phi,\gamma} A_{l\gamma}}.$$

1312   Since the first term of this expression is a constant,

$$1313 \qquad \bar{\phi}_{typ} = e^{\frac{1}{2}\left(1 - \rho_{\gamma,\phi}^2\right)\sigma_{A_{l\phi}}^2} \int_0^\infty e^{b_{\phi,\gamma} A_{l\gamma}}\, g_{typ}(\gamma)\, d\gamma.$$

1314   Substituting $g_{typ}(\gamma) = \frac{1}{\bar{\gamma}}\gamma\, g(\gamma)$, and replacing $g(\gamma)$ by the corresponding log-normal pdf yields

$$1315 \qquad \int_0^\infty e^{b_{\phi,\gamma} A_{l\gamma}}\, g_{typ}(\gamma)\, d\gamma = \frac{1}{\bar{\gamma}\sigma\sqrt{2\pi}} \int_0^\infty e^{-\frac{A^2}{2\sigma^2} + bA}\, d\gamma$$

1316   where we simplified the notation for brevity, using $\sigma^2 = \sigma_{A_{l\gamma}}^2$, $b = b_{\phi,\gamma}$, and $A = A_{l\gamma}$.

1317   Next, we change variable, using $d\gamma = e^A dA$, and adjust the bounds accordingly,

$$1318 \qquad \frac{1}{\bar{\gamma}\sigma\sqrt{2\pi}} \int_{-\infty}^\infty e^{-\frac{A^2}{2\sigma^2} + bA}\, e^A dA = \frac{1}{\bar{\gamma}\sigma\sqrt{2\pi}} \int_{-\infty}^\infty e^{-\frac{A^2}{2\sigma^2} + (1+b)A}\, dA$$

1319   Solving the integral term in Mathematica-online yields

$$1320 \qquad \frac{1}{\bar{\gamma}\sigma\sqrt{2\pi}} \int_{-\infty}^\infty e^{-\frac{A^2}{2\sigma^2} + (1+b)A}\, dA = \frac{1}{\bar{\gamma}}\, e^{\frac{1}{2}\sigma_{A_{l\gamma}}^2 (1+b_{\phi,\gamma})^2}$$

1321

$$1322 \qquad \bar{\phi}_{typ} = e^{\frac{1}{2}(1-\rho_{\gamma,\phi}^2)\sigma_{A_{l\phi}}^2} \frac{1}{\bar{\gamma}} e^{\frac{1}{2}\sigma_{A_{l\gamma}}^2(1+b_{\phi,\gamma})^2}$$

1323

$$1324 \qquad \bar{\phi}_{typ} = \frac{1}{\bar{\gamma}} e^{\frac{1}{2}\left[(1-\rho_{\gamma,\phi}^2)\sigma_{A_{l\phi}}^2 + \sigma_{A_{l\gamma}}^2(1+b_{\phi,\gamma})^2\right]}$$

1325

$$1326 \qquad R_0 = c\,\bar{\gamma}\,\bar{\phi}_{typ} = c\,e^{\frac{1}{2}\left[(1-\rho_{\gamma,\phi}^2)\sigma_{A_{l\phi}}^2 + \sigma_{A_{l\gamma}}^2(1+b_{\phi,\gamma})^2\right]}$$

1327

1328   Using $\bar{\gamma} = e^{\frac{1}{2}\sigma_{A_{l\gamma}}^2}$ and $\bar{\phi} = e^{\frac{1}{2}\sigma_{A_{l\phi}}^2}$ this simplifies to Equations 14 and 16 of the main text,

$$1329 \qquad \bar{\phi}_{typ} = \bar{\phi}e^{\sigma_{A_{l\gamma}A_{l\phi}}}$$

$$1330 \qquad R_0 = c\,\bar{\gamma}\,\bar{\phi}\,e^{\sigma_{A_{l\gamma}A_{l\phi}}}$$

1331 Further simplification follows from expressing $\bar{\gamma}$, $\bar{\phi}$ and $e^{\sigma_{A_{l\gamma}A_{l\phi}}}$ in terms of variances and
1332 covariances of γ, φ and α.

$$1333 \qquad \bar{\gamma} = e^{\frac{1}{2}\sigma_{A_{l\gamma}}^2}$$

$$1334 \qquad \phi_i = \frac{\varphi_i}{\alpha_i} = e^{(A_{l\varphi,i}-A_{l\alpha,i})} = e^{A_{l\phi,i}}$$

1335 where $A_{l\phi,i} = A_{l\varphi,i} - A_{l\alpha,i}$, which is the breeding value for the logarithm of lifetime infectivity, with

$$1336 \qquad var(A_{l\phi,i}) = \sigma_{A_{l\varphi}}^2 - 2\sigma_{A_{l\varphi}A_{l\alpha}} + \sigma_{A_{l\alpha}}^2$$

1337 From the properties of the log-normal distribution,

$$1338 \qquad \bar{\phi} = e^{\frac{1}{2}\sigma_{A_{l\phi}}^2} = e^{\frac{1}{2}(\sigma_{A_{l\varphi}}^2 - 2\sigma_{A_{l\varphi}A_{l\alpha}} + \sigma_{A_{l\alpha}}^2)}$$

1339 Furthermore,

$$1340 \qquad e^{\sigma_{A_{l\gamma}A_{l\phi}}} = e^{(\sigma_{A_{l\gamma}A_{l\varphi}} - \sigma_{A_{l\gamma}A_{l\alpha}})}$$

1341 Substitution of the expressions for $\bar{\gamma}$, $\bar{\phi}$ and $e^{\sigma_{A_{l\gamma}A_{l\phi}}}$ into $R_0 = c\,\bar{\gamma}\,\bar{\phi}\,e^{\sigma_{A_{l\gamma}A_{l\phi}}}$ yields

$$1342 \qquad R_0 = c\,e^{\frac{1}{2}\sigma_{A_{l\gamma}}^2}\,e^{\frac{1}{2}(\sigma_{A_{l\varphi}}^2 - 2\sigma_{A_{l\varphi}A_{l\alpha}} + \sigma_{A_{l\alpha}}^2)}\,e^{(\sigma_{A_{l\gamma}A_{l\varphi}} - \sigma_{A_{l\gamma}A_{l\alpha}})}$$

$$1343 \qquad R_0 = c\,e^{\frac{1}{2}(\sigma_{A_{l\gamma}}^2 + \sigma_{A_{l\varphi}}^2 + \sigma_{A_{l\alpha}}^2 + 2\sigma_{A_{l\gamma}A_{l\varphi}} - 2\sigma_{A_{l\varphi}A_{l\alpha}} - 2\sigma_{A_{l\gamma}A_{l\alpha}})}$$

$$1344 \qquad R_0 = c\,e^{\frac{1}{2}\sigma_{A_{lR_0}}^2}$$

1345 The right-hand side of this expression is identical to the mean genotypic value for R_0 (Equation 12).

                                                       Appendix 2

   Numerical solution to find the endemic equilibrium prevalence with heterogeneity

1348   To find the endemic prevalence, P, we partition the population into types, i, and numerically solve the
1349   expressions

1350
$$P_i = \frac{\mathcal{R}_{0,i}P}{\mathcal{R}_{0,i}P+1}$$

1351   and

1352
$$\mathcal{R}_{0,i} = \frac{c\gamma_i\bar{\varphi}_{\text{inf}}}{\alpha_i}$$

for P. Here we develop this numerical solution for the case where susceptibility, infectivity and recovery
follow a log-normal distribution, assuming separable mixing (see Appendix 1).

As can be seen from the expression for $\mathcal{R}_{0,i}$, the equilibrium prevalence for a type depends on
both its susceptibility (γ_i) and its recovery rate (α_i). Individuals with above-average susceptibility are
over-represented among the infecteds in the equilibrium, whereas individuals with above-average
recovery are under-represented. Hence, as can be seen from the expression for $\mathcal{R}_{0,i}$, the partitioning into
types should be based on γ_i/α_i. Therefore we define

1360
$$\theta_i = \frac{\gamma_i}{\alpha_i} = e^{A_{\theta,i}}$$

1361
$$A_{\theta_i} = A_{\gamma,i} - A_{\alpha,i}$$

1362
$$\theta \sim Lnorm(\mu_{A_\theta} = 0, \sigma^2_{A_\theta} = \sigma^2_{A_\gamma} - 2\sigma_{A_\gamma A_\alpha} + \sigma^2_{A_\alpha})$$

1363   To numerically solve the two equations given above, we also need $\bar{\varphi}_{\text{inf}}$. The $\bar{\varphi}_{\text{inf}}$ will depends on the
1364 θ_i of the infecteds when infectivity is correlated to susceptibility and/or recovery. Hence, we need the
1365   distribution of $\varphi|\theta$, which follows from

1366
$$\sigma_{A_\theta A_\varphi} = \sigma_{A_\gamma A_\varphi} - \sigma_{A_\alpha A_\varphi}$$

1367
$$b_{\varphi\theta} = \frac{\sigma_{A_\theta A_\varphi}}{\sigma^2_{A_\theta}}$$

1368
$$\rho_{\varphi\theta} = \sigma_{A_\theta A_\varphi}/\sigma_{A_\theta}\sigma_{A_\varphi}$$

1369
$$E(A_\varphi|A_\theta) = b_{\varphi\theta}A_\theta = \mu_{\varphi|\theta}$$

1370
$$\text{var}(A_\varphi|A_\theta) = (1 - \rho^2_{\varphi\theta})\sigma^2_{A_\varphi} = \sigma^2_{\varphi|\theta}$$

1371   so that

1372
$$\varphi|\theta \sim Lnorm(\mu_{\varphi|\theta} = b_{\varphi\theta}A_\theta, \sigma^2_{\varphi|\theta} = (1 - \rho^2_{\varphi\theta})\sigma^2_{A_\varphi})$$

1373   From the log-normal distribution:

1374
$$E(\varphi|\theta) = e^{\mu_{\varphi|\theta} + \frac{1}{2}\sigma^2_{\varphi|\theta}}$$

1375   Hence, we can partitioning θ into classes i, with

1376
$$\mathcal{R}_{0,i} = c\,\theta_i\,\bar{\varphi}_{\text{inf}}$$

1377
$$\bar{\varphi}_{\text{inf}} = \frac{1}{P}\sum_i f_i\,P_i\,E(\varphi_i|\theta_i)$$

1378
$$P_i = \frac{\mathcal{R}_{0,i}P}{\mathcal{R}_{0,i}P + 1}$$

1379
$$P = \sum_i f_i P_i$$

1380   where f_i is the fraction of individuals of type i, $f_i = N_i/N$, and P_i is the prevalence in type i, $P_i = I_i/N_i$.

1381   The numerical solution follows from iterating on these four equations. An R-code is in Supplementary

1382   Material 1.

1383

1384

1385

Appendix 3 - Methods for simulation of epidemics and validation of prevalence and genotypic value for individual disease status

We simulated endemics according to standard epidemiological theory to validate the numerical solution of the endemic prevalence (Equations 20a,b) and the genotypic values for binary disease status (Equation 24). We considered two compartments of individuals, susceptible individuals (S) and infected (I) individuals, and a so-called stochastic SIS-model where susceptible individuals can become infected, and infected individuals can recover and then immediately become susceptible again (Weiss and Dishon, 1971). For simplicity, we simulated genetic variation in susceptibility only, with $\gamma_i \sim \text{Lognormal}(0, \sigma^2_{A_{l\gamma}})$.

To limit Monte-Carlo error, we simulated a relatively large population of $N = 2{,}000$ genetically unrelated individuals for a total of 300,000 events (infection or recovery). We used a burn-in of 100,000 events before recording data on individual binary disease status. Hence, in the recorded data, the average individual experienced 100 events (50 infections and 50 recoveries).

The endemic was started by infecting a proportion $P_0 = 1\text{-}1/c$ of the individuals, chosen at random. Subsequently, we sampled events (infection or recovery) and the individual involved using Gillespie's algorithm (Gillespie, 1977). For each infected individual, the probability of recovery was proportional to the recovery rate, α. For susceptible individual i the probability of infection was proportional to $c\gamma_i I/N$, I/N denoting the fraction of the population that is infected. Probabilities were accumulated over all individuals and scaled to a sum of 1 by dividing them by their sum. Finally, the specific event was sampled by drawing a random number, say x, from a standard uniform distribution and finding the event and the corresponding individual belonging to the probability interval $[x_l, x_h]$, where $x_l < x < x_h$. The disease status of that individual and I were updated before sampling the next event. The time of each event was not simulated. After 300,000 events, prevalence was calculated as the disease status averaged over the entire population, and also by individual, discarding the burn-in period. The regression coefficient of average individual disease status on G_y was also estimated.

Additive genetic variance in log-susceptibility was $\sigma^2_{A_{l\gamma}} = 0.3^3$. Three scenarios were considered, differing in contact rate: $c = 1.22$ giving $P = 0.2$, $c = 2$ giving $P = 5$ and $c = 5.15$ giving $P = 0.8$. Those combinations of $\sigma^2_{A_{l\gamma}}$, c and P were found by numerically solving Equations 20a&b. The actual prevalences observed in the simulations were equal to these numerical solutions.

References

Gillespie, Daniel T. (1977). "Exact Stochastic Simulation of Coupled Chemical Reactions". The Journal of Physical Chemistry. 81 (25): 2340–2361.

Weiss, G. H., & Dishon, M. (1971). On the asymptotic behavior of the stochastic and deterministic models of an epidemic. Mathematical Biosciences, 11(3-4), 261-265.

Appendix 4. Additive genetic variance in log-normal traits.

We assumed log-normally distributed genotypic values for susceptibility, infectivity and recovery, also resulting in a log-normal distribution for G_{R_0} and for $1 - G_P$. Hence, genetic effects are additive on the log-scale, but taking the exponent introduces some non-additive genetic variance on the actual scale. Here we derive the fraction of the variance that is additive on the actual scale.

Because all genetic effects had a mean of zero on the log-scale, the problem is equivalent to finding the fraction of additive variance in $z = e^x$, where $x \sim N(\mu = 0, \sigma^2)$. From the properties of the log-normal distribution, $E(z) = e^\mu e^{\sigma^2/2}$. With a small change $d\mu$, the mean of z becomes $e^{d\mu} e^{\sigma^2/2}$. Hence, the mean of z changes by an amount $e^{\frac{\sigma^2}{2}}(e^{d\mu} - 1)$. Since $\lim_{d\mu \to 0} e^{d\mu} = 1 + d\mu$, this change corresponds to $e^{\frac{\sigma^2}{2}} d\mu$. Thus the least-squares linear regression coefficient of z on x equals

$$b_{z,x} = e^{\sigma^2/2}.$$

For example, the linear regression coefficient of the genotypic value for prevalence (Equation 26b) on the breeding value for the logarithm of R_0 equals $b_{G_P, A_{lR_0}} = c^{-1} \exp\left(\frac{1}{2}\sigma^2_{A_{lR_0}}\right)$. Thus the additive effect for z equals

$$A_z = e^{\sigma^2/2} x,$$

and additive variance in z equals

$$\sigma^2_{A_z} = \sigma^2 e^{\sigma^2}.$$

The total variance in z follows from the properties of the log-normal distribution,

$$\sigma^2_z = (e^{\sigma^2} - 1)e^{\sigma^2}.$$

The additive fraction of the variance in z, therefore, equals

$$f_{\sigma^2_{A_z}} = \sigma^2 / (e^{\sigma^2} - 1)$$

Figure A3.1 illustrates that the additive fraction of the variance in z approaches 1 when σ^2 goes to zero. For $\sigma^2 = 0.5^2$, ~88% of the variance in z is additive. Variances on the log scale larger than 0.5^2 are unrealistic (see main text). This indicates that at least 88% of the genetic variance in susceptibility, infectivity, recovery, R_0 and prevalence is additive when they follow a log-normal distribution.

1450

1451

1452 **Figure A4.1** – The additive fraction of the variance in traits following a log-normal distribution. sdx

1453 denotes the standard deviation on the normal scale.

1454

Appendix 5. Breeding value for individual disease status *vs.* breeding value for prevalence, without genetic variation in infectivity

Without genetic variation in infectivity we have $\varphi_i = \varphi = 1$, because the scale is included in the effective contact rate c. From Equation 24, the genotypic value for individual binary disease status is,

$$G_{y,i} = \frac{\mathcal{R}_{0,i} P}{\mathcal{R}_{0,i} P + 1}$$

where, from Equation 20b,

$$\mathcal{R}_{0,i} = c\gamma_i / \alpha_i$$

From Equation 26a, the genotypic value for prevalence is,

$$G_{P,i} = 1 - \frac{1}{G_{R_0,i}}$$

where, from Equation 8,

$$G_{R_0,i} = c\gamma_i / \alpha_i$$

Hence, without genetic variation in infectivity, $\mathcal{R}_{0,i}$ and $G_{R_0,i}$ are identical, and we will use the symbol $G_{R_0,i}$ in the following.

The linear approximation of the relationship between G_y and G_P follows from a comparison of their first derivatives with respect to G_{R_0},

$$\frac{dG_p}{dG_{R_0}} = \frac{1}{G_{R_0}^2}$$

$$\frac{dG_y}{dG_{R_0}} = \frac{P(G_{R_0} P + 1) - G_{R_0} P^2}{(G_{R_0} P + 1)^2}$$

$$= \frac{P}{(G_{R_0} P + 1)^2}$$

Substituting Equation 3, assuming limited heterogeneity, yields

$$\frac{dG_y}{dG_{R_0}} = \frac{G_{R_0} - 1}{G_{R_0}^3}$$

Hence,

$$\frac{dG_y}{dG_{R_0}} \bigg/ \frac{dG_p}{dG_{R_0}} = \frac{G_{R_0}}{G_{R_0} - 1} = P$$

Therefore, for a small change in an individual's genotypic value for R_0, the change in its genotypic value for binary disease status is only a fraction P of the change in its genotypic value for prevalence,

$$dG_y / dG_p = P$$

1486   Hence, when expressed relative to their mean, G_y and G_p differ approximately by a factor P (see also

1487   Figure 4 in Bijma 2020). This result is approximate, because the true relationship is non-linear and the

1488   expression $P_{eq} = 1 - 1/R_0$ is approximate with variation among individuals. For realistic magnitudes

1489   of the genetic variance, however, the non-linearity is limited. Note that the above derivation does not

1490   require the assumption of a log-normal distribution of susceptibility and recovery.

1491   So far, this appendix has considered genotypic values. However, the factor P also applies on the

1492   level of the breeding values, which can be shown as follows. The above derivation is based on

1493   derivatives with respect to genotypic value for R_0, say dx/dG_{R_0}, where x is G_y or G_P. Since the

1494   relationship between P an y arises entirely via G_{R_0}, we can translate the results to the breeding values

1495   using

1496
$$\frac{dx}{dA_{lR_0}} = \frac{dx}{dG_{R_0}} \frac{dG_{R_0}}{dA_{lR_0}}$$

1497   The latter, $\frac{dG_{R_0}}{dA_{lR_0}}$, is the same for x is G_y or x is G_P, so that the ratio dA_y/dA_p is the same as dG_y / dG_p.

1498   Hence, we also find

1499
$$dA_y / dA_p = P.$$

1500

1501

1502

 Appendix 6 Methods for observed response to selection

First a base population was generated of $N = 4000$ unrelated individuals, with genetic variation in

susceptibility only. No distinction was made between males and females. For each individual, breeding

values for the logarithm of susceptibility were sampled from $A_{l\gamma} \sim N(0, 0.3^2)$, and individual

susceptibility was calculated as $\gamma_i = e^{A_{l\gamma,i}}$. The expected prevalence for the base generation was

calculated as $P_0 = 1 - 1/c$, with a c of either 2 or 10, and the initial disease status of base generation

individuals was sampled at random from $Bernoulli(P_0)$.

Next, an endemic was simulated following methods described in Appendix 3, for a total of

15,000 events (sum of infections and recoveries), consisting of a burn-in of 10,000 events and 5,000

recorded events. The 4,000 individuals were ordered based on their mean individual disease status over

the 5,000 recorded events (so based on 1.25 events on average per individual), and the 2000 individuals

with the lowest values were selected as parents of the next generation (corresponding to a selected

proportion of 0.5).

Selected parents were mated at random. Each pair of parents produced two offspring, resulting

in $N = 4,000$ offspring. Offspring inherited the breeding value for the logarithm of susceptibility in a

Mendelian fashion; $A_{l\gamma,\text{offspring}} = \frac{1}{2}A_{l\gamma,parent1} + \frac{1}{2}A_{l\gamma,parent2} + N\left(0, \frac{1}{2}\sigma^2_{A_{l\gamma}}\right)$. The initial disease

status of offspring (*i.e.,* at the start of the burn-in period of their generation) was sampled at random

from $Bernoulli(P_{\text{offspring}})$, where $P_{\text{offspring}}$ denotes the expected prevalence in the offspring

generation, calculated as $P_{\text{offspring}} = \max\left[1 - \frac{1}{c\,e^{\bar{A}_{l\gamma}}}; 0.02\right]$. The 0.02 guaranteed an average of at least

80 infected individuals at the start of the endemic in any generation, also when the expected prevalence

was zero (*i.e*, when $1 - \frac{1}{c\,e^{\bar{A}_{l\gamma}}} \leq 0$). Then an endemic was started, as described above for the base

generation, *etc*. This process was repeated until the number of infected individuals dropped to zero,

implying extinction of the infection.

1526

 Supplementary material

1528

1529 **Supplementary Material 1:** R-code to numerically find the endemic equilibrium prevalence.

1530 See file "Supplementary Material 1 - Numerical Solution Prevalence Heterogeneity.R"

1531

1532

1533 **Supplementary Material 2:** Additional results for the endemic equilibrium prevalence.

1534 See file "Supplementary Material 2 - Effect of heterogeneity on endemic prevalence.xlsx". Note that,
1535 when $c < 2$, prevalence is (much) higher than $1 - 1/c$, while prevalence is only a little lower than $1 -$
1536 $1/c$ when $c > 2$. Hence, for $c > 2$, $P = 1 - 1/c$ is a useful predictor.

1537

1538

1539 **Supplementary Material 3:** Validation of Equation 34

1540 Equation 34 states that, in the absence of genetic variation in infectivity, the breeding value for
1541 individual disease status is given by

$$A_y \approx P(1 - P)A_{lR_0}$$

where P denotes the endemic prevalence and A_{lR_0} the breeding value for the logarithm om R_0. We
validated this expression using stochastic simulation of endemics following standard epidemiological
theory (as outlined in Appendix 3). We simulated a population of $N = 10,000$ individuals, for a total of
500,000 events (sum of infections and recoveries), using the first 100,000 events as burn in. We
considered input values of $\sigma^2_{A_{l\gamma}} = 0.1^2$, 0.3^2 and 0.5^2. We did not simulate genetic variation in the
recovery rate, because this has the identical effect as genetic variation in susceptibility (Equations 21
and 22). We also did not simulate genetic variation in infectivity, since the expression we intend to
validate here refers to the case without genetic variation in infectivity (In other words, the infectivity
component has to be left out of the A_{lR_0} in Equation 34). We considered a prevalence ranging from 0.1
to 0.9, with steps of 0.1, and found the values for the contact rate that correspond to these prevalences
numerically from solving Equations 20a and b (Table S3.1; R-code in Supplementary Material 1).

1554

1555 **Table S3.1** - Contact rates required to find a certain prevalence, for 3 values of $\sigma^2_{A_{l\gamma}}$.

P	0.1^2	0.3^2	0.5^2
0.1	1.108	1.073	1.010
0.2	1.247	1.218	1.163
0.3	1.424	1.404	1.362
0.4	1.665	1.652	1.628
0.5	2.000	2.000	2.000
0.6	2.502	2.523	2.560
0.7	3.340	3.393	3.495
0.8	5.015	5.135	5.374
0.9	10.040	10.364	11.025

1556

1557   By definition, the individual breeding value equals the regression of individual trait value on
1558 additive genetic effects. Validation, therefore, focussed on the comparison of the $P(1-P)$ term in the
1559 above expression for A_y to the regression coefficient of individual disease status on A_{lR_0} $(b_{y,A_{lR_0}})$
1560 estimated from the simulated data. Table S3.2 shows close agreement between these two parameters.

1561

1562 **Table S3.2** - Comparison of estimated regression coefficient (bhat) of individual disease status on
1563 individual breeding value for the logarithm of R_0 with $P(1-P)$. For three levels of genetic variance in
1564 log(R_0). P_desired denotes the desired prevalence; P_realized denotes the realized prevalence obtained
1565 using the contact rates given in Table S3.1.

	v(A_log_R0)			v(A_log_R0)			v(A_log_R0)		
	0.1^2	0.1^2	0.1^2	0.3^2	0.3^2	0.3^2	0.5^2	0.5^2	0.5^2
P_desired	P_realized	bhat	P(1-P)	P_realized	bhat	P(1-P)	P_realized	bhat	P(1-P)
0.10	0.10	0.09	0.09	0.10	0.09	0.09	0.11	0.10	0.10
0.20	0.20	0.16	0.16	0.20	0.16	0.16	0.20	0.16	0.16
0.30	0.30	0.21	0.21	0.31	0.21	0.21	0.30	0.20	0.21
0.40	0.40	0.25	0.24	0.40	0.23	0.24	0.40	0.23	0.24
0.50	0.50	0.25	0.25	0.50	0.24	0.25	0.50	0.24	0.25
0.60	0.60	0.23	0.24	0.60	0.23	0.24	0.60	0.23	0.24
0.70	0.70	0.21	0.21	0.70	0.21	0.21	0.70	0.20	0.21
0.80	0.80	0.16	0.16	0.80	0.16	0.16	0.80	0.15	0.16
0.90	0.90	0.09	0.09	0.90	0.09	0.09	0.90	0.09	0.09

1566
1567
1568