Zebrafish heme oxygenase 1a is necessary for normal development and macrophage migration

Kaiming Luo[1,2], Masahito Ogawa[2], Anita Ayer[2,3], Warwick J Britton[1,4], Roland Stocker[2,3], Kazu

Kikuchi[2,5], Stefan H Oehlers[1,6]*

[1] Tuberculosis Research Program at the Centenary Institute, The University of Sydney,

Camperdown, NSW 2050, Australia

[2] Victor Chang Cardiac Research Institute, Darlinghurst NSW 2010, Australia

[3] The Heart Research Institute, Newtown, NSW 2042, Australia

[4] Department of Clinical Immunology, Royal Prince Alfred Hospital, Camperdown, NSW 2050,

Australia

[5] National Cerebral and Cardiovascular Center, Suita, Osaka 564-8565, Japan

[6] The University of Sydney, Discipline of Infectious Diseases & Immunology and Marie Bashir

Institute, Camperdown, NSW 2006, Australia

Corresponding author email: stefan.oehlers@sydney.edu.au

Abstract

Heme oxygenase function is highly conserved between vertebrates where it plays important roles in

normal embryonic development and controls oxidative stress. Expression of the zebrafish heme

oxygenase 1 genes are known to be responsive to oxidative stress suggesting a conserved

physiological function. Here we generate a knockout allele of zebrafish *hmox1a* and characterize

the effects of *hmox1a* and *hmox1b* loss on embryonic development. We find that loss of *hmox1a* or

hmox1b causes developmental defects in only a minority of embryos, in contrast to *Hmox1* gene

deletions in mice that causes loss of most embryos. Using a tail wound inflammation assay we find

a conserved role for *hmox1a*, but not *hmox1b*, in normal macrophage migration to the wound site.

Together our results indicate zebrafish *hmox1a* has clearly a partitioned role from *hmox1b* that is
more consistent with conserved functions of mammalian Heme oxygenase 1.

Introduction

Heme oxygenases are a highly conserved family of proteins that catalyze the breakdown of heme to
carbon monoxide (CO), ferrous iron and biliverdin [1, 2]. In mammals, there are two isoforms of
heme oxygenase (HMOX): the inducible HMOX1, and the constitutively expressed HMOX2 [3].
Both isoforms contain a conserved 24-amino acid sequence known as the "heme binding pocket",
within which a conserved histidine residue acts as the ligand for heme iron [4, 5]. Heme is used as
substrate and cofactor for both isoforms, but the physiological properties and regulation of the two
proteins are different [4].

Under basal conditions in mammals, HMOX1 is highly expressed in tissues such as spleen, liver,
and bone marrow, while it is almost undetectable in other tissues [6]. Expression of HMOX1 is
induced by multiple stresses including oxidative stress, heme, iron starvation, inflammatory
cytokines, and physical tissue injury [7, 8]. Conversely, mammalian HMOX2 is constitutively and
highly expressed in the brain and testes and does not generally respond to stress conditions [4, 9,
10].

The role of HMOX1 in regulating the immune response has been demonstrated in models of tissue
injury or different diseases such as ischemic lung/liver injury, atherosclerosis, and non-
hemochromatosis liver diseases [11-14]. In 1999, the first case of HMOX1 deficiency was reported
in a young boy. He showed severe inflammatory phenotypes and accelerated atherosclerosis. The
patient died when he was only six years old [15]. This case suggests a critical role of Hmox1 in
protecting humans against inflammation and atherosclerotic vascular disease. Consistent with this,
irradiated mice transplanted with bone marrow cells from *Hmox1$^{-/-}$* mice show enhanced

atherosclerotic lesions together with a greater macrophage content and increased inflammatory

cytokines including monocyte chemotactic protein 1 [16].

The whole genome duplication event in the teleost lineage resulted in two homologs (*hmox1a* and

hmox1b) of human *HMOX1* in zebrafish genome. Both genes are inducible under oxidative stress

conditions. However, *hmox1a* has been shown to be expressed at a significantly higher level during

the early developmental stage and in response to oxidative stress conditions than *hmox1b* [17, 18].

Until now, no *hmox1a* mutant zebrafish allele has been generated to study the function of heme

oxygenase in zebrafish. Using a combination of CRISPR-Cas9 and TALEN mutagenesis

techniques, we have studied the function of zebrafish *hmox1a* in development and leukocyte

biology.

Methods

<u>Zebrafish husbandry</u>

Adult zebrafish were maintained at Centenary Institute and embryos were obtained by natural

spawning followed and raised in E3 media at 28-32°C (Sydney Local Health District AWC

Approvals: 16-037 and 17-036). All work with the mutant allele and the *Tg(mpeg1:gfp)* transgenic

line were performed in a WT TU/AB genetic background, CRISPR work with the

Tg(mfap4:turquoise) transgenic line was performed in a WT AB genetic background.

<u>Generation of a *hmox1a* mutant allele by TALEN mutagenesis</u>

The CHOPCHOP web tool was used to predict TALEN pairs targeting *hmox1a*. The selected

TALEN information is listed in Supplementary Table 1. Constructs were assembled using Golden

Gate assembly and cloned into pCS2TAL3DDD and pCS2TAL3RRR vectors [19-21].

Genome editing was performed as described [21]. Briefly, TALENs, used to generate mutant

zebrafish, were constructed using the Golden Gate assembly method [19, 20]. mRNAs were

synthesized using the Transcription Kits (Invitrogen) and diluted to 50 ng/μL for microinjection.

Samples were prepared immediately before each injection. mRNAs were injected into zebrafish

embryos at the early one-cell stage to 10% of the volume of the cell (~1 nL). Injected embryos were

raised until they reached adulthood and then outcrossed to wild-type fish. Embryos from these

crosses were used for screening to identify mutants. The resulting mutant line is *hmox1a^vcc42*.

After initial genotyping characterisation, experimental work was carried out on embryos derived

from F3-F5+ generation adults.

Genotyping

For *hmox1a^vcc42* genotyping, a piece of fin was cut off from larval or adult zebrafish for DNA

extraction. PCR was performed with the *hmox1a* genotyping primers described in Supplementary

Table 2 using an annealing temperature of 60°C and an extension time of 30 seconds. PCR products

were digested by *HinfI* restriction enzyme at 37°C for 8-24 h and the digests were resolved by

agarose gel electrophoresis.

Phenotyping

Embryos showing no eyes, severe cyclopisa, small eyes, small trunk somites and heart, reduced

notochord or missing floor plate were described as 'abnormal'. Embryos showing no gross

abnormalities were described as 'normal'.

CRISPR/Cas9 Gene Editing Technique

The gRNA target sites for each gene were designed using https://www.crisprscan.org, gRNA

oligonucleotide sequences are listed in Supplementary Table 3.   The gRNA templates were

amplified by PCR with scaffold reverse primer then were transcribed with HiScribe™ T7 High

Yield RNA Synthesis Kit (NEB) [22]. One cell stage embryos were injected with 1 nL of a pre-

incubated mixture containing 200 ng/μL of the four gRNAs, or eight gRNAs in the double

knockdown, and 2 ng/μL Cas9.

Imaging

Imaging was carried out on Leica M205FA and DM6000B, and Deltavision Elite microscopes as

previously described [23-25].

Drug treatments

Copper-protoporphyrin IX (CuPP) (Frontier) and tin-protoporthyrin IX (SnPP) (Frontier) were

dissolved in 0.05M NaOH and then diluted in pH 8.0 Tris-HCl solution to a stock concentration of

10 mM. All compounds were used at a final concentration of 10 μM. Embryos were treated with

drugs within 3 h of egg fertilization and changed every three days for developmental studies or

immediately after tail wounding and not changed for the duration of the macrophage migration

assays.

Quantitative Real–time PCR

Total RNA was extracted from homogenates using Trizol (Thermofisher) and cDNA was

synthesized with the High Capacity cDNA Synthesis Kit (Applied Biosystems). qPCR reactions

were performed on a LightCycler® 480 System. Gene expression was quantified by the delta–delta

C_T method normalized to host *bact*. Sequences of primers are listed in Supplementary Table 2.

Tail wounding assay

Caudal fin amputation was performed on 4 dpf embryos. Zebrafish larvae were anaesthetized using

tricaine for wounding and imaging experiments. Embryos were cut posterior to the notochord and

then recovered to fresh E3 and kept at 28ºC. Macrophages were imaged at 6 and 24 h post

wounding (hpw) after tail-fin incision of fluorescent reporter zebrafish larvae.

<u>Statistics</u>

All ANOVA and Student's *t* tests as appropriate for the number of comparisons were carried out

using Graphpad Prism. Each data point indicates a single animal unless otherwise stated.

Results

<u>Generation of zebrafish *hmox1a*^{vcc42} mutant allele</u>

The traditional method to establish a mutant animal model is to induce a premature stop codon

caused by nucleotide deletion. In several cases however, phenotypes in knockout mutant animals

using this method have been shown to lack specificity compared to antisense knockdown methods

[26-29]. Recently, it was determined that nonsense-mediated, decay-induced genetic compensation

can cause different phenotypes between knockout mutants and knockdown methods [26]. Animal

models established with in-frame mutation or mutant targeting at the gene promoter area have a

reduced probability of genetic compensation [26]. To minimize the potential effects of genetic

compensation by the paralogs *hmox1b*, *hmox2a*, and *hmox2b*, we generated an in-frame mutation

allele of *hmox1a* targeting the ATG start codon in exon 2 (Fig. 1A).

An in-frame ATG was found in exon 3 at amino acid 37, which was predicted to be active when the

initial ATG was deleted. The translated amino acid fragment from the second ATG start site in the

mutant mRNA sequence was predicted to be non-functional as the heme-binding histidine is absent

in the predicted amino acid sequence (Fig. 1A).

To generate the mutant allele, TALEN-injected embryos were raised to adulthood and three F0 fish

with mutations in somatic tissue were individually outcrossed with wild-type (WT) fish. After

sequencing of F1 fish, we identified an in-frame 26 bp deletion mutant *hmox1a* allele with deletion

of the original start codon, resulting in a predicted mutant protein lacking the functional histidine

(Fig. 1B). The mutant allele was designated as *hmox1a^{vcc42}*, and offspring were propagated for

analysis (Fig. 1C).

<u>Characterization of *hmox1a^{vcc42}* expression</u>

Genomic sequencing revealed the 'AG' splice acceptor site preceding exon 2 was deleted in

hmox1a^{vcc42}. Because there was no splice acceptor attached at the beginning of exon 2 to define the

termination of an intron in front of it, the mutant exon 2 would be treated as a part of a long intron

till the next splice acceptor before exon 3 (Fig. 2A). Thus, we hypothesized that mature mutant

*hmox1a^{vcc42}*mRNA would skip exon 2.

To explore this hypothesis, RNA was isolated from homozygous *hmox1a$^{vcc42/vcc42}$* and WT zebrafish

and cDNA was synthesized via reverse transcription. Primer F1 targeting the start of exon 1 was

used as forward primer and primer R1 targeting the end of exon 7 was used as reverse primer (Fig.

2A). The mutant *hmox1a* exon 1-7 amplicon was slightly smaller than WT as expected from the 41

bp size of exon 2 (Fig. 2B). The exon skipping in *hmox1a* mRNA was further confirmed by

sequencing the cDNA amplicon (Fig. 2C).

To confirm the effect of the *hmox1a^{vcc42}* allele on *hmox1a* transcripts, qPCR analysis was performed

to examine the expression of sections of the *hmox1a* mRNA. Primers spanning the exon 1-2 and 2-3

junctions were chosen for the first analysis (F2/R2; Fig. 2A). As expected, *hmox1a* exon 2 junction-

containing mRNA was almost undetectable in the mutants (Fig. 2D).

Conversely, it was hypothesized that other exon sequences, including exon 1 and exon 3-7, should

still maintained in *hmox1a^{vcc42}* transcripts. To investigate this hypothesis, primers (F3/R3; Fig. 2A)

targeting exon 4 and exon 5 regions were used for qPCR analysis at the same time. Mutant exons 4

and 5 of *hmox1a* mRNA were sustained and not degraded (Fig. 2E), indicating that nonsense-

mediated decay is not triggered by the *hmox1a^{vcc42}* allele.

<u>Morphologic analysis of *hmox1a$^{vcc42/vcc42}$* zebrafish</u>

The percentage of homozygous *hmox1a$^{vcc42/vcc42}$* embryos generated from *hmox1a^{vcc42}* heterozygote

in crosses slightly decreased from 3 to 7 dpf and further gradually decreased to 30 dpf; from 30 dpf

onwards, the percentage of homozygous mutant embryos was stable (Table 1).

To investigate the reason for the loss of homozygous mutants, 3 dpf embryos from sibling matched

hmox1a$^{WT/WT}$ and *hmox1a$^{vcc42/vcc42}$* adult in crosses were collected for morphologic analysis.

Embryos showing no eye, severe cyclopia or small eyes, trunk somites and heart, reduced

notochord and missing floor plate were described as 'abnormal' (Fig. 3A). Embryos that showed no

gross abnormalities were described as 'normal'. While WT embryos did not display abnormal

morphology (0%, n= 368), around 20% of *hmox1a$^{vcc42/vcc42}$* embryos displayed an abnormal

phenotype with shorter length at 3 dpf (Fig. 3B). The percentage of dead *hmox1a$^{vcc42/vcc42}$* embryos

at 11 dpf correlated with the proportion of abnormal *hmox1a$^{vcc42/vcc42}$* embryos at 3 dpf (Fig. 3C,

Table 2). This analysis suggests that a minority of zebrafish *hmox1a$^{vcc42/vcc42}$* embryos undergo an

aberrant morphogenetic process and are unable to survive embryogenesis.

It is known that embryo morphogenetic development is not only affected by gene expression but

also depends on maternal factors that are required for processes prior to the activation of the zygotic

genome. To identify whether *hmox1a* is a maternal factor, embryos from *hmox1a$^{vcc42/vcc42}$* females

crossed with *hmox1a$^{+/vcc42}$* males and *hmox1a$^{+/vcc42}$* females crossed with *hmox1a$^{vcc42/vcc42}$* males

were screened separately. No differences were observed between these crosses (Fig. 3D),

suggesting that the morphological phenotype was independent of the paternal genotype.

The 20% of *Hmox1*-deficient mice that survive until adulthood display smaller size and less body

weight [30]. To investigate whether zebrafish *hmox1a* deficiency also affects adult morphology,

hmox1a$^{vcc42/vcc42}$ adult zebrafish were assessed for morphology, body length and weight. In contrast

to *Hmox1$^{-/-}$* mice, *hmox1a$^{vcc42/vcc42}$* adult zebrafish developed without gross abnormalities compared

to their WT clutch mates (Fig. 3E). Body length and weight was also comparable between

hmox1a$^{vcc42/vcc42}$ and WT clutch mate adults (Fig. 3F). These results indicate zebrafish *hmox1a* may

have a different impact on embryogenesis and development compared to mouse *Hmox1*, or that

genetic compensation by another Hmox family member plays an important compensatory role

during the development of *hmox1a$^{vcc42/vcc42}$* zebrafish.

<u>Zebrafish *hmox1b* partially compensates for loss of *hmox1a* during embryonic development</u>

As a consequence of the whole genome duplication event in zebrafish, there are additional paralogs

hmox1b, *hmox2a*, and *hmox2b* that may increase the survival of *hmox1a$^{vcc42/vcc42}$* zebrafish

compared to mice, where *Hmox2* is the only related gene to *Hmox1*.

We first examined the expression of *hmox1b*, *hmox2a* and *hmox2b* in *hmox1a$^{vcc42/vcc42}$* zebrafish.

Significant upregulation of *hmox1b* was detected in phenotypically normal *hmox1a$^{vcc42/vcc42}$*

embryos compared to WT embryos and *hmox1b* expression was even higher in abnormal

hmox1a$^{vcc42/vcc42}$ embryos than the other phenotypically normal embryos of either genotype (Fig.

4A). The zebrafish *hmox2* genes, *hmox2a* and *hmox2b*, were expressed at a slightly higher level in

abnormal *hmox1a$^{vcc42/vcc42}$* embryos relative to WT embryos, and there was no change in the

expression of either *hmox2a* or *hmox2b* in normal homozygous *hmox1a$^{vcc42/vcc42}$* embryos (Fig. 4A).

Upregulation of *hmox1b*, but not of *hmox2a* or *hmox2b*, was also seen in *hmox1a* crispants (Fig.

4B).

Knockdown of *hmox1a* caused the appearance of visually similar developmental defects to those

seen in homozygous *hmox1a$^{vcc42/vcc42}$* embryos however there was a basal rate of pre-3 dpf embryo

death caused by the microinjection that obscured any statistically significant difference in rate of

developmental defects in *hmox1a* crispants (Fig. 4C, Table 2). Knockdown of *hmox1b* also caused

some developmental defects similar to the loss of *hmox1a* but at a statistically similar rate to the

embryo death in scrambled controls and overall developmental defects in *hmox1a* crispants (Fig.

4C, Table 2). Double knockdown of *hmox1a* and *hmox1b* by CRISPR-Cas9 increased the rate of

developmental defects compared to scramble control embryos across 3-7 dpf and increased the rate

of developmental defects compared to *hmox1b* knockdown embryos at 3 dpf (Fig. 4C, Table 2).

Treatment of developing WT embryos with the Hmox inhibitor tin protoporphyrin (SnPP) caused

increased mortality after 9 dpf compared to copper protoporphyrin (CuPP) controls (Fig. 4D, Table

2). Together, these results demonstrate a conserved role for zebrafish Hmox1a and Hmox1b in

embryonic development.

<u>Conservation of Hmox1a regulation of macrophage migration</u>

To investigate the role of *hmox1a* in macrophage biology, we crossed the *hmox1a^{vcc42}* allele into the

Tg(mpeg1:GFP)^{vcc7} background.

As *Hmox1* has been reported to be essential for hematopoiesis [31, 32], the total fluorescent area of

macrophages was measured by fluorescence microscopy in 4 dpf *Tg(mpeg1:GFP)^{vcc7}* zebrafish

embryos, where macrophages are marked by GFP. No difference was observed between the WT

control and *hmox1a$^{vcc42/vcc42}$* larvae (Fig. 5A).

Chemokine production plays a crucial role in the recruitment of leukocytes to sites of inflammation.

The monocyte chemoattractant protein-1 (MCP-1) and its receptor CCR2 are regulated by HMOX1

in mouse models of mycobacterial infection [33, 34]. To determine if the relationship between

Hmox1 and macrophage chemoattractant production is conserved in zebrafish, we performed tail

wounding of zebrafish embryos (Fig. 5B). The expression of the chemokine *mcp-1* and receptor

ccr2 were examined at 6 h post wounding (hpw) by qPCR. Expression of *mcp-1* was reduced while

ccr2 expression was increased in *hmox1a$^{vcc42/vcc42}$* larvae compared to WT littermates (Fig. 5B).

To investigate the functional consequence of reduced *mcp-1* expression, we quantified macrophage

migration to the wound. There was decreased macrophage accumulation at wound area in

hmox1a$^{vcc42/vcc42}$ homozygous larvae compared to their WT littermates at 24 hours post wounding

(Fig. 5D).

CRISPR/Cas9 technology was used to knockdown *hmox1a* expression in *Tg(mfap4:turquiose)^{xt12}*

zebrafish embryos, where the *mfap4* promoter drives macrophage-specific expression comparable

to the *mpeg1* promoter [35]. The number of wound-associated *mfap4*:Turquiose$^+$ cells was

significantly reduced in *hmox1a* crispants compared to scramble control larvae (Fig. 5E).

To confirm these genetic results, the HMOX inhibitor, SnPP, was used to treat tail wounded WT

Tg(mfap4:turquiose)^{xt12} larvae. SnPP decreased macrophage number at the wound site compared to

CuPP-treated controls (Fig. 5F).

As *hmox1b* knockdown had produced some of the developmental defects seen in *hmox1a* depletion,

we examined the effect of *hmox1b* knockdown on macrophage migration to a wound. Knockdown

of *hmox1b* did not affect macrophage migration (Fig. 5G).

These results demonstrate depletion of *hmox1a*, but not *hmox1b*, decreases macrophage migration

in zebrafish.

Discussion

We report the generation and characterization of a zebrafish with a *hmox1a* mutant allele with approximately 20% lethality in homozygotes during embryonic development. Morphologic analysis of *hmox1a$^{vcc42/vcc42}$* embryos revealed the occurrence of developmental disorders from 3 dpf that progressed to lethality within 14 dpf. These developmental defects were reproduced in double *hmox1a* and *hmox1b* crispants, and mortality was phenocopied by Hmox1 inhibitor treatment. Using live imaging and gene expression analyses we demonstrate conservation of the link between *hmox1a* and macrophage migration in zebrafish. This *hmox1a^{vcc42}* allele provides a platform to investigate the role of *hmox1a* in response to different environmental stresses in zebrafish.

The first *Hmox1$^{-/-}$* mouse model was established by Poss *et al* in 1997 with total deletion of exons 3 and 4, and partial deletion of exon 5. About 80% of *Hmox1$^{-/-}$* mice succumbed to prenatal lethality and premature mortality [30]. Our *hmox1a^{vcc42}* allele, which only contains deletion of exon 2, caused disorders of development and premature mortality in about 20% of *hmox1a$^{vcc42/vcc42}$* embryos. This ~20% mortality rate was similar in *hmox1a* crispants suggesting that the different survival rates between *Hmox1$^{-/-}$* mice and *hmox1a*-depleted zebrafish is not caused by residual peptide translated from the *hmox1a^{vcc42}* transcript. A possible explanation for the difference in survival is the high expression of compensatory *hmox1b,* which exists in zebrafish but not in mice. Our double knockdown experiment where depletion of *hmox1a* and *hmox1b* by CRISPR-Cas9 mutagenesis increased the rate of developmental abnormalities compared to control and *hmox1b*-depleted, but not *hmox1a*-depleted, embryos suggest *hmox1a* has a more important role in zebrafish development than *hmox1b*.

The upregulation of *hmox1b* in *hmox1a$^{vcc42/vcc42}$* embryos might be caused by genetic compensation resulting from the genomic editing in the *hmox1a* gene locus. Although genetic compensation has

been described for a long time [36, 37], the nonsense-mediated decay pathway mechanism has only

been recently described [26, 38]. Mutant alleles that avoid induction of the premature stop codon,

such as the *hmox1a^vcc42* allele, have been reported to reduce the probability of inducing genetic

compensation [39, 40]. We also observed upregulation of *hmox1b* in *hmox1a* crispants. This

suggests the upregulation of *hmox1b* in *hmox1a^vcc42/vcc42* embryos is more likely to be caused by the

stress within embryos secondary to the loss of Hmox1a function. Although we observed some

developmental defects in *hmox1b* crispants, the lack of statistical significance compared to scramble

control embryos and lack of significant additional developmental defects in double knockdown

embryos compared to *hmox1a* knockdown alone suggests a biologically insignificant function of

hmox1b in zebrafish embryogenesis.

About 80% of *hmox1a^vcc42/vcc42* larvae survived until adulthood without gross abnormalities at 90

dpf. This is in accordance with the mouse study where, in the minority of surviving *Hmox1*-

deficient mice, no difference in body weight was reported between *Hmox1*-deficient homozygous

and heterozygous mice until the age of 20 weeks when body size started to be lost in *Hmox1*-

deficient mice [30]. Further morphological analysis of *hmox1a^vcc42/vcc42* zebrafish over 90 dpf is

necessary to investigate the conservation of this effect in zebrafish.

It has been reported that expression of MCP-1, also known as CCL2, was significantly increased in

the liver, spleen, bronchoalveolar lavage fluid (BALF) and serum samples of *M. avium* infected

Hmox1^-/- mice compared to WT controls [34, 41]. However, inhibition of HMOX1 in *M. avium*-

infected RAW 264.7 cells reduced the expression of *MCP-1* compared to untreated cells [34]. The

different results from reported studies may be due to the examination of MCP-1 expression at

different levels since in the two experiments using *Hmox1^-/-* mice, MCP-1 expression was analyzed

at protein level [34, 41], but in the macrophage infection experiment, the gene expression of *MCP-1*

was analyzed [34]. In our study, decreased *mcp-1* transcript levels were detected in tail wounded

hmox1a$^{vcc42/vcc42}$ larvae, which was consistent with the *in vitro* macrophage study [34]. Taken

together, these results suggest that *mcp-1* expression is dependent on *hmox1a* expression in

zebrafish and raise the possibility that macrophage function is altered by increased alternative

activation of macrophages driven by the Ccl2/Ccr2 axis.

Using live imaging in zebrafish embryos, we provide direct evidence linking Hmox1a to the

regulation of macrophage migration. We found a macrophage migration defect in *hmox1a$^{vcc42/vcc42}$*

larvae that was recapitulated in *hmox1a* crispants and SnPP-treated larvae, but not in *hmox1b*

crispants. This was similar to the smaller effect of *hmox1b* depletion on development and provides

further evidence that *hmox1a* provides the most conserved functions expected of heme oxygenase 1

in zebrafish embryos. Further work with loss of function alleles of the zebrafish *hmox* family are

necessary to determine which gene or genes mediate the expected functions of heme oxygenase 1 in

other biological processes.

Acknowledgements

We thank Drs Elinor Hortle and Pradeep Cholan for zebrafish infection and analysis methods

training, Drs Angela Kurz and Kristina Jahn of Sydney Cytometry for assistance with imaging

equipment, the Victor Chang Cardiac Research Institute BioCORE staff for maintenance of

zebrafish, and members of the Tuberculosis Research Program at the Centenary Institute for helpful

discussion.

Funding: This work was supported by a joint Postgraduate Scholarship from the Chinese

Scholarship Council (No. 201608320222) and the University of New South Wales to KL; the

University of Sydney Fellowship G197581, NSW Ministry of Health under the NSW Health Early-

Mid Career Fellowships Scheme H18/31086 to SHO; the NHMRC Centre of Research Excellence

in Tuberculosis Control (APP1153493) to WJB; NHMRC Project Grant (APP1130247) to KK.

References

1.   Fraser, S.T., et al., *Heme Oxygenase-1: A Critical Link between Iron Metabolism,*
367   *Erythropoiesis, and Development.* Adv Hematol, 2011. **2011**: p. 473709.
368   2.   Noguchi, M., T. Yoshida, and G. Kikuchi, *Specific requirement of NADPH-cytochrome c*
369   *reductase for the microsomal heme oxygenase reaction yielding biliverdin IX alpha.* FEBS
370   Lett, 1979. **98**(2): p. 281-4.
371   3.   Cruse, I. and M.D. Maines, *Evidence suggesting that the two forms of heme oxygenase are*
372   *products of different genes.* J Biol Chem, 1988. **263**(7): p. 3348-53.
373   4.   Ayer, A., et al., *Heme oxygenases in cardiovascular health and disease.* Physiol Rev, 2016.
374   **96**(4): p. 1449-508.
375   5.   Dunn, L.L., et al., *New insights into intracellular locations and functions of heme*
376   *oxygenase-1.* Antioxid Redox Signal, 2014. **20**(11): p. 1723-42.
377   6.   Tenhunen, R., H.S. Marver, and R. Schmid, *Microsomal heme oxygenase. Characterization*
378   *of the enzyme.* J Biol Chem, 1969. **244**(23): p. 6388-94.
379   7.   Ryter, S.W., J. Alam, and A.M. Choi, *Heme oxygenase-1/carbon monoxide: from basic*
380   *science to therapeutic applications.* Physiol Rev, 2006. **86**(2): p. 583-650.
381   8.   Dwyer, B.E., et al., *Transient induction of heme oxygenase after cortical stab wound injury.*
382   Brain Res Mol Brain Res, 1996. **38**(2): p. 251-9.
383   9.   Sun, Y., M.O. Rotenberg, and M.D. Maines, *Developmental expression of heme oxygenase*
384   *isozymes in rat brain. Two HO-2 mRNAs are detected.* J Biol Chem, 1990. **265**(14): p. 8212-
385   7.
386   10.   Maines, M.D., *The heme oxygenase system: a regulator of second messenger gases.* Annu
387   Rev Pharmacol Toxicol, 1997. **37**: p. 517-54.
388   11.   Fujita, T., et al., *Paradoxical rescue from ischemic lung injury by inhaled carbon monoxide*
389   *driven by derepression of fibrinolysis.* Nat Med, 2001. **7**(5): p. 598-604.
390   12.   Devey, L., et al., *Tissue-resident macrophages protect the liver from ischemia reperfusion*
391   *injury via a heme oxygenase-1-dependent mechanism.* Mol Ther, 2009. **17**(1): p. 65-72.
392   13.   Hou, W.H., et al., *Iron increases HMOX1 and decreases hepatitis C viral expression in*
393   *HCV-expressing cells.* World J Gastroenterol, 2009. **15**(36): p. 4499-510.
394   14.   Ono, K., et al., *A promoter variant of the heme oxygenase-1 gene may reduce the incidence*
395   *of ischemic heart disease in Japanese.* Atherosclerosis, 2004. **173**(2): p. 315-9.
396   15.   Yachie, A., et al., *Oxidative stress causes enhanced endothelial cell injury in human heme*
397   *oxygenase-1 deficiency.* J Clin Invest, 1999. **103**(1): p. 129-35.
398   16.   Orozco, L.D., et al., *Heme oxygenase-1 expression in macrophages plays a beneficial role*
399   *in atherosclerosis.* Circ Res, 2007. **100**(12): p. 1703-11.
400   17.   Holowiecki, A., B. O'Shields, and M.J. Jenny, *Characterization of heme oxygenase and*
401   *biliverdin reductase gene expression in zebrafish (Danio rerio): Basal expression and*
402   *response to pro-oxidant exposures.* Toxicol Appl Pharmacol, 2016. **311**: p. 74-87.
403   18.   Sonnack, L., et al., *Concentration dependent transcriptome responses of zebrafish embryos*
404   *after exposure to cadmium, cobalt and copper.* Comp Biochem Physiol Part D Genomics
405   Proteomics, 2017. **24**: p. 29-40.
406   19.   Cermak, T., et al., *Efficient design and assembly of custom TALEN and other TAL effector-*
407   *based constructs for DNA targeting.* Nucleic Acids Res, 2011. **39**(12): p. e82.
408   20.   Sakuma, T., et al., *Efficient TALEN construction and evaluation methods for human cell and*
409   *animal applications.* Genes Cells, 2013. **18**(4): p. 315-26.
410   21.   Sugimoto, K., et al., *Zebrafish FOXP3 is required for the maintenance of immune tolerance.*
411   Dev Comp Immunol, 2017. **73**: p. 156-162.
412   22.   Wu, R.S., et al., *A Rapid Method for Directed Gene Knockout for Screening in G0*
413   *Zebrafish.* Dev Cell, 2018. **46**(1): p. 112-125 e4.

23.   Black, H.D., et al., *The cyclic nitroxide antioxidant 4-methoxy-TEMPO decreases*
mycobacterial burden in vivo through host and bacterial targets. Free Radic Biol Med,
2019. **135**: p. 157-166.
24.   Cheng, T., et al., *High content analysis of granuloma histology and neutrophilic*
inflammation in adult zebrafish infected with Mycobacterium marinum. Micron, 2020. **129**:
p. 102782.
25.   Matty, M.A., S.H. Oehlers, and D.M. Tobin, *Live Imaging of Host-Pathogen Interactions in*
Zebrafish Larvae. Methods Mol Biol, 2016. **1451**: p. 207-23.
26.   Rossi, A., et al., *Genetic compensation induced by deleterious mutations but not gene*
knockdowns. Nature, 2015. **524**(7564): p. 230-3.
27.   Kok, F.O., et al., *Reverse genetic screening reveals poor correlation between morpholino-*
induced and mutant phenotypes in zebrafish. Dev Cell, 2015. **32**(1): p. 97-108.
28.   Kurreck, J., *Antisense technologies. Improvement through novel chemical modifications.* Eur
J Biochem, 2003. **270**(8): p. 1628-44.
29.   Prakash, T.P., *An overview of sugar-modified oligonucleotides for antisense therapeutics.*
Chem Biodivers, 2011. **8**(9): p. 1616-41.
30.   Poss, K.D. and S. Tonegawa, *Heme oxygenase 1 is required for mammalian iron*
reutilization. Proc Natl Acad Sci U S A, 1997. **94**(20): p. 10919-24.
31.   Abraham, N.G., *Molecular regulation--biological role of heme in hematopoiesis.* Blood
Rev, 1991. **5**(1): p. 19-28.
32.   Szade, K., et al., *Heme oxygenase-1 deficiency triggers exhaustion of hematopoietic stem*
cells. EMBO Rep, 2020. **21**(2): p. e47895.
33.   Deshmane, S.L., et al., *Monocyte chemoattractant protein-1 (MCP-1): an overview.* J
Interferon Cytokine Res, 2009. **29**(6): p. 313-26.
34.   Regev, D., et al., *Heme oxygenase-1 promotes granuloma development and protects against*
dissemination of mycobacteria. Lab Invest, 2012. **92**(11): p. 1541-52.
35.   Walton, E.M., et al., *The Macrophage-Specific Promoter mfap4 Allows Live, Long-Term*
Analysis of Macrophage Behavior during Mycobacterial Infection in Zebrafish. PLoS One,
2015. **10**(10): p. e0138949.
36.   Somlo, M., et al., *The pho1 mutation. A frameshift, and its compensation, producing altered*
forms of physiologically efficient ATPase in yeast mitochondria. Eur J Biochem, 1985.
150(1): p. 89-94.
37.   Carelle-Calmels, N., et al., *Genetic compensation in a human genomic disorder.* N Engl J
Med, 2009. **360**(12): p. 1211-6.
38.   El-Brolosy, M.A. and D.Y.R. Stainier, *Genetic compensation: A phenomenon in search of*
mechanisms. PLoS Genet, 2017. **13**(7): p. e1006780.
39.   Ferec, C. and G.R. Cutting, *Assessing the Disease-Liability of Mutations in CFTR.* Cold
Spring Harb Perspect Med, 2012. **2**(12): p. a009480.
40.   Genschel, J. and H.H. Schmidt, *Mutations in the LMNA gene encoding lamin A/C.* Hum
Mutat, 2000. **16**(6): p. 451-9.
41.   Silva-Gomes, S., et al., *Heme catabolism by heme oxygenase-1 confers host resistance to*
Mycobacterium infection. Infect Immun, 2013. **81**(7): p. 2536-45.
456

Figure legends

Figure 1. Creation of the *hmox1a^{vcc42}* allele. A. Alignment of *HMOX1* homologs. Dark grey

background indicates amino acid residues that are identified in all three proteins (Uniprot). Light

grey background indicates amino acid residues that are similar in all four proteins (Uniprot). The

purple background indicates the functional histidine. Similarity, identity and functional histidine

were predicted by Uniprot database (https://www.uniprot.org/align/). **B.** Partial sequence of the

hmox1a^{vcc42} allele. TALEN mRNA recognition sequences are blue, and the *HinfI* site in the spacer

region is highlighted in green. Exons are indicated by grey-filled boxes with numbers. Black trace

indicates guanine, blue trace indicates cytosine, green trace indicates adenine and red trace indicates

thymine. **C.** Representative results from PCR-based genotyping using DNA extracted from the fins

of *hmox1a^{vcc42}* adult zebrafish followed by *HinfI* digests.

Figure 2. Characterization of the *hmox1a^{vcc42}* allele mRNA transcript. A. Schematic diagram of

the (predicted) splicing process. Primers used for whole-transcript PCR amplification are indicated

as F1 and R1. Primers F2/R2 and F3/R3 were used for qPCR analysis. **B.** DNA gel electrophoresis

visualization of the exon 1-7 amplicon of mutant *hmox1a* mRNA. **C.** Alignment of the mRNA

transcript of the mutant *hmox1a^{vcc42}* allele sequence (bottom) with WT zebrafish *hmox1a* sequence

(top). **D.** Quantification of *hmox1a* exon 2 mRNA by RT-qPCR analysis. **E.** Quantification of

hmox1a^{vcc42} exons 4-5 mRNA by RT-qPCR analysis. Statistical testing by Student's *t* test, error

bars represent one standard deviation, and each data point represents a biological replicate of pooled

embryos.

Figure 3. Morphologic analysis of *hmox1a^{vcc42/vcc42}* embryos. A. Images of 3 dpf embryos from a

hmox1a^{+/vcc42} intercross. Abnormal embryos displayed apparent shorter lengths. Scale bar

represents 0.5 mm. **B.** Quantification of 3 dpf embryo length in WT and *hmox1a^{vcc42/vcc42}* normal

and abnormal embryos. Statistical testing carried out by ANOVA. Data is pooled from 3 WT and 4

hmox1a^{vcc42/vcc42} replicates. **C.** Proportions of abnormal embryo morphology and death during

development. Raw counts are available in Table 2. Statistical testing was performed using Chi-

squared test on raw counts with abnormal and dead embryos treated as one group. **D.** Proportions of

abnormal embryos scored at 3 dpf from female *hmox1a^{+/vcc42}*; male *hmox1a^{vcc42/vcc42}* (F: +/vcc42; M:

vcc42/vcc42) and female *hmox1a*$^{vcc42/vcc42}$; male *hmox1a*$^{+/vcc42}$ (F: vcc42/vcc42; M: +/vcc42)

crosses. Statistical testing was performed using *t* test, and each data point represents a biological

replicate of pooled embryos from an individual crossing. **E.** Representative images of adult (90 dpf)

WT and *hmox1a*$^{vcc42/vcc42}$ zebrafish from an *hmox1a*$^{+/vcc42}$ intercross. Scale bar represents 0.5 cm. **F.**

Quantification of the length (left) and weight (right) of adult (90 dpf) *hmox1a*^{vcc42} zebrafish from an

hmox1a$^{+/vcc42}$ intercross. Statistical testing was performed using *t* test, and error bars represent

SEM.

Figure 4. *hmox1a* and *hmox1b* have non-redundant roles in zebrafish development. A.

Quantification of *hmox1b*, *hmox2a* and *hmox2b* expression in 3 dpf WT and *hmox1a*$^{vcc42/vcc42}$

normal and abnormal embryos. Statistical testing was performed using the one-way ANOVA, error

bars represent one standard deviation, and each data point represents a biological replicate of at

least 10 pooled embryos. **B.** Quantification of *hmox1b*, *hmox2a* and *hmox2b* expression in 3 dpf

scramble control and *hmox1a* crispant normal and abnormal embryos. Statistical testing was

performed using the one-way ANOVA, error bars represent one standard deviation, and each data

point represents a biological replicate of at least 10 pooled embryos. **C.** Quantification of abnormal

embryo morphology and death during double *hmox1a* and *hmox1b* crispant development. Statistical

testing was performed using Chi-squared test on raw counts with abnormal and dead embryos

treated as one group. **D.** Quantification of abnormal embryo morphology and death during

development in embryos treated with SnPP. Raw counts for panels C and D are available in Table

2.

Figure 5. Zebrafish *hmox1a* aids macrophage migration. A. Quantification of total macrophages

of *hmox1a*$^{vcc42/vcc42}$, *hmox1a*$^{+/vcc42}$ and WT larvae. Statistical testing was performed by one-way

ANOVA. **B.** Images represent embryos at 4 dpf (top) and macrophage recruitment to wound site

(bottom) at 6 hpw, red box indicates area measured for macrophage recruitment in panels D-G. **C.**

Relative quantity of *mcp-1* and *ccr2* transcripts measured by qPCR at 6 hpw, and each datapoint

represents biological replicate of a pool of at least 10 embryos. **D.** Quantification of macrophage

accumulation at wound site by normalized fluorescent area. Data are pooled from two biological

replicates. Statistical testing was performed by one-way ANOVA. **E**. Quantification of

macrophages at wound site by fluorescent area in scramble control larvae and *hmox1a* crispants. **F**.

Quantification of macrophages at wound site by fluorescent area in PPIX or SnPP treated zebrafish

larvae. **G.** Quantification of macrophages at wound site by fluorescent area in scramble control

larvae and *hmox1b* crispants. Statistical testing was performed by *t* test unless otherwise stated,

error bars represent one standard deviation, and data are from one experiment representative of at

least two biological replicates unless otherwise stated.

A

Human HMOX1	---[M]ERPQPDSMPQ DLSEALKEATKE VHTQAENAEFMR NFQKGQVTRDGF KLVMASLYHI	57
Zebrafish Hmox1a	[M]DSTKSKAAENTGS DLSEQIKAVTKDSH VRAENTQLMLSY QKGQITQTQYKLLLCSLYEI	60
Zebrafish Hmox1b	-----[M]EPQITDS DLSEQIKAATKDSH VRAENTELMLSY QRGRVTLQQYKLLLCSLYKI	54

* = Heme-binding histidine position, ▽ = Methionine (from in-frame ATG)

Human HMOX1	YVALEEEIERNKE SPVFAPVYFPEEL HRKAALEQDLAF WYGPRWQEVIPYTPAMQRYVKR	117
Zebrafish Hmox1a	YRALEEELDRNADHPAVQ PIYFPQELARLEALGQDLEHFFGPWRKRITVPAATHRYAQR	120
Zebrafish Hmox1b	YEALEEVLDSNASHDAVA PIYFPQELSRLPALRSDLQHFYGRGWREQMQVPDAALAYAQR	114

Human HMOX1	LHEVGRTEPELLVAHAYTRYLGDLSGGQVLKKIAQKALDLPSSGEGLAFFTFPNIASATK	177
Zebrafish Hmox1a	LREIGKSSPELLVAHAYTRYLGDLSGGQVLGKITQKSLGLT-GNKGILFFSFPGVTSANR	179
Zebrafish Hmox1b	LQQVGREHPEYLVAHAYTRYLGDLSGGQVLGRITQRSLGLQ-DGEGLSFFSFPNISSPKL	173

Human HMOX1	FKQLYRSRMNSLEMTPAVRQRVIEEAKTAFLLNIQLFEELQELLTHDTKDQSPSRAPGLR	237
Zebrafish Hmox1a	FKQLYRSRMNSIEFTEQKRQEALDEAVRAFEFNIDVFDDLQKMLSITEEASSDK-----G	234
Zebrafish Hmox1b	FKQLYRSRMNSITLTEEQRRAVLEEANTAFQLNIQVFDELQECVSCVSETEPAV-----R	228

Human HMOX1	QRASNKVQDSAPVETPRGKPPLNT-RSQAPLLRWVLTLSFLVATVAVGLYAM	288
Zebrafish Hmox1a	NEAASQS---------LSKTFSS----SPA-LQFALGVGITLATVGMGVYAF	272
Zebrafish Hmox1b	QRHSPQTQSTAA-DVGVCEPLQGSSSLMPRLLRTLLGVCVLL-AVG-MVYAF	277

▢ Methionine (from original ATG)   * Heme-binding histidine   ▽ Methionine (from in-frame ATG)

■ Fully conserved amino acid residues   ▨ Strongly conserved amino acid residues

▨ Weakly conserved amino acid residues

B

hmox1a

GCAA TCTGTTTTTTTAGAG ATG GACTC CCACCAAAAGC AAAGCGGCAGAGAACA CTGGC
Left TALEN / *Hin*fl / Right TALEN

M D S T S K A A E N T G R

hmox1a^vcc42

26 bp deletion

GCAATCTGTTT------------------------------AAAGCGGCAGAGAACACTGGC

C

*Hin*fl digest

200 bp — Uncleaved
100 bp — Cleaved

WT   +/vcc42   vcc42/vcc42