Genome archaeology of two laboratory *Salmonella enterica enterica* sv Typhimurium

Julie Zaworski[*†], Oyut Dagva[*§], Anthony W Kingston[*‡], Alexey Fomenkov[*], Richard D.

Morgan[*], Lionello Bossi[§], Elisabeth A. Raleigh[*] **orcid.org/0000-0003-0745-9925**

[*]New England Biolabs, Ipswich MA, 01938-2723

[§]University Paris-Saclay, CEA, CNRS, Institute for Integrative Biology of the Cell (I2BC),

91198, Gif-sur-Yvette, France.

[†]julie.zaworski@protonmail.com, Optilian, 92010 Montrouge, France; ORCID orcid.org/0000-

0003-4744-7208

[§]oyutdagva@gmail.com, Université de Lorraine, INRAE, DynAMic, F-54000 Nancy, France;

ORCID orcid.org/0000-0001-6337-7639

[‡]awkingston@gmail.com, Department of Biology and Mathematics, D'Youville College,

Buffalo, NY 14201

Reference numbers: data available in public repositories

ER3625: CP067091-CP067092

SUB8846059: ER3625 full genome

STK005: CP067397-CP067398

SUB8846115: STK005 (reisolate LB5000)

Running Head: *Salmonella* LB5000 and ER3625 genomes

Keywords: *Salmonella* LB5000, prophages, hybrid genome, transduction scars, H antigen, Gifsy

prophages, 6-methyladenine

Corresponding author:

Elisabeth A. Raleigh

Research Department

New England Biolabs

240 County Road

Ipswich, MA 01935

978 380 7238

raleigh@neb.com

Abstract

The Salmonella research community has used strains and bacteriophages over decades,

exchanging useful new isolates among laboratories for study of cell surface antigens, metabolic

pathways and restriction-modification studies. Here we present the sequences of two laboratory

Salmonella strains (STK005, an isolate of LB5000; and its descendant ER3625). In the ancestry

of LB5000, segments of ~15 and ~42 kb were introduced from *Salmonella enterica* sv Abony

103 into *Salmonella enterica* sv Typhimurium LT2, forming strain SD14; this strain is thus a

hybrid of *S. enterica* isolates.   Strains in the SD14 lineage were used to define flagellar antigens

from the 1950s to the 1970s, and to define three restriction-modification systems from the 1960s

to the 1980s. LB5000 was also used as host in phage typing systems used by epidemiologists. In

the age of cheaper and easier sequencing, this resource will provide access to the sequence that

underlies the extensive literature.

44   **INTRODUCTION**

45   Bacteria, phages and conjugal plasmids are in a constant arms race to develop

respectively defense against DNA invasion and methods of evading defenses. Bacteria express

variable surface features to thwart entry (e. g. flagellae, outer membrane proteins, O-antigen

sugar chains) and internal limitation mechanisms to abort establishment (e. g. restriction systems

and abortive-infection suicide factors) (DORON *et al.* 2018; AZAM AND TANJI 2019). Phages and

plasmids may defeat host restriction-modification (RM) systems with antirestriction proteins,

which frequently block type I DNA restriction-modification enzymes (e. g. (ZAVILGELSKY AND

RASTORGUEV 2009; SERFIOTIS-MITSA *et al.* 2010; BALABANOV *et al.* 2012; ROBERTS *et al.* 2012;

PIYA *et al.* 2017; GONZALEZ-MONTES *et al.* 2020). Low-specificity DNA modification enzymes

expressed only for the entering molecule are also found in prophages (MURRAY *et al.* 2018) and

plasmids (FOMENKOV *et al.* 2020).

Classical studies of horizontal exchange processes within and between genera frequently

employed strains in the ancestry of LB5000, particularly for study of cell surface antigens and

restriction systems. Early hybrid serovar ancestors were constructed by transduction from *S.*

enterica sv Abony into *S. enterica* sv Typhimurium. These studies defined flagellar antigen

types, the phenomenon of flagellar phase variation (JOYS AND STOCKER 1965; JOYS AND

STOCKER 1969) and the use of O-antigen as phage receptor (ORNELLAS AND STOCKER 1974;

STOCKER *et al.* 1980). Three restriction systems were identified using this hybrid lineage

(BULLAS AND COLSON 1975). In LB5000 itself, the three restriction systems of strain LT2 were

inactivated (BULLAS AND RYU 1983), allowing use for epidemiologic purposes as a host for

phages in the Anderson phage typing system (SCHMIEGER 1999). Intergeneric conjugal transfer

of chromosomal markers was studied using *Escherichia coli* K-12 derivatives and *Salmonella*

enterica sv Typhimurium lysotypes LT2 and LT7, using F' plasmids and integrated F factors

(SANDERSON 1987). LB5000 and relatives are particularly useful as an intermediate host in

molecular genetic applications, moving constructions between strains and genera (SANDERSON

AND ROTH 1988; SEGALL AND ROTH 1989; CHAUDHURI *et al.* 2009; LERCHE *et al.* 2017; YOON *et*

al. 2018; GRIEWISCH *et al.* 2020; JONES-CARSON *et al.* 2020).

Genome sequences are available for LT2 (MCCLELLAND *et al.* 2001) and LT7

(ZAWORSKI *et al.* 2021), but none within this hybrid lineage.

Here we present annotated sequences of two *S. enterica* sv Typhimurium variant

genomes: STK005 (our isolate of LB5000, from Anca Segall's G251), and ER3625, a derivative

of STK005 built in our lab for studies of HGT events. The LB5000 genome carries scars from

genetic steps in its descent from LT2, including the replacement of two flagellum-encoding

segments of the LT2 genome by segments of *S. enterica* sv Abony, with concomitant loss of

prophage Fels2. Recombination events involving two prophages document intralineage allele

shuffling. Mutations linked to the amino acid requirement of the strain were identified as well.

 MATERIAL AND METHODS

Salmonella **growth**: For each strain, a single colony was grown in RB (10 g soy peptone, 5 g

yeast extract, 5 g NaCl per liter) with the appropriate antibiotics (kanamycin 40 µg/ml,

streptomycin 100 µg/ml) overnight at 37˚C with 250 rpm agitation.

DNA preparation, sequencing and de novo assembly

DNA extraction: STK005 and ER3625 were prepared for PacBio RSII or Sequel sequencing.

gDNA was extracted with the Monarch Genomic DNA purification kit (#T3010 New England

Biolabs; Ipswich, MA, USA). For Oxford Nanopore sequencing of STK005, gDNA was

extracted from 1 ml of overnight culture with the kit Monarch HMW DNA Extraction kit

(T#3060 New England Biolabs; Ipswich, MA, USA) following the manufacturer instructions for

High Molecular Weight DNA extraction from gram-negative bacteria. After the extraction, the

HMW DNA was homogenized following by heat and pipetting before QC. A clean peak > 60 kb

was found on analysis with the Genomic DNA ScreenTape® (Agilent Technology, Santa Clara,

CA).

Library preparation: For the PacBio RSII sequencing platform, STK005 SMRTbell libraries

were constructed from genomic DNA samples sheared to between 10 and 20 kb using the G-

tubes protocol (Covaris; Woburn, MA, USA), end repaired, and ligated to PacBio hairpin

adapters. Incompletely formed SMRTbell templates and linear DNAs were digested with a

combination of Exonuclease III and Exonuclease VII (New England Biolabs; Ipswich, MA,

USA). DNA qualification and quantification were performed using the Qubit fluorimeter

(Invitrogen, Eugene, OR) and 2100 Bioanalyzer (Agilent Technology, Santa Clara, CA). The

SMRTbell library was prepared according to PacBio sample preparation protocol sequenced with

C4-P6 chemistry with a 300 min collection time. A total of 751,460 polymerase reads with N50

length of 17,626 bp was obtained (approximately 360X coverage) before filtering.

ER3625 libraries were sequenced using the PacBio Sequel I platform. Briefly, SMRTbell

libraries were constructed from genomic DNA samples following the PacBio protocol for Sequel

using the kit 100-938-900. DNA qualification and quantification were performed using the Qubit

fluorimeter (Invitrogen, Eugene, OR) and 2100 Bioanalyzer (Agilent Technology, Santa Clara,

CA). The resulting library, of average size 11,640 bp, was prepared for binding following the

PacBio guidelines generated by SMRT Link and run on a Sequel I machine.

The STK005 Nanopore library was prepared with 1.7 µg of HMW DNA following the

Genomic DNA by ligation (SQK-LSK109 protocol) from Oxford Nanopore. The adapter ligation

was performed for 1 h instead of 10 min. The flow cell was primed and loaded with ~100 ng of

library following the same documentation. After a 21h run on a second-use FLO-MIN106 flow

cell with the GridIONxs apparatus, 449K reads were obtained, with 27.39 kb read length

basecalled N50. The basecalling was performed on the fly with the high-accuracy basecalling

model from Guppy (version 3.2.10)

Sequence assembly: The STK005 (PacBio RSII reads) were assembled using RS_HGAP3

program with parameters: minimum subread length 5,000 bp, minimum polymerase quality 0.8,

Minimum polymerase read length 10,000 bp, 5 Mbp genome size and other parameters as default

settings. The three resulting contigs, of size 4,802,499 bp, 103,465 bp and 8,138 bp, were

circularized with Circlator (HUNT *et al.* 2015) with CCS generated with parameters minimum

full passes 3, minimum predicted accuracy 95, minimum read length of insert 1000 bp and the

other parameters as default. The smallest contig was a subset of the chromosome contig.

PacBio polishing for STK005 was done by running the RS_Reseq program twice, with

Minimum subread length 500, minimum polymerase read quality 75 and then 80, and 1500 bp

for Minimum polymerase read length. The final assembly is composed of a 4,796,208 bp

chromosome and a 93,938 bp plasmid.

The STK005 Nanopore reads were corrected and filtered using Canu (KOREN *et al.*

2017), the output corrected reads were then trimmed and assembled with Canu (version 1.4). A

first correction step was run with parameters minReadLength = 10000, stopOnReadQuality=false

and other as default. Then trimming and assembly were run with genomeSize= 5m and other

parameters as default. Two contigs of length 162,546 and 4,863,042 bp corresponding to the

plasmid and chromosome.

In parallel, the reads were mapped on the PacBio de novo assembly using minimap2

(version 2.17) with default parameters (LI 2018). In both cases, no inversion in between the

Gifsy prophages was observed.

PacBio Sequel assembly of ER3625 used the RS_HGAP4 program with parameters: 5

Mb expected genome size, minimum mapped length 5 kb, minimum mapped concordance 70

and other parameters as default settings. The 2 contigs obtained were circularized with Circlator

(version 1.5.5). Polishing was done by running the RS_Reseq program with minimum mapped

length 500, yielded 19 corrections. The final assembly is composed of a 4,796,355 bp

chromosome (coverage 541x) and a 93,939 bp plasmid (2,247x coverage).

Modification motifs: DNA motifs and degree of modification were generated using InterPulse

Duration (IPD) Ratios analysed with RS_Modification_and_Motif_Analysis from PacBio as in

(FLUSBERG *et al.* 2010; CLARK *et al.* 2012).

Genome annotation: Genbank deposits STK005 (chromosome CP067397, plasmid CP067398)

and ER3625 (chromosome CP067091 and plasmid CP067092) genomes were annotated with the

Genbank PGAP pipeline (TATUSOVA *et al.* 2016; HAFT *et al.* 2018). In-process analysis

employed annotations in Geneious (2018 version in steps: Menu Align and Assemble; Align

Whole Genomes; Tab Alignment View; Save as editable copy; Menu Annotate and Predict;

Annotate from Database) using LT2 [NC_003197.2] and Abony [NZ_CP007534.1] and segment

extractions from them as reference genomes in a local database folder.

Data availability

Supplementary material is available at figshare DOI... Table S1 "Pedigree LB5000 STK005" traces

the descent of LB5000 and some relatives through the literature since 1959; with wild ancestor,

donor strain, recipient strain, renaming, construction reference and mode of construction. File S1

"STK005 variants" contains 5 Excel spreadsheets documenting positions at which STK005

sequence differs from the genome sequences representing two ancestral components. For three

genome segments derived from LT2, variant nucleotide positions of LT2 (NC_003197.2) are

listed, with consequences for amino acid changes; for two short segments derived *Salmonella*

enterica serovar Abony SW803 (H1, 15.4 kb and H2, 41.7 kb), all annotated loci are listed as well,

since few positions vary from the Abony reference. The reference segments H1 and H2 were

extracted from *Salmonella enterica subsp. enterica* serovar Abony str. 0014 - NZ_CP007534.1:

for transduction 1 of Table S1, H1--nt 3956473-3942278 (reversed); for transduction 2, H2--nt

4713943-4725447(catenated)1-18493 (the segment crosses the numerical origin, so is catenated).

File S2 "Analysis of 11 kb high-similarity region in the Gifsy prophages" describes and displays

alignments (referred to as "Embedded figs") that resolve allele configurations in Gifsy prophages.

File S3 "STK005 genotype inferences from sequence" describes the recorded genotype; explains

assignment of flagellar genotype; describes the protocol used to generate the variant list found in

File S1; lists protein changes that account for the genotype; and points out some additional possible

changes. File S4 "PCR survey for 42bp insertion in isolates of *S. enterica* sv Typhimurium str

LT2" demonstrates the insertion is present in Gifsy-1 and conserved in most LT2 strains.

All the sequencing data can be found in the BioProject PRJNA605961. The genome

sequences of STK005 and ER3625 are available from GenBank with the accession

numbers (CP067397-CP067398) and (CP067091-CP067092), respectively. All the raw data are

available under the SRA ID SRR13390529, (SRR13404095, SRR13404094, SRR13404093,

SRR13404092, SRR13404091) and SRR13787361 for ER3625, STK005 RSII and STK005

Nanopore respectively. The methylation data are available as supplementary files on NCBI under

the IDs SUPPF_0000003906 and SUPPF_0000003907. Both strains are available from New

England Biolabs.

183   **RESULTS AND DISCUSSION**

Strain engineering scars: *Salmonella enterica* subsp. *enterica* serovar Abony DNA in

LB5000

The engineering history of LB5000 (Figure 1) includes two transductions from *Salmonella*

serovar Abony strain 803 (SPICER AND DATTA 1959) (see details and references in Table S1).

These two transductions left scars in the form of high-divergence segments in the genome

surrounding the flagellar loci H1 and H2 (Figure 2A). In *Salmonella enterica*, both H1 and H2

flagellar antigens are highly variable in wild lineages in the population; in addition, each

individual lineage is capable of switching from expressing its H1 antigen to its H2 or the reverse,

a phenomenon known as phase variation (SIMON *et al.* 1980). This case of phase variation

involves reversible site-specific rearrangement of regulatory signals that enable expression from

only one of two distant loci. In LB5000, both of these loci derive from *S. enterica* sv Abony.

H1, phase 1 flagellin region transduction patch: The H1 phenotype depends on the allelic state

of the gene *fliC*, within the H1 box of Figure 2A. This patch is the result of a first transduction

mediated by P22: recipient strain LT2 SL270 replaced the H1-i 1.2 allele of LT2 with H1-b from

S. Abony SW803 (SPICER AND DATTA 1959; JOYS AND STOCKER 1969) (see Table S1). Allele

H1-b was bequeathed to LB5000 ancestor SD7. The highly divergent interval (LT2 locus_ID

STM1946 (*uvrC*) to STM1961 (*fliD*)) includes two unshared IS insertions (IS3 in STK005,

IS200 in LT2) and encompasses the *fliC* flagellin gene. In contrast, this region in LB5000

(JJB81_10090 to JJB81_10010; nt 2037077 -> 2052486) aligns with only three mismatches to *S.*

Abony (SEEA0014_RS19195 to SEEA0014_RS19270; nt 3941779 -> 3957191); see below and

File S1.

H2, phase 2 flagellin region transduction patch: The H2 phenotype is determined by the

allelic state of the gene *fljB*, a *fliC* homolog. When the adjacent regulatory gene *fljA* is expressed,

it mediates repression of distant H1 expression; when the local DNA segment is inverted, *fljA*

and *fljB* are not expressed, and repression of distant gene *fliC* is lifted. Wild type for this LT2

isolate is H2-1.2; *S.* Abony SW803 was H2-enx. H2-enx was bequeathed to LB5000 ancestor

SD14.

These genetic changes are reflected in a second divergent region of the alignment (H2

black rectangle in Figure 2A; closer view in Figure 2B). Genes present in LT2 but absent in

STK005 include the Fels2 prophage (STM2694 to STM2740) and the flanking genes STM2741

to STM2769. Present but divergent are genes JJB81_13790 to JJB81_13790 when aligned with

LT2 locus_ID STM2679 and STM3779. We infer that this divergent region represents genomic

replacement of 106 kb of *S.* sv Typhimurium LT2 by 41 kb of *S.* sv Abony 803 during the

second transduction (Figure 1, panel A). Confirmation of the inference is illustrated in Figure

2C: 41 kb of our sequence is very similar (3 insertions, 14 mismatches) to a 41 kb segment of

NZ_CP007534 (sequence of *S. enterica* sv Abony strain 0014 (NCTC 6017; ATCC BAA-2162)

(TIMME *et al.* 2013); see File S1 sheet 4). Both flanks of the long deletion show high divergence

from LT2, and the *fljB* region is 100% identical to the sequenced *S.* sv Abony genome. Prophage

Fels-2 is not present in Abony serovar isolates, consistent with the present configuration.

hin gene, STM2772, JJB81_13890: The Hin protein mediates inversion of a segment including

its own gene and a promoter of *fliC* (H2, phase 2) and the *fljA* (repressor of *fliC* (H1, phase 1))

(SIMON *et al.* 1980). The STK005 *hin* homolog (JJB81_13890) is within the H2 transduction

patch. It is similar to the Abony SEEA0014-RS00045 gene except for the 3 single nucleotide

insertions; these insertions are all shared among LT2, STK005 and ER3625. In addition to the

closer identity to *S. enterica* sv Abony, the *hin* gene has suffered an inversion during the descent

of STK005 from LT2 (also found in ER3625). Considering the expected protein activity, this

reorganization is evidence of expected function.

ER3625 engineering and variation from STK005: ER3625 was engineered in our laboratory to

create a non-pathogenic recipient (HOISETH AND STOCKER 1981) with no Mrr restriction, to

pursue work on R-M systems in HGT using the system previously used in our laboratory

(KINGSTON *et al.* 2015; KINGSTON *et al.* 2017). Strain engineering required several steps with a

variety of methods (Figure 1B).

For future work we wish to verify that no off-target changes had occurred during lamda-

Red-mediated genome engineering or accompanying transduction. The *mrr* engineered locus

carries the designed sequence (*npt* flanked by FRT sites replacing the gene). The *aroA* gene was

deleted, but the scar of the FRT site recombination is aberrant. This DNA segment is annotated

by NCBI pipeline as JJB80_05020 (DUF1317 domain-containing protein) and JJB80_05025

(DUF1382 protein family). No SNPs were identified in genes between STK005 and ER3625 for

either the chromosome or the plasmid.

Structural complexity and intralineage prophage recombination

Gifsy-1 and Gifsy-2 are two LT2 prophages located between 2,728,977 and 2,776,819

(STM2584 to STM2636) and between 1,098,231 and 1,143,702 (STM1005 to STM1056)

respectively. These two related prophages are patchily similar to each other; while overall DNA

identity is 44.1%, this ~42 kb region is interspersed with highly divergent segments and

unrelated insertions (Figure 3). Of concern are two long regions of high identity: a ~12 kb stretch

of > 98% identity in the regions expressed early in lytic infection: recombination, regulators

related to CI, CII and Cro of lambda, and replication; and an 8 kb patch of ~ 78% identity at the

other end of prophages.

These long regions with very high identity are of concern for both technical and

biological reasons. First, long repeats are an assembly challenge even with long-read

technologies. Second, recombination within the host may generate new gene combinations that

are then transmissible to new hosts on prophage induction. Extensive mosaicism among tailed

phages is well known (GROSE AND CASJENS 2014); and creation of a new mosaic by

recombinational marker rescue from defective prophages has been noted before (DE PAEPE *et al.*

2014). Evidence of recombination within a laboratory lineage was found in *E. coli* ER2796

(ANTON *et al.* 2015).

Resolution of assembly errors: De novo assembly of the STK005 genome initially yielded

(artifactual) chromosome inversion. On examination of reads aligned with the two Gifsy regions,

many poorly or partially mapped reads and an important number of SNPs were seen, suggesting

imperfect mapping. A reassembly using only long CCS (> 5 kb; 47.7x coverage) was then

created with HGAP3, then circularized with Circlator. This assembly did not show a genomic

inversion relative to LT2. Further polishing to resolve SNPs used RS_reseq with CCS > 500 bp

(220.9x coverage). This time, few reads were mismapped at the Gifsy regions, confirming the

absence of the inversion in STK005 genome.

In parallel, Oxford Nanopore was used to generate long reads that could cover 40-45 kb

regions and reinforce confidence in the structural organization of the assembly. Reads >10 kb

were used to generate a de novo assembly with Canu (version 1.4). This assembly agreed with

the final PacBio assembly, i.e. without inversion of the genome. Mapping of the Nanopore reads

on the PacBio assembly confirmed the structure.

Prophage recombination: This 12 kb region of near identity codes for proteins involved in

homologous recombination, as well as regulation and replication. The recombination proteins are

related to those of the *E. coli* Rac prophage RecET recombination cluster; also present are

homologs of Lambda Orf and Rap, which also affect recombination. RecE and RecT provide

homologous recombination systems that can substitute for the host RecABCD system (LEMIRE *et*

al. 2008; MURPHY 2016). Rap (originally named NinG) facilitates recombination near linear

ends via the Lambda Red pathway (TARKOWSKI *et al.* 2002), acting as a structure-specific

nuclease (SHARPLES *et al.* 2004).

Three segments within this long region posed interpretation problems: assembly error or

recombination during the lineage leading to STK005. Our interpretation is that the reference LT2

sequence is in error in one case, and both reciprocal recombination and gene conversion have

affected this region (see Figure 3 and Files S2 and S4).

First is the appearance of a novel allele of Gifsy-1 *recET*, including a reading-frame-

preserving insertion of 42 nt (14 aa) in the middle of the RecE CDS and 9 SNPs flanking it (File

S2 Embedded fig2). The Gifsy-1 allele of *recET* found here appears to be ancestral: it is also

found in the Gifsy-1 sequence of *S. enterica* sv Typhimurium strain 14028S (Genbank

CP034479.1; File S2 Embedded fig2). (FIGUEROA-BOSSI *et al.* 2001; LEMIRE *et al.* 2007). Strain

14028S was used extensively in laboratory studies due to much more virulent properties in mice

(e.g. (FIGUEROA-BOSSI *et al.* 1997)). PCR tests of 19 LT2 isolates or derivatives confirm the

presence of the insertion in all of them except strains specifically cured of Gifsy-1 (File S4

Embedded fig1 and Embedded table1). A recent, independently-determined LT2 sequence

(CP014051) was obtained using PacBio and Illumina technology. This deposit agrees with our

sequence of STK005 across this region.

Second, there has been a reciprocal exchange of the sequence between Gifsy-1 and Gifsy-

2 prophages (File S2 Embedded fig3A, 3B). This exchange includes the position of the 18-amino

acid (51 bp) indel variation in CDSs encoding a DNA break and rejoin activity illustrated in

Figure 3B above.

Third, gene conversion has resulted in loss of the original Gifsy-2 allele of Rap (NinG),

replaced by Gifsy-1 information without change in the original Gifsy-1 (File S2 Embedded

fig4A, 4B).

Finding evidence of recombination activity within the high-identity region is interesting.

Though normally silent as prophages, both Gifsy-borne LT2 homologs of RecET can be

activated by mutation to mediate recombination in vivo (LEMIRE *et al.* 2008). Action of these

phage recombination systems may promote the mosaic genomic relationships among phages that

has been found (MARTINSOHN *et al.* 2008). The two events found here are compatible with that

role.

Recorded genotype and point mutations

Metabolic markers: LB5000 was originally described as auxotrophic for methionine,

tryptophan and leucine with the genotype *metA22 metE551 trpC2 ilv-452*. We identified

mutations likely responsible for the reported phenotypes; we also found mutations in other genes

involved in amino acid synthesis, see File S3.

Concerning methionine production, changes are found in *metA* (*metA22*) and *metE*

(*metE551*). In *met*A (STM4182 in LT2 and JJB80_21020 in STK005), a 10 nt deletion at the

protein C-terminus truncates it by 9 amino acids. A SNP in *metE* (G877A) leads to G293R

protein substitution. The *trpC2* allele carries nt change G545A leading to G->D amino acid

substitution. For isoleucine and leucine auxotrophies, SNPs were found in several genes. Coding

changes were a G > A mutation at position 418 in *ilvC* (G140S amino acid change) and a C to T

transition in *leuA* at position 424 (D142N).

Considering other amino acid pathways, *proB* and *proC* of proline synthesis have

mutations respectively C > T at position 535 (P179S) and C > T at position 317 (A106V).

However, our auxotrophy tests found that the strain requires methionine, valine/isoleucine and

tryptophan/phenylalanine/tyrosine but can grow without addition of proline in the media (data

not shown). The synthesis of proline may be mediated via *putA*, which has two silent mutations.

Genotype and phenotype of methyltransferases and their associates: DNA methylation

motifs and level of methylation mediated by R-M systems and orphan MTases were analyzed

with the PacBio RSII. At least six methylation motifs are known in ancestral strain *Salmonella*

enterica serovar Typhimurium LT2 ((ROBERTS *et al.* 2015); REBASE Organism number 18099).

Three motifs result from enterobacterial "orphan" methyltransferases (M; M.SenLT2 Dam,

M.SenLT2 Dcm and M.SenLT2IV; (ROBERTS *et al.* 2015)), and three are associated with

restriction-modification (RM) phenomena (M.SenLT2I, M.SenLT2II and M.SenLT2III). Two of

these RM systems have been well characterized. Characterization began with a transfer isolate

SD14 (Stocker lab strain SL1027). M.SenLT2I (known in the literature as the StyLT or StyLTI

RM system (COLSON AND COLSON 1971)) is a Type III enzyme, and M.SenLT2II (StySB or

StyLTII (FULLER-PACE *et al.* 1984)) is Type I. The third system, RM.SenLT2III (StySA), is

known to confer m6A modification (HATTMAN *et al.* 1976) but its protection/restriction

mechanism of action remains unclear.

The LB5000 RM genotype is listed as *hsdL6 hsdSA29 hsdSB121* for StyLT, StySA and

StySB. We identified appropriate mutations consistent with the R^-M^+ phenotype (Table 1 for M

activity; File S3 for genotype discussion). For StyLT and StySB, sense mutations leading to

amino acid changes were identified in the endonuclease genes, as well as silent mutations. Two

G->A changes (position 393 and 2,749 of STM0358) are present in the *res* gene of StyLT, the

second one leading to a D->N amino acid change in the C-terminal domain. For StySB, two G-

>A changes (position 2,769 and 2,473 of STM4526) are present in *hsdR*, one resulting in an R-

>C amino acid change in the C-terminal domain.

The LB5000 StySA R^-M^+ phenotype was reported to be unstable (TSAI *et al.* 1989), and

we found our STK005 isolate to be phenotypically StySA R^-M^-. The genetic position of StySA

was mapped close to and counterclockwise of the *hsdSB* locus (COLSON AND VAN PEL 1974); we

identified a methyltransferase homolog (JJB81_22590) in an appropriate location, but found no

coding change relative to STM4495. The genetic requirements for action of this system are under

study.

In Table 1, we also find partial methylation for the conserved orphan methyltransferase

M.SenLT2IV, as previously observed in three *Salmonella* species (PIRONE-DAVIES *et al.* 2015).

Methylation of m5C by M.Dcm was not determined here, due to technical limitations of Pacific

Bioscience long-read sequencing in detection of m5C (FLUSBERG *et al.* 2010).

ACKNOWLEGDGMENTS

We thank Stanley Maloy and Anca Segall (San Diego State University) for *Salmonella* strains,

phages and advice and training in their use. We thank Chloé Baum et Giron Koetsier for

technical assistance with the Oxford Nanopore sequencing and high molecular weight DNA

extraction.

New England Biolabs provided support for JZ, OD, TK, AF, RM and EAR. The authors declare

no conflicts of interest.

369 Figure 1: Strain pedigree charts. A. LB5000 B. ER3625. Strain names are boxed. To the right of each construction

370 step (arrows) are the methods used; to the left of each box is the genotype designation assigned as a result of the

371 step. In some cases, a genotype marker was discovered during later investigation. The different strain

372 modifications were done by mutagenesis (random by use of NG), transduction (use of phages) or transformation

373 (plasmids with induction). For more details and references on these steps, refer to Table S1.

Figure 2: Recombination scars. Panel A is a Mauve alignment of the STK005 and LT2 chromosomes. Blue stars

represent IS sequences not shared between the sequences. The flagellar regions H1 and H2 are boxed. Light pink

color is for high similarity areas; the white area is a missing segment; and dark red lines dip where mismatches

occur. Panel B is a close-up of the H2 region. Panel C is a Mauve alignment of S. Abony H2 (fla) region with STK005.

In panel B and C, the yellow and green rectangles below the alignment are landmarks, respectively gene STM2679

(JJB80-13795) and STM2780 (JJB80-13930).

Figure 3: Pairwise alignment of the Gifsy-1 and Gifsy-2 prophages of LT2. Panel A: alignment of the whole

prophages. Panel B: close-up of STM2631 and STM1010, revealing a 51 bp indel variation. The identity rows:

green=identical bases; yellow=lower identity; red=very low local identity. The sequences Gifsy-1 and Gifsy-2 are in

black, white and gray: black for identical bases, white for SNPs, gray for insertion/deletion (indel) variation

(corresponding gaps shown with a thin line).

LITERATURE CITED

Anton, B. P., E. F. Mongodin, S. Agrawal, A. Fomenkov, D. R. Byrd *et al.*, 2015 Complete

    Genome Sequence of ER2796, a DNA Methyltransferase-Deficient Strain of *Escherichia*

    coli K-12. PLoS One 10**:** e0127446.

Azam, A. H., and Y. Tanji, 2019 Bacteriophage-host arm race: an update on the mechanism of

    phage resistance in bacteria and revenge of the phage with the perspective for phage

    therapy. Applied Microbiology and Biotechnology 5**:** 6694-6611.

Balabanov, V. P., V. Y. Kotova, G. Y. Kholodii, S. Z. Mindlin and G. B. Zavilgelsky, 2012 A

    novel gene, *ardD*, determines antirestriction activity of the non-conjugative transposon

    Tn5053 and is located antisense within the *tniA* gene. FEMS Microbiology Letters 337**:**

    55-60.

Bullas, L. R., and C. Colson, 1975 DNA restriction and modification systems in *Salmonella*. III.

SP, a *Salmonella potsdam* system allelic to the SB system in S*almonella typhimurium*.

Mol Gen Genet 139**:** 177-188.

Bullas, L. R., and J. I. Ryu, 1983 *Salmonella typhimurium* LT2 strains which are r⁻ m⁺ for all

three chromosomally located systems of DNA restriction and modification. J Bacteriol

156**:** 471-474.

Chaudhuri, R. R., S. E. Peters, S. J. Pleasance, H. Northen, C. Willers *et al.*, 2009

Comprehensive Identification of *Salmonella enterica* Serovar Typhimurium Genes

Required for Infection of BALB/c Mice. PLoS Pathogens 5**:** e1000529.

Clark, T. A., I. A. Murray, R. D. Morgan, A. O. Kislyuk, K. E. Spittle *et al.*, 2012

Characterization of DNA methyltransferase specificities using single-molecule, real-time

DNA sequencing. Nucleic Acids Research 40**:** e29-e29.

Colson, C., and A. M. Colson, 1971 A new *Salmonella typhimurium* DNA host specificity. J Gen

Microbiol 69**:** 345-351.

Colson, C. C., and A. A. Van Pel, 1974 DNA restriction and modification systems in Salmonella.

I. SA and SB, two *Salmonella typhimurium* systems determined by genes with a

chromosomal location comparable to that of the Escherichia coli *hsd* genes. Molecular

Genetics and Genomics 129**:** 325-337.

De Paepe, M., G. Hutinet, O. Son, J. Amarir-Bouhram, S. Schbath *et al.*, 2014 Temperate phages

acquire DNA from defective prophages by relaxed homologous recombination: the role

of Rad52-like recombinases. PLoS genetics 10**:** e1004181.

Doron, S., S. Melamed, G. Ofir, A. Leavitt, A. Lopatina *et al.*, 2018 Systematic discovery of

antiphage defense systems in the microbial pangenome. Science 359.

Figueroa-Bossi, N., S. Uzzau, D. Maloriol and L. Bossi, 2001 Variable assortment of prophages

provides a transferable repertoire of pathogenic determinants in Salmonella. Molecular

Microbiology 39**:** 260-271.

Figueroa-Bossi, N. N., E. E. Coissac, P. P. Netter and L. L. Bossi, 1997 Unsuspected prophage-

like elements in Salmonella typhimurium. Molecular Microbiology 25**:** 161-173.

Flusberg, B. A., D. R. Webster, J. H. Lee, K. J. Travers, E. C. Olivares *et al.*, 2010 Direct

detection of DNA methylation during single-molecule, real-time sequencing. Nat

Methods 7**:** 461-465.

Fomenkov, A., Z. Sun, I. A. Murray, C. Ruse, C. McClung *et al.*, 2020 Plasmid replication-

associated single-strand-specific methyltransferases. Nucleic Acids Research 48**:** 12858-

12873.

Fuller-Pace, F. V., L. R. Bullas, H. Delius and N. E. Murray, 1984 Genetic recombination can

generate altered restriction specificity. Proc. Natl. Acad. Sci. USA 81**:** 6095-6099.

González-Montes, L., I. del Campo, M. P. Garcillán-Barcia, F. de la Cruz and G. Moncalian,

2020 ArdC, a ssDNA-binding protein with a metalloprotease domain, overpasses the

recipient *hsdRMS* restriction system broadening conjugation host range. PLoS genetics

16**:** e1008750-1008722.

Griewisch, K. F., J. G. Pierce and J. R. Elfenbein, 2020 Genetic Determinants of *Salmonella*

Resistance to the Biofilm-Inhibitory Effects of a Synthetic 4-Oxazolidinone Analog.

Applied and Environmental Microbiology 86.

Grose, J. H., and S. R. Casjens, 2014 Understanding the enormous diversity of bacteriophages:

the tailed phages that infect the bacterial family Enterobacteriaceae. Virology 468-470**:**

421-443.

Haft, D. H., M. DiCuccio, A. Badretdin, V. Brover, V. Chetvernin *et al.*, 2018 RefSeq: an update

on prokaryotic genome annotation and curation. Nucleic Acids Res 46**:** D851-D860.

Hattman, S., S. Schlagman, L. Goldstein and M. Frohlich, 1976 *Salmonella typhimurium* SA

host specificity system is based on deoxyribonucleic acid-adenine methylation. J

Bacteriol 127**:** 211-217.

Hoiseth, S. K., and B. A. Stocker, 1981 Aromatic-dependent S*almonella typhimurium* are non-

virulent and effective as live vaccines. Nature 291**:** 238-239.

Hunt, M., N. D. Silva, T. D. Otto, J. Parkhill, J. A. Keane *et al.*, 2015 Circlator: Automated

circularization of genome assemblies using long sequencing reads. Genome Biology 16**:**

2601.

Jones-Carson, J., A. Yahashiri, J.-S. Kim, L. Liu, L. F. Fitzsimmons *et al.*, 2020 Nitric oxide

disrupts bacterial cytokinesis by poisoning purine metabolism. Science Advances 6**:**

eaaz0260.

Joys, T. M., and B. A. Stocker, 1965 Complementation of non-flagellate *Salmonella* mutants. J

Gen Microbiol 41**:** 47-55.

Joys, T. M., and B. A. Stocker, 1969 Recombination in H1, the gene determining the flagellar

antigen-i of *Salmonella typhimurium*; mapping of H1 and fla mutations. J Gen Microbiol

58**:** 267-275.

Kingston, A. W., C. Ponkratz and E. A. Raleigh, 2017 Rpn (YhgA-Like) Proteins of *Escherichia*

coli K-12 and Their Contribution to RecA-Independent Horizontal Transfer. J Bacteriol

199**:** e00787-00716-00717.

Kingston, A. W., C. Roussel-Rossin, C. Dupont and E. A. Raleigh, 2015 Novel *recA*-

Independent Horizontal Gene Transfer in Escherichia coli K-12. PloS one 10**:** e0130813.

Koren, S., B. P. Walenz, K. Berlin, J. R. Miller, N. H. Bergman *et al.*, 2017 Canu: scalable and

accurate long-read assembly via adaptive k-mer weighting and repeat separation.

Genome Res 27**:** 722-736.

Lemire, S., N. Figueroa-Bossi and L. Bossi, 2007 Prophage contribution to *Salmonella* virulence

and diversity, pp. 159-191 in *Horizontal Gene Transfer in the Evolution of Bacterial*

Pathogenesis, edited by H. S. M. Hensel. Cambridge University Press, Cambridge,

England.

Lemire, S., N. Figueroa-Bossi and L. Bossi, 2008 A singular case of prophage complementation

in mutational activation of *recET* orthologs in *Salmonella enterica* serovar Typhimurium.

J Bacteriol 190**:** 6857-6866.

Lerche, M., H. Sandhu, L. Flöckner, M. Högbom and M. Rapp, 2017 Structure and Cooperativity

of the Cytosolic Domain of the CorA Mg2+ Channel from *Escherichia coli.* Structure 25**:**

1175-1186.e1174.

Li, H., 2018 Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics 34**:** 3094-

3100.

Martinsohn, J. T., M. Radman and M. A. Petit, 2008 The λ Red proteins promote efficient

recombination between diverged sequences: implications for bacteriophage genome

mosaicism. PLoS Genet 4**:** e1000065.

McClelland, M., K. E. Sanderson, J. Spieth, S. W. Clifton, P. Latreille *et al.*, 2001 Complete

genome sequence of *Salmonella enterica* serovar Typhimurium LT2. Nature 413**:** 852-

856.

Murphy, K. C., 2016 λ Recombination and Recombineering. EcoSal Plus 7.

Murray, I. A., R. D. Morgan, Y. Luyten, A. Fomenkov, I. R. Correa, Jr. *et al.*, 2018 The non-
specific adenine DNA methyltransferase M.EcoGII. Nucleic Acids Res 46**:** 840-848.

Ornellas, E. P., and B. A. D. Stocker, 1974 Relation of lipopolysaccharide character to P1
sensitivity in *Salmonella typhimurium*. Virology 60**:** 491-502.

Pirone-Davies, C., M. Hoffmann, R. J. Roberts, T. Muruvanda, R. E. Timme *et al.*, 2015
Genome-wide methylation patterns in *Salmonella enterica* Subsp. *enterica* Serovars.
PLoS One 10**:** e0123639.

Piya, D., L. Vara, W. K. Russell, R. Young and J. J. Gill, 2017 The multicomponent
antirestriction system of phage P1 is linked to capsid morphogenesis. Mol Microbiol 105**:**
399-412.

Roberts, G. A., A. S. Stephanou, N. Kanwar, A. Dawson, L. P. Cooper *et al.*, 2012 Exploring the
DNA mimicry of the Ocr protein of phage T7. Nucleic Acids Res 40**:** 8129-8143.

Roberts, R. J., T. Vincze, J. Posfai and D. Macelis, 2015 REBASE--a database for DNA
restriction and modification: enzymes, genes and genomes. Nucleic Acids Res 43**:** D298-
299.

Sanderson, K. E., 1987 F-mediated conjugation, F strains, and Hfr strains of *Salmonella*
typhimurium and *Salmonella abony*, pp. 2406-2412, edited by F. C. Neidhardt, R. Curtiss
III, J. L. Ingraham, E. C. Lin, K. B. Low *et al.* pdfs.semanticscholar.org, Washington,
DC.

Sanderson, K. E., and J. R. Roth, 1988 Linkage map of S*almonella typhimurium*, edition VII.
Microbiological reviews 52**:** 485-532.

Schmieger, H., 1999 Molecular survey of the *Salmonella* phage typing system of Anderson. J
Bacteriol 181**:** 1630-1635.

Segall, A. M., and J. R. Roth, 1989 Recombination between homologies in direct and inverse

orientation in the chromosome of *Salmonella*: intervals which are nonpermissive for

inversion formation. Genetics 122**:** 737-747.

Serfiotis-Mitsa, D., A. P. Herbert, G. A. Roberts, D. C. Soares, J. H. White *et al.*, 2010 The

structure of the KlcA and ArdB proteins reveals a novel fold and antirestriction activity

against Type I DNA restriction systems in vivo but not in vitro. Nucleic Acids Research

38**:** 1723-1737.

Sharples, G. J., F. A. Curtis, P. McGlynn and E. L. Bolt, 2004 Holliday junction binding and

resolution by the Rap structure-specific endonuclease of phage λ. J Mol Biol 340**:** 739-

751.

Simon, M., J. Zieg, M. Silverman, G. Mandel and R. Doolittle, 1980 Phase variation: evolution

of a controlling element. Science 209**:** 1370-1374.

Spicer, C. C., and N. Datta, 1959 Reversion of transduced antigenic characters in *Salmonella*

typhimurium. J Gen Microbiol 20**:** 136-143.

Stocker, B., B. M. Males and W. Takano, 1980 *Salmonella typhimurium* mutants of RfaH-

phenotype: genetics and antibiotic sensitivies. Microbiology 116**:** 17-24.

Tarkowski, T. A., D. Mooney, L. C. Thomason and F. W. Stahl, 2002 Gene products encoded in

the *ninR* region of phage lambda participate in Red-mediated recombination. Genes Cells

7**:** 351-363.

Tatusova, T., M. DiCuccio, A. Badretdin, V. Chetvernin, E. P. Nawrocki *et al.*, 2016 NCBI

prokaryotic genome annotation pipeline. Nucleic Acids Research 44**:** 6614-6624.

Timme, R. E., J. B. Pettengill, M. W. Allard, E. Strain, R. Barrangou *et al.*, 2013 Phylogenetic

diversity of the enteric pathogen *Salmonella enterica* subsp. *enterica* inferred from

genome-wide reference-free SNP characters. Genome biology and evolution 5**:** 2109-

2123.

Tsai, S. P., R. J. Hartin and J. Ryu, 1989 Transformation in restriction-deficient *Salmonella*

typhimurium LT2. J Gen Microbiol 135**:** 2561-2567.

Yoon, W., Y. Yoo, Y. S. Chae, S. H. Kee and B. M. Kim, 2018 Therapeutic advantage of

genetically engineered *Salmonella typhimurium* carrying short hairpin RNA against

inhibin alpha subunit in cancer treatment. Annals of Oncology 29**:** 2010-2017.

Zavilgelsky, G. B., and S. M. Rastorguev, 2009 Antirestriction proteins ArdA and Ocr as

efficient inhibitors of Type I restriction-modification enzymes. Molecular Biology 43**:**

241-248.

Zaworski, J., C. McClung, C. Ruse, P. R. Weigele, R. W. Hendrix *et al.*, 2021 Genome analysis

of *Salmonella enteric*a serovar Typhimurium bacteriophage L, indicator for StySA

(StyLT2III) restriction-modification system action. G3 Genes|Genomes|Genetics 11.

Table 1. DNA modification systems, motifs and degree of modification

SYSTEM[A]	MOTIF[B]	MOD TYPE[C]	% M[D] STK005	% M[D] ER3625
M.SenLBI (StyLT, StyLTI)	5' CAG**A**G 3'	m6A	99.9	99.9
M.SenLBII (StySB)	5' G**A**GNNNNNNRTAYG 3'	m6A	99.1	98.3
	3' CTCNNNNNNY**A**TRC 5'		100	100
SenLBORF22590P (StySA)	5' GATC**A**G 3'	m6A	0.3	0
M.SenLBIV	5' **A**TGCAT 3'	m6A	24.9	38.2
M.SenLBDam	5' G**A**TC 3'	m6A	99.9	99.9
M.SenLBDcm	5' C**C**WGG 3'	(m5C)	(1.0)	(0.3)
SenLBORF23430P	ND			

[a]System: name used in REBASE for MTases of LB5000 (REBASE organism 45877) followed by (names most

encountered in the RM literature). Numeral order and modification motifs for ER3625 (REBASE organism 45876)

are the same, except M.SenER3625 rather than M.SenLB. These systems are descendants of the LT2 systems

(REBASE organism 18099). SenLBORF22590P and SenER3625ORF22595 are present but inactive. They correspond

to M.SenLT2III. Plasmid-borne SenLBORF23430P and Sen362523455 descends from M.SenLT2ORF505, a

homolog of nonspecific MTase M.EcoGIX (Fomenkov supplementary Table 7).

[b]Motif: recognition sequence with modified base in bold. Sites shown once are symmetric, or asymmetric sites

methylated only on one strand; sites shown twice are asymmetric sites methylated on both strands.

[c] Mod Type: m6A=N6-methyladenine; m5C=5 methylcytosine. Dcm (m5C) motif is inferred from the presence

of the *dcm* gene; m5C is not efficiently detected by this approach.

[d]%M: (# motif sites modified (IPD >2, 25X coverage)/ motif occurrences)x100.