Somite morphogenesis is required for axial blood vessel formation

Eric Paulissen, Nicholas J. Palmisano, Joshua Waxman, & Benjamin L. Martin*

Department of Biochemistry and Cell Biology
Stony Brook University
Stony Brook, NY 11794-5215

* Correspondence: benjamin.martin@stonybrook.edu

Summary statement

Retinoic acid induced somite morphogenesis generates a midline cavity that accommodates migrating angioblasts, which form the axial blood vessels.

ABSTRACT

Angioblasts that form the major axial blood vessels of the dorsal aorta and cardinal vein migrate towards the embryonic midline from distant lateral positions. Little is known about what controls the precise timing of angioblast migration and their final destination at the midline. Using zebrafish, we found that midline angioblast migration requires neighboring tissue rearrangements generated by somite morphogenesis. The somitic shape changes cause the adjacent notochord to separate from the underlying endoderm, creating a ventral midline cavity that provides a physical space for the angioblasts to migrate into. The anterior to posterior progression of midline angioblast migration is facilitated by retinoic acid induced anterior to posterior somite maturation and the subsequent progressive opening of the ventral midline cavity. Our work demonstrates a critical role for somite morphogenesis in organizing surrounding tissues to facilitate notochord positioning and angioblast migration, which is ultimately responsible for creating a functional cardiovascular system.

Keywords: Retinoic acid, dorsal aorta, cardinal vein, somite, morphogenesis, ventral midline cavity, notochord endoderm separation, vasculogenesis, angioblasts, mesoderm, zebrafish

INTRODUCTION

Early organismal development relies on a variety of tissues that collectively organize into a functional body plan. In the trunk of the vertebrate embryo, many tissues are differentiating at once, and often interact with one another to influence their definitive structure (McMillen and Holley, 2015). The axial (notochord), paraxial (somites), and lateral (angioblasts and other tissues) mesoderm begins to be specified during gastrulation, and new cells are added to these tissues from posteriorly localized progenitor cells as they expand after gastrulation (Kimelman, 2016; Martin, 2016; Martin and Kimelman, 2012; Row et al., 2016). By the end of embryogenesis, the notochord is a prominent midline tissue of the trunk, and provides structural support to the body and signals that pattern adjacent tissues (Balmer et al., 2016; Glickman et al., 2003; Kimmel et al., 1995). Pairs of somites flank either side of the notochord and give rise to tissues that include the skeletal muscle, tendons, and bone (Tani et al., 2020). Ventral to the notochord and between the somites, the axial vasculature structures of the dorsal aorta and posterior cardinal vein distribute blood throughout the embryo (Hogan and Bautch, 2004; Isogai et al., 2001).

During somitogenesis, newly formed somites undergo a maturation event in which they change their morphology from a cuboidal shape to a chevron shape, and extend in the dorsal-ventral axis (Kimmel et al., 1995; Tlili et al., 2019). Many cellular rearrangements and mechanical stresses contribute to making the final definitive somite shape (Hollway et al., 2007; Leal et al., 2014; Tlili et al., 2019; Yin and Solnica-Krezel, 2007; Yin et al., 2018; Youn and Malacinski, 1981). At the same time that somites are changing shape, endothelial progenitors called angioblasts arise in the lateral plate mesoderm and migrate to the midline of the embryo to form the axial vasculature (Jin et al., 2005). Interestingly, angioblast migration and new somite formation occur in an anterior to posterior progression, with a wave of angioblast migration happening slightly after the wave of somitogenesis (Jin et al., 2005; Kohli et al., 2013; Yabe and Takada, 2016). Despite these two events occurring in close temporal and physical proximity to one another, it is not clear how they influence one another during this developmental stage. While some evidence has shown somite-related defects on blood vessel patterning, much of this research focused on angiogenic sprouting or defects in arterial-venous specification, well after angioblast migration and somitogenesis is completed (Lawson et

al., 2002; Shaw et al., 2006; Therapontos and Vargesson, 2010; Torres-Vázquez et al.,

2004).

Loss of function mutation of the notochord specifying gene *noto* gene results in non-

autonomous migration defects of angioblasts as they move to the midline. (Fouquet et

al., 1997; Helker et al., 2015). In the absence of *noto,* the notochord adopts a somitic

fate (Fouquet et al., 1997; Talbot et al., 1995). The requirement of notochord

specification for angioblast migration suggested that the notochord may act to attract

angioblasts to the midline. This model was further explored when a notochord-derived

secreted factor named *apela* (also known as *toddler/elabela*) was discovered (Chng et

al., 2013; Freyer et al., 2017; Helker et al., 2015; Pauli et al., 2014). This small peptide is

expressed in the notochord and when mutated caused angioblast migration defects.

However, *apela* loss of function also causes defects in mesoderm and endoderm

formation prior to notochord and angioblast specification, via gastrulation defects (Freyer

et al., 2017; Norris et al., 2017; Pauli et al., 2014). Similarly, *noto* loss of function causes

morphological changes that could interfere with angioblast migration, including a broad

expansion of somite tissue near the developing blood vessels (Halpern et al., 1995).

Thus, although it appears that mutations that affect notochord development can disrupt

angioblast migration, the exact mechanism of this effect is not clear.

As new somites form during body axis extension, they secrete all-trans retinoic acid

(RA). RA is a metabolic derivative of vitamin A, and a series of alcohol and aldehyde

dehydrogenases convert Vitamin A into RA in specific locations of the embryo including

newly formed somites (Duester, 2008). RA acts as a morphogen with broad roles during

vertebrate embryogenesis. Some of these include the development of the limb, skeleton,

hindbrain, and heart (Emoto et al., 2005; Heine et al., 1986; Lohnes et al., 1994;

Mendelsohn et al., 1994; Niederreither et al., 2001; Sandell et al., 2007). Although

involved in many processes, RA has a particular influence over the maturation of

mesoderm and somitic tissue (Hamade et al., 2006; Janesick et al., 2018, 2014; Li et al.,

2015). While RA is notably involved in ensuring the bilateral symmetry of the somites, it

is ultimately dispensable for axis elongation in the zebrafish (Berenguer et al., 2018;

Bernheim and Meilhac, 2020; Hamade et al., 2006; Kumar and Duester, 2014). Based

on the timing of somite derived RA signaling activity relative to when angioblasts migrate

to the midline, we speculated it may be involved in midline angioblast migration.

Here we show that RA mediated somite maturation is required for the proper formation

of the axial vasculature through a non-autonomous role in inducing morphological

changes in surrounding tissues. A process we call notochord-endoderm separation

(NES) occurs prior to angioblast migration, wherein the dorsal translocation of the

notochord during development leads to its separation from the endoderm along the

dorsal-ventral axis to generate a transient cavity we refer to as the ventral midline cavity

(VMC). The angioblasts migrate toward and eventually into the VMC after NES. The

induction of NES and the VMC is somite maturation dependent, and a delay or failure in

NES can cause systemic angioblast migration defects. This evidence places somite-

maturation as a critical event that is required for NES and the development of the axial

vasculature.

RESULTS

Retinoic acid promotes convergence of angioblasts to the midline and the

formation of the axial vasculature

To determine if RA signaling impacts angioblast migration, we made time-lapse movies

of *tg(kdrl:GFP)* embryos, which express GFP in angioblasts (Jin et al., 2005). Embryos

were mounted at the 10 somite-stage and their migration was observed over a 3-hour

time period. Wild-type angioblasts migrated to the midline in a manner consistent with

previously described research (Figure 1A, Supplemental Movie 1) (Helker et al., 2015;

Jin et al., 2005; Kohli et al., 2013). To determine what effect the loss of RA would have

on the migrating angioblasts, we generated *aldh1a2* mutants with the *tg(kdrl:GFP)*

angioblast reporter in the background. Time-lapse imaging of these embryos shows

defects in angioblast migration to the midline (Figure 1B, Supplemental Movie 2), which

was also confirmed by in-situ hybridization (Figure S1). Three other methods of

disrupting RA signaling caused the same phenotype. Loss of RA by treatment with N,N-

diethylaminobenzaldehyde (DEAB) (an inhibitor of the Aldh family, including Aldh1a2

(Morgan et al., 2015)), the small molecule retinoic acid receptor (RAR) alpha and

gamma inhibitor (BMS453), or expression of a dominant negative RARa using the

Tg(hsp70l:EGFP-dnHsa.RARA) transgenic line (hereafter referred to as *HS:dnRAR)* all

caused midline migration defects (Skvarca et al., 2019) (Figure S1). Angioblasts in these

embryos show delayed migration with disorganized anterior to posterior processivity. To

determine if addition of RA accelerated angioblast migration as well as anterior to

posterior processivity, we compared the angioblast migration patterns of 10 somite stage

tg(kdrl:GFP) embryos treated with either DMSO (vehicle) or 0.1 µM RA at the tailbud

stage (Figure 1C,1D, respectively). Angioblasts in the DMSO treated embryos migrate

normally, but the angioblasts treated with RA had prematurely initiated migration and

arrived earlier to the midline during development (Figure 1C and Figure 1D, respectively,

Supplemental Movie 3). To determine the rate at which angioblasts reach the midline,

we quantified the percentage of GFP fluorescence at the midline in relation to the total

amount of fluorescence for the both wild-type and *aldh1a2 -/-*. We defined angioblast

fluorescence as reaching the midline if they are settled beneath the notochord. The

aldh1a2 -/- embryos show very low percentage fluorescence at the midline, relative to

total fluorescence, compared to wild-type embryos 4 hours after observation, when

angioblast migration is largely complete (Figure 1E). On the other hand, fluorescence

was higher at the midline in embryos treated with RA than in DMSO treated control

embryos (Figure 1F). This shows that RA is necessary and sufficient for promoting

angioblast migration to the midline. To determine at which stage RA is needed for

angioblasts to migrate to the midline, we administered DEAB at different stages of

development. Inhibition of RA at tailbud stage caused the angioblast migration defect,

but treatment at the 5-somite stage resulted in normal migration (Figure 1H and 1I,

respectively), thus establishing the critical developmental window for RA signaling

regulation of midline angioblast migration.

Figure 1.Retinoic acid is required prior to segmentation for angioblast migration.

 (**A-D**) Dorsal view of *tg(kdrl:eGFP)* embryos with representative images taken every 60 minutes over the course of 180 minutes, in (**A**) untreated (N=12, see Supplemental Movie 1), (**B**) aldh1a2 -/- mutants (N=12, see Supplemental Movie 2), (**C**) DMSO treated (N=10), or 0.1 μM RA treated embryos (N=8, see Supplemental Movie 3). The anterior of the embryo is at the top and red arrows indicate the midline. (**E, F**) Graph of fluorescence intensity as angioblasts arrive at the midline for (**E**) wild-type and *aldh1a2 -/-* and for (**F**) DMSO and 0.1 μM RA. Y-axis indicates percentage of fluorescent intensity at midline relative to the fluorescence intensity of whole embryo. Time is measured on the X-axis. N=8 for all conditions, error bars indicate standard deviation. (**G-I**) time-lapse images of *tg(kdrl:eGFP)* embryos treated with either (**G**) DMSO vehicle (n=10) or (**H**) 20 μM DEAB at the 5 somite stage (n=6), and (**I**) 20 μM DEAB at tailbud stage (n=14). Scale bars, 50 μm.

Retinoic acid is required cell non-autonomously for angioblast migration.

To determine if RA functions cell autonomously to induce angioblast migration, we

utilized the transgenic zebrafish line *Tg(hsp70l:id3-2A-NLS-KikGR),* which carries a heat

shock inducible construct that overexpresses *id3* after a temperature shift (Row et al.,

2018). We previously showed that overexpression of *id3* from this line can cause

transplanted cells from a donor embryo to faithfully adopt an endothelial fate in a wild-

type host embryo. (Row et al., 2018). We utilized this by crossing *Tg(hsp70l:id3-2A-NLS-*

KikGR)), hereafter referred to *HS:id3*, to *HS:dnRAR,* to generate donor embryos

wherein transplanted cells would be targeted to the endothelium and express dominant

negative retinoic acid receptors (RARA). Cells positive for both transgenic constructs

were transplanted into *tg(kdrl:eGFP)* host embryos and these cells exhibited normal

migration to the midline (Figure 3A, Supplemental Movie 4). In addition, we utilized a RA

reporter line, *tg(RDBD,5XUAS:GFP),* to determine which tissues were subject to RA

signaling (Mandal et al., 2013). We performed immunohistochemistry against GFP and

the transcription factor Etv2, which labels the angioblasts (Figure 3B) (Sumanas et al.,

2005). The GFP expression indicates tissues that have been exposed to RA. These

tissues included the adaxial region of the somites, the notochord, and epidermis (Figure

3B, white arrow, white arrowhead, and gold arrow, respectively). The primitive

angioblasts, labeled with anti-Etv2 staining, do not overlap with GFP staining (Figure 3B,

red arrows). Together these results indicate that angioblasts are not receiving an RA

signal and do not require RA signaling for migration when surrounded by a wild-type

environment.

We investigated which tissue was responsible for the angioblast migration defect based

on the observed activity of the reporter. Given its proximity to the midline, we chose to

Figure 2. Retinoic acid signaling is not present in the endothelium and is required in the somites for angioblast migration.

(**A**) Representative images of a time-lapse of *tg(HS:dnRAR)*, *tg(HS:id3)* cells transplanted into a wild-type embryo. Cells labeled in magenta indicate migrating cells. These cells migrate along with the host angioblasts labeled by *tg(kdrl:gfp)*, indicating normal migration patterns in absence of cell-autonomous RA depletion conditions (n=3) (see Supplemental Movie 4). (**B**) DAPI stained section of 13-somite-stage *tg(RDBD,5XUAS:GFP)* embryo, immunostained for GFP and Etv2 (n=12). GFP signal is present in adaxial region of the somite, the notochord, and the epidermis (white arrow, white arrowhead, and gold arrows, respectively). The red arrows indicate areas of Etv2 staining. (**C**) GFP channel of image in (B). (**D**) Etv2 channel of image in (B). Note the lack of overlap between Etv2 staining and GFP staining. (**E**) Representative images of a time-lapse of wild-type cells transplanted into a wild-type notochord. (**F**) Representative images of a time-lapse of *tg(HS:dnRAR)* cells transplanted into a wild-type notochord. (**G**) Representative images of a time-lapse of wild-type cells transplanted into a wild-type somite. (**H**) Representative images of a time-lapse of *tg(HS:dnRAR)* cells transplanted into a wild-type somite. White arrows indicate angioblast migration defects. (**I**) Quantification of midline fluorescence, as a percentage of total, for notochord transplants at 180'. The difference for wild-type and *tg(HS:dnRAR)* were not statistically significant (P=0.254). (**J**) Quantification of midline fluorescence, as a percentage of total, for somite transplants at 180'. The difference for wild-type and *tg(HS:dnRAR)* were statistically significant (P<0.0001). Scale bars, 50 μm.

test whether the notochord or somitic mesoderm contributed to the defect. To test

notochord contribution, we transplanted cells from a wild-type donor to a *tg(kdrl:eGFP)*

host embryo to target the midline progenitors (Row et al., 2016). Host embryos with wild-

type cells transplanted to the notochord displayed normal angioblast migration to the

midline (Figure 2E). Similarly, transplants of *tg(HS:dnRAR)* cells into the notochord

showed normal midline angioblast migration(Figure 2F). In addition, in-situ hybridizations

against a known notochord-secreted angioblast chemoattractant *apela,* as well as the

receptors *aplnra* and *aplnrb,* show little change in RA depleted conditions (Figure

Supplement 2A-2F). This indicates that the lack of RA signaling in notochord cells is not

causing the angioblast migration defects. We then transplanted cells to target the somitic

mesoderm. Wild-type cells transplanted into the somitic mesoderm showed normal

angioblast migration to the midline (Figure 2G). However, when we transplanted

tg(HS:dnRAR) cells into the somitic mesoderm, angioblasts showed migratory defects in

the region of the host embryo in which they were transplanted (Figure 2H).

Quantification of fluorescence percentage at the midline, in the chimeric regions of the

notochord, showed no statistically significant difference compared to controls (Figure 2I).

However, quantification of fluorescence percentage in the chimeric regions of the

somites showed statistical significance between wild-type and *tg(HS:dnRAR)* (Figure

2J). This indicates that the somites are the principal tissue required for RA signaling

mediated midline angioblast migration.

The somitic mesoderm is required for angioblast migration to the midline.

To follow up the somite and notochord targeted transplant results, we examined midline

angioblast migration in embryos where these tissues are absent. Previous studies have

investigated genes linking the notochord to angioblast migration. Mutants of the genes

noto and *apela* were found to be notochord specific genes required for proper angioblast

migration (Cleaver and Krieg, 1998; Fouquet et al., 1997; Helker et al., 2015). The *noto*

gene induces notochord, and the notochord in turn secretes Apela which attracts the

angioblasts to the midline. Noto is required for the formation of the notochord, and in the

absence of Noto, the midline cells adopt a somitic fate (Talbot et al., 1995).

However, it is not clear if midline convergence was delayed or absent in notochord-less

embryos. In *noto* mutants, some angioblasts reach the midline at later developmental

stages, and resolve into a single blood vessel (Fouquet et al., 1997). We injected a *noto*

morpholino that faithfully phenocopies the *noto* mutant into *tg(kdrl:GFP)* embryos

(Ouyang et al., 2009). Control morphant embryos show normal angioblast migration

(Figure 3A), whereas *noto* morphants show slowed angioblast migration (Figure 3B).

However, midline convergence still occurs in both conditions (Figure 3C). This implies

that the notochord is ultimately dispensable for the formation of the VMC and midline

positioning of angioblasts.

Having ruled out the role of notochord tissue, we next investigated whether the somitic

mesoderm is required for NES and the formation of the VMC. Previous studies have

shown that the t-box transcription factor *tbx16* is required for proper somite formation

(Amacher et al., 2002; Goto et al., 2017; Griffin et al., 1998; Kimmel et al., 1989;

Manning and Kimelman, 2015; Row et al., 2011). In the absence of Tbx16, cells that

would normally join the paraxial mesoderm fail to do so, causing a deficiency in the trunk

somitic mesoderm, but leaving the endothelium intact (Thompson et al., 1998). We

utilized *tbx16* mutants to determine whether the lack of somites caused angioblast

migration defects, similar to RA loss of function. We examined midline angioblast

migration in *tbx16* mutant embryos with the *tg(kdrl:GFP)* transgene in the background. In

wild-type sibling embryos, angioblast migration progresses normally to the midline

(Figure 3D). However, in *tbx16* -/- embryos, angioblast migration does not occur, and

they remain in their positions in the lateral plate (Figure 6E). Fluorescence quantification

indicated that angioblasts never reach the midline in any embryo (Figure 3F). To

determine the effect of either notochord or somite loss on the definitive vasculature, we

performed transverse sections of *tg(kdrl:GFP)* embryos injected with control, *noto,* and

tbx16 morpholinos and stained with DAPI. Control morpholino showed normal formation

of the dorsal aorta and common cardinal vein (Figure 3G). Embryos injected with *noto*

morpholino have angioblasts located at the midline that appear to have resolved into one

blood vessel structure (Figure 3H). Embryos injected with *tbx16* morpholinos, however,

show two distinct vessels located on either side of the notochord (Figure 3I). This

indicates that the somites are critical for angioblast convergence at the midline, while the

notochord is ultimately dispensable.

Figure 3. The somitic mesoderm, not the notochord, is required for midline convergence of angioblasts.

(**A**) Time-lapse imaging of *tg(kdrl:eGFP)* embryos injected with control morpholino over a 240' period. (**B**) *tg(kdrl:eGFP)* embryos injected with *noto* morpholino over a 240' period. Red arrows indicate the midline. (**C**) Graph of embryos with midline angioblasts in *noto* and control morphants after 240'. N=15 for both conditions (**D**) Time-lapse imaging of wild-type*, tg(kdrl:eGFP)* embryos over a 240' period. (**E**) Timelapse imaging of *tbx16 -/-, tg(kdrl:eGFP)* embryos over a 240' period. Red arrows indicate the midline. (**F**) Graph of embryos with midline angioblasts in wild-type and *tbx16 -/-* after 240' N =18 for both conditions. (**G**) Section of wild-type*, tg(kdrl:eGFP)* embryos over a 240' period. (**H**) Section of 24 hpf *tg(kdrl:eGFP)* embryos injected *noto* morpholino and stained with DAPI. (**I**) Section of 24 hpf *tg(kdrl:eGFP)* embryos injected *tbx16* morpholino. Blood vessels are labeled in green and nuclei labeled in grey. Scale bars, 50 μm.

Retinoic acid induces a dorsal translocation of the notochord away from the

underlying endoderm.

Given the non-autonomous midline angioblast migration defects in RA loss of function

embryos, we examined if RA manipulation altered the normal development of non-

angioblast midline tissues. We utilized embryos labeled with a cell surface marker,

tg(ubb:lck-mng) (Adikes et al., 2020)*,* and found the relative location of notochord at the

midline changes based on retinoic acid activity. To measure this, embryos were treated

with either DMSO, DEAB, or RA at the tailbud stage. The embryos were then fixed,

deyolked, and imaged in the trunk at roughly the 5^{th} somite (Figure 4A-C). The relative

location of the notochord in DMSO treated embryos was in close proximity to the ventral

underlying endoderm at the 12-somite stage, but that distance increased by the 15-

somite stage (Figure 4A). In DEAB treated embryos, the notochord remains ventrally

localized at both the 12-somite stage and 15-somite stage (Figure 4B). When the same

experiment was done with embryos exposed to exogenous RA, the notochord was

prematurely dorsally localized at the 12-somite stage and extended even further by the

15-somite stage (Figure 4C).

We then determined the dynamics of the notochord displacement in real time. Using DIC

images, we were able to overlay the lateral trunk with the notochord. We then observed

the notochord displacement in a 250' time-lapse for DMSO treated wild-type, *aldh1a2 -/-*

with DMSO, and exogenous RA (Figure 4D, 4E, and 4F, respectively). We quantified this

data by measuring the distance of the ventral aspect of the notochord to the yolk cells of

295 the embryo (yellow brackets). We found that *aldh1a2 -/-* embryos showed delayed dorsal

296 displacement relative to DMSO in both the timing of dorsal movement and the

297 magnitude of that movement (Figure 4G).

299 Previous work showed that as somites mature, they extend in the both the dorsal and

300 ventral axis (Tlili et al., 2019), and based on this we hypothesized this extension was

301 concurrent with the dorsal translocation of the notochord. However, we wanted to rule

302 out the effect of forces on the anterior-most and posterior-most regions of the notochord.

303 The notochord is a rigid structure that is anchored in the tailbud near the notochord

304 primordia, and extends into the head mesoderm medial to the otic vesicle (Kimmel et al.,

305 1995). To eliminate the contribution other tissues could have on NES, we generated

306 trunk explants from embryos containing both *tg(actc1b:gfp)* and *tg(tbxta:kaede)*

307 transgenese that contained roughly 10 somites of the embryo. The 10-somite stage

308 embryos were sectioned near the somite borders of the 1st and 10th somite. We

309 performed live imaging of these trunk explants, focusing on the immature somites of the

310 posterior region (white boxes) (Figure 4H-4J, Supplemental Movie 5). In order to better

311 visualize the notochord, we separated the *tg(actc1b:gfp)* and *tg(tbxta:kaede)* signals into

312 separate channels. This distinguished the notochord reporter *tg(tbxta:kaede)* (Figure

313 4H'-4J'), from the somite reporter *tg(actc1b:gfp)* (Figure 4H"-4J"). The notochord reporter

314 showed clear dorsal movement away from the auto-fluorescent yolk (Figure 4H'-J'). The

315 somites also showed the corresponding expansion along the dorsal-ventral axis (Figure

316 4H"-4J"). Together, this data indicates that the proper development of the midline is both

317 retinoic acid-dependent and requires only factors localized to the trunk of the embryo.

Figure 4: RA signaling promotes dorsal translocation of the notochord

(**A-C**) Fluorescent images of fixed (A) DMSO treated (B) DEAD treated and (C) RA treated *tg(ubb:lck-mng)* embryo trunks at the 12- and 15-somite stage. Yellow brackets indicate the distance between the notochord and underlying yolk cell. (**D-F**) DIC time-lapse images of the notochord and zebrafish trunk. Focal planes of the notochord and somites were overlaid to show the relative location to one another. Yellow brackets indicate the distance of the ventral notochord to the ventral-most portion of the embryo in (**D**) DMSO treated embryos, (**E**) *aldh1a2 -/-* embryos treated with DMSO, or (**F**) 0.1 μM RA treated embryos. (**G***)* Quantification of dorsal translocation of notochord from the ventral tissue. Black lines indicate the median and interquartile range. Wild-type are indicated by brown squares, while *aldh1a2 -/-* and 0.1 μM RA embryos are shown by red circles and blue triangles, respectively. At t=0, DMSO vs. RA is P<.00001. At 150', DMSO is P<.00001 and P=.005 vs. *aldh1a2 -/-* and RA treatment, respectively. At 250', DMSO is P=.0001 and .054 vs. *aldh1a2 -/-* and RA treatment, respectively. (**H-J**) Time-lapse image of a trunk explant from *tg(actc1b:gfp), tg(tbxta:kaede)* embryo at (**H**) 0', (**I**) 60', and (**J**) 120'. (**H'-J'**) Time-lapse image of explant showing *tg(tbxta:kaede)* only. (**H''-J''**) Time-lapse image of explant showing *tg(actc1b:gfp)* only. Time-lapse shows the notochord adjacent the yolk at time 0', but moves dorsally from the yolk at time 120'. Scale bars, 25 μm.

Notochord-endoderm separation facilitates the midline convergence of

angioblasts

We speculated that RA dependent changes at the midline could be the cause of the

angioblast migration defect. In order to test this, we generated a heat-shock inducible

transgenic reporter of F-actin, *tg*(*hsp70l:lifeact-mScarlet*), which labels filamentous actin

throughout the embryo and allows visualization of all cells (Riedl et al., 2008). We

hereafter refer to *tg*(*hsp70l:lifeact-mScarlet)* as *tg(HS:lifeact).* We then crossed this

reporter to the *tg(kdrl:eGFP)* line to label the angioblasts. We treated embryos with

either DMSO (Figure 5A-5C), exogenous RA (Figure 5D and 5E), or DEAB (Figure 5F-

5I). Embryos were sectioned at the region of the 5^{th} somite (Figure 5A-5I). Embryos that

were treated with DMSO showed normal midline angioblast migration patterns from the

12-somite stage, 15-somite stage, and 18-somite stage (Figure 5A-5C). As angioblasts

approach the midline, a gap appears between notochord/hypochord and the underlying

endoderm (Figure 5A). This gap is filled by angioblasts by the 15-somite stage,

separating the notochord from the endoderm (Figure 5B). We call the process of dorsal

translocation of the notochord the notochord-endoderm separation (NES). The transient

opening that forms from NES is referred to as the ventral midline cavity (VMC). As

angioblast migration continues, additional angioblasts occupy the VMC until the 18-

somite stage, when angioblast migration is largely complete for the 5^{th} somite region of

the trunk (Figure 5C).

To determine if activation of retinoic acid is sufficient to accelerate angioblast migration

and NES concurrently, we treated embryos with exogenous RA and sectioned them at

the 12-somite and 15-somite stage. Embryos sectioned at the 12-somite stage showed

premature angioblast migration to the midline and NES, indicating that angioblast

migration and NES are responsive to exogenous RA signal (Figure 5D). Angioblasts

continue to be at the midline at 15-somite stage (Figure 5E). 18-somite stage embryos

were not available as posterior elongation discontinued in prolonged RA exposure, as

previously shown (Martin and Kimelman, 2010). To determine whether angioblast

migration and NES was attenuated in RA depletion conditions embryos were treated

with DEAB and sectioned at the 12-somite, 15-somite, 18-somite, and 21-somite stage

(Figure 5F-5I respectively). At the 12-somite stage, the angioblasts are localized in the

lateral plate mesoderm (Figure 5F). Focusing on the midline of the embryo, we observe

that the notochord remains flush against the endoderm with no visible VMC or NES

(Figure 5G). At the 15-somite stage, we observe some midline movement of the

angioblasts, however the notochord remains flush against the endoderm and no

angioblasts have been able to reach the midline (Figure 5G). During the 18-somite

stage, a small VMC has formed between the notochord and endoderm. The leading

angioblasts have partially migrated into this VMC, however the bulk of the angioblasts

still reside outside of the midline (Figure 5H). By the 21-somite stage, the angioblasts

have largely resolved to the midline, although with some angioblasts still residing outside

the midline (Figure 5I).

Given the pronounced delays in NES formation for anterior regions in RA depleted

embryos, we suspected that NES was a consequence of the known processive somite

maturation that begins in the anterior regions and progresses posteriorly (Stern and

Piatkowska, 2015). We sectioned 15-somite stage *tg*(*HS:lifeact*)*, tg(kdrl:egfp)* embryos

in the anterior and posterior regions corresponding to the 5^{th} and 12^{th} somite (Figure 5J).

In more immature 12th somite, angioblasts reside in their lateral regions and NES is not

pronounced (white dashed lines outline the somites, and yellow brackets show the

distance from the notochord to the ventral endoderm) (Figure 5K). In the 5th somite of

the same embryo, angioblasts reside at the midline and NES is very pronounced,

indicating NES occurs anterior to posteriorly (Figure 5L). The lack of angioblasts at the

12th somite, and low NES, is similar to DEAB treated embryos sectioned at the more

mature 5th somite position (Figure 5M).

To quantify how the dorsal translocation of the notochord was correlated with

angioblasts arriving to the midline, We created a two variable graph to measure both the

distance of the notochord to the ventral endoderm and the percentage of angioblasts

arriving at the midline for a given somite. For this analysis, we used 15-somite stage

embryos sectioned at the 5th somite and 12th somite. We expect that immature somites

in the posterior would have little midline migration of angioblasts and the anterior, with

more mature somites, would have significant midline migration. We measured the

distance in microns on the x-axis and the percentage of midline angioblasts on the y-axis

(Figure 5O-5P). In DMSO treated embryos, embryos sectioned at the 5th somite grouped

together, showing both significant notochord-endoderm distance and angioblasts

residing at the midline (Figure 5O, blue dots in the orange box). Embryos sectioned at

the 12th somite show short notochord-endoderm distance and little angioblast migration

towards the midline (Figure 5O, red squares in grey box). When comparing angioblasts

migration, the 5th somite and 12th somite regions were significantly different ($P > .0001$).

The corresponding sectioning of DEAB treated embryos shows a difference compared

to wild-type in the 5th somite region. Embryos sectioned at the 5th somite have a small

distance from the notochord to endoderm, as well as no angioblast migration to the

midline (Figure 5P, blue dots in orange box). Interestingly, the notochord-endoderm

distance/migration percentage ratio strongly resembled the more immature somites. The

angioblast migration percentage was not statistically significant between the 5th and 12th

somite in RA depletion conditions ($P = .33$). This indicates that the midline exists in a

more immature state in RA depleted conditions, and the more mature state provides

space in which the angioblasts can migrate into.

Figure 5. Retinoic acid mediates a notochord-endoderm separation required for terminal angioblast migration.

(**A-I**) Embryos generated by crossing *tg(HS:lifeact)* to *tg(kdrl:eGFP)* to label both actin and angioblasts. Yellow arrowheads indicate the midline while white arrows indicate angioblasts. A magnified image includes a white dashed line to indicate notochord-endoderm separation. (**A-C)**) DMSO treated embryos sectioned at the 5^{th} somite during the (**A**) 12-somite stage (n=8), (**B**) 15-somite stage (N=8), and (**C**) 18-somite stage (N=6). (**D-E**) 0.1 µM RA treated embryos sectioned at the 5^{th} somite during the (**D**) 12-somite stage (N=8) and (**E**) 15-somite stage (n=8). (**F-I**) 20 µM DEAB treated embryos sectioned at the 5^{th} somite during the (**F**) 12-somite stage (N=7), (**G**) the 15-somite stage (N=8), (**H**)18-somite stage (N=6), and (**I**) 21-somite stage (N=6). (**J-L**) 15 somite-stage embryos generated by crossing *tg(HS:lifeact)* to *tg(Kdrl:eGFP).* White dashed lines indicate outline of somites while yellow brackets indicate distance of notochord to ventral endoderm. (**J**) Schematic showing the sectioned region for experiments in (**K-O**). Red line indicates sectioning at 5^{th} somite and the blue line indicates sectioning at the 12^{th} somite. (**K**) A DMSO embryo sectioned at the 12^{th} somite. Note the lack of midline angioblasts and small distance between the notochord and the ventral endoderm. (**L**) A DMSO-treated embryo sectioned at the 5^{th} somite. Note the angioblasts at the midline and larger distance from the notochord to the ventral endoderm. (**M**) A DEAB embryo sectioned at the 5^{th} somite. Note the lack of midline angioblasts and small distance between the notochord and the ventral endoderm. (**N**) A two variable graph showing percentage of angioblast fluorescence at the midline compared to distance between the notochord and the ventral endoderm. Blue dots indicate DMSO treated, 15-somite stage embryos sectioned at the 5^{th} somite and red squares are the same embryos sectioned at the 12^{th} somite (N=8). (**O**) A two variable graph showing percentage of angioblast fluorescence at the midline compared to distance between the notochord and the ventral endoderm. Blue dots indicate DEAB treated, 15-somite stage embryos sectioned at the 5^{th} somite and red squares are the same embryos sectioned at the 12^{th} somite (N=8). Scale bars, 50 µm.

Retinoic acid induces changes in the definitive vasculature.

Pharmacological inhibition of RA signaling has previously been shown to effect the

zebrafish vasculature, notably causing a smaller than normal dorsal aorta (Pillay et al.,

2016). To confirm this effect in the *aldh1a2 -/-* mutant, we sectioned 24 hpf *tg(kdrl:GFP)*

and *aldh1a2 -/-* embryos, stained the sections with DAPI, and imaged them using

spinning disk confocal microscopy (Figure 6A, 6B). Images show that the dorsal aorta

and posterior cardinal vein in wild-type embryos are roughly equivalent in size (Figure

6A, white arrows). However, in the *aldh1a2 -/-* embryos, the size of the dorsal aorta is

reduced but remained lumenized (Figure 6B, white arrows). We confirmed this

observation by in-situ hybridization of the arterial marker *cldn5b* and the venous marker

dab2, which showed similar effects (Figure S2A-D) (Casie Chetty et al., 2017). To

determine if this trend extended beyond the 24 hpf developmental stage, we imaged the

trunk of 5 dpf *tg(kdrl:GFP)* embryos along with *tg(kdrl:GFP), aldh1a2* -/- siblings (Figure

6C,6D). The *tg(kdrl:GFP)* embryos alone showed no vascular defects and normal dorsal

aorta and posterior cardinal vein (Figure 6C, white arrow and yellow arrowhead

respectively). However, the vasculature of the *aldh1a2* -/- siblings show a large reduction

of the size of the dorsal aorta (Figure 6D, white arrow). In addition, it appears that the

posterior cardinal vein of *aldh1a2* -/-increased in size compared to its sibling (Figure 6D,

yellow arrowhead). To determine if the size change was the result of a reduced cell

number in the DA in response to RA loss, we utilized the transgenic line *tg(kdrl:nls-*

GFP), which labels the nucleus of endothelial cells to count the number of cells in each

blood vessel over a 225 µM horizontal section of a 24 hpf zebrafish embryo. Embryos

were treated with either DMSO, or with DEAB to inhibit RA. We see reduced cell number

in the DEAB treated artery versus the DMSO treated artery (Figure 6E). This

corresponds to an increase number of cells in the veins during DEAB treatment

compared to DMSO. While we cannot definitively say the migration defects caused the

arterial reduction, it indicates that RA is required for the proper formation of the definitive

vasculature.

Figure 6. Retinoic acid loss contributes to changes in the definitive vasculature, resulting in large veins and small arteries.
(**A**) A 5 dpf *tg(kdrl:eGFP)* embryo. White arrows indicate the dorsal aorta and yellow arrowheads indicate the cardinal vein. (**B**) 5 dpf *tg(kdrl:eGFP), aldh1a2-/-* embryo. White arrows indicate reduced size of dorsal aorta. (**C-D**) 25 hpf *tg(kdrl:eGFP).* embryo labeled with DAPI . (**C**) Wild-type embryos show lumenized blood vessels and normal sized dorsal aorta. (**D**) Labeled *aldh1a2 -/-, tg(kdrl:eGFP)* sibling shows small lumenized dorsal aorta with large posterior cardinal vein. (**E**) Quantification of DMSO (n=12) or 20 µM DEAB (n=12) treated *tg(kdrl:nls-eGFP)* embryos. Nuclei were counted in each structure over a 225 µM long transverse section of the zebrafish trunk. Two-tailed *P*-values for the unpaired t-test of wild-type vs 20 µM DEAB treated embryos are <.00001 and .0002 for arteries and veins, respectively. Scale bars, 50 µm.

Retinoic acid induced somite morphogenesis facilitates notochord-endoderm

separation and facilitates midline angioblast migration.

While NES and midline formation were shown to be highly correlated, we previously

showed that cell autonomous RA depletion within the somite caused angioblast

migration defects. Therefore, we sought to determine if RA depletion within the somites

caused morphological changes that compromised NES and angioblast migration to the

midline. During normal development, newly-born somites initially lay flat and extend

farther in the medial-lateral axis, demonstrated by a section of 15-somite stage embryo

at the 12th somite (Figure 7A, white dashed lines) (Tlili et al., 2019). However, as somites

mature, they will narrow in the medial-lateral axis and extend in the dorsal-ventral axis,

as demonstrated by a section of the 5th somite of a 15-somite stage embryo (Figure 7B,

white dashed line) (Tlili et al., 2019). We therefore sought to determine if somite

maturation could be affected cell autonomously in RA depletion conditions. We therefore

transplanted either wild-type or *HS:dnRAR* cells into *tg(ubb:lck-mng)* embryos. Embryos

were then sectioned at roughly the 5th somite region for 12-somite stage embryos

(Figure 7C and 7D). Embryos that were transplanted with wild-type cells showed little

difference with the contralateral somite in terms of shape and maturation (Figure 7C,

white dashed lines). However, embryos transplanted with *HS:dnRAR* cells showed a

contralateral defect in the somite containing donor cells, with the somite extending

farther in the medial-lateral axis while the contralateral somite extends farther in the

dorsal-ventral axis (Figure 7D, white dashed lines). To illustrate this effect across

multiple experimental embryos, we show overlapping shapes of chimeric wild-type

somites, the chimeric *HS:dnRAR* somites, and the contralateral somites from *HS:dnRAR*

chimeras (Figure 7E). The overlapping images show a consistent trend of a shallow

dorsal-ventral axis in the chimeric *HS:dnRAR* somites compared to the wild-type

somites. Together, this indicates that RA is required cell autonomously for somite

maturation and dorsal-ventral extension.

We speculate that the failure of somites to mature was the result of reduced cell

movements within the somite. To confirm this, mosaically labeled cells were generated

by transplanting rhodamine dextran labeled cells from a wild-type donor into a wild-type

host. The embryos were treated with either DMSO or DEAB and time-lapse imaged over

the course of 4 hours (Figure 7F and Figure 7G, respectively). Cells in DMSO treated

embryos showed broad movements consistent with that of somite rotation, where cells

located in the anterior of the somite will translocate to the posterior of the somite

(Hollway et al., 2007) (Figure 7F, white arrows). However, cells in DEAB treated

embryos show little intratissue displacement and cells largely remained static (Figure

7G, white arrows). To quantify this, we utilized *tg(hsp70I:CAAX-mCherry-2A-NLS-*

KikGR) embryos to observe cell movement across the entirety of the somite. Embryos

were heat shocked during shield stage and subsequently treated with either DMSO or

DEAB and their somites were time-lapse imaged for 4 hours. Using Fiji Trackmate

software (Tinevez et al., 2017), we quantified the displacement and speed of cell

movements within the 2nd somite (Figure 7H and 7I). Cells in DMSO treated embryos

showed significant amounts of displacement throughout the somite (Figure 7H).

However the DEAB cells showed less displacement over time (Figure 7H). Quantification

of speed in the same manner showed no statistically significant difference between

DMSO and DEAB treatment (Figure 7I). A sample of somite cell tracks shows little

directional movement in loss of RA conditions compared to DMSO, indicating cells are

not moving in a manner consistent with reshaping of the somite, as well as suggesting a

role for RA signaling in somite rotation (Figure 7J and 7K, Supplemental Movie 6 and 7,

respectively). The Cxcr4a/Cxcl12a signaling axis has been shown to be responsible for

controlling cell movements in the somite. In-situ hybridization for *cxcr4a* shows a loss of

expression in RA depletion conditions and gain of expression in RA addition conditions

(Hollway et al., 2007) (Figure S5D and S5E).

We speculated that the ventral expansion of the somite that occurs during somite cell

rearrangement and maturation was responsible for NES. At the 10 somite stage,

embryos have their notochord adjacent to the endoderm and level to ventral-most edge

of the somite (Figure 7L). As the embryo develops, the somite expands ventrally (black

bracket). As the ventral distance of the somite extends relative to the dorsal part of the

notochord (yellow bracket), it induces space that the angioblasts occupy (blue brackets).

Starting from the 10 somite stage, we measured the distance of ventral expansion

relative to the distance between the notochord and the endoderm every hour for 4 hours.

A simple linear regression of the data, indicated by the black line, shows a slope of

0.002 and y-intercept of 1.04 (Figure 7M). The flat slope indicates little variance across

time, and the y-intercept near 1 indicates that NES is equivalent to ventral expansion of

the somite. The 1:1 correlation of ventral somite expansion and notochord displacement,

as well as the *tbx16* loss of function NES defect, provides strong evidence that somite

morphogenesis is critical for NES.

Finally, we investigated a methodology to track all cellular movements within an embryo

to co-visualize somite morphogenesis along with angioblast migration and notochord-

endoderm separation. We used a live explant method to monitor migration in a

transverse section view. Live *tg(kdrl:GFP)* embryos, injected with *mcherry-caax* mRNA,

were sectioned at the 10-somite stage at the 4th somite and mounted to observe a

transverse section of the trunk. We were able to observe migrating angioblasts, as well

as somite morphogenesis and midline NES in the explant (Figure 7O-7P, Supplemental

Video 8). At the 180' time-point, angioblasts were not at the midline and the hypochord

remained flush with the endoderm (Figure 7O, Supplemental Video 8). At this time-point

the somites have begun to narrow in the medial-lateral axis. By the 240' stage, the

hypochord is separated away from the endoderm (Figure 7P). and shortly after this, the

angioblasts are able to complete their migration to the midline by the 270' time-point

(Figure 7P). Similar to the results seen in fixed sections, live explants of embryos treated

with DEAB show angioblast migration defects and lack of NES (Supplemental Video 9).

Figure 7. Retinoic acid controls intrasomitic cellular movements and somite shape changes that contribute to angioblast convergence to the midline.

(**A**) Image and schematic of newly born 12th somite using *tg(HS:lifeact), tg(kdrl:egfp)* embryo as reference. Dashed lines indicate the somite and notochord. (**B**) Schematic of more mature 5th somite (at the 12-somite stage) of *tg(HS:lifeact), tg(kdrl:egfp)* embryo as reference. Dashed lines indicate somite and notochord. (**C**) Transplant of wild-type cells into *tg(ubb:lck-mng)*. Dashed lines indicate somite shape. (**D**) Transplant of *tg(HS:dnrar)* cells into *tg(ubb:lck-mng)*. Dashed lines indicate somite shape. Note the immature shape of the somite containing transplanted cells. (**E**) Overlapping somite shapes of either WT→WT, *tg(HS:dnrar)*→WT, or the contralateral somite *tg(HS:dnrar)*→WT. (**F-G**) Time-lapse images of transplanted rhodamine-dextran labeled cells in DMSO and DEAB embryos at 12 somite stage. (**F**) DMSO treated embryos over 180'. White arrows indicate labeled cells (**G**) 20 µM DEAB treated embryos over 180'. Scale bars, 50 µm. Retinoic acid depleted embryos show little cellular movement during somite development. (**H**) Quantification of nuclei displacement in the 2nd somite of DMSO and DEAB embryos over a 4 hour time period. Displacement is defined as the distance between the beginning location and end location of the nuclei within the somite 2. (**I**) Quantification of speed in somite 2, defined as average units over 4 hours in 10 minute increments. (**J**) Sample of tracks from DMSO treated *tg(hsp70l:CAAX-mCherry-2A-NLS-KikGR)* embryos. (**K**) Sample tracks of DEAB treated *tg(hsp70l:CAAX-mCherry-2A-NLS-KikGR)* embryos. Tracks sort from red to blue, with red being the highest displacement. Scale bars, 20 µm (**L**) Diagram of ventral somite expansion and notochord to endoderm separation. Ventral somite expansion is calculated by ventral somite length (black bracket) – the notochord diameter (Yellow Bracket). Notochord/endoderm separation is shown in the blue bracket. (**M**) Quantification of the ventral expansion of the somite over notochord/endoderm separation in wild-type embryos. The linear regression indicated by the black line, where slope = 0.002, shows ventral somite expansion correlates with notochord/endoderm separation. (**N**) A two variable graph showing percentage of angioblast fluorescence at the midline compared to distance of ventral somite expansion beneath the notochord. Blue dots indicate DEAB treated, 15-somite stage embryos sectioned at the 5th somite and red squares are the same embryos sectioned at the 12th somite (N=8). Scale bars, 50 µm. (**O**) Live section of *mcherry-caax* injected *tg(kdrl:egfp)* embryos over 120'. Yellow arrowheads indicate the midline while white arrows indicate angioblasts. (**P**) Same live section as in (N), except at 240' focused at the midline. Note the separation of the hypochord and endoderm prior to midline fusion of angioblasts. Scale bars, 50 µm.

DISCUSSION

Here, we established a cell nonautonomous angioblast migration model wherein shape changes in the somitic mesoderm create a physical space for angioblasts to migrate into and develop into the axial vasculature. The emergence of the transient VMC is a necessary requirement for contralateral angioblasts to complete their migration and coalesce at the midline. We observed a defect in somitic mesoderm development in RA depletion conditions, which delayed somite maturation, intrasomite cellular movement, and significantly impeded NES. Conversely, we found that addition of RA accelerated somite maturation and NES, along with angioblast migration. We were able to confirm that the somitic mesoderm was required for angioblast migration by examining *tbx16* mutants, which have a lack of somites caused by EMT defects (Goto et al., 2017; Ho and Kane, 1990; Manning and Kimelman, 2015; Row et al., 2011). Loss of function of *tbx16* prevented angioblast migration completely and resulted in bifurcated blood vessel formation. These changes in vascular morphology indicate that the somitic mesoderm is a requirement for angioblast migration to the midline. It is notable that in amniotes, such as the mouse and quail, the dorsal aorta and cardinal vein are initially bifurcated in a similar manner to zebrafish *tbx16* mutants prior to fusion at the midline (Drake and Fleming, 2000; Pardanaud et al., 1996, 1987). It is possible that evolutionary differences in the somitic mesoderm between teleosts and amniotes could play a role in the differences in vascular development.

Very little is known about the process of NES. Our observations indicate that its initiation correlates very closely with angioblast migration, with angioblasts migrating into the VMC seemingly as soon as it appears in both wild-type and RA depletion conditions. Given that these blood vessels become lumenized, and need to achieve a sufficient volume for blood transport, it is plausible that they require morphological changes within the ventral part of the embryo to accommodate for their size. Delays in somite maturation prevent angioblasts from reaching the midline in a normal developmental time frame. It is possible that the consequences of this migration defect results in the malformed blood vessel structure in RA depleted embryos, which have small arteries and larger veins. Prior work showed that the most lateral angioblasts, which migrate later than more medial angioblasts, preferentially adopt a venous fate over an arterial one (Kohli et al., 2013). This implies that somite maturation is a key factor in determining the

proper distribution of endothelial progenitors to the arteries and veins, by essentially

timing angioblast migration and altering cell fate decision patterns. However, we cannot

definitively determine if this is solely from angioblast migration defects.

We found that the formation of the VMC does not necessarily require the notochord

tissue itself. In *noto* morphants, which causes the notochord to adopt a somitic fate,

there is still a ventral cavity and angioblasts are able to arrive at the midline. This implies

that the notochord is ultimately dispensable for the completion of angioblast migration,

even though its loss can produce migration defects (Helker et al., 2015; Sumoy et al.,

1997). The migratory defects could be the result of reduced notochord-secreted factors

such as *apela,* which greatly reduce but do not completely abolish migration (Helker et

al., 2015). Another factor could be that signals from the notochord, such as Hedgehog,

influence somite development which in turn facilitates angioblast migration (Blagden et

al., 1997; Hinits et al., 2009; Yin et al., 2018). While the fate of the midline tissue does

not need to be notochord for NES to occur, it is likely that mechanical coupling between

the somites and midline tissue is required such that somite morphogenesis drives the

separation of the midline mesoderm away from the underlying endoderm. In *noto* loss of

function, the somitic tissue that forms at the midline in place of notochord is continuous

with the flanking somites. In wild-type embryos, there is evidence of mechanical

interactions of the somite and notochord during convergent extension, which are

required for the remodeling of the adaxial cells, the medial most cells in the somite (Yin

and Solnica-Krezel, 2007).

The phenotypes observed in both the somites and endothelium after loss of RA signaling

are not the result of a simple deficit of mesoderm. RA signaling loss of function affects

somite maturation and morphogenesis, but does not prevent them from forming, and yet

the migration of the angioblasts is still disrupted. The direct transcriptional targets of RA

signaling that induce somite maturation are unknown, but it is clear that RA signaling

controls the timing and progress of somite shape changes as they mature. Cell

autonomous effects of RA depletion on the somites, coupled with angioblast migration

defects in RA depleted somites, indicate RA directly controls somite shape changes that

facilitate migration. The fact that somites can adopt mature shapes in explants, even in

absence of adjacent somites, indicates this control is local to individual somites. It is

also noteworthy that loss of retinoic acid signaling activity in a relatively few number of

adaxially localized cells in our chimeric somites were sufficient to cause somite shape

and angioblast migration defects. This implies that the somite cells most medial to the

midline could be the population required for the ventral expansion of the somite and

NES. It is not clear if NES is required for the initiation of angioblast migration, as some

midline-directed cellular movement occurs in DEAB treated embryos before NES is

visibly seen. It is possible that the somite itself could facilitate the angioblasts movement

to the midline, or simply provide an opening through which angioblasts can reach the

midline. The fact that angioblasts reside adjacent to the notochord in *tbx16* mutants, but

do not migrate beneath them, implies angioblasts are not sufficient to drive NES or their

coalescence at the midline. It is likely that ventral expansions of the somite drives the

notochord away from the endoderm. Together, this indicates that somitic maturation

requires tight spatial and temporal control to facilitate the concomitant development of

the endothelium.

Materials and Methods

Generation of the *tg(hsp70l:lifeact-mscarlet)* transgenic line

This transgenic line was generated using the plasmid pNJP002 (*hsp70l:lifeact-mScarlet*)

and *tol2* transgenesis (Kawakami, 2004; Kikuta and Kawakami, 2009). pNJP002 was

generated using PCR amplification paired with Gibson assembly cloning. A restriction

digest was performed on a tol2-hsp70 plasmid (Row et al., 2016) using the restriction

enzymes BamHI-HF and ClaI (NEB bio labs). PCR amplification of plasmid hCCR4

(lifeact-mScarlet) was performed using forward primer

CAAGCTACTTGTTCTTTTTGCAGGATCCATGGGCGTGGCCGACTTG and reverse

primer TTCGTGGCTCCAGAGAATCGATTCACTTGTACAGCTCGTCCATGC. Gibson

Assembly (NEBbuilder HiFi DNA Assembly, NEB bio labs) was performed on restriction

digested hCCR2 and PCR amplified *lifeact-mscarlet* to generate plasmid pNJP002. The

tg(hsp70l:lifeact-mscarlet) line was generated by injecting 25 picograms of pNJP002 with

25 picograms of *tol2* mRNA using methods described previously (Row et al., 2016).

In-situ hybridization and immunohistochemistry

Whole-mount in situ hybridization was performed as previously described (Griffin et al.,

1995). An antisense RNA probe was synthesized as previously described for *etv2*

(Sumanas et al., 2005). The *cldn5b, cxcr4a, apela, aplnra, aplnrb, cxcl12a* and *dab2* DIG

probes were generated by *taq* polymerase-generated PCR fragments integrated into a

pCRII vector using the Topo-TA Cloning kit (Thermo Fisher). Immunohistochemistry was

performed by fixing embryos overnight in 4% Paraformaldehyde (Sigma) dissolved in

phosphate buffered saline containing 0.1% Tween-20 (PBST) at 4°C. Permeabilization

was performed by the placing the embryos in PBS containing 2% Triton X-100 (Sigma)

for 1 hour with agitation. Embryos were washed 3x with PBST and blocked in a blocking

solution (PBST, 10% goat serum, 1% BSA) for 2 hours at room temperature. Primary

antibodies, Etv2 Rabbit Polyclonal and GFP mouse monoclonal, were diluted to 1:500 in

blocking solution and incubated overnight at 4°C. Embryos were washed 5x with PBST

for 20 mins each wash. Embryos were then incubated in blocking solution for 2 hours;

and then incubated in the secondary antibodies Alexa Fluor 488 and Alexa Fluor 568

diluted 1:1000 in blocking solution at 4°C overnight. DAPI stains were done in a 10

µg/mL dilution in PBST for at least 1 hour with agitation.

Histological analysis

Sections were fixed overnight in 4% Paraformaldehyde (Sigma) dissolved in phosphate

buffered saline containing (PBS) at 4°C. Nuclei labeling was done for whole embryos in

a 10 µg/mL dilution of DAPI in PBST for at least 1 hour with agitation. Sections, after

fixation and DAPI staining, were done in PBST with a 0.15mm microknife (Fine Science

Tools), at the indicated somite location for embryos up to 24 hpf. At 24 hpf or older,

embryos were sectioned at approximately the 7th somite. Sections were mounted in 1%

Agarose in PBS in a in a 35mm glass bottom dish with uncoated #1.5 coverslip and

20mm glass diameter (MatTek).

Microinjections

A mix of two *tbx16* morpholinos (MO1: AGCCTGCATTATTTAGCCTTCTCTA (1.5 ng)

and MO2: GATGTCCTCTAAAAGAAAATGTCAG (0.75 ng)) were prepared as previously

reported and injected into *tg(kdrl:GFP)* embryos (Row et al., 2016). 2.5 ng of control

morpholino and *noto* morpholino were injected into *tg(kdrl:GFP)* embryos (Ouyang et al.,

2009). 200 ng of *mcherry-caax* mRNA was injected into *tg(kdrl:GFP)* embryos.

Microscopy and Imaging

DIC and fluorescent time-lapse images of wild-type, *noto* morphants, *aldh1a2* mutants*,*

tbx16 mutants, and drug treated embryos were performed using a Leica DMI6000B

inverted microscope. Wild-type and control embryos were siblings of the morphants and

mutants. Live embryos were mounted in 1% low-melt agarose in embryo media

containing 1x tricaine (25x stock 0.4g/l;Pentair,TRS1) in a 35mm glass bottom dish with

uncoated #1.5 coverslip and 20mm glass diameter (MatTek). In-situ hybridization

experiments were imaged using a M165FC microscope (Leica) equipped with an Infinity

3 camera (Lumenera). Embryos were mounted in either horizontal or flat mounted

configuration in 70% glycerol on glass slides.

Fluorescent images of control, *tbx16, and noto morphant* (Figure 3) cross sections were

imaged on a custom assembled spinning disk confocal microscope consisting of an

automated Zeiss frame, a Yokogawa CSU-10 spinning disc, a Ludl stage controlled by a

Ludl MAC6000 and an ASI filter turret mated to a Photometrics Prime 95B camera. All

other cross sections, including explants, were performed on a custom-assembled

spinning disk confocal microscope consisting of a Zeiss Imager A.2 frame, a Borealis

modified Yokogawa CSU-10 spinning disc, ASI 150uM piezo stage controlled by an

MS2000, and ASI filter wheel, and a Hamamatsu ImageEM x2 EMCCD camera

(Hamamatsu C9100-23B). These microscopes were controlled with Metamorph

microscope control software (V7.10.2.240 Molecular Devices), and laser power levels

were set in Vortran's Stradus VersaLase 8 software. Images were processed in Fiji.

Drug treatments

RA was depleted in embryos using N,N-diethylaminobenzaldehyde (DEAB) and

BMS453. Stock concentrations of 20 mM DEAB in DMSO were diluted to working

concentrations of 20 µM in embryo media. Stock concentrations of 2 mM BMS453 in

DMSO were diluted to working concentrations of 2 µM in embryo media. Treatments

were done on shield stage embryos unless otherwise noted. All-trans RA was used to

activate the RA pathway. Stock concentrations of All-trans RA were diluted from 1 mM

stock to 0.1 µM working concentrations in embryo media. All treatments for all-trans RA

were performed in tailbud stage.

Generation of explants for imaging

Transgenic or mutant embryos were grown to the 10 somite stage and then transferred

to Modified Barth's Saline (MBS) (Sigma). In this medium, embryos were anesthetized

with 1 x Tricaine (25x stock 0.4g/l; Pentair, TRS1) and had their chorions manually

removed by forceps. The embryos were then sectioned with a 0.15mm microknife (Fine

Science Tools). Excess yolk was removed with the microknife, leaving some yolk to

prevent injury to the endoderm, and the embryo was then transferred to 35 mm glass-

bottom dish (Matek) with a fire polished pipette. 1.3 % low gelling temperature agarose

in MBS was heated to 40°C and placed over the embryo in the proper orientation. The

agarose was allowed to solidify and the explant was time-lapsed using fluorescent and

DIC microscopy.

Quantification and Statistical Analysis

When determining the fluorescence of angioblasts at the midline of the embryo, we

utilized the integrated density feature of Fiji software for image analysis. We calculated

the corrected total fluorescence of the area at the midline and of the whole embryo,

using the notochord as a reference for the midline. Corrected total fluorescence for the

midline was calculated as: (mean gray value of midline x area of midline) - (mean gray

value of background x area of midline). Corrected total fluorescence for the embryo was

calculated: (mean gray value of embryo x area of embryo) - (mean gray value of

background x area of embryo). Percentages for fluorescent intensity were calculated as

(corrected total fluorescence at the midline) / (corrected total fluorescence of the whole

embryo). Tracks for cells were made using the Trackmate plug-in for Fiji (Tinevez et al.,

2017). We utilized a Laplacian of Gaussian detector and a Simple LAP Tracker to

generate the tracks and analysis. P-values for indicated figures were generated using

two-tailed unpaired student t-tests. Graphs were generated using Graphpad Prism 8.4.2

for dot plots, and Excel for line graphs. The black lines indicate the median and

interquartile ranges, and were generated using the Prism software. Simple linear

regression was also determined using Graphpad software.

Acknowledgments

We thank David Matus for critical review of the manuscript, as well as members of the

Martin and Matus labs for helpful comments. We thank Jesus Torres Vazquez for

sending the kdrl transgenic reporter lines. We thank Neal Bhattacharji and Stephanie

Flanagan for excellent zebrafish care. We thank Tianying Chen for assistance in data

acquisition. We thank Rebecca Adikes, Thom Geer and Nobska Imaging for microscopy

support. The illustration of the 15-somite stage zebrafish embryo was created by

Biorender.com. This work was supported by the NSF (IOS 1452928) and NIH NIGMS

(R01GM124282).

References

Adikes RC, Kohrman AQ, Martinez MAQ, Palmisano NJ, Smith JJ, Medwig-Kinney TN,

Min M, Sallee MD, Ahmed OB, Kim N, Liu S, Morabito RD, Weeks N, Zhao Q,

Zhang W, Feldman JL, Barkoulas M, Pani AM, Spencer SL, Martin BL, Matus DQ.

2020. Visualizing the metazoan proliferation-quiescence decision in vivo. *Elife* **9**:1–

74. doi:10.7554/eLife.63265

Amacher SL, Draper BW, Summers BR, Kimmel CB. 2002. The zebrafish T-box genes

no tail and spadetail are required for development of trunk and tail mesoderm and

medial floor plate. *Development* **129**:3311–3323.

Balmer S, Nowotschin S, Hadjantonakis AK. 2016. Notochord morphogenesis in mice:

Current understanding and open questions. *Dev Dyn* **245**:547–557.

doi:10.1002/dvdy.24392

Berenguer M, Lancman JJ, Cunningham TJ, Dong PDS, Duester G. 2018. Mouse but

not zebrafish requires retinoic acid for control of neuromesodermal progenitors and

body axis extension. *Dev Biol* **441**:127–131. doi:10.1016/j.ydbio.2018.06.019

Bernheim S, Meilhac SM. 2020. Mesoderm patterning by a dynamic gradient of retinoic

acid signalling: Retinoic acid patterning. *Philos Trans R Soc B Biol Sci* **375**:1809.

doi:10.1098/rstb.2019.0556rstb20190556

Blagden CS, Currie PD, Ingham PW, Hughes SM. 1997. Notochord induction of

zebrafish slow muscle mediated by sonic hedgehog. *Genes Dev*.

doi:10.1101/gad.11.17.2163

Casie Chetty S, Rost MS, Enriquez JR, Schumacher JA, Baltrunaite K, Rossi A, Stainier

DYR, Sumanas S. 2017. Vegf signaling promotes vascular endothelial

differentiation by modulating etv2 expression. *Dev Biol* **424**:147–161.

doi:10.1016/j.ydbio.2017.03.005

Chng SC, Ho L, Tian J, Reversade B. 2013. ELABELA: A hormone essential for heart

development signals via the apelin receptor. *Dev Cell* **27**:672–680.

doi:10.1016/j.devcel.2013.11.002

Cleaver O, Krieg P a. 1998. VEGF mediates angioblast migration during development of

the dorsal aorta in Xenopus. *Development* **125**:3905–14.

Drake CJ, Fleming PA. 2000. Vasculogenesis in the day 6.5 to 9.5 mouse embryo.

Blood **95**:1671–9. doi:10.1016/j.stem.2010.07.016

Duester G. 2008. Retinoic Acid Synthesis and Signaling during Early Organogenesis.

Cell. doi:10.1016/j.cell.2008.09.002

Emoto Y, Wada H, Okamoto H, Kudo A, Imai Y. 2005. Retinoic acid-metabolizing

enzyme Cyp26a1 is essential for determining territories of hindbrain and spinal cord

in zebrafish. *Dev Biol*. doi:10.1016/j.ydbio.2004.11.023

Fouquet B, Weinstein BM, Serluca FC, Fishman MC. 1997. Vessel patterning in the

embryo of the zebrafish: Guidance by notochord. *Dev Biol* **183**:37–48.

doi:10.1006/dbio.1996.8495

Freyer L, Hsu CW, Nowotschin S, Pauli A, Ishida J, Kuba K, Fukamizu A, Schier AF,

Hoodless PA, Dickinson ME, Hadjantonakis AK. 2017. Loss of Apela Peptide in

Mice Causes Low Penetrance Embryonic Lethality and Defects in Early

Mesodermal Derivatives. *Cell Rep* **20**:2116–2130. doi:10.1016/j.celrep.2017.08.014

Glickman NS, Kimmel CB, Jones MA, Adams RJ. 2003. Shaping the zebrafish

notochord. *Development* **130**:873–887. doi:10.1242/dev.00314

Goto H, Kimmey SC, Row RH, Matus DQ, Martin BL. 2017. FGF and canonical Wnt

signaling cooperate to induce paraxial mesoderm from tailbud neuromesodermal

progenitors through regulation of a two-step epithelial to mesenchymal transition.

Development **144**:1412–1424. doi:10.1242/dev.143578

Griffin KJ, Amacher SL, Kimmel CB, Kimelman D. 1998. Molecular identification of

spadetail: regulation of zebrafish trunk and tail mesoderm formation by T-box

genes. *Development* **125**:3379–3388.

Halpern ME, Thisse C, Ho RK, Thisse B, Riggleman B, Trevarrow B, Weinberg ES,

Postlethwait JH, Kimmel CB. 1995. Cell-autonomous shift from axial to paraxial

mesodermal development in zebrafish floating head mutants. *Development*

121:4257–4265. doi:10.1016/0168-9525(96)81431-4

Hamade A, Deries M, Begemann G, Bally-Cuif L, Genêt C, Sabatier F, Bonnieu A,

Cousin X. 2006. Retinoic acid activates myogenesis in vivo through Fgf8 signalling.

Dev Biol **289**:127–140. doi:10.1016/j.ydbio.2005.10.019

Heine UI, Roberts AB, Munoz EF, Roche NS, Sporn MB. 1986. Effects of retinoid

deficiency on the development of the heart and vascular system of the quail

embryo. *Virchows Arch B Cell Pathol Incl Mol Pathol*. doi:10.1007/BF02889897

Helker CSM, Schuermann A, Pollmann C, Chng SC, Kiefer F, Reversade B, Herzog W.

2015. The hormonal peptide Elabela guides angiob lasts to the midline during

vasculogenesis. *Elife* **4**:1–13. doi:10.7554/eLife.06726

Hinits Y, Osborn DPS, Hughes SM. 2009. Differential requirements for myogenic

regulatory factors distinguish medial and lateral somitic, cranial and fin muscle fibre

populations. *Development* **136**:403–414. doi:10.1242/dev.028019

Ho RK, Kane DA. 1990. Cell-autonomous action of zebrafish spt-1 mutation in specific

mesodermal precursors. *Nature*. doi:10.1038/348728a0

Hogan KA, Bautch VL. 2004. Blood Vessel Patterning at the Embryonic Midline. *Curr

Top Dev Biol* **62**:55–85.

Hollway GE, Bryson-Richardson RJ, Berger S, Cole NJ, Hall TE, Currie PD. 2007.

Whole-Somite Rotation Generates Muscle Progenitor Cell Compartments in the

Developing Zebrafish Embryo. *Dev Cell* **12**:201–219.

doi:10.1016/j.devcel.2007.01.001

Isogai S, Horiguchi M, Weinstein BM. 2001. The vascular anatomy of the developing

zebrafish: An atlas of embryonic and early larval development. *Dev Biol* **230**:278–

301. doi:10.1006/dbio.2000.9995

Janesick A, Nguyen TTL, Aisaki K -i., Igarashi K, Kitajima S, Chandraratna RAS, Kanno

J, Blumberg B. 2014. Active repression by RAR signaling is required for vertebrate

axial elongation. *Development* **141**:2260–2270. doi:10.1242/dev.103705

Janesick A, Tang W, Shioda T, Blumberg B. 2018. RARγ is required for mesodermal

gene expression prior to gastrulation in xenopus. *Dev* **145**:2–9.

doi:10.1242/dev.147769

Jin SW, Beis D, Mitchell T, Chen JN, Stainier DYR. 2005. Cellular and molecular

analyses of vascular tube and lumen formation in zebrafish. *Development*

132:5199–5209. doi:10.1242/dev.02087

Kawakami K. 2004. Transgenesis and Gene Trap Methods in Zebrafish by Using the

Tol2 Transposable Element. *Methods Cell Biol* **77**:201–222. doi:10.1016/S0091-

679X(04)77011-9

Kikuta H, Kawakami K. 2009. Transient and stable transgenesis using tol2 transposon

vectors. *Methods Mol Biol* **546**:69–84. doi:10.1007/978-1-60327-977-2_5

Kimelman D. 2016. Tales of tails (and trunks): forming the posterior body in vertebrate

embryos David Kimelman. *Curr Top Dev Biol* **116**:517–536.

doi:10.1016/bs.ctdb.2015.12.008.Tales

Kimmel CB, Ballard WW, Kimmel SR, Ullmann B, Schilling TF. 1995. Stages of

embryonic development of the zebrafish. *Dev Dyn* **203**:253–310.

doi:10.1002/aja.1002030302

Kimmel CB, Kane DA, Walker C, Warga RM, Rothman MB. 1989. A mutation that

changes cell movement and cell fate in the zebrafish embryo. *Nature*.

doi:10.1038/337358a0

Kohli V, Schumacher JA, Desai SP, Rehn K, Sumanas S. 2013. Arterial and Venous

Progenitors of the Major Axial Vessels Originate at Distinct Locations. *Dev Cell*

25:196–206. doi:10.1016/j.devcel.2013.03.017

Kumar S, Duester G. 2014. Retinoic acid controls body axis extension by directly

repressing Fgf8 transcription. *Dev* **141**:2972–2977. doi:10.1242/dev.112367

Lawson ND, Vogel AM, Weinstein BM. 2002. Sonic hedgehog and vascular endothelial

growth factor act upstream of the Notch pathway during arterial endothelial

differentiation. *Dev Cell* **3**:127–136. doi:10.1016/S1534-5807(02)00198-3

Leal MA, Fickel SR, Sabillo A, Ramirez J, Vergara HM, Nave C, Saw D, Domingo CR.

2014. The Role of Sdf-1α signaling in Xenopus laevis somite morphogenesis. *Dev*

Dyn. doi:10.1002/dvdy.24092

Li J, Yue Y, Dong X, Jia W, Li K, Liang D, Dong Z, Wang X, Nan X, Zhang Q, Zhao Q.

2015. Zebrafish foxc1a plays a crucial role in early somitogenesis by restricting the

expression of aldh1a2 directly. *J Biol Chem* **290**:10216–10228.

doi:10.1074/jbc.M114.612572

Lohnes D, Mark M, Mendelsohn C, Dolle P, Dierich A, Gorry P, Gansmuller A, Chambon

P. 1994. Function of the retinoic acid receptors (RARs) during development. (I)

Craniofacial and skeletal abnormalities in RAR double mutants. *Development*

120:2723–2748.

Mandal A, Rydeen A, Anderson J, Sorrell MRJ, Zygmunt T, Torres-Vázquez J, Waxman

JS. 2013. Transgenic retinoic acid sensor lines in zebrafish indicate regions of

available embryonic retinoic acid. *Dev Dyn* **242**:989–1000. doi:10.1002/dvdy.23987

Manning AJ, Kimelman D. 2015. Tbx16 and Msgn1 are required to establish directional

cell migration of zebrafish mesodermal progenitors. *Dev Biol* **406**:172–185.

doi:10.1016/j.ydbio.2015.09.001

Martin BL. 2016. Factors that coordinate mesoderm specification from neuromesodermal

progenitors with segmentation during vertebrate axial extension. *Semin Cell Dev*

Biol **49**:59–67. doi:10.1016/j.semcdb.2015.11.014

Martin BL, Kimelman D. 2012. Canonical Wnt Signaling Dynamically Controls Multiple

Stem Cell Fate Decisions during Vertebrate Body Formation. *Dev Cell* **22**:223–232.

doi:10.1016/j.devcel.2011.11.001

Martin BL, Kimelman D. 2010. Brachyury establishes the embryonic mesodermal

progenitor niche. *Genes Dev* **24**:2778–2783. doi:10.1101/gad.1962910

McMillen P, Holley SA. 2015. The tissue mechanics of vertebrate body elongation and

segmentation. *Curr Opin Genet Dev*. doi:10.1016/j.gde.2015.02.005

Mendelsohn C, Lohnes D, Decimo D, Lufkin T, LeMeur M, Chambon P, Mark M. 1994.

Function of the retinoic acid receptors (RARs) during development. (II) Multiple

abnormalities at various stages of organogenesis in RAR double mutants.

Development **120**:2749–2771.

Morgan CA, Parajuli B, Buchman CD, Dria K, Hurley TD. 2015. N,N-

diethylaminobenzaldehyde (DEAB) as a substrate and mechanism-based inhibitor

for human ALDH isoenzymes. *Chem Biol Interact* **234**:18–28.

doi:10.1016/j.cbi.2014.12.008

Niederreither K, Vermot J, Messaddeq N, Schuhbaur B, Chambon P, Dollé P. 2001.

Embryonic retinoic acid synthesis is essential for heart morphogenesis in the

mouse. *Development* **128**:1019–1031.

Norris ML, Pauli A, Gagnon JA, Lord ND, Rogers KW, Mosimann C, Zon LI, Schier AF.

2017. Toddler signaling regulates mesodermal cell migration downstream of nodal

signaling. *Elife* **6**:e22626. doi:10.7554/eLife.22626

Ouyang X, Shestopalov IA, Sinha S, Zheng G, Pitt CLW, Li WH, Olson AJ, Chen JK.

2009. Versatile synthesis and rational design of caged morpholinos. *J Am Chem*

Soc **131**:13255–13269. doi:10.1021/ja809933h

Pardanaud L, Altmann C, Kitos P, Dieterlen-Lievre F, Buck CA. 1987. Vasculogenesis in

the early quail blastodisc as studied with a monoclonal antibody recognizing

endothelial cells. *Development* **100**:339–349.

Pardanaud L, Luton D, Prigent M, Bourcheix LM, Catala M, Dieterlen-Lièvre F. 1996.

Two distinct endothelial lineages in ontogeny, one of them related to hemopoiesis.

Development **122**:1363–1371.

Pauli A, Norris ML, Valen E, Chew GL, Gagnon JA, Zimmerman S, Mitchell A, Ma J,

Dubrulle J, Reyon D, Tsai SQ, Joung JK, Saghatelian A, Schier AF. 2014. Toddler:

An embryonic signal that promotes cell movement via apelin receptors. *Science*

(80-) **343**. doi:10.1126/science.1248636

Pillay LM, Mackowetzky KJ, Widen SA, Waskiewicz AJ. 2016. Somite-derived retinoic

acid regulates zebrafish hematopoietic stem cell formation. *PLoS One.*

doi:10.1371/journal.pone.0166040

Riedl J, Crevenna AH, Kessenbrock K, Yu JH, Neukirchen D, Bista M, Bradke F, Jenne

D, Holak TA, Werb Z, Sixt M, Wedlich-Soldner R. 2008. Lifeact: A versatile marker

to visualize F-actin. *Nat Methods* **5**:605–607. doi:10.1038/nmeth.1220

Row RH, Maître JL, Martin BL, Stockinger P, Heisenberg CP, Kimelman D. 2011.

Completion of the epithelial to mesenchymal transition in zebrafish mesoderm

requires Spadetail. *Dev Biol* **354**:102–110. doi:10.1016/j.ydbio.2011.03.025

Row RH, Pegg A, Kinney B, Farr GH, Maves L, Lowell S, Wilson V, Martin BL. 2018.

BMP and FGF signaling interact to pattern mesoderm by controlling basic helix-

loop-helix transcription factor activity. *Elife* **7**:e31018. doi:10.7554/eLife.31018

Row RH, Tsotras SR, Goto H, Martin BL. 2016. The zebrafish tailbud contains two

independent populations of midline progenitor cells that maintain long-term germ

layer plasticity and differentiate in response to local signaling cues. *Dev*.

doi:10.1242/dev.129015

Sandell LL, Sanderson BW, Moiseyev G, Johnson T, Mushegian A, Young K, Rey JP,

Ma JX, Staehling-Hampton K, Trainor PA. 2007. RDH10 is essential for synthesis of

embryonic retinoic acid and is required for limb, craniofacial, and organ

development. *Genes Dev* **21**:1113–1124. doi:10.1101/gad.1533407

Shaw KM, Castranova DA, Pham VN, Kamei M, Kidd KR, Lo BD, Torres-Vasquez J,

Ruby A, Weinstein BM. 2006. Fused-Somites-Like Mutants Exhibit Defects in Trunk

Vessel Patterning. *Dev Dyn* **235**:1753–1760. doi:10.1002/dvdy.20814

Skvarca LB, Han HI, Espiritu EB, Missinato MA, Rochon ER, McDaniels MD, Bais AS,

Roman BL, Waxman JS, Watkins SC, Davidson AJ, Tsang M, Hukriede NA. 2019.

Enhancing regeneration after acute kidney injury by promoting cellular

dedifferentiation in zebrafish. *DMM Dis Model Mech*. doi:10.1242/dmm.037390

Stern CD, Piatkowska AM. 2015. Multiple roles of timing in somite formation. *Semin Cell*

Dev Biol **42**:134–139. doi:10.1016/j.semcdb.2015.06.002

Sumanas S, Jorniak T, Lin S. 2005. Identification of novel vascular endothelial-specific

genes by the microarray analysis of the zebrafish cloche mutants. *Blood* **106**:534–

541. doi:10.1182/blood-2004-12-4653

Sumoy L, Bennett Keasey J, Dittman TD, Kimelman D. 1997. A role for notochord in

axial vascular development revealed by analysis of phenotype and the expression

of VEGR-2 in zebrafish flh and ntl mutant embryos. *Mech Dev* **63**:15–27.

doi:10.1016/S0925-4773(97)00671-0

Talbot WS, Trevarrow B, Halpern ME, Melby AE, Farr G, Postlethwait JH, Jowett T,

Kimmel CB, Kimelman D. 1995. A homeobox gene essential for zebrafish

notochord development. *Nature* **378**:150–157. doi:10.1038/378150a0

Tani S, Chung U il, Ohba S, Hojo H. 2020. Understanding paraxial mesoderm

development and sclerotome specification for skeletal repair. *Exp Mol Med*

52:1166–1177. doi:10.1038/s12276-020-0482-1

Therapontos C, Vargesson N. 2010. Zebrafish notch signalling pathway mutants exhibit

trunk vessel patterning anomalies that are secondary to somite misregulation. *Dev*

Dyn **239**:2761–2768. doi:10.1002/dvdy.22410

Thompson MA, Ransom DG, Pratt SJ, MacLennan H, Kieran MW, Detrich HW, Vail B,

Huber TL, Paw B, Brownlie AJ, Oates AC, Fritz A, Gates MA, Amores A, Bahary N,

Talbot WS, Her H, Beier DR, Postlethwait JH, Zon LI. 1998. The cloche and

spadetail genes differentially affect hematopoiesis and vasculogenesis. *Dev Biol*.

doi:10.1006/dbio.1998.8887

Tinevez JY, Perry N, Schindelin J, Hoopes GM, Reynolds GD, Laplantine E, Bednarek

SY, Shorte SL, Eliceiri KW. 2017. TrackMate: An open and extensible platform for

single-particle tracking. *Methods* **115**:80–90. doi:10.1016/j.ymeth.2016.09.016

Tlili S, Yin J, Rupprecht JF, Mendieta-Serrano MA, Weissbart G, Verma N, Teng X,

Toyama Y, Prost J, Saunders TE. 2019. Shaping the zebrafish myotome by

intertissue friction and active stress. *Proc Natl Acad Sci U S A* **116**:25430–25439.

doi:10.1073/pnas.1900819116

Torres-Vázquez J, Gitler AD, Fraser SD, Berk JD, Pham VN, Fishman MC, Childs S,

Epstein JA, Weinstein BM. 2004. Semaphorin-plexin signaling guides patterning of

the developing vasculature. *Dev Cell* **7**:117–123. doi:10.1016/j.devcel.2004.06.008

Yabe T, Takada S. 2016. Molecular mechanism for cyclic generation of somites:

Lessons from mice and zebrafish. *Dev Growth Differ* 31–42. doi:10.1111/dgd.12249

Yin C, Solnica-Krezel L. 2007. Convergence and extension movements affect dynamic

notochord-somite interactions essential for zebrafish slow muscle morphogenesis.

Dev Dyn **236**:2742–2756. doi:10.1002/dvdy.21295

Yin J, Lee R, Ono Y, Ingham PW, Saunders TE. 2018. Spatiotemporal Coordination of

FGF and Shh Signaling Underlies the Specification of Myoblasts in the Zebrafish

959   Embryo. *Dev Cell* **46**:735-750.e4. doi:10.1016/j.devcel.2018.08.024

960 Youn BW, Malacinski GM. 1981. Somitogenesis in the amphibian Xenopus laevis:

961   scanning electron microscopic analysis of intrasomitic cellular arrangements during

962   somite rotation. *J Embryol Exp Morphol* **64**:23–43.

964 **Supplemental Figures**

Figure S1. Three alternative methods of RA signaling disruption inhibit midline angioblast migration.
Angioblasts are labelled by *etv2* in-situ hybridization in 13-somite stage embryos (black arrows) after (**A**) DMSO, (**B**) BMS453, or (**C**) DEAB treatment, or after bud stage heat-shock in (**D**) wild-type and (**E**) *HS:dnrar* embryos. Embryos are shown from a dorsal view with anterior to the top.

Figure S2. Apela/Aplnr axis is not altered by changes in retinoic acid signaling
In situ hybridization of 10-somite stage embryos. (**A-B**) In-situ hybridization against *aplnra*. (**A**) DMSO treated embryos. (**B**) DEAB treated embryos. (**C-D**) In-situ hybridization against *aplnrb*. (**C**) DMSO treated embryos (**D**) DEAB treated embryos. (**E-F**) In-situ hybridization against *apela*. (**E**) DMSO treated embryos (**F**) DEAB treated embryos

dab2	*cldn5b*

wild type
- A — 12/12
- B — 12/12

aldh1a2 -/-
- C — 10/12
- D — 11/13

Figure S3. Effects of RA depletion on arterial and venous markers.
(A, C) *dab2* and (B, D) *cldn5b* in-situ hybridization labels the veins and arteries, respectively in (**A-B**) wild-type or (**C-D**) aldh1a2 -/- embryos. Embryos are shown from a lateral view with anterior to the left.

wild type	*tbx16 -/-*		*wild type*	*tbx16 -/-*

dab2
- A — 24/24
- B — 20/24

cln5b
- C — 20/20
- D — 18/19

Figure S4. Bifurcated angioblasts in *tbx16* mutants preferentially join the venous population. (A, B) *dab2* and (C, D) *cldn5b* in-situ hybridization labels the veins and arteries, respectively (black arrows) in (**A,C**) wild-type and (**B,D**) *tbx16 -/-* embryos. Embryos are shown from a dorsal view with anterior to the top.

Figure S5. Retinoic acid signaling affects the expression of genes associated with somitic movements.
(**A-C**) In-situ hybridization for *cxcr4a and* (**D-F**) *cxcl12a* in (**A, D**) DMSO, (**B, E**) 20 µM DEAB, or (**C, F**) 0.1 µM RA treated embryos. Embryos are shown from a dorsal view with anterior to the top.

Movie S1. Time-lapse fluorescent imaging of *tg(kdrl:eGFP)* embryo at the 10

**somite-stage*.* *tg(kdrl:eGFP)* marks angioblasts as they migrate to the midline.

Angioblasts display the anterior posterior processivity while coalescing at the midline.

Frame rate= one image/5 min. Run time= 235 min.

Movie S2. Time-lapse fluorescent imaging of *tg(kdrl:eGFP), aldh1a2 -/- embryo at

the 10 somite-stage. *tg(kdrl:eGFP)* marks angioblasts as they migrate to the midline.

Angioblasts lose anterior to posterior processivity and show disrupted migration. Frame

rate= one image/5 min. Run time= 235 min.

Movie S3. Time-lapse fluorescent imaging of a 10 somite-stage *tg(kdrl:eGFP)* **embryo with addition of 0.1 µM RA at the tailbud stage.** *tg(kdrl:eGFP)* marks angioblasts as they migrate to the midline. Angioblasts accelerate their processivity towards the midline. Frame rate= one image/5 min. Run time= 235 min.

Movie S4. Time-lapse fluorescent imaging of *HS:id3, HS;dnRAR* **cells in a** *tg(kdrl:eGFP)* **host.** Cells expressing *HS:id3, HS;dnRAR* migrate to the midline along with angioblasts in a *tg(kdrl:eGFP)* host. Cells with dnRAR migrate faithfully with angioblasts. Frame rate= one image/5 min. Run time= 180 min.

Movie S5. Time-lapse fluorescent imaging of *tg(tbxta:kaedae), tg(actc1b:gfp)* **trunk explant.** Trunk explant shows notochord displacement away from ventral-most portion of somites. Frame rate= one image/15 min. Run time= 285 min.

.

Movie S6. Time-lapse confocal fluorescent imaging of a *HS:mcherry-caax-p2a-NLS-kikume* **embryo in DMSO treatment.** *HS:mcherry-caax-p2a-NLS-kikume* marks nuclei within the somite. Sample tracks ranging from red (most displacement) to blue (least displacement) show broad movement within the somite. Frame rate= one image/5 min. Run time= 240 min.

Movie S7. Time-lapse confocal fluorescent imaging of a *HS:mcherry-caax*-p2a-*NLS-kikume* **embryo in DEAB treatment.** *HS:mcherry-caax-p2a-NLS-kikume* marks nuclei within the somite. Sample tracks ranging from red (most displacement) to blue (least displacement) show little directional movement in 20 µM DEAB treatment conditions. Frame rate= one image/5 min. Run time= 240 min.

Movie S8. Time-lapse fluorescent movie of *mcherry-caax* **mRNA injected** *tg(kdrl:eGFP)* **explant at the 10 somite-stage**. *tg(kdrl:eGFP)* marks angioblasts as they migrate to the midline and *mcherry-caax* marks the cell surfaces. A trunk explant, imaged around the 5th somite, shows normal migration of the angioblasts to the midline. Frame rate= one image/5 min. Run time= 280 min.

1021

1022   **Movie S9. Time-lapse fluorescent movie of *mcherry-caax* mRNA injected**

1023   ***tg(kdrl:eGFP)* explant at the 10 somite-stage, treated with 20 µM DEAB.**

1024   *tg(kdrl:eGFP)* marks angioblasts as they migrate to the midline. The trunk explant shows

1025   angioblast migration defects seen in fixed DEAB sections. Frame rate= one image/5 min.

1026   Run time= 175 min.

1027